Long-Term Oral Tamoxifen Administration Decreases Brain Derived Neurotrophic Factor in the Hippocampus of Female Long-Evans Rats

Laura E. Been[1], Amanda R. Halliday[1], Sarah M. Blossom[1], Elena M. Bien[1], Anya G. Bernhard[1], Grayson E. Roth[1], Karina I. Domenech Rosario[1], Karlie B. Pollock[1], Petra E. Abramenko[1], Leily M. Behbehani[1], Gabriel J. Pascal[1], and Mary Ellen Kelly[1,2]

[1] Haverford College, Department of Psychology and Neuroscience Program, Haverford, PA, USA

[2] University of Pennsylvania, Neuroscience Program, Philadelphia, PA, USA

Abstract

Tamoxifen is a selective estrogen receptor modulator (SERM) that is commonly used as an adjuvant drug therapy for estrogen-positive breast cancers. While this drug is effective at reducing the rate of cancer recurrence, many patients report unwanted cognitive and affective side effects such as brain fog, confusion, memory impairment, anxiety, and depression. Despite this, the impacts of chronic tamoxifen exposure on the brain are poorly understood, and rodent models of tamoxifen exposure do not replicate the chronic oral administration seen in patients. We therefore used long-term *ad lib* consumption of medicated food pellets in adult female rats to model chronic tamoxifen exposure in a clinically-relevant way. Gonadally-intact adult female Long-Evans Hooded rats consumed tamoxifen medicated food pellets for approximately 12 weeks while control animals received standard chow. At the conclusion of the experiment, animals were euthanized, and blood and brain samples were collected for analyses. Blood tamoxifen levels were measured using a novel ultra-performance liquid chromatography-tandem mass spectrometry assay, which found that this administration paradigm produced serum levels of tamoxifen similar to those in human patients. In the brain, brain derived neurotrophic factor (BDNF) was visualized in the hippocampus using immunohistochemistry and quantified using background-subtracted optical densitometry. Chronic oral tamoxifen treatment resulted in a decrease in BDNF expression across several regions of the hippocampus. Together, these findings provide a novel method of modeling and measuring chronic oral tamoxifen exposure, and suggest a putative mechanism by which tamoxifen may cause cognitive and behavioral changes reported by patients.

Introduction

Tamoxifen is a common adjuvant drug therapy used to decrease the recurrence risk of estrogen receptor-positive breast cancers (Jebahi et al., 2021; Kadakia & Henry, 2015; Novick et al., 2020). The mechanism of action of tamoxifen depends on its location in body tissue and the local estradiol environment (Novick et al., 2020). Generally, tamoxifen acts as an antagonist in the presence of endogenous estradiol (*e.g.*, in ovary-intact, pre-menopausal females), but is agonistic and protects against the effects of estrogen depletion in the absence of endogenous estradiol (Azizi-Malekabadi et al., 2015; Novick et al., 2020). Thus, it has been termed a selective estrogen receptor modulator (SERM). Patients typically take tamoxifen orally for 5-10 years following or concurrent with primary cancer treatment (Kadakia & Henry, 2015).

While tamoxifen is effective at decreasing the risk of cancer recurrence due to its antagonistic effects in breast tissue (Kadakia & Henry, 2015), people undergoing chronic tamoxifen treatment report unwanted psychological side effects such as anxiety, depression, brain fog, confusion, and memory impairment (Jebahi et al., 2021; Lee et al., 2016; Novick et al., 2020; Underwood et al., 2018). Long-term tamoxifen therapy has been shown to decrease performance on visuospatial, visual, and verbal memory tasks (Boele et al., 2015; Castellon et al., 2004; Lejbak et al., 2010; Underwood et al., 2018), decision making tasks (X. Chen et al., 2014; Lejbak et al., 2010), and overall executive function (Boele et al., 2015; Jenkins et al., 2004). However, it has been difficult to separate the effects of tamoxifen from the effects of primary cancer treatments, other adjuvant therapies such as aromatase inhibitors (Breckenridge et al., 2012) and the psychosocial effects of being a cancer survivor (Ahles et al., 2012).

Therefore, more research is needed to fully understand the effects of tamoxifen on the brain and cognition.

  To understand the impact of tamoxifen on the brain and behavior, some researchers have turned to rodent models. Like in people, tamoxifen treatment in rodents can lead to affective and cognitive changes; however, the impact of tamoxifen on cognitive-behavioral outcomes varies depending on the hormone status of the subjects (Novick et al., 2020). For example, tamoxifen administration to gonad-intact female rodents has been shown to increase anxiety- and depressive-like behaviors (Azizi-Malekabadi et al., 2015), decrease memory consolidation and retrieval ability on step-down avoidance tasks (D. Chen et al., 2002), and decrease novel object recognition ability (Valvassori et al., 2017). In ovariectomized rodents, on the other hand, tamoxifen administration is protective against anxiety- and depressive-like behaviors (Azizi-Malekabadi et al., 2015) and improves working memory ability (Velázquez-Zamora et al., 2012). Because tamoxifen is more commonly used in pre-menopausal patients (Kadakia & Henry, 2015), its effects on gonadally-intact female rodents are of particular interest. Notably, previous rodent studies have not modeled chronic oral tamoxifen use and instead have used a variety of tamoxifen administration methods including long-term injections (Cechinel-Recco et al., 2012), short-term or one-time injections (D. Chen et al., 2002; Kight & McCarthy, 2017; Velázquez-Zamora et al., 2012), and short-term gavages (Valvassori et al., 2017). This study aimed to fill in gaps in the literature by testing the effects of long-term oral tamoxifen administration on neurobiology in intact female rats.

The hippocampus is a strong putative candidate to mediate some of the cognitive side effects of tamoxifen therapy for several reasons. First, the hippocampus expresses ⬜- and β-estrogen receptors (ER⬜ and ERβ, respectively) across its various subregions (Frick et al., 2015; Harte-Hargrove et al., 2013; Spencer et al., 2008; Woolley, 1998). Research in rodents has demonstrated that estradiol fluctuations impact hippocampal neurophysiology (Liu et al., 2008; Woolley, 1998), which in turn affects depressive- and anxiety-like behaviors (Green & Galea, 2008; Walf & Frye, 2006) as well as performance on spatial memory tasks (Frick et al., 2015; Spencer et al., 2008). Hippocampal subregions such as the dentate gyrus (DG), Cornu Ammonis 1 (CA1), and Cornu Ammonis 3 (CA3) are of particular interest because they mediate spatial and emotional memory (Luine & Frankfurt, 2013; Spencer et al., 2008), which are impaired after long-term tamoxifen use (Novick et al., 2020). Within hippocampal neurons, Brain-Derived Neurotrophic Factor (BDNF) may play an important role in mediating tamoxifen's effects on cognition and emotion. Estrogen receptor activation influences BDNF transcription via estrogen response elements (EREs) that bind to the *BDNF* gene (Harte-Hargrove et al., 2013; Scharfman & MacLusky, 2006; Sohrabji & Lewis, 2006; Spencer et al., 2008). Indeed, fluctuating levels of estradiol during the estrous cycle lead to predictably fluctuating levels of BDNF expression; these changes are particularly salient in the hippocampus (Harte-Hargrove et al., 2013; Luine & Frankfurt, 2013; Scharfman & MacLusky, 2014). The downstream effects on BDNF are important for overall cognitive function. Mature BDNF binds to trkB receptors, which activates signaling cascades that are important for processes such as synaptogenesis and long-term potentiation (LTP) in the hippocampus (Harte-Hargrove et al., 2013). Finally,

estradiol has been shown to mediate both synaptogenesis (Sato et al., 2007) and LTP (Spencer et al., 2008) through its interactions with BDNF.

Taken together, this evidence suggests tamoxifen's impact on cognition and emotion may be mediated by BDNF expression in the hippocampus. We therefore hypothesized that long-term oral tamoxifen administration would lead to decreased BDNF expression in the hippocampus. To test this hypothesis, we established a model of long-term oral tamoxifen administration in adult, intact female Long-Evans rats. Using a novel ultra-performance liquid chromatography-tandem mass spectrometry (LC-MS/MS) detection method, we found that this long-term oral administration produced serum levels of tamoxifen comparable to those from Tamoxifen administration in people (Farrar & Jacobs, 2022). Brain tissue analyses found that tamoxifen-treated animals had significantly lower BDNF expression in the DG, dorsal CA3, and CA1. These results provide an important first step in characterizing the effect of long-term oral tamoxifen administration on the hippocampus, and may increase our understanding of the cognitive and affective side effects associated with long-term tamoxifen use.

Methods

Subjects

For all experiments, adult female Long-Evans Hooded rats were purchased from Charles River Laboratories (Wilmington, Massachusetts) at 3-4 months of age. Animals were housed in pairs in standard cages (17.7 x 9.4 in x 8.26 in; Allentown LLC, Allentown, NJ, USA) containing standard TekFresh cellulose low-dust rat bedding (Teklad 7099; Inotiv, West Lafayette, IN, USA), and a Plexiglas tube (Bio-Serv,

Flemington, NJ, USA) and wooden block (Bio-Serv, Flemington, NJ, USA) for enrichment. Subjects were kept on a 12-hour light-dark cycle with *ad libitum* food and water access.

Tamoxifen Self-Administration

Animals were divided into two experimental groups: tamoxifen (n = 20) and control (n = 16). Animals in the tamoxifen group consumed tamoxifen ad libitum for 10-13 weeks via medicated food pellets (Envigo, Indianapolis, IN) that were custom developed to simulate the standard dose of tamoxifen prescribed to premenopausal women diagnosed with breast cancer (20 mg/kg daily) while control animals received an ingredient-matched standard chow (16% protein, 55% carb, 3.4% fat) that did not contain Tamoxifen (Envigo) (Farrar & Jacobs, 2022).

Body weight and condition were monitored daily throughout the drug administration period. Animals in the tamoxifen group lost weight relative to controls (see results), so tamoxifen food pellets were mashed in a flavor-enhanced, nutritionally-fortified gel to enhance palatability (DietGel Recovery, Clear H20, Portland, ME, USA).

Blood Analysis

Following 10-13 weeks of access to medicated or control pellets, animals were euthanized via intracardial perfusion (see below). Blood samples were collected from the inferior vena cava immediately prior to perfusion and stored in heparin-coated vacutainer blood collection tubes (BD Medical, Franklin Lakes, NJ, USA) on ice until centrifugation. Samples were centrifuged at 2300 g at 4°C for 20 min and serum was stored at -80□ until assay.

An ultra-performance liquid chromatography-tandem mass spectrometry assay with reversed-phase chromatographic separation employing a Waters XBridge C18 (100 x 2.1 mm, 100 Å, 3.5 μm) and a runtime of 4.5 minutes was used to quantify

tamoxifen levels. Rat plasma samples (100 µL) were extracted with acetonitrile containing tamoxifen-d5 (5 ng/mL) and 0.1% formic acid. Two µL of the extract were injected onto the UPLC-MS/MS system for analysis.

Tamoxifen and tamoxifen-d5 were separated by ultra-performance liquid chromatography (UPLC) and detected using a triple quadrupole mass spectrometer (API4000). Tandem mass spectrometry (MS-MS) parameters were optimized in the positive ionization mode and the multiple reaction monitoring (MRM) transition of m/z 372.3 → 72.0 was used for tamoxifen analysis. Tamoxifen-d5 (m/z 377.3 → 72.0) was used as an internal standard. A calibration curve for tamoxifen was prepared in human plasma and was linear over the range of 0.1 – 250 ng/mL (r^2 > 0.99, **Figure 2, Table 1)**. Calibration standards and blanks in human plasma were extracted by protein precipitation with 400 µL of 5 ng/mL of tamoxifen-d5 in acetonitrile. Five µL of extract was injected for LC-MS/MS analysis. Tamoxifen and tamoxifen-d5 were separated on an XBridge C18 column (100 x 2.1 mm, 100 Å, 3.5 µm) using 0.1 % formic acid in water and 0.1 % formic acid in acetonitrile as aqueous and organic mobile phases with a flow rate of 0.5 mL/min. Analytes were separated using a gradient elution, with retention time of 2.68 min for tamoxifen and tamoxifen-d5. The LC-MS Tamoxifen assay used here was based on assays initially developed and validated in human plasma (Bobin-Dubigeon et al., 2019; Rama Raju et al., 2015). Therefore, as a confirmatory assay, human plasma and rat plasma were compared using 100 ng/mL concentrations of tamoxifen and found to be comparable (**Table 2**).

Tissue Histology and Immunohistochemistry

At the conclusion of the experiment, subjects were sacrificed by intracardial perfusion. Briefly, subjects were given an overdose of sodium pentobarbital (Beuthanasia-D, 22 mg/100 g body weight, Merck Animal Health, Madison, NJ, USA) and then transcardially perfused with approximately 250 mL of 25mM phosphate buffered saline (PBS) followed by approximately 250 mL of 4% paraformaldehyde in PBS. Brains were immediately extracted and post-fixed in 4% paraformaldehyde overnight and then cryoprotected for 48 h in 30% sucrose in PBS. Coronal sections (35 μm) were sectioned at -20°C on a cryostat (Leica Biosystems, Wetzlar, Germany), collected in a 1:6 series, and stored in cryoprotectant until immunohistochemical processing.

For immunohistochemistry, hippocampal sections were removed from cryoprotectant and rinsed 5 x 5 minutes in 25 mM PBS. After PBS washes, sections were incubated in hydrogen peroxide (1:100) for 15 minutes to reduce endogenous peroxidase activity. Sections were then washed 5 x 5 minutes in 25 mM PBS and before incubation in a rabbit monoclonal primary antibody against brain-derived neurotrophic factor (1:3000, Cat # ANT-010, Alomone Labs, Jerusalem, Israel) in 0.4% Triton-X 100 in PBS for 24 hours at room temperature. This antibody has previously been validated for use in rodent tissue (Louveau et al., 2015; Taliaz et al., 2010).

After incubation in the primary antibody, sections were rinsed in PBS (5 x 5 minutes) and incubated for one hour in a biotinylated secondary antibody (1:600, Biotin-SP AffiniPure Goat Anti-Rabbit IgG, Jackson Immunoresearch Labs; 111-065-003) in 0.4% Triton-X 100 in PBS. Sections were then rinsed in PBS (5 x 5 minutes) and incubated in avidin-biotin complex with horseradish peroxidase (per manufacturer's

instructions, Vectastain Elite ABC-HRP Kit, Vector Laboratories, Burlingame, CA, USA).
After 4 x 5 minute washes in PBS, sections were washed 2 x 5 minutes in a sodium
acetate buffer and then incubated for 10 minutes in 3,3'-diaminobenzidine HCl (0.2
mg/mL) in a nickel sodium acetate buffer (0.025 g Nickel II Sulfate/mL NaOAc) and
hydrogen peroxide (0.83 µL/mL) solution.

Stained tissue was then washed in sodium acetate buffer (2 x 5 min), washed in
PBS (3 x 5 min) mounted on glass slides (SuperFrost Plus Microscope Slides,
ThermoFisher Scientific, Waltham, MA, USA) and left to air-dry overnight. After drying,
slides were dehydrated with alcohols, cleared with xylenes and coverslipped with
Permount (ThermoFisher Scientific, Waltham, MA, USA).

BDNF Densitometry Analysis in Hippocampus

Images were acquired at 10X magnification using a Nikon E200 brightfield
microscope with a color camera and Spot Basic software (Diagnostic Instruments,
Sterling Heights, MI, USA). Five evenly-spaced template images (S1, S2, S3, S4, S5)
were selected from the rat brain atlas (Paxinos & Watson, 2013) to capture the rostral to
caudal extent of the hippocampus. The CA3 was split into dorsal and ventral regions for
imaging and analysis in order to capture it fully. Twenty images were collected for each
animal: five images per hippocampus region (DG, CA1, dCA3, vCA3).

Densitometry analyses were performed by experimenters who were blind to the
condition of the subject using ImageJ software (imagej.net/software/fiji; Schindelin et al.,
2012). Mean values for background optical density were collected from 36x36 pixel
regions that did not contain BDNF-ir cells using the "mean gray value" tool. Mean optical
density values for regions of interest (ROIs) were calculated after tracing the

hippocampus region of interest with the polygon tool and analyzing with the "mean gray value" tool (**Figure 4**). Background optical density was subtracted from region of interest optical density using the following equation: $(OD) = \log (255 / (OD_{ROI} - OD_{background}))$.

Statistical Analysis

All statistics were run using Jamovi (www.jamovi.org) with alpha values of 0.05. Body weight was compared between groups using independent samples t-tests. Two-way ANOVAs were used to look for interactions and main effects of drug condition (tamoxifen vs. control) and section level (1-5) on BDNF-ir density within each of the four ROIs (DG, CA1, vCA1, and CA3). Individual brain regions were analyzed separately. Significant effects were explicated using Fisher's LSD post-hoc comparisons. Individual samples were excluded from analysis if tissue damage precluded OD analysis, or if there was no adequate match for the template atlas plate. The relationship between serum tamoxifen levels and BDNF-ir density was examined using simple correlations.

Results

Body Weight

Animals in both groups had significant body weight changes during the course of the experiment. Control animals gained weight ($t(30) = 6.14$, $p < .001$) relative to baseline, whereas animals receiving tamoxifen (TAM) lost weight relative to baseline ($t(30) = 2.31$, $p = .028$, **Figure 1**).

Plasma Tamoxifen Levels

Tamoxifen was not found in any control samples. In tamoxifen-treated animals, plasma concentrations ranged from 10.7 to 48.3 ng/mL ($M = 28.1$, $SD = 11.2$, **Table 3**). Representative chromatograms can be found in **Figure 2**.

BDNF Immunoreactivity

For each brain region of interest, the density of BDNF-immunoreactive (ir) neurons in the DG, dCA3, vCA3, and CA1 was analyzed and compared between groups. In the DG, there was a significant main effect of drug condition on the density of BDNF-ir cells ($F(1, 146) = 9.454$, $p = 0.003$). Post-hoc tests revealed that BDNF-ir density was lower in tamoxifen-treated animals than control animals in Section 4 ($p = 0.03$, **Figure 3A**). In dCA3, there was a significant main effect of drug condition on the density of BDNF-ir cells ($F, 1, 144) = 11.94$, $p < 0.001$). Post-hoc tests revealed that BDNF-ir was lower in tamoxifen-treated animals than control animals in Section 3 ($p < 0.01$, **Figure 3B**). There was also a significant main effect of section on the density of BDNF-ir cells in dCA3 ($F(4, 144) = 3.302$, $p = 0.01$). Post-hoc tests revealed that Section 2 differed from Section 4 ($p = 0.02$) and Section 5 ($p = 0.03$), and Section 3 differed from Section 4 ($p < 0.01$) and Section 5 ($p < 0.01$). In vCA3, there was no effect of drug condition ($F(1, 145) = 0.071$, $p = 0.79$) or section ($F(1,145) = 0.87$, $p = 0.48$) on the density of BDNF-ir cells (**Figure 3C**). Finally, in CA1, there was a main effect of drug condition on the density of BDNF-ir cells ($F(1, 146) = 8.886$, $p = 0.003$). Post hoc tests revealed that BDNF-ir density was lower in tamoxifen-treated animals than control animals in section 2 ($p = 0.04$) and section 4 ($p = 0.02$, **Figure 3D**).

Animals' plasma tamoxifen concentration at the time of sacrifice did not correlate significantly with BDNF-ir cells in the DG ($r = -.4221$, $p = .2244$, **Figure 4A**) or CA1 ($r = -.2053$, $p = .5693$, **Figure 4B**). In both dorsal and ventral CA3 (**Figure 4C and 4D**), there was a trend towards increased plasma tamoxifen concentration being associated with

decreased BDNF-ir cells, although these did not reach significance (dCA3 r = -.6112, p = .0604; vCA3 r = -.6168, p = .0575).

Discussion

Here we demonstrate for the first time that long-term oral tamoxifen administration decreases BDNF expression in the hippocampus of female Long-Evans Hooded rats. Notably, we show that a novel administration method, long-term *ad lib* exposure to medicated food pellets, produces plasma tamoxifen levels that are similar to a human dose (20 mg/kg; Farrar & Jacobs, 2022). Further, we used a novel LC-MS/MS assay to detect tamoxifen levels in rat plasma. Together, these results provide a new way to model and measure the effects of long-term oral tamoxifen administration on the brain, which may be particularly useful for investigating mechanisms of cognitive and behavioral changes reported by patients taking Tamoxifen.

Voluntary oral administration via medicated food pellets is a clinically relevant, non-invasive, and effective method for modeling the effects of tamoxifen in rodents. Indeed, compared with systemic injections (Cechinel-Recco et al., 2012; D. Chen et al., 2002; Kight & McCarthy, 2017; Velázquez-Zamora et al., 2012); or oral gavage (Valvassori et al., 2017), tamoxifen oral self-administration is an attractive alternative in that it more closely models the metabolism of tamoxifen use in patient populations and eliminates stress associated with restraint, repeated injections, or gavage as potential confounding variables. Nonetheless, there are some challenges with this method. First, there was variability in individual animals' plasma tamoxifen concentrations. These differences likely resulted from differences in voluntary consumption across animals, both throughout the experiment and specifically on the day of euthanasia. Blood draws

were collected approximately 60 minutes after the last tamoxifen pellets were presented, but the amount of medicated pellet consumed, or the timing of consumption within the one-hour window, were not measured. Future studies should collect data about volume and timing of food intake, particularly prior to blood collection.

In addition, tamoxifen-fed animals lost weight across the course of the experiment, while control animals gained weight across the course of the experiment. It is unclear if the weight loss in the tamoxifen group was due to the tamoxifen food pellets being less palatable than standard chow, or because decreased appetite or weight loss is an effect of tamoxifen treatment. Presenting the tamoxifen pellets as a mash in a flavor-enhanced gel increased consumption, suggesting that palatability was at least part of the issue. Short-term tamoxifen injections induce anorexia in rodents (Gray et al., 1993; López et al., 2006; Wade & Heller, 1993; Walker et al., 2011, p. 2006); as such, an anorexic effect of tamoxifen cannot be ruled out. A similar study in mice found that 15 days of exposure to tamoxifen chow also led to weight loss, particularly in female subjects; animals did not consistently eat TAM chow for the first 5 days of exposure (Smith et al., 2022). While oral gavage would introduce stress and is therefore not recommended, future experiments may wish to explore mild food deprivation or pair feeding tamoxifen-treated and control animals to eliminate the effects of body weight as a potential confound on dependent variables.

Our finding that long-term oral tamoxifen administration decreased BDNF expression in the hippocampus may point towards a putative mechanism underlying cognitive side effects that patients taking tamoxifen report. Given the known association of estrogen receptor activation and BDNF transcription (Harte-Hargrove et al., 2013;

Scharfman & MacLusky, 2006; Sohrabji & Lewis, 2006), it is possible that tamoxifen's action at estrogen receptors in the brain could disrupt BDNF transcription, altering plasticity and neuronal stability in the hippocampus. These changes to hippocampal function could in turn impair cognitive function and lead to clinical symptoms like brain fog, memory deficits, and alterations in mood/anxiety. In support of this idea, Smith et al. (2022) found that short term TAM administration impacts hippocampal neurogenesis. Specifically TAM injection is associated with decreased progenitor cell proliferation in the dentate gyrus, although TAM ingestion is associated with increased neuronal differentiation in the dentate gyrus. Thus, our finding that long-term oral tamoxifen administration decreased BDNF density in the current study may be due to a decrease in cell proliferation, or conversely, decreases in BDNF following TAM treatment could contribute to changes in neurogenesis.

Future studies using this model of long-term oral tamoxifen administration should include behavioral assays of cognition, mood, and anxiety. While we were unable to collect a complete behavioral cohort due to laboratory shutdown during the Covid-19 pandemic, preliminary data from our lab suggests that tamoxifen-fed animals show more anxiety-like behavior in the open field compared to controls, but no deficits in elevated plus or novel object recognition. Our finding that long-term tamoxifen administration decreases BDNF in the DG, CA1, and dCA3 suggests that pattern separation, spatial memory, associative memory, and episodic memory processes could be particularly affected (Bartsch et al., 2011; Cherubini & Miles, 2015; Kesner, 2007; Langston et al., 2010). Disparate findings in the dorsal vs. ventral CA3 subregions further point towards tamoxifen's effects on cognitive function. For example, there is

some literature to suggest that the dorsal region is responsible for cognition and information processing, while the ventral region is more closely associated with stress and affect (Fanselow & Dong, 2010). A full behavioral profile of animals during long-term oral tamoxifen treatment, such as further assays of cognition (e.g., object-in-place, attentional set shifting), anxiety-like behaviors (e.g., defensive burying, light-dark box) and learned helplessness or anhedonia (e.g., forced swim, tail suspension, splash test, sucrose preference) would be useful to characterize tamoxifen's effects on behavior in this model, and may suggest additional brain regions for exploration. Further, while this study focused on the hippocampus, it is likely that tamoxifen impacts other estrogen-sensitive brain regions that are associated with cognition, anxiety, and/or mood disorders, such as the nucleus accumbens and amygdala. Tamoxifen could disrupt BDNF transcription in these regions by interacting with estrogen receptors via the same putative mechanism that we have identified in the hippocampus.

In addition to a novel administration paradigm, we also used a novel, sensitive, and reproducible method for the analysis of tamoxifen that is suitable for rat plasma. This will be useful to future researchers interested in the physiological effects of oral tamoxifen administration. While not significant, we did observe a trend towards correlation between plasma tamoxifen concentration and BDNF density in dCA3 and vCA3, supporting tamoxifen's potential role in decreasing BDNF transcription in the hippocampus. It is worth noting that tamoxifen is also commonly used in neuroscience research as a tool to induce Cre mechanisms. While these research applications are markedly different in that the doses are generally much higher and administration is acute (often a single injected dose) (Donocoff et al., 2020; Heffner, 2011) our research

suggests that tamoxifen administration affects BDNF expression, and researchers using tamoxifen in their work should consider its potential off-target effects on neurobiology.

Tamoxifen remains the gold-standard medication for preventing recurrence of ER+ breast cancers. Its ongoing use as an adjuvant therapy will likely continue despite common affective and cognitive side effects; tamoxifen treatments will enable breast cancer survivors to live longer and healthier lives. Further identifying and understanding the neural mechanisms underlying these side effects, however, could lead to promising interventions that would mitigate unwanted psychological changes without risking cancer recurrence. It is important that researchers and clinicians continue to explore ways to improve psychiatric health in at-risk and under-researched patient populations. Cancer survivors, women and other gender minorities in particular, are particularly vulnerable to a range of affective, anxiety, and cognitive disorders (Ahles et al., 2012; Di Giacomo et al., 2016; Fafouti et al., 2010). It is imperative that we find ways to support all aspects of their health as they navigate treatment.

Acknowledgements

This work was funded in part by Haverford College Department of Psychology. The authors wish to thank Shanice Edwards, Christina Vedar, Abigail Cohen, Athena Zuppa, and Ganesh Moorthy from the Children's Hospital of Philadelphia Research Institute Bioanalytical Core for developing and conducting the mass spectrometry assays. Some of the mass spectrometry data were previously presented as a poster at the American Society for Mass Spectrometry Conference on Mass Spectrometry and Allied Topics in

Philadelphia, PA in November of 2021. The authors also wish to thank Dr. Louise Charkoudian for her assistance interpreting the LC-MS data.

References

Ahles, T. A., Root, J. C., & Ryan, E. L. (2012). Cancer- and Cancer Treatment–Associated Cognitive Change: An Update on the State of the Science. *Journal of Clinical Oncology*, *30*(30), 3675–3686. https://doi.org/10.1200/JCO.2012.43.0116

Azizi-Malekabadi, H., Pourganji, M., Zabihi, H., Saeedjalali, M., & Hosseini, M. (2015). Tamoxifen antagonizes the effects of ovarian hormones to induce anxiety and depression-like behavior in rats. *Arquivos de Neuro-Psiquiatria*, *73*, 132–139. https://doi.org/10.1590/0004-282X20140221

Bartsch, T., Döhring, J., Rohr, A., Jansen, O., & Deuschl, G. (2011). CA1 neurons in the human hippocampus are critical for autobiographical memory, mental time travel, and autonoetic consciousness. *Proceedings of the National Academy of Sciences*, *108*(42), 17562–17567. https://doi.org/10.1073/pnas.1110266108

Bobin-Dubigeon, C., Campone, M., Rossignol, E., Salaun, E., Amiand, M.-B., & Bard, J.-M. (2019). New UPLC–MS/MS assay for the determination of tamoxifen and its metabolites in human plasma, application to patients. *Future Science OA*, *5*(5), FSO374. https://doi.org/10.2144/fsoa-2018-0113

Boele, F. W., Schilder, C. M. T., de Roode, M.-L., Deijen, J. B., & Schagen, S. B. (2015). Cognitive functioning during long-term tamoxifen treatment in postmenopausal women with breast cancer. *Menopause*, *22*(1), 17–25. https://doi.org/10.1097/GME.0000000000000271

Breckenridge, L. M., Bruns, G. L., Todd, B. L., & Feuerstein, M. (2012). Cognitive limitations associated with tamoxifen and aromatase inhibitors in employed breast cancer survivors. *Psycho-Oncology*, *21*(1), 43–53.

https://doi.org/10.1002/pon.1860

Castellon, S. A., Ganz, P. A., Bower, J. E., Petersen, L., Abraham, L., & Greendale, G. A. (2004). Neurocognitive Performance in Breast Cancer Survivors Exposed to Adjuvant Chemotherapy and Tamoxifen. *Journal of Clinical and Experimental Neuropsychology*, *26*(7), 955–969. https://doi.org/10.1080/13803390490510905

Cechinel-Recco, K., Valvassori, S. S., Varela, R. B., Resende, W. R., Arent, C. O., Vitto, M. F., Luz, G., de Souza, C. T., & Quevedo, J. (2012). Lithium and tamoxifen modulate cellular plasticity cascades in animal model of mania. *Journal of Psychopharmacology*, *26*(12), 1594–1604. https://doi.org/10.1177/0269881112463124

Chen, D., Wu, C. F., Shi, B., & Xu, Y. M. (2002). Tamoxifen and toremifene cause impairment of learning and memory function in mice. *Pharmacology Biochemistry and Behavior*, *71*(1), 269–276. https://doi.org/10.1016/S0091-3057(01)00656-6

Chen, X., Li, J., Chen, J., Li, D., Ye, R., Zhang, J., Zhu, C., Tian, Y., & Wang, K. (2014). Decision-making impairments in breast cancer patients treated with tamoxifen. *Hormones and Behavior*, *66*(2), 449–456. https://doi.org/10.1016/j.yhbeh.2014.07.005

Cherubini, E., & Miles, R. (2015). The CA3 region of the hippocampus: How is it? what is it for? how does it do it? *Frontiers in Cellular Neuroscience*, *9*. https://www.frontiersin.org/articles/10.3389/fncel.2015.00019

Di Giacomo, D., Cannita, K., Ranieri, J., V, C., Passafiume, D., & Ficorella, C. (2016). Breast cancer and psychological resilience among young women. *Journal of Psychopatology*, *3*, 1–5.

Donocoff, R. S., Teteloshvili, N., Chung, H., Shoulson, R., & Creusot, R. J. (2020).

    Optimization of tamoxifen-induced Cre activity and its effect on immune cell

    populations. *Scientific Reports*, *10*(1), Article 1. https://doi.org/10.1038/s41598-

    020-72179-0

Fafouti, M., Paparrigopoulos, T., Zervas, Y., Rabavilas, A., Malamos, N., Liappas, I., &

    Tzavara, C. (2010). Depression, Anxiety and General Psychopathology in Breast

    Cancer Patients: A Cross-sectional Control Study. *In Vivo*, *24*(5), 803–810.

Fanselow, M. S., & Dong, H.-W. (2010). Are The Dorsal and Ventral Hippocampus

    functionally distinct structures? *Neuron*, *65*(1), 7.

    https://doi.org/10.1016/j.neuron.2009.11.031

Farrar, M. C., & Jacobs, T. F. (2022). Tamoxifen. In *StatPearls*. StatPearls Publishing.

    http://www.ncbi.nlm.nih.gov/books/NBK532905/

Frick, K. M., Kim, J., Tuscher, J. J., & Fortress, A. M. (2015). Sex steroid hormones

    matter for learning and memory: Estrogenic regulation of hippocampal function in

    male and female rodents. *Learning & Memory*, *22*(9), 472–493.

    https://doi.org/10.1101/lm.037267.114

Gray, J. M., Schrock, S., & Bishop, M. (1993). Estrogens and antiestrogens: Actions

    and interactions with fluphenazine on food intake and body weight in rats.

    American Journal of Physiology-Regulatory, Integrative and Comparative

    Physiology, *264*(6), R1214–R1218.

    https://doi.org/10.1152/ajpregu.1993.264.6.R1214

Green, A. D., & Galea, L. A. M. (2008). Adult hippocampal cell proliferation is

    suppressed with estrogen withdrawal after a hormone-simulated pregnancy.

Hormones and Behavior, 54(1), 203–211.
https://doi.org/10.1016/j.yhbeh.2008.02.023

Harte-Hargrove, L. C., MacLusky, N. J., & Scharfman, H. E. (2013). Brain-derived
neurotrophic factor–estrogen interactions in the hippocampal mossy fiber
pathway: Implications for normal brain function and disease. *Neuroscience, 239*,
46–66. https://doi.org/10.1016/j.neuroscience.2012.12.029

Heffner, C. (2011, July 25). *Intraperitoneal Injection of Tamoxifen for Inducible Cre-
Driver Lines*. The Jackson Laboratory. https://www.jax.org/research-and-
faculty/resources/cre-repository/tamoxifen

Jebahi, F., Sharma, S., Bloss, J. E., & Wright, H. H. (2021). Effects of tamoxifen on
cognition and language in women with breast cancer: A systematic search and a
scoping review. *Psycho-Oncology, 30*(8), 1262–1277.
https://doi.org/10.1002/pon.5696

Jenkins, V., Shilling, V., Fallowfield, L., Howell, A., & Hutton, S. (2004). Does hormone
therapy for the treatment of breast cancer have a detrimental effect on memory
and cognition? A pilot study. *Psycho-Oncology, 13*(1), 61–66.
https://doi.org/10.1002/pon.709

Kadakia, K. C., & Henry, N. L. (2015). Adjuvant Endocrine Therapy in Premenopausal
Women with Breast Cancer. *Clinical Advances in Hematology & Oncology⃞:
H&O, 13*(10), 663–672.

Kesner, R. P. (2007). A behavioral analysis of dentate gyrus function. In H. E.
Scharfman (Ed.), *Progress in Brain Research* (Vol. 163, pp. 567–576). Elsevier.
https://doi.org/10.1016/S0079-6123(07)63030-1

Kight, K. E., & McCarthy, M. M. (2017). Sex differences and estrogen regulation of

    BDNF gene expression, but not propeptide content, in the developing

    hippocampus. *Journal of Neuroscience Research*, *95*(1–2), 345–354.

    https://doi.org/10.1002/jnr.23920

Langston, R. F., Stevenson, C. H., Wilson, C. L., Saunders, I., & Wood, E. R. (2010).

    The role of hippocampal subregions in memory for stimulus associations.

    Behavioural Brain Research, *215*(2), 275–291.

    https://doi.org/10.1016/j.bbr.2010.07.006

Lee, P. E., Tierney, M. C., Wu, W., Pritchard, K. I., & Rochon, P. A. (2016). Endocrine

    treatment-associated cognitive impairment in breast cancer survivors: Evidence

    from published studies. *Breast Cancer Research and Treatment*, *158*(3), 407–

    420. https://doi.org/10.1007/s10549-016-3906-9

Lejbak, L., Vrbancic, M., & Crossley, M. (2010). Endocrine therapy is associated with

    low performance on some estrogen-sensitive cognitive tasks in postmenopausal

    women with breast cancer. *Journal of Clinical and Experimental*

    Neuropsychology, *32*(8), 836–846. https://doi.org/10.1080/13803391003596389

Liu, F., Day, M., Muñiz, L. C., Bitran, D., Arias, R., Revilla-Sanchez, R., Grauer, S.,

    Zhang, G., Kelley, C., Pulito, V., Sung, A., Mervis, R. F., Navarra, R., Hirst, W.

    D., Reinhart, P. H., Marquis, K. L., Moss, S. J., Pangalos, M. N., & Brandon, N. J.

    (2008). Activation of estrogen receptor-β regulates hippocampal synaptic

    plasticity and improves memory. *Nature Neuroscience*, *11*(3), Article 3.

    https://doi.org/10.1038/nn2057

López, M., Lelliott, C. J., Tovar, S., Kimber, W., Gallego, R., Virtue, S., Blount, M.,

Vázquez, M. J., Finer, N., Powles, T. J., O'Rahilly, S., Saha, A. K., Diéguez, C., & Vidal-Puig, A. J. (2006). Tamoxifen-Induced Anorexia Is Associated With Fatty Acid Synthase Inhibition in the Ventromedial Nucleus of the Hypothalamus and Accumulation of Malonyl-CoA. *Diabetes*, *55*(5), 1327–1336. https://doi.org/10.2337/db05-1356

Louveau, A., Nerrière-Daguin, V., Vanhove, B., Naveilhan, P., Neunlist, M., Nicot, A., & Boudin, H. (2015). Targeting the CD80/CD86 costimulatory pathway with CTLA4-Ig directs microglia toward a repair phenotype and promotes axonal outgrowth. *Glia*, *63*(12), 2298–2312. https://doi.org/10.1002/glia.22894

Luine, V., & Frankfurt, M. (2013). Interactions between estradiol, BDNF and dendritic spines in promoting memory. *Neuroscience*, *239*, 34–45. https://doi.org/10.1016/j.neuroscience.2012.10.019

Novick, A. M., Scott, A. T., Epperson, C. N., & Schneck, C. D. (2020). Neuropsychiatric Effects of Tamoxifen: Challenges and Opportunities. *Frontiers in Neuroendocrinology*, *59*, 100869. https://doi.org/10.1016/j.yfrne.2020.100869

Paxinos, G., & Watson, C. (2013). *The Rat Brain in Stereotaxic Coordinates* (7th ed.). Academic Press. https://www.elsevier.com/books/the-rat-brain-in-stereotaxic-coordinates/paxinos/978-0-12-391949-6

Rama Raju, K. S., Taneja, I., Singh, S. P., Tripathi, A., Mishra, D. P., Hussain, K. M., Gayen, J. R., Singh, S. K., & Wahajuddin, M. (2015). Simultaneous determination of centchroman and tamoxifen along with their metabolites in rat plasma using LC-MS/MS. *Bioanalysis*, *7*(8), 967–979. https://doi.org/10.4155/bio.14.253

Sato, K., Akaishi, T., Matsuki, N., Ohno, Y., & Nakazawa, K. (2007). β-Estradiol induces

synaptogenesis in the hippocampus by enhancing brain-derived neurotrophic factor release from dentate gyrus granule cells. *Brain Research*, *1150*, 108–120. https://doi.org/10.1016/j.brainres.2007.02.093

Scharfman, H. E., & MacLusky, N. J. (2006). Estrogen and brain-derived neurotrophic factor (BDNF) in hippocampus: Complexity of steroid hormone-growth factor interactions in the adult CNS. *Frontiers in Neuroendocrinology*, *27*(4), 415–435. https://doi.org/10.1016/j.yfrne.2006.09.004

Scharfman, H. E., & MacLusky, N. J. (2014). Differential regulation of BDNF, synaptic plasticity and sprouting in the hippocampal mossy fiber pathway of male and female rats. *Neuropharmacology*, *76*, 696–708. https://doi.org/10.1016/j.neuropharm.2013.04.029

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D. J., Hartenstein, V., Eliceiri, K., Tomancak, P., & Cardona, A. (2012). Fiji: An open-source platform for biological-image analysis. *Nature Methods*, *9*(7), Article 7. https://doi.org/10.1038/nmeth.2019

Smith, B. M., Saulsbery, A. I., Sarchet, P., Devasthali, N., Einstein, D., & Kirby, E. D. (2022). Oral and Injected Tamoxifen Alter Adult Hippocampal Neurogenesis in Female and Male Mice. *ENeuro*, *9*(2), ENEURO.0422-21.2022. https://doi.org/10.1523/ENEURO.0422-21.2022

Sohrabji, F., & Lewis, D. K. (2006). Estrogen–BDNF interactions: Implications for neurodegenerative diseases. *Frontiers in Neuroendocrinology*, *27*(4), 404–414. https://doi.org/10.1016/j.yfrne.2006.09.003

Spencer, J. L., Waters, E. M., Romeo, R. D., Wood, G. E., Milner, T. A., & McEwen, B.
S. (2008). Uncovering the Mechanisms of Estrogen Effects on Hippocampal
Function. *Frontiers in Neuroendocrinology*, *29*(2), 219–237.
https://doi.org/10.1016/j.yfrne.2007.08.006

Taliaz, D., Stall, N., Dar, D. E., & Zangen, A. (2010). Knockdown of brain-derived
neurotrophic factor in specific brain sites precipitates behaviors associated with
depression and reduces neurogenesis. *Molecular Psychiatry*, *15*(1), 80–92.
https://doi.org/10.1038/mp.2009.67

The jamovi project (2022). jamovi (Version 2.3) [Computer Software]. Retrieved from
https://www.jamovi.org

Underwood, E. A., Rochon, P. A., Moineddin, R., Lee, P. E., Wu, W., Pritchard, K. I., &
Tierney, M. C. (2018). Cognitive sequelae of endocrine therapy in women treated
for breast cancer: A meta-analysis. *Breast Cancer Research and Treatment*,
168(2), 299–310. https://doi.org/10.1007/s10549-017-4627-4

Valvassori, S. S., Borges, C. P., Varela, R. B., Bavaresco, D. V., Bianchini, G., Mariot,
E., Arent, C. O., Resende, W. R., Budni, J., & Quevedo, J. (2017). The different
effects of lithium and tamoxifen on memory formation and the levels of
neurotrophic factors in the brain of male and female rats. *Brain Research
Bulletin*, *134*, 228–235. https://doi.org/10.1016/j.brainresbull.2017.08.006

Velázquez-Zamora, D. A., Garcia-Segura, L. M., & González-Burgos, I. (2012). Effects
of selective estrogen receptor modulators on allocentric working memory
performance and on dendritic spines in medial prefrontal cortex pyramidal
neurons of ovariectomized rats. *Hormones and Behavior*, *61*(4), 512–517.

https://doi.org/10.1016/j.yhbeh.2012.01.010

Wade, G. N., & Heller, H. W. (1993). Tamoxifen mimics the effects of estradiol on food

intake, body weight, and body composition in rats. *American Journal of*

Physiology-Regulatory, Integrative and Comparative Physiology, *264*(6), R1219–

R1223. https://doi.org/10.1152/ajpregu.1993.264.6.R1219

Walf, A. A., & Frye, C. A. (2006). A Review and Update of Mechanisms of Estrogen in

the Hippocampus and Amygdala for Anxiety and Depression Behavior.

Neuropsychopharmacology, *31*(6), Article 6.

https://doi.org/10.1038/sj.npp.1301067

Walker, E. A., Foley, J. J., Clark-Vetri, R., & Raffa, R. B. (2011). Effects of repeated

administration of chemotherapeutic agents tamoxifen, methotrexate, and 5-

fluorouracil on the acquisition and retention of a learned response in mice.

Psychopharmacology, *217*(4), 539–548. https://doi.org/10.1007/s00213-011-

2310-8

Woolley, C. S. (1998). Estrogen-mediated structural and functional synaptic plasticity in

the female rat hippocampus. *Hormones and Behavior*, *34*(2), 140–148.

https://doi.org/10.1006/hbeh.1998.1466

Figures and Tables

Figure 1. Body Weight. Body weight was tracked throughout the course of the experiment. Rats fed Tamoxifen chow (TAM) significantly lost weight, whereas rats fed standard control chow (CON) significantly gained weight. Data presented as mean ± standard error, * p < 0.05; *** p < 0.001.

Figure 2. Mass Spectrometry Analysis of Tamoxifen. Tamoxifen and tamoxifen-d5

were separated by ultra-performance liquid chromatography (UPLC) and detected using

a triple quadrupole mass spectrometer. Tandem mass spectrometry (MS-MS)

parameters were optimized in the positive ionization mode and the multiple reaction

monitoring (MRM) transition of m/z 372.3 72.0 was used for tamoxifen analysis. Tamoxifen-d5 (m/z 377.3 72.0) was used as an internal standard. A) A calibration curve for tamoxifen was prepared in human plasma and was linear over the range of 0.1 – 250 ng/mL. Analytes were separated using a gradient elution, with retention time of 2.68 min for tamoxifen and tamoxifen-d5. B) Representative chromatogram from Tamoxifen-treated animal of Tamoxifen (left) and Tamoxifen-d5 (right). C) Representative chromatogram from control-treated rat (standard chow) of Tamoxifen (left) and Tamoxifen-d5 (right) in plasma.

Figure 3. BDNF Expression in Hippocampal Subregions. For each brain region of interest, immunohistochemical localization of BDNF was quantified using background-subtracted optical density in five sections capturing the rostral to caudal extent of the hippocampus. Significant interactions and/or main effects of drug condition (Tamoxifen, TAM vs. Control, CON) and section number (1-5) on BDNF optical density were detected using two-way ANOVAs. A) In DG, there was a significant main effect of drug condition on BDNF-ir density. Post-hoc tests revealed that BDNF-ir density was lower in

TAM animals than CON animals in Section 4. B) In dCA3, there was also a significant main effect of drug condition on the density of BDNF-ir cells. Post-hoc tests revealed that BDNF-ir was lower in TAM animals than CON animals in Section 3. C) In CA1, there was a main effect of drug condition on the density of BDNF-ir cells. Post hoc tests revealed that BDNF-ir density was lower in TAM animals than CON animals in section 2. D) Finally, in vCA3, there were no interactions or main effects of drug condition or section level on BDNF immunoreactivity. Templates atlas sections and representative tracing for densitometry analyses shown below graphs. Data presented as mean ± standard error, asterisks indicate significant omnibus tests, * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$. Pound signs indicate significant post-hoc differences between drug conditions, # $p < 0.05$; ## $p < 0.01$; ### $p < 0.001$

Figure 4. Plasma Tamoxifen and Brain BDNF Correlations. The relationship

between plasma tamoxifen levels and BDNF expression in hippocampal subregions

was investigated using simple correlations. In all four brain regions of interest, plasma

tamoxifen was negatively correlated with brain BDNF levels. Although none of these

correlations reached significance, trends towards significance were detected in C) dCA3

and D) vCA3. Pearson's correlation coefficient (r) and p values for each brain area

presented.

Table 1. Summary of Tamoxifen calibration curve in plasma. A calibration curve for tamoxifen was prepared in human plasma and was linear over the range of 0.1 – 250 ng/mL with coefficient of regression, $r^2 > 0.99$. Data presented as Mean, Standard Deviation (SD), Accuracy (%), and Coefficient of Variation (CV %).

Calibration Curve (n=6)	Tamoxifen Concentration (ng/mL)									
	0.1	0.5	1	2.5	5.0	10	25	50	100	250
1	0.108	0.538	1.00	2.61	5.11	10.5	29.4	46.4	108	270
2	0.117	0.533	1.11	2.61	4.84	10.3	27.4	48.4	107	253
3	0.104	0.484	0.95	2.69	4.79	10.2	29.2	54.0	109	247
4	0.090	0.479	0.88	2.08	4.25	9.55	25.2	47.9	95.6	266
5	0.091	0.566	0.922	2.49	5.06	10.5	27.9	52.9	92.1	234
6	0.092	0.467	0.841	2.18	4.25	9.38	24.8	46.5	97.2	230
Mean	0.100	0.511	0.949	2.44	4.72	10.1	27.3	49.4	101	250
SD	0.011	0.040	0.096	0.253	0.382	0.487	1.95	3.29	7.35	16.3
Accuracy (%)	100	102	94.9	97.7	94.3	101	109	98.7	101	100
CV (%)	11.1	7.79	10.1	10.4	8.09	4.84	7.15	6.66	7.25	6.52

Table 2. Human plasma and rat plasma comparison. The Tamoxifen assay was initially developed in human plasma. Human plasma and rat plasma were compared using 100 ng/mL concentrations of tamoxifen. Results were comparable between human and rat plasma. Concentration (ng/mL), Data presented as Concentration (ng/mL), Mean, Standard Deviation (SD), Coefficient of Variation (% CV), and Accuracy (%).

	Concentration (ng/mL)	Mean ± SD	% CV	% Accuracy
Human Plasma	100	101 ± 7.35	7.25	101
Rat Plasma	100	97.62 ± 2.01	2.06	97.6

Table 3. Plasma Tamoxifen Concentration. Tamoxifen levels (ng/mL) in TAM-treated

rats ($n = 10$).

Subject	Measured Tamoxifen Concentration (ng/mL)
TAM 1	48.3
TAM 2	33.8
TAM 3	26.5
TAM 4	35.0
TAM 5	22.0
TAM 6	34.2
TAM 9	16.3
TAM 10	10.7
TAM 11	35.0
TAM 12	19.3