Synchronized LFP rhythmicity in the social brain reflects the context of social encounters

Alok Nath Mohapatra[1,*], David Peles[1], Shai Netser[1], Shlomo Wagner[1,*]

[1] Sagol Department of Neurobiology, Faculty of Natural Sciences, University of Haifa, Haifa, Israel

Short title: LFP rhythmicity in the social brain reflects the social context

*Corresponding author: Alok Nath Mohapatra, Sagol Department of Neurobiology, Faculty of

Natural Sciences, University of Haifa, Mt. Carmel, Haifa 3498838, Israel.

Email: thinkalok@gmail.com

Abstract

Mammalian social behavior is highly context-sensitive. Yet, little is known about the mechanisms that modulate social behavior according to its context. Recent studies have revealed a network of mostly limbic brain regions, here termed the "social brain", which regulates social behavior. We hypothesized that coherent theta and gamma rhythms reflect the organization of the social brain regions into functional networks in a context-dependent manner. To test this concept, we simultaneously recorded extracellular activity from multiple social brain regions in mice performing three social discrimination tasks. Local field potential (LFP) rhythmicity across all tasks was dominated by a general internal state. However, during stimulus investigation LFP rhythmicity was sensitive to stimulus characteristics. Specifically, the pattern of LFP coherence between the various regions reflected mainly the social context. Moreover, we found the ventral dentate gyrus to play a pivotal role in coordinating the context-specific rhythmic activity in the network.

Keywords: social context, social discrimination, social brain, theta rhythmicity, in vivo electrophysiology, local field potential

Brain-region name abbreviations

AcbC: Accumbens nucleus, core
AcbSh: Accumbens nucleus, shell
AhiAL: Amygdalo-hippocampal area, anterolateral part
BLA: Basolateral amygdaloid nucleus
BMP: Basomedial amygdaloid nucleus, posterior part
DMD: Dorsomedial hypothalamic nucleus, dorsal part
EA: Extended amygdala
IL: Infralimbic prefrontal cortex
LS: Lateral septum
MeAD: Medial amygdaloid nucleus, anterodorsal
Pir: Piriform cortex
PLH: Peduncular part of the lateral hypothalamus
PrL: Prelimbic prefrontal cortex
PVN: Paraventricular hypothalamic nucleus
vCA1: Field CA1 of the hippocampus, ventral part
vDG: Dentate gyrus, ventral part
VP: Ventral pallidum

Introduction

Mammalian social behavior is highly complex and dynamic, involving multiple types of distinct, sometimes even opposing, interactions between partners. Indeed, the identity of a partner can completely change the nature and trajectory of social actions taken by an individual [1]. In addition to these complexities, social interactions are highly dependent upon the social context. For example, humans will most likely respond differently to a hand placed upon their shoulder from behind if this happens in a frightening context, such as in a dark alley in a foreign city, then if the same contact occurs in a cocktail party. Presently, little is known of the brain mechanisms and neural circuits that encode the context of social encounters and change responses to social cues accordingly.

In the last two decades, studies have begun to reveal the brain circuits that sub-serve various types of social behavior [for recent review papers see for example 2, 3-5]. Such studies exposed the involvement of a vast network of limbic brain regions, here termed the "social brain" [6], in processing social sensory cues and regulating mammalian social behavior [7, 8]. These include striatal regions, such as the nucleus accumbens core (AcbC) and shell (AcbSh), the prelimbic (PrL) and infralimbic (IL) prefrontal cortical areas, several hippocampal and septal areas and multiple amygdaloid and hypothalamic nuclei [9-12]. Many of these areas are highly interconnected in a bidirectional manner [13-17], and some were shown to be involved in various, at times opposing, types of social behavior [9, 18-21]. It remains, however, unclear how this intricate network of brain areas generates the large repertoire of distinct types of social behavior. Recent studies using multi-site brain recordings from behaving animals have demonstrated that system-level neural activity in sub-networks of the social brain predicts individual social preferences [22] and decision-making [23] better than does local neural activity at any single brain region. These results thus suggest that coding of the various aspects of social behavior in the brain should be considered at the system level.

Oscillatory neural activity, mostly in the theta (4-12 Hz) and gamma (30-80 Hz) bands, was reported in many cortical and sub-cortical brain regions in various species [24-26], with its power being shown to intensify during demanding cognitive functions, such as learning [27-29] and social communication [30-32]. Furthermore, abnormal theta and gamma rhythms have been reported in multiple neurodevelopmental disorders [33-35], such as autism spectrum disorder (ASD) [36, 37]. Accordingly, one prominent hypothesis states that coherent manifestation of these

rhythms can dynamically coordinate the activity of neural ensembles dispersed over multiple brain

regions and link them into *ad hoc* functional networks [38].

In the present study, we hypothesized that coherent theta and gamma rhythms couple various

regions of the social brain into functional networks in a social context-dependent manner. In other

words, distinct social contexts dictate different patterns of coordinated rhythmic activity of

dispersed social brain neuronal ensembles, which in turn sub-serve context-dependent processing

of social cues and consequent behavioral responses. To test this hypothesis, we recorded

extracellular electrical activity simultaneously from multiple regions of the social brain in mice

performing three distinct binary social discrimination tasks (contexts). The same type of social

stimulus served as either preferred or less-preferred stimulus in all three tasks. Using this design,

we could link distinct patterns of rhythmic neural activity across the social brain to either stimulus

identity or its valence, or to the social context. Our results reveal that the pattern of coordinated

oscillatory activity (coherence) in the network is strongly correlated with the social context and

carries information that may be used to discriminate between distinct, albeit similar, social

contexts. Further, we revealed that the ventral dentate gyrus (vDG), an area previously linked to

contextual information [39, 40], seems to be involved in coordinating the coherent activity among

the various regions of the social brain.

Results

Analyzing the behavior of CD1 male mice during three distinct binary social discrimination

tasks

Using custom-built electrode arrays (EAr) we simultaneously acquired local field potential (LFP)

signals from up to 16 brain regions at a time (cumulative count: 18 regions; Figure S1A and Table

S1) during interactions of adult male mice (subjects; n=14) with various stimuli [41]. We aimed

to sample widespread social-behavior associated regions in the cortex (prefrontal and piriform),

striatum (nucleus accumbens and ventral pallidum), hippocampus (e.g. dentate gyrus and CA1),

septal nuclei (latera septum), amygdala (e.g. basolateral and medial) and hypothalamus (e.g.

dorsomedial and paraventricular nuclei). The location of each electrode was verified *post mortem*

[41], and since the targeting accuracy was limited, not all brain regions were recorded in each

subject (see Fig. S1A and Table S1 for details). For social contexts,, we employed three distinct

binary social discrimination tasks [46, 47], namely the social preference (SP), (Fig 1A), emotional-

state preference (EsP) (Fig. 1E) and sex preference (SxP) (Fig 1I) tasks [46]. (See timeline in Fig.

S1B). Each task comprised a five min-long baseline period involving empty chambers located at

opposite corners of the arena, followed by a five min-long encounter period, when a distinct

stimulus was introduced into each chamber [42]. Mice performing the SP task tended to interact

with social stimuli (conspecifics) for significantly more time than with objects throughout the

encounter period (Fig. 1B-D). Similarly, mice performing the EsP task preferred to interact with

socially isolated rather than group-housed stimuli (Fig. 1E-H), while mice performing the SxP task

tended to interact more with female than with male stimuli (Fig 1I-L). Thus, in each task, the

subjects discriminated between a preferred and a less-preferred stimulus. Importantly, the same

type of stimulus (a group-housed male mouse) that was the preferred stimulus in the SP task was

the less-preferred stimulus in the other two tasks. Therefore, this set of tasks allowed us to analyze

brain-wide neural activity patterns in association with either the type of stimulus (i.e., a group-

housed male vs. an object/group-housed female/isolated male) or its valence (i.e., preferred vs.

less-preferred), or the social context (i.e., SP, EsP or SxP task). It should be noted, that in our

hands ICR female mice do not discriminate between group-housed and isolated stimuli, hence we

conducted this study using male subjects only.

We further compared multiple behavioral parameters across the various tasks. There were no

significant differences between the tasks in terms of total time dedicated to stimuli investigation

(Fig. 1M), the total number of transitions made by the subjects between the two stimuli (Fig. 1N)

or the distance traveled by the subjects during a task (Fig. 1O). Nonetheless, the preference

(reflected by the relative discrimination index, RDI) between the two stimuli was lower in the EsP

task, as compared to the SP task (Fig. 1P). Overall, subject behavior was similar across the various

tasks.

Different tasks elicit different profiles of rhythmic LFP signals in multiple brain regions

The power spectral density (PSD) profiles of LFP signals recorded during the encounter period

(Fig. 2A-B), differed among the various tasks performed by the same subject in a brain region-

specific manner (Fig. S1C). For quantitative comparison, we calculated the mean theta (θP) and

gamma (γP) power separately for the baseline and encounter periods of each task for each brain

region. While the mean power during baseline across all regions did not significantly differ

between the tasks (Fig. 2C-D), the change in power during the encounter for both theta ($\Delta\theta$P) and

gamma ($\Delta\gamma$P) rhythms was highest for the SP task, compared to the other two (Fig 2E-F). Thus,

despite the generally similar behavior exhibited by subjects across the tasks (Fig. 1), their system-

level brain LFP signals significantly and consistently differed in power across tasks. Specifically,

despite involving only one social stimulus, the SP task induced the strongest LFP rhythmicity.

When considering each brain region separately, we found that in almost all cases, the mean power

of both rhythms was enhanced during the encounter period, as compared to baseline. Notably, the

mean power change differed significantly among the various tasks and specific regions (Fig. 2G-

H). The similar patterns of $\Delta\theta$P and $\Delta\gamma$P across the various regions suggested the existence of a

link between them. Accordingly, we found statistically significant correlation (Pearson's, r>0.25,

$p<0.05$) between $\Delta\theta$P and $\Delta\gamma$P for the SP and SxP tasks, while borderline significant correlation

(r=0.45, $p=0.059$) was observed for the EsP task (Fig. 2K-M). Thus, when measured over the

course of the entire session, both rhythms seem to be driven by the same process.

To examine the temporal dynamics of LFP rhythmicity during the various tasks, we plotted $\Delta\theta$P

and $\Delta\gamma$P as a function of time for each task and brain region. In accordance with our previous study

in rats [32], we found that both $\Delta\theta$P and $\Delta\gamma$P began to rise several seconds before stimulus

introduction, peaked within 20 s from this point and gradually declined in all brain regions and

tasks (Fig. S2). Thus, the dynamics of LFP rhythmicity across the session were similar among the

various tasks and did not seem to reflect the behavioral dynamics (shown in Fig. 1D, H, L). We

also found no significant correlation (Pearson's, $p>0.05$) between the mean power change and

speed of the subject during any task for either $\Delta\theta$P or $\Delta\gamma$P (Fig. S3A-F).

Overall, these results suggest that theta and gamma rhythmicity during the encounter period are

driven by an internal brain state that shows similar temporal dynamics across tasks, independent

of the behavioral dynamics.

LFP power changes during stimulus investigation are differentially modulated across brain

regions and tasks

Despite the uniform dynamics of LFP rhythmicity in the social brain during the encounter period,

it may be differentially modulated during specific behavioral events, such as stimulus

investigation. We thus examined the possibility that during investigation bouts, $\Delta\theta$P and $\Delta\gamma$P

(henceforth termed $^{\Delta}\theta$P and $^{\Delta}\gamma$P) differ between the various stimuli and tasks. As exemplified by

signals recorded from the amygdalo-hippocampal area (AhiAl) shown in Fig. 3A-F, a Z-score

analysis revealed elevation in theta power during investigation bouts towards social but not the

object stimuli in the SP task, during investigation of both stimulus types in the EsP task and during

investigation of female but not male stimuli in the SxP task. Very similar results were obtained for

gamma power (Fig. S4A-F). This analysis thus suggests a bias in the response towards specific

stimuli, in a task-specific manner. Interestingly, even though the same type of stimulus (a group-

housed male) was used in all tasks, this stimulus elicited a clear elevation in LFP rhythmicity only

during the SP and EsP tasks, but not during the SxP task. These results suggest that at least for the

AhiAl, the change in LFP power was not dictated by either the stimulus type nor by its valence.

To explore the stimulus-specific bias in LFP power change during each task, we calculated the

difference in $^\Delta\theta$P and $^\Delta\gamma$P between the two stimuli, separately for each brain region. Since a

possible bias of LFP rhythmicity of a given brain region may be associated with a behavioral bias

towards a specific stimulus, we examined the correlation between the two variables. To this end,

we correlated the $^\Delta\theta$P bias (preferred minus less-preferred) to the RDI values of each task. We

found a negative correlation in a specific set of brain regions (i.e, extended amygdala (EA) and

lateral septum (LS),) only for the SP task. In contrast, a positive correlation was found in a distinct

set of brain regions (nucleus accumbens shell (AcbSh), AhiAl, ventral pallidum (VP) and

dorsomedial hypothalamic nucleus (DMD)) during the EsP and SxP tasks. Specifically, the VP

exhibited a very strong and highly significant linear correlation with RDI values during both the

EsP and SxP tasks (Fig 3G). These results suggest a link between stimulus-specific bias in $^\Delta\theta$P and

behavioral preference in a task-specific manner.

To further explore this link, we plotted $^\Delta\theta$P bias across all tasks on a 3D plot, separately for each

brain region. We found that almost no region showed bias towards the object stimulus in the SP

task, with the various regions being equally distributed between the two stimuli in the EsP and SxP

tasks (Fig. 4B). In contrast, when $^\Delta\gamma$P was analyzed (Fig. 4C), we observed an opposite picture.

Here, almost all brain regions exhibited stronger responses to the grouped and male stimuli in the

EsP and SxP tasks, yet were rather equally distributed between the two stimuli in the SP task.

Thus, for $^\Delta\gamma$P, most brain regions (14/18) were equally divided between those biased towards less-

preferred stimuli (i.e., object+grouped+male) and those biased towards the type of stimulus used

in all three tasks (i.e, social+grouped+male). The probability of such an arrangement to occur by

chance is smaller than 0.001 (1-binomial test) for each of these two groups. The results thus suggest

that a bias in gamma power is mostly associated with the characteristics (i.e, valence or type) of

the stimulus. They also generally demonstrate opposite stimulus-dependent bias patterns between the theta and gamma rhythms during stimulus investigation, in contrast to their significant correlation when measured during the entire encounter period (Fig. 2E-G). This implies the existence of an independent active state in the social brain during stimulus investigation. Notably, of all brain regions considered, the vDG stood out as the only region biased to the combination of object/isolated/female stimuli. Moreover, this region showed an strong bias for all these stimuli in both $^{\Delta}\theta P$ and $^{\Delta}\gamma P$ (Fig. 3H-I). These results suggest a unique position for the vDG in the social brain, as also supported by results presented below.

Social encounters modulate coherence between brain regions in a social context-dependent manner

Synchronous activity (coherence) enhances effective communication between neuronal groups in different brain regions and dynamically binds them into functional networks [38]. We, therefore, examined the coherence of LFP rhythmicity between each pair of brain regions recorded by us, in both the theta and gamma bands. During the baseline period, the mean theta coherence (θCo) between all pairs of brain regions (99 pairs with ≥ 5 sessions from at least two subjects in all three tasks, see Table S2)) was similar across all tasks (Fig. 4A). Thus, the subjects displayed similar global brain synchronization while exploring the arena without stimuli, in all tasks. However, the change in theta coherence ($\Delta\theta Co$) during the encounter period differed significantly between tasks. While almost all pairs of brain regions exhibited increased θCo during the SP task, we observed significantly milder increases during EsP and SxP tasks, with many paired regions showing reduced θCo (Fig. 4B, E-G). Similar relationships among tasks were observed for changes in gamma coherence (γCo), although here the general tendency was one of decreased coherence during the encounter period (Fig. 4C-G). Notably, there was almost no correlation between the baseline period coherence and the change in coherence during the encounter period for any task (apart a weak correlation for SP gamma coherence; r=-0.21, p=0.0111; not shown). This suggests that the encounter-induced coherence change represents an internal state, independent of the resting state. Finally, when calculating the correlations in $\Delta\theta Co$ across all paired regions between the various tasks, we found a statistically significant high correlation between SxP and EsP, while no correlation was found between SP and SxP. A milder but significant correlation was found between SP and EsP (Fig. 4H). In contrast, all correlations were found significant for $\Delta\gamma Co$. These

results suggest a gradual shift from SP to EsP and to SxP when in $\Delta\theta$Co brain pattern is measured

across the whole encounter.

We, therefore, examined the encounter-induced coherence changes for each brain region

separately, by comparing the change in coherence between a given region and all other regions

across tasks in the theta (Fig. S5A) and gamma (Fig. S5B) bands. We found that differences

between tasks were brain-region specific, with most (13/18) regions showing significant

differences (after FDR corrections) between at least two tasks in $\Delta\theta$Co and one third (6/18)

showing differences in $\Delta\gamma$Co. Notably, in all cases, we found significantly higher coherence

changes during the SP task, as compared to at least one task, and in many cases, to both other tasks.

Thus, a subset of the recorded brain regions displayed differential changes in theta or gamma

coherence among the social contexts, with this change being majorly increased in the SP task, as

compared to the other two tasks.

Brain-wide coherence changes during investigation bouts reflect the social context

To explore possible modulation of LFP coherence during investigation bouts, we calculated the

mean $^\Delta\theta$Co between each pair of brain regions during all investigation bouts towards a given

stimulus, similarly to how we analyzed the power changes (Fig. 3). Since data had been collected

for a large number of brain-region pairs (99 pairs), we focused on pairs showing a mean coherence

change that crossed a cutoff value ± 1.5*standard deviation (SD) for each stimulus (about 20% of

the pairs). When plotting the bias in $^\Delta\theta$Co and $^\Delta\gamma$Co between the two stimuli in each task on a 3D

plot (Fig. S6A-B), we observed a much wider distribution than was found for power (Fig. 3H-I),

suggesting distinct principles of distribution. To further explore this possibility, we examined all

pairs of brain region that passed the aforementioned threshold, separately for each stimulus (Fig.

5A). Surprisingly, the three stimuli of the same type (i.e, social, grouped, male) did not share even

a single pair of brain regions that passed the $^\Delta\theta$Co cutoff value. Similarly, the preferred stimuli

(i.e, social, isolated, female) also did not share even a single pair among them. In contrast, multiple

pairs of brain regions shared similar changes in theta coherence between both stimuli used in each

task (Fig. 5A). For example, CeA-PrL and MeAD-VP showed increased coherence for both social

and object stimuli, BLA-LS and EA-Acbc showed increased coherence for both isolated and

grouped stimuli and Pir-AcbC and EA-AcbC exhibited the strongest increase in theta coherence

for both male and female stimuli. Similar results were observed for $^\Delta\gamma$Co (Fig. S6C). Thus, changes

in coherence during stimulus investigation seem to be dictated by the social context rather than by

stimulus characteristics, such as its type or valence.

For quantitative examination of this possibility, we calculated the correlation across all brain

regions for either $^{\Delta}\theta$Co (Fig. 5B) and $^{\Delta}\gamma$Co (Fig. 5C), between pairs of stimuli which share common

context, type or valence. We found strong and highly significant correlations between all pairs of

stimuli used in the same task (sharing context). In contrast, among the three stimuli of the same

type, only grouped (ESP) and male (SxP) showed significant correlation. Similarly, among

preferred stimuli, only isolated (EsP) and female (SxP) showed significant correlations. Notably,

in both of these cases the correlation was weaker than the correlation between any pair of stimuli

sharing the same context (Fig. 5B). Similar results were found for $^{\Delta}\gamma$Co (Fig. 5C). Thus, coherence

changes during stimulus investigation in both bands had the strongest association with the context

of the social interaction, relative to any characteristic of the stimulus.

Finally, we employed a Decision trees (multi-class Random forest) classifier to examine if $^{\Delta}\theta$Co

and $^{\Delta}\gamma$Co contain information which may be used to discriminate between the various contexts or

stimuli. First, we validated that the model achieved good (~60%) and significant accuracy in

predicting the social stimulus vs. object in the SP task using either $^{\Delta}\theta$Co or $^{\Delta}\gamma$Co. Notably, the

classification of the object vs. social was not accurate, suggesting that the presence of the social

stimulus (social context) mask the object classification (Fig. S7A-B). Then, we used the same

model for predicting the social context (SP, EsP and SxP) and found that using $^{\Delta}\theta$Co (Fig. 5D),

but not $^{\Delta}\gamma$Co (Fig. 5E), allowed the model to predict the right context better than any other context,

and that this prediction was the only one achieving more than a chance level (33.3%) accuracy

(although only SxP classification was statistically significant). In contrast, the same model worked

poorly for predicting stimulus identity among all six stimuli (Fig. S7C-D). Using the LFP theta

power ($^{\Delta}\theta$P) for predicting the social context by the same model achieved good and significant

classification only for the SP (Fig. S7E), while using both theta power and coherence allowed

accurate prediction of both SP and SxP contexts (Fig. S7F). These results suggest that LFP

rhythmicity in the theta range, especially the coherence between the various brain regions, is

informative regarding the social context of the animal more than regarding the identity or valence

of the social stimuli.

Analysis of Granger causality suggests that specific brain regions serve as hubs

The coherent LFP rhythmicity in the social brain can be dominated by specific regions serving as

hubs, thereby preceding other regions in terms of rhythmic neural activity. To identify hub

candidate regions, we first selected brain regions which are statistically over-represented (see

Methods) among pairs of regions exhibiting strong (mean $\pm$ 1.5*SD) bias in any task, separately

for $^\Delta\theta$Co (Fig. 6A) and $^\Delta\gamma$Co (Fig. 6B). We then examined the dependence of LFP rhythmicity of

each of these regions in terms of preceding rhythmicity of other regions, by calculating the change

in Granger causality (GC) [43] during the encounter period (Fig. 6C-E), as compared to the

baseline. We found distinct patterns of statistically significant changes in GC (encounter vs.

baseline periods) between the various tasks (Fig.6-H). Some regions, however, presented

significant GC changes in all tasks, suggesting that they might function as hubs. For example, the

vDG and AcbC participated in significant GC changes in both theta and gamma rhythms in all

tasks. At the same time, PrL and AhiAl were explicitly involved in theta GC changes in all tasks.

Interestingly, theta GC changes from vDG to AhiAl increased during a SP task but decreased

during a EsP task, while theta GC changes from PrL to vDG decreased during both EsP and SxP

tasks. These results suggest that these brain regions dictate LFP rhythmicity in the social brain

during social investigation.

To further explore this possibility, we have calculated the difference in GC change during

encounter between the two directions (from area 1 to area 2 and vice versa), for all couples of brain

regions across all tasks and rhythms (Fig. S8A-C). After applying FDR correction for multiple

comparisons, we found only vDG to LS, for gamma rhythmicity of the EsP task, which was

significantly higher in the vDG-LS direction than in the opposite direction (Fig. S8D).

Context-specific synchronization of LFP rhythmicity in the ventral dentate gyrus with precise

behavioral events

To further examine the candidate hub regions, we exploited our ability to determine the exact

timing of each investigation bout to quantify the synchronized modulation of LFP rhythmicity,

relative to these events. Thus, we compared the modulation of theta and gamma power in all

regions associated with significant GC changes (Fig. 6) relative to a defined battery of specific

behavioral events (Fig. 7A). These events included the beginning and end of investigation bouts

towards specific stimuli, as well as transitions between stimuli. We found a main effect in ANOVA

for multiple events, although in most cases, none of the regions showed significance *in post-hoc*

analysis (see Table S3). One region, the vDG, did, however, exhibit significant differences

between stimuli. The vDG displayed significantly higher theta and gamma powers at the end of

investigation bouts of social stimuli, as compared to object stimuli, specifically in the SP task (Fig.

7B-G and S7A-B). The same region also exhibited decreased theta and gamma powers at the

beginning of transitions from isolated to grouped stimuli, as compared to non-transitional bouts,

specifically in the EsP task (Fig. 7H-M and S7C-D).

These results, together with those shown in Fig. 3B-C and Fig. 6, suggest that the vDG may

function as a hub in the social brain network by coordinating rhythmic neural activity of the

network in a social context-dependent manner.

Discussion

In this study, we used multi-site electrophysiological recordings from the murine social brain to

seek system-level neural correlates of three distinct aspects of social interaction, namely, the type

of the social stimulus, its relative valence (preference) and the social context. To distinguish

between these three aspects, we relied on three social discrimination tasks (i.e., SP, EsP, and SxP)

in which male mice clearly prefer one of two distinct stimuli. This design enabled us to employ

the same type of social stimulus, a novel group-housed male mouse, in all three tasks, with this

stimulus being the preferred stimulus in the SP task and the less-preferred stimulus in the other

two tasks. Importantly, all three tasks took place in the same experimental arena, which enables

uniform interactions between the subject and the stimuli, i.e., stimulus investigation by the subject

[42]. Consequently, as much as we could measure, subject behavior was almost identical in all

three tasks. Therefore, behavioral differences cannot explain the significant differences in the

patterns of rhythmic LFP signals observed among the different tasks.

We analyzed LFP signals at three different time resolutions, specifically, across the whole session,

during stimulus investigation, and during specific behavioral events, such as at the beginning and

end of investigation bouts. When analyzing the power of both theta and gamma rhythms over an

entire session, some aspects seemed to be dictated by a general internal state. In accordance with

previous studies by us and others [31, 32], virtually all brain regions exhibited higher level of theta

and gamma power during the encounter period, as compared to the baseline period. Our

observation that the level of enhanced power was both brain region- and task-specific strongly

suggests that the power elevation was not caused by enhanced electrical noise or any other artifact

but rather by a genuine internal state of the animal. The uniform dynamics of both theta and gamma

power changes across all brain regions and tasks during the encounter period further supports the

existence of a general internal state which is independent of behavioral dynamics or context.

Notably, we observed significant correlations across brain regions between theta and gamma

power changes in all tasks, suggesting that both rhythms are similarly influenced by the internal

state. In agreement with our previous studies in both rats and mice [30, 32], theta and gamma

power maintained their high levels for a time, even after removal of the stimuli from the arena (not

shown), further supporting an encounter-induced general internal state, which slowly fades away.

This state did not seem to be caused by subject movement, as we found no correlation between

subject speed and changes in theta or gamma power for any brain region.

While the dynamics of the internal state seemed to be similar across the distinct contexts, other

aspects of the general (session-wise) changes in theta and gamma power exhibited context-specific

characteristics. For example, the general changes in both power and coherence were highest in the

SP task, suggesting a higher level of the internal state. Assuming that the general state reflects

social motivation, these results are somewhat surprising, given how the SP task involved only one

social stimulus and reasoning that among the various stimuli tested, the female would be the most

attractive to the male subjects. Our interpretation is that the SP task is simpler in terms of social

motivation, as it requires the animal to choose between an inanimate object and a conspecific,

while the other two tasks involve two social stimuli, thereby presenting the subject with a more

challenging dilemma. The higher confidence of the subject during the SP task is in accord with the

simpler pattern of theta coherence changes observed during this task (seen as a general increase

across almost all brain region pairs). Overall, these results suggest that the internal state level may

distinguish between some contexts, which is in accordance with the ability of the Random forest

model to predict only the SP context based on the arousal-induced LFP power. Nevertheless, the

changes in theta and gamma power across the encounter period did not differ between the EsP and

SxP tasks, and thus cannot be the sole basis for the context-specific responses to social cues.

Notably, a recent paper [22] that employed similar recordings during the SP test, used the power,

coherence and GC data (termed *Electome network*) from various regions of the social brain in a

machine-learning model to discriminate between social and object investigation. In accordance

with our results, this study reported that the model's precision was higher for the social than for

the object, thus suggesting that the social stimulus masks the object, which may be attributed to
the context effect.

Analysis of the power change, specifically during stimulus investigation, yielded a different
picture than did session-wide analysis. First, we found no correlation between theta and gamma
power changes during these periods, suggesting a distinct state of active sensing which
characterizes stimulus investigation. Moreover, although both theta and gamma power changes
across brain regions showed bias to specific combinations of stimuli, they did so in distinct
manners. While theta power was biased towards the preferred stimulus in the SP task, with almost
no region (other than hippocampal areas) showing a higher level during investigation of object
stimuli, the gamma power was clearly biased towards the less-preferred stimuli in the EsP and SxP
tasks (grouped and male stimuli), while showing a mixed preference between stimuli in the SP
task. Thus, as related to gamma power, the social brain may be divided between regions associated
with the valence of stimulus (biased towards less-preferred stimuli) and brain regions associated
with the type of stimulus. It should be noted that theta rhythmicity is thought to reflect top-down
processes, such as arousal and attention, which are regulated by brain wide-active neuromodulators
and recruit distributed brain networks [48-52]. In contrast, gamma rhythmicity is considered a
bottom-up process [53, 54], associated with the synchrony of local inhibitory networks [53-56].
This distinction may explain why theta and gamma rhythms reflect stimulus characteristics in an
opposing manner.

In accordance with our hypothesis, that coherent theta and gamma rhythms couple various regions
of the social brain in a social context-dependent manner, we found that the correlation in the
coherence change during stimulus investigation was strongest between the two stimuli in each
task, even the EsP and SxP tasks. In contrast, there were weaker correlations, if any, among the
three stimuli of the same type (social, grouped, male) or the preferred stimuli (social, isolated,
female). The fact that the same correlation pattern was observed for the coherence of both theta
and gamma rhythms supports the validity and significance of the observation. Moreover, using a
Decision Tree classifier, we demonstrated that the theta coherence between the recorded areas
could generate predictions regarding the social context, but not the specific stimulus, which are
accurate above the chance level. The limited accuracy of the model may be attributed to the
restricted number of recorded regions. Thus, we expect that a more comprehensive analysis of the
coherence within the social brain will be able to generate highly accurate prediction of the social

context. Moreover, GC analysis, representing causal time relationships between various brain
regions, also suggests distinct patterns of changes across the various contexts. Altogether, these
results are in accordance with the idea that the social brain processes information during stimulus
investigation in a context-dependent manner dictated by the context-dependent pattern of
coherence within the network. Such a mechanism may explain how the same stimulus induces
distinct patterns of brain activity in different social contexts, which then elicits distinct behavioral
responses to a stimulus.

Finally, the coherence changes and GC analyses led us to identify a small subset of brain regions
that seem highly influential within the network during the various tasks. Of these, the vDG and
AcbC were involved in significant GC changes during all tasks in both the theta and gamma bands,
and thus may serve as hubs that influence the activities of other regions. Analysis of LFP power
in relation to a battery of specific behavioral events demonstrated that while the small group of
brain regions considered showed differential responses as a whole, the vDG was the only region
that alone showed statistically significant responses. Together with its strong bias towards specific
stimuli, as demonstrated for both theta and gamma power during investigation bouts (Fig. 4B-C),
these results suggest a role for the vDG in orchestrating neural activity across the social brain
during social behavior. This conclusion agrees with previous studies reporting a central role of the
dentate gyrus in social behavior [57-59], and specifically in social discrimination [60, 61]. Notably,
multiple studies have implicated the DG in coding contextual changes. For example, DG neurons
were shown to rapidly detect and encode contextual changes [62], while knocking out NMDA
receptors specifically in DG granule cells abolished the ability of mice to distinguish between two
similar contexts [40]. Moreover, hypothalamic supramammillary neurons projecting to the DG
were shown to be activated by contextual novelty [39], while the activity of ventral hippocampal
neurons was shown to process information in a social context-sensitive manner [63]. These studies
are thus in line with our findings regarding the involvement of the vDG in context-dependent social
behavior.

In conclusion, our results suggest that the distribution of LFP rhythmic activity in the social brain
and, most specifically, the synchronization between the various regions is context-specific and
may thus mediate context-specific processing of social information, leading to social context-
dependent social responses and behavior.

Methods

Animals

Adult male and female CD1 mice (12-14 weeks old) were acquired from Envigo (Rehovot, Israel). All mice were housed in groups of 3-5 in a dark/light 12-hour cycle (lights on at 7 pm), with *ad libitum* food and water. Following surgery, implanted mice were housed in isolation so as to not disturb the implanted EAr. Experiments were performed in the dark phase of the dark/light cycle in a sound- and electromagnetic noise-attenuated chamber. All experiments were approved by the Institutional Animal Care and Use Committee of the University of Haifa (Ethical approval #616/19).

Surgery

Mice were anesthetized using isoflurane (induction 3%, 0.5%-0.8% maintenance in 200mL/min of air; SomnoSuite) and placed over a custom-made heating pad (37°C) in a standard stereotaxic device (Kopf Instruments, Tujunga, CA). Two burr holes were drilled for placing the ground and reference wires (silver wire, 127 µm, 300-500 Ω; A-M Systems, Carlsborg, WA). Two watch screws (0-80, 1/16", M1.4) were inserted into the temporal bone. The coordinates for Prl (AP= 2mm, ML= -0.3, DV= -1.8), AcbC (AP= 1, ML= -2.3, DV= -4.7), Pir (AP= -2, ML= -3.3, DV = -5) and CA1 (AP= -3, ML= -3.3, DV = -4.7) were indicated over the left hemisphere using a marker. The skull covering these marked coordinates was removed using a dental drill, and the exposed brain was kept moist with cold, sterile saline. We custom-designed the EA [41] from 16 individual 50 µm formvar-insulated tungsten wires (50-150 kΩ, #CFW2032882; California Wire Company). Before implantation, the EAr was dipped in DiI (1,1'-Dioctadecyl-3,3,3',3'-tetramethylindocarbocyanine perchlorate; 42364, Sigma-Aldrich) to visualize electrode locations *post-mortem*. The reference and ground wires were inserted into their respective burr holes. The EAr was lowered onto the surface of the exposed brain using a motorized manipulator (MP200; Sutter instruments). The dorsoventral coordinates were marked using the depth of the electrode targeting the PVN (AP= -1 mm, ML= -0.3), which was lowered slowly to -4.7 mm. The EA and exposed skull with the screws were secured with dental cement (Enamel plus, Micerium). Mice were sub-cutaneously injected with Baytril (5mg/kg; Bayer) and Norocarp (5 mg/kg; Carprofen, Norbrook Lab) post-surgery and allowed to recover for three days.

Electrophysiological and video recordings

Following brief exposure to isoflurane, subjects were attached to the headstage (RHD 32 ch, #C3314, Intan Technologies) through a custom-made Omnetics to Mill-Max adaptor (Mill-Max model 852-10-100-10-001000). Behavior was recorded using a monochromatic camera (30 Hz, Flea3 USB3, Flir) placed above the arena. Electrophysiological recordings were made with the RHD2000 evaluation system using an ultra-thin SPI interface cable connected to the headstage board (RHD 16ch, #C3334, Intan Technologies). Electrophysiological recordings (sampled at 20 kHz) were synchronized with recorded video using a TTL trigger pulse and by recording camera frame strobes.

Experiment design

We recorded the behavior and neural activity of 14 males in the SP task, 13 males in the EsP task, and 11 males in the SxP task (Table S1), while targeting 18 distinct brain regions. All the stimuli used for the tasks were unfamiliar to the subject mice. In experiments, the mice were briefly exposed to isoflurane, and the EAr was connected to the evaluation system. After 10 minutes of habituation, the recordings started in the arena (30 x 22 x 35 cm) with empty triangular chambers (12 cm isosceles, 35 cm height), as previously described [42]. The triangular chambers had one face ending with metal mesh (18 mm x 6 cm; 1 cm x 1 cm holes) through which the mice interacted with the stimuli. The test was divided into two 5 min periods, a baseline period (pre-encounter) and a period of encounter with the stimuli. The stimuli for the SP task were a novel group-housed male mouse (social) and a Lego toy (object). For the ESP task, isolated (7-14 days) male and group-housed male mice served as stimuli. Finally, for the SxP task, group-housed male and female mice were used as stimuli. Each subject was evaluated for three sessions of each task. The subjects first performed SP and free interactions, with 10 min between these tasks for three sessions, and then EsP and SxP tasks were performed likewise. Each day four sessions were recorded, two in the morning and two in the afternoon, six hours apart (See Fig. S1B). The free interaction data were not used in this study. We excluded sessions from further evaluations when there was a removal of the headstage from the EA by subjects or in a case of a missing video recording from a session . This accounts for the unequal number of sessions and subjects across tasks.

Histology

Subjects were transcardially perfused, and their brains was kept cold in 4% paraformaldehyde for 48 h. Brains were sectioned (50 µm) horizontally (VT 1200s, Leica). Electrode marks were visualized (DiI coated, Red) against DAPI-stained sections with an epifluorescence microscope (Ti2 eclipse, Nikon). The marks were used to locate the respective brain regions, based on the mouse atlas. Out of all implanted electrodes (256), 9% (23 electrodes from 14 mice) were found to be mistargeted and 36% (93) were non-functional (Table S1).

Behavioral analysis

Subject behavior was tracked using the TrackRodent algorithm [42] for tethered mice. Further parameters of behavior, like duration of interaction, interaction bouts, distance traveled by the subjects, subject speed, transitions between stimuli, and RDI values were calculated from the tracked results with custom codes written in MATLAB 2020a.

Electrophysiological data analysis

Only brain regions recorded for more than 5 sessions across at least 3 mice were analyzed. All signals were analyzed with codes custom-written in MATLAB 2020a. We excluded the signals recorded during 30 seconds around stimulus removal and insertion times, to avoid any effect of this action. First, the signals were down-sampled to 5 kHz and low-pass filtered to 300 Hz using a Butterworth filter. The power and time for the different frequencies were estimated using the 'spectrogram' function in MATLAB with the following parameters: Discrete prolate spheroidal sequences (dpss) window = 2 s; overlap = 50% ; increments = 0.5 Hz; and time bins = 0.5 s. The power of each frequency band (theta: 4-12 Hz and gamma: 30-80 Hz) was averaged for both the baseline and encounter periods (5-min long each). Changes in theta ($\Delta\theta P$) and gamma ($\Delta\gamma P$) powers for each brain region were defined as the mean difference in power between the encounter and baseline periods (Fig. 2C-D, H-I). For Z-score analysis of $^\Delta\theta P$ and $^\Delta\gamma P$ during investigation bouts for a given stimulus we used the pre-bout 5 s period as baseline, and averaged the Z-score across all bouts with the same stimulus in each session. Notably, throughout the study we have analyzed only investigation bouts that were longer than 2 s, for two reasons: 1) only >2 s bouts showed statistically significant differences between the stimuli in the various tasks and 2) only >2 s bouts allow a reliable calculation of theta coherence. LFP power ($^\Delta\theta P$ and $^\Delta\gamma P$) for specific bouts

with each stimulus was estimated by calculating the difference between the average power per

second during an investigation bout (which was longer than 2 s) during the encounter period and

the average power per second for investigation of both empty chambers in the baseline period of

the same session, followed by averaging these values over all sessions (Fig. 4).

Coherence analysis

We used the 'mscohere' function of MATLAB to estimate coherence values using Welch's

overlapped averaged periodogram method. The magnitude-squared coherence between two

signals, x, and y, was defined as follows:

542
$$Coherence_{xy} = \frac{S_{xy}}{\sqrt{S_{xx} S_{yy}}}$$

where S_{xy} is the cross-power spectral density of x and y, S_{xx} is the power spectral density of x

and S_{yy} is the power spectral density of y. All coherence analysis was quantified between brain

regions pairs involved in at least five sessions of behavior tasks. Coherence for the baseline period

was quantified as the average coherence of all brain region pairs for each context (Fig. 5A and F).

Changes in coherence (ΔθCo and ΔγCo) during the encounter period (Fig. 5B and G) between a

pair of brain regions were calculated as follows:

550
$$Change\ in\ Coherence = \frac{\mu\ (Coherence_{encounter} - Coherence_{baseline})}{\sigma\ (Coherence_{encounter} - Coherence_{baseline})}$$

where, Coherence$_{encounter}$ is the absolute coherence value between a pair of regions within a

frequency   band during whole encounter period. Coherence$_{baseline}$ is the absolute coherence value

between a pair of regions within a frequency band during an entire encounter period. The change

in coherence for specific bouts ($^{\Delta}$θCo and $^{\Delta}$γCo) to each stimulus was estimated by calculating the

difference between the average coherence per second during an investigation bout (≥2 s) in the

encounter period and the average coherence per second for investigation with both empty

chambers during the baseline period of the same session, followed by averaging these values over

all sessions (Fig. 6A and S5C). Brain regions that displayed higher frequencies of crossing the

threshold of mean ± 1.5 SD $^{\Delta}$θCo and $^{\Delta}$γCo, based on a binomial distribution test, were considered

to be hubs in the coherent social brain (Fig. 7A-B).

Inter-regional pairwise conditional Granger causality

We employed the multi-variate GC toolbox [43] to calculate GC values separately for baseline and

encounter periods between brain regions separately for each task and rhythm. To this end, we

selected brain regions most represented among brain region pairs that crossed the mean ± 1.5*SD

threshold for the difference in coherence change between preferred and less preferred stimuli in

any task, separately for $^{\Delta}\theta$Co and $^{\Delta}\gamma$Co. For GC analysis, LFP signals were measured at a reduced

sampling rate of 500 Hz. We used the "tsdata_to_infocrit" function to determine the model order

of the vector autoregressive (VAR) model. The median model order for all three tasks was 38

(Bayesian information criterion). To further fit the VAR model to our multi-session, multivariate

LFP data, the "tsdata_to_var" function of LWR (Levinson-Whittle recursion) in the regression

mode and a median model order of 38 was used separately for the baseline and encounter periods

of each task. Next, we estimated the autocovariance sequence of the fitted VAR model with the

"var_to_autocov" function. To maximize the computational efficiency of the function, an

acmaxlags of 1500 was chosen. This process did not violate the autocovariance VAR model, as

was estimated by the "var_info" function. Finally, we calculated the pairwise conditional

frequency-domain multivariate GC matrix using the "autocov_to_spwcgc" function, and summed

the GC for the relevant frequency band (theta or famma) using the "smvgc_to_mvgc" function.

Neural responses to behavioral events

We divided all investigation bouts into specific behavioral events, such as the start and end of an

investigation bout or transition from one stimulus to the other (Fig. 7A, nine distinct types). We

aligned LFP power and behavior events for each stimulus by calculating mean power 5 s before

and 5 s after the beginning (or end) of all investigation bouts (0.5 s bins) in a session. Furthermore,

for each bout, the mean power was normalized using Z-score analysis, where a pre-bout duration

of 5 s served as baseline (Table S2).

Statistical analysis

Statistical analysis was performed using GraphPad Prism 9.5. To test for normal distribution of

the data, we used the Kolmogorov-Smirnov and Shapiro-Wilk tests. Table S3 summarizes the

specific tests conducted for each figure. A paired t-test or Wilcoxon matched-pairs signed rank

test was used to compare different stimuli or conditions for the same group. An unpaired t-test or

Mann-Whitney test was performed to compare a parameter between distinct groups. For
comparison among multiple groups and parameters, ANOVA (normal distribution), Welch's
ANOVA (assuming unequal variance), and Kruskal-Wallis test (non-normal distribution) were
applied to the data. If a main effect or interaction were found in the tests above, Šídák's test,
Dunnett's T3 test or Dunn's *post-hoc* multiple comparison corrections were applied. Repeated
measures ANOVA or a Friedman test was used to compare multiple groups and parameters with
repeated variables. When main effects were observed in above tests, Šídák's or Dunn's test were
used for multiple comparisons corrections, respectively. Additionally, for comparison of two
factors and the interaction between them, from multiple groups and parameters where one of the
factors has repeated measurements, was performed using two-way ANOVA (no missing variables)
or mixed-models ANOVA (Restricted maximum likelihood model, REML). The ANOVA tests
were followed by Šídák's multiple comparison test if main effects or interactions were found. The
association between two groups or parameters was compared with either Pearson's or Spearman's
tests. A binomial distribution test was performed to compare the probability of brain regions
representing above-chance levels for specific parameters.
608

Decision tree classifier model
Data normalization – subtracting the mean value for each brain regions pair per mouse:
The data from two mice (total 14) were ignored as they had less than 40 recorded brain regions
pairs (out of 99). The mean value of each pair was computed and subtracted for each mouse
separately. This helped to reduce the variability of the measurements across mice and improved
classification accuracy. To reduce over-representation of a single stimulus in the computation of
the mean value for a pair, we first averaged the mean value per stimulus (for a specific mouse) and
then subtracted the average of these means.
Averaging bouts:
The average bout for each stimulus was computed for each session.
Data Imputation:
For each mouse, a slightly different set of brain areas were recorded due to slight inaccuracies in
placing the electrodes and slight difference in the individual mouse anatomy. This resulted in
missing entries from some of the brain regions pairs. We used a data imputation strategy to restore
these missing entries. Note that before this step, we subtracted the mean value per brain region per

mouse and averaged all the bouts from the same stimuli of the same session. The imputation algorithm is based on the MICE algorithm [44]. It is an iterative algorithm. In each iteration, it estimates the missing entries by a linear combination of (some of) the other entries. The used data imputation algorithm was defined as follows:

1. For each missing value of brain pair i in bout b, ($bp_{i,b}$), replace $bp_{i,b}$ with the average value of bp_i across the valid values bp_i (from all bouts of all mice with a valid measurement of bp_i).

2. For each bp_i ($order\ of\ brain\ pairs\ is\ randomized$):

   a. Randomly choose a set of predicting brain pairs $\{bp_j\}$ such that $bp_i \notin \{bp_j\}$ and $|\{bp_j\}| < 0.5*$number_of_equations. Where the number_of_equations equals to the number of (averaged) bouts from all the mice (66 predicting brain pairs as the number of bouts in our dataset is 131 average bouts).

   b. Compute linear regression coefficients {aj} (by least square method) to minimize:

$$\underset{\{a_j\}}{\mathrm{argmin}} \sum_b \left(bp_{i,b} - \left(a_0 + \sum_j a_j bp_{j,b} \right) \right)^2$$

   c. For each bout b in which $bp_{i,b}$ was not measured in-vivo, replace it with its estimation: $a_0 + \sum_j a_j bp_{j,b}$

3. Repeat steps 2 for 20 iterations.

Code was implemented in Matlab 2021a.

Classification and computation of confusion matrixes

We used Matlab's *TreeBagger()* function to train a multi-class Random forest classifier for discriminating between a pair of stimuli (social vs object) or contexts (3 classes) or between stimuli (6 classes) or between. We used 80 random trees (a parameter of the TreeBagger function). We used cross-validation with "one mouse leave out" strategy to compute a confusion matrix for each mouse based on a training set that includes examples from all of the other mice. We balanced the training set to have the same number of examples from each class by randomly removing some of the training examples. Since both the balancing and the Random forest algorithm have a random component, we repeated the estimation of confusion matrixes 100 times (for each mouse) to better

estimate the confusion matrixes. We then summed up all of the confusion matrixes (totally 1200

confusion matrixes: 12 mice and 100 confusion matrixes per mouse) and computed for each pair

of classes (i,j) the percent of cases where the prediction was i for bouts of class j.

Statistical Analysis

All tests were corrected for multiple comparisons using FDR corrections [45]. To compute p-

values, we used the average (over 100 iterations) confusion matrix for each mouse (totally 12

confusion matrixes in which each cell i,j represents the % of predictions of class i for bouts of class

j) and compare those with a set of random confusion matrixes which were generated by the same

procedure except for replacing the trained classifier with a random classifier. This random

classifier generated random labels with uniform distribution. To better describe the distribution of

the random confusion matrixes, we generated 83 random confusion matrixes per mouse (each one

of them is an average of 100 confusion matrixes). Then, the p-value for each cell in the confusion

matrix was computed separately by comparing the 12 values from the confusion matrix of the

trained classifier to the 83*12=996 values from the confusion matrixes of the random classifier

using Mann-Whitney U test. In case a mouse did not have a bout from a specific class, this mouse

was ignored in the computation of the p-value for the cells of this ground truth class.

Author contributions

A.N.M.: Formal analysis, Investigation, Methodology, Validation, Visualization, Writing - original draft, and Writing - review & editing; D.P.: Formal analysis, and software.; S.N.: Data curation, Project administration, Software, Validation, Visualization, Writing - original draft, and Writing - review & editing. S.W.: Conceptualization, Funding acquisition, Project administration, Resources, Supervision, Writing - original draft, and Writing - review & editing

Declaration of interests

The authors declare no competing interests.

Acknowledgments

We thank Boris Shklyar, Head of the Bio-imaging Unit and Dr. Maya Lalzar, Head of the Bioinformatics Unit of the Faculty of Natural Sciences of the University of Haifa for their guidance and technical assistance. We also thank Eng. Alex Bizer, the experimental systems engineer of the Faculty of Natural Sciences of the University of Haifa, for their help.

Funding

This study was supported by ISF-NSFC joint research program (grant No. 3459/20 to SW), the Israel Science Foundation (grants No. 1361/17 and 2220/22 to SW), the Ministry of Science, Technology and Space of Israel (Grant No. 3-12068 to SW) and the United States-Israel Binational Science Foundation (grant No. 2019186 to SW).

References

1. Adolphs R. Conceptual challenges and directions for social neuroscience. Neuron. 2010;65(6):752-67. Epub 2010/03/30. doi: 10.1016/j.neuron.2010.03.006. PubMed PMID: 20346753; PubMed Central PMCID: PMCPMC2887730.

2. Ford CL, Young LJ. Translational opportunities for circuit-based social neuroscience: advancing 21st century psychiatry. Curr Opin Neurobiol. 2021;68:1-8. Epub 2020/12/02. doi: 10.1016/j.conb.2020.11.007. PubMed PMID: 33260106; PubMed Central PMCID: PMCPMC8160019.

3. Kohl J, Dulac C. Neural control of parental behaviors. Curr Opin Neurobiol. 2018;49:116-22. Epub 2018/02/27. doi: 10.1016/j.conb.2018.02.002. PubMed PMID: 29482085; PubMed Central PMCID: PMCPMC6029232.

4. McKinsey G, Ahmed OM, Shah NM. Neural control of sexually dimorphic social behaviors. Curr Opin Physiol. 2018;6:89-95. Epub 2019/09/20. doi: 10.1016/j.cophys.2018.08.003. PubMed PMID: 31535059; PubMed Central PMCID: PMCPMC6750220.

5. Wei D, Talwar V, Lin D. Neural circuits of social behaviors: Innate yet flexible. Neuron. 2021;109(10):1600-20. Epub 2021/03/12. doi: 10.1016/j.neuron.2021.02.012. PubMed PMID: 33705708; PubMed Central PMCID: PMCPMC8141016.

6. Insel TR, Fernald RD. How the brain processes social information: searching for the social brain. Annu Rev Neurosci. 2004;27:697-722. Epub 2004/06/26. doi: 10.1146/annurev.neuro.27.070203.144148. PubMed PMID: 15217348.

7. Dickinson SY, Kelly DA, Padilla SL, Bergan JF. From Reductionism Toward Integration: Understanding How Social Behavior Emerges From Integrated Circuits. Front Integr Neurosci. 2022;16:862437. Epub 2022/04/19. doi: 10.3389/fnint.2022.862437. PubMed PMID: 35431824; PubMed Central PMCID: PMCPMC9010670.

8. Goodson JL. The vertebrate social behavior network: evolutionary themes and variations. Horm Behav. 2005;48(1):11-22. Epub 2005/05/12. doi: 10.1016/j.yhbeh.2005.02.003. PubMed PMID: 15885690; PubMed Central PMCID: PMCPMC2570781.

9. Dai B, Sun F, Tong X, Ding Y, Kuang A, Osakada T, et al. Responses and functions of dopamine in nucleus accumbens core during social behaviors. Cell Rep. 2022;40(8):111246. Epub 2022/08/25. doi: 10.1016/j.celrep.2022.111246. PubMed PMID: 36001967; PubMed Central PMCID: PMCPMC9511885.

721 10. Felix-Ortiz AC, Tye KM. Amygdala inputs to the ventral hippocampus bidirectionally

722   modulate social behavior. J Neurosci. 2014;34(2):586-95. Epub 2014/01/10. doi:

723   10.1523/JNEUROSCI.4257-13.2014. PubMed PMID: 24403157; PubMed Central PMCID:

724   PMCPMC3870937.

725 11. Kingsbury L, Huang S, Raam T, Ye LS, Wei D, Hu RK, et al. Cortical Representations of

726   Conspecific Sex Shape Social Behavior. Neuron. 2020;107(5):941-53 e7. Epub 2020/07/15.

727   doi: 10.1016/j.neuron.2020.06.020. PubMed PMID: 32663438; PubMed Central PMCID:

728   PMCPMC7486272.

729 12. Leroy F, Park J, Asok A, Brann DH, Meira T, Boyle LM, et al. A circuit from

730   hippocampal CA2 to lateral septum disinhibits social aggression. Nature.

731   2018;564(7735):213-8. Epub 2018/12/07. doi: 10.1038/s41586-018-0772-0. PubMed PMID:

732   30518859; PubMed Central PMCID: PMCPMC6364572.

733 13. Kohl J, Babayan BM, Rubinstein ND, Autry AE, Marin-Rodriguez B, Kapoor V, et al.

734   Functional circuit architecture underlying parental behaviour. Nature. 2018;556(7701):326-

735   31. Epub 2018/04/13. doi: 10.1038/s41586-018-0027-0. PubMed PMID: 29643503; PubMed

736   Central PMCID: PMCPMC5908752.

737 14. Scheggia D, La Greca F, Maltese F, Chiacchierini G, Italia M, Molent C, et al. Reciprocal

738   cortico-amygdala connections regulate prosocial and selfish choices in mice. Nat Neurosci.

739   2022;25(11):1505-18. Epub 2022/10/26. doi: 10.1038/s41593-022-01179-2. PubMed PMID:

740   36280797; PubMed Central PMCID: PMCPMC7613781.

741 15. Swanson LW, Hahn JD, Sporns O. Structure-function subsystem models of female and

742   male forebrain networks integrating cognition, affect, behavior, and bodily functions. Proc

743   Natl Acad Sci U S A. 2020;117(49):31470-81. Epub 2020/11/25. doi:

744   10.1073/pnas.2017733117. PubMed PMID: 33229546; PubMed Central PMCID:

745   PMCPMC7733829.

746 16. Twining RC, Vantrease JE, Love S, Padival M, Rosenkranz JA. An intra-amygdala

747   circuit specifically regulates social fear learning. Nat Neurosci. 2017;20(3):459-69. Epub

748   2017/01/24. doi: 10.1038/nn.4481. PubMed PMID: 28114293; PubMed Central PMCID:

749   PMCPMC5323274.

750 17. Yamamoto R, Ahmed N, Ito T, Gungor NZ, Pare D. Optogenetic Study of Anterior

751   BNST and Basomedial Amygdala Projections to the Ventromedial Hypothalamus. Eneuro.

752   2018;5(3). Epub 2018/07/05. doi: 10.1523/ENEURO.0204-18.2018. PubMed PMID:

753   29971248; PubMed Central PMCID: PMCPMC6027956.

18.   Dolen G, Darvishzadeh A, Huang KW, Malenka RC. Social reward requires coordinated

755   activity of nucleus accumbens oxytocin and serotonin. Nature. 2013;501(7466):179-84. doi:

756   10.1038/nature12518. PubMed PMID: 24025838; PubMed Central PMCID:

757   PMCPMC4091761.

19.   Felix-Ortiz AC, Burgos-Robles A, Bhagat ND, Leppla CA, Tye KM. Bidirectional

759   modulation of anxiety-related and social behaviors by amygdala projections to the medial

760   prefrontal cortex. Neuroscience. 2016;321:197-209. Epub 2015/07/25. doi:

761   10.1016/j.neuroscience.2015.07.041. PubMed PMID: 26204817; PubMed Central PMCID:

762   PMCPMC4721937.

20.   Huang WC, Zucca A, Levy J, Page DT. Social Behavior Is Modulated by Valence-

764   Encoding mPFC-Amygdala Sub-circuitry. Cell Rep. 2020;32(2):107899. Epub 2020/07/16.

765   doi: 10.1016/j.celrep.2020.107899. PubMed PMID: 32668253; PubMed Central PMCID:

766   PMCPMC7410267.

21.   Reis FM, Lee JY, Maesta-Pereira S, Schuette PJ, Chakerian M, Liu J, et al. Dorsal

768   periaqueductal gray ensembles represent approach and avoidance states. Elife. 2021;10. Epub

769   2021/05/07. doi: 10.7554/eLife.64934. PubMed PMID: 33955356; PubMed Central PMCID:

770   PMCPMC8133778.

22.   Mague SD, Talbot A, Blount C, Walder-Christensen KK, Duffney LJ, Adamson E, et al.

772   Brain-wide electrical dynamics encode individual appetitive social behavior. Neuron.

773   2022;110(10):1728-41 e7. Epub 2022/03/17. doi: 10.1016/j.neuron.2022.02.016. PubMed

774   PMID: 35294900; PubMed Central PMCID: PMCPMC9126093.

23.   Schaich Borg J, Srivastava S, Lin L, Heffner J, Dunson D, Dzirasa K, et al. Rat

776   intersubjective decisions are encoded by frequency-specific oscillatory contexts. Brain Behav.

777   2017;7(6):e00710. Epub 2017/06/24. doi: 10.1002/brb3.710. PubMed PMID: 28638715;

778   PubMed Central PMCID: PMCPMC5474713.

24.   Buzsaki G, Draguhn A. Neuronal oscillations in cortical networks. Science.

2004;304(5679):1926-9. doi: 10.1126/science.1099745. PubMed PMID: 15218136.

25.   Harris AZ, Gordon JA. Long-range neural synchrony in behavior. Annu Rev Neurosci.

2015;38:171-94. Epub 2015/04/22. doi: 10.1146/annurev-neuro-071714-034111. PubMed

PMID: 25897876; PubMed Central PMCID: PMCPMC4497851.

26.   Uhlhaas PJ, Pipa G, Lima B, Melloni L, Neuenschwander S, Nikolic D, et al. Neural

synchrony in cortical networks: history, concept and current status. Front Integr Neurosci.

2009;3:17. Epub 2009/08/12. doi: 10.3389/neuro.07.017.2009. PubMed PMID: 19668703;

PubMed Central PMCID: PMCPMC2723047.

27.   Bocchio M, Nabavi S, Capogna M. Synaptic Plasticity, Engrams, and Network

Oscillations in Amygdala Circuits for Storage and Retrieval of Emotional Memories. Neuron.

2017;94(4):731-43. Epub 2017/05/19. doi: 10.1016/j.neuron.2017.03.022. PubMed PMID:

28521127.

28.   Chen S, Tan Z, Xia W, Gomes CA, Zhang X, Zhou W, et al. Theta oscillations

synchronize human medial prefrontal cortex and amygdala during fear learning. Sci Adv.

2021;7(34). Epub 2021/08/20. doi: 10.1126/sciadv.abf4198. PubMed PMID: 34407939;

PubMed Central PMCID: PMCPMC8373137.

29.   Taub AH, Perets R, Kahana E, Paz R. Oscillations Synchronize Amygdala-to-Prefrontal

Primate Circuits during Aversive Learning. Neuron. 2018;97(2):291-8 e3. Epub 2018/01/02.

doi: 10.1016/j.neuron.2017.11.042. PubMed PMID: 29290553.

30.   John SR, Dagash W, Mohapatra AN, Netser S, Wagner S. Distinct Dynamics of Theta

and Gamma Rhythmicity during Social Interaction Suggest Differential Mode of Action in the

Medial Amygdala of Sprague Dawley Rats and C57BL/6J Mice. Neuroscience. 2022;493:69-

80. Epub 2022/05/02. doi: 10.1016/j.neuroscience.2022.04.020. PubMed PMID: 35490969.

31.   Kuga N, Abe R, Takano K, Ikegaya Y, Sasaki T. Prefrontal-amygdalar oscillations

related to social behavior in mice. Elife. 2022;11. Epub 2022/05/18. doi:

10.7554/eLife.78428. PubMed PMID: 35580019; PubMed Central PMCID:

PMCPMC9113747.

32.   Tendler A, Wagner S. Different types of theta rhythmicity are induced by social and

fearful stimuli in a network associated with social memory. Elife. 2015;4. doi:

10.7554/eLife.03614. PubMed PMID: 25686218; PubMed Central PMCID:

PMCPMC4353977.

33.   Buzsaki G, Watson BO. Brain rhythms and neural syntax: implications for efficient
coding of cognitive content and neuropsychiatric disease. Dialogues Clin Neurosci.
2012;14(4):345-67. Epub 2013/02/09. doi: 10.31887/DCNS.2012.14.4/gbuzsaki. PubMed
PMID: 23393413; PubMed Central PMCID: PMCPMC3553572.
34.   Cannon J, McCarthy MM, Lee S, Lee J, Borgers C, Whittington MA, et al.
Neurosystems: brain rhythms and cognitive processing. Eur J Neurosci. 2014;39(5):705-19.
Epub 2013/12/18. doi: 10.1111/ejn.12453. PubMed PMID: 24329933; PubMed Central
PMCID: PMCPMC4916881.
35.   Uhlhaas PJ, Roux F, Rodriguez E, Rotarska-Jagiela A, Singer W. Neural synchrony and
the development of cortical networks. Trends Cogn Sci. 2010;14(2):72-80. doi:
10.1016/j.tics.2009.12.002. PubMed PMID: 20080054.
36.   Geschwind DH, Levitt P. Autism spectrum disorders: developmental disconnection
syndromes. Curr Opin Neurobiol. 2007;17(1):103-11. Epub 2007/02/06. doi:
10.1016/j.conb.2007.01.009. PubMed PMID: 17275283.
37.   Lazaro MT, Taxidis J, Shuman T, Bachmutsky I, Ikrar T, Santos R, et al. Reduced
Prefrontal Synaptic Connectivity and Disturbed Oscillatory Population Dynamics in the
CNTNAP2 Model of Autism. Cell Rep. 2019;27(9):2567-78 e6. Epub 2019/05/30. doi:
10.1016/j.celrep.2019.05.006. PubMed PMID: 31141683; PubMed Central PMCID:
PMCPMC6553483.
38.   Fries P. Rhythms for Cognition: Communication through Coherence. Neuron.
2015;88(1):220-35. Epub 2015/10/09. doi: 10.1016/j.neuron.2015.09.034. PubMed PMID:
26447583; PubMed Central PMCID: PMCPMC4605134.
39.   Chen S, He L, Huang AJY, Boehringer R, Robert V, Wintzer ME, et al. A hypothalamic
novelty signal modulates hippocampal memory. Nature. 2020;586(7828):270-4. Epub
2020/10/02. doi: 10.1038/s41586-020-2771-1. PubMed PMID: 32999460.
40.   McHugh TJ, Jones MW, Quinn JJ, Balthasar N, Coppari R, Elmquist JK, et al. Dentate
gyrus NMDA receptors mediate rapid pattern separation in the hippocampal network.
Science. 2007;317(5834):94-9. Epub 2007/06/09. doi: 10.1126/science.1140263. PubMed
PMID: 17556551.

41.   Mohapatra AN, Netser S, Wagner S. Modular Electrode Array for Multi-site

Extracellular Recordings from Brains of Freely Moving Rodents. Curr Protoc.

2022;2(5):e399. Epub 2022/05/11. doi: 10.1002/cpz1.399. PubMed PMID: 35536117.

42.   Netser S, Haskal S, Magalnik H, Bizer A, Wagner S. A System for Tracking the

Dynamics of Social Preference Behavior in Small Rodents. J Vis Exp. 2019;(153). doi:

10.3791/60336. PubMed PMID: 31814614.

43.   Barnett L, Seth AK. The MVGC multivariate Granger causality toolbox: a new approach

to Granger-causal inference. J Neurosci Methods. 2014;223:50-68. Epub 2013/11/10. doi:

10.1016/j.jneumeth.2013.10.018. PubMed PMID: 24200508.

44.   Azur MJ, Stuart EA, Frangakis C, Leaf PJ. Multiple imputation by chained equations:

what is it and how does it work? Int J Methods Psychiatr Res. 2011;20(1):40-9. Epub

2011/04/19. doi: 10.1002/mpr.329. PubMed PMID: 21499542; PubMed Central PMCID:

PMCPMC3074241.

45.   Boca SM, Leek JT. A direct approach to estimating false discovery rates conditional on

covariates. PeerJ. 2018;6:e6035. Epub 2018/12/26. doi: 10.7717/peerj.6035. PubMed PMID:

30581661; PubMed Central PMCID: PMCPMC6292380.

46.   Jabarin R, Levy N, Abergel Y, Berman JH, Zag A, Netser S, et al. Pharmacological

modulation of AMPA receptors rescues specific impairments in social behavior associated

with the A350V Iqsec2 mutation. Transl Psychiatry. 2021;11(1):234. Epub 2021/04/24. doi:

10.1038/s41398-021-01347-1. PubMed PMID: 33888678; PubMed Central PMCID:

PMCPMC8062516.

47.   Netser S, Meyer A, Magalnik H, Zylbertal A, de la Zerda SH, Briller M, et al. Distinct

dynamics of social motivation drive differential social behavior in laboratory rat and mouse

strains. Nat Commun. 2020;11(1):5908. Epub 2020/11/22. doi: 10.1038/s41467-020-19569-0.

PubMed PMID: 33219219; PubMed Central PMCID: PMCPMC7679456.

48.   Clayton MS, Yeung N, Kadosh RC. The roles of cortical oscillations in sustained

attention. Trends in Cognitive Sciences. 2015;19(4):188-95. doi: 10.1016/j.tics.2015.02.004.

PubMed PMID: WOS:000352674600006.

49.   Fiebelkorn IC, Kastner S. A Rhythmic Theory of Attention. Trends in Cognitive

Sciences. 2019;23(2):87-101. doi: 10.1016/j.tics.2018.11.009. PubMed PMID:

WOS:000455717200005.

50.   Helfrich RF, Breska A, Knight RT. Neural entrainment and network resonance in support

of top-down guided attention. Curr Opin Psychol. 2019;29:82-9. Epub 2019/01/29. doi:

10.1016/j.copsyc.2018.12.016. PubMed PMID: 30690228; PubMed Central PMCID:

PMCPMC6606401.

51.   Karakas S. A review of theta oscillation and its functional correlates. Int J Psychophysiol.

2020;157:82-99. Epub 2020/05/20. doi: 10.1016/j.ijpsycho.2020.04.008. PubMed PMID:

32428524.

52.   Knyazev GG. Motivation, emotion, and their inhibitory control mirrored in brain

oscillations. Neurosci Biobehav R. 2007;31(3):377-95. doi: 10.1016/j.neubiorev.2006.10.004.

PubMed PMID: WOS:000246316100006.

53.   Buzsaki G, Wang XJ. Mechanisms of gamma oscillations. Annu Rev Neurosci.

2012;35:203-25. Epub 2012/03/27. doi: 10.1146/annurev-neuro-062111-150444. PubMed

PMID: 22443509; PubMed Central PMCID: PMCPMC4049541.

54.   Headley DB, Pare D. In sync: gamma oscillations and emotional memory. Front Behav

Neurosci. 2013;7:170. Epub 2013/12/10. doi: 10.3389/fnbeh.2013.00170. PubMed PMID:

24319416; PubMed Central PMCID: PMCPMC3836200.

55.   Benchenane K, Tiesinga PH, Battaglia FP. Oscillations in the prefrontal cortex: a

gateway to memory and attention. Curr Opin Neurobiol. 2011;21(3):475-85. Epub

2011/03/25. doi: 10.1016/j.conb.2011.01.004. PubMed PMID: 21429736.

56.   Palva JM, Palva S. Functional integration across oscillation frequencies by cross-

frequency phase synchronization. Eur J Neurosci. 2018;48(7):2399-406. Epub 2017/11/03.

doi: 10.1111/ejn.13767. PubMed PMID: 29094462.

57.   Cai Y, Tang X, Chen X, Li X, Wang Y, Bao X, et al. Liver X receptor beta regulates the

development of the dentate gyrus and autistic-like behavior in the mouse. Proc Natl Acad Sci

U S A. 2018;115(12):E2725-E33. Epub 2018/03/07. doi: 10.1073/pnas.1800184115. PubMed

PMID: 29507213; PubMed Central PMCID: PMCPMC5866608.

58.   Doucette E, Merfeld E, Leblanc H, Monasterio A, Cincotta C, Grella SL, et al. Social

behavior in mice following chronic optogenetic stimulation of hippocampal engrams.

Neurobiol Learn Mem. 2020;176:107321. Epub 2020/11/10. doi: 10.1016/j.nlm.2020.107321.

PubMed PMID: 33164892; PubMed Central PMCID: PMCPMC7708439.

59.   Leung C, Cao F, Nguyen R, Joshi K, Aqrabawi AJ, Xia S, et al. Activation of Entorhinal
Cortical Projections to the Dentate Gyrus Underlies Social Memory Retrieval. Cell Rep.
2018;23(8):2379-91. Epub 2018/05/24. doi: 10.1016/j.celrep.2018.04.073. PubMed PMID:
29791849.
60.   Cope EC, Waters RC, Diethorn EJ, Pagliai KA, Dias CG, Tsuda M, et al. Adult-Born
Neurons in the Hippocampus Are Essential for Social Memory Maintenance. Eneuro.
2020;7(6). Epub 2020/10/17. doi: 10.1523/ENEURO.0182-20.2020. PubMed PMID:
33060182; PubMed Central PMCID: PMCPMC7768285.
61.   Li J, Sun X, You Y, Li Q, Wei C, Zhao L, et al. Auts2 deletion involves in DG
hypoplasia and social recognition deficit: The developmental and neural circuit mechanisms.
Sci Adv. 2022;8(9):eabk1238. Epub 2022/03/03. doi: 10.1126/sciadv.abk1238. PubMed
PMID: 35235353; PubMed Central PMCID: PMCPMC8890717.
62.   Leutgeb JK, Leutgeb S, Moser MB, Moser EI. Pattern separation in the dentate gyrus and
CA3 of the hippocampus. Science. 2007;315(5814):961-6. Epub 2007/02/17. doi:
10.1126/science.1135801. PubMed PMID: 17303747.
63.   Wu WY, Yiu E, Ophir AG, Smith DM. Effects of social context manipulation on dorsal
and ventral hippocampal neuronal responses. Hippocampus. 2023. Epub 2023/02/16. doi:
10.1002/hipo.23507. PubMed PMID: 36789678.
919

Figures

Figure 1. Similar behavior of subject mice across three binary social discrimination tasks.

A. A scheme of the arena during SP task session. The two stimuli types are indicated on the right
side.

B. Heat maps of investigation bouts made by the subjects toward each of the stimuli (stimulus
type is noted above) across the five min-long encounter period of the SP task, with color-
coding of the investigation bout duration (see scale on the right of the panel). Each line
represents a distinct session.

C. Mean (±SEM) time dedicated by a subject for investigating each stimulus during the SP task
sessions shown in B. Wilcoxon matched pairs signed rank test, n = 33 sessions, W = -495,
****$p<0.0001$.

D. As in C, plotted vs. time using one min bins.

E-H. As in A-D, for the EsP task. Paired t-test, n = 28 sessions, t (27) =2.374, *p = 0.025.

I-L.   As in A-D, for the SxP task. Paired t-test, n = 26 sessions, t (25) =5.75, ****$p<0.0001$.

M. Mean (±SEM) total time dedicated by a subject to investigate both stimuli during the encounter

stage of each task.

N. Mean (±SEM) number of transitions between stimuli made by the subject during the encounter

period of each task.

O. Mean (±SEM) distance traveled by the subjects during the encounter stage of each task.

P. Mean (±SEM) RDI for each task. Kruskal-Wallis test, n = 3 tests, 87 sessions, H = 8.509, $p =$

0.0142; Dunn's *post-hoc* test, *$p<0.05$.

Figure 2. Brain region- and context-specific changes in the levels of theta and gamma power during a social encounter.

A. Color-coded spectrogram of LFP signals recorded in the AcbSh during the baseline (left) and encounter (right) periods of SP task conducted by a subject. The black line at 300 s represent the time of stimuli introduction into the arena. The color-coding scale is shown on the right.

B. PSD profiles of the baseline (blue) and encounter (red) periods of the example shown in A. The gray areas mark the theta and gamma ranges. The inset shows the theta range in higher resolution.

C. Mean (±SEM) theta power (θP) during the baseline period for each brain region in the three contexts Kruskal-Wallis test, n = 3 tests, 87 sessions, H = 2.725, P = 0.2561.

D. As in C, for gamma power (γP; Welch's ANOVA, W (DFn, DFd) = 0.9496(2,33.84), P = 0.2561)

E. Mean (±SEM) $\Delta\theta$P, averaged across all brain regions, for each task. W (DFn, DFd) = 14.67 (2, 31.80), p <0.0001. Dunnett's T3 multiple comparisons test, SP vs. EsP, p = 0.0018; SP vs. SxP, p = 0.0002; EsP vs. SxP, P = 0.2653.

F. As in G, for $\Delta\gamma$P. W (DFn, DFd) = 5.134 (2, 33.65), p = 0.0113. SP vs. EsP, p = 0.0531; SP vs. SxP, p = 0.0127; EsP vs. SxP, p = 0.7467.

G. Mean (±SEM) change in theta power ($\Delta\theta$P) during the encounter period, relative to the baseline period for each brain region in the three contexts (2-way ANOVA. Contexts: F (2, 659) = 3.838, p = 0.0220; Brain regions: F (17, 659) = 1.727, p = 0.0341; Interaction: F (34, 659) = 0.4548, p = 0.9970)

H. As in E, for change in gamma power ($\Delta\gamma$P; 2-way ANOVA. Contexts: F (2, 659) = 1.459, P = 0.2333; Brain regions: F (17, 659) = 2.050, P = 0.0076; Interaction: F (34, 659) = 0.5732, P = 0.9764).

I. Super-imposed traces of $\Delta\theta$P averaged across all brain regions for the SP (blue), EsP (green) and SxP (red) tasks. Time 0 min represents the time of stimuli insertion.

J. As in I, for $\Delta\gamma$P.

K. Mean $\Delta\gamma$P as a function of mean $\Delta\theta$P during the SP task, for each brain region. Pearson's correlation coefficient (r) and significance (p) are given.

L. As in K, for the EsP task.

M. As in K, for the SxP task. #p= 0.053, *p<0.05, **p<0.01, ***p<0.001. See also Figs. S1, S2.

Figure 3. Stimulus- and task-specific LFP power changes during investigation bouts.

A. Heat maps of average theta power in AhiAl, before and during social investigation bouts made by AhiAl-Implanted subjects with social (above) and object (below) stimuli during SP task sessions (n=17 sessions). Each row represents the mean Z-score of all bouts in a single session (using 0.5 s bins). Time '0' represents the beginning of the bout. The color code scale is on the right.

B. As in A, for investigation bouts of isolated (above) and grouped (below) social stimuli during EsP task sessions (n=14 sessions).

C. As in A, for investigation bouts of female (above) and male (below) social stimuli during SxP task sessions (n=13 sessions).

D. Mean (±SEM) Z-score trace of the data shown in A for both stimuli.

E. As in D. for the data shown in B.

F. As in D, for the data shown in C.

G. Correlation between mean change in theta power during investigation bouts ($^{\Delta}\theta P$) in specific brain regions and RDI values during the various tasks. Only statistically significant linear correlations are shown. Note that one outlier SP session which made the correlation even stronger was excluded (Fig. S4G)

H. A 3D plot of the mean difference between preferred and less-preferred stimuli in $^{\Delta}\theta P$. Each circle represents a given brain region, color- and shape-coded according to the combined bias across all tasks. See legend of the color and shape code of the distinct combinations below.

I. As in B, for $^{\Delta}\gamma P$.

See also Fig. S4.

Figure 4. Social encounters modulate the coherence between brain regions in a social context-

dependent manner.

A. Mean theta coherence during the baseline period for each task, across all (n=99) pairs of brain

regions recorded during all tasks (Kruskal-Wallis test, H = 0.75, $P = 0.687$).

B. As in A, for mean normalized change in theta coherence ($\Delta\theta$Co) during the encounter period.

Note the significant difference between the SP and other tasks (****p<0.0001, Dunn's *post-hoc*

test following the main effect in a Kruskal-Wallis test).

C-D. As in A and B, for gamma coherence (***p<0.001, ****p<0.0001).

E. Color-coded matrix of the mean normalized $\Delta\theta$Co (upper left) and $\Delta\gamma$Co (lower right) values

for all pairs of brain regions in the SP task. Empty spots represent brain region pairs with fewer

than five recorded sessions. Black spots separate between the $\Delta\theta$Co and $\Delta\gamma$Co matrices.

F. As in E, for EsP.

G. As in E, for SxP.

H. Coefficients and significance of Pearson's correlations of $\Delta\theta$Co across all coupled brain regions

for each pair of tasks (****p<0.0001, FDR adjusted).

I. As in H, for $\Delta\gamma$Co.

See also Fig. S5.

1015

Figure 5. Coherence changes during social investigation are informative regarding the social context.

A. Distribution of changes in theta coherence during investigation bouts ($^\Delta\theta$Co) between each pair of brain regions, plotted separately for each stimulus used in the SP (blue), EsP (green) and SxP (red) tasks. The names of brain region pairs which passed the mean cutoff value ± 1.5*SD are labeled, with those showing similarly high $^\Delta\theta$Co values for both stimuli of the same task in bold.

B. Spearman's correlation coefficients of mean $^\Delta\theta$Co across all paired brain regions, for couples of stimuli which were either used in the same task (left, brown bars), of the same type (middle, gray) or having the same valence (right, yellow). The correlated two stimuli are denoted by asterisk below each bar, while the statistical significance of the correlation is marked above the bars.

C. As in B, for $^\Delta\gamma$Co.

D. A color-coded confusion matrix for a multi-class Random forest classifier employed for predicting the social context from $^\Delta\theta$Co values across all brain regions and stimuli. The scale of the accuracy's color code is shown to the right. The percentage of cases a label was predicted for each ground truth are marked in the middle of each spot. *p<0.05, Mann-Whitney test, FDR adjusted.

E. As in D, for $^\Delta\gamma$Co.

See also Figs. S6, S7.

Figure 6. Distinct patterns of changes in Granger causality (GC) during the encounter period among tasks.

A. Schematic representation of the brain regions over-represented among the pairs that exhibited strong (mean $\pm 1.5*$SD) theta coherence bias towards one of the stimuli in any task. The regions are color-coded according to the task in which they were over-represented.

B. As in A, for gamma coherence.

C. Schematic representation of significant changes (encounter vs. baseline) in theta band GC during a SP task, among the regions listed in A. The direction of the GC changes in shown by a black arrow (top to bottom on the right and bottom to top on the left).

D. As in C, for EsP.

E. AS in C, for SxP.

F-H. As in C-E, for gamma band

See also Fig. S8

Figure 7. Context-specific differences in vDG LFP rhythmicity in specific behavioral events

A. Color-coded scheme of specific behavior event types, with ↓ indicating the beginning of an ,
   while **⅃** indicating the end of an investigation bout, and ***** indicating the beginning of a bout
   after transition between stimuli or repeated investigation of the same stimulus. Events showing
   significant differences in vDG LFP power are highlighted in yellow.

B. Super-imposed traces of the mean (±SEM) Z-score of changes in vDG theta power at the end
   of long bouts towards either a social stimulus (pink) or an object stimulus (purple) in the SP
   task. Time 0 represents the end of the bout, while the 5 s period before time 0 was considered
   as baseline. $*p<0.05$, Student's t-test between the mean Z-score values averaged over the last 3
   s of the traces.

C. As in B, for gamma power in the vDG during SP tasks.

D-E. As in B-C, for the EsP task.

F-G. as in B-C, for the SxP task.

H-I. As in B-C, for changes in LFP power at the beginning of repeated vs. transitional (between
   stimuli) investigation bouts of social and object stimuli across SP task sessions.

 J-K. As in H-I, for the EsP task. $**p<0.01$ by a Mann-Whitney test following the main effect in
   ANOVA.

L-M. As in H-I, for the SxP task.

See also Fig. S9.