Reproductive success mediates the effects of climate change and grassland management on
plant populations dynamics

Martin Andrzejak[1, 2, 3], Tiffany M. Knight[1, 2, 3], Carolin Plos[2, 3], Lotte Korell[1, 2, 3]

1. *Department of Community Ecology, Helmholtz Centre for Environmental Research*

GmbH–UFZ, Halle (Saale), Germany.

2. Institute of Biology, Geobotany and Botanical Garden, Martin Luther University Halle-

Wittenberg, *Germany*

3. *German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig,*

Germany

Open research statement

The data is currently available at figshare:

https://figshare.com/projects/Population_growth_rate_climate_change_and_land_use/157925

and the R code is currently available on GitHub:

https://github.com/Martin19910130/Climate_change_and_land_use_population_growth_rate

Upon acceptance the data will be available at figshare:

https://figshare.com/projects/Population_growth_rate_climate_change_and_land_use/157925

and the R code on GitHub:

https://github.com/Martin19910130/Climate_change_and_land_use_population_growth_rate

Abstract

Climate change is one of the largest threats to grassland plant species, which can be modified by land management. Although climate change and land management can separately and interactively influence plant demography this has been rarely considered within one experimental set-up. We used a large-scale experiment to quantify the effects of grassland management, climate change and their joint effect on the demography and population growth rate of 11 native plant species. We parameterized integral projection models with four years of demographic data to project the population growth rate. We hypothesized, plants would perform better in ambient than in the future climate treatment that creates hotter and drier summer conditions and that plant performance in grazing vs. mowing would vary across species and depend on their traits. Due to extreme drought events, over half of our study species went quasi extinct, which highlights how extreme climate events can influence long term experimental results. Of the persistent species, only one supported our expectations, and the rest either had higher population growth rates in the future climate treatment or showed no significant difference in population growth between climate treatments. Species with shorter flowering durations performed better in the mowing treatment while those with longer flowering durations performed better in the grazing treatment. The population growth rates of these species were sensitive to changes in vital rates related to reproduction and recruitment. Depending on the species we found effects of land management and climate change on population growth rates but we did not find strong support for interactive effects among both factors. Experiments combined with measuring plant demographic responses provide a way to isolate the effects of different drivers on the long-term persistence of species, and to identify the demographic vital rates that are critical to manage in the future. Our study suggests that it will become increasingly difficult

44 to maintain species with preferences for moister soil conditions, that traits such as flowering

45 duration might predict responses to management, and that vital rates such as reproduction and

46 recruitment are disproportionately important.

60 **Keywords**

61 Integral projection model, life table response experiment, sheep grazing, mowing, grassland

62 management, climate change experiment, plant population dynamics, elasticity analysis

Introduction

Grasslands cover approximately 40% of the Earth's surface (Bardgett et al., 2021) and provide important ecosystem services to society, such as food for livestock and humanity, pollination and biodiversity (Biurrun et al., 2021; Feurdean et al., 2018). In Central Europe, semi-natural grasslands were historically managed using low-intensity (i.e. extensive) mowing or grazing, which prevents their succession into forests (Poschlod et al., 1998; Tälle et al., 2015). These different types of management affect which types of plants can persist. For example, grazing favors plants with traits related to higher grazing resistance (e.g. hairiness) or grazing tolerance (e.g. resprouting capacity) (Bernhardt-Römermann et al., 2011; Evju et al., 2009; Kahmen et al., 2002). Anthropogenic climate change may modify the effects that management has on the demography and persistence of plant species (Bardgett et al., 2021; Kariyeva et al., 2012; Redlich et al., 2022). For example, individuals that currently avoid loss of reproduction from mowing by flowering before or after mowing events might shift in their phenology in the future due to climate change, which could lead to mismatches between flowering timing and mowing events that harm plant reproductive success (Cleland et al., 2006; Gordo & Sanz, 2010).

In central Europe, the climate is projected to become warmer, wetter in spring and fall, and drier in summer in the future (Döscher et al., 2002; Jacob & Podzun, 1997; Rockel et al., 2008; Wagner et al., 2013). Drought-tolerant species should be best able to survive the drier summers expected in the future (Belovsky & Slade, 2021). Species that flower earlier and across a longer time period might be favored in the future, because they can take best advantage of the wet spring conditions and because the long flowering duration might buffer their populations from complete reproductive failure during periods of extreme weather. This leads to the expectation that drought-tolerant plants with an early flowering start combined with a long flowering period

should be best able to avoid losing reproductive success under future climate conditions in the

presence of grazing and mowing land management types.

In order to understand population dynamics of plant species, demographic studies quantify vital

rates such as survival, growth and reproductive output across the entire life cycle of the plant and

project long-term population growth rate using structured population models (Caswell, 2000;

Crone et al., 2011; Easterling et al., 2000). A common approach to analyzing plant demographic

data are Integral Projection models (IPMs), which consider a relationship between continuous

plant size and vital rates and can also incorporate discrete development stages (Childs et al.,

2003; Jacquemyn et al., 2010; Yule et al., 2013). Retrospective analyses of IPMs, such as life

table response experiments, decompose the contribution of individual vital rates to the changes in

population growth rate between treatments. Vital rates that are more responsible for the change

are those that, either change dramatically between the treatments and/or those that the population

growth rate is sensitive to changes in (Rees & Ellner, 2009).

Many demographic studies have been conducted to assess how the demography of species

responds to climate (Ehrlen, 2019; Ehrlén et al., 2016; Ehrlén & Morris, 2015; Lemmer et al.,

2021; Morris et al., 2020). Most studies focus on natural variation across space and time in

climate conditions and measure plant demographic responses (Compagnoni et al., 2021). Such

observational studies are limited in their ability to assess cause and effect relationships, as there

are often other important factors that co-vary with climate. A few studies have experimentally

assessed how climate affects plant demography (Compagnoni & Adler, 2014; Lemmer et al.,

2021; Lyu & Alexander, 2022; Töpper et al., 2018; Williams et al., 2007). However, most

climate change experiments consider extreme manipulations of temperature and precipitation,

making it hard to predict how plants might respond under realistic future climate change

scenarios (Korell et al., 2020).

We have a unique opportunity to study how a realistic future climate scenario as well as two

different grassland management types (extensive grazing vs. mowing) influence the demography

and population dynamics of plants using the Global Change Experimental Facility (GCEF) in

Germany. Our study builds on the research of Lemmer and colleagues (Lemmer et al., 2021),

which discovered that land management types and climate change interactively affected the

population growth rate of the drought-tolerant grass, *Bromus erectus* L., and that this interaction

was primarily due to higher seedling recruitment of plants under the future climate and mowing

management treatment combination. Lemmer et al. (2020) measured the demography across a

short time period (one transition, two years) and concentrated on a single species. In this study,

we expand the scope to understand the demography and population responses to climate and land

management types across multiple transitions along a longer time period and multiple plant

species. Demographic data was collected over the course of five years (2018 – 2022) for eleven

(Table 1) grassland plant species that represent a range of different plant statures (tall vs. small),

life forms (grasses, herbs) and life histories (biennials vs perennials). In general, we expected i)

the effects of land management types will vary across species and depend on their traits, ii)

plants show higher population growth rates under ambient compared to future climate conditions

and iii) the effects of climate treatments and land management types on the population growth

rate will be interactive.

Methods

Study site

We conducted our research in the Global Change Experimental Facility (GCEF), which was

established in 2013. The experiment is part of the research station Bad Lauchstädt (51°22060 N,

11°50060 E, 118 m a.s.l.). It is designed to simulate a realistic future climate scenario combined

with the manipulation of different land management types, including extensively managed

grasslands. The future climate treatment is based on a mean of 12 regional climate models for the

region (Schädler et al., 2019).

The GCEF consists of 10 main units, five with ambient climate and five with future climate

conditions. Precipitation in the future climate treatment is manipulated seasonally: increases in

precipitation by +10 % compared to the ambient climate in spring and fall and reduced

precipitation by -20 % compared to the ambient climate in summer (for more details see

Schädler et al. 2019). The minimum temperature in the future climate treatment is increased by

about +1.5°C by passive nighttime warming (Schädler et al. 2019).

Each main unit is divided into five subunits that are 16m x 24m in size, each representing a

different land management. Each subunit has 14m x 19.5m available for measurements and the

remainder is left as an edge area, where no experiments or measurements can be undertaken

(Appendix S1, Figure S1). Our study was established in two of the five land managements, the

extensive meadow (mown one or two times per year) and the extensive pasture (grazed by 20

sheep for 24 hours twice to three times per year). These two land management types were

initially sown in 2013 with 56 plant species native to Germany (Schädler et al., 2019). This

design leads to the following treatment combinations in our experiment: (1) ambient grazing, (2)

ambient mowing, (3) future grazing and (4) future mowing.

Study species

In 2018, we chose eleven different species (Table 1) for our demographic study based on the

criteria that they were abundant enough for demographic data collection (minimum number of

individuals in 2018 was 50 per treatment combination), and that they were species that are

typically associated with mesic or dry grasslands. The species covered a range of growth forms,

plant families and life histories (biennial vs. perennial). Every plant species in this study is native

to Germany and was sown into the GCEF experiment when it was established.

Demographic data collection

In April 2018, we established one transect on all our study subunits resulting in 20 transects (five

replicates per treatment combination). Our goal was to find a minimum of 10 individuals per

transect, to reach a minimum of 50 individuals per treatment combination for every species.

Along each transect, 50 x 50 cm subplots were established. We assigned three of the subplots to

a fixed positions along each transect (0– 0.50 m; 3 – 3,50 m; 6-6.50 m). The location of other

subplots were chosen along the transect at strategic locations to capture individuals of species

that were needed to achieve our target of 50 individuals per species and transect. Following this

procedure, most transects had eight subplots. This leads to a different overall subplot number for

each species, with rarer species having more subplots than common species. Each subplot was

marked with a land marking (pipe with red plastic top) in the bottom left and a nail with a white

plastic ring the top right corner.

In April 2018, we measured the size of each individual target species in each subplot using either

basal area or number of leaves (Table 1). Basal area (cm^2) measured as the product of the

horizontal width and length at the base of each individual. Length was measured as the longest

lateral dimension of each plant, and width was measured perpendicular to the length. For rosette

species (e.g. *Plantago lanceolata, Tragopogon orientalis*), we used the number of leaves as

proxy for their size. We recorded the location of each individual within the 50 x 50 cm subplot,

using an XY-coordinate system so that individuals could be followed over the years. In rare

cases if we spotted individuals of our rarer species just outside of our subplot, we would measure

those individuals and track their position relative to the subplot (one of the coordinates would

exceed 50 cm).

From 2019 until 2022, we went to the field five times a year. In April, we re-located individuals

from the previous year and measured their size, identified new individuals and measured their

size, and counted the number of new seedlings (for details on how we identified seedlings see

Appendix S3, Table S2). Depending on the phenology of the species and the management events

(see Table 1, Table S1), we counted flowers and seed heads of all reproductive individuals in

each transect. Further we collected two seed heads per subunit and species from individuals that

were outside of our transects but within the same subunits. Number of seeds in each seed head

were counted in the lab. In late autumn (October/November), we counted the number of

seedlings in our subplots, as seedlings of these species can germinate in spring or in fall.

From 2019 onwards, we decided to mark individuals of the very abundant species *Bromus*

erectus with ID's as that helped separating and finding individuals. To mark the individuals, we

used rings made of wire with a tag with a running number on it and put it around the base of the

individual. We left enough space for the individual to grow. Special care was taken to not harm

any of the vegetation during that process.

The multiple droughts Germany experienced during several of the years of this study prevented

more frequent mowing and grazing than would otherwise be incorporated into the treatment plan

(Schädler et al., 2019). During the time of our experiment, the extensive meadows were mown

once per year in early summer time from 2018 – 2020 and two times (early and late summer) in

2021. The extensively grazed subplots were grazed twice per year in spring and early summer

from 2018-2020, and three times in 2021 (spring, early and late summer). For a detailed look at

the management timing, see Appendix S2, Table S1.

For information on traits for our species, we used Ellenberg values (Ellenberg et al., 1991), the

Bioflor database (https://wiki.ufz.de/biolflor/index.jsp) and our own measurements of phenology.

For drought tolerance, we used categorical Ellenberg values (Ellenberg et al., 1991, Table 1). For

grazing and mowing resistance, we used categorical values from the Bioflor database (Table 1).

In 2020, we collected phenology data on our target species in the GCEF between April and

December within the central 3 x 3 m vegetation survey plot of each subunit where no

experiments or other disturbances (besides the management events) occur. This provides

information on the month in which flowering began and the month it ended, taking all

individuals inside the central vegetation plot into account.

Quasi extinction

Some of the species declined so significantly over the course of our demographic study, that we considered them quasi extinct and no longer continued to collect demographic data. We defined quasi extinction from a treatment combination as fewer than 25 individuals total and fewer than 10 flowering individuals, thus those species treated the species as quasi extinct.

Life cycle stages and vital rates

For all species, we modeled a year-to-year life cycle (April to April) with two stage classes. A discrete class for seedlings and a continuous class for plants (Figure 1). Most of our species do not form long lasting seed banks, and thus we do not have a seed bank class. *Plantago lanceolata* does have a long-lived seed bank (Chen, 2018), but most of the seeds of this species germinate immediately and therefore demographic studies on this species do not typically incorporate the seed bank stage (e.g., www.plantpopnet.com).

We modeled each continuous state of the IPM as a function of the natural logarithm of the size of an individual. We used generalized linear models to fit all continuous variables. Survival is a function of whether or not an individual (i) was still alive at t1, dependent on its log transformed size (z) at t0 (Eq. 1), modeled as a Bernoulli process with probability of survival S_{t1} (Eq. 2).

[1] $$S_{i,t1} \sim Bernoulli\left(\hat{S}_{t1}\right)$$

[2] $$logit\left(\hat{S}_{t1}\right) = \alpha + \beta \, log_e(z)$$

In this function α is the intercept and β the slope of the curve.

Growth of plants $G_{i,t1}$ is described as the normally distributed change in z and log transformed

size at t1 (z') of a surviving individual plant (i) (Eq. 3). It was modeled as a linear function of z'

of surviving individuals depending on z, where the intercept is defined as α^G, the slope as β^G

(Eq. 4) and the standard deviation as σ_G (Eq. 3).

[3] $G_{i,t1} \sim Normal\left(\hat{G}_{t1}, \sigma_G\right)$

[4] $\hat{G}_{i,t1} = \alpha^G + \beta^G \, log_e(z)$

Similar to survival, we modeled reproduction probability ($P_{i,t0}$) as Bernoulli process (Eq. 5).

Here we tested how the probability an individual i reproduces at time t0 depends on z. The

intercept is defined as α_t^P and the slope as β^P using a logit link function (Eq. 6)

[5] $P_{i,to} \sim Bernoulli\left(\hat{P}_t\right)$

[6] $logit\left(\hat{P}_{to}\right) = \alpha_t^P + \beta^P \, log_e(z).$

For every reproductive plant, we calculated the number of seeds it produced (Eq. 7). The number

of seeds produced by an individual I at t0 was a function of z, modeled as a Poisson distribution.

The intercept was defined as α_t^F and slope β^F (Eq. 8)

[7] $F_{i,to} \sim Poisson\left(\hat{F}_{to}\right)$

[8] $\hat{F}_t = \alpha_t^F + \beta^F \, log_e(z).$

Recruitment, the emerged seedlings per total number of seeds, was separated into fall - ($\theta_{f,\,j,\,t0}$)

and spring recruitment ($\theta_{s,\,j,\,t0}$). To calculate recruitment, we summed the total number of seeds

per subplot ($S_{j,t}$, Eq. 9). Where $S_{i,t1}$ is the total number of seeds produced by one individual (n) in

a subplot at time t1.

[9] $S_{j,t} = \sum_{1}^{n} S_{i,t1}.$

We then divided the result of Eq. 9 by the number of seedlings found in that subplot in fall t0

($Rf_{j, to}$, Eq. 10) and spring t1 ($Rs_{j,t1}$, Eq. 11).

[10] $\theta_{f,j,t} = \hat{R}f_{j,to} / S_{j,t}.$

[11] $\theta_{s,j,t+1} = \hat{R}s_{j,t1} / S_{j,t}.$

Establishment (B_j) is the proportion of seedlings that survive to become plants from t0 to t1 in

each subplot j. We calculated the sum of seedlings in spring and fall t0 ($Rsum_{j, t0}$) of a subplot

and divided that number by the number of new individuals at t1 ($Ni_{j, t1}$) in the same subplot.

[12] $B_j = Ni_{j,t1} / Rsum_{j,t0}$

We calculated the log size distribution of new plants η which is defined as the normally

distributed size of individuals that entered the continuous plant stage in year t1 (Eq. 12). We used

the mean ($log_e(\hat{\eta}_{t1})$) and standard deviation (σ_η) of this size in our final parametrization of the

IPM.

[13] $log_e(\eta_{i,t1}) \sim Normal\left(log_e(\hat{\eta}_{t1}), \sigma_\eta\right)$

Effects of treatments on vital rates

For each vital rate (survival, growth, reproduction probability, number of seeds, recruitment,

establishment, size distribution of new plants), we generated three independent pipelines. The

first pipeline differentiated between the climate treatment and thus we calculated the parameters

for ambient and future climate, respectively (Table 2). The second pipeline differentiated

between land management types (mowing and grazing). In the third pipeline, we differentiated

between all four treatment combinations to also account for possible interactions among

treatments. In each of those models, we dropped year, to make the models more robust (but see

Appendix S5 for the "yearly" models).

Integral projection model (IPM)

For each species and pipeline, we created an IPM to calculate the influence of climate and

management on the long-term population growth (λ). In every IPM, we calculate all possible

transitions from z to z' at. The change in number of plants from one year to the other is described

by:

[14] $$n(z',t1) = M(t0)B\eta(z') + \int_{L}^{U} S(z)G(z',z) + P(z)F(z)\theta_f B\eta(z')n(z,t0)dz$$

Where the vector n(z', t1) is the number of plants at size z' at time t1. In this function, the first

term represents the recruitment of spring seedlings entering the adult plant class. This is based on

the number of spring seedlings at t0, M(t0), the seedling establishment, B, and the size

distribution of new plants η(z'). The second kernel (or surface) describes the growth and survival

of plants from t0, n(z, t0) to t1, n(z', t1). It is defined as an integral with in the upper limit U,

which is the biggest size observed and the lowest size L, observed for the species in our study.

The integrals were evaluated across 200 equally spaced size bins using the midpoint rules as a

200 x 200 matrix.

The recruitment of spring seedlings from one year to the next is described by:

[15] $M(t + 1) = \int_L^U P(z)F(z)\theta_s \, dz.$

As we did not distinguish the new plants we see in April as those that were seedlings from fall

vs. spring, we use the same establishment estimate for seedlings (B) twice in the model.

Effects of treatments on population growth rate

To determine the effect of the climate treatment on λ, we calculated λ for the ambient and future

treatments and then calculated an effect size by subtracting λ ambient from λ future. Positive

values show a species´ preference for ambient climate conditions whereas negative values show

a species´ preference for future climate conditions. Likewise, for land management types, we

subtracted λ grazed from λ mowing to calculate an effect size. Here, positive values show a

species preference for the grazing treatment whereas negative values show a species favors the

mowing treatment.

To test whether traits influenced the effect size of land management types on λ, we ran simple

linear models using the effect size as response variable and the following traits as explanatory

variable: flowering start month, duration of flowering, grazing resistance and mowing resistance.

Because flowering time, start and end could differ between treatments due to the effect of the

land management type we decided to not calculate a mean across those.

To quantify uncertainty in estimates of λ, we bootstrapped our data a thousand times and

calculated the λ each time. To test for significant differences in λ between treatments and

treatment combinations, we used permutation (randomization) tests (N = 1000 permutations).

For each treatment combination, we calculated sensitivities and elasticities of λ to kernel element

perturbations.

Life table response experiment (LTRE)

In order to decompose the influences of vital rates on the difference in λ, across treatment

combinations, we conducted Life Table Response Experiments (LTRE) for each pairwise

treatment combination. The difference in λ, $\Delta\lambda$, is:

[17] $\quad\quad\quad\quad \Delta\lambda = \lambda^{treatment\ combination\ 1} - \lambda^{treatment\ combination\ 2}$

The contribution of each vital rate to $\Delta\lambda$, $\widetilde{\delta}^{\ \alpha i}$, is:

[18] $\quad\quad\quad\quad \widetilde{\delta}^{\ \alpha i} = \sum_{i}^{14}(a_i^{treatment\ combination\ 1} - a_i^{treatment\ combination\ 2})\frac{\partial\lambda}{\partial\alpha_i}$

Here α_i is one of the fourteen vital rates that are included in the IPMs, while the sensitivity of λ to

each vital rate is defined as $\frac{\partial\lambda}{\partial\alpha_i}$. Vital rates will have a strong contribution to the observed

difference in λ between treatment combinations if the vital rates change strongly in magnitude

between the two treatment combinations and/or if λ is sensitive to changes in that vital rate. To

facilitate the understandability, we combined the vital rates with respect to six demographic

processes: survival, growth, reproduction, recruitment, establishment (Table 2). After this we

scaled $\widetilde{\delta}^{\ \alpha_i}$ to 1 (to display proportional influence).

Analysis

All analysis were carried out in R (R Core Team, 2022). Figures were made using the ggplot2

package (Wickham, 2016). The integral projection model was run and calculated using the ipmr

packages (Levin et al., 2021). Other packages used were: openxlsx (Schauberger & Walker,

2021) to load the xlsx files into R, dplyr (Wickham et al., 2022) for structuring data and piping

and the package parallel (R Core Team, 2022) to decrease the running time of R scripts.

Results

Quasi extinction

The following six species went quasi extinct over the course of the experiment: *Anthoxanthum*

odoratum, Crepis biennis, *Lotus corniculatus*, *Lychnis flos-cuculi* and *Trifolium pratense*. While

there is too small of a sample size (N=11 species) for statistical analysis, it is notable that the

species that went quasi extinct were also the ones that were less drought tolerant based on their

Ellenberg values (Table 1) and that the first few years of our study were extraordinarily dry for

our region (Boergens et al., 2020).

Management effects on population growth rate

Tragopogon orientalis had significantly higher λ in mowing compared to grazing management

(permutation test p = 0.042, Figure 2e), whereas all other species did not show a significant main

effect of management on λ (Figure 2a-d).

Traits related to grazing resistance and mowing resistance did not explain the effect size of land

management type on λ (Appendix S4, Figure S2). However, flowering duration was positively

related to the effect size of land management type on λ; species with lower flowering duration

(i.e. *Plantago lanceolata, Bromus erectus, Tragopogon orientalis*) performed better in the

mowing treatment while those with longer flowering duration (*Dianthus carthusianorum*,

Scabiosa ochroleuca) performed better in the grazing treatment (t = 6.087, p < 0.001, r^2 = 0.8;

Figure. 3a). The start of flowering was a poor predictor of management preferences (Figure 3b, t

= -0.206, p = 0.842, r^2 = -0.1191).

of flowering for each species. The line indicates the trend from the linear model. Symbols show the

management treatment in which phenology was measured.

Climate effects on population growth rate

B. erectus had a significantly higher λ in future compared to ambient climate (permutation test, p

= 0.023, Figure 4a), and the other four species did not show significant differences in λ with

regard to the climate treatment (Figure 4b-e).

Effect of climate and land management type on population growth rate

B. erectus had a significantly higher λ in the ambient mowing than in the ambient grazing

treatment combination (permutation test, p = 0.001). In *S. ochroleuca,* λ was significantly higher

in future compared to ambient climate in the mowing land management type (future mowing >

ambient mowing; permutation test, p = 0.011) but not in grazing land management (permutation

test, p > 0.05, Figure 5d). The λ of *T. orientalis* was significantly higher in ambient compared to

future climate only in the grazing management treatment (ambient grazing > future grazing;

permutation test, p = 0.002) and λ was higher in mowing compared to grazing management in

both climate treatments (permutation test, ambient grazing > ambient mowing, p = 0.048, future

grazing > future mowing, 0.023 Figure 5e). For all other species (*B. erectus*, *D. carthusianorum*

and *P. lanceolata*) the lines in the interaction plots crossed but they were not significantly

different from each other due to the large variability around the mean (Figure 5a-c).

Sensitivity

Across all species and treatments, λ was most sensitive to changes in reproduction parameters

(Figure 6). While the magnitude differed between treatments and species (e.g. *S. ochroleuca*

highest in grazing and future *D. carthusianorum* highest in ambient).

Life Table Response Experiment

The higher λ of *B. erectus* in future compared to ambient climate was primarily due to increases

in reproduction and recruitment of this species under future climate (Figure 7c and 7d). The

higher λ of *S. ochroleuca* in future compared to ambient climate in the mowing management

treatment was primarily due to increases in the reproduction of this species in the future mowing

treatment (Figure 7d). The higher λ of *T. orientalis* in ambient compared to future climate in the

grazing management treatment was primarily due to increases in plant survivorship (Figure 7c),

whereas the higher λ in this species in mowing compared to grazing in future climate treatment

was due to higher survival and reproduction (Figure 7b).

Discussion

Our study aimed to examine whether plant species interactively respond to climate and land

management types in their demography and population dynamics. However, in the early years of

our study, six of our eleven focal species went quasi extinct from the experiment, likely due to

the extreme drought conditions that occurred in our region in 2018 and 2019. This is becoming

an increasing phenomenon in experimental global change research—even the control treatments

experience extreme climate events because these are now more common in our rapidly changing

world (IPCC, 2014; Jentsch et al., 2007; Orlowsky & Seneviratne, 2012; Tebaldi et al., 2006). Of

the remaining five species, we found that flowering duration is an important trait that predicts

responses to land management types, with shorter duration species having higher λ in mowing

management and longer duration species having higher λ in grazing management. In contrast to

our expectation that plants would perform better in ambient compared to future climate

conditions, we found that two species (*B. erectus* and *S. ochroleuca*) have higher λ in the future

climate treatment and only *T. orientalis* shows the expected higher λ in ambient climate. We

suggest this is likely due to the high drought tolerance of *B. erectus* and *S. ochroleuca*. We did

not find strong support that the effects of climate and land management types were interactive

for our study species, and it is unclear if this is because these interactive effects do not occur or

because we have low power to detect them.

We expected that plants species in the study region (Central Germany) would perform best in

ambient climate, as they would have difficulty dealing with the dry conditions expected in the

summer under future climate. However, our results do not support this hypothesis. The first year

of our demographic study, 2018, coincided with the driest summer for Germany since start of

recording (Schuldt et al., 2020). This drought had dramatic and long-lasting effects on the plant

communities across the entire GCEF experiment. Many of the species that went quasi-extinct

were those that are indicative of moist soil conditions. Four of the five species that remained are

indicative of drier habitat conditions and commonly occur in semi-dry or dry meadows (*B.*

erectus, *D. carthusianorum* and *S. ochroleuca*) or are known to be highly generalized with

regards to the moisture of the habitat (*P. lanceolata*). *B. erectus* and *S. ochroleuca* were also the

two species that performed better in future experimental climate conditions. This result matches

previous experimental studies (Lemmer et al., 2021; Moser et al., 2011; Poniatowski et al., 2018)

that show high performance of *B. erectus* in hot and dry summer conditions. Our LTRE results

revealed that the higher λ of *B. erectus* in future climate was due primarily to the higher

reproduction and recruitment in this treatment (see also Lemmer et al. 2021). The earlier

flowering of *B. erectus* compared to other species could explain its relative advantage in future

climate conditions; *B. erectus* is able to take advantage of the additional water resources in

spring and reproduce before the hot and dry summer conditions in the future climate treatment.

Only one of our remaining species, *T. orientalis,* is indicative of moist to moderately dry soil

conditions (Sebald et al., 1990; Ellenberg et al., 1991; Bomble, 2013) and managed to persist in

our study. This species did indeed show the expected higher performance in the ambient

compared to future climate treatment. We suggest that we may have found more support for our

hypothesis if we had been able to study the demography of more typical species from moister

grassland communities as we initially planned. Our study highlights the importance of historical

legacies in shaping experimental results and support projections that communities may shift

towards more drought tolerant species under future climate conditions (Belovsky & Slade, 2021;

Craine et al., 2013).

Surprisingly, most of our study species did not show strong differences in their population

growth across land management types. The only exception was the biennial species, *T.*

orientalis, which performed significantly better in the mowing compared to grazing management

treatment, primarily due to higher survival and reproduction in the mowing treatment. The

vegetative parts of *T. orientalis* are relatively soft and easy for sheep to chew (personal

observation) and the large flower heads may attract sheep. We also observed that the flower buds

were eaten by other wild animals, such as rabbits (personal observation). This species adds to

several studies that showed that grazing has a negative effect on plant population dynamics

(Hansen & Wilson, 2006; Jacquemyn et al., 2012; van der Meer et al., 2014). For example, in

Primula veris, grazing was found to decrease population growth rate, primarily through its

negative effect on reproduction (Jacquemyn et al., 2012).

Our results show that the effect of management on λ depended on flowering duration. Plants that

flowered a relatively short duration showed higher λ in mowing (e.g. *T. orientalis*), whereas

those flowering across a longer duration showed higher λ in the grazing management. This could

be because plant populations that flower across a long time period are buffered from losing all of

their reproductive success from disturbances that occur in a short time interval, such as grazing.

However, we have to be cautious when interpreting these results as in our study years the

management was mostly applied in a lower frequency than usual; most years had one mowing

and two grazing events. This was because mowing and grazing were not possible when there was

too little biomass regrowth, which occurred during the dry study years. Other studies have also

shown that low frequency (once a year) and late season mowing benefits the reproduction of

plants (Jantunen et al., 2007; Johansen et al., 2019). Across all of our study species, we found

that the population growth rates were most sensitive to reproduction. Thus, it may be a general

finding that management treatments should be timed to interfere as little as possible with peak

flowering and fruiting (Nakahama et al., 2016; Reisch & Poschlod, 2009).

We expected that interaction effects of treatments on λ would be common in our study and that

each species would have a treatment combination that was most optimal for λ. However, we had

little power to test for these interactions due to the high variation across individuals in the

sensitive vital rates of seed production and recruitment. For example, the high variation in λ of *S.*

ochroleuca in the ambient-grazing treatment combination is likely due to very high seed

production (>2000 seeds) of three individuals in this treatment combination (Appendix S6).

Similarly, the high variation in λ in the ambient treatment for *D. carthusianorum* is likely due to

high seed production of a few individuals in the ambient mowing treatment (Appendix S6) and

high recruitment of new spring seedlings in a few plots in the ambient grazing treatment

(Appendix S7). Finally, while *P. lanceolata* appears overall have higher λ in ambient compared

to future climate treatments, high variance in the future-mowing treatment combination, which is

likely due to two reproductive individuals with outlier high seed production, may prevent the

detection of an effect. However, it is noteworthy that across all species in this study many of the

plants with outlier seed production were in the ambient climate treatment, suggesting that the

future climate treatment might negate the positive effects that a preferred management treatment

might have on the reproduction and population growth rate of these grassland species.

In demographic studies of both plants and animals, reproductive output is known to be skewed

across individuals, with only a few individuals producing many offspring (van Daalen & Caswell,

2020). The ability to distinguish deterministic treatments effects from stochasticity among

individuals would require much higher sample sizes than is available in our GCEF experiment.

While there are many advantages of conducting multi-species demographic studies like this one

and of conducting demographic research in a controlled experimental setting, one distinct

disadvantage is the effort and sample size required to adequately detect small changes in

sensitive vital rates (McMahon et al., 2019).

It was surprising to find in our LTRE that changes in reproduction and recruitment often

contributed most to the observed responses of λ to management and climate. Usually in longer

lived species, λ is highly sensitive to changes in survival (Weppler, Stoll and Stöcklin, 2006;

Kuss et al., 2008) and thus even small changes in survival across treatments should contribute

significantly to changes in λ. In our case, we find that λ is highly sensitive to recruitment. This

might be due to the relatively high λ of our perennial species; species with high λ are often more

sensitive to reproduction and recruitment than to survivorship (Franco & Silvertown, 2004; Ramula

et al., 2008). Further, many of our species were able to maintain their survival across treatments,

but had high changes in their reproduction and recruitment.

Conclusions

Our study is a first step towards answering important yet unanswered question on how climate

change affects population dynamics of grassland species and how this effect is modified by

different land management types (Ehrlen, 2019). Our results are optimistic for our four more

drought-tolerant species characteristic for dry or semi-dry grasslands, showing persistent or

growing populations in both management types (grazing vs. mowing) and under future climate

conditions as well as after natural extreme drought events. However, less drought-tolerant

species either went quasi extinct early in the study or were projected to decline in the future,

indicating that populations cannot persist in the presence of future climate conditions. Our results

suggest that it will be more difficult to create and maintain especially mesophilic grasslands

composed of species with preference for moister soil conditions in the future. Furthermore, we

suggest that management in future climate should carefully be timed outside of peak flowering

and fruiting, so that as many species as possible can reproduce and persist in managed

grasslands.

Acknowledgements

This research was funded by the Alexander von Humboldt Foundation (Alexander von Humboldt Professorship of TMK) and the Helmholtz Recruitment Initiative of the Helmholtz Association (TMK). The authors gratefully acknowledge the support of iDiv funded by the German Research Foundation (DFG-FZT 118). We appreciate the Helmholtz Association, the Federal Ministry of Education and Research, the State Ministry of Science and Economy of Saxony-Anhalt, and the State Ministry for Higher Education, Research and the Arts Saxony, which fund the Global Change Experimental Facility (GCEF) project. We thank the staff of the Bad Lauchstädt Experimental Research Station (especially Ines Merbach and Konrad Kirsch), for their work in in maintaining the plots and infrastructures of the GCEF, and François Buscot, Stefan Klotz, Thomas Reitz, and Martin Schädler for their role in setting up the GCEF. We thank Julia Lemmer, Christian Savona, Simon Bitzan, Georg Küstner, Lina Lüttgert and Amibeth Thompson for their help with the fieldwork.

Author contributions

The study was designed by LK (leading) and MA and TMK (supporting). Fieldwork was designed and carried out by LK and MA (both leading). The data for the phenology of plants was collected by CP (leading). Analysis was carried out by MA (leading) with support of the other co-authors. The manuscript was written by MA (leading) and TMK and LK (supporting). All authors commented and improved the manuscript.

Conflict of interest

No Author has a conflict of interest

References

Bardgett, R. D., Bullock, J. M., Lavorel, S., Manning, P., Schaffner, U., Ostle, N., Chomel, M.,
Durigan, G., L. Fry, E., Johnson, D., Lavallee, J. M., le Provost, G., Luo, S., Png, K.,
Sankaran, M., Hou, X., Zhou, H., Ma, L., Ren, W., … Shi, H. (2021). Combatting global
grassland degradation. *Nature Reviews Earth and Environment*, *2*(10), 720–735.
https://doi.org/10.1038/s43017-021-00207-2

Belovsky, G. E., & Slade, J. B. (2021). Climate change and primary production: Forty years in a
bunchgrass prairie. *PLoS ONE*, *15*(12 December).
https://doi.org/10.1371/journal.pone.0243496

Bernhardt-Römermann, M., Gray, A., Vanbergen, A. J., Bergès, L., Bohner, A., Brooker, R. W.,
De Bruyn, L., De Cinti, B., Dirnböck, T., Grandin, U., Hester, A. J., Kanka, R., Klotz, S.,
Loucougaray, G., Lundin, L., Matteucci, G., Mészáros, I., Oláh, V., Preda, E., … Stadler, J.
(2011). Functional traits and local environment predict vegetation responses to disturbance:
A pan-European multi-site experiment. *Journal of Ecology*, *99*(3), 777–787.
https://doi.org/10.1111/j.1365-2745.2011.01794.x

Biurrun, I., Pielech, R., Dembicz, I., Gillet, F., Kozub, Ł., Marcenò, C., Reitalu, T., Van
Meerbeek, K., Guarino, R., Chytrý, M., Pakeman, R. J., Preislerová, Z., Axmanová, I.,
Burrascano, S., Bartha, S., Boch, S., Bruun, H. H., Conradi, T., De Frenne, P., … Dengler,
J. (2021). Benchmarking plant diversity of Palaearctic grasslands and other open habitats.
Journal of Vegetation Science, *32*(4). https://doi.org/10.1111/jvs.13050

Boergens, E., Güntner, A., Dobslaw, H., & Dahle, C. (2020). Quantifying the Central European
Droughts in 2018 and 2019 With GRACE Follow-On. *Geophysical Research Letters*,
47(14). https://doi.org/10.1029/2020GL087285

Bomble, W. F. (2013). *Tragopogon-Bocksbart (Asteraceae) in Nordrhein-Westfalen*.

Caswell, H. (2000). *Matrix population models (Vol.1)* (Vol. 1).

Chen, S. (2018). *Modelling plant populations - do clonality and seed ecology matter?* .

Childs, D. Z., Rees, M., Rose, K. E., Grubb, P. J., & Ellner, S. P. (2003). Evolution of complex
flowering strategies: An age- and size-structured integral projection model. *Proceedings of*
the Royal Society B: Biological Sciences, *270*(1526), 1829–1838.
https://doi.org/10.1098/rspb.2003.2399

Cleland, E. E., Chiariello, N. R., Loarie, S. R., Mooney, H. A., & Field, C. B. (2006). Diverse
responses of phenology to global changes in a grassland ecosystem. *Proceedings of the*
National Academy of Sciences of the United States of America, *103*(37), 13740–13744.
https://doi.org/10.1073/pnas.0600815103

Compagnoni, A., & Adler, P. B. (2014). Warming, competition, and Bromus tectorum
population growth across an elevation gradient. *Ecosphere*, *5*(9).
https://doi.org/10.1890/ES14-00047.1

Compagnoni, A., Levin, S., Childs, D. Z., Harpole, S., Paniw, M., Römer, G., Burns, J. H., Che-
Castaldo, J., Rüger, N., Kunstler, G., Bennett, J. M., Archer, C. R., Jones, O. R., Salguero-
Gómez, R., & Knight, T. M. (2021). Herbaceous perennial plants with short generation time
have stronger responses to climate anomalies than those with longer generation time.
Nature Communications, *12*(1). https://doi.org/10.1038/s41467-021-21977-9

Craine, J. M., Ocheltree, T. W., Nippert, J. B., Towne, E. G., Skibbe, A. M., Kembel, S. W., &
Fargione, J. E. (2013). Global diversity of drought tolerance and grassland climate-change
resilience. *Nature Climate Change*, *3*(1), 63–67. https://doi.org/10.1038/nclimate1634

Crone, E. E., Menges, E. S., Ellis, M. M., Bell, T., Bierzychudek, P., Ehrlén, J., Kaye, T. N.,
Knight, T. M., Lesica, P., Morris, W. F., Oostermeijer, G., Quintana-Ascencio, P. F.,
Stanley, A., Ticktin, T., Valverde, T., & Williams, J. L. (2011). How do plant ecologists use
matrix population models? *Ecology Letters*, *14*(1), 1–8. https://doi.org/10.1111/j.1461-
0248.2010.01540.x

Döscher, R., Willén, U., Jones, C., Rutgersson, A., Meier, H. E. M., Hansson, U., & Graham, L.
P. (2002). The development of the regional coupled ocean-atmosphere model RCAO.
Boreal Environment Research, *7*(3), 183–192.

Easterling, M. R., Ellner, S. P., & Dixon, P. M. (2000). Size-Specific Sensitivity: Applying a
new sturctured population model. *Ecology*, *81*(3), 694–708.

Ehrlen, J. (2019). Climate change in grasslands - demography and population dynamics. In D. J.
Gibson & J. A. Newman (Eds.), *Grasslands and climate change* (pp. 172–187).

Ehrlén, J., & Morris, W. F. (2015). Predicting changes in the distribution and abundance of
species under environmental change. In *Ecology Letters* (Vol. 18, Issue 3, pp. 303–314).
https://doi.org/10.1111/ele.12410

Ehrlén, J., Morris, W. F., von Euler, T., & Dahlgren, J. P. (2016). Advancing environmentally
explicit structured population models of plants. *Journal of Ecology*, *104*(2), 292–305.
https://doi.org/10.1111/1365-2745.12523

Ellenberg, H., Weber, H. E., Düll, R. , Wirth, V., Werner, & W. Paulissen, D. (1991).
Zeigerwerte von Pflanzen in Mitteleuropa (3rd ed.). Erich Goltze KG.

Evju, M., Austrheim, G., Halvorsen, R., & Mysterud, A. (2009). Grazing responses in herbs in
relation to herbivore selectivity and plant traits in an alpine ecosystem. *Oecologia*, *161*(1),
77–85. https://doi.org/10.1007/s00442-009-1358-1

Feurdean, A., Ruprecht, E., Molnár, Z., Hutchinson, S. M., & Hickler, T. (2018). Biodiversity-
rich European grasslands: Ancient, forgotten ecosystems. *Biological Conservation*,
228(May), 224–232. https://doi.org/10.1016/j.biocon.2018.09.022

Franco, M., & Silvertown, J. (2004). A Comparative Demography of Plants Based upon Elasticities of Vital Rates. In *Source: Ecology* (Vol. 85, Issue 2).

Gordo, O., & Sanz, J. J. (2010). Impact of climate change on plant phenology in Mediterranean ecosystems. *Global Change Biology*, *16*(3), 1082–1106. https://doi.org/10.1111/j.1365-2486.2009.02084.x

Hansen, M. J., & Wilson, S. D. (2006). Is management of an invasive grass Agropyron cristatum contingent on environmental variation? *Journal of Applied Ecology*, *43*(2), 269–280. https://doi.org/10.1111/j.1365-2664.2006.01145.x

IPCC. (2014). IPCC, 2014: Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, R.K. Pachauri and L.A. Meyer (eds.)]. *IPCC, Geneva, Sw*, 151.

Jacob, D., & Podzun, R. (1997). Sensitivity studies with the regional climate model REMO. *Meteorology and Atmospheric Physics*, *63*(1–2), 119–129. https://doi.org/10.1007/BF01025368

Jacquemyn, H., Brys, R., Davison, R., Tuljapurkar, S., & Jongejans, E. (2012). Stochastic LTRE analysis of the effects of herbivory on the population dynamics of a perennial grassland herb. *Oikos*, *121*(2), 211–218. https://doi.org/10.1111/j.1600-0706.2011.19774.x

Jacquemyn, H., Brys, R., & Jongejans, E. (2010). Size-dependent flowering and costs of reproduction affect population dynamics in a tuberous perennial woodland orchid. *Journal of Ecology*, *98*(5), 1204–1215. https://doi.org/10.1111/j.1365-2745.2010.01697.x

Jantunen, J., Saarinen, K., Valtonen, A., & Saarnio, S. (2007). Flowering and seed production success along roads with different mowing regimes. *Applied Vegetation Science*, *10*(2), 285. https://doi.org/10.1658/1402-2001(2007)10[285:faspsa]2.0.co;2

Jentsch, A., Kreyling, J., & Beierkuhnlein, C. (2007). A new generation of climate-change experiments. *Frontiers in Ecology and the Environment*, *5*, 365–374.

Johansen, L., Westin, A., Wehn, S., Iuga, A., Ivascu, C. M., Kallioniemi, E., & Lennartsson, T. (2019). Traditional semi-natural grassland management with heterogeneous mowing times enhances flower resources for pollinators in agricultural landscapes. *Global Ecology and Conservation*, *18*, e00619. https://doi.org/10.1016/j.gecco.2019.e00619

Kahmen, S., Poschlod, P., & Schreiber, K. F. (2002). Conservation management of calcareous grasslands. Changes in plant species composition and response of functional traits during 25 years. *Biological Conservation*, *104*(3), 319–328. https://doi.org/10.1016/S0006-3207(01)00197-5

Kariyeva, J., van Leeuwen, W. J. D., & Woodhouse, C. A. (2012). Impacts of climate gradients on the vegetation phenology of major land use types in Central Asia (1981-2008). *Frontiers of Earth Science*, *6*(2), 206–225. https://doi.org/10.1007/s11707-012-0315-1

Korell, L., Auge, H., Chase, J. M., Harpole, S., & Knight, T. M. (2020). We need more realistic
climate change experiments for understanding ecosystems of the future. *Global Change*
Biology, *26*(2), 325–327. https://doi.org/10.1111/gcb.14797

Kuss, P., Rees, M., Ægisdóttir, H. H., Ellner, S. P., & Stöcklin, J. (2008). Evolutionary
demography of long-lived monocarpic perennials: A time-lagged integral projection model.
Journal of Ecology, *96*(4), 821–832. https://doi.org/10.1111/j.1365-2745.2008.01374.x

Lemmer, J., Andrzejak, M., Compagnoni, A., Knight, T. M., & Korell, L. (2021). Climate
change and grassland management interactively influence the population dynamics of
Bromus erectus (Poaceae). *Basic and Applied Ecology*, *56*, 226–238.
https://doi.org/10.1016/j.baae.2021.06.012

Levin, S. C., Childs, D. Z., Compagnoni, A., Evers, S., Knight, T. M., & Salguero-Gómez, R.
(2021). ipmr: Flexible implementation of Integral Projection Models in R. *Methods in*
Ecology and Evolution, *12*(10), 1826–1834. https://doi.org/10.1111/2041-210X.13683

Lyu, S., & Alexander, J. M. (2022). Competition contributes to both warm and cool range edges.
Nature Communications, *13*(1), 1–9. https://doi.org/10.1038/s41467-022-30013-3

McMahon, S. M., Arellano, G., & Davies, S. J. (2019). The importance and challenges of
detecting changes in forest mortality rates. *Ecosphere*, *10*(2).
https://doi.org/10.1002/ecs2.2615

Morris, W. F., Ehrlén, J., Dahlgren, J. P., Loomis, A. K., & Louthan, A. M. (2020). *Biotic and*
anthropogenic forces rival climatic/abiotic factors in determining global plant population
growth and fitness. 117(2), 1107–1112. https://doi.org/10.5061/dryad.95x69p8fm

Moser, B., Fridley, J. D., Askew, A. P., & Grime, J. P. (2011). Simulated migration in a long-
term climate change experiment: Invasions impeded by dispersal limitation, not biotic
resistance. *Journal of Ecology*, *99*(5), 1229–1236. https://doi.org/10.1111/j.1365-
2745.2011.01841.x

Nakahama, N., Uchida, K., Ushimaru, A., & Isagi, Y. (2016). Timing of mowing influences
genetic diversity and reproductive success in endangered semi-natural grassland plants.
Agriculture, Ecosystems and Environment, *221*, 20–27.
https://doi.org/10.1016/j.agee.2016.01.029

Orlowsky, B., & Seneviratne, S. I. (2012). Global changes in extreme events: Regional and
seasonal dimension. *Climatic Change*, *110*(3–4), 669–696. https://doi.org/10.1007/s10584-
011-0122-9

Poniatowski, D., Hertenstein, F., Raude, N., Gottbehüt, K., Nickel, H., & Fartmann, T. (2018).
The invasion of Bromus erectus alters species diversity of vascular plants and leafhoppers
in calcareous grasslands. *Insect Conservation and Diversity*, *11*(6), 578–586.
https://doi.org/10.1111/icad.12302

Poschlod, P., Kiefer, S., Tränkle, U., Fischer, S., & Bonn, S. (1998). Plant species richness in
calcareous grasslands as affected by dispersability in space and time. *Applied Vegetation*
Science, *1*(1), 75–91. https://doi.org/10.2307/1479087

R Core Team. (2022). *R: A language and environment for statistical computing. R Foundation*
for Statistical Computing. https://www.r-project.org/

Ramula, S., Knight, T. M., Burns, J. H., & Buckley, Y. M. (2008). General guidelines for
invasive plant management based on comparative demography of invasive and native plant
populations. *Journal of Applied Ecology*, *45*(4), 1124–1133. https://doi.org/10.1111/j.1365-
2664.2008.01502.x

Redlich, S., Zhang, J., Benjamin, C., Dhillon, M. S., Englmeier, J., Ewald, J., Fricke, U., Ganuza,
C., Haensel, M., Hovestadt, T., Kollmann, J., Koellner, T., Kübert-Flock, C., Kunstmann,
H., Menzel, A., Moning, C., Peters, W., Riebl, R., Rummler, T., … Steffan-Dewenter, I.
(2022). Disentangling effects of climate and land use on biodiversity and ecosystem
services—A multi-scale experimental design. *Methods in Ecology and Evolution*, *13*(2),
514–527. https://doi.org/10.1111/2041-210X.13759

Rees, M., & Ellner, S. P. (2009). Integral projection models for populations in temporally
varying environments. In *Ecological Monographs* (Vol. 79, Issue 4).

Reisch, C., & Poschlod, P. (2009). Land use affects flowering time: Seasonal and genetic
differentiation in the grassland plant scabiosa columbaria. *Evolutionary Ecology*, *23*(5),
753–764. https://doi.org/10.1007/s10682-008-9270-4

Rockel, B., Will, A., & Hense, A. (2008). The regional climate model COSMO-CLM (CCLM).
Meteorologische Zeitschrift, *17*(4), 347–348. https://doi.org/10.1127/0941-2948/2008/0309

Schädler, M., Buscot, F., Klotz, S., Reitz, T., Durka, W., Bumberger, J., Merbach, I., Michalski,
S. G., Kirsch, K., Remmler, P., Schulz, E., & Auge, H. (2019). Investigating the
consequences of climate change under different land-use regimes: a novel experimental
infrastructure. *Ecosphere*, *10*(3). https://doi.org/10.1002/ecs2.2635

Schauberger, P., & Walker, A. (2021). *Openxlsx: Read, Write and Edit xlsx Files.* (4.2.5).

Schuldt, B., Buras, A., Arend, M., Vitasse, Y., Beierkuhnlein, C., Damm, A., Gharun, M.,
Grams, T. E. E., Hauck, M., Hajek, P., Hartmann, H., Hiltbrunner, E., Hoch, G., Holloway-
Phillips, M., Körner, C., Larysch, E., Lübbe, T., Nelson, D. B., Rammig, A., … Kahmen, A.
(2020). A first assessment of the impact of the extreme 2018 summer drought on Central
European forests. *Basic and Applied Ecology*, *45*, 86–103.
https://doi.org/10.1016/j.baae.2020.04.003

Sebald, O., Seybold, S., & Philippi, G. (1990). *Die Farn- und Blütenpflanzen Baden-*
Württembergs (Vol. 1). E. Ulmer.

Tälle, M., Fogelfors, H., Westerberg, L., & Milberg, P. (2015). The conservation benefit of
mowing vs grazing for management of species-rich grasslands: A multi-site, multi-year

field experiment. *Nordic Journal of Botany*, *33*(6), 761–768.
https://doi.org/10.1111/njb.00966

Tebaldi, C., Hayhoe, K., Arblaster, J. M., & Meehl, G. A. (2006). Going to the extremes: An
intercomparison of model-simulated historical and future changes in extreme events. In
Climatic Change (Vol. 79, Issues 3–4, pp. 185–211). Springer Netherlands.
https://doi.org/10.1007/s10584-006-9051-4

Töpper, J. P., Meineri, E., Olsen, S. L., Rydgren, K., Skarpaas, O., & Vandvik, V. (2018). The
devil is in the detail: Nonadditive and context-dependent plant population responses to
increasing temperature and precipitation. *Global Change Biology*, *24*(10), 4657–4666.
https://doi.org/10.1111/gcb.14336

van Daalen, S. F., & Caswell, H. (2020). Variance as a life history outcome: Sensitivity analysis
of the contributions of stochasticity and heterogeneity. *Ecological Modelling*, *417*.
https://doi.org/10.1016/j.ecolmodel.2019.108856

van der Meer, S., Dahlgren, J. P., Mildén, M., & Ehrlén, J. (2014). Differential effects of
abandonment on the demography of the grassland perennial Succisa pratensis. *Population
Ecology*, *56*(1), 151–160. https://doi.org/10.1007/s10144-013-0400-7

Wagner, S., Berg, P., Schädler, G., & Kunstmann, H. (2013). High resolution regional climate
model simulations for Germany: Part II-projected climate changes. *Climate Dynamics*,
40(1–2), 415–427. https://doi.org/10.1007/s00382-012-1510-1

Weppler, T., Stoll, P., & Stöcklin, J. (2006). The relative importance of sexual and clonal
reproduction for population growth in the long-lived alpine plant Geum reptans. *Journal of
Ecology*, *94*(4), 869–879. https://doi.org/10.1111/j.1365-2745.2006.01134.x

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag.

Wickham, H., François, R., Henry, L., & Müller, K. (2022). *dplyr: A Grammar of Data
Manipulation* (1.0.9).

Williams, A. L., Wills, K. E., Janes, J. K., vander Schoor, J. K., Newton, P. C. D., & Hovenden,
M. J. (2007). Warming and free-air CO_2 enrichment alter demographics in four co-
occurring grassland species. *New Phytologist*, *176*(2), 365–374.
https://doi.org/10.1111/j.1469-8137.2007.02170.x

Yule, K. M., Miller, T. E. X., & Rudgers, J. A. (2013). Costs, benefits, and loss of vertically
transmitted symbionts affect host population dynamics. *Oikos*, *122*(10), 1512–1520.
https://doi.org/10.1111/j.1600-0706.2012.00229.

756 **Tables**

757 **Table 1** Description and ecology of each plant species in the study as extracted from the

758 following sources: https://wiki.ufz.de/biolflor/index.jsp* and Ellenberg et al., 1991**.

Species name	Growth form	Plant family	Life history*	Ellenberg**	Start of flowering	End of flowering	Duration of flowering	Unit of measure	Grazing resistance*	Mowing tolerance*
Anthoxanthum odoratum L.	Grass	Poaceae	p[1]	Ind[4]	April	April	1	Basal	5[10]	7[15]
Bromus erectus Huds.	Grass	Poaceae	p	3[5]	May	June	2	Basal	4[11]	5[16]
Crepis biennis L.	Forb	Asterace	b[2]	6[6]	June	August	3	Number	2[12]	6[17]
Dianthus carthusianorum L.	Forb	Caryoph	p	3	May	October	6	Basal	4	3[18]
Lotus corniculatus L.	Legume	Fabacea	p	4[7]	June	July	2	Basal	4	6
Lychnis flos-cuculi L.	Forb	Caryoph	p	7[8]	NA	NA	NA	Basal	2	4[19]
Medicago falcata (L.) Arcang	Legume	Fabacea	p	3	June	July	2	Basal	2	5
Plantago lanceolata L	Forb	Plantagi	p	ind	May	July	3	Number	6[13]	7
Scabiosa ochroleuca L.	Forb	Dipsacac	p	3	June	October	5	Basal	NA[14]	NA
Tragopogon orientalis L.	Forb	Asterace	h[3]	5[9]	June	August	3	Number	NA	NA
Trifolium pratense L.	Legume	Fabacea	p	5	May	May	1	Basal	4	7

[1]perennial and iteroparous, [2]biennial, [3]pluriennial-hapaxanthic, [4]Indifferent, [5]species is more

common at dry sites, avoids wet habitats, [6]prefers semi wet to wet locations, [7]usually more on

dry habitats, able to grow on wet sites, [8]prefers wet locations, without waterlogging, [9]prefers

intermediate locations, [10]moderately tolerant to grazing, [11]sensitive/moderately tolerant to

grazing, [12]intolerant/sensitive to grazing, [13]moderately tolerant/well tolerant to grazing, [14]no

information available, [15]well tolerant to mowing, [16]moderately tolerant to mowing, [17]moderately

to well tolerant to mowing, [18]sensitive to mowing, [19]sensitive to moderately tolerant to mowing,

Table 1. All vital rates and parameters that where used and extracted to calculate the integral

projection model (IPM)

Vital rates	Abbreviation	Life-cycle stage	Parameters	Distribution	LTRE stage
Plant survival	S	Survival	Intercept (Int), Slope	Bernoulli	Survival
Plant growth	G	Growth	Intercept (Int), Slope, SD	Normal	Growth
Reproduction probability	P	Reproduction	Intercept (Int), Slope	Bernoulli	Reproduction
Number of seeds per reproductive plant	F	Reproduction	Intercept (Int), Slope	Poisson	Reproduction
Fall recruitment	θf	Recruitment	Mean		Recruitment
Spring recruitment	θs	Recruitment	Mean		Recruitment
Seedling establishment	B	Establishment	Mean		Establishment
Size distribution of new plants	η	Establishment	Mean, SD		Establishment

Figure captions

Figure 1 Life cycle diagram of the study species. The diagram shows every vital rate

incorporated in the integral projection model (IPM) and the abbreviation used.

Figure 2 Log population growth rate (λ) for each of the five study species (a-e) considering the

two different land management types (grazing and mowing). Displayed is the mean $\pm$ one

standard deviation.

Figure 3 Relationship between the effect size of management on population growth rate (λ

grazing – λ mowing) plotted against a) The mean duration of flowering for each species and b)

the mean beginning

Figure 4 Bootstrapped logged λ plotted against the two climate treatments (ambient and future)

for the five different study species (a-e). Displayed is the mean logged λ of the 1000 repetitions,

with the standard deviation.

Figure 5 Effect of climate (ambient and future) and land management type (grazing and

mowing) on population growth rate (λ) of the five study species (a-e). Displayed is the mean and

the standard deviation.

Figure 6 Sensitivity of pairwise treatment combinations for every vital rate included in the IPM.

Each species is color-coded.

Figure 7 Results for the Life Table Response Experiment (LTRE). Each species is displayed in a

different color. a) Ambient climate treatment and the effect of the two land management types on

the parameters. b) Future climate treatment and the effect of the two land management types on

the parameters. c) Grazing land management type and the effect of climate on the parameters. d)

Mowing land management type and the effect of climate on the life stages. For a better overview,

we summed the parameters according to their corresponding live stage.

Figures

Figure 1

Spring seedling discrete

P, reproduction probability
F, Seeds per reproductive plant
θs, Spring recruitment

B, Seedling survival
η, Size distribution of new plants

Plant continuous

P, Reproduction probability
F, Seeds per reproductive plant
θf, Fall recruitment
B, Seedling survival
η, Size distribution of new plants

S, Plant survival
G, Plant growth

819   Figure 2

822   Figure 3

825   Figure 4

828 Figure 5

833 Figure 6

839   Figure 7