MitoSort: robust demultiplexing of pooled single-cell genomics data using

endogenous germline mitochondrial variants

Zhongjie Tang[1, #], Weixing Zhang[1, #], Peiyu Shi[1, #], Sijun Li[1], Xinhui Li[1], Yicong Xu[1], Yaqing

Shu[2], Jin Xu[1, *]

[1]State Key Laboratory of Biocontrol, School of Life Sciences, Sun Yat-Sen University,

Guangzhou, China

[2]Department of Neurology, The Third Affiliated Hospital of Sun Yat-sen University,

Guangzhou, China

[#] These authors contributed equally to the work.

[*]Correspondence should be addressed to:

Jin Xu, PhD. School of Life Sciences, Sun Yat-Sen University, No. 135 Xingang Xi

Road, Guangzhou, Guangdong, China. Email: xujin7@mail.sysu.edu.cn.

Abstract

Multiplexing across donors has emerged as a popular strategy to increase throughput,

reduce costs, overcome technical batch effect and improve doublet detection in single-

cell genomic studies. Using endogenous genetic barcodes eliminates the need for

additional experimental processing steps. The unique features of mtDNA make mtDNA

variants a more computationally efficient and robust choice for endogenous barcodes

compared to genome variants. Here we developed MitoSort, a method that uses mtDNA

germline variants to assign cells to their donor of origin and identify cross-genotype

doublets robustly and efficiently. Our study validated MitoSort's performance by both

in silico data and cell hashing data and demonstrated its use in various single-cell

sequencing approaches.

Main

The advent of single-cell sequencing techniques has expanded the repertoire of

detectable modalities within a single cell(1, 2). For instance, DOGMA-seq(3) enables

the simultaneous measurement of gene expression, chromatin accessibility, and protein

levels in individual cells, providing a powerful tool for the comprehensive exploration

of biological processes. However, the high costs of multimodal single-cell techniques

limit their application in large cohort studies. To meet the growing demand for profiling

large-scale samples, sample multiplexing has become a popular strategy that not only

reduces costs but also overcomes technical batch effects and improves doublet

detection(4). To avoid additional experimental processing steps, several computational

tools have been proposed to harness genome variants for demultiplexing cells from

different individuals and identifying cross-genotype doublets(5–9). This process

typically requires significant computational resources, even with reference genotypes,

due to the large size of the human genome. In contrast, mtDNA variants offer a more

computationally efficient and robust option, given their hundreds of copies per cell and

the small size of the mtDNA genome(10, 11). Moreover, mtDNA variants can be

captured with high coverage at single-cell resolution in multimodal single-cell

sequencing techniques such as mtscATAC-seq(12), ASAP-seq(3), DOGMA-seq(3), as

well as full-length single-cell RNA-seq (e.g., Smart-seq(13)). Despite the benefits of

using mtDNA variants for demultiplexing samples and identifying doublets, few

software options are currently available. To fill the gap, we developed MitoSort, a

software that exploits mtDNA germline variants exclusively to demultiplex samples

from different individuals and detect cross-genotype doublets efficiently and accurately

(Fig.1a).

The MitoSort pipeline is mainly composed of four steps: MT reads realignment, variant

calling, doublet identification and demultiplexing (Fig.1b). Briefly, to reduce false-

positive variants resulting from STAR(14) alignments artifacts, MT reads are realigned

with GATK(15) to facilitate accurate variant calling. VarScan(16) is then used to

identify putative single-nucleotide polymorphisms (SNPs) in mtDNA. An SNP-by-cell

matrix was generated after filtering low-frequency variants. By the initial K-means

clustering, sample-specific germline variants are identified, allowing for the precise

clustering of cells based on unique mitochondrial haplotypes. To detect doublets, we

model a cell's allele counts as being drawn from a beta-binomial distribution, with

parameters derived from one or two clusters. After removing doublets, cells are

assigned to their donor of origin by K-means clustering again. Additionally, our model

estimates the most likely number of pooled individuals, which is helpful when clinical

information is lost (Methods). It only takes 10 minutes for demultiplexing a mixture of

~10000 cells by MitoSort (the step of MT reads realignment was not involved in the

estimated runtime).

We first used *in silico* data with known cell origins to evaluate our method. To create

synthetic datasets, we collected eight published mtscATAC-seq datasets from

genetically distinct donors (detailed information is shown in Extended Data Table1).

The impact of factors affecting the classification performance of MitoSort was assessed.

Among these factors, the number of pooled samples, the number of cells for each

sample, and the percentage of doublets had little effect on the classification

performance of MitoSort (Fig.1c-e, Extended Data Fig.1a-f). We consistently obtained

high accuracy of origin assignment and doublet identification for 2- to 8-mixed samples

(500 cells per sample, 2000 mtDNA reads per cell, and 8% of cells as doublets) (Fig.1c).

Besides, when we synthetically mixed 8 samples (2000 mtDNA reads per cell and 8%

of cells as doublets) and varied the number of cells from 100 to 600 per sample, we

found the true positive rate (TPR) of both donor assignment and doublet identification

was consistently achieved at a rate of 1, even when the number of cells was as low as

100 (Extended Data Fig.1e). Only a small number of singlets were erroneously

classified as doublets, constituting less than 0.3% of all cases (Extended Data Fig.1f).

In addition, MitoSort achieved perfect performance regardless of the percentage of

simulated doublets (Fig.1e Extended Data Fig.1g-h). We found that the classification

performance of MitoSort was only sensitive to the sequencing depth of mitochondrial

genome (Fig.1f). Specifically, an increase in the number of mapped reads (ranging from

250 to 2000) resulted in a higher true positive rate (TPR) and a lower false discovery

rate (FDR), leading to improved accuracy (Fig.1f, Extended Data Fig.1g-h). It is worth

noting that when the number of reads mapped to the mitochondrial genome reached

approximately 1000 (roughly four-fold coverage), MitoSort was able to achieve perfect

cell assignment to their respective donors and accurately detect cross-genotype doublets.

Collectively, these data show that MitoSort is a highly accurate and scalable method

for demultiplexing samples from different individuals and identifying cross-genotype

doublets using mtDNA germline variants.

We next evaluated our method by two published datasets that used cell-hashing

technology to mark sample origins of cells prior to multiplexing. The DOGMA-seq

dataset consisted of activated and stimulated T cells from two healthy donors(17), while

the ASAP-seq dataset comprised NK cells from four donors(18). MitoSort was able to

assign cells to their respective donors accurately in both datasets, demonstrating high

consistency with the known cell hashing tags (Fig. 2a-b). MitoSort achieved high

accuracy of origin assignment (>= 98%) when the sequencing depth of mitochondrial

genome was not less than 4X (Extended Data Fig.2). These results further suggest the

robustness of MitoSort in assigning cells from different donors using mtDNA germline

variants.

Compared to droplet-based single-cell RNA sequencing techniques that typically

sequence the 3' ends of RNAs, full-length single-cell RNA-seq protocols offer

improved coverage of mitochondrial transcripts and facilitate variant calling for larger

parts of the mtDNA genome (Extended Data Fig.3a). Consequently, it is feasible to

demultiplex full-length single-cell RNA-seq data using MitoSort. We evaluated the

performance of MitoSort using the Smart-seq2 dataset comprising human cells from

four donors and the HEK293T cell line(19). We also benchmarked our method against

souporcell(8), one of the best computational tools for demultiplexing pooled single-cell

RNA-seq data using genome variants. The classification results obtained by MitoSort

were highly consistent with those produced by souporcell (Fig.2c). Cells that were

assigned as to the same donors shared the same mitochondrial haplotype (Extended

Data Fig.3b). Furthermore, MitoSort was faster than souporcell in processing the same

dataset (Fig.2d). These results also demonstrate that MitoSort can accurately

demultiplex pooled datasets and applied to variant single-cell genomics libraries,

provided that the mitochondrial sequence can be readout.

In the end, we conducted a case study in which we multiplexed B cells from four normal

individuals and four patients with auto-immune diseases to profile gene expression,

chromatin accessibility, and mitochondrial DNA (Figure 2e). The scATAC-seq library

contained an average of 25% mtDNA reads per cell, which allowed accurate mtDNA

variant calling and demultiplexing (Extended Data Fig.4a). Using MitoSort, we could

clearly classify 16472 cells into eight singlet populations, as well as doublet groups

based on the allele frequency of unique mtDNA germline variants for each individual

(Figure 2f). Doublet is another important issue to resolve in multiplexing datasets,

which become increasingly prevalent with the growing number of captured cells.

Traditional methods for doublet detection, which rely on the transcriptional profiles[20],

may not effectively address this issue when the pooled cells lack cell type diversity. In

contrast, mitochondrial variant-based approaches hold promise as an effective strategy

for identifying cross-genotype doublets and mitigating the effects of doublets in

multiplexed datasets, particularly when dealing with a large number of pooled samples.

With MitoSort, we detected 6.4% of cell barcodes as cross-genotype doublets involved

more than one MT genotype (Figure 2f, Extended Data Fig.4b-c). Further quantification

metrics showed an expected higher number of RNA molecules and chromatin

fragments in doublets (Extended Data Fig.4d). Following the removal of low-quality

cells and cross-genotype doublets, we obtained 12943 high-quality B cell singlets.

High-dimensional clustering of multi-omic profiles identified 5 clusters, which could

be annotated as naïve B cells and memory B cells based on known markers (Fig.2g,

Extended Data Fig.4e-f). Using mtDNA SNP information, we can match donors with

MitoSort assigned labels. Among the four patients with auto-immune diseases, two

received treatments with rituximab (RTX). We found that most memory B cells were

depleted in these two patients treated with RTX (Figure 2h-i), which was consistent

with the function of RTX(21). These data show that MitoSort can be used to accurately

demultiplex samples of individuals under different pathologies, which facilitates

downstream analyses without technical batch effect.

In conclusion, the unique properties of mitochondrial DNA (mtDNA) make it a more

efficient and robust choice for endogenous genetic barcodes in sample demultiplexing

and the detection of cross-genotype doublets. However, only a few software options are

currently available for this purpose. MitoSort fills this gap by providing an accurate and

scalable method for demultiplexing pooled single-cell genomics data (mtscATAC-seq,

ASAP-seq, DOGMA-seq, Smart-seq) across donors and identifying cross-genotype

doublets. Nonetheless, the matrilineal inheritance of mitochondria poses a limitation to

this approach, as it is challenging to distinguish individuals with haploidentical mtDNA.

Methods

Realignment for Mitochondrial reads

To correct potential mapping errors around indels, the reads mapped to mitochondrial

genome were realigned locally using GATK version 3.5(15).

Variant calling and germline mutation selection

VarScan2(16) was employed with the parameters "–min-var-freq 0.01," "min-depth 8,"

and "–min-reads2 2" to call variants on the mitochondrial genome using BAM files

containing read pairs realigned to the mitochondrial genome. Since mitochondrial reads

from the BAM file were contributed by multiple individuals, we retained variants with

allele frequencies ranging from 1% to 99% as potential germline variants for each

individual (1% < bulk VAF < 99%).To determine the number of reference alleles and

alternate alleles in potential germline variants for each cell, we developed an in-house

Python script (available at https://github.com/tangzhj/MitoSort) to generate two

matrices, namely, Ref.matrix and Alt.matrix. For each potential germline variant per

cell, the allele frequency was calculated as the ratio of alternate counts to all counts and

stored in the Frequency.matrix. The Frequency.matrix contained the allele frequency

for each potential germline variant, including sample-specific germline variants,

common germline variants, and widely spread somatic variants.

To ensure that only germline variants were included in our analysis, a straightforward

filtering criterion was employed. Specifically, we selected variants where the

proportion of cells having a variant allele frequency (VAF) above 99% among those

with a non-zero VAF was higher than 50% . This criterion was chosen to ensure the

selection of high-confidence germline variants with a high VAF in most cells, while

simultaneously excluding low-coverage germline variants and somatic variants with

low VAF.

Identification of specific germline variants for each individual

To identify sample-specific germline variants while excluding common germline

variants that could potentially interfere with subsequent doublet detection, we utilized

the robust information on germline variants in each sample. First, we performed initial

clustering using the k-means algorithm from the sklearn library, with parameters

n_clusters = pooled_sample_number, max_iter=1000, tol=1e-5, algorithm='full', and

n_init = 10, based on the information obtained from the Frequency.matrix. Within each

cell cluster, variants where the ratio between cells with VAF>99% and cells with

VAF>0 was higher than 0.5 were assigned as a variant set. Subsequently, common

variants that occurred in more than two sets were removed to obtain the final set of

specific germline variants for each individual. The approach we described above

allowed us to identify the cell clusters of each individual, as well as the sample-specific

germline variants for the first time, thereby providing an efficient and robust method

for downstream analysis such as doublet detection.

Doublet detection

Definitions:

K: the number of clusters defined by the above steps.

C: the number of cells. Lower-case c is used for indexing and referring to a specific cell

barcode.

G: the number of sample-specific germline variants sets. Lower-case g is used to

represent all the specific germline variants in a cluster, equal to G_i (*i* means cluster *i*).

g,c means these cluster-specific germline variants with observed data in cell *c*.

A: allele counts. Ag, c is a tuple of size 2 with the first number representing the number

of reference alleles and the second number representing the number of alternate alleles

observed in the cluster-specific germline variants in cell *c*.

We already defined the initial clusters of cells and their corresponding cluster-specific

germline variants. Our next objective was to identify doublets between each cluster.

For cluster-specific germline variants in each single cell, the observation of alternative

allele counts follows a binomial distribution, with *p* being the probability of alternative

allele frequency, $A_{g,c,1}$ being the alternative allele counts and N being the total counts

for cluster-specific germline variants.

Equation (1):

$$P\{X = A_{g,c,1}\} = \binom{N}{A_{g,c,1}} p^{A_{g,c,1}} (1 - p)^{A_{g,c,0}}$$

217 For a given cell, the probability of alternative alleles (*p*) or BAF at sample-specific

218 germline variants can be modeled by a beta prior.

219 The Beta prior model for *p* can be tuned to reflect the relative prior plausibility of each *p*.

220 Equation (2): prior model:

$$f(p) = \frac{1}{B(\alpha, \beta)} p^{\alpha - 1} (1 - p)^{\beta - 1}$$

222   Upon observing data $X=x$ where $x \in \{0, 1,\ldots, n\}$, the likelihood function of p, obtained

223   by plugging x into the Binomial pmf equation (1), we used equation (3):

$$L(p|X = A_{g,c,1}) = \binom{N}{A_{g,c,1}} p^{A_{g,c,1}}(1-p)^{A_{g,c,0}}$$

225   Via Bayes' Rule, the conjugate Beta prior combined with the Binomial data model

226   produce a Beta posterior model for posterior probability of cells belong to each

227   cluster(p). Putting equation (2) and equation (3) together, the posterior pdf is descried

228   by equation (4):

$$f(p|X = A_{g,c,1}) \propto f(p)L(p|X = A_{g,c,1}) \propto p^{(\alpha+A_{g,c,1})-1}(1-p)^{(\beta+A_{g,c,0})-1}$$

230   For each cluster, α and β can be estimated using the allele counts across all the sample-
231   specific germline variants in one cluster.

233    $\alpha = 1 +$ mean of alternate counts across variants in G_i of K_i, and $\beta = 1$. (for cluster).

234   For a cell c belongs to first-fitted cluster i (K_i), and $g=G_i$, the p_i was described by

235   equation (5):

$$f(p_i|X = A_{g,c,1}, A_{g,c,1} + A_{g,c,0}, c \in K_i, g = G_i) \propto B(\alpha + A_{g,c,1}, \beta + A_{g,c,0})$$

237   When the cell c belongs to first-fitted cluster i (K_i) but $g=G_j$, p_j was described by

238   equation (6):

$$f(p_j|X = A_{g,c,1}, A_{g,c,1} + A_{g,c,0}, c \in K_i, g = G_j) \propto B(\alpha + A_{g,c,1}, \beta + A_{g,c,0})$$

Then the P_i was p_1 for the first-fitted cluster and the highest P_j was selected as p_2 for

the second fitted cluster. We estimate p as the mode of the posterior distribution.

When the cell is a singlet, p_1 will be dramatically higher than any P_j.

To give a reasonable threshold, we treated p_1, p_2 as two-dimensional data. We treated

$p_1>0.99$ and $p_2<0.01$ as "Singlet" and treated $p_1<0.8$ or $p_2>0.2$ as "Doublet" at the

beginning, this cutoff can be set by users. To predict the class of unlabeled cells, we

use the 'KNeighborsClassifier' method, which calculates the distance between the

labeled cell point and all other unlabeled cell points in the dataset. The k=5 nearest

neighbors are identified based on the distance metric, and the majority class of the k=5

nearest neighbors is used to predict the class of the unlabeled cells. It should be noted

that if p_1 is greater than 0.99, we recommend not considering p_2, as high values of p_2

may be caused by potential somatic mutations.

Demultiplexing pooled samples

After removing doublets that were identified in the previous step, we further clustered

the remaining singlets using the k-means clustering from the sklearn library with

parameters   n_clusters   =   pooled_sample_number,   max_iter=1000,   tol=1e-5,

algorithm='full', and n_init = 10. The value of n_clusters parameter was set to

correspond to the number of pooled samples. In cases where the exact number of pooled

samples was unknown, we also provided the silhouette method to estimate the optimal

number of clusters. The silhouette score reaches its global maximum at the optimal

number of clusters, allowing for demultiplexing of pooled samples accurately.

Simulation of *in silico* data

In this study, we simulated mixtures of mtscATAC-seq data obtained from the 10x

Genomics platform using custom Python scripts. Following the alignment of reads to

the human reference genome hg38 using 10x Genomics Cell Ranger ATAC(22),

mitochondrial reads were merged into a bam file with cell barcodes prefixed to

distinguish individuals. Next, reads were realigned using GATK(15). In total, we had

52283 cells across 8 samples. To assess the performance of MitoSort systematically,

we generated synthetic mixtures into BAM files by random sampling using a range of

parameter choices, including the number of reads mapped to mitochondrial genome

(ranging from 250 to 2000), the number of pooled samples (ranging from 2 to 8), the

number of cells for each sample (ranging from 100 to 600), and the percentage of

doublets (1%, 5%, 10%, and 15%). Doublets were simulated by randomly choosing two

cell barcodes and merging their read counts as a new cell. For each parameter set,

simulations were repeated five times to account for variability. The MitoSort pipeline,

including variant calling, doublet identification and demultiplexing, was ran on each

simulated BAM file. We evaluated the performance of MitoSort by comparing its

classification results with the true labels, and computed accuracy, true positive rate

(TPR), and false discovery rate (FDR).

Evaluation of MitoSort using cell hashing data

For multiplexed DOGMA-seq dataset comprising activated and stimulated T cells from

two healthy donors(17), we first downloaded its HTO count data (GSM6032896) and

ATAC raw data (SRX14779147). HTO count data containing hashtag reads were used

to assign each cell a hashtag ID by the function HTODemux() from Seurat(23) package.

Raw reads in ATAC data were aligned to the human reference genome (hg38) using

Cell Ranger ATAC(22). Based on the generated BAM file, we ran MitoSort pipeline,

including MT reads realignment, variant calling, doublet identification and

demultiplexing. Cell barcodes classified as singlets by hashtag and MitoSort were

compared to evaluate the consistency of assignment results obtained from both

approaches.

For multiplexed ASAP-seq dataset comprising NK cells from four donors(18), we first

downloaded mitochondrial genotyping data and hashtag-based sample identity

information (GSM6413442). We then used MitoSort to identify doublets and

demultiplex samples based on the mitochondrial genotyping data. Finally, we compared

the donor assignment results obtained from MitoSort with those derived from the

hashtag data.

Analysis of multiplexed Smart-seq2 data

We obtained raw Smart-seq2 data for the Human Cell Atlas (HCA) benchmarking

sample(19) from SRR9621775 and processed it using zUMIs

(https://github.com/sdparekh/zUMIs), which involved reads filtering, mapping to the

human reference genome (hg38), and quantification of gene expression. Based on

species annotation from the original paper, human cells including PBMCs from four

donors and the HEK293T cell line were retained for downstream analysis. Next, we

used MitoSort and soupercell, respectively for demultiplexing human cells. The donor

assignment results and run time of MitoSort and soupercell were compared to evaluate

their performance.

Isolation of PBMCs and B cells

The study protocol was approved by the local ethics committee of Sun Yat-sen

University (#2015/521), and written informed consent was obtained from all donors.

All experiments and protocols for human studies involving human samples were

performed in accordance with the relevant guidelines and regulations. To obtain

peripheral blood mononuclear cells (PBMCs), the peripheral blood was diluted 1:1.5 in

0.9% physiological saline and cells were isolated with a Lymphoprep™ density

gradient (density 1.077 g/mL) according to the manufacturer's instructions

(STEMCELL). B cells were isolated using Miltenyi CD19 MicroBeads (#130-050-301)

as described by the manufacturer. Cells were counted using Trypan Blue exclusion

under a hemocytometer.

Single cell multi-omics library construction and sequencing

Single-cell multi-omics libraries were generated using the 10X Chromium Controller

and the Chromium Next GEM Single Cell Multiome ATAC + Gene Expression kit

according to the manufacturer's instructions. DNA LoBind tubes (1.5 ml; Eppendorf)

were used to wash cells in PBS and in downstream processing steps. To retain mtDNA,

permeabilization was performed using LLL conditions, as described in mtscATAC-

seq(3, 12) with 10 mM Tris-HCl pH 7.4, 10 mM NaCl, 3 mM $MgCl_2$, 0.1% NP40 and

1% BSA. After washing, cells were fixed with 1% formaldehyde (ThermoFisher no.

28906) in PBS for 10 minutes at room temperature, followed by quenching with glycine

solution to a final concentration of 0.125 M. After washing twice in PBS via

centrifugation at 400g, 5 min, 4 °C, cells were treated with lysis buffer (10 mM Tris-

HCl pH 7.4, 10 mM NaCl, 3 mM $MgCl_2$, 0.1% NP40, 1% BSA) for 3 minutes, followed

by adding 1 ml of chilled wash buffer (10 mM Tris-HCl pH 7.4, 10 mM NaCl, 3 mM

$MgCl_2$, 1% BSA) and inversion before centrifugation at 400g, 5 min, 4 °C. The

supernatant was discarded, and cells were diluted in 1× Diluted Nuclei buffer (10X

Genomics) before counting using Trypan Blue and a Countess II FL Automated Cell

Counter. Briefly, after tagmentation, the cells were loaded on a Chromium Controller

Single-Cell instrument to generate single-cell Gel Bead-In-Emulsions (GEMs)

followed by PCR as described in the 10X protocol. The final libraries were quantified

using a Qubit dsDNA HS Assay kit (Invitrogen) and a High Sensitivity DNA chip run

on a Bioanalyzer 2100 system (Agilent). Libraries were sequenced on illumina

NovaSeq 6000 at Berry Genomics.

Analysis of multi-omics data of multiplexed B cells

Raw data of multiome were preprocessed and aligned to the human reference genome

(hg38) using 10x Genomics Cell Ranger arc. The resulting BAM file from ATAC data

was subjected to the MitoSort pipeline for doublet identification and sample

demultiplexing. The cell-by-gene and cell-by-peak count matrices were further

processed using the R package Seurat(23) and Signac(24). High-quality cells with less

than 5000 RNA molecules, mitochondrial RNA content <10%, 200-3500 expressed

genes, 2000-50000 fragments, TSS enrichment score >= 5, nucleosome signal <1, and

the percentage of reads in peaks >= 50 were retained for downstream analysis.

Subsequently, weighted nearest neighbor (WNN) analysis was performed on both RNA

and ATAC-seq modalities, resulting in 10 clusters. Cells were then annotated into four

cell types using the following marker genes: B cells (BANK1, MS4A1); Plasma

(MZB1); T cells (CD3D, CD3E); monocytes (CD14, S100A9). After removing cross-

genotype doublets assigned by MitoSort, we obtained a total of 12943 B cells. Next,

WNN analysis was applied to the subclustering analysis of multiome data of B cells,

which yielded 5 clusters. B cells were further divided into two subtypes using the

following marker genes: naive B cells (TCL1A, IL4R); memory B cells (CD27). Based

on the donor origins of cells obtained from MitoSort, the percentage of each cluster in

each sample was calculated.

Code availability

MitoSort is freely available at https://github.com/tangzhj/MitoSort.

Contributions

J. X. designed and conceived the study. Z.T., P.S. collected and analyzed the data. X.L.,

S.L. and Y.X. supported the data analysis. Y.S. and W.Z. collected samples. W.Z.

generated the single cell multi-omic library. J. X., Z.T. and P.S. wrote the manuscript

with inputs from all authors. All authors read and approved the final manuscript.

Funding

This work was supported by National Key R&D Program of China (2021YFA1102100),

National Natural Science Foundation of China (32070644), Guangdong Basic and

Applied Basic Research Foundation (2019A1515110387, 2019B1515130004) to J.X.

Competing interests

The authors declare no conflict of interest

Fig.1 Workflow and evaluation of MitoSort.

a, Schematic showing the purpose of MitoSort.

b, Overview of MitoSort pipeline. Firstly, mitochondrial reads are realigned and

putative single-nucleotide polymorphisms (SNPs) are called. Subsequently, count

matrices of reference and alternate alleles are generated for each cell, where each row

represents a cell and each column represents a variant. Following that, sample-specific

germline variants are identified for each individual. Cross-genotype doublets are then

detected by a beta-binomial model. Lastly, singlets are demultiplexed for downstream

analyses.

c-f, Systematic assessment of MitoSort performance on simulated data using a range of

parameter choices, five replicate runs are used to ensure the consistency of the results.

c, Classification accuracy of MitoSort on simulated datasets comprising 2-8 mixed

samples, each with 500 cells, an 8% doublet rate, and 2000 MT reads per cell. **d**,

Classification accuracy of MitoSort on simulated datasets containing eight pooled

samples, each with varying numbers of cells, an 8% doublet rate and 2000 MT reads

per cell. **e**, Classification accuracy of MitoSort on simulated datasets containing eight

pooled samples, each with 500 cells, 2000 MT reads per cell and varying doublet rates.

f, Classification accuracy of MitoSort on simulated datasets containing eight pooled

samples, each with 500 cells, an 8% doublet rate, and varying numbers of mitochondrial

(MT) reads per cell.

Fig.2 Application of MitoSort to real multiplexed samples

a, Percentage of barcodes shared between hashtag-based (rows) and MitoSort-based

(columns) assignments in multiplexed DOGMA-seq data consisting of two donors.

b, Percentage of barcodes shared between hashtag-based (rows) and MitoSort-based

(columns) assignments in multiplexed ASAP-seq data consisting of four donors.

c, Percentage of barcodes shared between soupercell-based (rows) and MitoSort-based

(columns) assignments in multiplexed Smart-seq2 data consisting of human cells from

four donors and the HEK293T cell line.

d, Bar plot showing running time of MitoSort and soupercell when processing Smart-seq2 data.

e, Schematic of experimental design. Peripheral blood mononuclear cells (PBMCs) from four normal individuals and four patients with antoimmune diseases were collected and pooled. Next, B cells were enriched to profile gene expression, chromatin accessibility, and mitochondrial DNA.

f, Heatmap showing the allele frequency of aggregated sample-specific mitochondrial germline mutations (rows) across cells (columns) sorted by MitoSort assignment. The number in the parentheses of each row name represents the number of sample-specific mitochondrial germline mutations for this sample.

g, UMAP of weighted-nearest neighbor (WNN) graph for RNA and ATAC-seq modalities showing cell type annotation for each cell.

h, UMAP of weighted-nearest neighbor (WNN) graph for RNA and ATAC-seq modalities showing cells per individual colored by cell number densities. Disease status and clinical information for each individual are shown in subtitles.

i, Bar plot showing the percentage of each cell cluster in each individual.

Extended Data Fig.1 Evaluation of MitoSort using in silico data.

a-b, The true positive rate (TPR) and false discovery rate (FDR) of donor assignment and doublet identification by MitoSort on simulated datasets containing eight pooled samples, each with 500 cells, an 8% doublet rate, and varying numbers of mitochondrial (MT) reads per cell.

c-d, The true positive rate (TPR) and false discovery rate (FDR) of donor assignment and doublet identification by MitoSort on simulated datasets comprising 2-8 mixed samples, each with 500 cells, an 8% doublet rate, and 2000 MT reads per cell.

e-f, The true positive rate (TPR) and false discovery rate (FDR) of donor assignment and doublet identification by MitoSort on simulated datasets containing eight pooled

samples, each with varying numbers of cells, an 8% doublet rate and 2000 MT reads

per cell.

g-h, The true positive rate (TPR) and false discovery rate (FDR) of donor assignment

and doublet identification by MitoSort on simulated datasets containing eight pooled

samples, each with 500 cells, 2000 MT reads per cell and varying doublet rates.

Extended Data Fig.2 Evaluation of MitoSort using cell hashing data.

a, Heatmap showing the allele frequency of sample-specific mitochondrial germline

mutations (columns) across cells (rows) with different sequencing depth of

mitochondrial genome (2-4X, 4-8X and >8X) in multiplexed DOGMA-seq data.

b, Percentage of barcodes with different sequencing depth of mitochondrial genome (2-

4X, 4-8X and >8X) shared between hashtag-based (rows) and MitoSort-based (columns)

assignments in multiplexed DOGMA-seq data.

c, Heatmap showing the allele frequency of sample-specific mitochondrial germline

mutations (columns) across cells (rows) with different sequencing depth of

mitochondrial genome (5-10X, 10-20X and >20X) in multiplexed ASAP-seq data.

d, Percentage of barcodes with different sequencing depth of mitochondrial genome (5-

10X, 10-20X and >20X) shared between hashtag-based (rows) and MitoSort-based

(columns) assignments in multiplexed ASAP-seq data.

Extended Data Fig.3 Application of MitoSort to multiplexed Smart-seq2/3 data

a, Mean sequencing coverage along the mitochondrial genome per cell for the Human

Cell Atlas (HCA) benchmarking data profiled by 10x 3' scRNA-seq and Smart-seq2.

b, Heatmap showing the allele frequency of sample-specific mitochondrial germline

mutations (columns) across cells (rows) in multiplexed Smart-seq2 data comprising

human cells from four donors and the HEK293T cell line.

Extended Data Fig.4 Analysis of multi-omics data of multiplexed B cells

a, Violin plot showing distribution of percentage of mtDNA reads per cell in the scATAC-seq library.

b, tSNE plot of cell barcodes based on mitochondrial germline mutation profiles. Each dot represents a cell barcode and is colored based on the singlet/doublet classification by MitoSort.

c, Bar plot showing proportion of cell barcodes that are classified as singlet and doublet by MitoSort.

d, Violin plots showing the distribution of the number of RNA molecules (top) and ATAC fragments (bottom) per cell in cell barcodes classified as singlet and doublet.

e, UMAP of weighted-nearest neighbor (WNN) graph for RNA and ATAC-seq modalities. Each dot represents a cell colored by its cluster.

f, Dot plot showing expression of marker genes in each cluster. Dot size indicates the proportion of cells expressing the gene and dot color represents the scaled average expression level.

Reference

1. Vandereyken,K., Sifrim,A., Thienpont,B. and Voet,T. (2023) Methods and applications for single-cell and spatial multi-omics. *Nat Rev Genet*, 10.1038/s41576-023-00580-2.

2. Dimitriu,M.A., Lazar-Contes,I., Roszkowski,M. and Mansuy,I.M. (2022) Single-Cell Multiomics Techniques: From Conception to Applications. *Front Cell Dev Biol*, **10**.

3. Mimitou,E.P., Lareau,C.A., Chen,K.Y., Zorzetto-Fernandes,A.L., Hao,Y., Takeshima,Y., Luo,W., Huang,T.S., Yeung,B.Z., Papalexi,E., *et al.* (2021) Scalable, multimodal profiling of chromatin accessibility, gene expression and protein levels in single cells. *Nat Biotechnol*, **39**.

4. Zhang,Y., Xu,S., Wen,Z., Gao,J., Li,S., Weissman,S.M. and Pan,X. (2022) Sample-multiplexing approaches for single-cell sequencing. *Cellular and Molecular Life Sciences*, **79**.

5. Kang,H.M., Subramaniam,M., Targ,S., Nguyen,M., Maliskova,L., McCarthy,E., Wan,E., Wong,S., Byrnes,L., Lanata,C.M., *et al.* (2018) Multiplexed droplet single-cell RNA-sequencing using natural genetic variation. *Nat Biotechnol*, **36**.

6. Xu,J., Falconer,C., Nguyen,Q., Crawford,J., McKinnon,B.D., Mortlock,S., Senabouth,A., Andersen,S., Chiu,H.S., Jiang,L., *et al.* (2019) Genotype-free demultiplexing of pooled single-cell RNA-seq. *Genome Biol*, **20**.

7. Huang,Y., McCarthy,D.J. and Stegle,O. (2019) Vireo: Bayesian demultiplexing of
pooled single-cell RNA-seq data without genotype reference. *Genome Biol*, **20**.
8. Heaton,H., Talman,A.M., Knights,A., Imaz,M., Gaffney,D.J., Durbin,R.,
Hemberg,M. and Lawniczak,M.K.N. (2020) Souporcell: robust clustering of
single-cell RNA-seq data by genotype without reference genotypes. *Nat*
Methods, **17**.
9. Han,S., Kim,K., Park,S., Lee,A.J., Chun,H. and Jung,I. (2022) scAVENGERS: a
genotype-based deconvolution of individuals in multiplexed single-cell ATAC-
seq data without reference genotypes. *NAR Genom Bioinform*, **4**.
10. Penter,L., Gohil,S.H. and Wu,C.J. (2022) Natural Barcodes for Longitudinal
Single Cell Tracking of Leukemic and Immune Cell Dynamics. *Front Immunol*,
12.
11. Kulhankova,L., Montiel González,D., Bindels,E., Kling,D., Kayser,M. and
Mulugeta,E. (2023) Single-cell transcriptome sequencing allows genetic
separation, characterization and identification of individuals in multi-person
biological mixtures. *Commun Biol*, **6**.
12. Lareau,C.A., Ludwig,L.S., Muus,C., Gohil,S.H., Zhao,T., Chiang,Z., Pelka,K.,
Verboon,J.M., Luo,W., Christian,E., *et al.* (2021) Massively parallel single-cell
mitochondrial DNA genotyping and chromatin profiling. *Nat Biotechnol*, **39**.
13. Ramsköld,D., Luo,S., Wang,Y.C., Li,R., Deng,Q., Faridani,O.R., Daniels,G.A.,
Khrebtukova,I., Loring,J.F., Laurent,L.C., *et al.* (2012) Full-length mRNA-Seq
from single-cell levels of RNA and individual circulating tumor cells. *Nat*
Biotechnol, **30**.
14. Dobin,A., Davis,C.A., Schlesinger,F., Drenkow,J., Zaleski,C., Jha,S., Batut,P.,
Chaisson,M. and Gingeras,T.R. (2013) STAR: Ultrafast universal RNA-seq
aligner. *Bioinformatics*, **29**.
15. McKenna,A., Hanna,M., Banks,E., Sivachenko,A., Cibulskis,K., Kernytsky,A.,
Garimella,K., Altshuler,D., Gabriel,S., Daly,M., *et al.* (2010) The genome
analysis toolkit: A MapReduce framework for analyzing next-generation DNA
sequencing data. *Genome Res*, **20**.
16. Koboldt,D.C., Zhang,Q., Larson,D.E., Shen,D., McLellan,M.D., Lin,L.,
Miller,C.A., Mardis,E.R., Ding,L. and Wilson,R.K. (2012) VarScan 2: Somatic
mutation and copy number alteration discovery in cancer by exome sequencing.
Genome Res, **22**.
17. Xu,Z., Heidrich-O'Hare,E., Chen,W. and Duerr,R.H. (2022) Comprehensive
benchmarking of CITE-seq versus DOGMA-seq single cell multimodal omics.
Genome Biol, **23**.
18. Rückert,T., Lareau,C.A., Mashreghi,M.F., Ludwig,L.S. and Romagnani,C. (2022)
Clonal expansion and epigenetic inheritance of long-lasting NK cell memory.
Nat Immunol, **23**.

19. Mereu,E., Lafzi,A., Moutinho,C., Ziegenhain,C., McCarthy,D.J., Álvarez-
Varela,A., Batlle,E., Sagar, Grün,D., Lau,J.K., *et al.* (2020) Benchmarking
single-cell RNA-sequencing protocols for cell atlas projects. *Nat Biotechnol*, **38**.

20. Xi,N.M. and Li,J.J. (2021) Benchmarking Computational Doublet-Detection
Methods for Single-Cell RNA Sequencing Data. *Cell Syst*, **12**.

21. Kneitz,C., Wilhelm,M. and Tony,H.P. (2002) Effective B cell depletion with
rituximab in the treatment of autoimmune diseases. *Immunobiology*, **206**.

22. Satpathy,A.T., Granja,J.M., Yost,K.E., Qi,Y., Meschi,F., McDermott,G.P.,
Olsen,B.N., Mumbach,M.R., Pierce,S.E., Corces,M.R., *et al.* (2019) Massively
parallel single-cell chromatin landscapes of human immune cell development
and intratumoral T cell exhaustion. *Nat Biotechnol*, **37**.

23. Hao,Y., Hao,S., Andersen-Nissen,E., Mauck,W.M., Zheng,S., Butler,A., Lee,M.J.,
Wilk,A.J., Darby,C., Zager,M., *et al.* (2021) Integrated analysis of multimodal
single-cell data. *Cell*, **184**.

24. Stuart,T., Srivastava,A., Madad,S., Lareau,C.A. and Satija,R. (2021) Single-cell
chromatin state analysis with Signac. *Nat Methods*, **18**.

Fig.1

Fig.2