Genetic evidence of a functional linkage between the RNA-dependent RNA polymerase

and the highly structured S fragment located at the 5′ end of the genome of foot-and-

mouth disease virus.

Joseph C. Ward[a], Lidia Lasecka-Dykes[b], Samuel J. Dobson[a], Sarah Gold[b], Natalie J.

Kingston[a], Morgan R. Herod[a], Donald P. King[b], Tobias Tuthill[b], David J. Rowlands[a#]

and Nicola J. Stonehouse[a#]

a School of Molecular and Cellular Biology, Faculty of Biological Sciences and Astbury

Centre for Structural Molecular Biology, University of Leeds, Leeds, United Kingdom.

b The Pirbright Institute, Ash Road, Pirbright, Surrey, GU24 ONF, United Kingdom.

Running Head: The S fragment in FMDV replication

#Address correspondence to Nicola J. Stonehouse, N.J.Stonehouse@leeds.ac.uk

David J. Rowlands, d.j.rowlands@leeds.ac.uk

Key Words: picornavirus, S fragment, SHAPE, replication, 5′ UTR, 3D[pol]

Abstract word count: 223

Text word count: 5731

Abstract

Secondary and tertiary RNA structures play key roles in genome replication of picornaviruses.

Complex, functional structures are particularly abundant in the untranslated regions, where

they are involved in initiation of translation, priming of new strand synthesis and genome

circularisation. The 5′ UTR of foot-and-mouth disease virus (FMDV) is predicted to end in a

c. 360 nucleotide-long stem-loop, termed the short (S) fragment. This structure is highly

conserved and essential for viral replication, but the precise function(s) is unclear. Here, the

validity of earlier structural predictions was strengthened by comparative genomic analyses

that confirmed structure conservation of S fragments from a wide range of field isolates. In

addition, we used selective 2′ hydroxyl acetylation analysed by primer extension (SHAPE) to

experimentally-determine the organisation of the structure of the genome. To examine the role

of S fragment stem-loop structure in virus replication, we introduced a series of deletions to

the distal and proximal regions. These truncations affected genome replication in a size-

dependent and, in some cases, host cell-dependent manner. Furthermore, during passage of

viruses incorporating the largest tolerated deletion from the proximal region of the S fragment

stem-loop, an additional mutation was selected in the viral RNA-dependent RNA polymerase,

$3D^{pol}$, which influenced the function of the enzyme. These data suggest that the S fragment and

$3D^{pol}$ interact in the formation of the FMDV replication complex.

Author summary
46
Foot-and-mouth disease (FMD) is a highly contagious viral disease. It is a constant threat to

the global livestock industry, as it remains endemic in many parts of the world. A deeper

understanding of the molecular mechanisms of FMDV replication could help facilitate the

development of future novel control measures for FMD. However, a number of features of the

viral RNA genome remain poorly understood, including the role of a highly structured region,

termed the S fragment. In this study, we examined the structure of the S fragment and

introduced a series of deletions. These affected the ability to replicate. Furthermore, mutations

which arose when one of the deletions was incorporated into virus indicates that the S fragment

RNA and the RNA-dependent RNA polymerase enzyme interact during viral replication.

Introduction

The *Picornaviridae* is a family of single-stranded positive sense RNA viruses with small genomes approximately 7.5 - 8 kb in length. Typically the genome comprises a major single open reading frame flanked by 5′ and 3′ untranslated regions (UTRs), however, additional small open reading frames have been identified in some cardioviruses and enteroviruses (1, 2). The large 5′ UTRs present in most picornaviruses range from ~800 to ~1300 nucleotides (nts) and have been predicted *in silico* by energy minimisation algorithms to comprise several distinct highly structured domains. Some of these RNA structures are common for all picornaviruses and have been well characterised, for instance an internal ribosome entry site (IRES) responsible for the initiation of translation, which comprises ~450 nts for most picornaviruses, although smaller IRES elements resembling those found in some flaviviruses were also found (3–5). The number and organisation of these structural domains varies amongst picornaviruses. For example, a small stem-loop, the cis-active replicative element (*cre*), which is essential for replication, is found in the 5′ UTR of *Aphthoviruses*, but within the coding region in *Enteroviruses* (6–8). All picornavirus genomes include a structured domain at the 5′ end, although the size, conformation and sequence varies in viruses from different genera. Enterovirus 5′ UTRs terminate in a ~80 nucleotide cloverleaf structure, while in *Aphthoviruses* and *Cardioviruses* the 5′ end forms an extended stem-loop (9–11). The size of this stem-loop varies markedly; 40 nts in *Hepatoviruses* and ~80 nts in *Cardioviruses,* whereas the largest (~360 nts) is present in the *Aphthoviruses* (12). The functions of some of these RNA structures remain unknown.

Here, we focus on the long 5′ terminal stem-loop (termed S fragment) of foot-and-mouth disease virus (FMDV), a pathogen of cloven-hoofed animals, which belongs to the genus *Aphthovirus*. FMDV is the causative agent of foot-and-mouth disease, a highly contagious and

economically important infection posing a constant thread to the global livestock industry. In

endemic countries FMD is controlled by vaccination and by movement restrictions, while

movement restrictions and slaughter were used when outbreaks occurred in non-endemic

countries (13). Control by vaccination is complicated by high antigenic variability and the

occurrence of asymptomatic carrier animals (14–16). FMDV has an 8.5 kb RNA genome

organised into L, P1-2A, P2 and P3 coding regions, flanked by the 5′ UTR described above

and a short 3′ UTR. L is a protease responsible for separating itself from the polyprotein and

cleaving key cellular proteins, P1 encodes the structural (capsid) proteins, while P2 and P3

encode non-structural proteins involved in genome replication. The latter include the RNA-

dependent RNA polymerase (RdRp, also termed $3D^{pol}$) and the protease $3C^{pro}$, as well as less

well-characterised proteins including 2C, a potential helicase (17–19). At approximately 1.3

kb, the 5′ UTR of FMDV is ~$1/7^{th}$ of the entire genome and is the longest 5′ UTR within the

family *Picornaviridae* (10). Its sequence comprises at least five structurally and functionally

distinct domains. The S fragment stem-loop is located at the 5′ terminus and is followed by a

long poly-C-tract of variable length (about 70 to 250 nts), 2-4 tandem repeated sequences

predicted to form pseudoknots, the small stem-loop *cre* involved in uridylation of the

replication primer peptide VPg and finally the IRES element responsible for protein translation

initiation (20). The functions of several of these domains are poorly understood (21, 22),

although we have recently described the importance of a sequence in the pseudoknot region for

virus assembly (23).

Whilst the S fragment is known to be essential for replication, its precise function remains

unclear. The secondary structure of the S fragment appears to be conserved, although its

sequence and length varies between different clades of FMDV (10, 24). Truncated S fragments

have been observed for some field isolates, each time retaining its long stem-loop structure

(10). It has been reported that while distal portions of this stem-loop structure are not essential

for genome replication, they play a role in the modulation of host-cell innate immune responses

(25, 26). A corresponding structure present at the 5′ end of poliovirus (PV) RNA, a highly

structured region called cloverleaf , is involved in the switch from protein translation to RNA

replication, in addition to recruiting host and viral proteins which contribute to RNA

circularisation during genome replication (9, 27). It is possible that the S fragment in FMDV

has similar roles and it is known to interact with the 3′ UTR, suggesting a role in genome

circularisation (28).

We used selective-2′ hydroxyl acylation analysed by primer extension (SHAPE) to provide

direct evidence to complement comparative bioinformatic analyses and *in silico* predictions of

the structure of the S fragment. We confirm that substantial deletions can be made to the distal

region of the S fragment stem-loop without seriously compromising *in vitro* replication,

although host cell differences were observed. We also show that proximal deletions are less

well tolerated, which correlates with our observation that the proximal part of the S fragment

shows higher conservation of nucleotide pairings. Furthermore, viruses reconstructed to

include the maximal viable proximal deletion can, during sequential passages, select a

compensatory mutation in the 3D^{pol}. This mutation is located in a highly conserved position

known to interact with template RNA, and granted faster replication of a mutant replicon

carrying this proximal S fragment deletion.

Material and Methods

Cells lines.

Baby hamster kidney (BHK)-21 and Madin-Darby bovine kidney (MDBK) cells were obtained

from the American Type Culture Collection (ATCC) (LGC Standard) and maintained in

Dulbecco's modified Eagle's Medium with glutamine (Sigma-Aldrich) supplemented with

10% foetal calf serum (FCS), 50 U/ml penicillin and 50 µg/ml streptomycin as previously

described (29).

Plasmid construction.

The FMDV ptGFP replicon plasmids along with the equivalent 3D^{pol} knock-out controls 3D^{pol}

-GNN have already been described (30, 31).

A variant of the ptGFP replicon was produced to allow for easy mutagenesis of the S-fragment.

An *Eag*I site was introduced at each end of the S-fragment by PCR mutagenesis, altering the

wild-type sequence from AAAGGGGGCATTA to AAACGGCCGATTA at the 5′ end and

GCGCCCGCCTTT to GCGCGGCCGTTT at the 3′ end. S-fragment sequences containing

deletions were chemically synthesised (Thermofisher) and inserted into the replicon vector

using complementary *Eag*I sites.

Mutation of residue I189L of 3D^{pol} in the ptGFP replicon was achieved by PCR mutagenesis

resulting in A7023C substitution. All primer sequences are available on request.

Protein purification.

The I189L mutation was introduced into the His-tagged 3D^{pol} expression clone pET28a-3D,

and recombinant protein was expressed and purified as previously described (32, 33).

In vitro transcription.

In vitro transcription reactions were performed as described previously (34, 35). Briefly, 5 µg of replicon plasmid was linearised with *Asc*I (NEB), purified by phenol-chloroform extraction, ethanol precipitated, redissolved in RNase-free water and used in a T7 *in vitro* transcription reaction. Reactions were incubated at 32°C for 4 hours, treated with 1.25 units of RQ1 DNase for 30 minutes at 37°C and RNA recovered using an RNA Clean & Concentrator-25 spin column kit (Zymo Research), following manufacturer's instructions. All transcripts were quantified by NanoDrop 1000 (Thermo Scientific) and RNA integrity assessed by MOPS-formaldehyde gel electrophoresis.

Replication assays.

BHK-21 and MDBK cell based replicon replication assays were performed in 24-well plates with 0.5 µg/cm^2 of RNA using Lipofectin transfection reagent (Life Technologies) as previously described (35). For every experiment, transfection was performed in duplicate and experiments biologically repeated as indicated. Replicon replication was assessed by live cell imaging using an IncuCyte Zoom Dual colour FLR, an automated phase-contrast and fluorescent microscope within a humidifying incubator. At hourly intervals up to a defined end point, images of each well were taken and used to enumerate the ptGFP positive cell count per well, using our established protocols (34, 35). Data are shown at 8 hours post transfection (when maximum replication was observed) on a linear scale.

For ribavirin sensitivity assays, BHK-21 cells were pre-treated with ribavirin (Sigma) for 7 hours prior to transfection.

SHAPE analysis.

RNA transcripts representing the entire 5′ UTR of FMDV (nucleotides 1 – 1581) were

prepared as above. A sample containing 12 pmol of transcribed RNA was heated to 95°C for

2 minutes before cooling on ice. RNA folding buffer (100 mM HEPES, 66 mM $MgCl_2$ and

100 mM NaCl) was added to the RNA and incubated at 37°C for 30 minutes. The folded

RNA was treated with 5 mM *N*-methyl isatoic anhydride (NMIA) or DMSO for 50 minutes at

37°C. The chemically modified RNA was ethanol precipitated and resuspended in 0.5x Tris-

EDTA (TE) buffer.

Hex or FAM fluorescent primers were bound to modified RNA by heating the reaction to 85°C

for 1 minute, 60°C for 10 minutes and 35°C for 10 minutes in a thermocycler. Reverse

transcription was continued using Superscript III (Invitrogen) following manufacturers

protocol with incubation at 52°C for 30 minutes.

Post-extension, cDNA:RNA hybrids were disassociated by incubation with 4M NaOH at 95°C

for 3 minutes before neutralisation with 2M HCl. Extended cDNA was ethanol precipitated

and resuspended in deionized formamide (Thermo Fisher). Sequencing ladders were similarly

produced using 6 pmol of RNA with the inclusion of 10 mM ddCTP in the reverse transcription

mix and using a differentially labelled fluorescent primer (either Hex or FAM). A sequencing

ladder was combined with NMIA or DMSO samples and dispatched on dry ice for capillary

electrophoresis (Dundee DNA seq) (23).

NMIA reactivity was used as constraints for RNA secondary structure prediction of the S

fragment using the Vienna RNA probing package and based on sequence of the replicon used

in this study (36).

Construction of recombinant viruses.

Replicons used here are based on plasmid pT7S3 which encodes a full-length infectious copy
of FMDV O1 Kaufbeuren (O1K) (37). To generate infectious viral genomes the reporter
sequence was removed from replicons by digestion with flanking *Psi*I and *Xma*I restriction
enzymes and replaced with the corresponding fragment from pT7S3 encoding the capsid
proteins. Full length viral RNA was transcribed using a T7 MEGAscript kit (Thermo Fisher
Scientific), DNase treated using TurboDNase (Thermo Fisher Scientific) and purified using a
MEGAclear Transcription Clean-Up kit (Thermo Fisher Scientific). RNA quality and
concentration were determined by denaturing agarose gel electrophoresis and Qubit RNA BR
Assay Kit (Thermo Fisher Scientific).

Virus recovery.

BHK-21 cells were transfected in 25 cm^2 flasks with 8 µg per flask of infectious clone-derived
RNA using TransIT transfection reagent (Mirus) as described previously (35). At full
cytopathic effect (CPE) or 24 hours post-transfection (whichever was earlier) cell lysates were
freeze-thawed and clarified by centrifugation. The clarified lysate (1 ml) was blind passaged
onto naïve BHK-21 cells and this was repeated for five rounds of passaging.

Sequencing of recovered virus.

Recovered viruses at passage 3, 4 and 5, were sequenced with an Illumina Miseq (Illumina)
using a modified version of a previously described PCR-free protocol (38). Total RNA was
extracted from clarified lysates using TRizol reagent (Thermo Fisher Scientific) and residual

genomic DNA was removed using DNA-free DNA removal Kit (Thermo Fisher Scientific)

prior to ethanol precipitation. Purified RNA was used in a reverse transcription reaction as

previously described (38, 39). Following reverse transcription, cDNA was purified and

quantified using a Qubit ds DNA HS Assay kit (Thermo Fisher Scientific) and a cDNA library

prepared using Nextera XT DNA Sample Preparation Kit (Illumina). Sequencing was carried

out on the MiSeq platform using MiSeq Reagent Kit v2 (300 cycles) chemistry (Illumina) and

paired-end sequencing.

FastQ files were quality checked using FastQC with poor quality reads filtered using the Sickle

algorithm (40). Host cell reads were removed using FastQ Screen algorithm and FMDV reads

assembled *de novo* into contigs using IDBA-UD (41). Contigs that matched the FMDV library

(identified using Basic Local Alignment Search Tool (BLAST)) were assembled into

consensus sequences using SeqMan Pro software in the DNA STAR Lasergene 13 package

(DNA STAR) (42). Finally, the filtered fastQ reads were aligned to the *de novo* constructed

consensus sequences using BWA-MEM algorithm incorporated into Burrows-Wheeler Aligner

(BWA), and mutations were visualised using the Integrative Genomics Viewer (IGV) (43–46).

Cell killing assays.

Virus titre was determined by plaque assay on BHK-21 cells as described before (47). BHK-

21 cells were seeded with 3 x10^4 cells/well in 96 well plates and allowed to settle overnight.

Cell monolayers were inoculated with each rescued virus at MOI of 0.01 PFU for 1 hour,

inoculum was removed and 150 µl of fresh GMEM (supplemented with 1% FCS) was added

to each well. Appearance of CPE was monitored every hour by live cell microscopy (contrast

phase) using the IncuCyte S3. CPE was observed as rounding of the cells and progress of

infection was monitored as a drop in confluency compared to the mock treated cells (i.e.,

treatment with uninfected cell lysate). Serial images of cells were analysed using IncuCyte S3

2018B software.

Sym/Sub polymerase activity assays

Sym/Sub RNA with sequence GCAUGGGCCC was synthesised (Sigma Aldrich) and 5′ end

labelled using γ^{32}P UTP (Perking Elmer) and T4 polynucleotide kinase (NEB) following

manufacturer's protocol. Labelled RNA was purified by ethanol precipitation and resuspended

in nuclease free water.

0.5 μM of end labelled RNA was incubated in polymerisation buffer (30 mM MOPS pH 7.0,

33 mM NaCl, 5 mM MgAc) and heated to 95°C for 2 minutes before chilling on ice. 2 μM of

recombinant 3D^{pol} was added to the RNA and incubated at 37°C for 10 minutes to promote

annealing. After annealing, 50 μM of rNTP was added to the reaction, addition of nucleotide

varied depending on experiment. Aliquots were taken periodically, and reactions stopped by

addition of 2x TBE-urea RNA loading dye (Thermo Fisher).

Extension products in loading dye were heated to 70°C for 5 minutes before loading on a 23%

denaturing polyacrylamide gel containing 7 M urea. Samples were electrophoresed until

sufficiently separated. After electrophoresis gels were fixed for 30 minutes in fixative solution

and exposed onto a phosphoscreen (48, 49).

Bioinformatic analysis

Full genome sequences of 118 FMDV isolates, representing all seven serotypes

(Supplementary Table S1), were downloaded from GenBank. The sequences were chosen

(based on the region encoding the VP1 protein) to represent the known genomic diversity of

FMDV across all seven serotypes. VP1 is the most variable part of the FMDV genome and

frequently used to calculate phylogenetic relationship of FMDV isolates.

The RNA structure of the S fragment was predicted as described before using the RNAalifold

program implemented in the ViennaRNA package (36, 50). For the covariance analysis, also

calculated using RNAalifold program, only those isolates were included which contained

complete sequence at the 5′ end of the genome. Data representing covariance were

superimposed onto the schematics of the S fragment RNA structure and visualised using Forna

tool implemented in the ViennaRNA Web Services (51). The sequence logo representing

amino acid conservation within positions 185 – 194 of the $3D^{pol}$ was prepared using a WebLogo

3.7.4 server and using 1123 FMDV $3D^{pol}$ sequences obtained from fmdbase.org (an FMDV

sequence database generated by the FAO World Reference Laboratory for FMD at The

Pirbright Institute) (52, 53).

Results

SHAPE analysis of the S fragment.

Based on computational folding, such as mFOLD algorithms, the S fragment is predicted to

form a single large hairpin stem-loop comprising approximately 360 nucleotides. Here, we use

selective 2′ hydroxyl acylation analysed by primer extension (SHAPE) to further investigate

the structure of this part of the FMDV genome. This approach relies on the formation of 2′-O-

adducts in single stranded and accessible regions of the RNA, which can then be detected by

reverse transcription.

RNA transcripts representing the full FMDV 5′ UTR were folded before incubation with 5 mM

NMIA. The modified RNAs were then used in reverse-transcription reactions containing

fluorescently end-labelled primers. Reverse transcription is terminated when the enzyme

reaches a nucleotide with a NMIA adduct, thus creating a series of cDNA fragments of different

lengths. These cDNA fragments were analysed by capillary gel electrophoresis alongside a

sequencing ladder to identify the sites of termination, indicating the locations of unpaired and

accessible nucleotides. The proximity of the poly-C-tract to the 3′ end of the S fragment limited

potential primer binding sites, resulting in unreliable data for the last 60 nucleotides of the S

fragment and such was excluded from the analysis (shown in grey in Figure 1A). The rest of

the S fragment was well covered and the SHAPE reactivities from 6 independent experiments

were used to complement the mFOLD algorithm analyses to generate a structural prediction of

this region (Figure 1).

In general, the SHAPE mapping data fitted well with the *in silico* predictions with the most

reactive residues coinciding with bulges within the predominately double stranded stem-loop

structure (Figure 1A). The locations of reactive residues can be seen in the NMIA reactivity

graph, where groups of nucleotides within bulges showed high reactivity (Figure 1B). However,

some predicted bulges showed little NMIA reactivity, suggesting steric hindrance possibly due

to higher order tertiary structure. Similarly, some predicted base-paired nucleotides were

NMIA reactive. These mostly occurred at the tops and bottom of bulges and may result from

nucleotides becoming transiently available for modification during 'breathing' (54). The 5′

terminal two nucleotides (uridines at position 1 and 2) were highly reactive and therefore likely

to be non-base paired.

The effects of deletions to the distal or proximal regions of the S fragment stem-loop on

replicon replication.

The SHAPE data and the structural prediction described in Figure 1 were used to design a

series of truncations to the S fragment, and the modified sequences were introduced into a

replicon. The replicon was based on an infectious clone of FMDV O1K in which the P1 region

of the genome is replaced with a ptGFP reporter (55) (Figure 2A). Deletions were designed to

end at either the top or bottom of bulges, helping ensure that the stem-loop hairpin structure

was maintained. To facilitate replacement of the S fragment with modified sequences *Eag*I

restriction sites were introduced at each end of the S fragment sequence. These sites required

minimal modification of the original sequence and showed no predicted change in RNA

structure, and the modified genome sequence is hereafter referred to as WT. Addition of the

*Eag*I site at the 5′ end of the S fragment resulted in a large drop in replication, however when

paired with the 3′ EagI site, restoring base pairing, replicative potential was drastically

improved, although still below that of the original O1K WT sequence (Figure S1). Deletions

of 70, 148, 246 and 295 nucleotides were introduced at the distal end (top) of the S fragment

stem-loop (constructs named D-70, D-148, D-246 and D-295, respectively) and deletions of

48, 97, 195 and 271 nucleotides were made to the proximal end (bottom) of the S fragment

(constructs referred to as P-48, P-97, P-195 and P-271, respectively) (Figure 2B). The truncated

sequences were chemically synthesised and introduced into the WT replicon via the *Eag*I sites,

meaning that the highly conserved 5′ and 3′ regions and essential UU nucleotides at positions

1 and 2 were maintained in all truncation mutants. The eight new replicon clones were

transcribed into RNAs and transfected into BHK-21 or MDBK cells, alongside WT and a $3D^{pol}$-

GNN negative control. The $3D^{pol}$-GNN replicon cannot replicate and so the ptGFP signal

produced represents translation of the input RNA. BHK-21 and MDBK cells are continuous

cell lines known to support replication of FMDV: BHK-21 cells are used for FMDV vaccine

production, whereas MDBK cells originate from a natural host of FMDV. Replication was

monitored by measuring reporter expression using an IncuCyte ZOOM live cell imaging

system, with analysis by our established protocols (performed 8 hours post transfection, when

replication has reached its peak (Figure S2)) (Figure 2).

As described previously and consistent with sequence information from natural virus isolates

(10, 26), the D-70 mutation caused no significant drop in replication in either cell line. However,

the replicon bearing the D-148 deletion, which removes the distal half of the S fragment stem-

loop, showed reduced replication in BHK-21 cells and none in MDBK cells (Figure 2C,D).

Innate immunity is compromised in BHK-21 cells but intact in MDBK cells (56, 57),

suggesting that the distal portion of the S fragment stem-loop plays a role in modulating a

competent host immune response, in agreement with previously published data (25, 58).

Replicons with the largest distal deletions, D-246 and D-295, did not replicate in either cell

line.

While the consequences of deleting sequences from the distal region of the S fragment stem-

loop have been reported (25), the effects of proximal deletions have not been investigated. The

two smallest proximal deletions (P-48 and P-97), showed modest but significant decreases in

replication compared to WT (1.7 and 1.4-fold, respectively) (Figure 2E,F. However, the larger

deletions, P-195 and P-271, completely ablated replication and reporter gene expression was

reduced to a level comparable to the GNN control. Interestingly, there was no significant

difference between the replication of the P-48 and P-97 deletions in either BHK-21 or MDBK

cells (Figure 2). Overall, the proximal part of the S fragment stem-loop appears to be more

sensitive to deletion in a cell type-independent manner, with truncations as small as 48

nucleotides having a significant effect on replication.

Earlier studies indicated that the proximal part of the S fragment stem-loop is more conserved

than the distal part (10). In recent years, advances in high throughput sequencing (HTS) have

led to a great increase in the number of FMDV full genome sequences available on public

domains, including sequences of viruses of SAT serotypes which were previously

underrepresented. This increase in the range of sequences available prompted us to revisit the

subject of the S fragment variability and we carried out a covariance analysis based on 118

FMDV isolates representing all seven serotypes. Figure 3 shows that there is more base-pairing

conservation in the proximal than in the distal parts of the S fragment stem-loop (Figure 3).

The base-pairing conservation in the proximal part of the otherwise very variable S fragment

stem-loop (low nucleotide identity of approximately 47%) suggests evolutionary pressure to

maintain the structure of this portion of the stem-loop and complements our observations from

the replicon experiments (Figure 2) (10, 59).

Viruses with proximal deletions of the S fragment replicate more slowly than WT and

can select for a mutation in 3D^{pol} during serial passage.

We investigated the consequences of proximal deletions to the S fragment stem-loop further

by introducing them into an FMDV O1K infectious clone. This enabled evaluation of

replication in the context of the entire genome and investigation of the possible selection of

compensatory or adaptive mutations during serial passage. Replicons bearing proximal

deletions of the S fragment stem-loop were converted into infectious clones by replacing the

ptGFP reporter sequence with the original O1K structural protein sequence. RNA transcripts

of the truncated S fragment clones and WT were transfected into BHK-21 cells. Supernatants

containing recovered virus were harvested and used to infect naïve BHK-21 cells for 5

continuous passages. Recovery of infectious virus from each construct was assessed by the

ability of 5th passage samples to induce CPE. As expected, CPE was induced and virus

recovered from the WT construct. Infectious virus was also recovered from P-48 and P-97 but

no virus was recovered from either P-195 or P-271 constructs.

The consequences of the P-48 and P-97 S fragment deletions for the dynamics of virus

replication were investigated by assessing the time taken to induce CPE in BHK-21 cell

monolayers after infection with the mutated viruses or WT at a low MOI of 0.01 PFU. The

integrity of the cell monolayers was assessed at hourly intervals over 63 hours using live cell

imaging monitored by Incucyte S3 (Figure 4A). Both P-48 and P-97 viruses developed CPE

more slowly than WT, as anticipated from the earlier replicon experiments (Figure 4A). For

comparison, while a D-148 virus bearing a larger deletion from the distal end of the S fragment

stem-loop replicated at a slower rate than the WT virus, D-148 replicated significantly faster

than viruses carrying deletions at the proximal end of the S fragment (i.e., P-48 and P-97;

Figure 4A); further confirming replicon experiments. Ability of the viruses to form plaques

was also assessed. Both, P-48 and P-97 mutants generated significantly smaller plaques when

comparing to WT and/or D-148 mutant virus (Figure 4B).

Sequencing of recovered viruses revealed one isolate with mutations in 2C, which were not

investigated further. In addition, a single mutation in the $3D^{pol}$ sequence was found in a high

proportion of the viral progeny of one of three P-97 viral replicates. The A7203C change

resulted in I189L mutation in the $3D^{pol}$ protein amino acid sequence. The evolution of this

mutation was investigated further by sequencing viruses from earlier passages. The proportion

of the I189L mutation in the population increased from 40% at passage 3 to 51% at passage 4,

and 61% at passage 5, suggesting it confers a selective advantage in the context of the proximal

S fragment stem-loop deletion and so is sequentially enriched. With four exceptions (two

isolates containing valine and two isolates containing threonine at this position) out of 1123

sequences of $3D^{pol}$ available, all known FMDV isolates have isoleucine at this position (Figure

S3), making the amino acid substitution for leucine novel. . This residue is known to interact

with template RNA (Figure 5A), playing a crucial role during viral replication (60). No such

mutations were observed in the WT or the P-48 mutant viruses after passage.

Effect of the 3D^{pol} I189L compensatory mutation on replicon replication.

The 3D^{pol} I189L mutation was introduced into WT, GNN and P-97 replicons by site-specific mutagenesis and the consequence for replication assessed by reporter gene expression following transfection into BHK-21 cells, as above. Introduction of 3D^{pol} I189L into the WT replicon had no significant effect on replication, however, a small but significant increase in replication was observed when it was introduced into the P-97 replicon (Figure 5B). As expected, I189L was unable to restore the activity of the GNN negative control.

I189L modifies 3D^{pol} activity.

The influence of the 3D^{pol} I189L mutation on polymerase function was investigated using Sym/Sub assay; a highly sensitive method for interrogating polymerase activity. Sym/Sub assays have been used to study PV replication and enable extension of a radiolabelled template to be measured at the single nucleotide level (48). The method uses a 10-nucleotide template oligonucleotide which base-pairs to create a small double stranded region with a four nucleotide 5′ overhang, thus allowing 3′ nucleotide addition. In the presence of rNTPs and recombinantly expressed 3D^{pol}, the addition of single or multiple nucleotides can be assessed over time by separating input template and elongation products by PAGE (Figure 6A). Initially, an assay using ATP only was undertaken to probe the ability of the enzyme to add a single nucleotide. WT and I189L 3D^{pol} proteins behaved similarly in the assay with no significant differences in the rate of addition of the single nucleotide (Figure 6B & C).

3D^{pol} activity was examined further by including all four rNTPs into the Sym/Sub assay. The design of the Sym/Sub oligonucleotide allows for elongation by a maximum of +4 nucleotides

to produce a fully double stranded product (Figure 7A). Both the WT and the I189L 3D^{pol}

enzymes produced the expected +4 product and, in addition, a longer product equivalent to +12

nucleotides was produced by both polymerases (as calculated by Rf value for migration

travelled) (Figure 7B). This is, reminiscent of the larger products seen in recombination assays

reported in the investigation of PV 3D^{pol} (48, 61). Although there was no measurable difference

in rate of addition of the four nucleotides by WT or mutant enzyme, I189L 3D^{pol} produced the

larger +12 product in greater quantity and faster (Figure 7C & D).

The functions of WT and I189L 3D^{pol} enzymes were dissected in finer detail by examining

incremental addition of rNTP combinations to Sym/Sub assay in the order ATP, UTP, GTP

and CTP. The reactions were incubated for 300 seconds (as above) before separation by

electrophoresis. Addition of ATP alone produced a distinct +1 band, as previously described

(Figure 8A). The addition of ATP and UTP together produced a +2 product in addition to lower

intensity bands corresponding to +3 and +4 nucleotides. Inclusion of ATP, UTP and GTP

resulted in a strong band equivalent to +3 product in addition to a minor +4 band. All four

rNTPs produced a +4 band, as expected (Figure 8A). The appearance of the additional bands

in reactions where some rNTPs were missing indicates misincorporation into the growing

strand. We examined the error-rates of WT and I189L 3D^{pol} by measuring the relative amount

of misincorporated +4 product and found no significant difference between the two enzymes

(Figure 8B).

Differences in misincorporation by the WT and I189L mutant polymerase were investigated

further by assessing the effects of ribavirin on replicons bearing either of the two enzymes as

increased misincorporation rates should result in increased sensitivity to the nucleoside

analogue. However, no differences were seen in the sensitivity of the WT, P-97 and P-97 3D^{pol}

I189L replicons in the presence of up to 1 mM ribavirin (Figure 8C). Therefore, it appears that

differences in error rate were unlikely to be responsible for the production of increased amounts

of the larger +12 product by the I189L 3D^{pol} (Figure 7C and Figure 8D).

Discussion

In comparison to other picornaviruses, the 5′ UTR of FMDV is uniquely large and complex,

comprising several distinct structural domains, the functions of most of which are poorly

understood. The first *c.* 360 nucleotides are predicted to form a single, long stem-loop called

the S fragment (22). Although supported by functional studies, to our knowledge the secondary

structure of the S fragment has not been determined biochemically (25). Here, using SHAPE

chemistry applied to a sequence of a well-studied FMDV isolate (O1 Kaufbeuren) (35), we

confirm that the S fragment folds into a single, long stem-loop containing several bulges

formed by unpaired nucleotides. The importance of some of these bulges for viral viability was

determined previously (25).

Despite the high sequence and length variability of the S fragment observed in FMDV field

isolates, most viruses maintain the full-length stem-loop (10). This suggests a strong

evolutionary pressure to maintain the structure. The S fragment is known to interact with viral

and cellular proteins and the 3′ UTR, likely facilitating circulation of the viral genome during

replication (64–66). However, there is evidence that some deletions to the distal part of the S

fragment are tolerated and a number of unrelated field isolates from different FMDV serotypes

(e.g., A, C and O) have been shown to carry deletions in their S fragments. These deletions

arose independently and range in length from few to 76 nucleotides (10, 26, 62).   Unlike

deletions found in other parts of the FMDV genome (e.g., deletions found in the region

encoding 3A), those found in the S fragment do not appear to be host specific, with viruses

carrying deletions within the S fragment being isolated from both cattle and pig (10, 26, 62,

63). A more recent study showed that deletion of 164 nucleotides from the distal part of the S

fragment can produce a viable, although attenuated virus, suggesting a role in evasion of innate

immunity, at least partially explaining the evolutionary trend to maintain the full structure of

the S fragment by field viruses (25). Here, we show that although deletion of 246 nucleotides

or more from the distal region on the S fragment is not tolerated mutants carrying deletions D-

70 and D-148 remain replication-competent.

The proximal part of the S fragment stem-loop shows higher sequence and nucleotide pairing

conservation compared to the distal region (Figure 3). Surprisingly, deletions to the proximal

part of the stem-loop (i.e., up to 97 nucleotides) are also tolerated, although to a lesser extent

than distal deletions, resulting in both attenuated replicons and viruses (cf P-48 and P-97

mutants versus D-148 mutant, Figure 2 and 4) (10). Interestingly, while the replication of D-

148 mutant replicon was more impaired in MDBK cells than in the BHK-21 cells, this cell-

specific difference was not seen for the viable replicons carrying deletions to the proximal part

of the S fragment stem-loop. Since, unlike BHK-21 cells, MDBK cells appear to have a

functional interferon pathway this might suggest that, although required for viral replication,

this proximal part of the S fragment stem-loop does not play a part in the evasion of the innate

immunity (56, 57).

Of three viral isolates carrying the maximum viable proximal deletion to the S fragment (P-

97), two resulted in selective enrichment of a compensatory mutation over serial passages. In

both cases, the mutations showed a trend towards fixation. One isolate developed a number of

mutations in 2C (data not shown), while another developed a single I189L mutation in a highly

conserved site of $3D^{pol}$ known to form hydrophobic contact with a nucleotide base within

template RNA during viral replication (60). Given the essential role of isoleucine 189 in viral

RNA replication, we investigated the effect of the I189L mutation on replication and RNA

strand synthesis. In fact, introduction of I189L into the P-97 replicon enhanced its replication,

suggesting an advantageous effect of this mutation. Although further biochemical analysis of

the I189L $3D^{pol}$ mutant did not show an altered rate of nucleotide incorporation when compared

to the WT polymerase, it did produce more of a 12-nucleotide long product. Such long products

were observed in similar reactions for the $3D^{pol}$ of PV and were the result of a nucleotide mis-

incorporation and template switch (48, 67, 68). While the 12-nucleotide long product most

likely occurred due to nucleotide misincorporation, there is no evidence for altered fidelity of

the I189L mutant. Whether this product occurred as the result of strand slippage, or a template

switch remains to be investigated. Nevertheless, development of the compensatory mutation

over a serial passage in a virus carrying maximal viable proximal deletion to the S fragment

structure, and the advantageous effect of this mutation on replicon replication suggests the

possibility of (direct or indirect) interaction between the S fragment and the viral polymerase

(or its precursor). Further studies are required to verify this interplay.

In conclusion, using biochemical methods, we confirmed previously proposed secondary

structure of the S fragment of FMDV. Despite its higher sequence and nucleotide pairing

conservation, small deletions to the proximal part of the S fragment stem-loop are viable,

although resulting in attenuated replicons and viruses. An advantageous compensatory

mutation within a highly conserved site of $3D^{pol}$ appeared during serial passage of a viral isolate

bearing the largest viable mutation to the proximal part of the S fragment, suggests potential

interplay between the S fragment and the viral polymerase.

Acknowledgements

This work was supported by funding from the Biotechnology and Biological Sciences Research

Council (BBSRC) of the United Kingdom (research grant BB/K003801/1). Additionally, The

Pirbright Institute receives grant-aided support from the BBSRC (projects BBS/E/I/00007035,

BBS/E/I/00007036 and BBS/E/I/00007037) and the UK Department for the Environment,

Food and Rural Affairs (Defra project SE2945).

1.   Lulla V, Dinan AM, Hosmillo M, Chaudhry Y, Sherry L, Irigoyen N, Nayak KM,

Stonehouse NJ, Zilbauer M, Goodfellow I, Firth AE. 2019. An upstream protein-

coding region in enteroviruses modulates virus infection in gut epithelial cells. Nat

Microbiol 4:280–292.

2.   Roos,' RP, Kong W-P, Semler2 BL. 1989. Polyprotein processing of Theiler's murine

encephalomyelitis virus. J Virol 63:5344.

3.   Martínez-Salas E, Francisco-Velilla R, Fernandez-Chamorro J, Lozano G, Diaz-

Toledano R. 2015. Picornavirus IRES elements: RNA structure and host protein

interactions. Virus Res 206:62–73.

4.   Lopez de Quinto S. 2002. IRES-driven translation is stimulated separately by the

FMDV 3'-NCR and poly(A) sequences. Nucleic Acids Res 30:4398–4405.

5.   Chard LS, Kaku Y, Jones B, Nayak A, Belsham GJ. 2006. Functional analyses of RNA

structures shared between the internal ribosome entry sites of hepatitis C virus and the

picornavirus porcine teschovirus 1 Talfan. J Virol 80:1271–1279.

6.   Mason PW, Bezborodova S V, Henry TM. 2002. Identification and characterization of

a cis-acting replication element (cre) adjacent to the internal ribosome entry site of

foot-and-mouth disease virus. J Virol 76:9686–94.

7.   Goodfellow IG, Kerrigan D, Evans DJ. 2003. Structure and function analysis of the

poliovirus cis-acting replication element (CRE). RNA 9:124–137.

8.   Goodfellow I, Chaudhry Y, Richardson A, Meredith J, Almond JW, Barclay W, Evans

DJ. 2000. Identification of a cis-acting replication element within the poliovirus coding

region. J Virol 74:4590–600.

9.   Barton DJ, O'Donnell BJ, Flanegan JB. 2001. 5' cloverleaf in poliovirus RNA is a cis-

acting replication element required for negative-strand synthesis. EMBO J 20:1439–

48.

10.   Carrillo C, Tulman ER, Delhon G, Lu Z, Carreno A, Vagnozzi A, Kutish GF, Rock

586   DL. 2005. Comparative genomics of foot-and-mouth disease virus. J Virol 79:6487–

587   504.

588   11.   Carocci M, Bakkali-Kassimi L. 2012. The encephalomyocarditis virus. Virulence

589   3:351–67.

590   12.   Drexler JF, Corman VM, Lukashev AN, Van Den Brand JMA, Gmyl AP, Brunink S,

591   Rasche A, Seggewi N, Feng H, Leijten LM, Vallo P, Kuiken T, Dotzauer A, Ulrich

592   RG, Lemon SM, Drosten C, Akoua-Koffi CG, Ameneiros RS, Goodman SM,

593   Godlevska E V., Tungaluna GCG, Görföl T, Hassanin A, Koivogui L, Krüger DH,

594   Lakim MB, Lavrenchenko L, Leroy EM, Matthee S, ter Meulen J, Adu-Sarkodie Y,

595   Oppong S, Preiser W, Pereira MJR, Reusken CBEM, Schneider J, Schmidt-Chanasit J,

596   Fichet-Calvet E, Seebens-Hoyer A, Gloza-Rausch F, Setién AA, Tschapka M, Wells

597   K, Wegner T, Wilkinson DA. 2015. Evolutionary origins of hepatitis A virus in small

598   mammals. Proc Natl Acad Sci U S A 112:15190–15195.

599   13.   Knight-Jones TJD, Rushton J. 2013. The economic impacts of foot and mouth disease

600   – What are they, how big are they and where do they occur? Prev Vet Med 112:161–

601   173.

602   14.   Mahapatra M, Parida S. 2018. Foot and mouth disease vaccine strain selection: current

603   approaches and future perspectives. Expert Rev Vaccines 17:577–591.

604   15.   Park J-H. 2013. Requirements for improved vaccines against foot-and-mouth disease

605   epidemics. Clin Exp Vaccine Res 2:8–18.

606   16.   Stenfeldt C, Eschbaumer M, Rekant SI, Pacheco JM, Smoliga GR, Hartwig EJ,

607   Rodriguez LL, Arzt J. 2016. The Foot-and-Mouth Disease Carrier State Divergence in

608   Cattle. J Virol 90:6344–64.

17.   Sweeney TR, Cisnetto V, Bose D, Bailey M, Wilson JR, Zhang X, Belsham GJ, Curry

S. 2010. Foot-and-mouth disease virus 2C is a hexameric AAA+ protein with a

coordinated ATP hydrolysis mechanism. J Biol Chem 285:24347–24359.

18. Teterina NL, Gorbalenya AE, Egger D, Bienz K, Rinaudo MS, Ehrenfeld E. 2006.

Testing the modularity of the N-terminal amphipathic helix conserved in picornavirus

2C proteins and hepatitis C NS5A protein. Virology 344:453–467.

19. Bienz K, Egger D, Troxler M, Pasamontes L. 1990. Structural organization of

poliovirus RNA replication is mediated by viral proteins of the P2 genomic region. J

Virol 64:1156–1163.

20. Black DN, Stephenson P, Rowlands DJ, Brown F. 1979. Sequence and location of the

poly C tract in aphtho- and cardiovirus RNA. Nucleic Acids Res 6:2381–2390.

21. Mellor EJC, Brown F, Harris TJR. 1985. Analysis of the Secondary Structure of the

Poly(C) Tract in Foot-and-Mouth Disease Virus RNAs. J Gen Virol 66:1919–1929.

22. Clarke BE, Brown AL, Currey KM, Newton SE, Rowlands DJ, Carroll AR. 1987.

Potential secondary and tertiary structure in the genomic RNA of foot and mouth

disease virus. Nucleic Acids Res 15:7067–7079.

23. Ward JC, Lasecka-Dykes L, Neil C, Adeyemi OO, Gold S, McLean-Pell N, Wright C,

Herod MR, Kealy D, Warner E, Jackson T, King DP, Tuthill TJ, Rowlands DJ,

Stonehouse NJ. 2022. The RNA pseudoknots in foot-and-mouth disease virus are

dispensable for genome replication, but essential for the production of infectious virus.

PLOS Pathog 18:e1010589.

24. Lasecka-Dykes L, Wright CF, Di Nardo A, Logan G, Mioulet V, Jackson T, Tuthill

TJ, Knowles NJ, King DP. 2018. Full Genome Sequencing Reveals New Southern

African Territories Genotypes Bringing Us Closer to Understanding True Variability

of Foot-and-Mouth Disease Virus in Africa. Viruses 10.

25. Kloc A, Diaz-San Segundo F, Schafer EA, Rai DK, Kenney M, de los Santos T,

Rieder E. 2017. Foot-and-mouth disease virus 5'-terminal S fragment is required for

replication and modulation of the innate immune response in host cells. Virology

512:132–143.

26. Valdazo-González B, Timina A, Scherbakov A, Abdul-Hamid N, Knowles NJ, King

DP. 2013. Multiple introductions of serotype O foot-and-mouth disease viruses into

East Asia in 2010–2011. Vet Res 44:76.

27. Xiang W, Harris K, Alexander L, Wimmer E. 1995. Interaction between the 5'-

terminal cloverleaf and 3AB/3CDpro of poliovirus is essential for RNA replication. J

Virol 69:3658–3667.

28. Serrano P, Pulido MR, Sáiz M, Martínez-Salas E. 2006. The 3' end of the foot-and-

mouth disease virus genome establishes two distinct long-range RNA-RNA

interactions with the 5' end region. J Gen Virol 87:3013–22.

29. Forrest S, Lear Z, Herod MR, Ryan M, Rowlands DJ, Stonehouse NJ. 2014. Inhibition

of the foot-and-mouth disease virus subgenomic replicon by RNA aptamers. J Gen

Virol 95:2649–57.

30. Tulloch F, Pathania U, Luke GA, Nicholson J, Stonehouse NJ, Rowlands DJ, Jackson

T, Tuthill T, Haas J, Lamond AI, Ryan MD. 2014. FMDV replicons encoding green

fluorescent protein are replication competent. J Virol Methods 209C:35–40.

31. Herod MR, Ferrer-Orta C, Loundras E-A, Ward JC, Verdaguer N, Rowlands DJ,

Stonehouse NJ. 2016. Both cis and trans Activities of Foot-and-Mouth Disease Virus

3D Polymerase Are Essential for Viral RNA Replication. J Virol 90:6864–6883.

32. Ferrer-Orta C, Arias A, Perez-Luque R, Escarmís C, Domingo E, Verdaguer N. 2004.

Structure of foot-and-mouth disease virus RNA-dependent RNA polymerase and its

complex with a template-primer RNA. J Biol Chem 279:47212–21.

33. Bentham M, Holmes K, Forrest S, Rowlands DJ, Stonehouse NJ. 2012. Formation of

Higher-Order Foot-and-Mouth Disease Virus 3Dpol Complexes Is Dependent on

Elongation Activity. J Virol 86:2371–2374.

34.   Herod MR, Tulloch F, Loundras E-A, Ward JC, Rowlands DJ, Stonehouse NJ. 2015.

Employing transposon mutagenesis to investigate foot-and-mouth disease virus

replication. J Gen Virol 96:3507–3518.

35.   Herod MR, Gold S, LaseckaDykes L, Wright C, Ward JC, McLean TC, Forrest S,

Jackson T, Tuthill TJ, Rowlands DJ, Stonehouse NJ. 2017. Genetic economy in

picornaviruses: Foot-and-mouth disease virus replication exploits alternative precursor

cleavage pathway. PLOS Pathog 13:e1006666.

36.   Lorenz R, Bernhart SH, Höner zu Siederdissen C, Tafer H, Flamm C, Stadler PF,

Hofacker IL. 2011. ViennaRNA Package 2.0. Algorithms Mol Biol 6:1–14.

37.   King AMQ, Blakemore WE, Ellard FM, Drew J, Stuart DI. 1999. Evidence for the role

of His-142 of protein 1C in the acid-induced disassembly of foot-and-mouth disease

virus capsids. J Gen Virol 80:1911–1918.

38.   Logan G, Freimanis GL, King DJ, Valdazo-González B, Bachanek-Bankowska K,

Sanderson ND, Knowles NJ, King DP, Cottam EM. 2014. A universal protocol to

generate consensus level genome sequences for foot-and-mouth disease virus and other

positive-sense polyadenylated RNA viruses using the Illumina MiSeq. BMC Genomics

15:828.

39.   Acevedo A, Andino R. 2014. Library preparation for highly accurate population

sequencing of RNA viruses. Nat Protoc 9:1760–1769.

40.   GitHub - najoshi/sickle: Windowed Adaptive Trimming for fastq files using quality.

41.   Peng Y, Leung HCM, Yiu SM, Chin FYL. 2012. IDBA-UD: a de novo assembler for

single-cell and metagenomic sequencing data with highly uneven depth.

Bioinformatics 28:1420–1428.

42.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment

686   search tool. J Mol Biol 215:403–410.

687   43.   Li H, Durbin R. 2010. Fast and accurate long-read alignment with Burrows–Wheeler

688   transform. Bioinformatics 26:589–595.

689   44.   Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G,

690   Mesirov JP. 2011. Integrative genomics viewer. Nat Biotechnol 29:24–26.

691   45.   Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler

692   transform. Bioinformatics 25:1754–1760.

693   46.   Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with

694   BWA-MEM.

695   47.   Jackson T, Sheppard D, Denyer M, Blakemore W, King AMQ. 2000. The Epithelial

696   Integrin αvβ6 Is a Receptor for Foot-and-Mouth Disease Virus. J Virol 74:4949.

697   48.   Arnold JJ, Cameron CE. 2000. Poliovirus RNA-dependent RNA polymerase

698   (3D(pol)). Assembly of stable, elongation-competent complexes by using a

699   symmetrical primer-template substrate (sym/sub). J Biol Chem 275:5329–36.

700   49.   de la Higuera I, Ferrer-Orta C, Moreno E, de Ávila AI, Soria ME, Singh K, Caridi F,

701   Sobrino F, Sarafianos SG, Perales C, Verdaguer N, Domingo E. 2018. Contribution of

702   a Multifunctional Polymerase Region of Foot-and-Mouth Disease Virus to Lethal

703   Mutagenesis. J Virol 92.

704   50.   Lasecka-Dykes L, Tulloch F, Simmonds P, Luke GA, Ribeca P, Gold S, Knowles NJ,

705   Wright CF, Wadsworth J, Azhar M, King DP, Tuthill TJ, Jackson T, Ryan MD. 2021.

706   Mutagenesis Mapping of RNA Structures within the Foot-and-Mouth Disease Virus

707   Genome Reveals Functional Elements Localized in the Polymerase (3Dpol)-Encoding

708   Region. mSphere 6.

51.   Kerpedjiev P, Hammer S, Hofacker IL. 2015. Forna (force-directed RNA): Simple and

effective online RNA secondary structure diagrams. Bioinformatics 31:3377–3379.

52. Schneider TD, Stephens RM. 1990. Sequence logos: a new way to display consensus

sequences. Nucleic Acids Res 18:6097.

53. Crooks GE, Hon G, Chandonia JM, Brenner SE. 2004. WebLogo: a sequence logo

generator. Genome Res 14:1188–1190.

54. Homan PJ, Favorov O V, Lavender CA, Kursun O, Ge X, Busan S, Dokholyan N V,

Weeks KM. 2014. Single-molecule correlated chemical probing of RNA. Proc Natl

Acad Sci U S A 111:13858–63.

55. Tulloch F, Pathania U, Luke GA, Nicholson J, Stonehouse NJ, Rowlands DJ, Jackson

T, Tuthill T, Haas J, Lamond AI, Ryan MD. 2014. FMDV replicons encoding green

fluorescent protein are replication competent. J Virol Methods 209:35–40.

56. Conzelmann KK. 2004. Reverse Genetics of Mononegavirales. Curr Top Microbiol

Immunol 283:1–41.

57. Schlender J, Bossert B, Buchholz U, Conzelmann K-K. 2000. Bovine Respiratory

Syncytial Virus Nonstructural Proteins NS1 and NS2 Cooperatively Antagonize

Alpha/Beta Interferon-Induced Antiviral Response. J Virol 74:8234.

58. Luna VER, Luk ADH, Tyring SK, Hellman JM, Lefkowitz SS. 1984. Properties of

bovine interferons. Exp 1984 4012 40:1410–1412.

59. Lasecka-Dykes L, Wright CF, Di Nardo A, Logan G, Mioulet V, Jackson T, Tuthill

TJ, Knowles NJ, King DP. 2018. Full genome sequencing reveals new southern african

territories genotypes bringing us closer to understanding true variability of foot-and-

mouth disease virus in Africa. Viruses 10.

60. Ferrer-Orta C, Arias A, Perez-Luque R, Escarmís C, Domingo E, Verdaguer N. 2004.

Structure of Foot-and-Mouth Disease Virus RNA-dependent RNA Polymerase and Its

Complex with a Template-Primer RNA. J Biol Chem 279:47212–47221.

61. Woodman A, Arnold JJ, Cameron CE, Evans DJ. 2016. Biochemical and genetic

analysis of the role of the viral polymerase in enterovirus recombination. Nucleic

Acids Res 44:6883–95.

62. Mohapatra JK, Pawar SS, Tosh C, Subramaniam S, Palsamy R, Sanyal A, Hemadri D,

Pattnaik B. 2011. Genetic characterization of vaccine and field strains of serotype A

foot-andmouth disease virus from India. Acta Virol 55:349–352.

63. Wright CF, Knowles NJ, Di Nardo A, Paton DJ, Haydon DT, King DP. 2013.

Reconstructing the origin and transmission dynamics of the 1967–68 foot-and-mouth

disease epidemic in the United Kingdom. Infect Genet Evol 20:230–238.

64. Lawrence P, Rieder E. 2009. Identification of RNA helicase A as a new host factor in

the replication cycle of foot-and-mouth disease virus. J Virol 83:11356–66.

65. Rodríguez-Pulido M, Borrego B, Sobrino F, Sáiz M. 2011. RNA Structural Domains

in Noncoding Regions of the Foot-and-Mouth Disease Virus Genome Trigger Innate

Immunity in Porcine Cells and Mice. J Virol 85:6492.

66. Rodríguez Pulido M, Serrano P, Sáiz M, Martínez-Salas E. 2007. Foot-and-mouth

disease virus infection induces proteolytic cleavage of PTB, eIF3a,b, and PABP RNA-

binding proteins. Virology 364:466–474.

67. Woodman A, Arnold JJ, Cameron CE, Evans DJ. 2016. Biochemical and genetic

analysis of the role of the viral polymerase in enterovirus recombination. Nucleic

Acids Res 44:6883–6895.

68. Arnold JJ, Cameron CE. 1999. Poliovirus RNA-dependent RNA polymerase (3D(pol))

is sufficient for template switching in vitro. J Biol Chem 274:2706–2716.

Figure 1. SHAPE analysis of the S fragment. (A) Superimposition of SHAPE reactivity on the *in silico* predicted secondary structure WT S fragment. Secondary structure was predicted using the Vienna RNA probing package and visualised using VARNA. NMIA reactivity is overlaid and represented on a colour scale from low (white) to high (red). Nucleotides for which there is no data are represented as grey. Coloured regions corresponding to the peaks in NIMIA reactivity are shown for ease of interpretation and localisation. **(B)** Individual nucleotide NMIA reactivity as analysed by SHAPE reactions and capillary electrophoresis. High reactivity indicates high probability of single-stranded regions i.e. non base-paired nucleotides. Data was analysed using QuSHAPE. Corresponding coloured regions are shown for some regions across **A** and **B** for ease of interpretation and localisation of SHAPE data to RNA structure. (n = 6, error bars represent SEM).

Figure 2. Truncations to the distal and proximal regions of the S fragment can impair replication.
(A) Cartoon schematic of FMDV O1K replicon with the region encoding the structural proteins replaced
with a GFP reporter. **(B)** Schematic representation of S fragment deletions. The maintained 3' and 5'
proximal (P) regions are represented by the dotted grey line and was constant in all deletions **(C-D)**
Replication of replicons with 70, 148, 246 and 295 nucleotides removed from the distal (D) region of
the S fragment was measured following transfection into BHK-21 cells or MDBK cells. Replication was
monitored by GFP expression using an IncuCyte, shown at 8 hours post-transfection alongside WT and
3D^{pol} inactivating mutant 'GNN' acting as a positive and negative control, respectively. **(E-F)** Replication
of replicons with 48, 97, 195 and 271 nucleotides removed from the proximal region of the S fragment
and measured as in (C). (n = 3, error bars represent SEM) ** P<0.01, **** P<0.0001.

Figure 3. The extent of nucleotide and pairing conservation within the S fragment of the FMDV. (A)
A schematic representation of conserved RNA structure of the S fragment in all 118 FMDV isolates.
Nucleotide positions which form conserved pairing in 116/118 FMDV isolates were colour-coded
according to number of pairing types (purple for 1 nucleotide pairing type at a given position and
yellow for 6 nucleotide pairing types; see the colour matrix included at the left bottom corner of the

figure). There are six possible nucleotide pairings: A-U, G-C, G-U, U-A, C-G and U-G. Nucleotide pairings
which were not conserved in three or more FMDV isolates, remained white. Due to the length of the
S fragment stem-loop, the resolution of the image was not sufficient for detailed view of the individual
nucleotide pairings and so the S fragment was artificially divided into two parts, proximal and distal,
and the images of their schematic RNA structures enlarged (**B-C**). **(B)** The proximal part of the S
fragment included nucleotide positions 1 to 90 and 272 to 364, **(C)** while the distal part included
nucleotide positions 90 - 272. Numbers represent nucleotide positions of the S fragment sequence.

801
Figure 4. Fitness of FMDV containing a truncation at the proximal end of the S fragment. (A)
Recovered WT, P-48, P-97 and D-148 virus populations were used to infect BHK-21 cells at an MOI of
0.01 PFU. CPE was monitored over time using live an IncuCyte S3 live cell imaging and is shown as a
drop in cell confluency; (n =3 error bars represent SEM). **(B)** Plaque sizes of the WT, P-48, P-97 and
D-148 variants grown on BHK-21 cells. Each wells containing plaques was scanned and the plaque size
was estimated in pixels. All plaques were counted to avoid bias in plaque selection. Grey dots
represent individual plaques, while black dot represents the mean; *** P < 0.001 when comparing to
plaque size induced by the WT variant.

Figure 5. Compensatory mutation I189L in the structure of 3D^{pol} and its effect on replication of FMDV replicons. (A) WT 3D^{pol} crystal structure with I189 residue highlighted in blue, shown with (bottom, left) and without bound RNA (top, left). The I189L residue mutation was modelled onto the WT 3D^{pol} crystal structure and highlighted in red (right). Structure shown with (bottom, right) and without (top, right) bound RNA. PDB file 1WNE. **(B)** The 3D^{pol} I189L mutation was introduced into WT, GNN and P-97 replicons, termed WT I189L, GNN I189L and P-97 I189L respectively. Replicon RNA was transcribed and transfected into BHK-21 cells. WT and 3D^{pol} GNN replicons were included as controls, with the latter for the level of input translation (n =3 error bars represent SEM, * P < 0.05).

Figure 6. **Single nucleotide addition to a Sym/Sub substrate by WT and I189L 3Dpol.** **(A)** Schematic
of the Sym/Sub experimental protocol. Radioactively end labelled RNA oligonucleotides are annealed
before addition of rNTPs and recombinant 3D^{pol}. **(B)** Extension of the 10mer Sym/Sub template with a
single nucleotide (ATP) using either WT or I189L FMDV 3D^{pol}. Aliquots of the reactions were taken at
20, 60, 120 and 300 seconds, RNA fragments separated by electrophoresis and visualised using a
phosphoimager. **(C)** Densitometry of +1 product is plotted as rate of addition of a single nucleotide
shown over time, %+1 product refers to the total amount of input Sym/Sub template that was
extended by 1 nucleotide.

Figure 7. Sym/Sub assays showing differences between WT and I189L 3D^{pol} in the production of a ~
+12 nucleotide product. (A) Schematic of the Sym/Sub experimental protocol using all four rNTPs. **(B)**
Extension of the 10mer Sym/Sub template with all four nucleotides using either WT or I189L
recombinant FMDV 3D^{pol}. Aliquots of the reactions were taken at 20, 60, 120 and 300 seconds, RNA
fragments separated by electrophoresis and visualised using a phosphoimager. **(C)** Densitometry of
the production of the +4 product by either WT or I189L 3D^{pol} over time. **(D)** Densitometry of the
production of the +12 product by either WT or I189L 3D^{pol}. Data are shown as the % of the input
Sym/Sub template elongating to the +4 or +12 products (n = 3, error bars represent SEM).

Figure 8. Misincorporation by WT and I189L 3D^{pol}. (A) Sym/Sub experiments were repeated with the addition of either ATP, ATP and UTP, ATP and UTP and GTP or all four rNTPs together. Reactions were continued for 300 seconds before electrophoresis and visualization. Misincorporation of GTP to generate a +4 product is highlighted by the red box. **(B)** Densitometry of the misincorporated GTP shows no difference in the rate of misincorporation by WT or I189L 3D^{pol}. **(C)** WT, P-97 and P-97 I189L replicons were transfected into BHK-21 cells in the presence of up to 1 mM of ribavirin. Replication was monitored by expression of the GFP reporter and shown here at 8 hours post-transfection. **(D)** Schematic for potential mechanism for the generation of a +12 product. Since misincorporation can occur at the fourth position allowing for the addition of a second guanosine residue, if templates were to separate and reanneal by either strand slippage or annealing of a new template, extension could result in the addition of 12 nucleotides (n = 3, error bars represent SEM).

Supplementary Figure S1. Effect of introduction of *Eag*I sites on replication. Number of GFP positive cells/well detected at 8 hours post-transfection of BHK-21 cells. WT represents the original O1K replicon sequence. 3D-GNN is the polymerase defective negative control. 5' *Eag*I replicon possesses mutations to the sequence to create an *Eag*I site at the 5' end of the S fragment. 3' *Eag*I possesses mutations to the sequence to create and *Eag*I site at the 3' end of the S fragment. Aerie shows replicon with both 5' and 3' sites modified and Aerie 3D-GNN comprises both *Eag*I sites and 3D-GNN mutation. n=2, error bars represent SEM.

866   **Supplementary Figure S2. Time course of FMDV replicon replication.** Number of GFP positive
867   cells/well detected at hourly time points post-transfection of WT and 3D-GNN replicons in BHK-21
868   cells. n=2, error bars represent SEM.

873 **Supplementary Figure S3. Conservation of the isoleucine at the position 189 (I189) within the 3D^pol**
874 **protein of FMDV.** Sequence logo based on 1123 sequences of FMDV 3D^pol available on fmdbase.org
875 and prepared using WebLogo 3 server. The x-axis represents amino acids positions of the 3D^pol protein
876 of FMDV, while the y-axis shows probability of a particular amino acid being present at a given position.
877 Amino acids are colour-coded according to their charge.
878

Supplementary Table S1

FMDV sequences selected from GenBank

Serotype:	Number of isolates:	GenBank accession numbers:
A	19	AY593788, MH053305, JF749843, HM854024, HQ832580, MH053306, KM268896, AY593802, KJ608371, MH053307, AY593751, AY593754, AY593761, AY593764, AY593766, AY593767, HM854022, AY593791, AY593794
Asia 1	12	AY593795, AY687334, DQ533483, DQ989306, DQ989315, DQ989319, EF149010, EF614458, HQ632774, JF739177, KM268898, MF782478
C	6	MH053308, KM268897, MH053309, AJ133357, MH053310, AJ007347
O	21	AY593819, MH053313, MH053311, MH053312, KF112885, KJ206909, HQ632769, HQ632771, KU291242, KR401154, GU384683, KF694737, AJ539140, MH053315, JX040491, MH053317, MH053318, MH053316, KJ560291, DQ404170, KU821591
SAT 1	21	AY593838, AY593845, MH053319, AY593844, JF749860, MH053321, AY593846, AY593839, AY593842, AY593841, AY593840, MH053322, AY593843, KM268899, MH053323, MH053324, MH053325, MH053326, MH053327, MW355668, MW355669
SAT2	19	MH053330, MH053332, MH053328, MH053329, JX014255, MH053333, AY593849, JX014256, MW355670 - MW355673, AY593847, MH053335, KM268900, JF749862, MH053336, MH053337, KU821592,
SAT3	20	AY593853, AY593851, MH053339, MH053340, MH053344, MH053343, AY593850, KJ820999, MH053341, MH053351, KX375417, KM268901, MH053350, MW355674 - MW355680