High prevalence of lipopolysaccharide mutants and R2-Pyocin susceptible variants in *Pseudomonas aeruginosa* populations sourced from cystic fibrosis lung infections.

Madeline Mei[a,b,c], Preston Pheng[a], Detriana Kurzeja-Edwards[a] & Stephen P. Diggle[a*]

[a]Center for Microbial Dynamics and Infection, School of Biological Sciences, Georgia Institute of Technology, Atlanta, Georgia, USA; [b]Department of Pediatrics, Division of Pulmonary, Allergy and Immunology, Cystic Fibrosis, and Sleep, Emory University School of Medicine, Atlanta, Georgia, USA; [c]Emory+Children's Center for Cystic Fibrosis and Airway Disease Research, Emory University School of Medicine, Atlanta, Georgia, USA.

Running title: High Frequency of R2-pyocin susceptible *Pseudomonas aeruginosa* in CF lungs

*Address correspondence to stephen.diggle@biosci.gatech.edu

Keywords: R-pyocin, bacteriocin, *Pseudomonas aeruginosa*, cystic fibrosis, lipopolysaccharide

Author contributions: M.M., and S.P.D. conceived the study and wrote the paper; M.M., P.P. and D.K.E. performed the experimental work and analyzed the data.

ABSTRACT

Chronic, highly antibiotic-resistant infections in cystic fibrosis (CF) lungs contribute to increasing morbidity and mortality. *Pseudomonas aeruginosa*, a common CF pathogen, exhibits resistance to multiple antibiotics, contributing to antimicrobial resistance (AMR). These bacterial populations display genetic and phenotypic diversity, but it is unclear how this diversity affects susceptibility to bacteriocins. R-pyocins, i.e. bacteriocins produced by *P. aeruginosa*, are phage tail-like antimicrobials. R-pyocins have potential as antimicrobials, however recent research suggests the diversity of *P. aeruginosa* variants within CF lung infections leads to varying susceptibility to R-pyocins. This variation may be linked to changes in lipopolysaccharide (LPS), acting as the R-pyocin receptor. Currently, it is unknown how frequently R-pyocin-susceptible strains are in chronic CF lung infection, particularly when considering the heterogeneity within these strains. In this study, we tested R2-pyocin susceptibility of 139 *P. aeruginosa* variants from 17 sputum samples of seven CF patients and analyzed LPS phenotypes. We found that 83% of sputum samples did not have R2-pyocin-resistant variants, while nearly all samples contained susceptible variants. there was no correlation between LPS phenotype and R2-pyocin susceptibility, though we estimate that about 76% of sputum-derived variants lack an O-specific antigen, 40% lack a common antigen, and 24% have altered LPS cores. The absence of a correlation between LPS phenotype and R-pyocin susceptibility suggests LPS packing density may play a significant role in R-pyocin susceptibility among CF variants. Our research supports the potential of R-pyocins as therapeutic agents, as many infectious CF variants are susceptible to R2-pyocins, even within diverse bacterial populations.

IMPORTANCE

Cystic fibrosis (CF) patients often experience chronic, debilitating lung infections caused by antibiotic-resistant Pseudomonas aeruginosa, contributing to antimicrobial resistance (AMR). The genetic and phenotypic diversity of P. aeruginosa populations in CF lungs raises questions about their susceptibility to non-traditional antimicrobials, like bacteriocins. In this study, we focused on R-pyocins, a type of bacteriocin with high potency and a narrow killing spectrum. Our findings indicate that a large number of infectious CF variants are susceptible to R2-pyocins, even within diverse bacterial populations, supporting their potential use as therapeutic agents. The absence of a clear correlation between lipopolysaccharide (LPS) phenotypes and R-pyocin susceptibility suggests that LPS packing density may play a significant role in R-pyocin susceptibility among CF variants. Understanding the relationship between LPS phenotypes and R-pyocin susceptibility is crucial for developing effective treatments for these chronic infections.

INTRODUCTION

Cystic fibrosis (CF) is a genetic disorder primarily characterized by chronic respiratory infections, leading to progressive loss of lung function and increased morbidity and mortality in affected individuals (1). The disease arises from a defective anion pump, known as the cystic fibrosis transmembrane regulator (CFTR), which results in an imbalance of chloride on epithelial cell surfaces (1). This imbalance causes mucus dehydration, making it too thick for proper clearing of airways along the respiratory tract in people with CF. Consequently, this creates an environment that promotes bacterial and fungal infections, which are notoriously challenging to treat.

Pseudomonas aeruginosa, a Gram-negative opportunistic bacterium, is the most prevalent bacterial pathogen that colonizes and persistently infects CF lungs (2-7). With a large genome, *P. aeruginosa* can regulate diverse metabolic processes, resist antibiotic treatment, and control various complex social behaviors that aid in its persistence in infections (8-16). Once established

in the CF lung, *P. aeruginosa* is extremely difficult to eradicate and often becomes multi-drug
resistant (MDR). Although *P. aeruginosa* inherently possesses many antibiotic-resistant
mechanisms, strains sourced from chronic lung infections are known to become particularly
problematic and resistant to multiple antibiotics over time (17-21). While recent data suggests that
the approval and use of the new triple CF therapeutic TRIKAFTA®
(elexacaftor/tezacaftor/ivacaftor) has decreased the overall prevalence of *P. aeruginosa* infection,
the 2019 CF Foundation Patient Registry Annual Report notes that the prevalence of MDR *P.*
aeruginosa infection remains constant (22). In the era of antibiotic resistance, new treatment
approaches are crucial against *P. aeruginosa* and other multi-drug resistant pathogens, especially
in the case of CF chronic lung infections.
78
One of the challenges in eradicating *P. aeruginosa* from CF lungs is likely due to the high degree
of phenotypic, and genomic diversity that evolves within *P. aeruginosa* populations over the
course of infection (6, 18, 19, 23, 24). While this heterogeneity is known to affect antibiotic
susceptibility, its impact on alternative therapeutic options, such as bacteriocins and
bacteriophage, remains poorly understood (6, 18, 19). As pathogens like *P. aeruginosa* become
increasingly difficult to eliminate, antimicrobials such as bacteriocins, may serve as useful
alternative therapies in combination with antibiotics and phage for more effective treatment.
86
Bacteriocins are narrow-spectrum, proteinaceous antimicrobials produced by one strain of
bacteria and active against other strains of the same or closely related species (25, 26). *P.*
aeruginosa produces various bacteriocins called pyocins, which are grouped based on similarities
in their physical and chemical properties, export mechanisms, and mode of action (27-31).
Pyocins produced by *P. aeruginosa* are classified as S-, R-, and F-pyocins, each with varying
antimicrobial spectra (30-34). R-pyocins are narrow-spectrum, contractile, phage tail-like
bacteriocins, further categorized into subtypes R1-R5 based on specificity to different receptors
on target cells, however R-types 2-4 are often considered as a single, functional subtype (30, 31,
35-38). The proposed mode of action involves the R-pyocin tail fiber foot binding to a
monosaccharide receptor on the lipopolysaccharide (LPS) (specific to each R-subtype), causing
the sheath to contract and push the tail spike and core to puncture the outer membrane, resulting
in membrane depolarization and eventual cell death (**Fig. S1**) (37, 39-44). R-pyocins have gained
interest for their potential to combat *P. aeruginosa* due to their powerful ability to kill target cells,
their inability to replicate without a DNA-containing head, and their well-documented mechanism
of action (30, 31, 41, 43-45). In this study, we focus on R-pyocins because of their promising
features as antimicrobial agents and the need for a better understanding of their effectiveness
against clinical strains of *P. aeruginosa*.
104
The LPS of *P. aeruginosa* not only acts as the receptor for R-pyocins, but importantly plays a
significant role in virulence and immune evasion. LPS is composed of three major regions: the
lipid A, the core oligosaccharide, and the O-specific antigen (OSA) (46). The core structure is
relatively conserved across *P. aeruginosa* strains and acts as a platform for the attachment of
various OSA side chains; Both OSA and common polysaccharide antigen (CPA) can be found in
P. aeruginosa LPS, but their presence and heterogeneity vary among different isolates or variants
even within the same strain ((46-51). The core structure of *P. aeruginosa* LPS is essential for
maintaining outer membrane integrity and plays a crucial role in bacterial survival, thus it is costly
to bacterial fitness to alter. LPS is composed of a conserved lipid A region, anchoring LPS to the
outer membrane, and a core oligosaccharide region that extends from lipid A (47, 52-54). This
core region serves as the attachment site for the OSA and CPA side chains, (as well as R-pyocins)
contributing to the overall heterogeneity of *P. aeruginosa* LPS (38, 47, 54-58). OSA structures
can differ in their length, sugar composition, and linkage patterns, which contributes to antigenic
variation and immune evasion strategies employed by the bacterium during infection (59-61). The

presence and composition of OSA and CPA side chains in *P. aeruginosa* LPS can vary
significantly between strains, thus the diverse repertoire of OSA structures can influence the
bacterium's interactions with not only other pathogens in infection, but also the host immune
system, allowing for evasion of immune surveillance and facilitating chronic infection
establishment in CF patients (59, 62-65). Moreover, certain *P. aeruginosa* isolates have been
reported to downregulate or even lose OSA expression during chronic CF infections, contributing
to increased adaptability and persistence in the host environment (51, 66-68)

During chronic CF infections, P. aeruginosa faces a challenging host environment characterized
by immune responses and antibiotic exposure. In this context, certain variants of *P. aeruginosa*
have been observed to downregulate or lose OSA expression (46, 51, 69). This phenomenon is
thought to provide a survival advantage by reducing recognition and clearance by the host
immune system. By modulating OSA expression, these isolates can persist and adapt more
effectively within the CF lung environment, leading to prolonged and difficult-to-treat infections.
The downregulation or loss of OSA expression by certain isolates during chronic CF infections
further underscores the bacterium's adaptability and persistence in the host environment (46, 69).
Understanding the structural and functional characteristics of *P. aeruginosa* LPS is crucial for
developing targeted therapeutic strategies to combat these challenging infections effectively, and
particularly when considering LPS-binding antimicrobials such as bacteriophage and R-type
bacteriocins. While there have been studies serotyping and R-pyocin typing panels of strains (38,
70, 71), there has not yet been a study examining LPS diversity among variants of the same strain
or comparing variants within multiple different strains and the resulting impacts on R-pyocin
susceptibility.

In previous research, we tested 20 individual *P. aeruginosa* variants from sputum samples
collected from three different individuals with cystic fibrosis (CF) (60 variants total, from 3 *P.*
aeruginosa populations). We found a variation in susceptibility to R2-pyocins within these
populations (51). Additionally, we isolated and compared the lipopolysaccharide (LPS)
phenotypes of three of these variants from the same sputum sample and patient, discovering that
each variant had different LPS glycoforms, which likely contributed to their varying susceptibility
(51). If R-pyocins are to be considered as a therapy, it is essential to further investigate their
efficacy against diverse clinical strains of *P. aeruginosa*, as the prevalence of R-pyocin-
susceptible strains in CF chronic lung infections is currently not well understood.

We previously found that R1-pyocin producers make up the majority of R-pyocin-typeable CF *P.*
aeruginosa strains, and other studies have shown that R1-type producers are often susceptible
to R2-pyocins (38, 51, 70). As a result, we expected a high frequency of *P. aeruginosa* strains
isolated from chronic lung infections with LPS mutations conferring R2-pyocin susceptibility.
Although we discovered no correlation between LPS phenotype and R2-pyocin susceptibility,
most *P. aeruginosa* strains isolated from CF lung infections were sensitive to R2-pyocins. Our
findings support the potential use of R-pyocins as alternative anti-pseudomonal treatments in
chronic infections.

RESULTS

The majority of *P. aeruginosa* populations in CF patients have variants susceptible to R2-
pyocins. To determine the prevalence of R2-pyocin susceptible variants among *P. aeruginosa*
populations from CF sputum samples, we analyzed up to 10 previously described variants (51)
from longitudinal samples of 7 individuals with CF (139 total variants collected from 17 sputum
samples or "populations") using the soft-agar overlay method (**Fig. S2**). For consistency, we used

PAO1Δ*rmlC,* an LPS core mutant (38), as an R2-pyocin susceptible control and PAO1 wt as a
resistant control in each experiment. Using R2-pyocins from PAO1 without the pf4 prophage
(PAO1ΔPf4) (72), we assessed the relative susceptibility of each CF variant and found that (i)
each population showed heterogeneity within the population, consistent with previous work using
a microtiter plate method (51), and (ii) nearly every *P. aeruginosa* population included variants
with some degree of susceptibility to R2-pyocins. In our sample, we found that only 3 out of 7
individuals had sputum containing R2-pyocin resistant variants (**Fig. 1**). We also discovered that
most populations contained R2-susceptible variants even across longitudinal sampling from the
same patient, although the degree of relative susceptibility and sensitivity to R2-pyocins fluctuated
over time (**Fig. 2**). Longitudinal sputum collections in our biobank showed no trend of variants
becoming R2-pyocin resistant over a period of approximately 2 years. To further confirm that
these results were R2-pyocin dependent, we included lysates from a PAO1ΔPf4ΔR mutant
(lacking both the prophage and a functional R2-pyocin) in each assay, ensuring no killing activity
when R2-pyocins were absent.
182
There is a high frequency of LPS mutants and diversity among LPS phenotypes in CF
strains of *P. aeruginosa*. To characterize the LPS phenotypes of R2-pyocin susceptible and
resistant variants in our biobank, we isolated LPS using a hot-aqueous phenol method (73) and
compared the LPS migration profiles of our clinical variants to those of isogenic LPS mutants of
our Nottingham PAO1 wild type strain. Out of 139 variants across all populations, we observed
that 40% lacked common antigen (CPA or A-band) (**Fig. 3A**) and 76% lacked O-specific antigen
(OSA or B-band) (**Fig. 3B**). Additionally, 24% presented altered LPS cores (**Fig. 3C**), as
determined by comparing to PAO1. We found that 13 out of the 17 sputum collections sampled
contained variants with diverse LPS profiles; approximately 76% of our *P. aeruginosa* populations
contained a mixture of variants with varying LPS phenotypes.
193
To assess the correlation between R2-pyocin susceptibility and specific LPS glycoform
phenotypes in CF strains, we conducted a one-tailed Spearman correlation analysis for each
phenotype individually. We assumed independence between the three glycoform phenotypes
(CPA+ or CPA-, OSA+ or OSA-, WT or altered core) for these analyses, even though some
dependencies may exist due to overlapping biosynthetic pathways and the essentiality of the LPS
core. We found no significant correlations specifically between the presence or absence of CPA
(**Fig. 3D**), OSA (**Fig. 3E**), or an altered core (**Fig. 3F**) (p > 0.05).
201
We further evaluated the relationship between LPS phenotype and R2-pyocin susceptibility by
assessing the LPS as a whole. Using a numbered scale, we assigned an LPS Score of 0 when a
variant presented as CPA-, OSA-, and had an altered core; this variant was treated as having 0/3
WT LPS characteristics. A score of 1 was assigned when a variant exhibited one of the three LPS
glycoforms as WT (presence of OSA, CPA, or unaltered core), a score of 2 when two of three
glycoforms were WT, and a score of 3 when a variant presented CPA+, OSA+, with a WT core.
Using this classification system, we also found no correlation with R2-pyocin susceptibility (Fig.
4). However, we did find that three R2-pyocin resistant variants were all CPA- and OSA-. Only
20% of variants from all sputum samples exhibited CPA+, OSA+, and WT LPS cores.
211
212
213   **DISCUSSION**
214
Cystic fibrosis (CF) chronic lung infections present a significant challenge due to their high
antibiotic resistance, difficulty in eradication, and potentially lethal nature (1). *P. aeruginosa*, a
common pathogen found in CF lungs, contributes to infection-related morbidity and mortality due

to its significant antibiotic resistance and ability to diversify over time (2-7). As a result, it is crucial
to explore alternative antimicrobial and therapeutic options for treating these chronic infections.

Bacteriocins, particularly R-pyocins, have recently gained attention for their high potency, narrow
spectrum, and anti-pseudomonal properties. However, the efficacy of R-pyocins and other phage-
like bacteriocins remains uncertain due to the diverse phenotypic behavior of *P. aeruginosa* in
chronic CF lung infections. Previous work demonstrated that *P. aeruginosa* populations display
varying susceptibility to R-pyocins, emphasizing the need to better understand the implications of
this heterogeneity on treatment outcomes (51). This heterogeneity is particularly important when
considering resistant variants in infection, as the prevalence of resistant variants and the benefits
of resistance-related phenotypes to survival in chronic infection remain unclear.

In a previous study, we tested the R2-pyocin susceptibility of 60 isolates from 3 patients
expectorated sputum samples, but the limited sample size did not provide a comprehensive
understanding of R-pyocin sensitivity and resistance behaviors over time or across more CF
patients. In the current study, a larger number of variants were tested, collected from multiple
longitudinal samples and more individuals with chronic CF lung infections. A total of 139 *P.
aeruginosa* variants were collected from 17 expectorated sputum samples (including longitudinal
collections) of 7 individuals with CF and were tested for R2-pyocin susceptibility and LPS
phenotypes.

The analysis of *P. aeruginosa* populations from CF lung infections revealed that (i) approximately
76% of sputum samples contained *P. aeruginosa* populations exhibiting heterogeneity among
LPS phenotypes; (ii) no correlation was found between R2-pyocin susceptibility and the presence
or absence of a specific LPS glycoform; and (iii) it was estimated that about 76% of the variants
sampled from sputum lacked O-specific antigen, 40% lacked common antigen, and 23% exhibited
altered LPS cores.

In previous work, R-pyocin susceptibility testing has typically focused on one variant per sputum
collection (38, 70). However, the current study expanded this approach by examining multiple
variants from a wide range of sputum samples from various individuals, including some
longitudinal samples. This comprehensive analysis revealed that nearly all bacterial populations
studied contained susceptible variants (16 out of 17 populations), with heterogeneity in
susceptibility observed among the susceptible variants (51). Only about 5% of the tested variants
were resistant to R2-pyocins, and these variants were obtained from only 4 sputum samples from
3 patients. No trends in R2-pyocin susceptibility were identified across longitudinal samples, and
susceptibility did not cluster by patient, but by longitudinal collection.

The high frequency of R2-pyocin susceptible strains in CF lung infections suggests that (i) there
is no strong selective pressure for *P. aeruginosa* to alter LPS in a way that leads to R2-pyocin
resistance, and (ii) R2-pyocins could be effective against *P. aeruginosa* for the majority of
individuals with CF lung infections, as 94% of individuals in the cohort were infected with R2-
pyocin susceptible variants.

Previous studies have shown there is some correlation between OSA serotype and R-pyocin
genotype, and though susceptibility to R1, R2, and R5 pyocins were tested for representative
strains of varying serotypes, no statistic was provided linking O-antigen serotype with R-pyocin
susceptibility (38, 70). Several specific residues on the outer core have been identified as binding
factors for each of the 3 functional subtypes of R-pyocins (38) and this implies that O-serotype
alone is not the sole determinant of R-pyocin susceptibility as the residues recognized are present

among all naturally occurring glycoforms of LPS regardless of serotype. It has not been shown that any of the serotypes have differently structured outer cores, thus in this work, we sought to first look at the presence or absence of the LPS "decorations" (OSA and CPA) and the impacts of variations in these phenotypes rather than correlate serotypes. This study is novel in examining LPS glycoform diversity (i.e. presence or absence of OSA and CPA) among variants of the same strain or comparing variants within multiple different strains and the resulting impacts on R-pyocin susceptibility.

In this study, LPS phenotypes were analyzed based on the presence or absence of the CPA, the OSA, and the altered or PAO1-like (WT) LPS core. No correlations were found between any of the phenotypes or between various combinations of LPS phenotypes and R2-pyocin susceptibility. However, some sputum samples contained variants with similar LPS phenotypes. For example, one sputum collection had all variants exhibiting the phenotype CPA+, OSA+, and presenting a WT core. A high percentage of variants lacked OSA (76%), while more than half maintained the CPA (40%).

The absence of correlation between specific LPS glycoform phenotypes suggests that other LPS-modifying behaviors, such as "packing density", may be more influential in determining R2-pyocin susceptibility. LPS packing density refers to the arrangement and density of lipopolysaccharide (LPS) molecules on the outer membrane of Gram-negative bacteria, which is also essential for the stability and integrity of the bacterial outer membrane (38, 74-76). Packing density determines how closely the LPS molecules are arranged on the surface of the membrane, which can influence several aspects of bacterial physiology, including the permeability of the outer membrane to antimicrobial agents, the accessibility of OSA side chains, affecting the recognition of bacterial pathogens by immune cells and the production of antibodies, biofilm formation, and the exposure and accessibility of residues targeted by bacteriophage and bacteriocins (including R-pyocins) }(74-76). Other groups have proposed the importance of LPS packing density and the impact on R-pyocin susceptibility (38, 50). One group found several genes responsible for outer membrane lipid homeostasis and LPS packing via TnSeq, including the *mla* and *lpt* transport pathways (50). Though there is some identification of pathways involved in maintaining outer membrane integrity, lipid homeostasis, and LPS arrangement, the regulation of these behaviors remain poorly understood.

In summary, the majority of *P. aeruginosa* strains isolated from CF lung infections appear to be susceptible to R2-pyocins, supporting the potential for R-pyocins as effective antimicrobials. This study also revealed no correlation between LPS phenotype and R2-susceptibility in strains evolved in chronic infection, indicating that further research is necessary to evaluate other LPS-modifying behaviors and packing density in the CF lung environment, as well as the implications of these behaviors for the use of LPS-binding phage and bacteriocins.

MATERIALS AND METHODS

Bacterial strains, media, and culture conditions. We collected expectorated sputum samples for this study in previous work from adult CF patients through Emory–Children's Center for Cystic Fibrosis and Airways Disease Research by the Cystic Fibrosis Biospecimen Laboratory with IRB approval (Georgia Tech approval H18220) (51). We isolated *P. aeruginosa* populations from each sputum sample using selective media (Pseudomonas Isolation Agar, Sigma-Aldrich) before isolating single colonies for further characterization. In total we studied 139 variants sourced from

CF chronic lung infections for R2-pyocin susceptibility and LPS phenotype. All *P. aeruginosa* lab
strains used in this work are listed and described in **Table 1**. All *P. aeruginosa* strains isolate from
CF sputum are listed and described in **Supplementary Dataset 1**. We grew all bacterial cultures
in lysogeny broth (LB) medium at 37°C with 200 rpm shaking agitation and used standard genetic
techniques for construction of *P. aeruginosa* mutants. The construction of plasmids and
generation of mutants are described below and all plasmids and primers used for mutant
generation can be found in **Table 1**.

Generation of mutants. We constructed the PAO1Δ*rmlC* mutant as follows: 600 bp DNA
sequences flanking the open reading frames of *rmlC* (PA5164) were PCR amplified using Q5
DNA polymerase (New England Biolabs, Ipswich, MA) (38). We then cloned these sequences
into the SphI-XbaI digested suicide vector pDM4 (50) by Gibson assembly using NEBuilder HiFi
Assembly master mix (New England Biolabs), transformed into *E. coli* S17-λpir by electroporation
and selected on LB agar plates supplemented with 30 µg/mL chloramphenicol. We verified cloned
inserts by colony PCR and Sanger sequencing (Eton Bioscience, NC). We introduced the deletion
construct into PAO1 wt by electroporation and strains carrying single crossover insertions of the
deletion constructs were selected on LB agar plates supplemented with 300 µg/mL
chloramphenicol (77, 78). Resistant colonies were cultured in LB without antibiotic and plated on
LB agar plates with 0.25% NaCl and 5% sucrose to select for loss of the construct (79). We
screened sucrose-resistant colonies for antibiotic sensitivity to ensure loss of the vector sequence
and were assessed for presence of the gene deletion by colony PCR and Sanger sequencing of
the PCR product. We constructed the PAO1ΔPf4ΔR mutant by deleting the tail fiber and
chaperone genes of the R-pyocin gene cassette (PA0620-PA0621) from the PAO1ΔPf4 strain as
described previously (51, 72).

Expression and collection of R-pyocins. We collected R-pyocins in lysates as previously
described (51). LB cultures of PAO1, PAO1ΔR (R-pyocin null mutant) (46), PAO1ΔPf4 (prophage
mutant) (72), and PAO1Δ*pf4* ΔR were inoculated at 1:100 from overnight planktonic cultures, and
grown to mid-logarithmic growth phase (approximately 4 h) in 10 mL of LB media. We then added
ciprofloxacin to each culture for a final concentration of 0.1 µg/mL to induce R-pyocin production
(51, 80, 81), and cultures were incubated for a further 3 h. Chloroform was used to lyse remaining
cells, before centrifuging lysates at ~3,300 x g for 10 min. The R-pyocin-containing supernatants
were separated and stored at 4°C.

Spot assay for Relative R-pyocin Susceptibility. Spot assays to assess R-pyocin activity were
conducted as previously described (51). We used overnight cultures of clinical *P. aeruginosa*
variants to inoculate 4 mL of cooled soft top (0.4%) agar at an optical density at 600 nm (OD$_{600}$)
of 0.01. This mixture was poured onto LB agar plates for an overlay of the indicator strain. We
extracted R2-pyocin lysates from PAO1, PAO1ΔPf4 and respective R-pyocin null mutants were
vortexed, then serial diluted 10-fold in phosphate buffered saline (PBS) before spotting 5 µL of
each dilution onto the soft-top overlay indicator strains. Spots were allowed to dry before
incubating plates at 37°C overnight. Clear zones of growth inhibition from PAO1ΔPf4 lysates were
used to determine R-pyocin dependent activity against the indicator strains, and we determined
relative susceptibility by counting the number of these serial dilutions of lysate indicating inhibition.
PAO1 wild type was used as an R2-resistant control strain and PAO1Δ*rmlC* was used as an R2-
susceptible control strain to standardize and quality control each experiment (38). We conducted
spot assays for R-pyocin susceptibility in triplicate.

LPS isolation and visualization. We isolated LPS of bacterial cultures and visualized as
described by previous methods (73). We used overnight bacterial cultures of 5 mL LB grown at
37°C at 200 rpm shaking. We then diluted overnight cultures 1:10 with LB to measure OD$_{600}$

before making a 1 mL suspension of bacteria adjusted to an OD_{600} of 0.5 and pelleting by
centrifugation at 10,600 x g for 10 min. The cell pellet was resuspended in 200 µL of 1x SDS-
buffer. Suspended bacterial cells were boiled for 15 min before we prepared LPS by hot aqueous-
phenol extraction as previously described but without the addition of DNase I and RNase solutions
(as deemed optional in the protocol) (73). We visualized samples using 15 µL of LPS preparation
on a 4%-20% Tris-glycine mini gel using the Pro-Q Emerald 300 Lipopolysaccharide Gel Stain Kit
(ThermoFisher) (73).

365
366 **Statistical analysis.** We performed all data analysis using Prism (Graphpad software, version
367 9).

368
369 **Data Availability.** We have made raw data and code available in the Dryad Digital Repository:
370 https://datadryad.org/stash/share/k9quE9rIzJ-gRXo42tF559U2MEpmXH6wC3emdpinkkM (82).

371
372 **Supplemental Material**

373
Supplementary Dataset 1. Detailed description of bacterial strains, average susceptibility and
LPS scoring used in this study.

Acknowledgements. We thank the Cystic Fibrosis and Airways Disease Research (CF-AIR) at
Emory University, for clinical CF populations of *P. aeruginosa*. We also thank Patrick Secor for
generously sharing the PAO1ΔPf4 strain with us. This work was supported by the Cystic Fibrosis
Foundation by grants (DIGGLE18I0 and DIGGLE20G0),the National Institutes of Health (NIH) for
a grant (R01AI153116) and the National Science Foundation (NSF) for a grant (2003721) to SPD.
This work was also supported by the NIH Vaccinology Training Program Grant (#T32AI074492)
to MM.

384
385
386 **Figure legends**

387
388 **Table 1.** Strains, primers and plasmids used in this study.

389
**Figure 1. R2-pyocin inhibition zones for multiple CF variants from different patient sputum
samples.** The inhibition zones caused by R2-pyocin applied to soft agar overlays containing
clinical variants from the CF sputum samples listed on the left. PAO1Δ*rmlC* serves as an R2-
susceptible control, and PAO1 wt as an R2-resistant control. Red boxes indicate patients with
populations containing resistant variants. Up to ten variants per sputum sample were tested. An
inhibition zone indicates R2-pyocin susceptibility, while the absence of a visible zone indicates
resistance.

397
**Figure 2. R2-pyocin susceptibility varies among clinical *P. aeruginosa* populations and
changes over time.** Relative susceptibility to R2-pyocins differs among variants in each
population and across populations. Susceptibility heterogeneity is observed across all five
patients and within the same patient's longitudinal samples. There is no trend of strains becoming
more resistant to R2-pyocins in longitudinally sampled populations, up to 2 years (consecutive
numbers with collection number in parentheses). Two sputum samples contained at least one
variant resistant to R2-pyocins (Relative Susceptibility = 0). Relative Susceptibility was
determined by counting the number of serial dilutions (including neat) where inhibition zones
appeared after R2-pyocins were applied to agar overlays. Means are shown with 95% confidence
interval error bars.

Figure 3. LPS phenotype heterogeneity and lack of correlation with R2-susceptibility. A one-tailed Spearman correlation analysis for each phenotype revealed no correlations between R2-pyocin susceptibility and the absence of (A) common antigen (CPA), (C) O-specific antigen (OSA), or (E) an altered core (p > 0.05). CPA was absent in 42.73% of variants (B), OSA was absent in 76.36%, and (F) 27.27% of variants had altered LPS cores. Each data point represents the average Relative Susceptibility of a variant, and variants are colored based on their longitudinal collection.

Figure 4. No correlation between combined LPS deficiency phenotypes and R2-pyocin susceptibility. An LPS Score of 0 to 3 was assigned to each variant from each sputum collection to assess the correlation between overall LPS phenotypes and R2-pyocin susceptibility. Variants from each population are shown as data points of the same color. No correlation was found between LPS Score and R2-pyocin susceptibility (p > 0.05).

Figure S1. R-pyocin binding to LPS receptor and puncturing target cell membrane. When released into the environment by producing cells, R-pyocins bind to target cells by recognizing core residues of the lipopolysaccharide (LPS) on the outer cell membrane. After tail fiber binding, the baseplate dissociates, initiating sheath contraction and driving the iron-tipped tube through the cell surface, killing the target bacterium.

Figure S2. R2-pyocin lysate collection and susceptibility testing procedure. R-pyocins are isolated from *P. aeruginosa* by growing strains to early logarithmic growth, adding sub-minimal inhibitory concentration (MIC) ciprofloxacin, and lysing cells with chloroform after several hours for maximum R-pyocin production. Cultures are then centrifuged, separating into phases, allowing for the collection of cell-free R-pyocin-containing supernatant (lysates) to be stored. R-pyocin lysates are serially diluted and spotted onto soft agar overlays to test the susceptibility of strains of interest.

References

1.   Elborn JS. 2016. Cystic fibrosis. Lancet 388:2519-2531.
2.   Hogardt M, Heesemann J. 2010. Adaptation of Pseudomonas aeruginosa during persistence in the cystic fibrosis lung. Int J Med Microbiol 300:557-62.
3.   Kerem E, Corey M, Gold R, Levison H. 1990. Pulmonary function and clinical course in patients with cystic fibrosis after pulmonary colonization with Pseudomonas aeruginosa. J Pediatr 116:714-9.
4.   Boucher JC, Yu H, Mudd MH, Deretic V. 1997. Mucoid Pseudomonas aeruginosa in cystic fibrosis: characterization of muc mutations in clinical isolates and analysis of clearance in a mouse model of respiratory infection. Infect Immun 65:3838-46.
5.   Winstanley C, O'Brien S, Brockhurst MA. 2016. Pseudomonas aeruginosa Evolutionary Adaptation and Diversification in Cystic Fibrosis Chronic Lung Infections. Trends in Microbiology 24:327-337.
6.   Williams D, Evans B, Haldenby S, Walshaw MJ, Brockhurst MA, Winstanley C, Paterson S. 2015. Divergent, coexisting Pseudomonas aeruginosa lineages in chronic cystic fibrosis lung infections. American journal of respiratory and critical care medicine 191:775-785.
7.   Workentine M, Sibley C, Glezerson B, Purighalla S, Norgaard-Gron J, Parkins M, Rabin H, Surette M. 2013. Phenotypic Heterogeneity of Pseudomonas aeruginosa Populations in a Cystic Fibrosis Patient. PLoS One 8:e60225.
8.   Stover CK, Pham XQ, Erwin AL, Mizoguchi SD, Warrener P, Hickey MJ, Brinkman FSL, Hufnagle WO, Kowalik DJ, Lagrou M, Garber RL, Goltry L, Tolentino E, Westbrock-

Wadman S, Yuan Y, Brody LL, Coulter SN, Folger KR, Kas A, Larbig K, Lim R, Smith K, Spencer D, Wong GKS, Wu Z, Paulsen IT, Reizer J, Saier MH, Hancock REW, Lory S, Olson MV. 2000. Complete genome sequence of Pseudomonas aeruginosa PAO1, an opportunistic pathogen. Nature 406:959-964.

9. Foweraker JE, Laughton CR, Brown DFJ, Bilton D. 2005. Phenotypic variability of Pseudomonas aeruginosa in sputa from patients with acute infective exacerbation of cystic fibrosis and its impact on the validity of antimicrobial susceptibility testing. Journal of Antimicrobial Chemotherapy 55:921-927.

10. Bradbury R, Tristram S, Roddam L, Reid D, Inglis T, Champion A. 2011. Antimicrobial susceptibility testing of cystic fibrosis and non-cystic fibrosis clinical isolates of Pseudomonas aeruginosa: a comparison of three methods. British Journal of Biomedical Science 68:1-4.

11. Morita Y, Tomida J, Kawamura Y. 2014. Responses of Pseudomonas aeruginosa to antimicrobials. Frontiers in Microbiology 4.

12. Smith RS, Iglewski BH. 2003. P. aeruginosa quorum-sensing systems and virulence. Curr Opin Microbiol 6:56-60.

13. Diggle SP, Griffin A, Campbell GS, West S. 2007. Cooperation and conflict in quorum-sensing bacterial populations, vol 450.

14. Pesci EC, Pearson JP, Seed PC, Iglewski BH. 1997. Regulation of las and rhl quorum sensing in Pseudomonas aeruginosa. J Bacteriol 179:3127-32.

15. D'Costa VM, McGrann KM, Hughes DW, Wright GD. 2006. Sampling the Antibiotic Resistome. Science 311:374-377.

16. Cao P, Fleming D, Moustafa DA, Dolan SK, Szymanik KH, Redman WK, Ramos A, Diggle FL, Sullivan CS, Goldberg JB, Rumbaugh KP, Whiteley M. 2023. A Pseudomonas aeruginosa small RNA regulates chronic and acute infection. Nature 618:358-364.

17. Bhagirath AY, Li Y, Somayajula D, Dadashi M, Badr S, Duan K. 2016. Cystic fibrosis lung environment and Pseudomonas aeruginosa infection. BMC pulmonary medicine 16:174-174.

18. Jorth P, Ehsan Z, Rezayat A, Caldwell E, Pope C, Brewington JJ, Goss CH, Benscoter D, Clancy JP, Singh PK. 2019. Direct Lung Sampling Indicates That Established Pathogens Dominate Early Infections in Children with Cystic Fibrosis. Cell reports 27:1190-1204.e3.

19. Darch S, McNally A, Harrison F, Corander J, Barr H, Paszkiewicz K, Holden S, Fogarty A, Crusz S, Diggle S. 2015. Recombination is a key driver of genomic and phenotypic diversity in a Pseudomonas aeruginosa population during cystic fibrosis infection. Scientific Reports (Nature Publisher Group) 5:7649.

20. Williams D, Fothergill JL, Evans B, Caples J, Haldenby S, Walshaw MJ, Brockhurst MA, Winstanley C, Paterson S. 2018. Transmission and lineage displacement drive rapid population genomic flux in cystic fibrosis airway infections of a Pseudomonas aeruginosa epidemic strain. Microbial Genomics.

21. Feliziani S, Luján AM, Moyano AJ, Sola C, Bocco JL, Montanaro P, Canigia LF, Argaraña CE, Smania AM. 2010. Mucoidy, quorum sensing, mismatch repair and antibiotic resistance in Pseudomonas aeruginosa from cystic fibrosis chronic airways infections. PLoS One 5.

22. Cystic Fibrosis Foundation Patient Registry. 2019. Annual Data Report. Cystic Fibrosis Foundation.

23. Clark S, Diaz Caballero J, Cheang M, Coburn B, Wang P, Donaldson S, Zhang Y, Liu M, Keshavjee S, Yau Y, Waters V, Elizabeth Tullis D, Guttman D, Hwang D. 2015. Phenotypic diversity within a Pseudomonas aeruginosa population infecting an adult with cystic fibrosis. Scientific Reports (Nature Publisher Group) 5:10932.

24. Roberts AE, Kragh KN, Bjarnsholt T, Diggle SP. 2015. The Limitations of In Vitro Experimentation in Understanding Biofilms and Chronic Infection. J Mol Biol 427:3646-61.

511   25.   Bradley DE. 1967. Ultrastructure of bacteriophage and bacteriocins. Bacteriological
512   reviews 31:230-314.
513   26.   Reeves P. 1972. The bacteriocins (molecular biology, biochemistry and biophysics).
514   Spring-Verlag, New York.
515   27.   Kageyama M, Egami F. 1962. On the purification and some properties of a pyocin, a
516   bacteriocin produced by Pseudomonas aeruginosa. Life Sciences 1:471-476.
517   28.   Holloway BW. 1969. Genetics of Pseudomonas. Bacteriological reviews 33:419-443.
518   29.   Kageyama M. 1975. Bacteriocins and bacteriophages in Pseudomonas aeruginosa.
519   Microbial drug resistance:291-305.
520   30.   Michel-Briand Y, Baysse C. 2002. The pyocins of Pseudomonas aeruginosa. Biochimie
521   84:499-510.
522   31.   Scholl D. 2017. Phage Tail–Like Bacteriocins. Annual Review of Virology 4:453-467.
523   32.   Jacob F. 1954. Induced Biosynthesis and Mode of Action of a Pyocine, an Antibiotic
524   obtained from Ps. pyocyanea. Ann Inst Pasteur 86:149-60.
525   33.   Ito S, Kageyama M. 1970. Relationship between pyocins and a bacteriophage in
526   Pseudomonas aeruginosa. The Journal of General and Applied Microbiology 16:231-240.
527   34.   Govan JRW. 1974. Studies on the Pyocins of Pseudomonas aeruginosa: Production of
528   Contractile and Flexuous Pyocins in Pseudomonas aeruginosa. Microbiology 80:17-30.
529   35.   Ikeda K, Egami F. 1969. Receptor Substance for Pyocin R I. Partial Purification and
530   Chemical Properties. The Journal of Biochemistry 65:603-609.
531   36.   Ikeda K, Egami F. 1973. Lipopolysaccharide of Pseudomonas aeruginosa with special
532   reference to pyocin R receptor activity. The Journal of General and Applied Microbiology
533   19:115-128.
534   37.   Ikeda K, Nishi Y. 1973. Interaction between pyocin R and pyocin R receptor. The Journal
535   of General and Applied Microbiology 19:209-219.
536   38.   Köhler T, Donner V, van Delden C. 2010. Lipopolysaccharide as Shield and Receptor for
537   R-Pyocin-Mediated Killing in Pseudomonas aeruginosa. Journal of Bacteriology
538   192:1921-1928.
539   39.   Higerd TB, Baechler CA, Berk RS. 1969. Morphological Studies on Relaxed and
540   Contracted Forms of Purified Pyocin Particles. Journal of Bacteriology 98:1378-1389.
541   40.   Yui C. 1971. Structure of Pyocin R I. Isolation of Sheath from Pyocin R by Alkali Treatment
542   and Its Properties. The Journal of Biochemistry 69:101-110.
543   41.   Ohsumi M, Shinomiya T, Kageyama M. 1980. Comparative Study on R-type Pyocins of
544   Pseudomonas aeruginosa. The Journal of Biochemistry 87:1119-1126.
545   42.   Uratani Y, Hoshino T. 1984. Pyocin R1 inhibits active transport in Pseudomonas
546   aeruginosa and depolarizes membrane potential. J Bacteriol 157:632-6.
547   43.   Ge P, Scholl D, Leiman PG, Yu X, Miller JF, Zhou ZH. 2015. Atomic structures of a
548   bactericidal contractile nanotube in its pre- and postcontraction states. Nature Structural
549   & Molecular Biology 22:377-382.
550   44.   Ge P, Scholl D, Prokhorov NS, Avaylon J, Shneider MM, Browning C, Buth SA, Plattner
551   M, Chakraborty U, Ding K, Leiman PG, Miller JF, Zhou ZH. 2020. Action of a minimal
552   contractile bactericidal nanomachine. Nature 580:658-662.
553   45.   Nakayama K, Takashima K, Ishihara H, Shinomiya T, Kageyama M, Kanaya S, Ohnishi
554   M, Murata T, Mori H, Hayashi T. 2000. The R-type pyocin of Pseudomonas aeruginosa is
555   related to P2 phage, and the F-type is related to lambda phage. Mol Microbiol 38:213-31.
556   46.   Huszczynski SM, Lam JS, Khursigara CM. 2019. The Role of Pseudomonas aeruginosa
557   Lipopolysaccharide in Bacterial Pathogenesis and Physiology. Pathogens 9:6.
558   47.   Lam J, Taylor V, Islam S, Hao Y, Kocíncová D. 2011. Genetic and Functional Diversity of
559   Pseudomonas aeruginosa Lipopolysaccharide. Frontiers in Microbiology 2.

560   48.   Hao Y, King JD, Huszczynski S, Kocíncová D, Lam JS. 2013. Five new genes are
561   important for common polysaccharide antigen biosynthesis in Pseudomonas aeruginosa.
562   mBio 4:e00631-12.

563   49.   Anbo M, Jelsbak L. 2023. A bittersweet fate: detection of serotype switching in
564   Pseudomonas aeruginosa. Microb Genom 9.

565   50.   Carim S, Azadeh AL, Kazakov AE, Price MN, Walian PJ, Lui LM, Nielsen TN, Chakraborty
566   R, Deutschbauer AM, Mutalik VK, Arkin AP. 2021. Systematic discovery of pseudomonad
567   genetic factors involved in sensitivity to tailocins. Isme j 15:2289-2305.

568   51.   Mei M, Thomas J, Diggle SP. 2021. Heterogenous Susceptibility to R-Pyocins in
569   Populations of Pseudomonas aeruginosa Sourced from Cystic Fibrosis Lungs. mBio 12.

570   52.   Lerouge I, Vanderleyden J. 2002. O-antigen structural variation: mechanisms and possible
571   roles in animal/plant-microbe interactions. FEMS Microbiol Rev 26:17-47.

572   53.   Byrd MS, Sadovskaya I, Vinogradov E, Lu H, Sprinkle AB, Richardson SH, Ma L, Ralston
573   B, Parsek MR, Anderson EM, Lam JS, Wozniak DJ. 2009. Genetic and biochemical
574   analyses of the Pseudomonas aeruginosa Psl exopolysaccharide reveal overlapping roles
575   for polysaccharide synthesis enzymes in Psl and LPS production. Molecular Microbiology
576   73:622-638.

577   54.   King JD, Kocíncová D, Westman EL, Lam JS. 2009. Review: Lipopolysaccharide
578   biosynthesis in Pseudomonas aeruginosa. Innate Immunity 15:261-312.

579   55.   de Kievit TR, Lam JS. 1994. Monoclonal antibodies that distinguish inner core, outer core,
580   and lipid A regions of Pseudomonas aeruginosa lipopolysaccharide. J Bacteriol 176:7129-
581   39.

582   56.   Kintz E, Goldberg JB. 2008. Regulation of lipopolysaccharide O antigen expression in
583   Pseudomonas aeruginosa. Future Microbiology 3:191-203.

584   57.   Poon KKH, Westman EL, Vinogradov E, Jin S, Lam JS. 2008. Functional Characterization
585   of MigA and WapR: Putative Rhamnosyltransferases Involved in Outer Core
586   Oligosaccharide Biosynthesis of Pseudomonas aeruginosa. Journal of Bacteriology
587   190:1857-1865.

588   58.   Lau PCY, Lindhout T, Beveridge TJ, Dutcher JR, Lam JS. 2009. Differential
589   Lipopolysaccharide Core Capping Leads to Quantitative and Correlated Modifications of
590   Mechanical and Structural Properties in Pseudomonas aeruginosa Biofilms. Journal of
591   Bacteriology 191:6618-6631.

592   59.   Maldonado RF, Sá-Correia I, Valvano MA. 2016. Lipopolysaccharide modification in
593   Gram-negative bacteria during chronic infection. FEMS Microbiol Rev 40:480-93.

594   60.   Pier GB. 2007. Pseudomonas aeruginosa lipopolysaccharide: a major virulence factor,
595   initiator of inflammation and target for effective immunity. Int J Med Microbiol 297:277-95.

596   61.   Kocíncová D, Lam JS. 2011. Structural diversity of the core oligosaccharide domain of
597   Pseudomonas aeruginosa lipopolysaccharide. Biochemistry (Mosc) 76:755-60.

598   62.   Wang L, Wang Q, Reeves PR. 2010. The variation of O antigens in gram-negative
599   bacteria. Subcell Biochem 53:123-52.

600   63.   Qin S, Xiao W, Zhou C, Pu Q, Deng X, Lan L, Liang H, Song X, Wu M. 2022.
601   Pseudomonas aeruginosa: pathogenesis, virulence factors, antibiotic resistance,
602   interaction with host, technology advances and emerging therapeutics. Signal Transduct
603   Target Ther 7:199.

604   64.   Di Lorenzo F, Silipo A, Bianconi I, Lore NI, Scamporrino A, Sturiale L, Garozzo D, Lanzetta
605   R, Parrilli M, Bragonzi A, Molinaro A. 2015. Persistent cystic fibrosis isolate Pseudomonas
606   aeruginosa strain RP73 exhibits an under-acylated LPS structure responsible of its low
607   inflammatory activity. Mol Immunol 63:166-75.

608   65.   del Campo R, Martínez E, del Fresno C, Alenda R, Gómez-Piña V, Fernández-Ruíz I,
609   Siliceo M, Jurado T, Toledano V, Arnalich F, García-Río F, López-Collazo E. 2011.

610   Translocated LPS might cause endotoxin tolerance in circulating monocytes of cystic
611   fibrosis patients. PLoS One 6:e29577.
612 66. Cigana C, Curcurù L, Leone MR, Ieranò T, Lorè NI, Bianconi I, Silipo A, Cozzolino F,
613   Lanzetta R, Molinaro A, Bernardini ML, Bragonzi A. 2009. Pseudomonas aeruginosa
614   exploits lipid A and muropeptides modification as a strategy to lower innate immunity
615   during cystic fibrosis lung infection. PLoS One 4:e8439.
616 67. Ciofu O, Mandsberg LF, Bjarnsholt T, Wassermann T, Høiby N. 2010. Genetic adaptation
617   of Pseudomonas aeruginosa during chronic lung infection of patients with cystic fibrosis:
618   strong and weak mutators with heterogeneous genetic backgrounds emerge in mucA
619   and/or lasR mutants. Microbiology (Reading) 156:1108-1119.
620 68. Giordano NP, Cian MB, Dalebroux ZD. 2020. Outer Membrane Lipid Secretion and the
621   Innate Immune Response to Gram-Negative Bacteria. Infect Immun 88.
622 69. Tognon M, Köhler T, Gdaniec BG, Hao Y, Lam JS, Beaume M, Luscher A, Buckling A,
623   van Delden C. 2017. Co-evolution with Staphylococcus aureus leads to
624   lipopolysaccharide alterations in Pseudomonas aeruginosa. The ISME Journal 11:2233-
625   2243.
626 70. Redero M, López-Causapé C, Aznar J, Oliver A, Blázquez J, Prieto AI. 2018. Susceptibility
627   to R-pyocins of Pseudomonas aeruginosa clinical isolates from cystic fibrosis patients.
628   Journal of Antimicrobial Chemotherapy 73:2770-2776.
629 71. Redero M, Aznar J, Prieto AI. 2020. Antibacterial efficacy of R-type pyocins against
630   Pseudomonas aeruginosa on biofilms and in a murine model of acute lung infection. J
631   Antimicrob Chemother doi:10.1093/jac/dkaa121.
632 72. Secor PR, Sweere JM, Michaels LA, Malkovskiy AV, Lazzareschi D, Katznelson E,
633   Rajadas J, Birnbaum ME, Arrigoni A, Braun KR, Evanko SP, Stevens DA, Kaminsky W,
634   Singh PK, Parks WC, Bollyky PL. 2015. Filamentous Bacteriophage Promote Biofilm
635   Assembly and Function. Cell host & microbe 18:549-559.
636 73. Davis MR, Jr., Goldberg JB. 2012. Purification and visualization of lipopolysaccharide from
637   Gram-negative bacteria by hot aqueous-phenol extraction. J Vis Exp doi:10.3791/3916.
638 74. Jacobson KH, Gunsolus IL, Kuech TR, Troiano JM, Melby ES, Lohse SE, Hu D, Chrisler
639   WB, Murphy CJ, Orr G, Geiger FM, Haynes CL, Pedersen JA. 2015. Lipopolysaccharide
640   Density and Structure Govern the Extent and Distance of Nanoparticle Interaction with
641   Actual and Model Bacterial Outer Membranes. Environ Sci Technol 49:10642-50.
642 75. Snyder S, Kim D, McIntosh TJ. 1999. Lipopolysaccharide bilayer structure: effect of
643   chemotype, core mutations, divalent cations, and temperature. Biochemistry 38:10758-
644   67.
645 76. Clifton LA, Ciesielski F, Skoda MW, Paracini N, Holt SA, Lakey JH. 2016. The Effect of
646   Lipopolysaccharide Core Oligosaccharide Size on the Electrostatic Binding of
647   Antimicrobial Proteins to Models of the Gram Negative Bacterial Outer Membrane.
648   Langmuir 32:3485-94.
649 77. Choi K-H, Schweizer HP. 2006. mini-Tn7 insertion in bacteria with single attTn7 sites:
650   example Pseudomonas aeruginosa. Nature Protocols 1:153.
651 78. HOLLOWAY BW. 1955. Genetic Recombination in Pseudomonas aeruginosa.
652   Microbiology 13:572-581.
653 79. Hmelo LR, Borlee BR, Almblad H, Love ME, Randall TE, Tseng BS, Lin C, Irie Y, Storek
654   KM, Yang JJ, Siehnel RJ, Howell PL, Singh PK, Tolker-Nielsen T, Parsek MR, Schweizer
655   HP, Harrison JJ. 2015. Precision-engineering the Pseudomonas aeruginosa genome with
656   two-step allelic exchange. Nature Protocols 10:1820-1841.
657 80. Brazas MD, Hancock REW. 2005. Ciprofloxacin Induction of a Susceptibility Determinant
658   in Pseudomonas aeruginosa. Antimicrobial Agents and Chemotherapy 49:3222-3227.
659 81. Penterman J, Singh PK, Walker GC. 2014. Biological cost of pyocin production during the
660   SOS response in Pseudomonas aeruginosa. J Bacteriol 196:3351-9.

661   82.   Mei M, Pheng P, Kurzeja-Edwards D, Diggle SP. 2023. High prevalence of
662   lipopolysaccharide mutants and R2-Pyocin susceptible variants in Pseudomonas
663   aeruginosa populations sourced from cystic fibrosis lung infections. Dryad. Dataset.
664   https://datadryad.org/stash/share/k9quE9rIzJ-
665   gRXo42tF559U2MEpmXH6wC3emdpinkkM.

Figure 1

Figure 2

Figure 3

Figure 4

Figure S1

Figure S2

Step 1. Isolate R-pyocins

Grow strains to early/pre-log phase

Add Ciprofloxacin to 0.1μg/mL

Grow strains to mid-late log phase

Add chloroform to lyse cells

Spin and store cell-free lysate at 4°C

Step 2. Add R-pyocins to CF isolates

Grow indicator strains to mid-log phase

Inoculate 0.4% agar overlay at an OD of 0.01

Using cell-free lysate...

Serial dilute lysate (in PBS)

Spot 5μL lysates onto overlay

Incubate plates overnight

Count serial dilutions with inhibition