Characterization of an Australian outbreak Japanese Encephalitis virus genotype 4 isolate in

mice and human cortical brain organoids

Wilson Nguyen[a], Romal Stewart[a], Bing Tang[a], Kexin Yan[a], Thibaut Larcher[b], Cameron Bishop[a],

Narayan Gyawali[a], Gregor Devine[a], Andreas Suhrbier[a,c], Daniel J. Rawle[a*].

[a] QIMR Berghofer Medical Research Institute, Brisbane, Queensland 4029, Australia.

[b] INRAE, Oniris, PAnTher, APEX, Nantes, France.

[c] GVN Center of Excellence, Australian Infectious Disease Research Centre, Brisbane,

Queensland, 4029 and 4072, Australia.

[*]Corresponding author. Email: Daniel.Rawle@qimrberghofer.edu.au

A.S. D.J.R. joint last.

ABSTRACT

Human infections with Japanese encephalitis virus (JEV) are a leading cause of viral

encephalitis, with ≈70,000 symptomatic cases (≈40% with long-term neurological sequelae) and

≈20,000 deaths reported annually, primarily in Asia and the Western Pacific. An outbreak of

JEV genotype 4 was also recently reported in Australia, with an isolate ($JEV_{NSW/22}$) obtained

from a stillborn piglet. Herein we characterize the neuropathology of $JEV_{NSW/22}$, JEV_{FU}

(genotype 2) and $JEV_{Nakayama}$ (genotype 3) in adult C57BL/6J wild-type mice, mice deficient in

interferon regulatory factor 7 ($IRF7^{-/-}$), and mice deficient in the type I interferon receptor

($IFNAR^{-/-}$). In C57BL/6J and $IRF7^{-/-}$ mice with lethal outcomes, viral antigen was detected by

immunohistochemistry in *inter alia* the cortex, thalamus and hippocampus. In these mice,

histological lesions included neuronal degeneration, neuronal vacuolation, perivascular cuffing,

leukocyte infiltrates, hemorrhage, and microgliosis, with apoptosis and astrocyte activation

detected by immunohistochemistry. Microgliosis and hemorrhage persisted in some surviving

mice ≈3-4 weeks post infection. JEV was universally lethal in $IFNAR^{-/-}$ mice by day 3 with

histological signs of brain hemorrhage, but produced no other detectable brain infection or

lesions, with NS1 readily detected in blood vessels, but not neurons, by immunohistochemistry.

All JEV isolates showed robust productive cytopathic infection of human cortical brain

organoids (hBOs), as well as a human neural progenitor cell line (RENcell VM). We thus

describe a new $IRF7^{-/-}$ mouse model for JEV, which shows increased penetrance of lethal

neuroinvasiveness and recapitulates many aspects human disease. Although less virulent in

$IRF7^{-/-}$ mice, $JEV_{NSW/22}$ retained the capacity to cause lethal encephalitis and to replicate and

destroy human neuronal cells.

Introduction

Japanese Encephalitis virus (JEV) is a single-stranded positive-sense RNA virus from the *Flaviviridae* family that is transmitted from amplifying hosts (primarily pigs and wading birds) via mosquitoes to humans [1]. While infection is usually asymptomatic, encephalitis can develop in ≈1 in 250 people, with ~30% of encephalitic cases becoming fatal, and 30-50% of non-fatal encephalitic cases retaining persistent neurological symptoms including seizures, speech impediments and paralysis [2,3]. JEV is the leading cause of viral encephalitis in Asia, with ~70,000 cases and ~20,000 deaths per annum [4]. After a bite from an infected mosquito, JEV replicates in peripheral blood monocytes producing a viremia that, in some cases, leads to virus crossing the blood brain barrier [5]. JEV primarily infects neurons in the brain [6] leading to uncontrolled inflammation (encephalitis) and neuronal cell death [2]. There are several available and effective vaccines against JEV [7], but there are no specific licensed treatments.

Efforts to develop treatments for JEV neuropathology are hindered by the difficulties in diagnosing early infection [8] before the virus has infected central nervous system (CNS) neurons; and once the virus has infected CNS neurons, treatments must also be able to effectively enter the brain. In addition, mouse models of JEV neuropathology are often physiologically inappropriate, using intracranial routes of infection or very young mice [9-11]. Well-characterized, adult mouse models with high penetrance of JEV neuropathology after peripheral inoculation would represent useful tools to study mechanisms of infection and disease and evaluate potential new interventions.

JEV exists as five genotypes (G1-G5) which are phylogenetically, antigenically and geographically distinct [12-14]. G3 was the main genotype endemic in Asia until 1990, after which genotype I has dominated since [15]. G2 has been identified in Malaysia, Indonesia, Papua New Guinea and Australian outbreaks from 1970-2000. G5 has been identified in China, and is now dominant in South Korea [16]. G4 was the least common genotype worldwide, having only been identified in mosquitoes from Indonesia and Papua New Guinea [8]. In February 2021, a fatal JEV infection occurred on the Tiwi Islands, 80 km off the coast of Darwin, Australia [17]. Sequencing identified that the virus belonged to the rare JEV G4 [17]. In 2022, a geographically widespread outbreak throughout most Australian states was attributed to JEV G4, which caused 44 confirmed human cases and 7 deaths [18]. Outbreaks occurred in several piggeries, causing high abortion and stillbirth rates in sows [19]. Based on proximity to piggeries, estimates indicate that

~740,000 people may be at risk of being infected by JEV in Australia [19]. Given that *Culex*

annulirostris mosquitoes, considered the primary vector for JEV in Australia [18,19], and

amplifying vertebrate hosts including wading birds and pigs are widespread, it is possible that

JEV may become, or is already, endemic in Australia [18]. Murray Valley Encephalitis virus

(MVEV) and Kunjin virus are phylogenetically closely related to JEV and have similar vector

and reservoir hosts, and are endemic to Australia [20].

Herein, we characterize a new JEV mouse model using IRF7[-/-] mice, which shows high

penetrance of CNS infections, as well as neuropathology that recapitulates many aspects seen in

human post mortem brain specimens. We also describe for the first time the behavior of a recent

JEV G4 isolate from the Australian outbreak in mice and in human cortical brain organoids.

Materials and methods

Ethics statement and regulatory compliance

All mouse work was conducted in accordance with the "Australian code for the care and use of

animals for scientific purposes" as defined by the National Health and Medical Research Council

of Australia. Mouse work was approved by the QIMR Berghofer Medical Research Institute

Animal Ethics Committee (P3746, A2108-612). Mice were euthanized using CO_2 or cervical

dislocation.

Breeding and use of GM mice was approved under a Notifiable Low Risk Dealing (NLRD)

Identifier: NLRD_Suhrbier_Oct2020: NLRD 1.1(a). Use of Imojev was approved under

NLRD_Suhrbier_Oct2019: NLRD2.1(d).

All infectious JEV work was conducted in a dedicated suite in a biosafety level-3 (PC3) facility

at the QIMR Berghofer MRI (Australian Department of Agriculture, Water and the Environment

certification Q2326 and Office of the Gene Technology Regulator certification 3445). All work

was approved by the QIMR Berghofer Medical Research Institute Safety Committee (P3746).

Cell lines and culture

Vero E6 (C1008, ECACC, Wiltshire, England; obtained via Sigma Aldrich, St. Louis, MO,

USA) and C6/36 cells (ATCC# CRL-1660) were cultured in medium comprising RPMI 1640

(Gibco), supplemented with 10% fetal bovine serum (FBS), penicillin (100 IU/ml)/streptomycin

(100 μg/ml) (Gibco/Life Technologies) and L-glutamine (2 mM) (Life Technologies). Cells

were cultured at 37°C and 5% CO_2. Cells were routinely checked for mycoplasma (MycoAlert

Mycoplasma Detection Kit MycoAlert, Lonza) and FBS was assayed for endotoxin

contamination before purchase [21].

RENcell VM Human Neural Progenitor Cell Line (Sigma Aldrich) were cultured in medium

comprising DMEM F-12 (Thermo Fisher Scientific), penicillin (100 IU/ml)/streptomycin

(100 µg/ml) (Gibco/Life Technologies), 20 ng/ml FGF (STEMCELL Technologies), 20 ng/ml

EGF (STEMCELL Technologies), and 2% B27 supplements (Thermo Fisher Scientific). Cells

were detached using StemPro Accutase Cell Dissociation Reagent (Thermo Fisher Scientific),

and were cultured in Matrigel (Sigma). For crystal violet staining of remaining cells after

infection, formaldehyde (7.5% w/v)/crystal violet (0.05% w/v) was added to wells overnight,

plates washed twice in water, and plates dried overnight.

Virus isolates and culture

$JEV_{Nakayama}$ (GenBank: EF571853), JEV_{FU} (GenBank: AF217620), and Imojev were a gift from

Assoc. Prof. Gregor Devine (QIMR Berghofer MRI). $JEV_{NSW/22}$ (GenBank: OP904182) was a

gift from Dr. Peter Kirkland (Elizabeth Macarthur Agriculture Institute, New South Wales,

Australia). $MVEV_{TC123130}$ (GenBank: JN119814.1) was obtained from the "Doherty Virus

Collection" currently held at QIMR Berghofer MRI. Virus stocks were generated by infection of

C6/36 cells (for JEV and MVEV) or Vero E6 cells (for Imojev and YFV 17D) at multiplicity of

infection (MOI) ≈0.1, with supernatant collected after ~5 days, cell debris removed by

centrifugation at 3000 x g for 15 min at 4°C, and virus aliquoted and stored at -80°C. Virus titers

were determined using standard $CCID_{50}$ assays (see below).

$CCID_{50}$ assays

Vero E6 cells were plated into 96 well flat bottom plates at $2x10^4$ cells per well in 100 µl of

medium. For tissue titrations, tissues were homogenized in tubes each containing 4 ceramic

beads twice at 6000 x g for 15 seconds, followed by centrifugation twice at 21,000 x g for 5 min

before 5 fold serial dilutions in 100 µl RPMI 1640 supplemented with 2% FBS. For cell culture

supernatant or mouse serum, 10 fold serial dilutions were performed in 100 µl RPMI 1640

supplemented with 2% FBS. A volume of 100 µl of serially diluted samples were added to each

well of 96 well plate containing Vero E6 cells and the plates cultured for 5-7 days at 37°C and

5% CO_2. The virus titer was determined by the method of Spearman and Karber.

Mouse infections

C57BL/6J mice were purchased from the Animal Resources Center, Canning Vale WA,

Australia. IFNAR[-/-] were kindly provided by Dr P. Hertzog (Monash University, Melbourne,

Australia). IRF7[-/-] mice were kindly provided by T. Taniguchi (University of Tokyo) [22-24].

C57BL/6N mice were purchased from The Jackson Laboratory (stock no. 005304).

C57BL/6NJ[ΔNnt8-12] were generated by The Australian Phenomics Network, Monash University,

Melbourne, Australia as described [25]. Mice used in this study were female, except IRF7[-/-] mice

where both males and females were used. The age/age range at infection is indicated in the

figure legends. Mice were sorted into groups so that each group had a similar mean age and age

distribution, and in the case of IRF7[-/-] mice, had equal numbers of males and females in each

group. All mice strains except C57BL/6J were bred in-house at QIMR Berghofer MRI, and mice

housing conditions were described previously [26]. Mice were infected subcutaneously (s.c.) at the

base of the tail with 100 µl of virus inoculum. Serum was collected via tail nicks in to

Microvette Serum-Gel 500 µl blood collection tubes (Sarstedt, Numbrecht, Germany). Mice

were weighed and monitored for disease manifestations and were humanely euthanized using

CO_2 based on a score card system (Supplementary Figure 2B). At necropsy, brain and spleens

were collected for virus titrations by $CCID_{50}$ assays and/or for histology.

Histopathology and immunohistochemistry

Brains and spleens were fixed in 10% formalin and embedded in paraffin. Human cortical brain

organoids were embedded in agarose by adding 4% melted agarose and incubating on ice to

solidify, prior to standard paraffin embedding. Sections were stained with H&E (Sigma Aldrich)

and slides were scanned using Aperio AT Turbo (Aperio, Vista, CA USA) and analyzed using

Aperio ImageScope software (LeicaBiosystems, Mt Waverley, Australia) (v10).

For anti-flavivirus NS1 immunohistochemistry using 4G4, sections were affixed to positively

charged adhesive slides and air-dried overnight at 37°C. Sections were dewaxed and rehydrated

through xylol and descending graded alcohols to water. Sections were transferred to Dako

Epitope Retrieval Solution (pH 9.0) and subjected to heat antigen retrieval (100°C for 20 min)

using the Biocare Medical de-cloaking chamber, and slides allowed to cool for 20 minutes before

transferring to TBS plus 0.025% Tween-20 (TBS-T). Endogenous mouse Ig was blocked by

incubating sections with donkey anti-mouse Fab fragments (Jackson Immunoresearch) diluted

1:50 in Biocare Medical Rodent block M for 60 minutes. Sections were washed three times in

TBS-T, then incubated with anti-mouse Fc for 15 minutes, before a further three washes in TBS-

T. Nonspecific antibody binding was inhibited by incubation with Biocare Medical Background

Sniper with 1% BSA and 20% donkey serum and 20% goat serum for 15 minutes. Primary

antibody 4G4 (mouse anti-flavivirus NS1 [27-29]) was diluted 1 in 4 with Da Vinci Green diluent

and applied to the sections overnight at room temperature. Sections were washed three times in

TBS-T and endogenous peroxidase was blocked by incubating slides in Biocare Medical

Peroxidased 1 for 5 minutes. Sections were washed three times in TBS-T and Perkin Elmer Opal

HRP Polymer or Perkin Elmer Goat anti-mouse HRP diluted 1:500 in TBST was applied for 60

minutes. Sections were washed three times in TBS-T and signal developed in Vector Nova Red

for 5 minutes after which they were washed three times in water. Sections were lightly

counterstained in Haematoxylin (program 7 Leica Autostainer), washed in water, dehydrated

through ascending graded alcohols, cleared in xylene, and mounted using DePeX or similar.

Apoptosis was detected using the ApopTag Peroxidase In Situ Apoptosis Detection Kit

(Merck Catalogue No. S7100) as per manufacturer's instructions.

For anti-GFAP IHC, antigen retrieval was performed in 0.1M citric acid buffer (pH 6.0) at

1005°C for 20 min. Endogenous peroxidase activity was blocked using 1.0% H_2O_2 and 0.1%

sodium azide in TBS for 10 min. Endogenous mouse Ig was blocked by incubating sections with

goat anti-mouse Fab fragments (Jackson Immunoresearch) diluted 1:50 in Biocare Medical

Mouse block M for 60 minutes. Nonspecific antibody binding was inhibited by incubation with

Biocare Medical Background Sniper + 2% BSA for 30 min. Mouse anti-GFAP clone GA-5

(Biocare Medical, CM065C), was diluted 1:250 in the above buffer and incubated on sections for

1 hr at room temperature. After washes, Perkin Elmer Goat anti-mouse HRP diluted 1:500 in

TBST was applied for 60 minutes. Nova Red development and counter staining was performed

as for 4G4 above.

Infection of human cortical brain organoids

hBOs were reprogrammed from adult dermal fibroblasts (HDFa, Gibco, C0135C) using the

CytoTune-iPS 2.0 Sendai Reprogramming Kit (Invitrogen, A16518) [30], and were grown using

the CelVivo Clinostar incubator (Invitro Technologies) as described [31]. On the day of infection,

~30-day-old hBOs were transferred from each Clinoreactor into an ultra-low-binding 24 well

plate (one hBO per well), and each hBO was infected with 10^5 CCID$_{50}$ of either JEV$_{Nakayama}$,

JEV$_{FU}$, JEV$_{NSW/22}$, MVEV$_{TC123130}$, Imojev [32], or YFV 17D [29] for ~4 hrs. For shorter term culture

(up to 4 days for viral growth kinetics), virus inoculum was removed and hBOs were washed

twice with media in the well, before 1 ml differentiation media was added to each well and the

24 well plate was placed within a humidified tissue culture incubator at 37ºC, 5% CO$_2$ for up to 4

days. For longer term culture (up to 11 dpi), hBOs were washed twice with media by

transferring them to an ultra-low-binding 6 well plate containing 5 ml of media, and then hBOs

were transferred into 50 mm LUMOX gas exchange dishes (SARSTEDT) (4 organoids per dish)

containing 7 ml of differentiation media, and placed within a humidified tissue culture incubator

at 37ºC, 5% CO$_2$ for up to 11 days. hBOs were imaged using an EVOS FL (Advanced

Microscopy Group), and organoid area was determined using Image J v1.53 [33].

Statistics

Statistical analyses of experimental data were performed using IBM SPSS Statistics for

Windows, Version 19.0 (IBM Corp., Armonk, NY, USA). The t-test was used when the

difference in variances was <4, skewness was > -2 and kurtosis was <2. Otherwise, the non-

parametric Kolmogorov-Smirnov test was used. Kaplan-Meier statistics were determined by

log-rank (Mantel-Cox) test. Area under the curve analysis was performed in GraphPad Prism 8.

Results

JEV$_{NSW/22}$ infection produces a viremia but is not lethal in C57BL/6J mice

The JEV$_{NSW/22}$ G4 virus was isolated from the brain of a stillborn piglet in New South Wales

(NSW), Australia in February 2022 (GenBank accession OP904182). The JEV$_{Nakayama}$ G3

prototype was isolated in Japan in 1935 (GenBank accession EF571853), and the JEV$_{FU}$ G2 was

isolated in Australia in 1995 [34,35] (GenBank accession AF217620). MVEV was isolated in

Australia in 1974 [36] (MVEV$_{TC123130}$, GenBank accession JN119814). The latter three viruses

were isolated from human patients. Brief descriptions of the viral isolates and their phylogenic

relationships with other flaviviruses are provided in Supplementary Figure 1.

Six week old adult C57BL/6J mice were infected subcutaneously (s.c.) with 5×10^5 CCID$_{50}$ of

the aforementioned viruses. Viremia for all four viruses were broadly similar, peaking 1 day

post infection (dpi) at 2-3 $\log_{10}CCID_{50}$/ml of serum, with nearly all mice showing no detectable

viremia by day 4 (Figure 1A-D). Infection appeared to stall weight gain for most mice until day

~10 (Figure 1E). Between 8 to 12 dpi, four mice (2 infected with $JEV_{Nakayama}$, 1 JEV_{FU} and 1

$MVEV_{TC123130}$) out of the total of 24 mice showed weight loss $\geq$20% and were euthanized

(Figure 1E, †). A further four mice (1 $JEV_{Nakayama}$, 2 JEV_{FU}, 1 $MVEV_{TC123130}$) lost >5% of their

body weight, but subsequently recovered. None of the $JEV_{NSW/22}$ infected mice lost more than

~3% of their body weight (Figure 1E).

Kaplan-Meier survival curves provided no significant differences for the different viral

isolates (Figure 1F). Mice that were euthanized also displayed varying levels of abnormal

posture (hunching), reduced activity, and fur ruffling, on the day of euthanasia (Supplementary

Figure 2A). For the 4 mice that were euthanized (Figure 1E, †) viremias were not overtly higher

(Figure 1A-D, open symbols); however, these mice had very high levels of brain infection ($\approx$8-9

$\log_{10}CCID_{50}$/mg) (Figure 1G). At the time of euthanasia, viral titers in the spleen in these mice

were below the level of detection (Figure 1G), consistent with the viremia data.

C57BL/6J mice infected with JEV or MVEV led to viral neuroinvasion, apoptosis and reactive

astrogliosis

The brains of C57BL/6J mice were analysed by immunohistochemistry (IHC) using the pan-

flavivirus anti-NS1 antibody (4G4). Staining was consistently seen in the cortex for JEV and

MVEV infected mice, with staining also seen in the thalamus (Figure 2A, Supplementary Figure

3A-C). Prevailing infection of the cerebral cortex and thalamus parallels IHC data from post-

mortem human brains [6,37]. In some mice, virus was detected in other brain regions, including the

hippocampus, anterior olfactory nucleus (Figure 2A, $JEV_{Nakayama}$), and caudate putamen

(Supplementary Figure 3B, JEV_{FU}). A JEV_{FU} infected mouse that lost ~15% body weight and

then recovered (Figure 1E) had residual virus staining in the cortex at day 29 post infection

(Supplementary Figure 3D), while a mouse that survived infection with minimal weight loss had

no detectable viral antigen staining (Supplementary Figure 3E). As expected, 4G4 staining was

associated with cells showing neuronal morphology (Figure 2A, Supplementary Figure 3D).

ApopTag staining was evident in areas that were positive for virus staining, particularly in the

cortex, and was again associated with cells showing neuronal morphology (Figure 2B and

Supplementary Figure 4A-C). Apoptotic cells were also evident in the mouse that showed ~15%

body weight loss and recovered (Supplementary Figure 4D). Mice with signs of disease from

JEV or MVEV infection (Supplementary Figure 2) thus showed high levels of viral antigen and

apoptosis in the brain. In addition, like in humans, animals can recover from brain infection,

despite apoptotic damage, although neurological sequelae may ensue.

Astrocytes are a type of glial cell that provides physical and chemical support to neurons.

Astrocytes become activated (reactive astrogliosis) in response to brain pathology, including

from JEV infection [38-40]. Reactive astrocytes are characterized by the upregulated expression of

glial fibrillary acidic protein (GFAP) and the growth/extension of cellular processes [41]. Brains

from mice with signs of disease from JEV or MVEV infection (Supplementary Figure 2) had

significant upregulation of GFAP positive astrocytes throughout the brain, including the heavily

infected areas of the cortex, thalamus and anterior olfactory nucleus (Figure 2C, Supplementary

Figure 5). GFAP is constitutively expressed in astrocytes of the hippocampus, corpus callosum,

and cerebral peduncle [42], consistent with staining seen in uninfected brains (Figure 2D, right).

Widespread reactive astrogliosis is thus a feature of lethal JEV and MVEV encephalitis in

C57BL/6J mice.

IFNAR$^{-/-}$ mice are a model of lethal JEV viremia without neuroinvasion

As JEV$_{NSW/22}$ did not lead to fatal neuroinvasion in C57BL/6J mice, type I interferon receptor

knockout (IFNAR$^{-/-}$) mice were evaluated. IFNAR$^{-/-}$ mice have been used extensively in studies

of pathogenic flaviviruses such as Zika virus (ZIKV), West Nile virus (WNV), and Yellow fever

virus (YFV), and generally provide a robust viremia and a lethal outcome [28,29,43,44]. Adult

IFNAR$^{-/-}$ mice were infected with $5x10^5$ CCID$_{50}$ of JEV$_{Nakayama}$, JEV$_{FU}$, or JEV$_{NSW/22}$ via s.c.

injection. A robust viremia developed, reaching 7-8 log$_{10}$CCID$_{50}$/ml of serum by 2 dpi (Figure

3A). JEV$_{NSW/22}$ infected mice had ≈2 log lower viremia compared to JEV$_{Nakayama}$ and JEV$_{FU}$ on 3

dpi (Figure 3A). Mice displayed a rapid loss of body weight, which was slightly delayed for

JEV$_{NSW/22}$ at 2 dpi (Figure 3B), requiring euthanasia 2/3 dpi (Figure 3C). Mice infected with any

of the JEV strains also displayed varying levels of abnormal posture (hunching), reduced

activity, and fur ruffling (Supplementary Figure 6A).

Consistent with the robust viremia (Figure 3A), spleen tissue titers reached ~6-9

log$_{10}$CCID$_{50}$/g at 2-3 dpi (Figure 3D). Similar levels were seen in brains (Figure 3E); however,

4G4 staining illustrated that brain cells were not infected (Figure 3F). Viremia and brain titers

correlated strongly (Supplementary Figure 6B), arguing that the brain titers (Figure 3E) arose

from virus in the blood vessels of the brain. Interestingly, anti-flavivirus NS1 antibody staining

was clearly present in blood vessels (Figure 3G,H), with extracellular JEV NS1 also found in

serum of JEV patients [45]. NS1 from JEV is reported to increase vascular leakage and may

contribute to mortality [46], although overt vascular leakage was not evident in H&E sections of

brains at 2-3 dpi. Neither apoptosis (Supplementary Figure 6C), nor significantly increased

staining for reactive astrocytes (Supplementary Figure 5, IFNAR$^{-/-}$) were apparent for brains of

JEV-infected IFNAR$^{-/-}$ mice.

JEV$_{NSW/22}$ is less virulent than JEV$_{Nakayama}$ and JEV$_{FU}$ in IRF7$^{-/-}$ mice

C57BL/6J mice had a relatively low viremia and low penetrance of lethal brain infection (Figure

1), and IFNAR$^{-/-}$ mice had a rapid and lethal viremia without brain infection (Figure 3). A model

with a higher penetrance of brain infection is clearly desirable, given encephalitis is the key

clinical concern for JEV infections. Interferon regulatory factor 7 knockout (IRF7$^{-/-}$ mice) were

thus chosen as these mice produced an intermediate phenotype for WNV [47], as well as another

arbovirus [24]. As lower inoculation doses of JEV were reported to increase mortality [48], adult

C57BL/6J and IRF7$^{-/-}$ mice were inoculated s.c. with JEV$_{NSW/22}$ at $5x10^4$, $5x10^3$ and $5x10^2$

CCID$_{50}$. Reducing the dose of JEV$_{NSW/22}$ for C57BL/6J mouse infection (compared to the $5x10^5$

dose used in Figure 1) did not provide significant changes to weight change or viremia

(Supplementary Figure 7A, C-E). For JEV$_{NSW/22}$ infection in IRF7$^{-/-}$ mice, the inoculation dose

of $5x10^3$ CCID$_{50}$ provided a peak viremia 2 dpi that was marginally higher than the other doses

tested (Figure 4C, Supplementary Figure 7F-G). One mouse infected with this dose had weight

loss >20% requiring euthanasia on 8 dpi, whereas the $5x10^2$ and $5x10^4$ infection doses caused

weight loss no greater than ≈10% (Supplementary Figure 7B). An inoculation dose of $5x10^3$

CCID$_{50}$ was thus chosen for IRF7$^{-/-}$ mice.

JEV$_{NSW/22}$, JEV$_{Nakayama}$ and JEV$_{FU}$ infection of IRF7$^{-/-}$ mice (dose $5x10^3$ CCID$_{50}$) resulted in

robust viremias (Figure 4A-C), that were higher than those seen in C57BL/6J mice, where the

dose was $5x10^5$ CCID$_{50}$ (Figure 1A-C). Area under the curve analysis indicated no significant

difference between the viremias for the three JEV strains in IRF7$^{-/-}$ mice. MVEV$_{TC123130}$

infection produced a viremia in all IRF7$^{-/-}$ mice, although area under the curve analysis indicated

viremias were significantly lower than JEV (Figure 4D).

Infection of IRF7$^{-/-}$ mice with the different JEV isolates and MVEV (n=10 mice per group)

illustrated a mean weight loss of 3-12% for all four groups (Figure 4E), with the exception of 15

mice that reached ethically defined end points for weight loss and/or disease manifestations

(Supplementary Figure 8). Kaplan-Meier curves illustrated significantly higher survival for

IRF7$^{-/-}$ mice infected with JEV$_{NSW/22}$ when compared with JEV$_{Nakayama}$ and JEV$_{FU}$, with only 1/10

IRF7$^{-/-}$ mice infected with JEV$_{NSW/22}$ requiring euthanasia (Figure 4F). MVEV$_{TC123130}$ showed

no lethality in IRF7$^{-/-}$ mouse (Figure 4F), consistent with the low viremia (Figure 4D).

Brain infection in IRF7$^{-/-}$ mice infected with JEV

Brains titers in euthanized IRF7$^{-/-}$ mice were ~4-9 log$_{10}$CCID$_{50}$/gram, while spleen titers were

<2-5 log$_{10}$CCID$_{50}$/g (Figure 4G). IHC staining for viral antigen confirmed JEV infection in the

brains of IRF7$^{-/-}$ mice (Figure 4H), with reduced levels of staining reflecting lower brain titers

(Supplementary Figure 9). Viral antigen staining was consistently present in the cortex, with

staining also seen in the medulla, thalamus, and hippocampus (Figure 4H-I, Supplementary

Figure 9). Brain regions showing ApopTag staining broadly overlapped with staining for viral

antigen (Figure 4H-I). Reactive astrogliosis was also evident (Supplementary Figure 5).

Overall, this suggests that JEV infection of IRF7$^{-/-}$ mice produced a consistent viremia, with

mice showing brain infection and apoptotic damage.

JEV causes severe histological lesions in C57BL/6J and IRF7$^{-/-}$ mouse brains

H&E staining of JEV infected mouse brains revealed a series of histopathological lesions,

predominantly neuronal degeneration, neuronal vacuolation, perivascular cuffing, leukocyte

infiltrates, hemorrhage, meningitis and microgliosis (Figure 5A). IHC for the microglial marker,

Iba1, confirmed the presence microgliosis (Figure 5B).

The presence of H&E lesions were scored for the different JEV isolates and mouse strains

(Supplementary Figure 10). C57BL/6J and IRF7$^{-/-}$ mice euthanized during acute infections had a

broadly similar incidence of the aforementioned lesions, whereas only hemorrhage was seen in

IFNAR$^{-/-}$ mice (Supplementary Figure 10, Acute). Neuronal degeneration, neuronal vacuolation,

hemorrhage and microgliosis was not seen in the n=1 JEV$_{NSW/22}$ infected IRF7$^{-/-}$ mouse that was

euthanized 8 dpi (Supplementary Figure 10, Acute). For C57BL/6J and IRF7$^{-/-}$ mice that

survived and were euthanized later (nominally chronic phase), only hemorrhage was consistently

observed (Supplementary Figure 10, Chronic).

Lesions in C57BL/6J and IRF7[-/-] mouse brains were associated with areas of high virus

infection, most prominently in the cortex (Supplementary Figure 11); however, lesions were also

found in other areas of the brain including where there was minimal viral antigen staining

(Supplementary Figure 12A). Some mice that lost >8% body weight and then recovered had

persistent lesions detectable at the latest time point sampled (up to day 32 for C57BL/6J and day

19 for IRF7[-/-]) (Supplementary Figure 12B), consistent with persistent neurological sequelae in

humans that survive infection. Mice that did not lose any body weight did not show any overt

brain lesions at the later time points, indicating that brain infection causing weight loss is

associated with persistent lesions.

Overall, the lesions in C57BL/6J and IRF7[-/-] mouse brains are consistent with H&E detectable

lesions in post-mortem human JEV infected brains [6,38,49-51].

Publically available RNA-Seq data of mouse brains infected with JEV

A publically available RNA-Seq data (raw read counts) of C57BL/6 brain infection with JEV

isolate P20778 (Vellore strain, genotype 3)[52] (GEO accession: GSE154002) was reanalyzed [53] to

provide a list of differentially expressed genes (DEGs) for infected brains versus uninfected

brains from C57BL/6 mice. GFAP was significantly up-regulated 5.4 fold by infection

(Supplementary Table 2, $\log_2$FC 2.4), consistent with the GFAP IHC (Figure 2C) and reactive

astrogliosis seen herein. Iba1 (Aif1) was also significantly up-regulated 5.5 fold (Supplementary

Table 2, $\log_2$FC 2.5), consistent with the H&E and IHC (Figure 5) and microgliosis observed in

our study. To provide potential insights into the nature of the cellular infiltrates (e.g.

perivascular cuffing and leukocyte infiltrates, Figure 5), immune cell type abundance estimates

were obtained from the aforementioned RNA-Seq expression data using SpatialDecon [54]. The

dominant cell types identified were CD4 T cells, NKT cells, monocytes/macrophages,

neutrophils, innate lymphoid cells, T regs, and CD8 T cells, with an increase in microglia cells

also identified (Supplementary Figure 13).

Productive replication of JEV in human cortical brain organoids

Human cortical brain organoids (hBOs) have been used to investigate the pathogenesis of

flaviviruses, including JEV and ZIKV [44,55,56]. We generated ~30 day old hBO from HDFa cells,

growing the "minibrains" in a CelVivo Clinostar incubator as described [31] (Figure 6A). hBOs

were infected with 10^5 $CCID_{50}$ of the indicated JEV and MVEV isolates, as well as (i) the Imojev

chimeric virus vaccine (previously called ChimeriVax-JE) comprising the prME genes of JEV on

the YFV 17D backbone [32,57] and (ii) the Yellow Fever live attenuated vaccine strain (YFV 17D)

[29], with wild-type YFV infection [58], and occasionally YFV 17D vaccination [59], able to cause

neuropathology. hBOs were fixed in formalin 4 dpi and IHC undertaken using the anti-NS1

monoclonal antibody 4G4 [29]. $JEV_{Nakayama}$ infected hBOs showed the most pronounced viral

antigen staining, with $JEV_{NSW/22}$, JEV_{FU} and $MVEV_{TC123130}$ also showing clear staining (Figure

6B). Viral antigen was primarily localized to the outer surface of the organoids (Figure 6B),

where cells are in direct contact with the culture medium. Viral antigen staining for Imojev and

YFV 17D infected hBOs was also seen, but was sparse (Figure 6B and Supplementary Figure

14A).

All viruses were able to productively infect hBOs, with $JEV_{Nakayama}$ infection generating

higher viral titers than Imojev at all time points (Figure 6C, $p<0.001$). By 96 h $JEV_{Nakayama}$ titers

were also significantly higher than those seen for $JEV_{NSW/22}$ (Figure 6C).

Over 11 days, uninfected hBOs grew in size by ~20% (Figure 6D, Uninfected). hBOs

infected with JEV isolates shrank in cross sectional area by ~5% to 15% by 11 dpi, although

when compared with uninfected organoids, this only approached significance for $JEV_{NSW/22}$

(Figure 6D; Supplementary Figure 14B-C). hBOs infected with $MVEV_{TC123130}$, Imojev and YFV

17D infected hBO did not shrink significantly when compared with uninfected controls, although

the $MVEV_{TC123130}$ data suggested a marginal reduction in area (Figure 6C; Supplementary Figure

14C). These data (Figure 6D) mirror viral replication data (Figure 6C) and likely reflect virus-

induced CPE.

All viruses replicated productively in a human neural progenitor cell line, RENcell VM, with

all viruses except Imojev causing fulminant CPE (Supplementary Figure 15A,B). Although

Vero E6 cells are widely used for flavivirus research, CPE induced by $JEV_{NSW/22}$ was

considerably less pronounced in these cells (Supplementary Figure 15C).

Discussion

Herein we provide a comprehensive *in vivo* and *in vitro* characterization of the genotype 4

JEV$_{NSW/22}$ isolate from the recent Australian outbreak, and illustrate mouse models of infection and CNS neuropathological manifestations that recapitulate many aspects of human and primates disease [6,37-40,60]. Like the proptypical JEV$_{FU}$ (G2) and JEV$_{Nakayama}$ (G3) isolates, JEV$_{NSW/22}$ was able productively to infect hBOs, and in IRF7$^{-/-}$ mice, retained the capacity for neuroinvasive infection, generation of severe histopathological lesions and induction of mortality, consistent with reported human fatalities in the 2022 Australian outbreak.

Serosurvey data, albeit limited, suggests the ratio of human symptomatic to asymptomatic infections is not particularly different for JEV$_{NSW/22}$. To date 45 clinical cases have been notified for the recent outbreak in Australia [61], with serosurveys in Victoria (n=820 participants) and New South Wales (NSW) (n=917 participants) reporting 3.3% and 8.7% of participants as seropositive for JEV, respectively [62,63]. Although somewhat dated, population data for the primary recruitment locations for the serosurveys is available from Australian Bureau of Statistics 2016, with Victorian recruitment locations providing a population total of 160,294 (Mildura, Lockington, Shepparton, Cobram, Yarrawonga, Rutherglen, Wodonga, Wangaratta, Rochester), and NSW locations a total of 68,431 (Balranald, Corowa, Dubbo, Griffith, Temora). As [0.033 x 160,294] + [0.087 x 68,431] / 250 = 45, the serosurvey data is consistent with the expected symptomatic to asymptomatic ratio of ≈1 in 250 for JEV and thus provides no compelling evidence for virulence differences for JEV$_{NSW/22}$ in human populations.

Mouse models of flavivirus pathogenesis frequently use IFNAR$^{-/-}$ mice [29,64,65] as flaviviruses often replicate poorly in wild-type mice as the ability of NS5 to suppress the antiviral type I IFN responses in humans often fails to operate in mice [66]. We show herein that IFNAR$^{-/-}$ mice are not a good model for JEV neuropathology, as they reach ethically defined endpoints before brain infection can occur. However IFNAR$^{-/-}$ mice are a good model of robust and lethal viremia, which may provide a useful and stringent model for vaccine testing. IRF7$^{-/-}$ mice do provide a new and useful model for JEV as they show high penetrance of brain infection and neuropathology after s.c. inoculation of JEV. IRF7$^{-/-}$ retain the ability to induce some type I IFN responses (primarily via IRF3) [24] that likely prevent early mortality, thereby allowing subsequent brain infection. JEV$_{NSW/22}$ showed lower levels of mortality than JEV$_{FU}$ and JEV$_{Nakayama}$ in IRF7$^{-/-}$ mice (Figure 4F). JEV$_{NSW/22}$ also showed less pronounced induction of CPE in Vero E6 cells (Supplementary Figure 15C), another system in which type I interferon responses are diminished. Vero E6 cells retain the ability to respond to IFNα/β, but cannot make type I IFNs

[67]. These results suggest $JEV_{NSW/22}$ is more sensitive to, or less able to suppress, type I IFN

responses. Inhibition of type I IFN responses is mediated by NS5 for inhibition of STAT2 and

NS3 [68] and subgenomic flavivirus RNA (sfRNA)/NS5 [44] for inhibition of STAT1. Although the

latter likely operates for WNV in mice [69] and is involved in promoting apoptosis [44], the

efficiency of these systems during JEV infection of humans and mice remains to be determined.

sfRNA is derived from the 3'UTR [44], where $JEV_{NSW/22}$ does show a small number of nucleic acid

changes that are unlikely to affect sfRNA production (Supplementary Table 1).

$JEV_{NSW/22}$ does not contain known attenuating mutations that would explain its reduced

virulence in IRF7[-/-] mice. There are 9 amino acids in the Imojev vaccine envelope gene that have

been associated with attenuated neurovirulence; F107, K138, V176, A177, G244, H264, M279,

V315 and R439 [70-80]. At these positions, $JEV_{Nakayama}$, JEV_{FU}, $JEV_{NSW/22}$, and $MVEV_{TC123130}$ all

have the same amino acids (L107, E138, T177, E244, Q264, K279, A315, and K439), except for

position 176 where $JEV_{Nakayama}$ has T176, and JEV_{FU}, $JEV_{NSW/22}$, and $MVEV_{TC123130}$ have I176

(Supplementary Table 1). $JEV_{NSW/22}$ retains E at position 138, and this amino acid has been

identified by several studies as a principal neurovirulence determinant [81-85], with a role in

neuronal cell binding hypothesized [86]. YFV 17D has a valine (V) at this residue, possibly

contributing to reduced neurovirulence [87]. NS1 and NS2A have been implicated in

neuroinvasion, but not neurovirulence [79]. Among the six changes in NS1 associated with

attenuation of neuroinvasiveness are R147H and R339M [79], of which H147 is present in both

$JEV_{Nakayama}$ and $JEV_{NSW/22}$, with K339 (a conserved substitution for R) present in $JEV_{NSW/22}$.

JEV non-structural protein 4B (NS4B) alone can induce apoptosis and encephalitis [88], however,

NS4B is completely conserved between $JEV_{Nakayama}$, JEV_{FU}, and $JEV_{NSW/22}$ (Supplementary

Table 1). prM has been reported to influence neuroinvasiveness of ZIKV [64] and JEV virulence

in mice [89]. $JEV_{NSW/22}$ has a number of unique changes in prM (Supplementary Table 1), although

their functional implications remain unclear. $JEV_{NSW/22}$ also has an additional N-linked

glycosylation site at position 175 in NS1 that is lacking in $JEV_{Nakayama}$ and JEV_{FU}

(Supplementary Table 1). However, this N-linked glycosylation site is reported to increase

WNV neuroinvasiveness in wild-type mice [90,91], a trend not seen in our JEV data (Figure 1F).

Thus $JEV_{NSW/22}$ shows no obvious sequence characteristics that can be ready associated with

reduced virulence.

Limitations of this study include the use of $JEV_{NSW/22}$, a pig isolate; however, there are only 4

amino acid differences between this isolate and a JEV G4 sequence from a human brain in the

Tiwi Islands (Northern Territory, Australia) in 2021 (Genbank accession OM867669} [17]). The

differences are; envelope-238 F vs. L, NS2A-71 I vs. T, NS2B-59 E vs. G, NS3-436 E vs. G. In

addition, $JEV_{Nakayama}$ was passaged in suckling mouse brains, which may contribute to the

increased neuroinvasion seen herein, although the adaptive mutations acquired during passaging,

if any, are currently unknown. Furthermore, JEV_{FU} has not been passaged in mice, but was still

more lethal than $JEV_{Nakayama}$ in IRF7$^{-/-}$ mice. Use of C57BL/6J mice that lack a functional

nicotinamide nucleotide transhydrogenase *(Nnt)* may be another issue, as background and *Nnt*

are able to affect viral immunopathogenesis [25,53]. However, we found that neither *Nnt* nor a

C57BL/6N genetic background significantly impacted JEV replication or immunopathology

(Supplementary Figure 16).

In conclusion, we show that $JEV_{NSW/22}$ retains capacity for robust replication in neurons, and

is capable of neuroinvasive infection, severe histopathological lesions and mortality in IRF7$^{-/-}$

mice, consistent with reported human fatalities in the 2022 Australian outbreak.

ACKNOWLEDGEMENTS

From QIMR Berghofer MRI we thank Dr I Anraku for managing the PC3 (BSL3) facility,

animal house staff for mouse breeding and agistment, Drs Anthony White and Lotta Oikari for

providing the iPSC line, Crystal Chang, Sang-Hee Park andAshwini Potadar for the histology

and immunohistochemistry, Dr Viviana Lutzky for proof reading. We thank Dr. Peter Kirkland

for supplying the $JEV_{NSW/22}$ isolate. We thank Dr. Andrii Slonchak for insights into whether

3'UTR changes may affect sfRNA production.

DATA AVAILABILITY

All data is provided in the manuscript and accompanying supplementary files.

FUNDING

A.S. holds an Investigator grant from the National Health and Medical Research Council

(NHMRC) of Australia (APP1173880). We also acknowledge the intramural grant from QIMR

Berghofer awarded to R.S. and D.J.R. for purchase of the CelVivo Clinostar incubator.

AUTHOR CONTRIBUTIONS

Conceptualization, D.J.R. Methodology, D.J.R., W.N., and A.S. Formal analysis, W.N., D.J.R.

and A.S. Investigation, W.N., B.T., R.S., K.Y., N.G., T.L., and D.J.R. Resources, R.S., N.G.,

G.D., and A.S. Data curation, W.N., D.J.R., and A.S. Writing, original draft – D.J.R. Writing,

review and editing, A.S and W.N. Visualization, D.J.R., W.N. and A.S. Supervision, D.J.R.,

G.D. and A.S. Project administration, D.J.R., G.D. and A.S. Funding acquisition, D.J.R, G.D.

and A.S.

DECLARATION OF INTERESTS

The authors declare no competing interests.

FIGURE LEGENDS

Figure 1. JEV and MVEV infection in C57BL/6J mice. (A-D) Female ≈6 week old C57BL/6J

mice were infected s.c. with 5×10^5 $CCID_{50}$ of the indicated viruses. Viremia for each of the n=6

per group over 6 days is shown. Horizontal lines show mean viremia $\pm$ SE. All mice recorded a

detectable viremia for at least one timepoint. Limit of detection is 2 $log_{10}CCID_{50}$/ml of serum.

Open symbols indicate the viremia of mice that were euthanized due to weight loss reaching the

ethically defined endpoint of $\geq$20%. (E) Percent body weight change of individual mice

compared to each mouse's weight on day zero. Four mice lost $\geq$20% body weight and were

euthanized (†). (F) Kaplan-Meier plot showing percent survival (n=6 for each virus/virus

isolate). (G) Viral tissue titers in brains and spleens of the four euthanized mice (n=2

$JEV_{Nakayama}$, n=1 JEV_{FU}, n=1 $MVEV_{TC123130}$). Tissue titers determined by $CCID_{50}$ assay (limit of

detection ~2 $log_{10}CCID_{50}$/g).

Figure 2. IHC for viral antigen, apoptosis and reactive astrocytes in JEV infected

C57BL/6J mouse brain. IHC of $JEV_{Nakayama}$ infected brain (required euthanasia on 9 dpi, brain

virus titer 9.2 $log_{10}CCID_{50}$/g). (A) Staining for flavivirus NS1 using the 4G4 monoclonal

antibody. High magnification images show cells with neuronal morphology in the cortex (left)

and hippocampus (right). The latter also shows staining of dendrites and axons (fibrilar patterns

above and densely staining cells). 4G4 staining of brains of the other mice requiring euthanasia

(marked by † in Figure 1E) is shown in Supplementary Figure 3. (B) ApopTag staining of the

same brain as in 'A'. High magnification images show cells with neuronal morphology in the

cortex (left) and apoptotic cells in the dentate gyrus (right). ApopTag staining of the brains of

the other mice requiring euthanasia is shown in Supplementary Figure 4. (C) Staining for GFAP,

a marker of reactive astrocytes, for the same brain as in 'A'. High magnification images show a

typical reactive astrocyte in the cortex (left) and reactive astrocytes in the thalamus (right).

GFAP IHC for the brains of the other mice requiring euthanasia is shown in Supplementary

Figure 5. (D) IHC negative controls; uninfected mouse brain stained with 4G4 (left), ApopTag

(middle), and anti-GFAP (right).

Figure 3. JEV lethal viremia in IFNAR$^{-/-}$ mice. Female IFNAR$^{-/-}$ mice (9-24 week old) were

infected s.c. with 5×10^5 CCID$_{50}$ of the indicated virus (n=4 per group). (A) Mean viremia

determined by CCID$_{50}$ assay (limit of detection 2 log$_{10}$CCID$_{50}$/ml). Statistics by t-tests. (B)

Mean percent body weight change compared to 0 dpi. Statistics 2 dpi JEV$_{NSW/22}$ versus

JEV$_{Nakayama}$ (Kolmogorov-Smirnov test, p=0.023) and for JEV$_{NSW/22}$ versus JEV$_{FU}$ (t-test,

p=0.008). (C) Kaplan-Meier plot showing percent survival. (D) Viral tissue titers in spleens

harvested at euthanasia (3 dpi for all mice except for 1 JEV$_{FU}$ mouse at 2 dpi), determined by

CCID$_{50}$ assay (limit of detection ~2 log$_{10}$ CCID$_{50}$/g). (E) Viral titers in brains. (F) IHC staining

for flavivirus NS1 using the 4G4 monoclonal antibody. (G, H) High magnification images from

F showing NS1 staining in blood vessels surrounding the pale grey, biconcave shaped, red blood

cells.

Figure 4. JEV neuroinvasive infection in IRF7$^{-/-}$ mice. IRF7$^{-/-}$ mice (15-48 weeks old) were

infected s.c. with 5×10^3 CCID$_{50}$ of the indicated virus. (A-D) Viremias for indicated virus

determined by CCID$_{50}$ assay; limit of detection is 2 log$_{10}$CCID$_{50}$/ml. Horizontal line indicates

mean at each time point, and error bars represent SE; n=4 mice per group. Unfilled symbols

represent data for mice that were euthanized due to acute disease manifestations (Supplementary

Figure 8). (E) Percent body weight change compared to 0 dpi. Four mice lost >20% body

weight were euthanized (marked by †) and are plotted individually. The weight change for the

remaining mice are shown as means $\pm$ SE. Data is from two independent experiments, n=10

mice per group. (F) Kaplan-Meier plot showing percent survival (n=10 per group, data from two

independent experiments). Statistics by log-rank (Mantel-Cox) tests. Symbols as for E. (G)

Brain and spleen tissue titers for 8 euthanized mice (n=3 JEV$_{Nakayama}$, n=4 JEV$_{FU}$, n=1

JEV$_{NSW/22}$). Tissue titers determined by CCID$_{50}$ assay (limit of detection ~2 log$_{10}$CCID$_{50}$/g). (H)

IHC of JEV$_{NSW/22}$ infected brain (euthanized on day 8, brain virus titer 6.3 log$_{10}$CCID$_{50}$/g) using

4G4 monoclonal antibody (top) or ApopTag (bottom). High magnification images of cortex

(left) and medulla (right). (I) IHC using 4G4 (left) and ApopTag IHC (right) for the JEV$_{Nakayama}$

infected brain with the highest virus titer (euthanized on day 7, brain virus titer 8.3

log$_{10}$CCID$_{50}$/g). IHC using 4G4 for the brains of two other euthanized mice are shown in

Supplementary Figure 9.

Figure 5. Histological lesions in brains of JEV infected mice. (A) Examples of indicated

lesions stained with H&E. Degeneration of neurons indicated by (i) pyknotic nuclei (black

unfilled arrow heads indicating condensation and fragmentation of nuclei, staining dark blue),

and (ii) hyper-eosinophilic cytoplasm of degenerating neurons in the cornus ammonis of the

hippocampus (area also staining for viral antigen Supplementary Figure 11). Neuronal

vacuolation indicated by fluid accumulation around the neurons (arrowheads). Perivascular

cuffing is indicated by leukocytes aggregating in blood vessels (arrowheads). Leukocyte

infiltrates (extravascular) are indicated by dashed ovals. Hemorrhagic lesions are indicated by

extravascular red blood cells (arrows). Microgliosis is indicated by accumulation of microglia,

which have an elongated rod-shaped nuclei (arrows). Meningitis is indicated by accumulation of

leukocytes around the meninges (arrowheads). Images of uninfected controls accompany each

image(s) of lesions. (B) IHC using anti-Iba1, a microglial marker.

Figure 6. Infection of human cortical brain organoids (hBOs). (A) Photograph of "mini-

brains" cultured in a rotating CelVivo Clinostar incubator. (B) IHC of viral antigen (4G4) for

hBOs at 4 dpi. Images are representative of n=4 hBOs for each group. Magnified images of

sparse Imojev and YFV 17D infected cells are shown in Supplementary Figure 13A. (B) Viral

growth kinetics determined by CCID$_{50}$ assays of culture supernatants at the indicated hours post

infection; limit of detection is 2 log$_{10}$CCID$_{50}$/ml. At all time points JEV$_{Nakayama}$ vs. Imojev, and

at 96 h JEV$_{Nakayama}$ vs. JEV$_{NSW/22}$ were significant (t tests, n=5 organoids per group). (C) Mean

percentage change in organoid area 11 dpi vs. 9 dpi for each organoid (n=8 for uninfected and

MVEV$_{TC123130}$, otherwise n=4). Statistics are by Kolmogorov-Smirnov test for uninfected versus

$JEV_{Nakayama}$, and t-test for uninfected versus JEV_{FU} or $JEV_{NSW/22}$.

References

607   1   van den Hurk, A. F., Ritchie, S. A. & Mackenzie, J. S. Ecology and Geographical Expansion of
608   Japanese Encephalitis Virus. *Annual Review of Entomology* **54**, 17-35,
609   doi:10.1146/annurev.ento.54.110807.090510 (2008).
610   2   Ashraf, U. *et al.* Pathogenicity and virulence of Japanese encephalitis virus: Neuroinflammation
611   and neuronal cell damage. *Virulence* **12**, 968-980, doi:10.1080/21505594.2021.1899674 (2021).
612   3   Solomon, T. *et al.* Japanese encephalitis. *Journal of Neurology, Neurosurgery &*
613   *Psychiatry* **68**, 405, doi:10.1136/jnnp.68.4.405 (2000).
614   4   Quan, T. M., Thao, T. T. N., Duy, N. M., Nhat, T. M. & Clapham, H. Estimates of the global burden
615   of Japanese encephalitis and the impact of vaccination from 2000-2015. *eLife* **9**, e51027,
616   doi:10.7554/eLife.51027 (2020).
617   5   Hsieh, J. T., Rathore, A. P. S., Soundarajan, G. & St John, A. L. Japanese encephalitis virus
618   neuropenetrance is driven by mast cell chymase. *Nat Commun* **10**, 706, doi:10.1038/s41467-
619   019-08641-z (2019).
620   6   Johnson, R. T. *et al.* Japanese encephalitis: immunocytochemical studies of viral antigen and
621   inflammatory cells in fatal cases. *Ann Neurol* **18**, 567-573, doi:10.1002/ana.410180510 (1985).
622   7   Hegde, N. R. & Gore, M. M. Japanese encephalitis vaccines: Immunogenicity, protective efficacy,
623   effectiveness, and impact on the burden of disease. *Hum Vaccin Immunother* **13**, 1-18,
624   doi:10.1080/21645515.2017.1285472 (2017).
625   8   Pham, D. *et al.* Emergence of Japanese encephalitis in Australia: a diagnostic perspective.
626   *Pathology* **54**, 669-677, doi:10.1016/j.pathol.2022.07.001 (2022).
627   9   Bharucha, T. *et al.* Mouse models of Japanese encephalitis virus infection: A systematic review
628   and meta-analysis using a meta-regression approach. *PLOS Neglected Tropical Diseases* **16**,
629   e0010116, doi:10.1371/journal.pntd.0010116 (2022).
630   10   Beasley, D. W. C. *et al.* Protection against Japanese encephalitis virus strains representing four
631   genotypes by passive transfer of sera raised against ChimeriVax™-JE experimental vaccine.
632   *Vaccine* **22**, 3722-3726, doi:https://doi.org/10.1016/j.vaccine.2004.03.027 (2004).
633   11   Tajima, S. *et al.* Growth, Pathogenesis, and Serological Characteristics of the Japanese
634   Encephalitis Virus Genotype IV Recent Strain 19CxBa-83-Cv. *Viruses* **15**, 239 (2023).
635   12   Gao, X. *et al.* Changing Geographic Distribution of Japanese Encephalitis Virus Genotypes, 1935-
636   2017. *Vector Borne Zoonotic Dis* **19**, 35-44, doi:10.1089/vbz.2018.2291 (2019).
637   13   Kang, B. K. *et al.* Comparison of the antigenic relationship between Japanese encephalitis virus
638   genotypes 1 and 3. *Clin Exp Vaccine Res* **5**, 26-30, doi:10.7774/cevr.2016.5.1.26 (2016).
639   14   Han, N. *et al.* Comparison of genotypes I and III in Japanese encephalitis virus reveals distinct
640   differences in their genetic and host diversity. *J Virol* **88**, 11469-11479, doi:10.1128/jvi.02050-14
641   (2014).
642   15   Mulvey, P. *et al.* The Ecology and Evolution of Japanese Encephalitis Virus. *Pathogens* **10**,
643   doi:10.3390/pathogens10121534 (2021).
644   16   Lee, A. R., Song, J. M. & Seo, S. U. Emerging Japanese Encephalitis Virus Genotype V in Republic
645   of Korea. *J Microbiol Biotechnol* **32**, 955-959, doi:10.4014/jmb.2207.07002 (2022).
646   17   Sikazwe, C. *et al.* Molecular detection and characterisation of the first Japanese encephalitis
647   virus belonging to genotype IV acquired in Australia. *PLoS Negl Trop Dis* **16**, e0010754,
648   doi:10.1371/journal.pntd.0010754 (2022).

18   Mackenzie, J. S., Williams, D. T., van den Hurk, A. F., Smith, D. W. & Currie, B. J. Japanese
Encephalitis Virus: The Emergence of Genotype IV in Australia and Its Potential Endemicity.
Viruses **14** (2022).
19   Yakob, L. *et al.* Japanese encephalitis emergence in Australia: the potential population at risk.
Clin Infect Dis, doi:10.1093/cid/ciac794 (2022).
20   Gyawali, N. *et al.* Neglected Australian Arboviruses Associated With Undifferentiated Febrile
Illnesses. *Front Microbiol* **10**, 2818, doi:10.3389/fmicb.2019.02818 (2019).
21   Johnson, B. J. *et al.* Heat shock protein 10 inhibits lipopolysaccharide-induced inflammatory
mediator production. *J Biol Chem* **280**, 4037-4047, doi:10.1074/jbc.M411569200 (2005).
22   Honda, K. *et al.* IRF-7 is the master regulator of type-I interferon-dependent immune responses.
Nature **434**, 772-777, doi:10.1038/nature03464 (2005).
23   Sato, M. *et al.* Distinct and essential roles of transcription factors IRF-3 and IRF-7 in response to
viruses for IFN-alpha/beta gene induction. *Immunity* **13**, 539-548, doi:10.1016/s1074-
7613(00)00053-4 (2000).
24   Rudd, P. A. *et al.* Interferon response factors 3 and 7 protect against Chikungunya virus
hemorrhagic fever and shock. *J Virol* **86**, 9888-9898, doi:10.1128/JVI.00956-12 (2012).
25   Rawle, D. J. *et al.* Widespread discrepancy in Nnt genotypes and genetic backgrounds
complicates granzyme A and other knockout mouse studies. *eLife* **11**, e70207,
doi:10.7554/eLife.70207 (2022).
26   Dumenil, T. *et al.* Warmer ambient air temperatures reduce nasal turbinate and brain infection,
but increase lung inflammation in the K18-hACE2 mouse model of COVID-19. *Science of The*
Total Environment **859**, 160163, doi:https://doi.org/10.1016/j.scitotenv.2022.160163 (2023).
27   Prow, N. A. *et al.* A vaccinia-based single vector construct multi-pathogen vaccine protects
against both Zika and chikungunya viruses. *Nat Commun* **9**, 1230, doi:10.1038/s41467-018-
03662-6 (2018).
28   Hobson-Peters, J. *et al.* A recombinant platform for flavivirus vaccines and diagnostics using
chimeras of a new insect-specific virus. *Science Translational Medicine* **11**, eaax7888,
doi:10.1126/scitranslmed.aax7888 (2019).
29   Yan, K. *et al.* A Yellow Fever Virus 17D Infection and Disease Mouse Model Used to Evaluate a
Chimeric Binjari-Yellow Fever Virus Vaccine. *Vaccines* **8** (2020).
30   Oikari, L. E. *et al.* Altered Brain Endothelial Cell Phenotype from a Familial Alzheimer Mutation
and Its Potential Implications for Amyloid Clearance and Drug Delivery. *Stem Cell Reports* **14**,
924-939, doi:10.1016/j.stemcr.2020.03.011 (2020).
31   Stewart, R. *et al.* Increased neurovirulence of omicron BA.5 over BA.1 in human brain organoids
and K18-hACE2 mice. PREPRINT (Version 1) Research Square https://doi.org/10.21203/rs.3.rs-
2702556/v1. (2023).
32   Guirakhoo, F. *et al.* Immunogenicity, genetic stability, and protective efficacy of a recombinant,
chimeric yellow fever-Japanese encephalitis virus (ChimeriVax-JE) as a live, attenuated vaccine
candidate against Japanese encephalitis. *Virology* **257**, 363-372, doi:10.1006/viro.1999.9695
(1999).
33   Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of image analysis.
Nature Methods **9**, 671-675, doi:10.1038/nmeth.2089 (2012).
34   Williams, D. T., Wang, L. F., Daniels, P. W. & Mackenzie, J. S. Molecular characterization of the
first Australian isolate of Japanese encephalitis virus, the FU strain. *J Gen Virol* **81**, 2471-2480,
doi:10.1099/0022-1317-81-10-2471 (2000).
35   Poidinger, M., Hall, R. A. & Mackenzie, J. S. Molecular Characterization of the Japanese
Encephalitis Serocomplex of the Flavivirus Genus. *Virology* **218**, 417-421,
doi:https://doi.org/10.1006/viro.1996.0213 (1996).

36 Williams, D. T. *et al.* The Molecular Epidemiology and Evolution of Murray Valley Encephalitis
Virus: Recent Emergence of Distinct Sub-lineages of the Dominant Genotype 1. *PLOS Neglected*
Tropical Diseases **9**, e0004240, doi:10.1371/journal.pntd.0004240 (2015).
37 Iwasaki, Y., Zhao, J. X., Yamamoto, T. & Konno, H. Immunohistochemical demonstration of viral
antigens in Japanese encephalitis. *Acta Neuropathol* **70**, 79-81, doi:10.1007/bf00689518 (1986).
38 German, A. C. *et al.* A preliminary neuropathological study of Japanese encephalitis in humans
and a mouse model. *Trans R Soc Trop Med Hyg* **100**, 1135-1145,
doi:10.1016/j.trstmh.2006.02.008 (2006).
39 Siva Venkatesh, I. P., Bhaskar, M. & Basu, A. Japanese encephalitis viral infection modulates
proinflammatory cyto/chemokine profile in primary astrocyte and cell line of astrocytic origin.
Metabolic Brain Disease **37**, 1487-1502, doi:10.1007/s11011-022-00991-w (2022).
40 Li, Y. *et al.* Microarray Analysis Identifies the Potential Role of Long Non-Coding RNA in
Regulating Neuroinflammation during Japanese Encephalitis Virus Infection. *Frontiers in*
Immunology **8**, doi:10.3389/fimmu.2017.01237 (2017).
41 Wilhelmsson, U. *et al.* Redefining the concept of reactive astrocytes as cells that remain within
their unique domains upon reaction to injury. *Proceedings of the National Academy of Sciences*
103, 17513-17518, doi:10.1073/pnas.0602841103 (2006).
42 Zhang, Z. *et al.* The Appropriate Marker for Astrocytes: Comparing the Distribution and
Expression of Three Astrocytic Markers in Different Mouse Cerebral Regions. *BioMed Research*
International **2019**, 9605265, doi:10.1155/2019/9605265 (2019).
43 Hazlewood, J. E. *et al.* The Chimeric Binjari-Zika Vaccine Provides Long-Term Protection against
ZIKA Virus Challenge. *Vaccines (Basel)* **10**, doi:10.3390/vaccines10010085 (2022).
44 Slonchak, A. *et al.* Zika virus noncoding RNA cooperates with the viral protein NS5 to inhibit
STAT1 phosphorylation and facilitate viral pathogenesis. *Science Advances* **8**, eadd8095,
doi:10.1126/sciadv.add8095 (2022).
45 Roberts, A., Kesarwani, V., Gupta, R. & Gandhi, S. Electroactive reduced graphene oxide for
highly sensitive detection of secretory non-structural 1 protein: A potential diagnostic biomarker
for Japanese encephalitis virus. *Biosens Bioelectron* **198**, 113837,
doi:10.1016/j.bios.2021.113837 (2022).
46 Puerta-Guardo, H. *et al.* Flavivirus NS1 Triggers Tissue-Specific Vascular Endothelial Dysfunction
Reflecting Disease Tropism. *Cell Reports* **26**, 1598-1613.e1598,
doi:https://doi.org/10.1016/j.celrep.2019.01.036 (2019).
47 Daffis, S. *et al.* Interferon regulatory factor IRF-7 induces the antiviral alpha interferon response
and protects against lethal West Nile virus infection. *J Virol* **82**, 8465-8475,
doi:10.1128/jvi.00918-08 (2008).
48 Aoki, K. *et al.* Type-I interferon response affects an inoculation dose-independent mortality in
mice following Japanese encephalitis virus infection. *Virology Journal* **11**, 105, doi:10.1186/1743-
422X-11-105 (2014).
49 Miyake, M. THE PATHOLOGY OF JAPANESE ENCEPHALITIS. A REVIEW. *Bull World Health Organ*
30, 153-160 (1964).
50 Waller, C. *et al.* Japanese Encephalitis in Australia — A Sentinel Case. *New England Journal of*
Medicine **387**, 661-662, doi:10.1056/NEJMc2207004 (2022).
51 Maamary, J. *et al.* New Detection of Locally Acquired Japanese Encephalitis Virus Using Clinical
Metagenomics, New South Wales, Australia. *Emerging Infectious Disease journal* **29**, 627,
doi:10.3201/eid2903.220632 (2023).
52 Tripathi, A. *et al.* Lack of Interferon (IFN) Regulatory Factor 8 Associated with Restricted IFN-
gamma Response Augmented Japanese Encephalitis Virus Replication in the Mouse Brain. *J Virol*
95, e0040621, doi:10.1128/JVI.00406-21 (2021).

53   Bishop, C. R. *et al.* Mouse models of COVID-19 recapitulate inflammatory pathways rather than
gene expression. *PLoS Pathog* **18**, e1010867, doi:10.1371/journal.ppat.1010867 (2022).

54   Danaher, P. *et al.* Advances in mixed cell deconvolution enable quantification of cell types in
spatial transcriptomic data. *Nat Commun* **13**, 385, doi:10.1038/s41467-022-28020-5 (2022).

55   Setoh, Y. X. *et al.* Determinants of Zika virus host tropism uncovered by deep mutational
scanning. *Nat Microbiol* **4**, 876-887, doi:10.1038/s41564-019-0399-4 (2019).

56   Zhang, B. *et al.* Differential antiviral immunity to Japanese encephalitis virus in developing
cortical organoids. *Cell Death & Disease* **9**, 719, doi:10.1038/s41419-018-0763-y (2018).

57   Chambers, T. J., Nestorowicz, A., Mason, P. W. & Rice, C. M. Yellow fever/Japanese encephalitis
chimeric viruses: construction and biological properties. *J Virol* **73**, 3095-3101,
doi:10.1128/jvi.73.4.3095-3101.1999 (1999).

58   Frassetto, F. P. & Rosemberg, S. Neuropathology of yellow fever autopsy cases. *Tropical*
Diseases, Travel Medicine and Vaccines **9**, 1, doi:10.1186/s40794-022-00187-1 (2023).

59   Cohen, M. *et al.* Case Report: Yellow Fever Vaccine-Associated Neurotropic Disease and
Associated MRI, EEG, and CSF Findings. *Front Neurol* **12**, 779014,
doi:10.3389/fneur.2021.779014 (2021).

60   Myint, K. S. *et al.* Neuropathogenesis of Japanese encephalitis in a primate model. *PLoS Negl*
Trop Dis **8**, e2980, doi:10.1371/journal.pntd.0002980 (2014).

61   Japanese encephalitis virus (JEV). Australian Governement Dept of Health and Aged Care.
https://www.health.gov.au/health-alerts/japanese-encephalitis-virus-jev/japanese-encephalitis-
virus-jev (accessed April 2023).

62   Serourvey for JEV in northern Victoria. https://www.health.vic.gov.au/infectious-
diseases/serosurvey-for-japanese-encephalitis-in-northern-victoria (accessed April 2023).

63   Summary of NSW JEV seosurvey results.
https://www.health.nsw.gov.au/environment/pests/vector/Documents/jev-serosurvey-
report.pdf (accessed April 2023).

64   Nakayama, E. *et al.* Neuroinvasiveness of the MR766 strain of Zika virus in IFNAR-/- mice maps
to prM residues conserved amongst African genotype viruses. *PLoS Pathog* **17**, e1009788,
doi:10.1371/journal.ppat.1009788 (2021).

65   Zheng, W. *et al.* Liver transcriptomics reveals features of the host response in a mouse model of
dengue virus infection. *Front Immunol* **13**, 892469, doi:10.3389/fimmu.2022.892469 (2022).

66   Grant, A. *et al.* Zika Virus Targets Human STAT2 to Inhibit Type I Interferon Signaling. *Cell Host*
Microbe **19**, 882-890, doi:10.1016/j.chom.2016.05.009 (2016).

67   Prow, N. A. *et al.* Lower temperatures reduce type I interferon activity and promote alphaviral
arthritis. *PLoS Pathog* **13**, e1006788, doi:10.1371/journal.ppat.1006788 (2017).

68   Setoh, Y. X. *et al.* Helicase Domain of West Nile Virus NS3 Protein Plays a Role in Inhibition of
Type I Interferon Signalling. *Viruses* **9**, 326 (2017).

69   Laurent-Rolle, M. *et al.* The NS5 protein of the virulent West Nile virus NY99 strain is a potent
antagonist of type I interferon-mediated JAK-STAT signaling. *J Virol* **84**, 3503-3515,
doi:10.1128/JVI.01161-09 (2010).

70   Yang, H. *et al.* A novel amino acid site closely associated with the neurovirulence of live,
attenuated Japanese encephalitis vaccine (SA14-14-2 strain). *Vaccine* **38**, 2636-2642,
doi:https://doi.org/10.1016/j.vaccine.2020.01.005 (2020).

71   Yun, S. I. *et al.* A molecularly cloned, live-attenuated japanese encephalitis vaccine SA14-14-2
virus: a conserved single amino acid in the ij Hairpin of the Viral E glycoprotein determines
neurovirulence in mice. *PLoS Pathog* **10**, e1004290, doi:10.1371/journal.ppat.1004290 (2014).

72   Arroyo, J. *et al.* Molecular basis for attenuation of neurovirulence of a yellow fever
Virus/Japanese encephalitis virus chimera vaccine (ChimeriVax-JE). *J Virol* **75**, 934-942,
doi:10.1128/jvi.75.2.934-942.2001 (2001).
73   Yang, D. *et al.* Characterization of live-attenuated Japanese encephalitis vaccine virus SA14-14-2.
Vaccine **32**, 2675-2681, doi:https://doi.org/10.1016/j.vaccine.2014.03.074 (2014).
74   Liu, X. *et al.* The structure differences of Japanese encephalitis virus SA14 and SA14-14-2 E
proteins elucidate the virulence attenuation mechanism. *Protein & Cell* **10**, 149-153,
doi:10.1007/s13238-018-0551-6 (2019).
75   Yang, J. *et al.* Envelope Protein Mutations L107F and E138K Are Important for Neurovirulence
Attenuation for Japanese Encephalitis Virus SA14-14-2 Strain. *Viruses* **9** (2017).
76   Ni, H., Chang, G. J., Xie, H., Trent, D. W. & Barrett, A. D. Molecular basis of attenuation of
neurovirulence of wild-type Japanese encephalitis virus strain SA14. *J Gen Virol* **76 (Pt 2)**, 409-
413, doi:10.1099/0022-1317-76-2-409 (1995).
77   Zhou, Y. *et al.* Mutation of I176R in the E coding region weakens Japanese encephalitis virus
neurovirulence, but not its growth rate in BHK-21 cells. *Arch Virol* **163**, 1351-1355,
doi:10.1007/s00705-018-3765-2 (2018).
78   Anwar, M. N. *et al.* Phenotypic and Genotypic Comparison of a Live-Attenuated Genotype I
Japanese Encephalitis Virus SD12-F120 Strain with Its Virulent Parental SD12 Strain. *Viruses* **12**
(2020).
79   Gromowski, G. D., Firestone, C. Y. & Whitehead, S. S. Genetic Determinants of Japanese
Encephalitis Virus Vaccine Strain SA14-14-2 That Govern Attenuation of Virulence in Mice. *J Virol*
89, 6328-6337, doi:10.1128/jvi.00219-15 (2015).
80   Monath, T. P. *et al.* Single mutation in the flavivirus envelope protein hinge region increases
neurovirulence for mice and monkeys but decreases viscerotropism for monkeys: relevance to
development and safety testing of live, attenuated vaccines. *J Virol* **76**, 1932-1943,
doi:10.1128/jvi.76.4.1932-1943.2002 (2002).
81   Chen, L. K. *et al.* Generation and characterization of organ-tropism mutants of Japanese
encephalitis virus in vivo and in vitro. *Virology* **223**, 79-88, doi:10.1006/viro.1996.0457 (1996).
82   Sumiyoshi, H., Tignor, G. H. & Shope, R. E. Characterization of a highly attenuated Japanese
encephalitis virus generated from molecularly cloned cDNA. *J Infect Dis* **171**, 1144-1151,
doi:10.1093/infdis/171.5.1144 (1995).
83   Zheng, X. *et al.* Acidity/Alkalinity of Japanese Encephalitis Virus E Protein Residue 138 Alters
Neurovirulence in Mice. *J Virol* **92**, doi:10.1128/jvi.00108-18 (2018).
84   Chiou, S. S. & Chen, W. J. Phenotypic changes in the Japanese encephalitis virus after one
passage in Neuro-2a cells: generation of attenuated strains of the virus. *Vaccine* **26**, 15-23,
doi:10.1016/j.vaccine.2007.10.047 (2007).
85   Zhao, Z. *et al.* Characterization of the E-138 (Glu/Lys) mutation in Japanese encephalitis virus by
using a stable, full-length, infectious cDNA clone. *J Gen Virol* **86**, 2209-2220,
doi:10.1099/vir.0.80638-0 (2005).
86   Wang, X. *et al.* Near-atomic structure of Japanese encephalitis virus reveals critical determinants
of virulence and stability. *Nature Communications* **8**, 14, doi:10.1038/s41467-017-00024-6
(2017).
87   Xia, Q. *et al.* Virulence and Cross-Protection Conferred by an Attenuated Genotype I-Based
Chimeric Japanese Encephalitis Virus Strain Harboring the E Protein of Genotype V in Mice.
Microbiology Spectrum **10**, e01990-01922, doi:10.1128/spectrum.01990-22 (2022).
88   Wang, Q. *et al.* Japanese Encephalitis Virus Induces Apoptosis and Encephalitis by Activating the
PERK Pathway. *J Virol* **93**, doi:10.1128/jvi.00887-19 (2019).

89   Tajima, S. *et al.* E and prM proteins of genotype V Japanese encephalitis virus are required for its
increased virulence in mice. *Heliyon* **5**, e02882, doi:10.1016/j.heliyon.2019.e02882 (2019).
90   Whiteman, M. C. *et al.* Multiple amino acid changes at the first glycosylation motif in NS1
protein of West Nile virus are necessary for complete attenuation for mouse neuroinvasiveness.
Vaccine **29**, 9702-9710, doi:10.1016/j.vaccine.2011.09.036 (2011).
91   Whiteman, M. C. *et al.* Development and characterization of non-glycosylated E and NS1 mutant
viruses as a potential candidate vaccine for West Nile virus. *Vaccine* **28**, 1075-1083,
doi:10.1016/j.vaccine.2009.10.112 (2010).

Figure 1

Figure 2

A Anti-flavivirus NS1 (virus infected cells)

JEV_Nakayama **brain titer 9.2 log$_{10}$CCID$_{50}$/g**
C57BL/6J mouse 9 dpi

Cortex

Hippocampus

Anterior olfactory nucleus

Thalamus

100 μm

2 mm

50 μm

B ApopTag (apoptosis)

Cortex

Dentate gyrus

Anterior olfactory nucleus

10 μm

2 mm

50 μm

C Anti -GFAP (reactive astrocytes)

Cortex

Thalamus

10 μm

2 mm

50 μm

D

Anti-flavivirus NS1

Uninfected C57BL/6J mouse brain

ApopTag

Anti-GFAP

2 mm

2 mm

2 mm

Cortex

Cortex

Cortex

100 μm

100 μm

100 μm

Figure 3

Figure 4

Figure 5

A

Neurons with pyknotic nuclei
Nakayama C57BL/6J 11 dpi

Hyper-eosinophilic neurons
Nakayama C57BL/6J 9 dpi

Neuronal vacuolation
Nakayama IRF7$^{-/-}$ 7 dpi

Uninfected control

Perivascular cuffing
NSW IRF7$^{-/-}$ 8 dpi

Uninfected control

Leukocyte infiltrates
Nakayama IRF7$^{-/-}$ 7 dpi

Uninfected control

Small hemorrhagic lesions
Nakayama C57BL/6J 9 dpi

Uninfected control

Meningitis
Nakayama IRF7$^{-/-}$ 7 dpi

Uninfected control

Microgliosis
Nakayama IRF7$^{-/-}$ 19 dpi (survived)

Uninfected control

B Microgliosis (Iba1 IHC)
Nakayama C57BL/6J 9 dpi

Uninfected control

Figure 6

Panel A: hBO "minibrains"; 20 rpm

Panel B: JEV_Nakayama; 0.5 mm; 10 µm; JEV_NSW/22; 0.5 mm; 10 µm; JEV_FU; MVEV_TC123130; Imojev; YFV17D; Uninfected

Panel C:
p<0.0001
p<0.001
p=0.008
p<0.001
p<0.001
p<0.001

Virus titer ($\log_{10}$CCID$_{50}$/ml) ± SE
Hours post infection

- JEV_Nakayama
- JEV_FU
- JEV_NSW/22
- MVEV_TC123130
- Imojev
- YFV-17D

Panel D:
p=0.055
p=0.009
p=0.01

% change in organoid area (day 11 vs day 0) ± SE

Uninfected, JEV_Nakayama, JEV_FU, JEV_NSW/22, MVEV_TC123130, Imojev, YFV-17D

A

Virus name	GenBank accession	JEV Genotype	Isolation year	Isolation source	Passage history
JEV$_{Nakayama}$	EF571853	3	1935	Human cerebrospinal fluid	>11 x suckling mouse brain
JEV$_{FU}$	AF217620	2	1995	Human serum	C6/36 (unknown number)
JEV$_{NSW2022}$	OP904182	4	2022	Stillborn piglet brain	1 x C6/36 1 x BHK-21
MVEV$_{TC123130}$	JN119814	N/A	1974	Human brain	Unknown
YFV 17D	MT107250	N/A	1927 (Original "Asibi" parent isolate)	Human serum (Original "Asibi" parent isolate)	53 x monkeys 18 x embryonic mouse tissue 50 x minced whole chicken embryo 152 x nervous tissue deprived chick embryo
JEV$_{prME}$ component in Imojev (SA$_{14}$-14-2 strain)	MK585066	3	1954 (Original SA-14 parent isolate)	*Culex pipiens* mosquito larvae (Original SA-14 parent isolate)	100 x primary hamster kidney cells 22 x suckling mice

B

Supplementary Figure 1. (A) Summary of virus isolates used in the study. (B) Phylogenetic tree for whole genome nucleotide sequence (left) and prME amino acid sequence (right). Phylogenetic trees were constructed after nucleotide or amino acid sequence alignment using MEGA-X (Molecular Evolutionary Genetics Analysis 10, Penn State University, State College, PA, USA) and the ClustalW plugin with default parameters. The phylogenetic tree was constructed using the Maximum Likelihood method and the General Time Reversible model (nucleotide sequence) or JTT matrix-based model (amino acid sequence). Whole genome sequence is not available for MVEV$_{TC123130}$. Whole genome sequence for Imojev was constructed by combining the prME sequence from JEV$_{SA14-14-2}$ with the remainder of the genome from YFV 17D.

Activity

Dpi	JEV Nakayama Cg1#2	Cg1#6	JEV FU Cg2#1	MVEV Cg1#3
0	0	0	0	0
1	0	0	0	0
2	0	0	0	0
3	0	0	0	0
4	0	0	0	0
5	0	0	0	0
6	0	0	0	0
7	0	0	0	0
8	0	0	1	0
9	1	0	■	1
10	■	0	■	■
11	■	1	■	■
12	■	■	■	■

Weight loss score

Dpi	JEV Nakayama Cg1#2	Cg1#6	JEV FU Cg2#1	MVEV Cg1#3
0	0	0	0	0
1	0	0	0	0
2	0	0	0	0
3	0	0	0	0
4	0	0	0	0
5	0	0	0	0
6	0	0	0	0
7	0	0	2	0
8	1	0	3	2
9	3	0	■	3
10	■	1	■	■
11	■	3	■	■
12	■	■	■	■

Posture

Dpi	JEV Nakayama Cg1#2	Cg1#6	JEV FU Cg2#1	MVEV Cg1#3
0	0	0	0	0
1	0	0	0	0
2	0	0	0	0
3	0	0	0	0
4	0	0	0	0
5	0	0	0	0
6	0	0	0	0
7	0	0	0	0
8	0	0	1	0
9	1	0	■	1
10	■	0	■	■
11	■	1	■	■
12	■	■	■	■

Fur ruffling

Dpi	JEV Nakayama Cg1#2	Cg1#6	JEV FU Cg2#1	MVEV Cg1#3
0	0	0	0	0
1	0	0	0	0
2	0	0	0	0
3	0	0	0	0
4	0	0	0	0
5	0	0	0	0
6	0	0	0	0
7	0	0	0	0
8	0	0	0	0
9	1	0	■	1
10	■	0	■	■
11	■	0	■	■
12	■	■	■	■

B

Criteria	Level 0	Level 1	Level 2	Level 3
Joint swelling	Normal	slight to moderate swelling	Moderate/pronounce swelling and/or erythema	Joint distortion or rigidity
Injection site reaction	No reaction	Mild scaring	Moderate ulceration	Severe ulceration
Posture	Normal	Hunching noted only at rest	Severe hunching, some impairment of normal movement	Wobbling, unable to maintain upright posture
Activity	Normal	Mild to moderately decreased	Stationary unless stimulated.	Moderate paralysis, usually more than 1 limb
Fur ruffling (fever sign)	Normal	Mild to moderate ruffling	Severe ruffling	Shivering
Hind leg weakness	Normal	Mild hind limb weakness/very mild and/or occasional gait alterations	Moderate hind limb weakness and occasional dragging of hind legs	Severe hind limb weakness/repeated dragging of hind legs
Weight loss	Normal	<10 %	<15%	≥20%

Supplementary Figure 2. Disease scores for the four mice that were euthanized.
(A) For the mice shown in Fig. 1E that were euthanized, the disease scores are shown.
(B) The mice were monitored daily using the score card. Any animal reaching a level of 3 in any single criteria were euthanized. If an animal reaches a grade of 2 in two or more criteria the animal will be euthanized.

Supplementary Figure 3. Viral antigen (NS1) staining in C57BL/6J mouse brains. IHC using a pan-flavivirus NS1 monoclonal antibody (4G4). (A-C) The brains from the 3 other C57BL/6J mice that required euthanasia in Fig. 1 (the fourth is shown in Fig. 2). (D) JEV$_{FU}$ infected C57BL/6J mice that lost ~15% body weight then recovered (from Fig. 1E). (E) A representative image of a mouse brain where infection did not lead to significant weight loss. (F) Uninfected control.

Supplementary Figure 4. ApopTag staining in C57Bl/6J mouse brains. Staining for apoptosis using Apoptag. (A-C) The brains from the 3 other C57BL/6J mice that required euthanasia in Figure 1. (D) The JEV_{FU} infected C57BL/6J mice that lost ~15% body weight then recovered (see Figure 1E). (E) Uninfected control.

C57BL/6J	**IRF7$^{-/-}$**	**IFNAR$^{-/-}$; all JEV$_{NSW/22}$**

MVEV

Nakayama

FU

Nakayama

Nakayama

Nakayama

Nakayama

NSW/22

Uninfected

Supplementary Figure 5. GFAP IHC for reactive astrocytes in mouse brains. Staining for reactive actrocyte using GFAP for the brains from C57BL/6J, IRF7$^{-/-}$ and IFNAR$^{-/-}$ mice that succumbed to the indicated virus. Uninfected mouse brains are also shown.

A

Activity

Dpi	FU Cg4# (1)	FU Cg4# (2)	FU Cg4# (3)	FU Cg4# (4)	FU Cg1# (1)	Nakayama Cg2# (1)	Nakayama Cg5# (1)	Nakayama Cg5# (2)	Nakayama Cg5# (3)	NSW/22 Cg3# (1)	NSW/22 Cg3# (2)	NSW/22 Cg3# (3)	NSW/22 Cg3# (4)
0	0	0	0	0	0	0	0	0	0	0	0	0	0
1	0	0	0	0	1	1	0	0	0	0	0	0	0
2	1	1	1	1	2	1	0	0	0	1	1	1	1
3	2	2	2	2	■	2	2	2	2	2	2	2	2

Weight loss score

Dpi	FU Cg4# (1)	FU Cg4# (2)	FU Cg4# (3)	FU Cg4# (4)	FU Cg1# (1)	Nakayama Cg2# (1)	Nakayama Cg5# (1)	Nakayama Cg5# (2)	Nakayama Cg5# (3)	NSW/22 Cg3# (1)	NSW/22 Cg3# (2)	NSW/22 Cg3# (3)	NSW/22 Cg3# (4)
0	0	0	0	0	0	0	0	0	0	0	0	0	0
1	1	1	1	0	0	0	0	0	0	0	0	0	0
2	2	1	2	2	2	1	1	1	1	0	0	0	0
3	2	1	2	2	■	3	2	2	2	2	1	2	1

Posture

Dpi	FU Cg4# (1)	FU Cg4# (2)	FU Cg4# (3)	FU Cg4# (4)	FU Cg1# (1)	Nakayama Cg2# (1)	Nakayama Cg5# (1)	Nakayama Cg5# (2)	Nakayama Cg5# (3)	NSW/22 Cg3# (1)	NSW/22 Cg3# (2)	NSW/22 Cg3# (3)	NSW/22 Cg3# (4)
0	0	0	0	0	0	0	0	0	0	0	0	0	0
1	0	0	0	0	0	0	0	0	0	0	0	0	0
2	1	1	1	1	1	1	1	1	0	0	0	0	0
3	2	2	2	2	■	2	2	2	2	2	2	2	2

Fur ruffling

Dpi	FU Cg4# (1)	FU Cg4# (2)	FU Cg4# (3)	FU Cg4# (4)	FU Cg1# (1)	Nakayama Cg2# (1)	Nakayama Cg5# (1)	Nakayama Cg5# (2)	Nakayama Cg5# (3)	NSW/22 Cg3# (1)	NSW/22 Cg3# (2)	NSW/22 Cg3# (3)	NSW/22 Cg3# (4)
0	0	0	0	0	0	0	0	0	0	0	0	0	0
1	0	0	0	0	0	0	0	0	0	0	0	0	0
2	1	1	1	1	0	0	0	0	0	0	0	0	0
3	1	1	1	1	■	0	1	1	1	0	0	0	0

B

$R^2 = 0.75$
p=0.0003

Brain virus titer ($log_{10}CCID_{50}$/g) vs Viremia ($log_{10}CCID_{50}$/ml)

C

$JEV_{NSW/22}$ brain IFNAR$^{-/-}$ mouse, 3 dpi, titer 6.6 log_{10} $CCID_{50}$/g

Anti-flavivirus NS1   ApopTag

Supplementary Figure 6. JEV infection of IFNAR$^{-/-}$ mice. (A) Disease scores as per Supplementary Figure 2B for mice shown in Figure 3. (B) Pearson correlation between brain titer (y-axis) from Figure 3E and viremia (x-axis) from Figure 3A. (B) IHC for flavivirus NS1 (left) or ApopTag (right) showing no detectable staining of brain cells. Images are representative of all IFNAR$^{-/-}$ brains and JEV isolates.

Supplementary Figure 7. JEV$_{NSW/22}$ dose range in C57BL/6J and IRF7$^{-/-}$ mice. Adult female C57BL/6J mice (15 weeks of age) or IRF7$^{-/-}$ mice (15-48 weeks of age) were infected s.c. with JEV$_{NSW/22}$ at the indicated CCID$_{50}$ doses of JEV$_{NSW/22}$. (A) Percent body weight change of individual C57BL/6J mice compared to each mouse's weight 0 dpi. (B) As for A but using IRF7$^{-/-}$ mice. The data for 5x10^3 is the same as Figure 4E, showing individual mice. (C-E) Viremia for C57BL/6J mice infected with the indicated doses, determined by CCID$_{50}$ assays with a limit of detection of ~2 log$_{10}$CCID$_{50}$/ml. Horizontal lines indicate mean at each time point, and error bars represent SE. (F-G) As for C-E but for IRF7$^{-/-}$ mice.

Supplementary Figure 8. Disease scores for IRF7-/- mice that were euthanized.
Mice described in Figure 4E, F. Scoring system shown in Supplementary Figure 2B.

Anti-flavivirus NS1 (4G4)

JEV$_{Nakayama}$ brain titer 3.9 log$_{10}$CCID$_{50}$/g

JEV$_{Nakayama}$ brain titer 6.6 log$_{10}$CCID$_{50}$/g

100 μm

Supplementary Figure 9. Viral antigen staining in IRF7$^{-/-}$ mouse brains. IHC using a pan-flavivirus NS1 monoclonal antibody (4G4) for the brains from the 2 other IRF7$^{-/-}$ mice that required euthanasia (Figure 3). Low levels of staining of neurons were found in the JEV$_{Nakayama}$ infected mice that had a low virus titer in the brain (3.9 log$_{10}$CCID$_{50}$/g) (left). Black arrowheads show small patches of staining in the cortex, hippocampus and thalamus (right).

Acute (euthanized due to weight loss/disease scores)

Mouse	C57BL/6J			IRF7-/-			IFNAR-/-		
JEV isolate	NSW	Nakayama	FU	NSW	Nakayama	FU	NSW	Nakayama	FU
Euthanasia		9-11 dpi	8 dpi	8 dpi	7-12 dpi	6-10 dpi	3 dpi	3 dpi	3 dpi
Neuronal degeneration /vacuolation	N/A	2/2	1/1	0/1	2/6	5/8	0/4	0/3	0/4
Perivascular cuffing	N/A	2/2	1/1	1/1	6/6	8/8	0/4	0/3	0/4
Hemorrhage	N/A	2/2	1/1	0/1	5/6	5/8	3/4	2/3	2/4
Leukocyte infiltrates	N/A	2/2	1/1	1/1	1/6	3/8	0/4	0/3	0/4
Microgliosis	N/A	2/2	1/1	0/1	3/6	6/8	0/4	0/3	0/4
Meningitis	N/A	2/2	1/1	1/1	5/6	6/8	0/4	0/3	0/3

Chronic (euthanized after recovery from acute phase)

Mouse	C57BL/6J			IRF7-/-		
JEV isolate	NSW	Nakayama	FU	NSW	Nakayama	FU
Euthanasia	32 dpi	32 dpi	32 dpi	19 dpi	19 dpi	
Neuronal degeneration /vacuolation	0/6	0/4	0/5	0/3	0/1	N/A
Perivascular cuffing	0/6	0/4	1/5	0/3	1/1	N/A
Hemorrhage	1/6	2/4	1/5	1/3	1/1	N/A
Leukocyte infiltrates	0/6	0/4	1/5	0/3	0/1	N/A
Microgliosis	0/6	0/4	2/5	1/3	1/1	N/A
Meningitis	0/6	0/4	0/5	0/3	1/1	N/A

Supplementary Figure 10. Lesion presence in H&E brain sections. For lesions described in Figure 5. H&E scoring of 0 indicates no overt presence of these lesions. N/A – not available; no NSW-infected C57BL/6J mice died of acute infection, and all FU infected IRF7[-/-] mice died of acute disease.

Supplementary Figure 11. Histopathological lesions overlapped with areas of virus infection. (A) Cortex regions from JEV infected C57BL/6J, IRF7$^{-/-}$, IFNAR$^{-/-}$ mice or uninfected mice. The cortex regions from JEV infected mice were heavily infected, with most neurons stained positive for viral antigen. H&E detectable signs of neuron degeneration/vacuolation were concentrated in regions with high staining for viral antigen. (B) Viral antigen staining (left) in hippocampus overlapped with histological signs of neuron degeneration/vacuolation (right) (see Figure 5).

A   IRF7$^{-/-}$ and C57BL/6J mice (n=18) that were euthanized due to ethically defined disease end points

	Neuron degeneration/ vacuolation	Perivascular cuffing	Hemorrhage	Leukocyte infiltrates	Microgliosis
Cerebral cortex	9/18	14/18	5/18	5/18	3/18
Hypothalamus	0/18	4/18	3/18	2/18	4/18
Thalamus	0/18	11/18	2/18	2/18	7/18
Hippocampus	0/18	5/18	0/18	0/18	4/18
Caudate putamen	1/18	9/18	1/18	2/18	3/18
Basal forebrain	0/18	6/18	2/18	0/18	0/18
Midbrain	0/18	7/18	1/18	0/18	3/18
Pons	0/18	4/18	3/18	1/18	1/18
Medulla	0/18	3/18	0/18	0/18	0/18
Cerebellum	0/18	3/18	1/18	0/18	0/18

B   IRF7$^{-/-}$ and C57BL/6J mice (n=19) that survived infection (taken on day 19 and day 32, respectively)

	Neuron degeneration/ vacuolation	Perivascular cuffing	Hemorrhage	Leukocyte infiltrates	Microgliosis
Cerebral cortex	0/19	2/19	2/19	1/19	4/19
Hypothalamus	0/19	0/19	2/19	0/19	0/19
Thalamus	0/19	0/19	0/19	0/19	2/19
Hippocampus	0/19	0/19	0/19	0/19	0/19
Caudate putamen	0/19	0/19	0/19	0/19	2/19
Basal forebrain	0/19	1/19	4/19	0/19	0/19
Midbrain	0/19	0/19	0/19	0/19	0/19
Pons	0/19	0/19	1/19	0/19	0/19
Medulla	0/19	0/19	0/19	0/19	0/19
Cerebellum	0/19	1/19	2/19	0/19	0/19
Anterior olfactory nucleus	0/19	0/19	1/19	0/19	0/19
Ventral striatum	0/19	0/19	2/19	0/19	0/19

Supplementary Figure 12. Histopathological lesions in specific brain regions for C57BL/6J and IRF7$^{-/-}$. Scoring only reflects presence or absence of lesions, and does not indicate severity of lesions. (A) Acute (B) Chronic.

Supplementary Figure 13. RNA-Seq read counts from brains of C57BL/6 JEV-infected and mock-infected mice obtained from the Gene Expression Omnibus (GEO accession: GSE154002) were normalised for sequencing depth and composition using DESeq2. An estimation of cell type abundances was performed on normalised read counts with the SpatialDecon package in R, using an adult mouse immune cell gene expression reference (Yoshida *et al.* 2019. Cell 176; 897-912.e20). Log_{10} abundance scores are shown for cell types that were significantly different between infected and mock-infected groups (t-test, p-value < 0.05), with cell types ordered from largest to smallest difference in mean abundance score between groups.

Supplementary Fig. 14. Human cortical brain organoids measurements and magnified virus staining. (A) Occasional cells staining positive for IHC for viral antigen in hBOs infected with Imojev or YFV 17D. (B) Light microscopy images of hBOs infected with $JEV_{Nakayama}$, JEV_{FU}, $JEV_{NSW2022}$, $MVEV_{TC123130}$, Imojev or YFV 17D at day 0 and day 11 post infection compared to uninfected hBOs. Scale bar is consistent across all images. (C) Change in hBO area on day 4, 7, 9 and 11 compared to day 0. n=8 for uninfected and $MVEV_{TC123130}$, n=4 for all others.

Supplementary Figure 15. Replication and CPE in RENcell VM neural progenitor and Vero E6 cell lines. (A) Viral growth kinetics in RENcell VM determined by $CCID_{50}$ assays of culture supernatant at the indicated times post infection. Data is the mean of 6 replicates per virus isolate across 2 independent experiments; (limit of detection is $2 \log_{10}CCID_{50}$/ml). (B) Images of crystal violet stained RENcell VM at 6 dpi (representative of n=3 per group). Less blue/violet stained cells indicates more viral CPE. (C) Vero cells were seeded at 2.5×10^5 cells per well in 24 well plates overnight at 37°C. Cells were infected at MOI=0.05 for 1 hr at 37°C before overlay media (0.375% w/v high viscosity carboxymethyl cellulose [CMC, Sigma-Aldrich]/RPMI 1640 substituted with 2% FCS) was added to each well. Plates were incubated for 5 days before the media was removed and monolayers were fixed and stained with 0.1% w/v crystal violet (Sigma-Aldrich) in formaldehyde (1% v/v) and methanol (1% v/v). Plates were washed in tap water and dried before images were taken. Images shown are representative of n=4 per group. CPE is less pronounced in $JEV_{NEW/22}$ infected cells, with plaques (arrows), rather than fulminant CPE, apparent at 5 dpi.

Supplementary Fig. 16. Neither mouse background nor *Nnt* significantly affected JEV neuropathogenesis. Female C57BL/6J (10-16 weeks old) (black circles), C57BL/6NJ*Nnt-/-* (blue squares), and C57BL/6NJ (red triangles) mice were infected s.c. with 5×10^5 CCID$_{50}$ JEV$_{Nakayama}$. (A) Average viremia as determined by CCID$_{50}$ assays; limit of detection for individual mice is 2 log$_{10}$CCID$_{50}$/ml. (B) Percent body weight change of individual mice compared to their weight 0 dpi. Six mice lost ≥20% body weight and required euthanasia (†). (C) Kaplan Myer plot showing percent survival (n=6 for each mouse strain). (D) Viral tissue titers in brains of the six euthanized mice (n=2 from each mouse strain). Tissue titers determined by CCID$_{50}$ assays (limit of detection ~2 log$_{10}$CCID$_{50}$/g). (E) Representative images of H&E stained sections of brains from mice that required euthanasia. No overt differences were identified between C57BL/6J, C57BL/6NJ*Nnt-/-*, or C57BL/6NJ mice.