Neurovirulence of the Australian outbreak Japanese Encephalitis virus genotype 4 is lower

compared to genotypes 2 and 3 in mice and human cortical brain organoids

Wilson Nguyen[a], Narayan Gyawali[a], Romal Stewart[a], Bing Tang[a], Abigail L. Cox[a], Kexin Yan[a],

Thibaut Larcher[b], Cameron Bishop[a], Nicholas Wood[c], Gregor J. Devine[a], Andreas Suhrbier[a,d#],

Daniel J. Rawle[a*#].

[a] QIMR Berghofer Medical Research Institute, Brisbane, Queensland 4029, Australia.

[b] INRAE, Oniris, PAnTher, APEX, Nantes, France.

[c] National Centre for Immunisation Research and Surveillance, Westmead, New South Wales,

Australia.

[d] GVN Center of Excellence, Australian Infectious Disease Research Centre, Brisbane,

Queensland, 4029 and 4072, Australia.

[*]Corresponding author. Email: Daniel.Rawle@qimrberghofer.edu.au

[#] should be considered joint last authors

SUMMARY

Background

Human infections with Japanese encephalitis virus (JEV) are a leading cause of viral

encephalitis. An unprecedented outbreak of JEV genotype 4 was recently reported in Australia,

with an isolate ($JEV_{NSW/22}$) obtained from a stillborn piglet brain.

Methods

Herein we compared the neuropathology of $JEV_{NSW/22}$, JEV_{FU} (genotype 2) and $JEV_{Nakayama}$

(genotype 3) in adult C57BL/6J wild-type mice, mice deficient in interferon regulatory factor 7

($Irf7^{-/-}$), and mice deficient in type I interferon receptor ($Ifnar^{-/-}$), as well as in human cortical

brain organoids (hBOs). Using human serum post-Imojev vaccination, we performed

neutralisation assays to determine $JEV_{NSW/22}$ susceptibility to vaccine responses.

Findings

In C57BL/6J and $Irf7^{-/-}$ mice with lethal outcomes, brain infection and histopathological lesions

recapitulated those seen in humans and primates. JEV was universally lethal in $Ifnar^{-/-}$ mice by

day 3 with histological signs of brain hemorrhage, but produced no other detectable brain

infection or lesions, with viral protein detected in blood vessels but not neurons. We thus

describe a new $Irf7^{-/-}$ mouse model for $JEV_{NSW/22}$, which had increased viremia compared to

C57BL/6J mice, allowing for lethal neuroinvasive infection in one mouse. Overall, $JEV_{NSW/22}$

was less neurovirulent than other JEV isolates in C57BL/6J and $Irf7^{-/-}$ mice, and was more

sensitive to type I interferon. All JEV isolates showed robust cytopathic infection of human

cortical brain organoids, albeit lower for $JEV_{NSW/22}$. We also show that Imojev vaccination in

humans induced neutralizing antibodies against $JEV_{NSW/22}$, with the level of cross-neutralisation

related to the conservation in envelope protein amino acid sequences for each isolate.

Interpretation

Our study establishes $JEV_{NSW/22}$ mouse models of infection, allowing for possible lethal

neuroinvasive infection that was rarer than for other JEV genotypes. JEV vaccination regimens

may afford protection against this newly emerged JEV genotype 4 strain, although neutralizing

antibody responses are sub-optimal.

Funding

QIMRB received a generous philanthropic donation from the Brazil Family Foundation awarded to D.J.R. to support Japanese Encephalitis virus research at QIMRB. A.S. holds an Investigator grant from the National Health and Medical Research Council (NHMRC) of Australia (APP1173880). We also acknowledge the intramural grant from QIMR Berghofer awarded to R.S. and D.J.R. for purchase of the CelVivo Clinostar incubator for producing human cortical brain organoids. The project "Japanese encephalitis vaccine via the intradermal route in children and adults (JEVID-2): A clinical trial comparing the immunogenicity and safety of Japanese encephalitis vaccine administered by subcutaneous and intradermal routes" being conducted by G.D., N.G., and N.W. was funded by the Sydney Children's Hospitals Network and New South Wales Health.

Keywords: Japanese encephalitis virus, mouse models, brain organoids, vaccine responses, virus outbreak.

Research in context

Evidence before the study

JEV from the historically rare genotype 4 recently emerged in Australia, causing an unprecedented outbreak, with 44 human cases and 7 fatalities. While a range of JEV mouse models have been reported, none of them infect adult mice with a genotype 4 isolate. The efficacy of current vaccines for this JEV genotype are also unclear.

Added value of this study

We establish well characterised adult and subcutaneously infected mouse models for JEV which recapitulate many aspects of human disease including lethal neuroinvasive infection and severe histopathological lesions. Prolonged viremia was significantly associated with lethal neuroinvasiveness in $Irf7^{-/-}$ mice. We demonstrate that a genotype 4 Australian isolate, $JEV_{NSW/22}$, exhibited markedly diminished lethal neuroinvasion compared to other JEV genotypes. Using serum from Imojev vaccine recipients, neutralizing antibodies against $JEV_{NSW/22}$ were present, albeit at sub-optimal titers.

Implications of all the available evidence

The establishment of well characterised adult mouse models of $JEV_{NSW/22}$ with rare neuropenetrance after peripheral inoculation that recapitulate human disease is an important tool that can now be deployed in pre-clinical studies and to understand disease pathogenesis. Our study suggests that new vaccines should be developed against circulating JEV strains for optimal neutralizing antibody responses.

Introduction

Japanese Encephalitis virus (JEV) is a single-stranded positive-sense RNA virus from the

Flaviviridae family that is transmitted from amplifying hosts (primarily pigs and wading birds)

via mosquitoes to humans (1). While infection is usually asymptomatic, encephalitis can

develop in ≈1 in 250 people, with ~30% of encephalitic cases becoming fatal, and 30-50% of

non-fatal encephalitic cases retaining persistent neurological symptoms including seizures,

speech impediments and paralysis (2, 3). JEV is the leading cause of viral encephalitis in Asia,

with ~70,000 cases and ~20,000 deaths per annum (4). After a bite from an infected mosquito,

JEV replicates in peripheral blood monocytes producing a viremia that, in some cases, leads to

virus crossing the blood brain barrier (5). JEV primarily infects neurons in the brain (6) leading

to uncontrolled inflammation (encephalitis) and neuronal cell death (2). There are several

available and effective vaccines against JEV (7), but there are no specific licensed treatments.

JEV exists as five genotypes (1 to 5) which are phylogenetically, antigenically and

geographically distinct (8-10). Genotype 3 was the main genotype endemic in Asia until 1990,

after which genotype I has dominated since (11). Genotype 2 has been identified in Malaysia,

Indonesia, Papua New Guinea and Australian outbreaks from 1970-2000. Genotype 5 has been

identified in China, and is now dominant in South Korea (12). Genotype 4 was the least

common genotype worldwide, having only been identified in mosquitoes from Indonesia and

Papua New Guinea (13). In February 2021, a fatal JEV infection occurred on the Tiwi Islands,

80 km off the coast of Darwin, Australia (14). Sequencing identified that the virus belonged to

the historically rare JEV genotype 4 (14). In 2022, a geographically widespread outbreak

throughout most Australian states was attributed to JEV genotype 4, which caused 44 confirmed

human cases and 7 deaths (15). Outbreaks occurred in several piggeries, causing high abortion

and stillbirth rates in sows (16). Based on proximity to piggeries, estimates indicate that

~740,000 people may be at risk of being infected by JEV in Australia (16). Given that *Culex*

annulirostris mosquitoes, considered the primary vector for JEV in Australia (15, 16), and

amplifying vertebrate hosts including wading birds and pigs are widespread, it is possible that

JEV may become, or is already, endemic in Australia (15). Murray Valley Encephalitis virus

(MVEV) and Kunjin virus are phylogenetically closely related to JEV and have similar vector

and reservoir hosts, and are endemic to Australia (17).

Efforts to develop treatments for JEV neuropathology are hindered by the difficulties in

diagnosing early infection (13) before the virus has infected central nervous system (CNS)

neurons; and once the virus has infected CNS neurons, treatments must also be able to

effectively enter the brain. While a range of JEV mouse models have been reported

(Supplementary Table 1) (18), there are no studies of genotype 4 in adult mice, with the only

study on genotype 4 JEV using 3-week-old mice (19). Well-characterized, adult mouse models

of recent genotype 4 isolates would represent useful tools to study mechanisms of infection and

disease and evaluate potential new interventions. Herein, we compare and contrast a JEV

genotype 4 isolate from the recent Australian outbreak ($JEV_{NSW/22}$) with historical JEV isolates

(genotypes 2 and 3) in three mouse strains, human cortical brain organoids and *in vitro*

neutralisation assays with human serum after JEV vaccination.

Materials and methods

Ethics statement and regulatory compliance

All mouse work was conducted in accordance with the "Australian code for the care and use of

animals for scientific purposes" as defined by the National Health and Medical Research Council

of Australia. Mouse work was approved by the QIMR Berghofer Medical Research Institute

(MRI) Animal Ethics Committee (P3746, A2108-612). Mice were euthanized using CO_2.

Breeding and use of GM mice was approved under a Notifiable Low Risk Dealing (NLRD)

Identifier: NLRD_Suhrbier_Oct2020: NLRD 1.1(a). Use of Imojev was approved under

NLRD_Suhrbier_Oct2019: NLRD2.1(d).

All infectious JEV work was conducted in a dedicated suite in a biosafety level-3 (PC3)

facility at the QIMR Berghofer MRI (Australian Department of Agriculture, Water and the

Environment certification Q2326 and Office of the Gene Technology Regulator certification

3445). All work was approved by the QIMR Berghofer MRI Safety Committee (P3746).

Human serum samples before and after Imojev vaccination were collected from 9 participants

with human research ethics approval from Sydney Children's Hospitals Network Human

Research Ethics Committee (2022/ETH02471 HCHNHREC) for an ongoing project led by Dr.

Nicholas Wood; "Japanese encephalitis vaccine via intradermal route in children and adults

(JEVID-2): Critical policy relevant research for Australia".

Human serum samples after Imojev vaccination were also collected from 10 participants with

approval from QIMR Berghofer MRI Human Research Ethics Committee (P3476) provided for

neutralisation assays (including MVEV cross-neutralisation).

Research with JEV were approved under the Queensland Biosecurity Act, Scientific Research

Permit (restricted matter) – Permit number PRID000916.

Cell lines and culture

Vero E6 (C1008, ECACC, Wiltshire, England; obtained via Sigma Aldrich, St. Louis, MO,

USA), BHK-21 (ATCC# CCL-10), and C6/36 cells (ATCC# CRL-1660) were cultured in

medium comprising RPMI 1640 (Gibco), supplemented with 10% fetal bovine serum (FBS),

penicillin (100 IU/ml)/streptomycin (100 µg/ml) (Gibco/Life Technologies) and L-glutamine

(2 mM) (Life Technologies). Cells were cultured at 37°C and 5% CO_2. Cells were routinely

checked for mycoplasma (MycoAlert Mycoplasma Detection Kit MycoAlert, Lonza) and FBS

was assayed for endotoxin contamination before purchase (20).

Mouse embryonic fibroblasts (MEFs) were either wild type or *Irf3/7*[-/-] and have been

described previously (21). MEFs were cultured in Dulbecco's Modified Eagle Medium

(DMEM) (Gibco), supplemented with 50µg/ml Penicillin/50µg/ml Streptomycin and 10% Fetal

Bovine Serum (FBS). Cells were cultured at 37°C and 5% CO_2. For virus growth kinetics,

MEFs were seeded at $2x10^5$ cells/ml in 12 or 24 well plates one day prior to infection at

multiplicity of infection (MOI) 0.1 of the indicated JEV or MVEV. After 1 hr incubation, MEFs

were washed twice with 1 ml PBS, and culture medium was added and sampled daily for virus

titrations by $CCID_{50}$ (see below).

RENcell VM Human Neural Progenitor Cell Line (Sigma Aldrich) were cultured in medium

comprising DMEM F-12 (Thermo Fisher Scientific), penicillin (100 IU/ml)/streptomycin

(100 µg/ml) (Gibco/Life Technologies), 20 ng/ml FGF (STEMCELL Technologies), 20 ng/ml

EGF (STEMCELL Technologies), and 2% B27 supplements (Thermo Fisher Scientific). Cells

were detached using StemPro Accutase Cell Dissociation Reagent (Thermo Fisher Scientific),

and were cultured in Matrigel (Sigma). For crystal violet staining of remaining cells after

infection, formaldehyde (7.5% w/v)/crystal violet (0.05% w/v) was added to wells overnight,

plates washed twice in water, and plates dried overnight.

Virus isolates and culture

JEV$_{Nakayama}$ (GenBank: EF571853), JEV$_{FU}$ (GenBank: AF217620), and Imojev were obtained

from A/Prof. Gregor Devine (QIMR Berghofer MRI). $JEV_{NSW/22}$ (GenBank: OP904182) was a

gift from Dr. Peter Kirkland (Elizabeth Macarthur Agriculture Institute, New South Wales,

Australia). $MVEV_{TC123130}$ (GenBank: JN119814.1) was obtained from the "Doherty Virus

Collection" currently held at QIMR Berghofer MRI. Virus stocks were generated by infection of

C6/36 cells (for JEV and MVEV) or Vero E6 cells (for Imojev and YFV 17D) at multiplicity of

infection (MOI) ≈0.1, with supernatant collected after ~5 days, cell debris removed by

centrifugation at 3000 x g for 15 min at 4°C, and virus aliquoted and stored at -80°C. Virus

stocks used in these experiments underwent less than three passages in our laboratory, with prior

passage history in Supplementary Figure 1. Virus titers were determined using standard $CCID_{50}$

assays (see below).

Validation of virus stock sequences

Viral RNA was extracted from virus stock culture supernatants using NucleoSpin RNA Virus kit

(Machery Nagel) as per manufacturer's instructions. cDNA was synthesized using ProtoScipt

First Strand cDNA Synthesis Kit (New England Biolabs) as per manufacturer's instructions.

PCR was performed using Q5 High-Fidelity 2X Master Mix (New England Biolabs) as per

manufacturer's instructions with the following primers; JEV envelope Forward 5'

GGAAGCATTGACACATGTGC 3' and Reverse 5' TCTGTGCACATACCATAGGTTGTG 3',

and MVEV envelope Forward 5' GAGCATTGACACATGCGCAAAG and Reverse 5'

TGTGCACATCCCATAAGTGGTTC 3'. PCR products were run on a 1% agarose gel and

DNA was extracted using Monarch DNA Gel Extraction Kit (New England Biolabs). DNA was

sequenced by Sanger sequencing using either the forward or reverse primer. Sequences of our

virus stocks matched the sequences on GenBank (Supplementary Figure 1).

$CCID_{50}$ assays

Vero E6 cells were plated into 96 well flat bottom plates at $2x10^4$ cells per well in 100 µl of

medium. For tissue titrations, tissues were homogenized in tubes each containing 4 ceramic

beads twice at 6000 x g for 15 seconds, followed by centrifugation twice at 21,000 x g for 5 min

before 5 fold serial dilutions in 100 µl RPMI 1640 supplemented with 2% FBS. For cell culture

supernatant or mouse serum, 10 fold serial dilutions were performed in 100 µl RPMI 1640

supplemented with 2% FBS. A volume of 100 µl of serially diluted samples were added to each

well of 96 well plate containing Vero E6 cells and the plates cultured for 5-7 days at 37°C and

5% CO_2. The virus titer was determined by the method of Spearman and Karber.

Mouse infections

C57BL/6J mice were purchased from the Animal Resources Center, Canning Vale WA,

Australia. *Ifnar$^{-/-}$* were kindly provided by Dr P. Hertzog (Monash University, Melbourne,

Australia). *Irf7$^{-/-}$* mice were kindly provided by T. Taniguchi (University of Tokyo) (22-24).

C57BL/6N mice were purchased from The Jackson Laboratory (stock no. 005304).

C57BL/6NJ$^{\Delta Nnt8-12}$ were generated by The Australian Phenomics Network, Monash University,

Melbourne, Australia as described (25). Mice used in this study were female, except *Irf7$^{-/-}$* mice

where both males and females were used. The age/age range at infection is indicated in the

figure legends. Mice were sorted into groups so that each group had a similar mean age and age

distribution, and in the case of *Irf7$^{-/-}$* mice, had equal numbers of males and females in each

group. All mice strains except C57BL/6J were bred in-house at QIMR Berghofer MRI, and mice

housing conditions were described previously (26). Mice were infected subcutaneously (s.c.) at

the base of the tail with 100 µl of virus inoculum (doses ranging from $5x10^2$ to $5x10^5$ as

indicated in the figure legends). Serum was collected via tail nicks in to Microvette Serum-Gel

500 µl blood collection tubes (Sarstedt, Numbrecht, Germany). Mice were weighed and

monitored for disease manifestations and were humanely euthanized using CO_2 based on a score

card system (Supplementary Figure 2B). At necropsy, brain and spleens were collected for virus

titrations by $CCID_{50}$ assays and/or for histology.

Histopathology and immunohistochemistry

Brains and spleens were fixed in 10% formalin and embedded in paraffin. Human cortical brain

organoids were embedded in agarose by adding 4% melted agarose and incubating on ice to

solidify, prior to standard paraffin embedding. Sections were stained with H&E (Sigma Aldrich)

and slides were scanned using Aperio AT Turbo (Aperio, Vista, CA USA) and analyzed using

Aperio ImageScope software (LeicaBiosystems, Mt Waverley, Australia) (v10). Leukocyte

infiltrates were quantified by measuring nuclear (strong purple staining) / cytoplasmic (total red

staining) pixel ratios in scanned H&E stained images, and was undertaken using Aperio Positive

Pixel Count Algorithm (Leica Biosystems) (27).

For anti-flavivirus non-structural protein 1 (NS1) immunohistochemistry using 4G4, sections
were affixed to positively charged adhesive slides and air-dried overnight at 37°C. Sections
were dewaxed and rehydrated through xylol and descending graded alcohols to water. Sections
were transferred to Dako Epitope Retrieval Solution (pH 9.0) and subjected to heat antigen
retrieval (100°C for 20 min) using the Biocare Medical de-cloaking chamber, and slides allowed
to cool for 20 minutes before transferring to TBS plus 0.025% Tween-20 (TBS-T). Endogenous
mouse Ig was blocked by incubating sections with donkey anti-mouse Fab fragments (Jackson
Immunoresearch) diluted 1:50 in Biocare Medical Rodent block M for 60 minutes. Sections
were washed three times in TBS-T, then incubated with anti-mouse Fc for 15 minutes, before a
further three washes in TBS-T. Nonspecific antibody binding was inhibited by incubation with
Biocare Medical Background Sniper with 1% BSA and 20% donkey serum and 20% goat serum
for 15 minutes. Primary antibody 4G4 (mouse anti-flavivirus NS1 (28-30)) was diluted 1 in 4
with Da Vinci Green diluent and applied to the sections overnight at room temperature. Sections
were washed three times in TBS-T and endogenous peroxidase was blocked by incubating slides
in Biocare Medical Peroxidased 1 for 5 minutes. Sections were washed three times in TBS-T
and Perkin Elmer Opal HRP Polymer or Perkin Elmer Goat anti-mouse HRP diluted 1:500 in
TBST was applied for 60 minutes. Sections were washed three times in TBS-T and signal
developed in Vector Nova Red for 5 minutes after which they were washed three times in water.
Sections were lightly counterstained in Haematoxylin (program 7 Leica Autostainer), washed in
water, dehydrated through ascending graded alcohols, cleared in xylene, and mounted using
DePeX or similar.
Apoptosis was detected using the ApopTag Peroxidase In Situ Apoptosis Detection Kit
(Merck Catalogue No. S7100) as per manufacturer's instructions.
For anti-GFAP IHC, antigen retrieval was performed in 0.1M citric acid buffer (pH 6.0) at
1005°C for 20 min. Endogenous peroxidase activity was blocked using 1.0% H_2O_2 and 0.1%
sodium azide in TBS for 10 min. Endogenous mouse Ig was blocked by incubating sections with
goat anti-mouse Fab fragments (Jackson Immunoresearch) diluted 1:50 in Biocare Medical
Mouse block M for 60 minutes. Nonspecific antibody binding was inhibited by incubation with
Biocare Medical Background Sniper + 2% BSA for 30 min. Mouse anti-GFAP clone GA-5
(Biocare Medical, CM065C), was diluted 1:250 in the above buffer and incubated on sections for
1 hr at room temperature. After washes, Perkin Elmer Goat anti-mouse HRP diluted 1:500 in

TBST was applied for 60 minutes. Nova Red development and counter staining was performed

as for 4G4 above.

Infection of human cortical brain organoids

hBOs were reprogrammed from adult dermal fibroblasts (HDFa, Gibco, C0135C) using the

CytoTune-iPS 2.0 Sendai Reprogramming Kit (Invitrogen, A16518) (31), and were grown using

the CelVivo Clinostar incubator (Invitro Technologies) as described (32). On the day of

infection, ~30-day-old hBOs were transferred from each Clinoreactor into an ultra-low-binding

24 well plate (one hBO per well), and each hBO was infected with 10^5 $CCID_{50}$ of either

$JEV_{Nakayama}$, JEV_{FU}, $JEV_{NSW/22}$, $MVEV_{TC123130}$, Imojev (33), or YFV 17D (30) for ~4 hrs. For

shorter term culture (up to 4 days for viral growth kinetics), virus inoculum was removed and

hBOs were washed twice with media in the well, before 1 ml differentiation media was added to

each well and the 24 well plate was placed within a humidified tissue culture incubator at 37ºC,

5% CO_2 for up to 4 days. For culture up to 11 dpi, hBOs were washed twice with media by

transferring them to an ultra-low-binding 6 well plate containing 5 ml of media, and then hBOs

were transferred into 50 mm LUMOX gas exchange dishes (SARSTEDT) (4 organoids per dish)

containing 7 ml of differentiation media, and placed within a humidified tissue culture incubator

at 37ºC, 5% CO_2 for up to 11 days. hBOs were imaged using an EVOS FL (Advanced

Microscopy Group), and organoid 2D image circumference was determined by drawing around

the edge of the organoid using Image J v1.53 (34).

Human serum neutralisation assays

Cohort 1 is human serum samples from 9 participants (age >50 years), and were collected ~28

days ($\pm$ 2 days) after vaccination with the Imojev vaccine (Sanofi-Aventis Australia). Serum

was also collected pre-Imojev vaccination for these same participants for baseline analysis.

Cohort 2 is human serum samples from 10 participants (age range 24 to 60 years), collected at

variable times post-vaccination (~ 2 months to 1 year). Pre-vaccination serum was not collected

for Cohort 2. Neutralizing antibodies against Imojev, $JEV_{Nakayama}$, JEV_{FU}, $JEV_{NSW/22}$, and

$MVEV_{TC123130}$ were measured by plaque reduction neutralization (PRNT) assay (35, 36). BHK-

21 cells were seeded at $1.6{\times}10^5$ cells per well in 24 well plates overnight at 37°C. Serum

samples were heat-inactivated (56°C, 30 min) and serially diluted 4-fold from 1:5 to 1:160 in

BHK-21 cell culture media. Serum was then incubated with 100–110 pfu of Imojev, $JEV_{Nakayama}$,

JEV_{FU}, $JEV_{NSW/22}$, and $MVEV_{TC123130}$ for 1 h at 37°C. Serum plus virus mixtures were then

added to BHK-21 cell monolayers and incubated for 1 hr at 37°C to enable non-neutralized virus

to adsorb to cells. Thereafter, 1 ml of 0.375% w/v carboxymethyl cellulose (CMC, Sigma-

Aldrich)/RPMI 1640 was added, and the plates were incubated at 37°C in a CO_2 incubator for

3 days for $JEV_{Nakayama}$ and 4 days for $JEV_{NSW/22}$. The CMC medium was then removed, and the

cell monolayers were fixed and stained with 0.1% w/v crystal violet (Sigma-Aldrich) in

formaldehyde (1% v/v) and methanol (1% v/v). Plate wells were washed with tap water, dried

and the plaques were counted. The $PRNT_{50}$ titer was interpolated from plaque count compared

to the average plaque count for the naïve or no serum control.

Envelope protein structure visualisations

JEV envelope protein structure was downloaded from the Protein Data Bank (PDB: 5WSN) (37).

Envelope structure visualisations were generated using PyMol Molecular Graphics System

(version 2.3.3; Schrodinger, NY, USA). Virus sequences were downloaded from GenBank (see

Supplementary Figure 1 for accession numbers) and aligned using Mega X (Molecular

Evolutionary Genetics Analysis 10, Penn State University, State College, PA, USA) and the

ClustalW plugin with default parameters. Differences in amino acids compared to Imojev were

colored according to structural conservation as described (38).

Statistics

Statistical analyses of experimental data were performed using IBM SPSS Statistics for

Windows, Version 19.0 (IBM Corp., Armonk, NY, USA). The t-test was used when the

difference in variances was <4, skewness was > -2 and kurtosis was <2. Otherwise, the non-

parametric Kolmogorov-Smirnov test or Mann-Whitney test was used. Paired t-test was used for

comparing neutralisation of different viruses with the same human serum sample. For matched

human serum samples, a paired non-parametric Wilcoxon matched-pairs signed rank test

(GraphPad Prism 8) was used since difference in variance was >4. Kaplan-Meier statistics were

determined by log-rank (Mantel-Cox) test. Area under the curve analyses were performed in

GraphPad Prism 8, with area under the curve values then compared by t-test. Correlation

analyses of $PRNT_{50}$ with envelope conservation used the non-parametric Spearman's rank-order

correlation.

Results

JEV$_{NSW/22}$ infection produces a viremia but is not lethal in C57BL/6J mice

The JEV$_{NSW/22}$ G4 virus was isolated from the brain of a stillborn piglet in New South Wales

(NSW), Australia in February 2022 (GenBank accession OP904182). The JEV$_{Nakayama}$ genotype

3 prototype was isolated in Japan in 1935 (GenBank accession EF571853), and the JEV$_{FU}$

genotype 2 was isolated in Australia in 1995 (39, 40) (GenBank accession AF217620). MVEV

was isolated in Australia in 1974 (41) (MVEV$_{TC123130}$, GenBank accession JN119814). The

latter three viruses were isolated from human patients. Brief descriptions of the viral isolates and

their phylogenic relationships with other flaviviruses are provided in Supplementary Figure 1.

Six week old adult C57BL/6J mice were infected subcutaneously (s.c.) with 5×10^5 CCID$_{50}$ of

the aforementioned viruses. Viremia for all four viruses were broadly similar, peaking 1 day

post infection (dpi) at 2-3 log$_{10}$CCID$_{50}$/ml of serum, with nearly all mice showing no detectable

viremia by day 4 (Figure 1A-D). Infection appeared to stall weight gain for most mice until day

~10 (Figure 1E). Between 8 to 12 dpi, four mice (2 infected with JEV$_{Nakayama}$, 1 JEV$_{FU}$ and 1

MVEV$_{TC123130}$) out of the total of 24 mice showed weight loss $\geq$20% and were euthanized

(Figure 1E, †). A further four mice (1 JEV$_{Nakayama}$, 2 JEV$_{FU}$, 1 MVEV$_{TC123130}$) lost >5% of their

body weight, but subsequently recovered. None of the JEV$_{NSW/22}$ infected mice lost more than

~3% of their body weight (Figure 1E). Kaplan-Meier survival curves provided no significant

differences for the different viral isolates (Figure 1F). Mice that were euthanized also displayed

varying levels of abnormal posture (hunching), reduced activity, and fur ruffling, on the day of

euthanasia (Supplementary Figure 2A). For the 4 mice that were euthanized (Figure 1E), these

mice had very high levels of brain infection ($\approx$8-9 log$_{10}$CCID$_{50}$/mg) (Figure 1G). At the time of

euthanasia, viral titers in the spleen in these mice were below the level of detection (Figure 1G),

consistent with the viremia data.

We then infected six week old adult C57BL/6J mice with 5×10^3 CCID$_{50}$ (s.c.) of the

aforementioned viruses to determine if reduced inoculation dose can increase viremia and

neuropenetrance, as has been reported previously (42). Compared to mice infected with 5×10^5

CCID$_{50}$, viremia in mice infected with 5×10^3 CCID$_{50}$ of virus was significantly higher at later

time points for JEV$_{Nakayama}$ (Figure 1A, day 2-3), JEV$_{FU}$ (Figure 1B, day 3-4), and MVEV$_{TC123130}$

(Figure 1D, day 2-3). For JEV$_{NSW/22}$, viremia at day 3 post infection was higher on average but

this did not reach statistical significance (Figure 1C). The viremia area under the curve was

significantly higher for JEV$_{Nakayama}$ inoculated with the lower dose (Figure 1A), but was not

significantly different for the other viruses. This coincided with an increase in mortality for

JEV$_{Nakayama}$ inoculated at the lower dose, but not for the other viruses (Figure 1H-I). Viral titers

in the brains or spleens of mice that succumbed to infection were not significantly different

between inoculum doses (Figure 1G). JEV$_{NSW/22}$ had a significantly lower viremia compared to

JEV$_{Nakayama}$ and JEV$_{FU}$ (Figure 1J), consistent with significantly lower mortality compared with

JEV$_{Nakayama}$ (Figure 1I), with no C57BL/6J mice infected with JEV$_{NSW/22}$ succumbing to infection

(Figure 1F and I). Overall, this suggests that JEV$_{NSW/22}$ was significantly less virulent in

C57BL/6J mice.

C57BL/6J mice infected with JEV or MVEV led to viral neuroinvasion, apoptosis and reactive

astrogliosis

The brains of C57BL/6J mice were analysed by immunohistochemistry (IHC) using the pan-

flavivirus anti-NS1 antibody (4G4). This antibody detects the viral non-structural protein 1,

which is a highly conserved multi-functional protein important for flavivirus replication (43).

Staining was consistently seen in the cortex for JEV and MVEV infected mice, with staining also

seen in the thalamus (Figure 2A, Supplementary Figure 3A-C). Prevailing infection of the

cerebral cortex and thalamus parallels IHC data from post-mortem human brains (6, 44). In

some mice, virus was detected in other brain regions, including the hippocampus, anterior

olfactory nucleus (Figure 2A, JEV$_{Nakayama}$), and caudate putamen (Supplementary Figure 3B,

JEV$_{FU}$). A JEV$_{FU}$ infected mouse that lost ~15% body weight and then recovered (Figure 1E)

had residual virus staining in the cortex at day 29 post infection (Supplementary Figure 3D),

while a mouse that survived infection with minimal weight loss had no detectable viral antigen

staining (Supplementary Figure 3E). As expected, 4G4 staining was associated primarily with

cells showing neuronal morphology (Figure 2A, Supplementary Figure 3D).

ApopTag staining was evident in areas that were positive for virus staining, particularly in the

cortex, and was again associated with cells showing neuronal morphology (Figure 2B and

Supplementary Figure 4A-C). Apoptotic cells were also evident in the mouse that showed ~15%

body weight loss and recovered (Supplementary Figure 4D). Mice with signs of disease from

JEV or MVEV infection (Supplementary Figure 2) thus showed high levels of viral antigen and

apoptosis in the brain. In addition, like in humans, animals can recover from brain infection,

despite apoptotic damage, although neurological sequelae may ensue.

Astrocytes are a type of glial cell that provides physical and chemical support to neurons.

Astrocytes become activated (reactive astrogliosis) in response to brain pathology, including

from JEV infection (45-47). Reactive astrocytes are characterized by the upregulated expression

of glial fibrillary acidic protein (GFAP) and the growth/extension of cellular processes (48).

Brains from mice with signs of disease from JEV or MVEV infection (Supplementary Figure 2)

had significant upregulation of GFAP positive astrocytes throughout the brain, including the

heavily infected areas of the cortex, thalamus and anterior olfactory nucleus (Figure 2C,

Supplementary Figure 5). GFAP is constitutively expressed in astrocytes of the hippocampus,

corpus callosum, and cerebral peduncle (49), consistent with staining seen in uninfected brains

(Figure 2D, right). Widespread reactive astrogliosis is thus a feature of lethal JEV and MVEV

encephalitis in C57BL/6J mice.

JEV$_{NSW/22}$ is less virulent than JEV$_{Nakayama}$ and JEV$_{FU}$ in Irf7$^{-/-}$ mice

C57BL/6J mice had a relatively low viremia and low penetrance of lethal brain infection for

JEV$_{NSW/22}$ (Figure 1). A model with a higher penetrance of brain infection is clearly desirable,

given encephalitis is the key clinical concern for JEV infections. Interferon regulatory factor 7

knockout (*Irf7$^{-/-}$* mice) were thus chosen as these mice enhanced neuroinvasion for WNV (50), as

well as another arbovirus (24). We first sought to optimize the inoculation dose for JEV$_{NSW/22}$ in

Irf7$^{-/-}$ mice by comparing 5×10^5, 5×10^4, 5×10^3, and 5×10^2 CCID$_{50}$. The inoculation dose of 5×10^3

CCID$_{50}$ provided the highest viremia area under the curve, which was significantly higher than

5×10^4 and 5×10^5 CCID$_{50}$ inoculation doses (Figure 3A). The highest infection dose (5×10^5) led

to the highest peak viremia on day 1 post infection, but was cleared significantly more quickly

than lower infection doses (Figure 3A, p=0.004). Our data agrees with the concept of an

'optimal dose' that is not too high as to excessively stimulate type-I IFN, but is high enough to

establish a robust viremia (42). One mouse infected with 5×10^3 CCID$_{50}$ JEV$_{NSW/22}$ succumbed to

infection (Figure 3C-D), while all other mice survived infection. An inoculation dose of 5×10^3

CCID$_{50}$ was thus chosen to compare JEV and MVEV strains in *Irf7$^{-/-}$* mice.

$JEV_{NSW/22}$, $JEV_{Nakayama}$ and JEV_{FU} infection of $Irf7^{-/-}$ mice (dose 5×10^3 $CCID_{50}$) resulted in

robust viremias, while $MVEV_{TC123130}$ viremia was significantly lower than JEV (Figure 3B,

$p<0.0001$). $JEV_{NSW/22}$ had a significantly lower viremia (area under the curve) compared to

JEV_{FU} (Figure 3E, $p=0.004$). Infection of $Irf7^{-/-}$ mice with the different JEV isolates and MVEV

(n=14 mice per group, n=56 total infected mice) illustrated a mean weight loss of 5-10% for

surviving mice across all four groups (Figure 3C), with the exception of 20 mice that reached

ethically defined end points for weight loss and/or disease manifestations (Supplementary Figure

6). Kaplan-Meier curves illustrated significantly higher survival for $Irf7^{-/-}$ mice infected with

$JEV_{NSW/22}$ or $MVEV_{TC123130}$ when compared with $JEV_{Nakayama}$ and JEV_{FU}, with only 1/14

$JEV_{NSW/22}$ or $MVEV_{TC123130}$ infected $Irf7^{-/-}$ mice requiring euthanasia (Figure 3D). These data are

consistent with the significantly lower viremias of $JEV_{NSW/22}$ and $MVEV_{TC123130}$ infected $Irf7^{-/-}$

mice (Figure 3B). On average, mice that succumbed to infection had a more robust viremia

(Figure. 3E), further implicating a higher viremia in a higher chance of lethal neuropenetrance in

$Irf7^{-/-}$ mice.

Brain titers in euthanized $Irf7^{-/-}$ mice were ~4-9 $\log_{10}CCID_{50}$/gram, while spleen titers were

<2-5 $\log_{10}CCID_{50}$/g (Figure 3F). IHC staining for viral antigen confirmed JEV infection in the

brains of $Irf7^{-/-}$ mice, with reduced levels of staining reflecting lower brain titers (Figure 3G-H,

Supplementary Figure 7). Viral antigen staining was consistently present in the cortex, with

staining also seen in the medulla, thalamus, and hippocampus (Figure 3G-H, Supplementary

Figure 7). Brain regions showing ApopTag staining broadly overlapped with staining for viral

antigen (Figure 3G-H). Reactive astrogliosis was also evident (Supplementary Figure 5).

Overall, this suggests that JEV infection of $Irf7^{-/-}$ mice produced a consistent viremia, with mice

showing brain infection and apoptotic damage.

$JEV_{NSW/22}$ and JEV_{FU} are more sensitive to type I IFN compared to $JEV_{Nakayama}$ and

$MVEV_{TC123130}$

Viremia and survival were compared between C57BL/6J and $Irf7^{-/-}$ mouse strains infected with

5×10^3 $CCID_{50}$ virus to determine whether differences in viremia may be associated with the

differences in viral neuropenetrance and survival. Area under the curve analyses showed a

significantly increased/prolonged viremia in $Irf7^{-/-}$ mice compared to C57BL/6J mice for all JEV

strains (Figure 4A-C). However, $MVEV_{TC123130}$ viremia was not significantly improved in $Irf7^{-/-}$

mice (Figure 4D). Kaplan-Meier survival curves indicated that JEV$_{FU}$ had a significantly

increased mortality in *Irf7$^{-/-}$* mice compared to C57BL/6J mice (Figure 4F), while there was no

significant difference between mouse strains for the other viruses (Figure 4E, G-H).

Differences in sensitivity to type I IFN between virus strains may explain the different

outcomes in C57BL/6J versus *Irf7$^{-/-}$* mice. To examine type I IFN sensitivity in the absence of *in*

vivo adaptive immune responses, mouse embryonic fibroblasts (MEFs) with (wild-type) and

without (*Irf3/7$^{-/-}$*) functional type I IFN production were infected with JEV$_{Nakayama}$, JEV$_{FU}$,

JEV$_{NSW/22}$ and MVEV$_{TC123130}$. In wild-type MEFs, JEV$_{FU}$ and JEV$_{NSW/22}$ had significantly lower

virus replication kinetics compared to JEV$_{Nakayama}$ and MVEV$_{TC123130}$ (Figure 4I). In *Irf3/7$^{-/-}$*

MEFs, the difference in replication between virus strains was diminished, with JEV$_{FU}$ not

significantly different to JEV$_{Nakayama}$ (Figure 4J). This suggests that JEV$_{FU}$ and JEV$_{NSW/22}$ were

more sensitive to type I IFN compared to JEV$_{Nakayama}$ and MVEV$_{TC123130}$, consistent with an

increase in mortality in *Irf7$^{-/-}$* mice (partially defective type I IFN responses) compared to

C57BL/6J mice for JEV$_{FU}$ and JEV$_{NSW/22}$ but not for JEV$_{Nakayama}$ or MVEV$_{TC123130}$.

Ifnar$^{-/-}$ mice are a model of lethal JEV viremia without neuroinvasion

As JEV$_{NSW/22}$ did not lead to fatal neuroinvasion in C57BL/6J mice, and fatal neuroinvasion in

Irf7$^{-/-}$ mice was rare, type I interferon receptor knockout (*Ifnar$^{-/-}$*) mice were evaluated. *Ifnar$^{-/-}$*

mice have been used extensively in studies of pathogenic flaviviruses such as Zika virus (ZIKV),

West Nile virus (WNV), and Yellow fever virus (YFV), and generally provide a robust viremia

and a lethal outcome (21, 29, 30, 51). Adult *Ifnar$^{-/-}$* mice were infected with 5×10^5 CCID$_{50}$ of

JEV$_{Nakayama}$, JEV$_{FU}$, JEV$_{NSW/22}$, or MVEV$_{TC123130}$ via s.c. injection. A robust viremia developed,

reaching 6-8 log$_{10}$CCID$_{50}$/ml of serum by 2 dpi (Figure 5A). JEV$_{NSW/22}$ infected mice had ≈2 log

lower viremia compared to JEV$_{Nakayama}$ and JEV$_{FU}$ on 3 dpi (Figure 5A). Mice displayed a rapid

loss of body weight, which was slightly delayed for JEV$_{NSW/22}$ and MVEV$_{TC123130}$ at 2 dpi

(Figure 5B), requiring euthanasia 2/3 dpi (Figure 5C). Mice infected with any of the JEV strains

also displayed varying levels of abnormal posture (hunching), reduced activity, and fur ruffling

(Supplementary Figure 8A).

Consistent with the robust viremia (Figure 5A), spleen tissue titers reached ~6-9

log$_{10}$CCID$_{50}$/g at 2-3 dpi (Figure 5D). Similar levels were seen in brains (Figure 5E); however,

4G4 staining illustrated that brain cells were not infected (Figure 5F). Viremia and brain titers

correlated strongly (Supplementary Figure 8B), arguing that the brain titers (Figure 5E) likely

arose from virus in the blood vessels of the brain. Interestingly, anti-flavivirus NS1 antibody

staining was clearly present in blood vessels (Figure 5G,H), with extracellular JEV NS1 also

found in serum of JEV patients (52). Secreted NS1 from JEV infected cells is reported to

increase vascular leakage and may contribute to mortality (53), although overt vascular leakage

of NS1 was not evident in IHC, with lack of type I IFN signalling potentially involved (54, 55).

Neither apoptosis (Supplementary Figure 8C), nor significantly increased staining for reactive

astrocytes (Supplementary Figure 5, *Ifnar*$^{-/-}$) were apparent for brains of JEV-infected *Ifnar*$^{-/-}$

mice. Therefore, *Ifnar*$^{-/-}$ mice infected with JEV or MVEV represent a model of lethal viremia

without viral neuroinvasion.

JEV causes severe histological lesions in C57BL/6J and Irf7*$^{-/-}$ *mouse brains

H&E staining of JEV infected mouse brains revealed a series of histopathological lesions,

predominantly neuronal degeneration with pyknotic nuclei (often associated with apoptosis),

neuronal vacuolation, perivascular cuffing, leukocyte infiltrates, hemorrhage, meningitis and

microgliosis (Figure 6A). IHC for the microglial marker, Iba1, confirmed the presence of

microgliosis (Figure 6B).

The presence of H&E lesions were scored for the different JEV isolates and mouse strains

(Supplementary Figure 9). C57BL/6J and *Irf7*$^{-/-}$ mice that were euthanized at the time when viral

neuroinvasion led to the criteria for humane euthanasia being met (full score cards provided in

Supplementary Figure 2 and 7) had a broadly similar incidence of the aforementioned lesions,

whereas only hemorrhage was seen in *Ifnar*$^{-/-}$ mice (Supplementary Figure 9, Acute). Leukocyte

infiltration was quantified by calculating the ratio of nuclear/cytoplasmic staining, as leukocytes

have a higher ratio of nuclear to cytoplasmic staining (27). This confirmed that C57BL/6J and

Irf7$^{-/-}$ mice that succumbed to infection had significantly increased leukocyte infiltrates

(Supplementary Figure 9B). For C57BL/6J and *Irf7*$^{-/-}$ mice that survived and were euthanized

later (nominally chronic phase), only hemorrhage was consistently observed (Supplementary

Figure 9, Chronic).

Lesions in C57BL/6J and *Irf7*$^{-/-}$ mouse brains were associated with areas of high virus

infection, most prominently in the cortex (Supplementary Figure 10); however, lesions were also

found in other areas of the brain including where there was minimal viral antigen staining

(Supplementary Figure 11A). Some mice that lost >8% body weight and then recovered had

persistent lesions detectable at the latest time point sampled (up to day 32 for C57BL/6J and day

21 for $Irf7^{-/-}$) (Supplementary Figure 11B), consistent with persistent neurological sequelae in

humans that survive infection. Mice that did not lose any body weight did not show any overt

brain lesions at the later time points, indicating that brain infection causing weight loss is

associated with persistent lesions.

Overall, the lesions in C57BL/6J and $Irf7^{-/-}$ mouse brains are consistent with H&E detectable

lesions in post-mortem human JEV infected brains (6, 45, 56-58). To provide additional

potential insights into the nature of the cellular infiltrates (e.g. perivascular cuffing and leukocyte

infiltrates, Figure 6), immune cell type abundance estimates were obtained from reanalyzed (59)

publically available JEV-infected mouse brain RNA-Seq expression data (60) using

SpatialDecon (61). The dominant cell types identified were CD4 T cells, NKT cells,

monocytes/macrophages, neutrophils, innate lymphoid cells, T regs, and CD8 T cells, with an

increase in microglia cells also identified (Supplementary Figure 12). GFAP was significantly

up-regulated 5.4 fold by infection (Supplementary Table 3, log_2FC 2.4), consistent with the

GFAP IHC (Figure 2C) and reactive astrogliosis seen herein. Iba1 (Aif1) was also significantly

up-regulated 5.5 fold (Supplementary Table 3, log_2FC 2.5), consistent with the H&E and IHC

(Figure 6) and microgliosis observed in our study.

Productive replication of JEV in human cortical brain organoids

Human cortical brain organoids (hBOs) have been used to investigate the pathogenesis of

flaviviruses, including JEV and ZIKV (21, 62, 63). We generated ~30 day old hBOs from adult

human dermal fibroblast (HDFa) cells, growing the hBOs in a CelVivo Clinostar incubator as

described (32) (Figure 7A). hBOs were infected with 10^5 $CCID_{50}$ of the indicated JEV and

MVEV isolates, as well as (i) the Imojev chimeric virus vaccine (previously called ChimeriVax-

JE) comprising the prME genes of the attenuated $JEV_{SA14-14-2}$ strain on the YFV 17D backbone

(33, 64) and (ii) the Yellow Fever live attenuated vaccine strain (YFV 17D) (30), with wild-type

YFV infection (65), and occasionally YFV 17D vaccination (66), able to cause neuropathology.

hBOs were fixed in formalin 4 dpi and IHC undertaken using the anti-NS1 monoclonal antibody

4G4 (30). $JEV_{Nakayama}$ infected hBOs showed the most pronounced viral antigen staining, with

$JEV_{NSW/22}$, JEV_{FU} and $MVEV_{TC123130}$ also showing clear staining (Figure 7B). Viral antigen was

primarily localized to the outer surface of the organoids (Figure 7B), where cells are in direct

contact with the culture medium. Viral antigen staining for Imojev and YFV 17D infected hBOs

was also seen, but was sparse (Figure 7B and Supplementary Figure 13A).

All viruses were able to productively infect hBOs, with $JEV_{Nakayama}$ infection generating

higher viral titers than Imojev at all time points (Figure 7C, p<0.001). By 96 h $JEV_{Nakayama}$ titers

were also significantly higher than those seen for $JEV_{NSW/22}$ (Figure 7C).

Over 11 days, uninfected hBOs grew in circumference by ~20% (Figure 7D, Uninfected).

hBOs infected with JEV isolates shrank in circumference by ~5% to 15% by 11 dpi, although

when compared with uninfected organoids, this only approached significance for $JEV_{NSW/22}$

(Figure 7D; Supplementary Figure 13B-C). hBOs infected with $MVEV_{TC123130}$, Imojev and YFV

17D infected hBO did not shrink significantly when compared with uninfected controls, although

the $MVEV_{TC123130}$ data suggested a marginal reduction in circumference (Figure 7C;

Supplementary Figure 13C). These data (Figure 7D) reflect differences in viral replication

(Figure 7C) and/or viral protein immunohistochemistry (Figure 7B) and likely reflect virus-

induced CPE.

All viruses replicated productively in a human neural progenitor cell line, RENcell VM, with

all viruses except Imojev causing fulminant CPE (Supplementary Figure 14A,B). Although

Vero E6 cells are widely used for flavivirus research, CPE induced by $JEV_{NSW/22}$ was

considerably less pronounced in these cells (Supplementary Figure 14C).

Human post Imojev-vaccination sera neutralizes JEV and MVEV with titers related to

envelope protein amino acid conservation

Imojev is one of two JEV vaccines available in Australia. The Imojev prME genes are derived

from the genotype 3 $JEV_{SA14-14-2}$ (33, 64) strain, which was attenuated via extensive *in vitro* and

in vivo passaging (Supplementary Figure 1). Most flavivirus neutralizing antibodies recognize

epitopes on the envelope protein, particularly in the putative receptor binding domain III (67).

$JEV_{Nakayama}$ and the $JEV_{SA14-14-2}$ component of the Imojev vaccine both belong to genotype 3, but

$JEV_{Nakayama}$ has 96.8% envelope protein amino acid identity to Imojev (Figure 8A). JEV_{FU} has

96.4% envelope protein identity, while $JEV_{NSW/22}$ has drifted further from the genotype 2 and 3

strains with 93.4% envelope protein identity (Figure 8A). $MVEV_{TC123130}$, which is the closest

phylogenetically related flavivirus to JEV (Supplementary Figure 1B), has 80.4% envelope

amino acid identity (Figure 1A). Alignment of the envelope amino acid differences for these

strains compared to Imojev reveal that a disproportionate number of the non-conservative

changes were in domain III (Figure 8A).

Serum neutralizing antibodies post-vaccination is currently viewed as the best measurable

correlate of vaccine protection for JEV (68). To determine if the Australian outbreak genotype 4

$JEV_{NSW/22}$ is neutralized by antibodies produced in response to Imojev vaccination, serum was

taken from two human cohorts. Cohort 1 was serum collected from individuals at approximately

28 days post-vaccination, while cohort 2 serum was collected at variable times post-vaccination

(2 months to 1 year). Ethical approvals for cohort 1 serum allowed for neutralisation assays

against the Imojev vaccine itself, $JEV_{Nakayama}$, JEV_{FU}, and $JEV_{NSW/22}$ (Figure 8B), while ethical

approval for cohort 2 additionally allowed for neutralisation assays against $MVEV_{TC123130}$

(Figure 8C). All serum samples from both cohorts had a measurable 50% neutralisation titer

($PRNT_{50}$) titer against $JEV_{Nakayama}$, JEV_{FU}, and $JEV_{NSW/22}$ (Figure 8B-C), indicating that

seroconversion provided neutralizing antibodies that cross-react between JEV genotypes 2, 3 and

4. In cohort 1, $PRNT_{50}$ titers were significantly lower against $JEV_{Nakayama}$ and $JEV_{NSW/22}$

compared to against Imojev (Figure 8A), suggesting key antigenic differences from the vaccine.

In cohort 2, $PRNT_{50}$ titers against $JEV_{Nakayama}$ were not significantly different compared to

Imojev, however $PRNT_{50}$ titers were significantly lower for $JEV_{NSW/22}$ (Figure 8C). Similar

conclusions were drawn when the raw percentage of plaque neutralisation at a high serum

dilution (1:160) was compared between virus strains (Supplementary Figure 15). With $PRNT_{50}$

data from both cohorts combined, the percentage amino acid identity of the envelope protein

compared to the Imojev vaccine significantly correlated with $PRNT_{50}$ titers (Figure 8D, black

line). The significance (p-value) and the correlation coefficient (rho), were similar when the

same analysis was conducted excluding the $PRNT_{50}$ data for $MVEV_{TC123130}$ (Figure 8D, brown

line). Overall, our data indicates that Imojev vaccination provided neutralizing antibodies

against $JEV_{NSW/22}$ in all individuals, but the level of cross-neutralisation were related to the

conservation in envelope protein amino acid sequences.

Discussion

Herein we provide a comprehensive *in vivo* and *in vitro* characterization of the genotype 4

$JEV_{NSW/22}$ isolate from the recent Australian outbreak, and illustrate mouse models of infection

and rare CNS neuropathological manifestations that recapitulate many aspects of human and
primates disease (6, 44-47, 69). The capacity of $JEV_{NSW/22}$ to cause lethal neuroinvasive
infection in mice was significantly diminished compared to $JEV_{Nakayama}$ and JEV_{FU}, with only
one $Irf7^{-/-}$ mouse succumbing to $JEV_{NSW/22}$ infection out of 63 infected C57BL/6J or $Irf7^{-/-}$ mice.
Such rare lethal neuroinvasion recapitulates what is seen in humans, with approximately 1 in 750
infections causing fatality (2, 3). Serosurvey data, albeit limited, suggests the ratio of human
symptomatic to asymptomatic infections is not particularly different for $JEV_{NSW/22}$. To date 45
clinical cases have been notified for the recent outbreak in Australia (70), with serosurveys in
Victoria (n=820 participants) and New South Wales (NSW) (n=917 participants) reporting 3.3%
and 8.7% of participants as seropositive for JEV, respectively (71, 72). Although somewhat
dated, population data for the primary recruitment locations for the serosurveys is available from
Australian Bureau of Statistics 2016, with Victorian recruitment locations providing a population
total of 160,294 (Mildura, Lockington, Shepparton, Cobram, Yarrawonga, Rutherglen,
Wodonga, Wangaratta, Rochester), and NSW locations a total of 68,431 (Balranald, Corowa,
Dubbo, Griffith, Temora). As [0.033 x 160,294] + [0.087 x 68,431] / 250 = 45, the serosurvey
data is consistent with the expected symptomatic to asymptomatic ratio of ≈1 in 250 for JEV and
thus provides no compelling evidence for overt virulence differences for $JEV_{NSW/22}$ in human
populations.
Our $Irf7^{-/-}$ mouse model of $JEV_{NSW/22}$ provides for a more robust viremia, and a slightly higher
chance of lethal neuroinvasive infection. Increased lethal neuro-penetrance in $Irf7^{-/-}$ mice was
associated with a prolonged viremia, possibly via increased inflammation-driven blood brain
barrier breakdown as a result (5, 73). The use of $Irf7^{-/-}$ mice to increase lethal neuroinvasive
infection compared to C57BL/6J was only suitable for JEV_{FU} and $JEV_{NSW/22}$. This was likely
due to higher sensitivity to type I IFN for these isolates, demonstrated using MEF cells,
compared to $JEV_{Nakayama}$ and $MVEV_{TC123130}$. The partially defective type I IFN responses in $Irf7^{-}$
$^{/-}$ mice (24) thus provides a benefit for JEV_{FU} and $JEV_{NSW/22}$ neuro-penetrance, but not for
$JEV_{Nakayama}$ or $MVEV_{TC123130}$. When type I IFN responses were completely defective ($Ifnar^{-/-}$
mice and $Irf3/7^{-/-}$ MEFs), differences between virus replication and/or lethality were minimal.
Overall, these results suggest that $JEV_{NSW/22}$ may be more sensitive to, or less able to suppress,
type I IFN responses. Inhibition of type I IFN responses is mediated by NS5 for inhibition of
STAT2 and NS3 (74) and subgenomic flavivirus RNA (sfRNA)/NS5 (21) for inhibition of

STAT1. Although the latter likely operates for WNV in mice (75) and is involved in promoting

apoptosis (21), the efficiency of these systems during JEV infection of humans and mice remains

to be determined. sfRNA is derived from the 3'UTR (21), where JEV$_{NSW/22}$ does show a small

number of nucleic acid changes, but which are unlikely to affect sfRNA production

(Supplementary Table 3). Mouse models of flavivirus pathogenesis frequently use *Ifnar*$^{-/-}$ mice

(30, 76, 77) as flaviviruses often replicate poorly in wild-type mice as the ability of NS5 to

suppress the antiviral type I IFN responses in humans often fails to operate in mice (78). We

show herein that *Ifnar*$^{-/-}$ mice are not a good model for JEV neuropathology, as they reach

ethically defined endpoints before brain infection can occur. However *Ifnar*$^{-/-}$ mice are a good

model of robust and lethal viremia, which may provide a useful and stringent model for vaccine

testing.

JEV$_{NSW/22}$ does not contain known attenuating mutations that would explain its reduced

virulence in mice. There are 9 amino acids in the Imojev vaccine envelope gene that have been

associated with attenuated neurovirulence; F107, K138, V176, A177, G244, H264, M279, V315

and R439 (79-89). At these positions, JEV$_{Nakayama}$, JEV$_{FU}$, JEV$_{NSW/22}$, and MVEV$_{TC123130}$ all

have the same amino acids (L107, E138, T177, E244, Q264, K279, A315, and K439), except for

position 176 where JEV$_{Nakayama}$ has T176, and JEV$_{FU}$, JEV$_{NSW/22}$, and MVEV$_{TC123130}$ have I176

(Supplementary Table 3). JEV$_{NSW/22}$ retains E at position 138, and this amino acid has been

identified by several studies as a principal neurovirulence determinant (90-94), with a role in

neuronal cell binding hypothesized (37). YFV 17D has a valine (V) at this residue, possibly

contributing to reduced neurovirulence (95). NS1 and NS2A have been implicated in

neuroinvasion, but not neurovirulence (88). Among the six changes in NS1 associated with

attenuation of neuroinvasiveness are R147H and R339M (88), of which H147 is present in both

JEV$_{Nakayama}$ and JEV$_{NSW/22}$, with K339 (a conserved substitution for R) present in JEV$_{NSW/22}$.

JEV non-structural protein 4B (NS4B) alone can induce apoptosis and encephalitis (96),

however, NS4B is completely conserved between JEV$_{Nakayama}$, JEV$_{FU}$, and JEV$_{NSW/22}$

(Supplementary Table 3). prM has been reported to influence neuroinvasiveness of ZIKV (76)

and JEV virulence in mice (97). JEV$_{NSW/22}$ has a number of unique changes in prM

(Supplementary Table 3), although their functional implications remain unclear. JEV$_{NSW/22}$ also

has an additional N-linked glycosylation site at position 175 in NS1 that is lacking in JEV$_{Nakayama}$

and JEV$_{FU}$ (Supplementary Table 3). However, this N-linked glycosylation site is reported to

increase WNV neuroinvasiveness in wild-type mice (98, 99), a trend not seen in our JEV data

(Figure 1F). Thus JEV$_{NSW/22}$ shows no obvious sequence characteristics that can be ready

associated with the reduced virulence in mice, and mutagenesis experiments would be required

to fully understand these differences.

All human participants vaccinated with the Imojev vaccine induced neutralizing antibodies to

JEV$_{NSW/22}$, suggesting that this vaccine, which is available in Australia, is likely to afford some

protection against Australian outbreak genotype 4 JEV. However, divergence of envelope

protein amino acid sequences from that of the Imojev vaccine affected the PRNT$_{50}$ titers,

although it is unclear how this may translate to the impact on vaccine efficacy, especially given

that *in vitro* neutralisation assays do not capture the full range of protective mechanisms

mobilised *in vivo* (100). Nonetheless, this provides a strong rationale for development of

updated JEV vaccines that use antigen sequences from currently circulating JEV strains, such as

genotype 4 in Australia (14), genotype 5 in Republic of Korea (12), and genotype 1 in most other

areas of South East Asia (101). Imojev vaccination also produced neutralizing antibodies against

MVEV$_{TC123130}$ which is consistent with previous studies using other JEV vaccines (102, 103),

and is consistent with cross-reactivity in serology-based diagnostic assays (13, 104). There is

also some evidence that JEV vaccination or infection provides partial cross-protection against

MVEV and vice versa (105-107).

Although one limitation of this study may be that JEV$_{NSW/22}$ was isolated from a pig, there are

only 4 amino acid differences between this isolate and a JEV G4 sequence from a human brain in

the Tiwi Islands (Northern Territory, Australia) in 2021 (Genbank accession OM867669 (14)).

The differences are; envelope-238 F vs. L, NS2A-71 I vs. T, NS2B-59 E vs. G, NS3-436 E vs. G.

In addition, JEV$_{Nakayama}$ was passaged in suckling mouse brains, which may contribute to the

increased neuroinvasion in C57BL/6J mice, although the adaptive mutations acquired during

passaging, if any, are currently unknown. Furthermore, JEV$_{FU}$ has not been passaged in mice,

but was still more lethal than JEV$_{Nakayama}$ in *Irf7$^{-/-}$* mice. Use of C57BL/6J mice that lack a

functional nicotinamide nucleotide transhydrogenase *(Nnt)* may be another issue, as background

and *Nnt* are able to affect viral immunopathogenesis (25, 59). However, we found that neither

Nnt nor a C57BL/6N genetic background significantly impacted JEV replication or

immunopathology (Supplementary Figure 16).

In conclusion, we show that JEV$_{NSW/22}$ has reduced neuropenetrance in mice but retains

capacity for rare lethal neuroinvasion, consistent with reported human fatalities in the 2022

Australian outbreak.

AUTHOR CONTRIBUTIONS

Conceptualization, D.J.R. Methodology, D.J.R., W.N., A.S, G.D., N.G., and N.W. Formal

analysis, W.N., D.J.R. and A.S. Investigation, W.N., B.T., R.S., A.C., K.Y., N.G., T.L., and

D.J.R. Resources, R.S., N.G., G.D., N.W., and A.S. Data curation, W.N., D.J.R., N.G., and A.S.

Writing, original draft – D.J.R. Writing, review and editing, A.S and W.N. Visualization,

D.J.R., W.N. and A.S. Supervision, D.J.R., G.D. and A.S. Project administration, D.J.R., G.D.,

N.W., and A.S. Funding acquisition, A.S., D.J.R, and G.D, and N.W.

DECLARATION OF INTERESTS

The authors declare no competing interests.

ACKNOWLEDGEMENTS

From QIMR Berghofer MRI we thank Dr I Anraku for managing the PC3 (BSL3) facility,

animal house staff for mouse breeding and agistment, Drs Anthony White and Lotta Oikari for

providing the iPSC line, Crystal Chang, Sang-Hee Park and Ashwini Potadar for the histology

and immunohistochemistry, Dr Viviana Lutzky for proof reading, and Dr. Gunter Hartel (Head

of Statistics, QIMRB) for assistance with statistics. We thank Dr. Peter Kirkland for supplying

the $JEV_{NSW/22}$ isolate. We thank Dr. Andrii Slonchak for insights into whether 3'UTR changes

may affect sfRNA production.

DATA AVAILABILITY

All data is provided in the manuscript and accompanying supplementary files.

References

1.   van den Hurk AF, Ritchie SA, Mackenzie JS. Ecology and Geographical Expansion of Japanese
Encephalitis Virus. Annual Review of Entomology. 2008;54(1):17-35.
2.   Ashraf U, Ding Z, Deng S, Ye J, Cao S, Chen Z. Pathogenicity and virulence of Japanese
encephalitis virus: Neuroinflammation and neuronal cell damage. Virulence. 2021;12(1):968-80.
3.   Solomon T, Dung NM, Kneen R, Gainsborough M, Vaughn DW, Khanh VT. Japanese encephalitis.
Journal of Neurology, Neurosurgery & Psychiatry. 2000;68(4):405.

4.   Quan TM, Thao TTN, Duy NM, Nhat TM, Clapham H. Estimates of the global burden of Japanese
encephalitis and the impact of vaccination from 2000-2015. eLife. 2020;9:e51027.
5.   Hsieh JT, Rathore APS, Soundarajan G, St John AL. Japanese encephalitis virus neuropenetrance
is driven by mast cell chymase. Nat Commun. 2019;10(1):706.
6.   Johnson RT, Burke DS, Elwell M, Leake CJ, Nisalak A, Hoke CH, et al. Japanese encephalitis:
immunocytochemical studies of viral antigen and inflammatory cells in fatal cases. Ann Neurol.
1985;18(5):567-73.
7.   Hegde NR, Gore MM. Japanese encephalitis vaccines: Immunogenicity, protective efficacy,
effectiveness, and impact on the burden of disease. Hum Vaccin Immunother. 2017;13(6):1-18.
8.   Gao X, Liu H, Li X, Fu S, Cao L, Shao N, et al. Changing Geographic Distribution of Japanese
Encephalitis Virus Genotypes, 1935-2017. Vector Borne Zoonotic Dis. 2019;19(1):35-44.
9.   Kang BK, Hwang JM, Moon H, Han SY, Kim JM, Yang DK, et al. Comparison of the antigenic
relationship between Japanese encephalitis virus genotypes 1 and 3. Clin Exp Vaccine Res. 2016;5(1):26-
30.
10.   Han N, Adams J, Chen P, Guo ZY, Zhong XF, Fang W, et al. Comparison of genotypes I and III in
Japanese encephalitis virus reveals distinct differences in their genetic and host diversity. J Virol.
2014;88(19):11469-79.
11.   Mulvey P, Duong V, Boyer S, Burgess G, Williams DT, Dussart P, et al. The Ecology and Evolution
of Japanese Encephalitis Virus. Pathogens. 2021;10(12).
12.   Lee AR, Song JM, Seo SU. Emerging Japanese Encephalitis Virus Genotype V in Republic of Korea.
J Microbiol Biotechnol. 2022;32(8):955-9.
13.   Pham D, Howard-Jones AR, Hueston L, Jeoffreys N, Doggett S, Rockett RJ, et al. Emergence of
Japanese encephalitis in Australia: a diagnostic perspective. Pathology. 2022;54(6):669-77.
14.   Sikazwe C, Neave MJ, Michie A, Mileto P, Wang J, Cooper N, et al. Molecular detection and
characterisation of the first Japanese encephalitis virus belonging to genotype IV acquired in Australia.
PLoS Negl Trop Dis. 2022;16(11):e0010754.
15.   Mackenzie JS, Williams DT, van den Hurk AF, Smith DW, Currie BJ. Japanese Encephalitis Virus:
The Emergence of Genotype IV in Australia and Its Potential Endemicity. Viruses [Internet]. 2022; 14(11).
16.   Yakob L, Hu W, Frentiu FD, Gyawali N, Hugo LE, Johnson B, et al. Japanese encephalitis
emergence in Australia: the potential population at risk. Clin Infect Dis. 2022.
17.   Gyawali N, Taylor-Robinson AW, Bradbury RS, Pederick W, Faddy HM, Aaskov JG. Neglected
Australian Arboviruses Associated With Undifferentiated Febrile Illnesses. Front Microbiol.
2019;10:2818.
18.   Bharucha T, Cleary B, Farmiloe A, Sutton E, Hayati H, Kirkwood P, et al. Mouse models of
Japanese encephalitis virus infection: A systematic review and meta-analysis using a meta-regression
approach. PLOS Neglected Tropical Diseases. 2022;16(2):e0010116.
19.   Tajima S, Maeki T, Nakayama E, Faizah AN, Kobayashi D, Isawa H, et al. Growth, Pathogenesis,
and Serological Characteristics of the Japanese Encephalitis Virus Genotype IV Recent Strain 19CxBa-83-
Cv. Viruses. 2023;15(1):239.
20.   Johnson BJ, Le TT, Dobbin CA, Banovic T, Howard CB, Flores Fde M, et al. Heat shock protein 10
inhibits lipopolysaccharide-induced inflammatory mediator production. J Biol Chem. 2005;280(6):4037-
47.
21.   Slonchak A, Wang X, Aguado J, Sng JDJ, Chaggar H, Freney ME, et al. Zika virus noncoding RNA
cooperates with the viral protein NS5 to inhibit STAT1 phosphorylation and facilitate viral pathogenesis.
Science Advances. 2022;8(48):eadd8095.
22.   Honda K, Yanai H, Negishi H, Asagiri M, Sato M, Mizutani T, et al. IRF-7 is the master regulator of
type-I interferon-dependent immune responses. Nature. 2005;434(7034):772-7.

23.   Sato M, Suemori H, Hata N, Asagiri M, Ogasawara K, Nakao K, et al. Distinct and essential roles
of transcription factors IRF-3 and IRF-7 in response to viruses for IFN-alpha/beta gene induction.
Immunity. 2000;13(4):539-48.
24.   Rudd PA, Wilson J, Gardner J, Larcher T, Babarit C, Le TT, et al. Interferon response factors 3 and
7 protect against Chikungunya virus hemorrhagic fever and shock. J Virol. 2012;86(18):9888-98.
25.   Rawle DJ, Le TT, Dumenil T, Bishop C, Yan K, Nakayama E, et al. Widespread discrepancy in Nnt
genotypes and genetic backgrounds complicates granzyme A and other knockout mouse studies. eLife.
2022;11:e70207.
26.   Dumenil T, Le TT, Rawle DJ, Yan K, Tang B, Nguyen W, et al. Warmer ambient air temperatures
reduce nasal turbinate and brain infection, but increase lung inflammation in the K18-hACE2 mouse
model of COVID-19. Science of The Total Environment. 2023;859:160163.
27.   Prow NA, Tang B, Gardner J, Le TT, Taylor A, Poo YS, et al. Lower temperatures reduce type I
interferon activity and promote alphaviral arthritis. PLoS Pathog. 2017;13(12):e1006788.
28.   Prow NA, Liu L, Nakayama E, Cooper TH, Yan K, Eldi P, et al. A vaccinia-based single vector
construct multi-pathogen vaccine protects against both Zika and chikungunya viruses. Nat Commun.
2018;9(1):1230.
29.   Hobson-Peters J, Harrison JJ, Watterson D, Hazlewood JE, Vet LJ, Newton ND, et al. A
recombinant platform for flavivirus vaccines and diagnostics using chimeras of a new insect-specific
virus. Science Translational Medicine. 2019;11(522):eaax7888.
30.   Yan K, Vet LJ, Tang B, Hobson-Peters J, Rawle DJ, Le TT, et al. A Yellow Fever Virus 17D Infection
and Disease Mouse Model Used to Evaluate a Chimeric Binjari-Yellow Fever Virus Vaccine. Vaccines
[Internet]. 2020; 8(3).
31.   Oikari LE, Pandit R, Stewart R, Cuní-López C, Quek H, Sutharsan R, et al. Altered Brain Endothelial
Cell Phenotype from a Familial Alzheimer Mutation and Its Potential Implications for Amyloid Clearance
and Drug Delivery. Stem Cell Reports. 2020;14(5):924-39.
32.   Stewart R, Ellis S, Yan K, Dumenil T, Tang B, Nguyen W, et al. Increased neurovirulence of
omicron BA.5 over BA.1 in human brain organoids and K18-hACE2 mice. PREPRINT (Version 1) Research
Square https://doi.org/10.21203/rs.3.rs-2702556/v1. 2023.
33.   Guirakhoo F, Zhang ZX, Chambers TJ, Delagrave S, Arroyo J, Barrett AD, et al. Immunogenicity,
genetic stability, and protective efficacy of a recombinant, chimeric yellow fever-Japanese encephalitis
virus (ChimeriVax-JE) as a live, attenuated vaccine candidate against Japanese encephalitis. Virology.
1999;257(2):363-72.
34.   Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. Nature
Methods. 2012;9(7):671-5.
35.   Nasveld PE, Ebringer A, Elmes N, Bennett S, Yoksan S, Aaskov J, et al. Long term immunity to live
attenuated Japanese encephalitis chimeric virus vaccine: randomized, double-blind, 5-year phase II
study in healthy adults. Hum Vaccin. 2010;6(12):1038-46.
36.   Furuya-Kanamori L, Gyawali N, Mills DJ, Mills C, Hugo LE, Devine GJ, et al. Immunogenicity of a
single fractional intradermal dose of Japanese encephalitis live attenuated chimeric vaccine. Journal of
Travel Medicine. 2023;30(2):taac122.
37.   Wang X, Li S-H, Zhu L, Nian Q-G, Yuan S, Gao Q, et al. Near-atomic structure of Japanese
encephalitis virus reveals critical determinants of virulence and stability. Nature Communications.
2017;8(1):14.
38.   Miyata T, Miyazawa S, Yasunaga T. Two types of amino acid substitutions in protein evolution.
Journal of Molecular Evolution. 1979;12(3):219-36.
39.   Williams DT, Wang LF, Daniels PW, Mackenzie JS. Molecular characterization of the first
Australian isolate of Japanese encephalitis virus, the FU strain. J Gen Virol. 2000;81(Pt 10):2471-80.

40.   Poidinger M, Hall RA, Mackenzie JS. Molecular Characterization of the Japanese Encephalitis
Serocomplex of the Flavivirus Genus. Virology. 1996;218(2):417-21.
41.   Williams DT, Diviney SM, Niazi A-u-R, Durr PA, Chua BH, Herring B, et al. The Molecular
Epidemiology and Evolution of Murray Valley Encephalitis Virus: Recent Emergence of Distinct Sub-
lineages of the Dominant Genotype 1. PLOS Neglected Tropical Diseases. 2015;9(11):e0004240.
42.   Aoki K, Shimada S, Simantini DS, Tun MMN, Buerano CC, Morita K, et al. Type-I interferon
response affects an inoculation dose-independent mortality in mice following Japanese encephalitis
virus infection. Virology Journal. 2014;11(1):105.
43.   Rastogi M, Sharma N, Singh SK. Flavivirus NS1: a multifaceted enigmatic viral protein. Virology
Journal. 2016;13(1):131.
44.   Iwasaki Y, Zhao JX, Yamamoto T, Konno H. Immunohistochemical demonstration of viral
antigens in Japanese encephalitis. Acta Neuropathol. 1986;70(1):79-81.
45.   German AC, Myint KS, Mai NT, Pomeroy I, Phu NH, Tzartos J, et al. A preliminary
neuropathological study of Japanese encephalitis in humans and a mouse model. Trans R Soc Trop Med
Hyg. 2006;100(12):1135-45.
46.   Siva Venkatesh IP, Bhaskar M, Basu A. Japanese encephalitis viral infection modulates
proinflammatory cyto/chemokine profile in primary astrocyte and cell line of astrocytic origin. Metabolic
Brain Disease. 2022;37(5):1487-502.
47.   Li Y, Zhang H, Zhu B, Ashraf U, Chen Z, Xu Q, et al. Microarray Analysis Identifies the Potential
Role of Long Non-Coding RNA in Regulating Neuroinflammation during Japanese Encephalitis Virus
Infection. Frontiers in Immunology. 2017;8.
48.   Wilhelmsson U, Bushong EA, Price DL, Smarr BL, Phung V, Terada M, et al. Redefining the
concept of reactive astrocytes as cells that remain within their unique domains upon reaction to injury.
Proceedings of the National Academy of Sciences. 2006;103(46):17513-8.
49.   Zhang Z, Ma Z, Zou W, Guo H, Liu M, Ma Y, et al. The Appropriate Marker for Astrocytes:
Comparing the Distribution and Expression of Three Astrocytic Markers in Different Mouse Cerebral
Regions. BioMed Research International. 2019;2019:9605265.
50.   Daffis S, Samuel MA, Suthar MS, Keller BC, Gale M, Jr., Diamond MS. Interferon regulatory factor
IRF-7 induces the antiviral alpha interferon response and protects against lethal West Nile virus
infection. J Virol. 2008;82(17):8465-75.
51.   Hazlewood JE, Tang B, Yan K, Rawle DJ, Harrison JJ, Hall RA, et al. The Chimeric Binjari-Zika
Vaccine Provides Long-Term Protection against ZIKA Virus Challenge. Vaccines (Basel). 2022;10(1).
52.   Roberts A, Kesarwani V, Gupta R, Gandhi S. Electroactive reduced graphene oxide for highly
sensitive detection of secretory non-structural 1 protein: A potential diagnostic biomarker for Japanese
encephalitis virus. Biosens Bioelectron. 2022;198:113837.
53.   Puerta-Guardo H, Glasner DR, Espinosa DA, Biering SB, Patana M, Ratnasiri K, et al. Flavivirus
NS1 Triggers Tissue-Specific Vascular Endothelial Dysfunction Reflecting Disease Tropism. Cell Reports.
2019;26(6):1598-613.e8.
54.   Baccala R, Welch MJ, Gonzalez-Quintial R, Walsh KB, Teijaro JR, Nguyen A, et al. Type I
interferon is a therapeutic target for virus-induced lethal vascular damage. Proceedings of the National
Academy of Sciences. 2014;111(24):8925-30.
55.   Iannacone M, Sitia G, Isogawa M, Whitmire JK, Marchese P, Chisari FV, et al. Platelets prevent
IFN-α/β-induced lethal hemorrhage promoting CTL-dependent clearance of lymphocytic
choriomeningitis virus. Proceedings of the National Academy of Sciences. 2008;105(2):629-34.
56.   Miyake M. THE PATHOLOGY OF JAPANESE ENCEPHALITIS. A REVIEW. Bull World Health Organ.
1964;30(2):153-60.
57.   Waller C, Tiemensma M, Currie BJ, Williams DT, Baird RW, Krause VL. Japanese Encephalitis in
Australia — A Sentinel Case. New England Journal of Medicine. 2022;387(7):661-2.

58.   Maamary J, Maddocks S, Barnett Y, Wong S, Rodriguez M, Hueston L, et al. New Detection of
Locally Acquired Japanese Encephalitis Virus Using Clinical Metagenomics, New South Wales, Australia.
Emerging Infectious Disease journal. 2023;29(3):627.
59.   Bishop CR, Dumenil T, Rawle DJ, Le TT, Yan K, Tang B, et al. Mouse models of COVID-19
recapitulate inflammatory pathways rather than gene expression. PLoS Pathog. 2022;18(9):e1010867.
60.   Tripathi A, Singh Rawat B, Addya S, Surjit M, Tailor P, Vrati S, et al. Lack of Interferon (IFN)
Regulatory Factor 8 Associated with Restricted IFN-gamma Response Augmented Japanese Encephalitis
Virus Replication in the Mouse Brain. J Virol. 2021;95(21):e0040621.
61.   Danaher P, Kim Y, Nelson B, Griswold M, Yang Z, Piazza E, et al. Advances in mixed cell
deconvolution enable quantification of cell types in spatial transcriptomic data. Nat Commun.
2022;13(1):385.
62.   Setoh YX, Amarilla AA, Peng NYG, Griffiths RE, Carrera J, Freney ME, et al. Determinants of Zika
virus host tropism uncovered by deep mutational scanning. Nat Microbiol. 2019;4(5):876-87.
63.   Zhang B, He Y, Xu Y, Mo F, Mi T, Shen QS, et al. Differential antiviral immunity to Japanese
encephalitis virus in developing cortical organoids. Cell Death & Disease. 2018;9(7):719.
64.   Chambers TJ, Nestorowicz A, Mason PW, Rice CM. Yellow fever/Japanese encephalitis chimeric
viruses: construction and biological properties. J Virol. 1999;73(4):3095-101.
65.   Frassetto FP, Rosemberg S. Neuropathology of yellow fever autopsy cases. Tropical Diseases,
Travel Medicine and Vaccines. 2023;9(1):1.
66.   Cohen M, Nguyen M, Nix CD, Case B, Nickerson JP, Bernard J, et al. Case Report: Yellow Fever
Vaccine-Associated Neurotropic Disease and Associated MRI, EEG, and CSF Findings. Front Neurol.
2021;12:779014.
67.   Pierson TC, Fremont DH, Kuhn RJ, Diamond MS. Structural insights into the mechanisms of
antibody-mediated neutralization of flavivirus infection: implications for vaccine development. Cell Host
Microbe. 2008;4(3):229-38.
68.   Hombach J, Solomon T, Kurane I, Jacobson J, Wood D. Report on a WHO consultation on
immunological endpoints for evaluation of new Japanese encephalitis vaccines, WHO, Geneva, 2–3
September, 2004. Vaccine. 2005;23(45):5205-11.
69.   Myint KS, Kipar A, Jarman RG, Gibbons RV, Perng GC, Flanagan B, et al. Neuropathogenesis of
Japanese encephalitis in a primate model. PLoS Negl Trop Dis. 2014;8(8):e2980.
70.   Japanese encephalitis virus (JEV). Australian Governement Dept of Health and Aged Care.
https://www.health.gov.au/health-alerts/japanese-encephalitis-virus-jev/japanese-encephalitis-virus-
jev (accessed April 2023).
71.   Serourvey for JEV in northern Victoria. https://www.health.vic.gov.au/infectious-
diseases/serosurvey-for-japanese-encephalitis-in-northern-victoria (accessed April 2023).
72.   Summary of NSW JEV seosurvey results.
https://www.health.nsw.gov.au/environment/pests/vector/Documents/jev-serosurvey-report.pdf
(accessed April 2023).
73.   Hsieh JT, St John AL. Japanese encephalitis virus and its mechanisms of neuroinvasion. PLoS
Pathog. 2020;16(4):e1008260.
74.   Setoh YX, Periasamy P, Peng NYG, Amarilla AA, Slonchak A, Khromykh AA. Helicase Domain of
West Nile Virus NS3 Protein Plays a Role in Inhibition of Type I Interferon Signalling. Viruses.
2017;9(11):326.
75.   Laurent-Rolle M, Boer EF, Lubick KJ, Wolfinbarger JB, Carmody AB, Rockx B, et al. The NS5
protein of the virulent West Nile virus NY99 strain is a potent antagonist of type I interferon-mediated
JAK-STAT signaling. J Virol. 2010;84(7):3503-15.

76.   Nakayama E, Kato F, Tajima S, Ogawa S, Yan K, Takahashi K, et al. Neuroinvasiveness of the
MR766 strain of Zika virus in IFNAR-/- mice maps to prM residues conserved amongst African genotype
viruses. PLoS Pathog. 2021;17(7):e1009788.
77.   Zheng W, Yan Q, Li Z, Wang X, Wu P, Liao F, et al. Liver transcriptomics reveals features of the
host response in a mouse model of dengue virus infection. Front Immunol. 2022;13:892469.
78.   Grant A, Ponia SS, Tripathi S, Balasubramaniam V, Miorin L, Sourisseau M, et al. Zika Virus
Targets Human STAT2 to Inhibit Type I Interferon Signaling. Cell Host Microbe. 2016;19(6):882-90.
79.   Yang H, Fan F, Liu L, Liu J, Sun Y, Xie A, et al. A novel amino acid site closely associated with the
neurovirulence of live, attenuated Japanese encephalitis vaccine (SA14-14-2 strain). Vaccine.
2020;38(11):2636-42.
80.   Yun SI, Song BH, Kim JK, Yun GN, Lee EY, Li L, et al. A molecularly cloned, live-attenuated
japanese encephalitis vaccine SA14-14-2 virus: a conserved single amino acid in the ij Hairpin of the Viral
E glycoprotein determines neurovirulence in mice. PLoS Pathog. 2014;10(7):e1004290.
81.   Arroyo J, Guirakhoo F, Fenner S, Zhang ZX, Monath TP, Chambers TJ. Molecular basis for
attenuation of neurovirulence of a yellow fever Virus/Japanese encephalitis virus chimera vaccine
(ChimeriVax-JE). J Virol. 2001;75(2):934-42.
82.   Yang D, Li X-F, Ye Q, Wang H-J, Deng Y-Q, Zhu S-Y, et al. Characterization of live-attenuated
Japanese encephalitis vaccine virus SA14-14-2. Vaccine. 2014;32(23):2675-81.
83.   Liu X, Zhao X, Na R, Li L, Warkentin E, Witt J, et al. The structure differences of Japanese
encephalitis virus SA14 and SA14-14-2 E proteins elucidate the virulence attenuation mechanism.
Protein & Cell. 2019;10(2):149-53.
84.   Yang J, Yang H, Li Z, Wang W, Lin H, Liu L, et al. Envelope Protein Mutations L107F and E138K Are
Important for Neurovirulence Attenuation for Japanese Encephalitis Virus SA14-14-2 Strain. Viruses
[Internet]. 2017; 9(1).
85.   Ni H, Chang GJ, Xie H, Trent DW, Barrett AD. Molecular basis of attenuation of neurovirulence of
wild-type Japanese encephalitis virus strain SA14. J Gen Virol. 1995;76 (Pt 2):409-13.
86.   Zhou Y, Wu R, Zhao Q, Chang YF, Wen X, Feng Y, et al. Mutation of I176R in the E coding region
weakens Japanese encephalitis virus neurovirulence, but not its growth rate in BHK-21 cells. Arch Virol.
2018;163(5):1351-5.
87.   Anwar MN, Wang X, Hameed M, Wahaab A, Li C, Sharma M, et al. Phenotypic and Genotypic
Comparison of a Live-Attenuated Genotype I Japanese Encephalitis Virus SD12-F120 Strain with Its
Virulent Parental SD12 Strain. Viruses [Internet]. 2020; 12(5).
88.   Gromowski GD, Firestone CY, Whitehead SS. Genetic Determinants of Japanese Encephalitis
Virus Vaccine Strain SA14-14-2 That Govern Attenuation of Virulence in Mice. J Virol. 2015;89(12):6328-
37.
89.   Monath TP, Arroyo J, Levenbook I, Zhang ZX, Catalan J, Draper K, et al. Single mutation in the
flavivirus envelope protein hinge region increases neurovirulence for mice and monkeys but decreases
viscerotropism for monkeys: relevance to development and safety testing of live, attenuated vaccines. J
Virol. 2002;76(4):1932-43.
90.   Chen LK, Lin YL, Liao CL, Lin CG, Huang YL, Yeh CT, et al. Generation and characterization of
organ-tropism mutants of Japanese encephalitis virus in vivo and in vitro. Virology. 1996;223(1):79-88.
91.   Sumiyoshi H, Tignor GH, Shope RE. Characterization of a highly attenuated Japanese encephalitis
virus generated from molecularly cloned cDNA. J Infect Dis. 1995;171(5):1144-51.
92.   Zheng X, Zheng H, Tong W, Li G, Wang T, Li L, et al. Acidity/Alkalinity of Japanese Encephalitis
Virus E Protein Residue 138 Alters Neurovirulence in Mice. J Virol. 2018;92(22).
93.   Chiou SS, Chen WJ. Phenotypic changes in the Japanese encephalitis virus after one passage in
Neuro-2a cells: generation of attenuated strains of the virus. Vaccine. 2007;26(1):15-23.

994. Zhao Z, Date T, Li Y, Kato T, Miyamoto M, Yasui K, et al. Characterization of the E-138 (Glu/Lys) mutation in Japanese encephalitis virus by using a stable, full-length, infectious cDNA clone. J Gen Virol. 2005;86(Pt 8):2209-20.

95. Xia Q, Zhang Y, Yang Y, Ma X, Guan Z, Zhang J, et al. Virulence and Cross-Protection Conferred by an Attenuated Genotype I-Based Chimeric Japanese Encephalitis Virus Strain Harboring the E Protein of Genotype V in Mice. Microbiology Spectrum. 2022;10(6):e01990-22.

96. Wang Q, Xin X, Wang T, Wan J, Ou Y, Yang Z, et al. Japanese Encephalitis Virus Induces Apoptosis and Encephalitis by Activating the PERK Pathway. J Virol. 2019;93(17).

97. Tajima S, Shibasaki KI, Taniguchi S, Nakayama E, Maeki T, Lim CK, et al. E and prM proteins of genotype V Japanese encephalitis virus are required for its increased virulence in mice. Heliyon. 2019;5(11):e02882.

98. Whiteman MC, Wicker JA, Kinney RM, Huang CY, Solomon T, Barrett AD. Multiple amino acid changes at the first glycosylation motif in NS1 protein of West Nile virus are necessary for complete attenuation for mouse neuroinvasiveness. Vaccine. 2011;29(52):9702-10.

99. Whiteman MC, Li L, Wicker JA, Kinney RM, Huang C, Beasley DW, et al. Development and characterization of non-glycosylated E and NS1 mutant viruses as a potential candidate vaccine for West Nile virus. Vaccine. 2010;28(4):1075-83.

100. Burton DR. Antiviral neutralizing antibodies: from in vitro to in vivo activity. Nature Reviews Immunology. 2023.

101. Li F, Feng Y, Wang G, Zhang W, Fu S, Wang Z, et al. Tracing the spatiotemporal phylodynamics of Japanese encephalitis virus genotype I throughout Asia and the western Pacific. PLOS Neglected Tropical Diseases. 2023;17(4):e0011192.

102. Broom AK, Wallace MJ, Mackenzie JS, Smith DW, Hall RA. Immunisation with gamma globulin to murray valley encephalitis virus and with an inactivated Japanese encephalitis virus vaccine as prophylaxis against australian encephalitis: evaluation in a mouse model. J Med Virol. 2000;61(2):259-65.

103. Bielefeldt-Ohmann H, Prow NA, Wang W, Tan CSE, Coyle M, Douma A, et al. Safety and immunogenicity of a delta inulin-adjuvanted inactivated Japanese encephalitis virus vaccine in pregnant mares and foals. Veterinary Research. 2014;45(1):130.

104. Rawle DJ, Nguyen W, Dumenil T, Parry R, Warrilow D, Tang B, et al. Sequencing of Historical Isolates, K-mer Mining and High Serological Cross-Reactivity with Ross River Virus Argue against the Presence of Getah Virus in Australia. Pathogens. 2020;9(10).

105. Lobigs M, Pavy M, Hall R. Cross-protective and infection-enhancing immunity in mice vaccinated against flaviviruses belonging to the Japanese encephalitis virus serocomplex. Vaccine. 2003;21(15):1572-9.

106. Williams DT, Daniels PW, Lunt RA, Wang LF, Newberry KM, Mackenzie JS. Experimental infections of pigs with Japanese encephalitis virus and closely related Australian flaviviruses. Am J Trop Med Hyg. 2001;65(4):379-87.

107. Lobigs M, Larena M, Alsharifi M, Lee E, Pavy M. Live chimeric and inactivated Japanese encephalitis virus vaccines differ in their cross-protective values against Murray Valley encephalitis virus. J Virol. 2009;83(6):2436-45.

FIGURE LEGENDS

Figure 1. JEV and MVEV infection in C57BL/6J mice. (A-D) Female ≈6 week old C57BL/6J mice were infected s.c. with 5×10^5 (black lines) or 5×10^3 (red lines) $CCID_{50}$ of the indicated viruses. Mean viremia for n=6 per group over 5 days is shown. Error bars represent standard error. All mice recorded a detectable viremia for at least one timepoint. Limit of detection is 2 $log_{10}CCID_{50}$/ml of serum. Statistics represent t-test or Kolmogorov-Smirnov test at the indicated timepoint or for area under the curve (AUC) values (see methods section). (E) Percent body weight change of individual mice after infection with the indicated virus at 5×10^5 $CCID_{50}$ compared to each mouse's weight on day zero. Four mice lost ≥20% body weight and were euthanized (†). (F) Kaplan-Meier plot showing percent survival (n=6 for each virus/virus isolate inoculated at 5×10^5 $CCID_{50}$). (G) Viral tissue titers in brains and spleens of seven euthanized mice at the time when the criteria for humane euthanasia was met (see Supplementary Figure 2) (n=4 $JEV_{Nakayama}$ – black circles, n=2 JEV_{FU} – blue square, n=1 $MVEV_{TC123130}$ green downward triangle). Tissue titers determined by $CCID_{50}$ assay (limit of detection ~2 $log_{10}CCID_{50}$/g). (H) Percent body weight change of individual mice after infection with the indicated virus at 5×10^3 $CCID_{50}$ compared to each mouse's weight on day zero. Six mice lost ≥20% body weight and were euthanized (†). (I) Kaplan-Meier plot showing percent survival (n=6 for each virus/virus isolate inoculated at 5×10^3 $CCID_{50}$). (J) Viremia comparing $JEV_{Nakayama}$ (black circles), JEV_{FU} (blue squares), $JEV_{NSW/22}$ (red triangles) and $MVEV_{TC123130}$ (green downward triangles) at 5×10^3 inoculation dose; data is a reanalysis of data presented in Figure 1A-D. Data is mean of n=6 per group and error bars represent standard error. Statistics are t-test for $JEV_{Nakayama}$ versus $JEV_{NSW/22}$ on day 2 and JEV_{FU} versus $JEV_{NSW/22}$ on day 3.

Figure 2. IHC for viral antigen, apoptosis and reactive astrocytes in JEV infected C57BL/6J mouse brain. IHC of $JEV_{Nakayama}$ infected brain (required euthanasia on 9 dpi, brain virus titer 9.2 $log_{10}CCID_{50}$/g), which is representative of all other C57BL/6J brains from infected mice requiring euthanasia (n=10). (A) Staining for flavivirus NS1 using the 4G4 monoclonal antibody. High magnification images show cells with neuronal morphology in the cortex (left) and hippocampus (right). The latter also shows staining of dendrites and axons (fibrilar patterns above and densely staining cells). 4G4 staining of brains of the other mice requiring euthanasia (marked by † in Figure 1E) is shown in Supplementary Figure 3. (B) ApopTag staining of the

same brain as in 'A'. High magnification images show cells with neuronal morphology in the

cortex (left) and apoptotic cells in the dentate gyrus (right). ApopTag staining of the brains of

the other mice requiring euthanasia is shown in Supplementary Figure 4. (C) Staining for GFAP,

a marker of reactive astrocytes, for the same brain as in 'A'. High magnification images show a

typical reactive astrocyte in the cortex (left) and reactive astrocytes in the thalamus (right).

GFAP IHC for the brains of the other mice requiring euthanasia is shown in Supplementary

Figure 5. (D) IHC negative controls; uninfected mouse brain stained with 4G4 (left), ApopTag

(middle), and anti-GFAP (right).

1080

Figure 3. JEV neuroinvasive infection in *Irf7$^{-/-}$* mice. (A) *Irf7$^{-/-}$* mice (15-48 weeks old) were

infected s.c. with 5×10^5, 5×10^4, 5×10^3 or 5×10^2 CCID$_{50}$ of JEV$_{NSW/22}$. Data is the mean of n=6

per group and error bars represent standard error. Statistics are a t-test of area under the curve

for the indicated comparisons. (B) *Irf7$^{-/-}$* mice (15-48 weeks old) were infected s.c. with 5×10^3 of

the indicated virus (n=8 for each virus). Statistics are a t-test of area under the curve for JEV$_{FU}$

versus JEV$_{NSW/22}$ or MVEV$_{TC123130}$. (C) Percent body weight change compared to 0 dpi. 20

mice lost >20% body weight or reached a disease score that required euthanized (marked by †)

and are plotted individually. The weight change for the remaining mice are shown as means $\pm$

SE. Data is from three independent experiments, total n=14 mice per group. (D) Kaplan-Meier

plot showing percent survival (n=14 per group, data from three independent experiments).

Statistics by log-rank (Mantel-Cox) tests. Symbols as for C. (E) Viremia of *Irf7$^{-/-}$* mice

averaged for mice with non-lethal outcomes (black circles, n=19), versus those with lethal

outcomes (brown squares, n=13). Statistics are comparing average viremia of mice with lethal

outcomes versus mice with non-lethal outcomes at each timepoint by t-test or Kolmogorov-

Smirnov test. T-test of area under the curve values for this comparison is also shown. (F) Brain

and spleen tissue titers for 13 euthanized mice at the time when the criteria for humane

euthanasia was met (see Supplementary Figure 6) (n=5 JEV$_{Nakayama}$, n=6 JEV$_{FU}$, n=1 JEV$_{NSW/22}$,

n=1 MVEV$_{TC123130}$). Tissue titers determined by CCID$_{50}$ assay (limit of detection ~2

log$_{10}$CCID$_{50}$/g). (G) IHC of JEV$_{NSW/22}$ infected brain (euthanized on day 8, brain virus titer 6.3

log$_{10}$CCID$_{50}$/g) using 4G4 monoclonal antibody (top) or ApopTag (bottom). High magnification

images of cortex (left) and medulla (right). (H) IHC using 4G4 (left) and ApopTag IHC (right)

for the JEV$_{Nakayama}$ infected brain with the highest virus titer (euthanized on day 7, brain virus

titer 8.3 $\log_{10}CCID_{50}$/g). IHC images are representative of n=20 brains of mice that succumbed

to infection (some other examples shown in Supplementary Figure 8).

1105

Figure 4. Comparisons of viremia and survival between C57BL/6J versus *Irf7*$^{-/-}$ mice, and

comparison of virus replication in wild-type versus *Irf3/7*$^{-/-}$ mouse embryonic fibroblasts

(MEFs). C57BL/6J or *Irf7*$^{-/-}$ mice were infected with $5x10^3$ $CCID_{50}$ of JEV$_{Nakayama}$, JEV$_{FU}$,

JEV$_{NSW/22}$, or MVEV$_{TC123130}$. Data shown in panels A-H are a re-analysis of data presented in

Figure 1A-D, Figure 1I, Figure 3B, and Figure 3D. (A-D) Female ≈6 week old C57BL/6J mice

(black lines) or *Irf7*$^{-/-}$ mice (red lines) were infected s.c. with $5x10^3$ $CCID_{50}$ of the indicated

viruses. Mean viremia for n=6 C57BL/6J mice and n=8 Irf7$^{-/-}$ mice per group over 5 days is

shown. Error bars represent standard error. Limit of detection is 2 $\log_{10}CCID_{50}$/ml of serum.

Statistics represent t-test or Kolmogorov-Smirnov test at the indicated timepoint or for area

under the curve (AUC) values (see statistics methods section). (E-H) Kaplan-Meier plot

showing percent survival for C57BL/6J mice (black line, n=6) and Irf7$^{-/-}$ mice (ref line, n=14)

infected with $5x10^3$ $CCID_{50}$ of the indicated virus. Statistics by log-rank (Mantel-Cox) tests. (I)

Wild type or (J) *Irf3/7*$^{-/-}$ MEFs were infected with JEV$_{Nakayama}$ (black circles), JEV$_{FU}$ (blue

squares), JEV$_{NSW/22}$ (red triangles), or MVEV$_{TC123130}$ (green downward triangles) at MOI 0.1.

Virus titer in the culture supernatant was monitored over 4 days. Data is the mean two

independent experiments with a total of n=6 replicates per group. Error bars represent standard

error. Statistics are by t-test or Kolmogorov-Smirnov test for the indicated comparisons.

1123

Figure 5. JEV and MVEV lethal viremia in *Ifnar*$^{-/-}$ mice. Female *Ifnar*$^{-/-}$ mice (9-24 week

old) were infected s.c. with $5x10^5$ $CCID_{50}$ of the indicated virus (n=4 for JEV$_{Nakayama}$, JEV$_{NSW/22}$

and MVEV$_{TC123130}$, and n=5 for JEV$_{FU}$). (A) Mean viremia determined by $CCID_{50}$ assay (limit

of detection 2 $\log_{10}CCID_{50}$/ml). Statistics by t-tests. (B) Mean percent body weight change

compared to 0 dpi. Statistics 2 dpi JEV$_{NSW/22}$ versus JEV$_{Nakayama}$ (Kolmogorov-Smirnov test,

p=0.023) and for JEV$_{NSW/22}$ versus JEV$_{FU}$ (t-test, p=0.008). (C) Kaplan-Meier plot showing

percent survival. (D) Viral tissue titers in spleens harvested at euthanasia (3 dpi for all mice

except for 1 JEV$_{FU}$ mouse at 2 dpi), determined by $CCID_{50}$ assay (limit of detection ~2 $\log_{10}$

$CCID_{50}$/g). (E) Viral titers in brains harvested at euthanasia. (F) IHC staining for flavivirus NS1

using the 4G4 monoclonal antibody. The brain shown was infected with JEV$_{Nakayama}$ (titer 7.2

log10 CCID50/g, 3 dpi). Staining was representative of all other JEV brains. (G, H) High

magnification images from F showing NS1 staining in blood vessels surrounding the pale grey,

biconcave shaped, red blood cells.

Figure 6. Histological lesions in brains of JEV infected mice. (A) Examples of indicated

lesions stained with H&E. Degeneration of neurons indicated by (i) pyknotic nuclei (black

unfilled arrow heads indicating condensation and fragmentation of nuclei, staining dark blue),

and (ii) hyper-eosinophilic cytoplasm of degenerating neurons in the cornus ammonis of the

hippocampus (area also staining for viral antigen Supplementary Figure 10). Neuronal

vacuolation indicated by fluid accumulation around the neurons (arrowheads). Perivascular

cuffing is indicated by leukocytes aggregating in blood vessels (arrowheads). Leukocyte

infiltrates (extravascular) are indicated by dashed ovals. Hemorrhagic lesions are indicated by

extravascular red blood cells (arrows). Microgliosis is indicated by accumulation of microglia,

which have an elongated rod-shaped nuclei (arrows). Meningitis is indicated by accumulation of

leukocytes around the meninges (arrowheads). Images of uninfected controls accompany each

image(s) of lesions. Histology scores for all mouse brains are shown in Supplementary Figure 9

and 10. (B) IHC using anti-Iba1, a microglial marker.

Figure 7. Infection of human cortical brain organoids (hBOs). (A) Photograph of "mini-

brains" cultured in a rotating CelVivo Clinostar incubator. (B) IHC of viral antigen (4G4) for

hBOs at 4 dpi. Images are representative of n=4 hBOs for each group. Magnified images of

sparse Imojev and YFV 17D infected cells are shown in Supplementary Figure 12A. (B) Viral

growth kinetics up to 4 dpi determined by $CCID_{50}$ assays of culture supernatants at the indicated

hours post infection; limit of detection is 2 $\log_{10}CCID_{50}$/ml. At all time points $JEV_{Nakayama}$ vs.

Imojev, and at 96 h $JEV_{Nakayama}$ vs. $JEV_{NSW/22}$ were significant (t tests, n=5 organoids per group).

(C) Mean percentage change in organoid area at 11 dpi vs. 9 dpi for each organoid (n=8 for

uninfected and $MVEV_{TC123130}$, otherwise n=4). Statistics are by Kolmogorov-Smirnov test for

uninfected versus $JEV_{Nakayama}$, and t-test for uninfected versus JEV_{FU} or $JEV_{NSW/22}$.

Figure 8. Human post Imojev-vaccination sera neutralizes JEV and MVEV with titers

related to envelope protein amino acid conservation. (A) Envelope protein (domains I, II, III

and STEM) amino acid sequences for Imojev, $JEV_{Nakayama}$, JEV_{FU}, $JEV_{NSW/22}$, and $MVEV_{TC123130}$

(refer to Supplementary Figure 1 for GenBank accession numbers). Sequences for isolates were

aligned using MEGA-X and the ClustalW plugin with default parameters. Colouring indicates

amino acid category compared to Imojev (orange = unique to Imojev, purple = identical, blue =

conservative amino acid difference, pink = non-conservative amino acid difference (38)).

Crystal structure of JEV envelope (PDB: 5WSN) with amino acid differences for $JEV_{Nakayama}$,

JEV_{FU}, $JEV_{NSW/22}$, and $MVEV_{TC123130}$ compared to Imojev coloured as described in the table.

Percentages indicate percent sequence identity relative to Imojev. (B) Human serum taken at day

0 and day 28 post-Imojev vaccination (n=9, cohort 1) was used in plaque reduction neutralisation

assays against $JEV_{Nakayama}$, JEV_{FU} and $JEV_{NSW/22}$, and the plaque reduction neutralisation 50%

titer ($PRNT_{50}$) was calculated. Mean and standard error are shown. Statistics are paired t-test

comparing Imojev with $JEV_{Nakayama}$ or $JEV_{NSW/22}$. (B) Human serum taken 2-12 months post-

Imojev vaccination (n=10, cohort 2) was used in plaque reduction neutralisation assays against

$JEV_{Nakayama}$, JEV_{FU}, $JEV_{NSW/22}$, and $MVEV_{TC123130}$ and the plaque reduction neutralisation 50%

titer ($PRNT_{50}$) was calculated. Mean and standard error are shown. Statistics are paired t-test

comparing $JEV_{Nakayama}$ with $JEV_{NSW/22}$. (D) $PRNT_{50}$ titers in 'B' and 'C' plotted against

percentage envelope protein amino acid identity in 'A'. Curve fit is shown, and statistics

calculated by Spearman correlation with p and rho values shown (black line represents all data,

brown line excludes $MVEV_{TC123130}$ from analysis).

Figure 1

Figure 2

A Anti-flavivirus NS1 (virus infected cells)

JEV$_{Nakayama}$ **brain titer 9.2 log$_{10}$CCID$_{50}$/g**
C57BL/6J mouse 9 dpi

Cortex

Hippocampus

Anterior olfactory nucleus

Thalamus

100 µm

2 mm

50 µm

B ApopTag (apoptosis)

Cortex

Dentate gyrus

Anterior olfactory nucleus

10 µm

2 mm

50 µm

C Anti -GFAP (reactive astrocytes)

Cortex

Thalamus

10 µm

2 mm

50 µm

D Uninfected C57BL/6J mouse brain

Anti-flavivirus NS1

ApopTag

Anti-GFAP

Cortex

Cortex

Cortex

2 mm

2 mm

2 mm

100 µm

100 µm

100 µm

Figure 3

Figure 4

Figure 5

Figure 6

A

Neurons with pyknotic nuclei
Nakayama C57BL/6J 11 dpi

Hyper-eosinophilic neurons
Nakayama C57BL/6J 9 dpi

Neuronal vacuolation
Nakayama *Irf7*⁻/⁻ 7 dpi

Uninfected control

Perivascular cuffing
NSW *Irf7*⁻/⁻ 8 dpi

Uninfected control

Leukocyte infiltrates
Nakayama *Irf7*⁻/⁻ 7 dpi

Uninfected control

Hemorrhagic lesions
Nakayama C57BL/6J 9 dpi

Uninfected control

Meningitis
Nakayama *Irf7*⁻/⁻ 7 dpi

Uninfected control

Microgliosis
Nakayama *Irf7*⁻/⁻ 19 dpi (survived)

Uninfected control

B Microgliosis (Iba1 IHC)
Nakayama C57BL/6J 9 dpi

Uninfected control

Figure 7

Figure 8

A

		Envelope		
	Domain I	Domain II	Domain III	Stem

ORANGE = unique to Imojev PURPLE = identical BLUE = conservative PINK = non-conservative

Imojev (100%) Nakayama (96.8%) FU (96.4%) NSW/22 (93.4%) MVEV (80.4%)

☐ Domain I ▨ Domain II ▨ Domain III ■ Stem

B Cohort 1

PRNT$_{50}$ titer, replicates and mean $\pm$ SE

p=0.049
p=0.012

Imojev JEV$_{Nakayama}$ JEV$_{FU}$ JEV$_{NSW/22}$ Baseline

C Cohort 2

PRNT$_{50}$ titer, replicates and mean $\pm$ SE

p=0.042

Imojev JEV$_{Nakayama}$ JEV$_{FU}$ JEV$_{NSW/22}$ MVEV$_{TC123130}$

D

PRNT$_{50}$ titer

● JEV$_{Nakayama}$
■ JEV$_{FU}$
▲ JEV$_{NSW/22}$
▼ MVEV$_{TC123130}$
◆ Imojev

p<0.0001
rho = 0.453

p<0.0001
rho = 0.447

% identity of envelope amino acid sequence compared to Imojev

A

Virus name	GenBank accession	JEV Genotype	Isolation year	Isolation source	Passage history
JEV$_{Nakayama}$	EF571853	3	1935	Human cerebrospinal fluid	>11 x suckling mouse brain
JEV$_{FU}$	AF217620	2	1995	Human serum	C6/36 (unknown number)
JEV$_{NSW2022}$	OP904182	4	2022	Stillborn piglet brain	1 x C6/36 1 x BHK-21
MVEV$_{TC123130}$	JN119814	N/A	1974	Human brain	Unknown
YFV 17D	MT107250	N/A	1927 (Original "Asibi" parent isolate)	Human serum (Original "Asibi" parent isolate)	53 x monkeys 18 x embryonic mouse tissue 50 x minced whole chicken embryo 152 x nervous tissue deprived chick embryo
JEV$_{prME}$ component in Imojev (SA$_{14}$-14-2 strain)	MK585066	3	1954 (Original SA-14 parent isolate)	*Culex pipiens* mosquito larvae (Original SA-14 parent isolate)	100 x primary hamster kidney cells 22 x suckling mice

B   Whole genome (nucleotide sequence)   prME (amino acid sequence)

Supplementary Figure 1. (A) Summary of virus isolates used in the study. (B) Phylogenetic tree for whole genome nucleotide sequence (left) and prME amino acid sequence (right). Phylogenetic trees were constructed after nucleotide or amino acid sequence alignment using MEGA-X (Molecular Evolutionary Genetics Analysis 10, Penn State University, State College, PA, USA) and the ClustalW plugin with default parameters. The phylogenetic tree was constructed using the Maximum Likelihood method and the General Time Reversible model (nucleotide sequence) or JTT matrix-based model (amino acid sequence). Whole genome sequence is not available for MVEV$_{TC123130}$. Whole genome sequence for Imojev was constructed by combining the prME sequence from JEV$_{SA14-14-2}$ with the remainder of the genome from YFV 17D.

A

Activity

Dpi	Cg1 #2	Cg1 #6	Cg1 #2	Cg1 #3	Cg1 #6	Cg1 #4	Cg2 #1	Cg3 #3	Cg1 #3	Cg4 #1
	JEV Nakayama						*JEV FU*		*MVEV*	
0	0	0	0	0	0	0	0	0	0	0
1	0	0	0	0	0	0	0	0	0	0
2	0	0	0	0	0	0	0	0	0	0
3	0	0	0	0	0	0	0	0	0	0
4	0	0	0	0	0	0	0	0	0	0
5	0	0	0	0	0	0	0	0	0	0
6	0	0	0	0	0	0	0	0	0	0
7	0	0	0	0	0	0	0	0	0	0
8	0	0	0	0	0	0	1	0	0	0
9	1	0	1	1	0	0	▪	0	1	1
10	▪	0	2	2	0	0	▪	1	▪	2
11	▪	1	▪	▪	0	0	▪	1	▪	▪
12	▪	▪	▪	2	0	▪	▪	▪	▪	▪
13	▪	▪	▪	▪	▪	▪	▪	1	▪	▪
14	▪	▪	▪	▪	▪	▪	▪	2	▪	▪

Weight loss score

Dpi	Cg1 #2	Cg1 #6	Cg1 #2	Cg1 #3	Cg1 #6	Cg1 #4	Cg2 #1	Cg3 #3	Cg1 #3	Cg4 #1
	JEV Nakayama						*JEV FU*		*MVEV*	
0	0	0	0	0	0	0	0	0	0	0
1	0	0	0	0	0	0	0	0	0	0
2	0	0	0	0	0	0	0	0	0	0
3	0	0	0	0	0	0	0	0	0	0
4	0	0	0	0	0	0	0	0	0	0
5	0	0	0	0	0	0	0	0	0	0
6	0	0	0	0	0	0	0	1	0	0
7	0	0	0	0	0	0	2	2	0	0
8	1	0	0	0	0	0	3	3	2	0
9	3	0	1	1	0	0	▪	3	▪	1
10	▪	1	3	3	0	0	▪	3	▪	3
11	▪	3	▪	1	0	▪	▪	3	▪	▪
12	▪	▪	▪	3	0	▪	▪	3	▪	▪
13	▪	▪	▪	▪	▪	▪	▪	2	▪	▪
14	▪	▪	▪	▪	▪	▪	▪	3	▪	▪

Posture

Dpi	Cg1 #2	Cg1 #6	Cg1 #2	Cg1 #3	Cg1 #6	Cg1 #4	Cg2 #1	Cg3 #3	Cg1 #3	Cg4 #1
	JEV Nakayama						*JEV FU*		*MVEV*	
0	0	0	0	0	0	0	0	0	0	0
1	0	0	0	0	0	0	0	0	0	0
2	0	0	0	0	0	0	0	0	0	0
3	0	0	0	0	0	0	0	0	0	0
4	0	0	0	0	0	0	0	0	0	0
5	0	0	0	0	0	0	0	0	0	0
6	0	0	0	0	0	0	0	0	0	0
7	0	0	0	0	0	0	0	0	0	0
8	0	0	0	0	0	0	1	0	0	0
9	1	0	1	1	0	0	▪	0	1	1
10	▪	0	1	1	0	0	▪	1	▪	1
11	▪	1	▪	▪	0	0	▪	1	▪	▪
12	▪	▪	▪	1	0	▪	▪	▪	▪	▪
13	▪	▪	▪	▪	▪	▪	▪	1	▪	▪
14	▪	▪	▪	▪	▪	▪	▪	2	▪	▪

Fur ruffling

Dpi	Cg1 #2	Cg1 #6	Cg1 #2	Cg1 #3	Cg1 #6	Cg1 #4	Cg2 #1	Cg3 #3	Cg1 #3	Cg4 #1
	JEV Nakayama						*JEV FU*		*MVEV*	
0	0	0	0	0	0	0	0	0	0	0
1	0	0	0	0	0	0	0	0	0	0
2	0	0	0	0	0	0	0	0	0	0
3	0	0	0	0	0	0	0	0	0	0
4	0	0	0	0	0	0	0	0	0	0
5	0	0	0	0	0	0	0	0	0	0
6	0	0	0	0	0	0	0	0	0	0
7	0	0	0	0	0	0	0	0	0	0
8	0	0	0	0	0	0	0	0	0	0
9	1	0	0	0	0	0	▪	0	1	0
10	▪	0	0	0	0	0	▪	0	▪	0
11	▪	0	▪	▪	0	0	▪	1	▪	▪
12	▪	▪	▪	0	0	▪	▪	0	▪	▪
13	▪	▪	▪	▪	▪	▪	▪	0	▪	▪
14	▪	▪	▪	▪	▪	▪	▪	3	▪	▪

B

Criteria	Score			
	0	1 (mild)	2 (moderate)	3 (severe)
Joint swelling	Normal	Slight to moderate swelling	Moderate/pronounce swelling and/or erythema	Joint distortion or rigidity
Injection site reaction	No reaction	Mild scaring	Moderate ulceration	Severe ulceration
Posture	Normal	Hunching noted only at rest	Severe hunching, some impairment of normal movement	Wobbling, unable to maintain upright posture
Activity	Normal	Mild to moderately decreased	Stationary unless stimulated.	Moderate paralysis, 1 or more limbs
Fur ruffling (fever sign)	Normal	Mild to moderate ruffling	Severe ruffling	Shivering
Hind leg weakness	Normal	Mild hind limb weakness/very mild and/or occasional gait alterations	Moderate hind limb weakness and occasional dragging of hind legs	Severe hind limb weakness/repeated dragging of hind legs
Weight loss	<10 %	10-15%	15-20%	≥20%

Supplementary Figure 2. Disease scores for the ten C57BL/6J mice that were euthanized.
(A) For the mice shown in Fig. 1F and I that were euthanized, the disease scores are shown. (B) The mice were monitored daily using the score card. Any animal reaching a level of 3 in any single criteria were euthanized. If an animal reaches a grade of 2 in two or more criteria the animal will be euthanized.

Supplementary Figure 3. Viral antigen (NS1) staining in C57BL/6J mouse brains. IHC using a pan-flavivirus NS1 monoclonal antibody (4G4). (A-C) The brains from the 3 other C57BL/6J mice that required euthanasia in Fig. 1 (the fourth is shown in Fig. 2). (D) JEV_{FU} infected C57BL/6J mice that lost ~15% body weight then recovered (from Fig. 1E). (E) A representative image of a mouse brain where infection did not lead to significant weight loss. (F) Uninfected control.

Supplementary Figure 4. ApopTag staining in C57Bl/6J mouse brains. Staining for
apoptosis using Apoptag. (A-C) The brains from the 3 other C57BL/6J mice that required
euthanasia in Figure 1. (D) The JEV$_{FU}$ infected C57BL/6J mice that lost ~15% body weight
then recovered (see Figure 1E). (E) Uninfected control.

C57BL/6J	**IRF7⁻ᐟ⁻**	**IFNAR⁻ᐟ⁻; all JEV_{NSW/22}**

MVEV

Nakayama

FU

Nakayama

Nakayama

Nakayama

Nakayama

NSW/22

Uninfected

Supplementary Figure 5. GFAP IHC for reactive astrocytes in mouse brains. Staining for reactive actrocyte using GFAP for the brains from C57BL/6J, IRF7⁻ᐟ⁻ and IFNAR⁻ᐟ⁻ mice that succumbed to the indicated virus. Uninfected mouse brains are also shown.

Supplementary Figure 6. Disease scores for IRF7-/- mice that were euthanized.
Mice described in Figure 3C, D. Scoring system shown in Supplementary Figure 2B.

Anti-flavivirus NS1 (4G4)

JEV$_{Nakayama}$ brain titer 3.9 log$_{10}$CCID$_{50}$/g

JEV$_{Nakayama}$ brain titer 6.6 log$_{10}$CCID$_{50}$/g

100 μm

Supplementary Figure 7. Viral antigen staining in IRF7$^{-/-}$ mouse brains. IHC using a pan-flavivirus NS1 monoclonal antibody (4G4) for the brains from 2 other IRF7$^{-/-}$ mice that required euthanasia (Figure 3). Low levels of staining of neurons were found in the JEV$_{Nakayama}$ infected mice that had a low virus titer in the brain (3.9 log$_{10}$CCID$_{50}$/g) (left). Black arrowheads show small patches of staining in the cortex, hippocampus and thalamus (right).

A

Activity

Dpi	FU Cg4# 1	Cg4# 2	Cg4# 3	Cg4# 4	Cg1# 1	Nakayama Cg2# 1	Cg5# 1	Cg5# 2	Cg5# 3	NSW/22 Cg3# 1	Cg3# 2	Cg3# 3	Cg3# 4	$MVEV_{TC123130}$ Cg3# 1	Cg3# 2	Cg3# 3	Cg3# 4
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1	0	0	0	0	1	1	0	0	0	0	0	0	0	0	0	0	0
2	1	1	1	1	2	1	0	0	0	1	1	1	1	0	0	0	0
3	2	2	2	2	■	2	2	2	2	2	2	2	2	1	1	2	1

Posture

Dpi	FU Cg4# 1	Cg4# 2	Cg4# 3	Cg4# 4	Cg1# 1	Nakayama Cg2# 1	Cg5# 1	Cg5# 2	Cg5# 3	NSW/22 Cg3# 1	Cg3# 2	Cg3# 3	Cg3# 4	$MVEV_{TC123130}$ Cg1# 1	Cg1# 2	Cg1# 3	Cg1# 4
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
2	1	1	1	1	1	1	1	1	0	0	0	0	0	1	0	0	0
3	2	2	2	2	■	2	2	2	2	2	2	2	2	2	2	1	2

Fur ruffling

Dpi	FU Cg4# 1	Cg4# 2	Cg4# 3	Cg4# 4	Cg1# 1	Nakayama Cg2# 1	Cg5# 1	Cg5# 2	Cg5# 3	NSW/22 Cg3# 1	Cg3# 2	Cg3# 3	Cg3# 4	$MVEV_{TC123130}$ Cg1# 1	Cg1# 2	Cg1# 3	Cg1# 4
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
2	1	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0
3	1	1	1	1	■	0	1	1	1	0	0	0	0	1	1	2	1

Weight loss score

Dpi	FU Cg4# 1	Cg4# 2	Cg4# 3	Cg4# 4	Cg1# 1	Nakayama Cg2# 1	Cg5# 1	Cg5# 2	Cg5# 3	NSW/22 Cg3# 1	Cg3# 2	Cg3# 3	Cg3# 4	$MVEV_{TC123130}$ Cg1# 1	Cg1# 2	Cg1# 3	Cg1# 4
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
2	2	1	2	2	2	1	1	1	1	0	0	0	0	0	0	0	0
3	2	1	2	2	■	3	2	2	2	2	1	2	1	2	2	2	2

B

$R^2 = 0.78$
$p<0.0001$

C

$JEV_{NSW/22}$ brain $IFNAR^{-/-}$ mouse, 3 dpi, titer 6.6 $\log_{10}$ $CCID_{50}$/g

Anti-flavivirus NS1   ApopTag

Supplementary Figure 8. JEV and MVEV infection of IFNAR$^{-/-}$ mice. (A) Disease scores as per Supplementary Figure 2B for mice shown in Figure 3. (B) Pearson correlation between brain titer (y-axis) from Figure 3E and viremia (x-axis) from Figure 3A. (B) IHC for flavivirus NS1 (left) or ApopTag (right) showing no detectable staining of brain cells. Images are representative of all IFNAR$^{-/-}$ brains and JEV isolates.

A Acute (euthanized due to weight loss/disease scores)

Mouse	C57BL/6J			*Irf7*[-/-]			*Ifnar*[-/-]		
JEV isolate	NSW	Nakayama	FU	NSW	Nakayama	FU	NSW	Nakayama	FU
Euthanasia		9-14 dpi	8-12 dpi	8 dpi	7-12 dpi	6-10 dpi	3 dpi	3 dpi	3 dpi
Neuronal degeneration /vacuolation	N/A	6/6	2/2	1/1	3/8	7/10	0/4	0/3	0/4
Perivascular cuffing	N/A	6/6	2/2	1/1	8/8	10/10	0/4	0/3	0/4
Hemorrhage	N/A	5/6	1/2	0/1	5/8	6/10	3/4	2/3	2/4
Leukocyte infiltrates	N/A	5/6	2/2	1/1	1/8	3/10	0/4	0/3	0/4
Microgliosis	N/A	6/6	2/2	1/1	5/8	7/10	0/4	0/3	0/4
Meningitis	N/A	6/6	2/2	1/1	6/8	8/10	0/4	0/3	0/3

B

C Chronic (euthanized after recovery from acute phase)

Mouse	C57BL/6J			*Irf7*[-/-]		
JEV isolate	NSW	Nakayama	FU	NSW	Nakayama	FU
Euthanasia	32 dpi	32 dpi	32 dpi	19-21 dpi	19-21 dpi	19-21 dpi
Neuronal degeneration /vacuolation	0/6	0/4	0/5	0/13	0/6	0/4
Perivascular cuffing	1/6	0/4	1/5	2/13	5/6	1/4
Hemorrhage	3/6	2/4	1/5	5/13	5/6	1/4
Leukocyte infiltrates	0/6	0/4	1/5	0/13	0/6	0/4
Microgliosis	0/6	0/4	2/5	2/13	5/6	2/4
Meningitis	0/6	0/4	0/5	1/13	0/6	1/4

Supplementary Figure 9. Lesion presence in H&E brain sections. A) Scoring for lesions described in Figure 6 for mice with acute disease. H&E scoring of 0 indicates no overt presence of these lesions. N/A – not available. B) Ratio of nuclear (blue/dark purple) to non-nuclear (red) staining of H&E stained brain sections (a measure of leukocyte infiltration). Data is the mean and standard error for n=30 infected and n=14 uninfected mouse brains from C57BL/6J and *Irf7*[-/-] mice that succumbed to JEV or MVEV infection (Fig. 1F, 1I and 3D). Statistics by t-test. C) As for 'A' but for mice that survived infection.

Supplementary Figure 10. Histopathological lesions overlapped with areas of virus infection. (A) Cortex regions from JEV infected C57BL/6J, IRF7$^{-/-}$, IFNAR$^{-/-}$ mice or uninfected mice. The cortex regions from JEV infected mice were heavily infected, with most neurons stained positive for viral antigen. H&E detectable signs of neuron degeneration/vacuolation were concentrated in regions with high staining for viral antigen. (B) Viral antigen staining (left) in hippocampus overlapped with histological signs of neuron degeneration/vacuolation (right) (see Figure 6).

A   IRF7$^{-/-}$ and C57BL/6J mice (n=27) that were euthanized due to ethically defined disease end points

	Neuron degeneration/ vacuolation	Perivascular cuffing	Hemorrhage	Leukocyte infiltrates	Microgliosis
Cerebral cortex	17/27	20/27	8/27	8/27	8/27
Hypothalamus	0/27	827	3/27	2/27	7/27
Thalamus	0/27	16/27	2/27	2/27	9/27
Hippocampus	0/27	10/27	0/27	1/27	8/27
Caudate putamen	1/27	16/27	1/27	5/27	7/27
Basal forebrain	0/27	12/27	2/27	0/27	0/27
Midbrain	0/27	7/27	1/27	0/27	5/27
Pons	0/27	6/27	6/27	1/27	1/27
Medulla	0/27	4/27	0/27	0/27	0/27
Cerebellum	0/27	6/27	1/27	0/27	0/27
Anterior olfactory nucleus	0/27	3/27	0/27	0/27	0/27
Ventral striatum	0/27	7/27	0/27	0/27	0/27

B   IRF7$^{-/-}$ and C57BL/6J mice (n=38) that survived infection (taken on day 19-21 and day 32, respectively)

	Neuron degeneration/ vacuolation	Perivascular cuffing	Hemorrhage	Leukocyte infiltrates	Microgliosis
Cerebral cortex	0/38	4/38	7/38	1/38	5/38
Hypothalamus	0/38	0/38	2/38	0/38	0/38
Thalamus	0/38	2/38	1/38	0/38	4/38
Hippocampus	0/38	0/38	0/38	0/38	0/38
Caudate putamen	0/38	4/38	0/38	0/38	5/38
Basal forebrain	0/38	1/38	5/38	0/38	0/38
Midbrain	0/38	0/38	0/38	0/38	2/38
Pons	0/38	0/38	3/38	0/38	1/38
Medulla	0/38	0/38	1/38	0/38	0/38
Cerebellum	0/38	1/38	2/38	0/38	0/38
Anterior olfactory nucleus	0/38	0/38	1/38	0/38	0/38
Ventral striatum	0/38	0/38	2/38	0/38	1/38

Supplementary Figure 11. Histopathological lesions in specific brain regions for C57BL/6J and IRF7$^{-/-}$. Scoring only reflects presence or absence of lesions, and does not indicate severity of lesions. (A) Acute (B) Chronic.

Supplementary Figure 12. RNA-Seq read counts from brains of C57BL/6 JEV-infected and mock-infected mice obtained from the Gene Expression Omnibus (GEO accession: GSE154002) were normalised for sequencing depth and composition using DESeq2. An estimation of cell type abundances was performed on normalised read counts with the SpatialDecon package in R, using an adult mouse immune cell gene expression reference (Yoshida *et al.* 2019. Cell 176; 897-912.e20). Log$_{10}$ abundance scores are shown for cell types that were significantly different between infected and mock-infected groups (t-test, p-value < 0.05), with cell types ordered from largest to smallest difference in mean abundance score between groups.

Supplementary Fig. 13. Human cortical brain organoids measurements and magnified virus staining. (A) Occasional cells staining positive for IHC for viral antigen in hBOs infected with Imojev or YFV 17D. (B) Light microscopy images of hBOs infected with JEV$_{Nakayama}$, JEV$_{FU}$, JEV$_{NSW2022}$, MVEV$_{TC123130}$, Imojev or YFV 17D at day 0 and day 11 post infection compared to uninfected hBOs. Scale bar is consistent across all images. (C) Change in hBO area on day 4, 7, 9 and 11 compared to day 0. n=8 for uninfected and MVEV$_{TC123130}$, n=4 for all others.

Supplementary Figure 14. Replication and CPE in RENcell VM neural progenitor and Vero E6 cell lines. (A) Viral growth kinetics in RENcell VM determined by $CCID_{50}$ assays of culture supernatant at the indicated times post infection. Data is the mean of 6 replicates per virus isolate across 2 independent experiments; (limit of detection is $2 \log_{10}CCID_{50}$/ml). (B) Images of crystal violet stained RENcell VM at 6 dpi (representative of n=3 per group). Less blue/violet stained cells indicates more viral CPE. (C) Vero cells were seeded at 2.5×10^5 cells per well in 24 well plates overnight at 37°C. Cells were infected at MOI=0.05 for 1 hr at 37°C before overlay media (0.375% w/v high viscosity carboxymethyl cellulose [CMC, Sigma-Aldrich]/RPMI 1640 substituted with 2% FCS) was added to each well. Plates were incubated for 5 days before the media was removed and monolayers were fixed and stained with 0.1% w/v crystal violet (Sigma-Aldrich) in formaldehyde (1% v/v) and methanol (1% v/v). Plates were washed in tap water and dried before images were taken. Images shown are representative of n=4 per group. CPE is less pronounced in $JEV_{NEW/22}$ infected cells, with plaques (arrows), rather than fulminant CPE, apparent at 5 dpi.

Supplementary Fig. 15. Percentage neutralisation of JEV and MVEV at serum dilution 1:160. (A) Human serum from cohort 1 (n=9) taken taken pre-vaccination (baseline) and day 28 post-Imojev vaccination was used in plaque reduction neutralisation assays against Imojev, $JEV_{Nakayama}$, $JEV_{NSW/22}$ and JEV_{FU}. (B) Human serum from cohort 2 (n=10) taken >2 months post-Imojev vaccination was used in plaque reduction neutralisation assays against Imojev, $JEV_{Nakayama}$, $JEV_{NSW/22}$ and JEV_{FU} and $MVEV_{TC123130}$. Individual data points in both A and B represent the mean percentage neutralisation at serum dilution 1:160 from duplicate wells in $PRNT_{50}$ assay. The mean of all individuals (n=9 from cohort 1 and n=10 from cohort 2) and standard error is shown. Statistics are by paired t-test.

Supplementary Fig. 16. Neither mouse background nor *Nnt* significantly affected JEV neuropathogenesis. Female C57BL/6J (10-16 weeks old) (black circles), C57BL/6NJ*Nnt-/-* (blue squares), and C57BL/6NJ (red triangles) mice were infected s.c. with 5×10^5 CCID$_{50}$ JEV$_{Nakayama}$. (A) Average viremia as determined by CCID$_{50}$ assays; limit of detection for individual mice is 2 log$_{10}$CCID$_{50}$/ml. (B) Percent body weight change of individual mice compared to their weight 0 dpi. Six mice lost $\geq$20% body weight and required euthanasia (†). (C) Kaplan Myer plot showing percent survival (n=6 for each mouse strain). (D) Viral tissue titers in brains of the six euthanized mice (n=2 from each mouse strain). Tissue titers determined by CCID$_{50}$ assays (limit of detection ~2 log$_{10}$CCID$_{50}$/g). (E) Representative images of H&E stained sections of brains from mice that required euthanasia. No overt differences were identified between C57BL/6J, C57BL/6NJ*Nnt-/-*, or C57BL/6NJ mice.