Title page

Title: The first two chromosome-scale genome assemblies of American hazelnut enable

comparative genomic analysis of the genus *Corylus*

Authors: Scott H. Brainard[1*], Dean M. Sanders[2], Tomas Bruna[3], Shengqiang Shu[3], Julie C.

Dawson[1]

Author affiliations

[1] Department of Horticulture, University of Wisconsin-Madison, Madison, WI 53706, USA

[2] University of Wisconsin Biotechnology Center, University of Wisconsin-Madison, Madison,

WI 53706, USA

[3] U.S. Department of Energy Joint Genome Institute, Berkeley, CA 94720, USA

[*]corresponding author: shbrainard@wisc.edu

ORCID IDs

S. H. Brainard: 0000-0001-7678-3716

D. M. Sanders: 0000-0002-4626-914X

T. Bruna: 0000-0001-9811-5532

S. Shu: 0000-0002-4336-8994

J. C. Dawson: 0000-0002-9907-8611

Abstract

The native shrub American hazelnut (*Corylus americana*) is currently used in breeding programs

that are aiming to develop commercially viable hazelnut varieties for the U.S. Upper Midwestern

U.S. This species provides significant ecological benefits as it is a perennial crop and well-

adapted to this region. Breeding cycles for perennial species are long, and may benefit from the

use of predictive methods such as genomic selection to reduce cycle time and increase the

efficiency of field trials. High-quality reference genome assemblies are very useful for the

implementation marker-assisted selection and genomic prediction, and we therefore developed

the first chromosome-scale reference assemblies for *C. americana*, using the accessions 'Rush'

and 'Winkler'. Initial draft assemblies were created using HiFi PacBio reads and Arima Hi-C

sequencing to assemble genomes into 11 pseudomolecules. We then utilized Oxford Nanopore

reads and a high-density genetic map in order to perform error correction. N50 scores were

calculated to be 31.9 Mb and 35.3 Mb for 'Rush' and 'Winkler', respectively, while 97.1% (for

'Winkler') and 90.2% (for 'Rush') of the total genome was assembled into the 11

pseudomolecules. Gene prediction was performed using both RNAseq libraries as well as protein

homology data. 'Rush' had a BUSCO score of 99.0 for its assembly and 99.0 for its annotation,

while 'Winkler' had corresponding scores of 96.9 and 96.5, indicating extremely high-quality

assemblies. These two independent, *de novo* assemblies enable unbiased assessment of structural

variation across the genome, as well as patterns of syntenic relationships within *C. americana*

and the *Corylus* genus. These assemblies are also an important first step in providing a resource

for using next-generation sequencing data in the improvement of *C. americana*. We demonstrate

this utility through the generation of high-density SNP marker sets from genotyping-by-

sequencing data for 1,343 *C. americana*, *C. avellana*, and *C. americana* x *C. avellana* hybrids, in

order to assess population structure in natural and breeding populations. Finally, the

transcriptomes of these assemblies, as well as several other recently published *Corylus* genomes,

were utilized to perform phylogenetic analysis of sporophytic self-incompatibility (SSI) in

hazelnut, providing further evidence of unique molecular pathways governing self-

incompatibility in *Corylus* not exhibited in other well-studied SSI systems. We hope these

assemblies will aide in the application of modern breeding methods to the development of

commercially viable hazelnut varieties for the U.S. Upper Midwest.

Keywords

Corylus americana, hazelnut, *de novo* genome assembly, long-read sequencing, comparative

genomics, population genetics, sporophytic self-incompatibility

Declarations

Funding: Sequencing and bioinformatics was supported by the Jewett Prize of the Arnold

Arboretum and USDA-NIFA SCRI Grant No. H007913501, with matching funds from the

Savanna Institute and The Grantham Foundation for the Protection of the Environment. Genome

annotation conducted at the U.S. Department of Energy Joint Genome Institute was supported by

the Office of Science of the U.S. Department of Energy under Contract No. DE-AC02-

05CH11231.

Authors' contributions: SHB and JCD designed the analysis. DMS and SHB carried out genome

assembly and SNP calling. SHB, TB and SS performed genome annotation. SHB performative

comparative genetic analyses and prepared the manuscript, with contributions from all authors.

Acknowledgments

We thank Al Kovalevski, Samridhi Chaturvedi and Ashely Yow for providing helpful advice in

the performance of phylogenetic analysis. In addition, Nahla Bassil, Shawn Mehlenbacher, Tom

Molnar, Malachi Persche, Lois Braun, and Mark Hamann assisted with the collection of tissue

samples used in this study. We thank Thomas Hickey and Marissa Nix as well as the staff of the

University of Wisconsin–Madison Biotechnology Center DNA Sequencing Facility and the

USDA National Clonal Germplasm Repository in Corvallis, OR for logistical support of the

research. We thank Chuck and Gerta Zinda for allowing us to study the *C. americana* planted on

their property in Barneveld, WI for our research.

Main Text

Introduction

Hazelnut (*Corylus* spp.) is a globally significant nut crop, with annual production of over 1

million tons, produced across 34 countries . This production is driven by a wide range of market

uses across diverse industries, including food products, pharmaceuticals, and dietary

supplements (Ceylan et al., 2022). Currently, commercial cultivation relies overwhelmingly on

cultivars of European hazelnut (*C. avellana*) adapted to Mediterranean climates (Di Lena et al.,

2022) with moderate winters to satisfy their chilling requirements (Mehlenbacher, 1991). The

distribution of hazelnut production reflects these climatic requirements, with major centers of

production in Turkey, Italy, and Spain, with Turkey alone producing over 65% of the global

supply (Semih Uzundumlu et al., 2022). In the U.S., this narrow range of adaptation currently

restricts production to relatively small regions such as the Willamette Valley in Oregon (Revord

et al., 2020).

As a woody perennial crop, hazelnuts have substantial potential to provide ecosystem services

such as reduced soil erosion and nutrient runoff (Demchik et al., 2014), while also sequestering

carbon in above and belowground biomass (Granata et al., 2020). Expanding the climatic

conditions under which commercially viable hazelnut varieties can be grown would not only

make production more resilient in the face of changing climates, but provide high-value

alternative crops to producers across wider geographies. One possible method for the

development of commercially viable hazelnut varieties for additional growing regions, is the

improvement of the native North American shrub *C. americana*, which possesses disease

resistance and cold hardiness traits lacking in most cultivated forms of European hazelnut

(Molnar et al., 2018). To this end, genomics-aided breeding of *C. americana*, utilizing methods

such as marker-assisted selection and genomic prediction, would be greatly aided by a high-

quality reference genome assembly. Such a resource would, for example, facilitate the rapid

identification and alignment of genetic polymorphisms across experimental and breeding

populations (Kang et al., 2016). In addition, well-annotated assemblies provide the opportunity

to investigate the genetic components underlying specific biomolecular pathways involved in

abiotic and biotic stress resistance, employing methods ranging from fine mapping of

quantitative trait loci (QTL) to gene editing (Bolger et al., 2017; Dmitriev et al., 2022).

To facilitate such applied and experimental approaches, this study reports the first chromosome-

scale reference assemblies for *C. americana* using two accessions, 'Rush' and 'Winkler', which

have historically been widely used in hazelnut breeding in the Eastern United States, largely as

sources of resistance to the endemic fungal pathogen *Anisogramma anomala* (Eastern Filbert

Blight; EFB) (Bhattarai et al., 2017). Constructed using a combination of long-read sequencing,

chromosome conformation capture, and genetic mapping, these chromosome-scale reference

assemblies are of high-quality and can immediately be deployed in advancing breeding and

genetic research objectives. Such potential is illustrated in this study through the identification of

high-density single nucleotide polymorphism (SNP) markers for over 1,343 hazelnut plants,

drawn from Midwestern natural and breeding populations. These markers provide a detailed

assessment of population structure in *C. americana* and interspecific hybrids between *C.*

americana and *C. avellana*. 20[th] century hazelnut breeding the Midwestern U.S. involved

interspecific hybridization with *C. avellana* (Weschcke, 1954; Rutter, 1987), but generations of

open-pollination and lack of detailed pedigree records obfuscate the genetic background of

current Midwestern varieties. An analysis of population structure helps to resolve this question.

Understanding the proportional representation of *C. americana* and *C. avellana* genetic

backgrounds in varieties which have been successful selected under Midwestern conditions will

help determine breeding strategies for key traits.

In addition, the last two years has seen the release of chromosome-scale genome assemblies for

several other *Corylus* species: *C. avellana* (cultivars 'Tombul' (Lucas et al., 2021) and 'Tonda

Gentile della Langhe' ('TGdL') (Pavese et al., 2021)), *C. heterophylla* (accessions from Siping

City, Jilin (Liu et al., 2021) and Yanqing, Beijing (Zhao et al., 2021)), and a wild specimen of *C.*

mandshurica (Li et al., 2021). Together with the two *C. americana* genomes reported here, this

study presents the first comparative genomic analyses across nearly half of the *Corylus* genus.

Comparative genetic analysis also is useful for analyzing specific traits, such as self-

incompatibility, an understanding of which is critical to both hazelnut breeding and production.

Plants have evolved numerous strategies to increase the frequency of outcrossing, ranging from

variation in the timing of floral development, to genetically regulated mechanisms. With respect

to the latter, two primary classes of self-incompatibility have been identified: gametophytic self-

incompatibility (GSI) and sporophytic self-incompatibility (SSI) (Silva & Goring, 2001). The

former is typified by *Rosacea* and *Solanaceae* species, wherein the haploid genotype of the male

gamete at an "S-locus" (often encoding an F-box (SLF/SFB) protein) is detected by the female

parent, which blocks the growth of the germinating pollen tube (Sijacic et al., 2004; Sassa,

2016). The latter is exemplified by *Brassicaceae* species, where the diploid genotype of the male

parent at the S-locus (often encoding a cysteine-rich protein (SCR/SP11)) mediates detection by

the female parent, and a suppression of pollen germination (Hiscock & McInnis, 2003). Self-

incompatibility in *Corylus* is a form of SSI, regulated by a single S-locus (Mehlenbacher &

Thompson, 1988), which fluorescent microscopic observation of pollen tube growth has been

demonstrated to include at least 33 alleles with 8 levels of linear dominance (Mehlenbacher,

1997, 2014). We pursue a phylogenetic approach to the analysis of SSI using the transcriptomes

reported here, and this analysis sheds additional light on the evolution of this mechanism in

Corylus.

Materials and methods

Plant material collection

Tissue was collected from *C. americana* accessions 'Rush' and 'Winkler' maintained at the

National Clonal Plant Germplasm Repository (NCGR) in Corvallis, Oregon (PI 557022 and PI

557019, respectively). 'Rush' is a specimen collected around 1900 by J.F. Jones in Lancaster,

Pennsylvania, while 'Winkler' was collected in 1910 by Wendell Williams in Danville, Iowa

(Molnar, 2011). These two varieties were historically significant in early- and mid-20[th] century

hazelnut breeding programs, and as such represent genetically relevant, pure *C. americana*

selections, which also reflect the wide geographic distribution of *C. americana* in the Eastern

U.S. Today, 'Rush' remains a widely used source of EFB resistance (Bhattarai et al., 2017).

Photos of the clones located at the NCGR are shown in Fig. 1A. Tissue was collected on April

13[th], 2020, from bushes that had been etiolated for three days prior to sampling in order to

minimize the concentration of volatiles and secondary metabolites in leaf tissue. Samples were

immediately flash-frozen using liquid nitrogen and stored at -80°C until DNA extraction.

Long-read sequencing

High-molecular weight DNA was extracted following the protocol described by Vaillancourt &

Buell (2019). In brief, leaf tissue was homogenized in liquid nitrogen, lysed with Carlson lysis

buffer, and purified with chloroform and Qiagen Genomic-tips (Qiagen N.V., Venlo, The

Netherlands). Purity of extracted DNA was assessed spectrophotometrically using a NanoDrop™

One (ThermoFischer Scientific, Waltham, Massachusetts). DNA was quantified a using Qubit™

dsDNA High Sensitivity kit (ThermoFisher Scientific, Waltham, Massachusetts), and diluted and

assessed for size using an Agilent FemtoPulse System (Santa Clara, California).

Pacific Biosciences HiFi libraries were prepared according to PN 101-853-100 Version 03

(Pacific Biosciences, Menlo Park, California). Modifications included shearing with a Covaris g-

TUBE (Covaris, Woburn, Massachusetts) and size selecting with BluePippin (Sage Science,

Beverly, MA). Libraries was sequenced on a PacBio Sequel II using the Sequel Polymerase

Binding Kit 2.2 at the University of Wisconsin-Madison Biotechnology Center's DNA

Sequencing Facility. Oxford Nanopore libraries were prepared following the Native Barcoding

Expansion protocol (Oxford Nanopore Technologies, Oxford, United Kingdom). The library was

sequenced on a R9.4.1 flowcell on an Oxford Nanopore PromethION, also at the UW-Madison

Biotechnology Center. Partway through the sequencing run, DNA was flushed with the Oxford

Nanopore Technologies Flow Cell Wash Kit (EXP-WSH004), and additional library was loaded.

Hi-C sequencing

Nuclei were extracted from flash-frozen leaf tissue using a Sigma CelLytic™ PN Plant Nuclei

Isolation/Extraction Kit (Sigma-Aldrich, Burlington, Massachusetts). Crosslinking was

performed following low-input protocols from Arima Genomics (Arima Genomics, Carlsbad,

California). Crosslinked nuclei were quantified using a Qubit™ dsDNA High Sensitivity kit,

samples were sheared to 600bp, and library preparation was performed using a KAPA® Hyper

Prep kit (Roche, Basel, Switzerland). The final library was assessed for quality using the Agilent

TapeStation System with a D1000 kit (Agilent Technologies, Santa Clara, California), and

paired-end, 150 bp paired-end sequences were generated using an Illumina NovaSeq 6000

(Illumina, San Diego, California).

RNA extraction, library preparation and sequencing

Prior to RNA extraction, Tissuelyser II adapters, safelock tubes, and 5mm stainless steel beads

were chilled at -80°C for 2 hours. Working on dry ice, leaf and kernel tissue was added to

prechilled tubes and disrupted using the Tissuelyser II (Qiagen N.V., Venlo, The Netherlands)

for 2 rounds at 30Hz for 1 minute each round. Samples were then processed using the Qiagen

Plant RNeasy (Qiagen N.V., Venlo, The Netherlands) with an on-column DNA digest. Total

RNA was assayed for purity and integrity using a NanoDrop One™ Spectrophotometer and

Agilent 2100 Bioanalyzer (Agilent Technologies, Santa Clara, California), respectively. RNA

libraries were prepared from samples that met the Illumina TruSeq® Stranded Total RNA

Sample Preparation Guide (15031048 E) input guidelines using the Illumina TruSeq® Stranded

Total (Plant) RNA Sample Preparation kit (Illumina Inc., San Diego, California, USA). For each

library preparation, cytoplasmic, mitochondrial and chloroplast ribosomal RNA was removed

using biotinylated target-specific oligos combined with paramagnetic beads tagged with

streptavidin. Following purification, the reduced RNA was fragmented using divalent cations

under elevated temperature. Fragmented RNA was copied into first stranded cDNA using

SuperScript II Reverse Transcriptase (Invitrogen, Carlsbad, California, USA) and random

primers. Second strand cDNA was synthesized using a modified dNTP mix (dTTP replaced with

dUTP), DNA Polymerase I, and RNase H. Double-stranded cDNA was cleaned with AMPure

XP Beads (1X) (Beckman Coulter, Brea, California,). The cDNA products were incubated with

Klenow DNA Polymerase to add a single "A" nucleotide to the 3' end of the blunt DNA

fragments. Unique dual indexes (UDI) were ligated to the DNA fragments and cleaned with two

rounds of AMPure XP beads (0.8X). Adapter-ligated DNA was amplified by PCR and cleaned

with AMPure XP beads (0.8X). Final libraries were assessed for size and quantity using an

Agilent DNA1000 chip and a Qubit® dsDNA HS Assay Kit (Invitrogen, Carlsbad, California,

USA), respectively. Libraries were standardized to 2 nM, and 150-bp paired-end sequencing was

performed on an Illumina NovaSeq 6000.

Genome assembly

An initial assembly was created using PacBio CCS reads with the program hifiasm v0.16.0-r369

(Cheng et al., 2021) on a Unix server with 160 cores and 3TB of RAM. This draft assembly

exhibited higher contiguity than references generated with Canu using ONT reads, HiCanu, and

Flye using either HiFi or ONT reads. The program purge_haplotigs v1.1.1 (Roach et al., 2018)

was then utilized to reduce artificial genome duplication caused by incorporation of non-

collapsed haplotigs in the final assembly. Hi-C reads were aligned, filtered, and binned using the

haplotig-purged assembly using the program juicer (Durand et al., 2016). Contigs were then

scaffolded and ordered using the program 3D-DNA (Dudchenko et al., 2017).

While integration of Hi-C data led to improved contiguity and reduced artificial duplication in

the 'Winkler' assembly, contigs in 'Rush' were fragmented following the use of juicer and 3D-

DNA. As a result, this step was omitted for 'Rush'. The assemblies were next iteratively polished

three times with Racon (Vaser et al., 2017), using the original PacBio reads. We then used a

recently constructed linkage map to detect erroneous inversions in the physical assemblies

(Brainard et al., 2023), using visual inspection of heatmaps of recombination frequencies in an F_1

population. The program Ragtag v2.1.0 (Alonge et al., 2022) was then used to correct observed

errors using ONT reads which overlapped any identified inversions. Initial quality assessments

were made using BUSCO v5.4.3 (Manni et al., 2021) with the eudicots_odb10 database, and

quast v5.2.0 (Mikheenko et al., 2018) with default parameters.

Genome annotation

Transcript assemblies for both 'Rush' and 'Winkler' were made from ~200M 2 x 150 bp

stranded paired-end Illumina RNAseq reads using the software PERTRAN, which conducts

genome-guided transcriptome short read assembly via GSNAP (Wu & Nacu, 2010) and builds

splice alignment graphs after alignment validation, realignment, and correction. These outputs

were subsequently used to construct 37,794 and 36,419 transcript assemblies for 'Rush' and

'Winkler', respectively, using the program PASA (Haas et al., 2003). Loci were determined by

transcript assembly alignments and/or EXONERATE alignments of proteins from *Mimulus*

guttatus, *Arabidopsis thaliana*, *Gossypium raimondii*, *Betula platyphylla*, *Carya illinoinensis*,

Castanea dentata, *Quercus rubra*, *Prunus persica*, *Fragaria vesca*, *Glycine max*, *Medicago*

truncatula, *Vitis vinifera*, *Liriodendron tulipifera*, *Juglans microcarpa*, *Populus trichocarpa*,

Beta vulgaris, *Solanum lycopersicum*, *Sorghum bicolor*, *Oryza sativa*, and Swiss-Prot (release

2022_04 of eukaryote proteomes), with up to 2k bp extension on both ends unless extending into

another locus on the same strand. Alignments were made to the respective genome assemblies

following softmasking for repetitive elements. The repeat library consisted of *de novo* repeats

identified by RepeatModeler2 (Flynn et al., 2020), generated using both the 'Rush' and

'Winkler' assemblies, as well as *C. americana* repeats identified in the databases RepBase and

Dfam. Gene models were predicted by homology-based predictors: FGENESH+ (Salamov &

Solovyev, 2000), FGENESH_EST (which is similar to FGENESH+, but uses EST to compute

splice site and intron input instead of protein/translated ORF), EXONERATE (Slater & Birney,

2005), PASA assembly ORFs (a homology constrained ORF finder), and AUGUSTUS (Stanke

et al., 2006) trained by the high confidence PASA assembly ORFs, and with intron hints from

short read alignments. The best-scored predictions for each locus were selected using multiple

positive factors including EST and protein support, and overlap with repeats as a negative factor.

The selected gene predictions were improved by PASA. The improvement included adding

UTRs, splicing correction, and adding alternative transcripts. PASA-improved gene model

proteins were subject to protein homology analysis to the above-mentioned proteomes to obtain a

Cscore and protein coverage. Cscore is a protein BLASTP score ratio to the mutual best hit

(MBH) BLASTP score, and protein coverage is the highest percentage of protein aligned to the

best homologs. PASA-improved transcripts were selected based on Cscore, protein coverage,

EST coverage, and their CDS overlap with repeats. The transcripts were selected if their Cscore

and protein coverage were ≥ 0.5 or if covered by ESTs. For gene models whose CDS were

overlapped by repeats by more than 20%, their Cscore was required to be at least 0.9, and

homology coverage at least 70%, in order to be selected. The selected gene models were subject

to Pfam analysis, and gene models whose proteins were more than 30% overlapped by Pfam TE

domains were removed as weak gene models. Incomplete gene models, low homology supported

gene models without full transcriptome support, short single exon (< 300 bp CDS) with neither

protein domains nor good expression, and repetitive gene models without strong homology

support were manually filtered out.

Functional annotation was performed using the software OmicsBox (BioBam Bioinformatics,

2019). First, NCBI Blast+ v2.10.0 (Altschul et al., 1990) was used to BLAST each gene against

the Viridiplantae taxon of the UniProtKB/Swiss-Prot v5 database (The UniProt Consortium,

2023), using a e-value threshold of 1e-3, a word size of 6, and an HSP cutoff length of 33. Gene

ontology (GO) terms were then mapped and annotated using the Blast2GO pipeline (Gotz et al.,

2008) and GOA v2022.08 (Huntley et al., 2015). Next, InterProScan v5.59-91.0 (Jones et al.,

2014) and eggNOG-v 5.0.2 (Huerta-Cepas et al., 2019) were used to add additional GO terms by

mapping protein domains to their respective databases. All annotations were subsequently

merged. Finally, tandem repeats were identified using TandemRepeatFinder (Benson, 1999)

with default parameters, and tRNAs were predicted using tRNAscan-SE (Chan & Lowe, 2019).

Genotyping by sequencing

To better compare genomes of *C. avellana* and *C. americana*, we used GBS to genotype a

population of interspecific hybrids and a population of *C. americana.* Tissue from 1,343 samples

of breeding lines from the Upper Midwest Hazelnut Development Initiative (UMHDI), Oregon

State University, in Corvallis, Oregon, and Rutgers University, in New Brunswick, New Jersey,

along with full-sib F_1 populations derived from controlled crosses between these varieties, and a

wild Midwestern population of *C. americana* sourced from the Wisconsin Department of Natural

Resources and planted in Barneveld, WI, was sampled following budbreak in May of 2020.

Genomic DNA was extracted and libraries for genotyping-by-sequencing were prepared using a

double digestion with the restriction enzymes *Nsi*I and *Bfa*I following the methodology described

by Elshire et al. (2011). This specific double digest was selected based on analysis of k-mer

distributions of sequence libraries generated with *ApeK*I alone, *Nsi*I and MspI, *Pst*I and *Bfa*I,

*Pst*I and *Msp*I, *ApeK*I and *Bfa*I, and *ApeK*I and *Msp*I. Illumina GBS barcodes and adapters were

ligated, and paired-end reads (2 x 150 bp, 10 million reads / sample) were generated using an

Illumina NovaSeq 6000.

Due to the high levels of synteny across the genomes described above, direct use of Illumina

sequence data to infer variable degrees of interspecific hybridization, using tools such as sppIDer

(Langdon et al., 2018), is limited. Biallelic SNPs were therefore identified using the

TASSEL GBSv2 pipeline (Bradbury et al., 2007) and filtered for missing data (< 10% across all

samples), minor allele frequency (> 0.05), linkage disequilibrium ($r^2 < 0.75$), and allele depth

(80% of samples with a depth > 8) using bcftools (Danecek et al., 2021). Population structure

analysis using principal components analysis (Price et al., 2006) has been shown to be a simple

and efficient alternative to more complex model-based approaches such as STRUCTURE

(Falush et al., 2007) and ADMIXTURE (Alexander et al., 2009). This filtered VCF file was

therefore converted to a Euclidean distance matrix using R (R Core Team, 2021), and multi-

dimensional scaling (which permits some missing data) was then performed to obtain measures

analogous to scores along the first two principal components of the distance matrix.

Comparative analyses of Corylus

Comparisons across the *Corylus* genus were made utilizing the five currently available

chromosome-scale genome assemblies, as well as the *C. americana* assemblies reported here for

'Rush' and 'Winkler'. This included two *C. avellana* assemblies (cultivars 'Tombul' (Lucas et

al., 2021) and 'Tonda Gentile della Langhe' (Pavese et al., 2021)), two *C. heterophylla*

assemblies (an accession from Siping City, Jilin (Liu et al., 2021) and an accession from

Yanqing, Beijing (Zhao et al., 2021)), and one *C. mandshurica* assembly (a wild specimen from

Xinglong (Li et al., 2021)). All of these assemblies were annotated using similar *ab initio*

prediction methods which combined RNAseq data and protein homology data, and thus their

transcriptomes were well-suited to comparative analyses with 'Rush' and 'Winkler'.

GENESPACE v0.94 (Lovell et al., 2022) was used to investigate genome-wide syntenic

relationships, with *Betula pendula* (silver birch) and *Malus domestica* (apple) included as

outgroups (gene models being obtained from https://genomevolution.org/coge/ and

https://www.rosaceae.org, respectively). In brief, GENESPACE utilizes the program

OrthoFinder (Emms & Kelly, 2019) to identify orthogroups from predicted gene models, and

then parses orthologs to define syntenic blocks across species using BLAST and MCScanX

(Wang et al., 2012). Genes in *C. americana* with no orthologous sequences in any other *Corylus*

species were also identified using OrthoFinder.

Phylogenetic analysis of sporophytic self-incompatibility

Sequences for three S-locus genes (MIK3 homologues) reported by Hou et al. (2022) were

obtained from the transcriptome reported in Zhao et al. (2021). NCBI BLAST v2.6.0+ (Altschul

et al., 1990) was used to identify homologous sequences in all *Corylus* species, as well as *B.*

pendula, *M. domestica*, and *Brassica oleracea*. These latter three species were included as

variably-related outgroups which also exhibit self-incompatibly: SSI in the case of *B. oleracea*

and *B. pendula*, and GSI in the case of *M. domestica*. The *B. oleracea* transcriptome was

obtained from http://brassicagenome.net/. Multiple sequence alignments were made using

muscle v5.1 (Edgar, 2021). These 84 alignments were then imported into MEGA11 (Tamura et

al., 2021) and used to build a phylogenetic tree using maximum likelihood and the Jones-Taylor-

Thornton matrix-based model (Jones et al., 1992). A bootstrap consensus tree was inferred from

100 replicates, in which branches present in less than 50% of these replicates were collapsed.

Trees were visualized and clades labeled using the program FigTree v1.4.4.

Results

Sequencing and assembly

Table 1 summarizes the raw sequence data used in assembling the two genomes. For 'Rush',

14.97 Gb of CCS sequence (44.32x coverage) and 13.60 Gb of ONT sequence (40.29x coverage)

were generated. For 'Winkler', 19.60 Gb of CCS sequence (58.02x) and 12.69 Gb of ONT

sequence (37.57x coverage) was generated. Short read sequencing of Arima Hi-C libraries

generated 78.7x and 114x coverage for 'Rush' and 'Winkler', respectively. While 'Winkler'

libraries therefore generated substantially more data across all sequencing platforms, in the case

of both accessions, coverage was judged to be more than sufficient for the assembly of a diploid

organism, assumed to have a relatively small genome size of ~350-370 Mb, based on

comparisons with previous *Corylus* assemblies (Li et al., 2021; Liu et al., 2021; Lucas et al.,

2021; Pavese et al., 2021; Zhao et al., 2021). The contact map for 'Winkler', showing contigs

scaffolded into chromosome-scale pseudomolecules, is shown in Fig. 1C (visualized using

Juicebox (https://github.com/aidenlab/Juicebox)).

Table 1. Summary of sequence data generated on PacBio, Oxford Nanopore and Illumina
sequencing platforms.

Sample	Sequence Type	Reads	Avg. length (bp)	Coverage
'Rush'	PacBio CCS	1,026,165	14,584	44.3x
'Winkler'	PacBio CCS	1,264,282	15,496	58.0x
'Rush'	ONT	581,538	22,053	40.3x
'Winkler'	ONT	616,903	21,814	37.6x
'Rush'	Arima Hi-C	88,601,323	2 x 150	78.7x
'Winkler'	Arima Hi-C	129,091,816	2 x 150	114.7x

Table 2 presents statistics summarizing the quality and contiguity of the 'Rush' and 'Winkler'

assemblies. While both genomes exhibit high quality and contiguity, with N50 scores > 30 Mb,

and over 90% of the genome assembled into 11 pseudomolecules, for both accessions there are

clear differences which reflect the divergence in assembly methods. Both assemblies are close in

length to previous reports of the size of *Corylus* species, however the 'Rush' assembly is over 50

Mb larger than the 'Winkler' assembly. This is likely due to unresolved duplication, apparent

also in Fig. 1D, due to the fact that Hi-C sequence data could not be included in the assembly,

which is also reflected in the many more contigs present in 'Rush' compared to 'Winkler'. While

including Hi-C data for 'Winkler' led to a reduction in total contigs from 398 (following hifiasm

and haplotig_purge) to the 264 reported in Table 1, including Hi-C data for 'Rush' led to an

expansion and fragmentation of contigs to 1,845. Benchmarking Universal Single Copy

Orthologs (BUSCOs) were calculated for 'Rush' and 'Winkler', and while both assemblies had

nearly identical complete and single copy BUSCOs represented (95.1% and 94.6%, respectively)

'Rush' contained more duplicates (3.8% vs. 1.4%). This finding, and the slightly larger

maximum contig size also suggest unresolved artificial duplication.

Table 2. Summary of statistics related to genome assembly. BUSCO score codes: C: complete;
S: complete and single copy; D: complete and duplicated; F: fragmented; M: missing.

Statistic	'Winkler'	'Rush'
Assembly size (bp)	337,645,099	388,211,906
Number of contigs	264	1118
GC content (%)	36.05	36.68
Assembly L50 (#)	5	5
Contig N50 (bp)	31,913,876	35,381,848
Maximum contig size (bp)	44,790,255	46,319,783
Genome in chromosomes (%)	97.1	90.2
BUSCO (C [S, D]; F; M) (%)	96 [94.6, 1.4]; 0.9; 3.1	98.9 [95.1, 3.8]; 0.2; 0.9

Both assemblies were also annotated for several genomic features: gene models were predicted

using a combination of RNAseq data and protein homology data; transposable elements were

identified using RepeatModeler; tandem repeats were identified using TandemRepeatFinder; and

tRNAs were predicted using tRNAscan-SE. These features are presented in a Circos-style plot in

Fig. 1B. Specific statistics related to the gene model predictions are summarized in Table 3.

Table 3. Summary of statistics for the annotation of 'Rush' and 'Winkler'.

Statistic	'Winkler'	'Rush'
Primary transcripts	23,331	24,562
Alternative transcripts	5,701	6,174
Median exon length (bp)	159	159
Median intron length (bp)	245	248
Average number of exons per gene	5.7	5.7

These annotations appear extremely similar in terms of the structure of the gene models which

were predicted. Slightly fewer genes were predicted in 'Winkler', which in this case is likely not

the consequence of unresolved duplication, as the genomes were soft masked for repetitive

elements. Finally, additional statistics related to the genome annotation are provided in Table 4,

which compares the 'Rush' and 'Winkler' assemblies to the five currently published

chromosome-scale *Corylus* assemblies.

Table 4. Comparison with annotations of previously published *Corylus* species. BUSCO scores
refer to all complete representatives, both single and duplicated.

Species and cultivar name or accession)	Genome size (11 chromosomes)	Repeat content	Predicted genes	Genome BUSCO	Annotation BUSCO
C. americana ('Winkler')	327.7	48.4%	23,331	96.9	96.5
C. americana ('Rush')	350.2	49.0%	24,562	99.0	99.0
C. mandshurica	367.7	68.7%	25,923	97.1	92.2
C. heterophylla (Beijing)	361.9	56.7%	27,591	93.5	92.0
C. heterophylla (Jilin)	342.9	58.0%	22,319	94.7	92.7
C. avellana ('TGdL')	373.1	41.5%	27,791	96.2	92.0
C. avellana ('Tombul')	369.8	38.1%	27,270	97.0	76.8

The total number of predicted genes are within the range previously reported in *Corylus*. In

addition, the BUSCO scores for 'Rush' and 'Winkler' are substantially higher than the scores

reported for any previous *Corylus* genome annotation, and are very close to the assembly

BUSCO scores. This suggests not only that these annotations are highly complete, but also

provides support for the accuracy of the predictive procedure described above.

Comparison across the Corylus *genus*

Comparisons across multiple species can be made simultaneously, or in a pairwise manner.

Using the five previously published *Corylus* assemblies, which contain comparable annotations

to the genome presented here, together with the closely related species *Betula pendula* and

outgroup *Malus domestica*, the GENESPACE package (Lovell et al., 2022) was utilized to

visualize syntenic relationships in both manners, as well as construct a pangenome assembly

anchored to *C. americana* 'Winkler'. Fig. 2 illustrates the general result that across these *Corylus*

genomes, there is almost universal synteny between all defined blocks, illustrating the high

degree of relatedness between each pair of species. Differences in pseudomolecule numbering is

most likely an artifact of chromosome numbering typically being made according to descending

physical contig size. *Corylus* chromosome sizes are generally very similar, and assemblies had

slight variations in contig lengths. Fig. 3 shows dot plots representing each pairwise comparisons

between syntenic blocks for each of the seven *Corylus* genomes. Despite the high degree of

synteny across all *Corylus* species on a chromosome scale, it is clear that *C. americana* exhibits

nearly perfect synteny with the two *C. avellana* assemblies. This decreases marginally when

compared against the two *C. heterophylla* assemblies, with small inversions evident on a number

of chromosomes. *C. mandshurica* very clearly exhibits the least amount of synteny, with large

inversions and rearrangements of syntenic blocks across most chromosomes.

With respect to *B. pendula*, this figure makes clear the specific large-scale rearrangements which

relate the 14 chromosomes of *B. pendula* to the 11 of *Corylus* spp. Specifically, chromosomes 1

and 3 in *B. pendula* were split to form chromosomes 1 and 4 in *Corylus* spp., with *B. pendula*

chromosome 11 also being fused with the end of chromosome 1 in *Corylus*. In addition,

chromosomes 4 and 6 in *B. pendula* together constitute chromosome 2 in *Corylus*, while

chromosomes 12 and 14 constitute chromosome 3 in *Corylus*.

Single copy orthologs unique to C. americana

It is also possible to use the orthogroups generated by OrthoFinder to determine, instead of

synteny, specifically those putative genes in *C. americana* which possess no ortholog in any

currently annotated *Corylus* species, nor the other two outgroups included in this analysis: *B.*

pendula and *M. domestica*. This subset of the transcriptome consists of 87 predicted single copy

orthologs present in both 'Rush' and 'Winkler', but absent in all other analyzed genomes. Many

of these genes logically did not return hits when the Viridiplantae database was queried using

NCBI-BLAST. Fig. 2B presents the GO terms associated with those genes which were able to be

annotated following the methods described above. Of particular note is the gene

CamerWinkler.08G009300, which is characterized as involved in "defense response to fungus".

Given *C. americana*'s high level of resistance to the endemic fungal pathogen EFB, this gene

would be a valuable targets for future functional characterization.

Contributions of C. americana *and* C. avellana *to current Midwest germplasm*

1,343 individual plants were sequenced using GBS, and SNP markers were called using

TASSEL. Sampled plants represented wild *C. americana*, cultivars from breeding programs in

Oregon, New Jersey, and Minnesota, and F_1 populations between these cultivars. A biplot for the

first two principal components of the distance matrix constructed using these markers is shown

in Fig. 4. This visualization makes evident the wide genetic diversity represented in current

Midwestern hazelnut varieties, and lends support to the significant *C. avellana* contribution to

specific selections, such as Rose9-2. Many currently named Midwestern varieties, on the other

hand, cluster extremely closely with 'Rush', 'Winkler', and wild *C. americana* from the DNR,

suggesting that while an interspecific cross may have occurred in the past, subsequent selection

of progeny greatly favored *C. americana* genetic contributions.

Phylogenetic analysis of sporophytic self-incompatibility

The locus regulating sporophytic self-incompatibility (SSI) in hazelnut has been fine-mapped in

C. avellana (Hill et al., 2021) and *C. heterophylla* x *C. avellana* interspecific hybrids (Hou et al.,

2022). These studies identified three MIK2 homologues believed to be responsible for SSI in

these two *Corylus* species. In order to examine their phylogenetic relationship, multiple sequence

alignment was performed between the top BLAST hit for each of these three genes and all

currently available *Corylus* transcriptomes. Outgroups were included as above for *B. pendula*

and *B. oleracea*, which both also exhibit SSI (Hynynen et al., 2010; Kitashiba & Nasrallah,

2014), as well as *M. domestica*, a more distantly-related species that is a well-studied example of

gametophytic self-incompatibility (Cheng et al., 2006; Minamikawa et al., 2010). This

phylogenetic tree is shown in Fig. 5, and illustrates the genus-specific evolution of SSI in

Corylus, including several clades containing orthologous sequences only found in *Corylus*

species.

Discussion

This assembly is a critical step in developing the use of next-generation sequencing data in the

improvement of *C. americana*, while also enabling comparisons of structural variation across

Corylus species.

Variable contiguity of two de novo *assemblies*

The fragmentation of the 'Rush' assembly by the 3D-DNA pipeline (using Arima Hi-C sequence

data) was unexpected, given the widespread and successful use of this method in building

chromosome-scale assemblies (Ghurye et al., 2019), and in particular, the substantial

improvements in contiguity that was observed when using this data to refine the initial 'Winkler'

assembly. Two possible explanations for this discrepancy suggest themselves. First, the 'Rush'

accession generated less absolute sequence compared to the 'Winkler' assembly on all

sequencing platforms, and as a result there was less sequence data to include in each

bioinformatic step. The raw assemblies generated by hifiasm were indeed also more contiguous

for Winkler compared to 'Rush'. Nevertheless, with coverage >40x for all sequence data types,

the failure of Hi-C sequence to improve contiguity would not appear to be solely a consequence

of a lack of data. A second likely compounding factor, however, is a difference in the relative

heterozygosity of 'Rush' and 'Winkler'. Heterozygosity is known to complicate phased genome

assembly algorithms which use HiFi and Hi-C sequence data (Garg, 2021). Using the 53,696

GBS-derived SNPs described above it was possible to directly estimate genome-wide

heterozygosity as a ratio of heterozygous to total polymorphic sites. By this metric, 22%

heterozygosity was observed for 'Rush', while only 17% of polymorphic sites were

heterozygous in 'Winkler', suggesting that this could have contributed to the challenges in

scaffolding 'Rush' using Hi-C data.

Regardless of the specific underlying factors complicating this stage of genome assembly, it is

clear that even in diploid species with relatively small genome sizes, integrating chromosomal

conformation capture sequencing data into genome assembly pipelines is not a universally

applicable method. At the same time, the use of high-coverage long-read sequence data,

combined with modern assemblers for scaffolding and polishing chromosome-scale genomes,

has substantial capacity to generate high-quality reference assemblies, even in outcrossing,

heterozygous organisms.

PCA-derived population structure

This study reports the first combined analysis of relatedness between commercial varieties of

hazelnuts from the three main breeding programs in the United States: Oregon State University,

Rutgers University, and the Upper Midwest Hazelnut Development Initiative. This evaluation of

population structure, by combining a diversity of clonal varieties with F_1 full-sib families

produced through multiple pairwise controlled crosses of these clones, help elucidate a long

outstanding question regarding the degree to which Midwest-bred hazelnut accessions remain

closely related to *C. avellana*. 20[th] century Midwest breeding populations included interspecific

crosses with *C. avellana*, but the precise frequency of such hybridization events, and the fact that

they were often followed by several generations of open pollination within *C. americana*

dominant breeding orchards, has left this question unresolved. Fig. 4 shows clearly that the

majority of these Midwest accessions are indeed very closely related to wild *C. americana*

sourced from the Department of Natural Resources, as well as 'Rush' and 'Winkler' themselves.

A clear outlier in this respect is the variety Rose9-2, which very closely resembles other F_1

interspecific crosses.

Cross-genus level analysis

Transcriptomic analysis of the seven *Corylus* genomes included in this study reveals a consistent

distribution of genes across the 11 chromosomes. This analysis considers nearly half of the

recognized species in *Corylus* (Thompson et al., 1996). The observed consistency in gene

distribution suggests a relatively recent common ancestor of these hazelnut species. This finding

is important for understanding the evolutionary relationships between these species and could

have implications for hazelnut breeding and genetic improvement programs by providing a

foundation for identifying useful traits and genes that are present in all species.

Attempts at genetically characterizing the SSI S-locus in *Corylus* have included hybridization-

based staining of genes identified in *Brassica oleracea* L., which showed irregular hybridization

(Hampson et al., 1996). Similarly, gene expression studies identified only 61% homology

between *B. oleracea* alleles and S alleles in *Corylus* (Torello Marinoni et al., 2009). Finally,

transcriptomic analysis of the fine-mapped S-locus region in multiple *Corylus* species suggest

Corylus may harbor a novel SSI molecular mechanism that differs from *Brassica* (Hill et al.,

2021; Hou et al., 2022). The analysis presented here provides additional evidence for

independent evolution of a sporophytic self-incompatibility mechanism that is unique to *Corylus,*

and not shared by other members of the *Betulaceae*, nor model species of SSI such as *Brassica*

oleracea.

Conclusion

High-quality, annotated, chromosome-scale genome assemblies are essential tools for utilizing

modern genetic methods in the investigation of the molecular control of important traits, as well

as the application of advanced breeding methods. These two assemblies provide an important

resource which we hope will enable the application of such methods to the study and

improvement of both *C. americana*, as well as interspecific hybrids which expand the range over

which hazelnuts can be cultivated. Our analysis demonstrates highly conserved genome-wide

synteny across *Corylus* species. As such, genomic tools, analyses and resources developed in

one species may be more broadly useful to breeding programs in other species within the genus.

Data availability

Genome assemblies, as well as short- and long-read sequence data (from Arima Hi-C, Oxford

Nanopore, PacBio, and RNAseq libraries) are available through the NCBI Genome & Sequence

Read Archive, respectively, under BioProject ID: PRJNA939214 BioSample accessions:

SAMN33458638 ('Rush'), SAMN33458639 ('Winkler'). Genome assemblies, GFF3 models of

predicted genes, functionally annotated peptide sequences, anchored pangenome, and VCF of

GBS-derived SNPs are also available at https://zenodo.org/deposit/7439335. Bioinformatic

pipelines for genome assembly and polymorphism identification are published at:

https://github.com/shbrainard/Camericana_pipelines.

References

Alexander, D.H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry in unrelated individuals. *Genome Research*, *19*, 1655–1664. https://doi.org/10.1101/gr.094052.109

Alonge, M., Lebeigle, L., Kirsche, M., Jenike, K., Ou, S., Aganezov, S., Wang, X., Lippman, Z.B., Schatz, M.C., & Soyk, S. (2022). Automated assembly scaffolding using RagTag elevates a new tomato system for high-throughput genome editing. *Genome Biology*, *23*, 1–19. https://doi.org/10.1186/s13059-022-02823-7

Altschul, S.F., Gish, W., Miller, W., Myers, E.W., & Lipman, D.J. (1990). Basic local alignment search tool. *Journal of Molecular Biology*, *215*, 403–410. https://doi.org/10.1016/S0022-2836(05)80360-2

Benson, G. (1999). Tandem repeats finder: a program to analyze DNA sequences.. *Nucleic Acids Research*, *27*, 573–580

Bhattarai, G., Mehlenbacher, S.A., & Smith, D.C. (2017). Eastern filbert blight disease resistance from Corylus americana 'Rush' and selection 'Yoder #5' maps to linkage group 7. *Tree Genetics & Genomes*, *13*, 1–10. https://doi.org/10.1007/s11295-017-1129-9

BioBam Bioinformatics. (2019). OmicsBox – Bioinformatics Made Easy

Bolger, M., Schwacke, R., Gundlach, H., Schmutzer, T., Chen, J., Arend, D., Oppermann, M., Weise, S., Lange, M., Fiorani, F., Spannagl, M., Scholz, U., Mayer, K., & Usadel, B. (2017). From plant genomes to phenotypes. *Journal of Biotechnology*, *261*, 46–52. https://doi.org/10.1016/j.jbiotec.2017.06.003

Bradbury, P.J., Zhang, Z., Kroon, D.E., Casstevens, T.M., Ramdoss, Y., & Buckler, E.S. (2007). TASSEL: software for association mapping of complex traits in diverse samples. *Bioinformatics*, *23*, 2633–2635. https://doi.org/10.1093/bioinformatics/btm308

Brainard, S.H., Dawson, J.C., Fischbach, J.A., & Braun, L.C. (2023). Improving selection efficiency in C. americana × C. avellana interspecific hybrids through the development of an indel-based genetic map. *Acta Horticulturae*, *In press*. https://doi.org/10.1101/2023.02.05.527175

Ceylan, F.D., Adrar, N., Bolling, B.W., & Capanoglu, E. (2022). Valorisation of hazelnut by-products: current applications and future potential. *Biotechnology and Genetic Engineering Reviews*, 1–36. https://doi.org/10.1080/02648725.2022.2160920

Chan, P.P., & Lowe, T.M. (2019). tRNAscan-SE: Searching for tRNA genes in genomic
sequences. *Methods in molecular biology*, *1962*, 1–14. https://doi.org/10.1007/978-1-
4939-9173-0_1

Cheng, H., Concepcion, G.T., Feng, X., Zhang, H., & Li, H. (2021). Haplotype-resolved de novo
assembly using phased assembly graphs with hifiasm. *Nature Methods*, *18*, 170–175.
https://doi.org/10.1038/s41592-020-01056-5

Cheng, J., Han, Z., Xu, X., & Li, T. (2006). Isolation and identification of the pollen-expressed
polymorphic F-box genes linked to the S-locus in apple (*Malus × domestica*). *Sexual*
Plant Reproduction, *19*, 175–183. https://doi.org/10.1007/s00497-006-0034-4

Danecek, P., Bonfield, J.K., Liddle, J., Marshall, J., Ohan, V., Pollard, M.O., Whitwham, A.,
Keane, T., McCarthy, S.A., Davies, R.M., & Li, H. (2021). Twelve years of SAMtools
and BCFtools. *GigaScience*, *10*, giab008. https://doi.org/10.1093/gigascience/giab008

Demchik, M., Fischbach, J., Kern, A., Lane, J., McCown, B., Zeldin, E., & Turnquist, K. (2014).
Selection of American hazelnut as a potential oilseed crop. *Agroforestry Systems*, *88*,
449–459. https://doi.org/10.1007/s10457-014-9704-7

Di Lena, B., Curci, G., Vergni, L., & Farinelli, D. (2022). Climatic Suitability of Different Areas
in Abruzzo, Central Italy, for the Cultivation of Hazelnut. *Horticulturae*, *8*, 1–16.
https://doi.org/10.3390/horticulturae8070580

Dmitriev, A.A., Pushkova, E.N., & Melnikova, N.V. (2022). Plant Genome Sequencing: Modern
Technologies and Novel Opportunities for Breeding. *Molecular Biology*, *56*, 495–507.
https://doi.org/10.1134/S0026893322040045

Dudchenko, O., Batra, S.S., Omer, A.D., Nyquist, S.K., Hoeger, M., Durand, N.C., Shamim,
M.S., Machol, I., Lander, E.S., Aiden, A.P., & Aiden, E.L. (2017). De novo assembly of
the Aedes aegypti genome using Hi-C yields chromosome-length scaffolds. *Science*,
356, 92–95. https://doi.org/10.1126/science.aal3327

Durand, N.C., Shamim, M.S., Machol, I., Rao, S.S.P., Huntley, M.H., Lander, E.S., & Aiden,
E.L. (2016). Juicer Provides a One-Click System for Analyzing Loop-Resolution Hi-C
Experiments. *Cell Systems*, *3*, 95–98. https://doi.org/10.1016/j.cels.2016.07.002

Edgar, R.C. (2021). High-accuracy alignment ensembles enable unbiased assessments of
sequence homology and phylogeny. *bioRxiv*, 1–30.
https://doi.org/10.1101/2021.06.20.449169

Elshire, R.J., Glaubitz, J.C., Sun, Q., Poland, J.A., Kawamoto, K., Buckler, E.S., & Mitchell,
S.E. (2011). A Robust, Simple Genotyping-by-Sequencing (GBS) Approach for High
Diversity Species. *PLOS ONE*, *6*, e19379. https://doi.org/10.1371/journal.pone.0019379

Emms, D.M., & Kelly, S. (2019). OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome Biology*, *20*, 238. https://doi.org/10.1186/s13059-019-1832-y

Falush, D., Stephens, M., & Pritchard, J.K. (2007). Inference of population structure using
multilocus genotype data: dominant markers and null alleles: TECHNICAL ARTICLE.
Molecular Ecology Notes, *7*, 574–578. https://doi.org/10.1111/j.1471-
8286.2007.01758.x

Flynn, J.M., Hubley, R., Goubert, C., Rosen, J., Clark, A.G., Feschotte, C., & Smit, A.F. (2020).
RepeatModeler2 for automated genomic discovery of transposable element families.
Proceedings of the National Academy of Sciences, *117*, 9451–9457.
https://doi.org/10.1073/pnas.1921046117

Garg, S. (2021). Computational methods for chromosome-scale haplotype reconstruction.
Genome Biology, *22*, 1–24. https://doi.org/10.1186/s13059-021-02328-9

Ghurye, J., Rhie, A., Walenz, B.P., Schmitt, A., Selvaraj, S., Pop, M., Phillippy, A.M., & Koren,
S. (2019). Integrating Hi-C links with assembly graphs for chromosome-scale assembly.
PLOS Computational Biology, *15*, e1007273.
https://doi.org/10.1371/journal.pcbi.1007273

Gotz, S., Garcia-Gomez, J.M., Terol, J., Williams, T.D., Nagaraj, S.H., Nueda, M.J., Robles, M.,
Talon, M., Dopazo, J., & Conesa, A. (2008). High-throughput functional annotation and
data mining with the Blast2GO suite. *Nucleic Acids Research*, *36*, 3420–3435.
https://doi.org/10.1093/nar/gkn176

Granata, M.U., Bracco, F., & Catoni, R. (2020). Carbon dioxide sequestration capability of
hazelnut orchards: daily and seasonal trends. *Energy, Ecology and Environment*, *5*, 153–
160. https://doi.org/10.1007/s40974-020-00161-7

Haas, B.J., Delcher, A.L., Mount, S.M., Wortman, J.R., Smith, R.K., Hannick, L.I., Maiti, R.,
Ronning, C.M., Rusch, D.B., Town, C.D., Salzberg, S.L., & White, O. (2003). Improving
the Arabidopsis genome annotation using maximal transcript alignment assemblies.
Nucleic Acids Research, *31*, 5654–5666. https://doi.org/10.1093/nar/gkg770

Hampson, C.R., Coleman, G.D., & Azarenko, A.N. (1996). Does the genome of *Corylus
avellana* L. contain sequences homologous to the self-incompatibility gene of Brassica?.
Theoretical and Applied Genetics, *93*, 759–764. https://doi.org/10.1007/BF00224073

Hill, R.J., Baldassi, C., Snelling, J.W., Vining, K.J., & Mehlenbacher, S.A. (2021). Fine mapping
of the locus controlling self-incompatibility in European hazelnut. *Tree Genetics &
Genomes*, *17*, 6. https://doi.org/10.1007/s11295-020-01485-5

Hiscock, S.J., & McInnis, S.M. (2003). Pollen recognition and rejection during the sporophytic
self-incompatibility response: Brassica and beyond. *Trends in Plant Science*, *8*, 606–613.
https://doi.org/10.1016/j.tplants.2003.10.007

Hou, S., Zhao, T., Yang, Z., Liang, L., Ma, W., Wang, G., & Ma, Q. (2022). Stigmatic
Transcriptome Analysis of Self-Incompatible and Compatible Pollination in *Corylus
heterophylla* Fisch. × *Corylus avellana* L.. *Frontiers in Plant Science*, *13*.
https://doi.org/10.3389/fpls.2022.800768

Huerta-Cepas, J., Szklarczyk, D., Heller, D., Hernández-Plaza, A., Forslund, S.K., Cook, H., Mende, D.R., Letunic, I., Rattei, T., Jensen, L.J., von Mering, C., & Bork, P. (2019). eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids Research*, *47*, D309–D314. https://doi.org/10.1093/nar/gky1085

Huntley, R.P., Sawford, T., Mutowo-Meullenet, P., Shypitsyna, A., Bonilla, C., Martin, M.J., & O'Donovan, C. (2015). The GOA database: Gene Ontology annotation updates for 2015. *Nucleic Acids Research*, *43*, D1057–D1063. https://doi.org/10.1093/nar/gku1113

Hynynen, J., Niemistö, P., Viherä-Aarnio, A., Brunner, A., Hein, S., & Velling, P. (2010). Silviculture of birch (*Betula pendula* Roth and *Betula pubescens* Ehrh.) in northern Europe. *Forestry: An International Journal of Forest Research*, *83*, 103–119. https://doi.org/10.1093/forestry/cpp035

Jones, D.T., Taylor, W.R., & Thornton, J.M. (1992). The rapid generation of mutation data matrices from protein sequences. *Bioinformatics*, *8*, 275–282. https://doi.org/10.1093/bioinformatics/8.3.275

Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H., Maslen, J., Mitchell, A., Nuka, G., Pesseat, S., Quinn, A.F., Sangrador-Vegas, A., Scheremetjew, M., Yong, S.-Y., Lopez, R., & Hunter, S. (2014). InterProScan 5: genome-scale protein function classification. *Bioinformatics*, *30*, 1236–1240. https://doi.org/10.1093/bioinformatics/btu031

Kang, Y.J., Lee, T., Lee, J., Shim, S., Jeong, H., Satyawan, D., Kim, M.Y., & Lee, S.-H. (2016). Translational genomics for plant breeding with the genome sequence explosion. *Plant Biotechnology Journal*, *14*, 1057–1069. https://doi.org/10.1111/pbi.12449

Kitashiba, H., & Nasrallah, J.B. (2014). Self-incompatibility in Brassicaceae crops: lessons for interspecific incompatibility. *Breeding Science*, *64*, 23–37. https://doi.org/10.1270/jsbbs.64.23

Langdon, Q.K., Peris, D., Kyle, B., & Hittinger, C.T. (2018). sppIDer: A Species Identification Tool to Investigate Hybrid Genomes with High-Throughput Sequencing. *Molecular Biology and Evolution*, *35*, 2835–2849. https://doi.org/10.1093/molbev/msy166

Li, Y., Sun, P., Lu, Z., Chen, J., Wang, Z., Du, X., Zheng, Z., Wu, Y., Hu, H., Yang, J., Ma, J., Liu, J., & Yang, Y. (2021). The *Corylus mandshurica* genome provides insights into the evolution of *Betulaceae* genomes and hazelnut breeding. *Horticulture Research*, *8*, 1–13. https://doi.org/10.1038/s41438-021-00495-1

Liu, J., Wei, H., Zhang, X., He, H., Cheng, Y., & Wang, D. (2021). Chromosome-Level Genome Assembly and HazelOmics Database Construction Provides Insights Into Unsaturated Fatty Acid Synthesis and Cold Resistance in Hazelnut (*Corylus heterophylla*). *Frontiers in Plant Science*, *12*. https://doi.org/10.3389/fpls.2021.766548

Lovell, J.T., Sreedasyam, A., Schranz, M.E., Wilson, M., Carlson, J.W., Harkess, A., Emms, D.,
Goodstein, D.M., & Schmutz, J. (2022). GENESPACE tracks regions of interest and
gene copy number variation across multiple genomes. *eLife*, *11*, e78526.
https://doi.org/10.7554/eLife.78526

Lucas, S.J., Kahraman, K., Avşar, B., Buggs, R.J.A., & Bilge, I. (2021). A chromosome□scale
genome assembly of European hazel (*Corylus avellana* L.) reveals targets for crop
improvement. *The Plant Journal*, *105*, 1413–1430. https://doi.org/10.1111/tpj.15099

Manni, M., Berkeley, M.R., Seppey, M., Simão, F.A., & Zdobnov, E.M. (2021). BUSCO
Update: Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic
Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. *Molecular Biology*
and Evolution, *38*, 4647–4654. https://doi.org/10.1093/molbev/msab199

Mehlenbacher, S. (1991). Chilling requirements of hazelnut cultivars. *Scientia Horticulturae*,
47, 271–282

Mehlenbacher, S.A. (1997). Revised dominance hierarchy for S-alleles in *Corylus avellana* L.:.
Theoretical and Applied Genetics, *94*, 360–366. https://doi.org/10.1007/s001220050424

Mehlenbacher, S.A. (2014). Geographic Distribution of Incompatibility Alleles in Cultivars and
Selections of European Hazelnut. *Journal of the American Society for Horticultural*
Science, *139*, 191–212. https://doi.org/10.21273/JASHS.139.2.191

Mehlenbacher, S.A., & Thompson, M.M. (1988). Dominance relationships among S-alleles in
Corylus avellana L.. *Theoretical and Applied Genetics*, 669–672

Mikheenko, A., Prjibelski, A., Saveliev, V., Antipov, D., & Gurevich, A. (2018). Versatile
genome assembly evaluation with QUAST-LG. *Bioinformatics*, *34*, i142–i150.
https://doi.org/10.1093/bioinformatics/bty266

Minamikawa, M., Kakui, H., Wang, S., Kotoda, N., Kikuchi, S., Koba, T., & Sassa, H. (2010).
Apple S locus region represents a large cluster of related, polymorphic and pollen-
specific F-box genes. *Plant Molecular Biology*, *74*, 143–154.
https://doi.org/10.1007/s11103-010-9662-z

Molnar, T.J. (2011). Corylus. In C. Kole (Ed.), *Wild Crop Relatives: Genomic and Breeding*
Resources (pp. 15–48). Springer Berlin Heidelberg, Berlin, Heidelberg.

Molnar, T.J., Honig, J.A., Mayberry, A., Revord, R.S., Lovell, S.T., Mehlenbacher, S.A., &
Capik, J.M. (2018). Corylus americana: a valuable genetic resource for developing
hazelnuts adapted to the eastern United States. *Acta Horticulturae*, *1226*, 115–122.
https://doi.org/10.17660/ActaHortic.2018.1226.16

Pavese, V., Cavalet-Giorsa, E., Barchi, L., Acquadro, A., Torello Marinoni, D., Portis, E., Lucas,
S.J., & Botta, R. (2021). Whole-genome assembly of *Corylus avellana* cv "Tonda Gentile
delle Langhe" using linked-reads (10X Genomics). *G3 Genes|Genomes|Genetics*, *11*,
jkab152. https://doi.org/10.1093/g3journal/jkab152

Price, A.L., Patterson, N.J., Plenge, R.M., Weinblatt, M.E., Shadick, N.A., & Reich, D. (2006).
Principal components analysis corrects for stratification in genome-wide association
studies. *Nature Genetics*, *38*, 904–909. https://doi.org/10.1038/ng1847

R Core Team. (2021). *R: A Language and Environment for Statistical Computing*. R Foundation
for Statistical Computing, Vienna, Austria.

Revord, R.S., Lovell, S.T., Capik, J.M., Mehlenbacher, S.A., & Molnar, T.J. (2020). Eastern
Filbert Blight Resistance in American and Interspecific Hybrid Hazelnuts. *Journal of the*
American Society for Horticultural Science, *145*, 162–173.
https://doi.org/10.21273/JASHS04732-19

Roach, M.J., Schmidt, S.A., & Borneman, A.R. (2018). Purge Haplotigs: allelic contig
reassignment for third-gen diploid genome assemblies. *BMC Bioinformatics*, *19*.
https://doi.org/10.1186/s12859-018-2485-7

Rutter, P.A. (1987). Badgersett Research Farm – plantings, projects and goal. *Northern Nut*
Growers Annual Reprot, *78*, 173–186

Salamov, A.A., & Solovyev, V.V. (2000). Ab initio Gene Finding in *Drosophila* Genomic DNA.
Genome Research, *10*, 516–522

Sassa, H. (2016). Molecular mechanism of the S-RNase-based gametophytic self-incompatibility
in fruit trees of *Rosaceae*. *Breeding Science*, *66*, 116–121.
https://doi.org/10.1270/jsbbs.66.116

Semih Uzundumlu, A., Kurtoğlu, S., & Şerefoğlu, C. (2022). The role of Turkey in the world
hazelnut production and exporting. *Emirates Journal of Food and Agriculture*, *34*, 117–
127. https://doi.org/10.9755/ejfa.2022.v34.i1.2810

Sijacic, P., Wang, X., Skirpan, A.L., Wang, Y., Dowd, P.E., McCubbin, A.G., Huang, S., & Kao,
T. (2004). Identification of the pollen determinant of S-RNase-mediated self-
incompatibility. *Nature*, *429*, 302–305. https://doi.org/10.1038/nature02523

Silva, N.F., & Goring, D.R. (2001). Mechanisms of self-incompatibility in flowering plants:.
Cellular and Molecular Life Sciences, *58*, 1988–2007.
https://doi.org/10.1007/PL00000832

Slater, G.S.C., & Birney, E. (2005). Automated generation of heuristics for biological sequence
comparison. *BMC Bioinformatics*, *6*, 1–11. https://doi.org/10.1186/1471-2105-6-31

Stanke, M., Schöffmann, O., Morgenstern, B., & Waack, S. (2006). Gene prediction in
eukaryotes with a generalized hidden Markov model that uses hints from external
sources. *BMC Bioinformatics*, *7*, 1–11. https://doi.org/10.1186/1471-2105-7-62

Tamura, K., Stecher, G., & Kumar, S. (2021). MEGA11: Molecular Evolutionary Genetics
Analysis Version 11. *Molecular Biology and Evolution*, *38*, 3022–3027.
https://doi.org/10.1093/molbev/msab120

The UniProt Consortium. (2023). UniProt: the Universal Protein Knowledgebase in 2023.
Nucleic Acids Research, *51*, D523–D531. https://doi.org/10.1093/nar/gkac1052

Thompson, M., Lagerstedt, H., & Mehlenbacher, S. (1996). Hazelnuts. In J. Janick & J. Moore
(Eds.), *Fruit breeding: Nuts* (pp. 125–184). Wiley, New York, NY.

Torello Marinoni, D., Beltramo, C., Akkak, A., Destefanis, M.L., Boccacci, P., & Botta, R.
(2009). Gene Expression and Sporophytic Self-Incompatibility in Hazelnut. *Acta*
Horticulturae, *845*, 227–232. https://doi.org/10.17660/ActaHortic.2009.845.31

Vaillancourt, B., & Buell, C.R. (2019). High molecular weight DNA isolation method from
diverse plant species for use with Oxford Nanopore sequencing 783159.
https://doi.org/10.1101/783159

Vaser, R., Sović, I., Nagarajan, N., & Šikić, M. (2017). Fast and accurate de novo genome
assembly from long uncorrected reads. *Genome Research*, *27*, 737–746.
https://doi.org/10.1101/gr.214270.116

Wang, Y., Tang, H., DeBarry, J.D., Tan, X., Li, J., Wang, X., Lee, T. -h., Jin, H., Marler, B.,
Guo, H., Kissinger, J.C., & Paterson, A.H. (2012). MCScanX: a toolkit for detection and
evolutionary analysis of gene synteny and collinearity. *Nucleic Acids Research*, *40*, e49–
e49. https://doi.org/10.1093/nar/gkr1293

Weschcke, C. (1954). *Growing Nuts in the North: A Personal Story of the Author's Experience*
of 33 Years with Nut Culture in Minnesota and Wisconsin. Webb Publishing Company.

Wu, T.D., & Nacu, S. (2010). Fast and SNP-tolerant detection of complex variants and splicing
in short reads. *Bioinformatics*, *26*, 873–881.
https://doi.org/10.1093/bioinformatics/btq057

Zhao, T., Ma, W., Yang, Z., Liang, L., Chen, X., Wang, G., Ma, Q., & Wang, L. (2021). A
chromosome-level reference genome of the hazelnut, *Corylus heterophylla* Fisch.
GigaScience, *10*, giab027. https://doi.org/10.1093/gigascience/giab027

Figure 1. A: Clockwise from top left: the typical involucres, growth habit, and nuts of *C. americana* 'Rush', held by the NCGR in Corvallis, Oregon; **B:** Circular plot of genomic features of 'Winkler'. From outside in: ideogram of the 11 chromosomes of *C. americana*; total gene content, as identified by gene prediction and annotation; transposable elements, as identified by RepeatModeler; heatmap of tandem repeats; line graph of tRNA content. **C:** Contact map showing aligned and sorted Arima Hi-C data for 'Winkler' as visualized in Juicebox; green squares indicate ordered and oriented contigs, and blue squares indicate chromosomes. **D:** Whole genome comparison of the 'Rush' and 'Winkler' chromosome-scale assemblies. Upper plot: Histograms of insertion/deletions, repeat expansions/contractions, and tandem repeat/expansions

843   as visualized by Assemblytics; lower plot: Dot plot of delta file generated by nucmer. While
844   small indels are evident, largely consistent synteny is observed across all 11 chromosomes. A
845   single sizeable inversion on chromosome 9 is highlighted in red.
846

Figure 2. A: Riparian plot generated by GENESPACE. Single copy orthologs were used to generate and visually compare synteny
between 'Rush' and 'Winkler', the five other published *Corylus* genomes, *Betula pendula* (a close relative in the *Betulaceae*), and the
outgroup *Malus domestica*. In addition to illustrating consistent genome-wide inter-species synteny across the 11 chromosomes of
Corylus, this plot clearly illustrates the large-scale relationships between these 11 chromosomes, and the 14 chromosomes of *B.*
pendula. **B:** Barchart of the biological process term associated with the single-copy orthologs identified in *C. americana* which have
no orthologous copies in any of the five other *Corylus* genomes analyzed in GENESPACE, nor *B. pendula* or *M. domestica*. Barchart
of the biological process term associated with the single-copy orthologs identified in C. americana which have no orthologous copies
in any of the five other *Corylus* genomes analyzed in GENESPACE, nor *B. pendula* or *M. domestica*.

Figure 3. Pairwise dotplots visualizing pairwise BLAST hits between C americana 'Winkler', and each of the other eight genomes included in the GENESPACE analysis. Target and query genes which were identified as belonging to the same orthogroup are plotted, and color coded by sequence similarity. **A:** 'Winkler' vs. 'Winkler'; **B:** 'Winkler' vs. 'Rush'; **C:** 'Winkler' vs. *C. avellana* 'Tombul'; **D:** 'Winkler' vs. *C. avellana* Tonda Gentille delle Longhe; **E:** 'Winkler' vs. C. heterophylla (Beijing); **F:** 'Winkler' vs. *C. heterophylla* (Jilin); **G:** 'Winkler' vs. *C. mandshurica*; **H:** 'Winkler' vs. *B. pendula*; **I:** 'Winkler' vs. *M. domestica*.

Figure 4. PCA biplot of SNPs identified in *C. americana*, *C. avellana* and interspecific hybrids. On the left, wild *C. americana*
sourced from the DNR clusters together with 'Rush' and 'Winkler', along with a number of Midwestern varieties and an F_1 family
produced through a cross between two of them (light and dark green dots, respectively). In the middle, an outlier Midwest variety
(Rose9-2) exhibits PC1 scores similar to an F_1 between Eric4-21 (a Midwest variety that appears similar to *C. americana*) x Jefferson
(a *C. avellana* variety from Oregon State University). On the right are *C. avellana* varieties from Oregon State University and Rutgers
University, along with an interspecific F_1 population between a Rutgers cultivar and Rose9-2. This plot illustrates that while most
Midwestern varieties appear to be genetically quite similar to *C. americana*, there is substantial diversity among them (more so than
the limited number of *C. avellana* varieties included in the analysis) with some appearing quite similar to F_1 interspecific F_1 hybrids –
a finding supported by the historical record of interspecific hybridization in Midwestern breeding efforts.
879

Figure 5. Phylogenetic tree showing the evolution of genes comprising the self-incompatibility
locus in *Corylus* across the seven available *Corylus* genomes, with *B. pendula*, *M. domestica*,
and *B. oleracea* included as outgroups. Node labels represent bootstrap values based on 100
permutations. Color overlays illustrate evident clades, and highlight the fact that several lineages
of S locus genes are entirely specific to *Corylus*.

889