Fossil calibrations for molecular analyses and divergence time estimation for true crabs (Decapoda: Brachyura)

Javier Luque[1,2,3,*], Heather Bracken-Grissom[3,4], Javier Ortega-Hernández[2], and Joanna Wolfe[2]

[1]Museum of Zoology, Department of Zoology, University of Cambridge, Downing Street, Cambridge CB2 3EJ, UK.

[2]Museum of Comparative Zoology and Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, MA 02138, USA.

[3]Department of Biological Sciences, Florida International University-Biscayne Bay Campus, North Miami, FL 33181, USA.

[4]Department of Invertebrate Zoology, National Museum of Natural History, Smithsonian Institution, 10th and Constitution Ave NW, Washington, DC 20560, USA.

*Corresponding author: jl2351@cam.ac.uk; jluque@fas.harvard.edu

ABSTRACT

True crabs, or Brachyura, comprise over 7,300 known species and are among the most ecologically dominant, economically significant, and popularly recognized group of extant crustaceans. There are over 3,000 fossil brachyuran species known from mid and upper Jurassic, Cretaceous, and Cenozoic deposits across the globe, many of preserved in exquisite detail, but the origins and early evolution of true crabs remain unresolved. This uncertainty hinders the identification of the stratigraphically earliest occurrence of major brachyuran groups in the fossil record, obscuring our understanding of their phylogenetic relationships and thus the ability to estimate divergence times to answer large scale macroevolutionary questions. Here, we present 36 vetted fossil node calibration points for molecular phylogenetic analysis of crabs (one Anomura and 35 Brachyura) and reassess the earliest occurrences of several key clades based on recent fossil discoveries or re-examination of previous studies. For each calibrated node, we provide the minimum and tip maximum ages for the stratigraphically oldest fossil that can be reliably assigned to the group. Disentangling the anatomical disparity of fossil forms and their phylogenetic relationships is crucial to recognize the earliest branching members among brachyuran groups. This represents a critical first step understanding the evolution of carcinization and decarcinization in true crabs, the appearance of key adaptations, and the transition from sea to land and freshwater. The identification of reliable fossils for deep time calibrations, both as tips and nodes, is pivotal to ensure not only precise but more accurate divergence time estimations when reconstructing the crab tree of life.

Keywords: Cretaceous, crown group, evolution, freshwater, stem group, phylogeny.

INTRODUCTION

The origins and evolution of true crabs, or Brachyura, have sparked the fascination of scientists and the public alike in the last decades, thanks to their astonishing anatomical diversity (disparity) and multiple convergent instances of gain/loss of the crab-like body plan (i.e., carcinization and decarcinization) (Scholtz, 2014; Luque et al., 2019b; Wolfe et al., 2021). With over 7,600 extant species and more than 3,000 fossil species known (see Ng et al., 2008; Schweitzer et al., 2010; Luque et al., 2019b; Poore and Ahyong, 2023, and references therein), brachyurans are one of the most speciose groups of crustaceans. The monophyly of Brachyura is widely accepted, based on morphological and molecular grounds, and they are considered reciprocal sister groups with Anomura (collectively known as false crabs and allies), both forming a clade referred to as Meiura (Scholtz and Richter, 1995; Schram and Dixon, 2004; Hegna et al., 2020).

The systematics and classification of brachyurans have changed considerably in the last few decades, especially concerning family-level ranks and above. New lines of evidence have become available in the form of comprehensive phylogenetic studies, new fossil discoveries, and the critical re-interpretations of previous ones using novel technologies. Molecular and morphological phylogenetics are key to better understanding the evolutionary relationships through common ancestry among and within brachyurans, allowing us to quantitatively examine previous views and hypotheses largely based on traditional alpha taxonomy. This is particularly important when considering problematic groups such as the so-called podotremes and the heterotreme eubrachyurans, now widely recognized as paraphyletic groups, as well as the relationships among primary freshwater groups (e.g., Tan et al., 2018; Wang et al., 2018; Luque et al., 2019b; Ma et al., 2019; Tang et al., 2020; Wang et al., 2020; Luque et al., 2021; Tsang et al., 2022; Wolfe et al., 2022; Zhang et al., 2022; Poore and Ahyong, 2023).

While molecular and morphological phylogenies provide different types of evidence for understanding evolutionary relationships among organisms, fossils offer critical anatomical, spatial, and temporal information inaccessible from extant species alone (Luque et al., 2019b). Thus, accurate calibrations in a phylogenetic framework require reliable fossils, both in terms of their systematic affinities and their chronostratigraphic occurrences, since the use of equivocal (e.g., superficially similar but unrelated) or poorly constrained temporal and stratigraphic fossils may lead to precise yet inaccurate dating (either as tips or nodes), divergence time estimates, and therefore may confound interpretations of macroevolutionary patterns and processes over time.

The fossil record of brachyurans crabs consists mainly of marine taxa, which tend to be the

most abundant and often well preserved. Conversely, fossils of non-marine crabs (i.e., terrestrial,

semi-terrestrial, and freshwater) are remarkably scarce and often fragmentary, largely due to the

relatively dynamic and high-energy environments they inhabit, the geochemistry of the substrates,

and scavenging of their corpses or even re-working and consumption of their own exuviae (e.g.,

Locatelli, 2013; Luque, 2017; Luque et al., 2018). Such biases limit the number of remains than can

fossilize, often restricted to the most resilient biomineralized body parts (e.g., claw dactyli and

propodi remains), with whole organisms and soft to lightly-biomineralized tissues only preserved

under exceptional conditions (Luque et al., 2021). These biases constrain the number of fossils

suitable as reliable calibration points in molecular estimations of time divergence, and consequently

the available fossil data that can be used to examine the evolution of terrestrialization and the

transition from marine to non-marine habitats (Tsang et al., 2014; Luque et al., 2021; Watson-Zink,

2021; Tsang et al., 2022; Wolfe et al., 2022). Understanding the early origins of marine and non-

marine brachyurans by means of their fossil record requires not only new paleontological

discoveries, but also a critical re-examination of previous records. This holds true for marine crab

fossils as well.

Molecular biologists face limitations when selecting fossils for internal node calibrations

related to evaluating specific details of such fossil occurrences, such as their chronostratigraphic and

lithostratigraphic context, and the reliability of their systematic placement, especially with

fragmentary material (Gandolfo et al., 2008; Parham et al., 2012; Wolfe et al., 2016). Among crabs,

convergence is ubiquitous, and there is a trove of extinct and extant groups that superficially may

resemble each other, particularly in their dorsal carapaces. This may lead to inappropriate selection

of fossils that are not in fact related, and introduce large errors and inaccuracies into the estimation

of divergence dates when they are used to calibrate particular nodes and branches.

Fossils collected *in-situ* are often assigned a tentative relative age based on the

chronostratigraphic age of the geological formation that contains them. However, geological

formations can span tens to hundreds of meters in thickness and hundreds of thousands to millions

of years in age, whereas the age of a unitary fossil sample represents a single point in time, making

it difficult to constrain its age with respect to the age of the entire lithostratigraphic unit. For

example, if a formation is known to be 'Eocene' in age, but its base and top have not been better

constrained via absolute (radiometric isotope) or relative (e.g., biostratigraphy, stratigraphic

position) dating, it means that it could, hypothetically, range chronostratigraphically anywhere from

56.0 to 33.9 Ma. Even if the relative age of such formation is further constrained, for example to

'lower Eocene (Ypresian)', the age bracket would range from 56.0 to 47.8 Ma, in which case the

fossil would be assigned these as its tip maximum and minimum ages (e.g., several Italian fossil

occurrences, see below). This can be circumvented when other components of the fossil

assemblages shed light on the biostratigraphic ranges of other micro and/or macrofossils (e.g., the

porcellanid crab in calibration 1 here), when reliable radioisotopic data are associated to the same

strata as the fossils (e.g., the pseudothelphusid freshwater crab in calibration 16 herein), or if

magnetostratigraphic information is available (e.g., the parthenopid crab in calibration 23 herein).

An issue that is often oversimplified, and sometimes hard to circumvent, is the extrapolation

of the age range of a formation in a specific locality to other geographically separated localities

where the same formation occurs. While, in principle, the horizontality and superposition of beds in

a normal sequence would allow for regional correlations, extrapolating ages is not always

straightforward. This is particularly important for fossils near the lower and upper boundaries of a

lithostratigraphic unit, principally due to the irregular geometry of the basins, unconformities and

other erosional surfaces, eustasy, differential subsidence, diachronism, lateral facial changes, and

pinching of the strata, to name some. Thus, assigning the age of a formation in one locality to a

fossil found in the same formation in a different locality would be better informed by gaining some

level of understanding of the local and regional geological and stratigraphic context.

Lastly, unless the ages near the top and/or the bottom of a sequence are known, an absolute

age from a sample coming from a bed within the stratigraphic interval can inform about the overall

age range of the formation, but does not represent the age of the formation *per se*, and therefore the

age of the fossils present in the formation, unless the fossils come from or near the dated bed itself

(as in the case in calibration 16 herein). Nonetheless, an approximated age bracket for a fossil

occurring between two dated layers is as good of an approximation as possible, given the available

data, in which case the minimum and tip maximum ages of the fossil will be informed and

constrained by such absolute ages (e.g., as in the pinnotherid crab in calibration 10 herein).

Here we present a set of 36 phylogenetically vetted fossil calibration points for molecular

analyses and divergence time estimations of true crabs (one anomuran outgroup, 35 brachyuran

ingroups) (based on the crown group topologies inferred by Wolfe et al., 2022) (Table 1), and

provide information on the type of material investigated, its repository and catalogue number,

phylogenetic justification, minimum and tip maximum ages, age justification, and a discussion,

following best practices (Parham et al., 2012). For crown Meiura (Anomura + Brachyura)

calibrations 1–7 (Table 1), we provide a soft maximum age of 191.8 Ma based on the estimated

maximum age for the brachyuran fossil *Eocarcinus praecursor* Withers, 1932, from the Early

Jurassic (Pliensbachian) of England. Soft maximum age of 100.9 Ma for crown Eubrachyura

calibrations 8–19 and 21–36 is based on the estimated maximum age for the crown eubrachyuran

fossil *Cretapsara athanata* Luque, *in* Luque et al., 2021, from the mid-Cretaceous (uppermost

Albian–lowermost Cenomanian) of Burma (Table 1). Soft maximum age of 97.0 Ma for crown

Portunoidea calibration 20 is based on the estimated maximum age for the stem group portunoid

fossil *Eogeryon elegius* Ossó, 2016, from the mid-Cretaceous (Cenomanian) of Spain (Luque et al.,

2021).

For each vetted calibration, we reassess the earliest occurrences of several key groups in

light of recent fossil findings and re-examination of previous ones. In some cases, the selection of a

specific fossil calibration point from the literature does not necessarily represent the oldest record

that has been putatively assigned to a given family, but the oldest reliable occurrence that can be

referred to the clade with optimal confidence in its morphology. Minimum and tip maximum ages

were either obtained from the ages reported in the original publications, or from stratigraphically

constrained absolute or relative ages from the same localities as the fossil material or localities

nearby, whenever possible. Numerical ages for geological time intervals follow the International

Chronostratigraphic Chart by the International Commission on Stratigraphy (Cohen et al., 2013;

updated 2023).

The identification and critical examination of key fossils occurrences is crucial to

disentangle the anatomical disparity of crabs and their phylogenetic relationships, to recognize the

lower limits of Brachyura, the origins of carcinization and decarcinization, the transition from sea to

land and freshwater, and thus to ensure not only precise but more accurate time calibrations—both

as tips and nodes—and divergence time estimates when reconstructing the crab tree of life.

Table 1. Summary list of 36 vetted fossils for molecular divergence time estimations of brachyuran crabs (one

anomuran outgroup, 35 brachyuran ingroups). Ages in millions of years (Ma).

Point	Placement	Superfamily	Family	Genus & species	Tip max. age (Ma)	Min. age (Ma)	Oldest fossil occurrence reported in:
1	Crown	Galatheoidea	Porcellanidae	*Vibrissalana jurassica* Robins and Klompmaker, 2019	152.0	145.0	Robins and Klompmaker (2019)
2	Crown	Dromioidea	Dynomenidae	*Graptocarcinus muiri* Stenzel, 1944	113.0	100.5	Stenzel (1944)
3	Crown	Dromioidea	Dromiidae	*Dromiopsis gigas* Forir, 1887	100.5	93.9	Forir (1887)
4	Stem	Homoloidea	Homolidae	*Doerflesia ornata* Feldmann and Schweitzer, 2009	152.0	145.0	Feldmann and Schweitzer (2009)
5	Stem	Palaeocorystoidea	Palaeocorystidae	*Joeranina kerri* Luque, Feldmann, Schweitzer, Jaramillo, and Cameron 2012	118.0	113.0	Luque et al. (2012)
6	Stem	Raninoidea	Lyreididae	*Marylyreidus punctatus* (Rathbun, 1935) (as *Notopocorystes*)	104.0	100.5	Rathbun (1935)
7	Crown	Cyclodorippoidea	Cymonomidae	*Cymonomus primitivus* Müller and Collins, 1991	37.71	33.9	Müller and Collins (1991)
8	Crown	Potamoidea	Potamidae	*Alontecarcinus buratoi* De Angeli and Caporiondo, 2019	41.2	37.71	De Angeli and Caporiondo (2019)
9	Crown	Ocypodoidea	Ocypodidae	*Afruca miocenica* (Artal, 2008)	15.97	13.82	Artal (2008),
10	Crown	Pinnotheroidea	Pinnotheridae	*Pinnixa* sp.	21.68	18.7	Luque et al. (2017)
11	Crown	Grapsoidea	Percnidae	*Percnon santurbanensis* Ceccon and De Angeli, 2019	33.9	27.82	Ceccon and De Angeli (2019)
12	Crown	Grapsoidea	Grapsidae	*Pachygrapsus hungaricus* Müller, 1974	13.82	12.7	Müller (1974)
13	Crown	Grapsoidea	Varunidae	*Brachynotus corallinus* Beschin, Busulini, De Angeli, and Tessier, 2007	56.0	47.8	Beschin et al. (2007)
14	Crown	Grapsoidea	Gecarcinidae	*Cardisoma guanhumi* Latreille, *in* Latreille, Le Peletier, Serville and Guérin, 1828.	0.129	0.0117	Collins and Donovan (1997)
15	Crown	Grapsoidea	Sesarmidae	indet.	23.03	23.0	Serrano-Sánchez et al. (2016)
16	Crown	Pseudothelphusioidea	Pseudothelphusidae	indet.	18.18	17.93	Luque et al. (2019a)
17	Crown	Trichodactyloidea	Trichodactylidae	indet.	43.44	40.94	Klaus et al. (2017)
18	Crown	Majoidea	Epialtidae + Mithracidae	*Micippa antiqua* Beschin, De Angeli, and Checchi, 2001	33.9	27.82	Beschin et al. (2001)
19	Stem	Portunoidea	Eogeryonidae	*Eogeryon elegius* Ossó, 2016	97.0	93.9	Ossó (2016)

20	Crown	Portunoidea	Geryonidae	*Chaceon helmstedtense* (Bachmayer and Mundlos, 1968) (as *Coeloma*?)	33.9	27.82	Bachmayer and Mundlos (1968)
21	Crown	Portunoidea	Portunidae: Thalamitinae	*Lessinithalamita gioiae* De Angeli and Ceccon, 2015	53.3	47.8	De Angeli and Ceccon (2015)
22	Crown	Portunoidea	Polybiidae	*Liocarcinus heintzi* Schweitzer and Feldmann, 2010	33.9	27.82	Schweitzer and Feldmann (2010a)
23	Crown	Parthenopoidea	Parthenopidae + Dairoididae	*Aragolambrus collinsi* Ferratges, Zamora, and Aurell, 2019	57.1	53.02	Ferratges et al. (2019)
24	Crown	Calappoidea	Calappidae	*Calappa zinsmeisteri* Feldmann and Wilson, 1988	41.2	33.9	Feldmann and Wilson (1988)
25	Crown	Cancroidea	Cancridae	*Anatolikos undecimspinosus* Schweitzer et al. 2006	56.0	41.2	Schweitzer et al. (2006)
26	Crown	Dorippoidea	N/A	*Bartethusa hepatica* Quayle and Collins, 1981	41.2	37.71	Quayle and Collins (1981)
27	Crown	Leucosioidea	Leucosiidae	*Typilobus alponensis* Beschin, De Angeli, and Zorzin, 2009	56.0	47.8	Beschin et al. (2009)
28	Crown	Goneplacoidea	Euryplacidae	*Chirinocarcinus wichmanni* (Feldmann et al., 1995) (as *Glyphithyreus*)	66.0	61.6	Feldmann et al. (1995)
29	Crown	Eriphioidea	Eriphiidae	*Eriphia verrucosa* (Forskal, 1775) (as *Cancer*)	9.3	4.1	Betancort et al. (2014)
30	Crown	Trapezioidea	Trapeziidae	*Archaeotetra lessinea* De Angeli and Ceccon, 2013	53.3	47.8	De Angeli and Ceccon (2013)
31	Crown	Eriphioidea	Oziidae	*Ozius collinsi* Karasawa, 1992	15.97	13.82	Karasawa (1992)
32	Crown	Pilumnoidea	Pilumnidae	*Glabropilumnus trispinosus* Beschin, Busulini, and Tessier *in* Beschin et al., 2016	56.0	47.8	Beschin et al. (2016a)
33	Crown	Goneplacoidea	Goneplacidae	*Carcinoplax temikoensis* Feldmann and Maxwell, 1990	39.1	34.6	Feldmann and Maxwell (1990)
34	Crown	Xanthoidea	Pseudorhombilidae	*Pseudorhombila patagonica* Glaessner, 1933	23.03	5.333	Glaessner (1933)
35	Crown	Xanthoidea	Panopeidae	*Panopeus incisus* Beschin, Busulini, De Angeli, and Tessier, 2007	56.0	47.8	Beschin et al. (2007)
36	Crown	Xanthoidea	Xanthidae	*Phlyctenodes edwardsi* Beschin, Busulini, and Tessier *in* Beschin et al., 2016	56.0	47.8	Beschin et al. (2016a)

FOSSIL CALIBRATIONS

1. Anomura: Galatheoidea: Porcellanidae (crown)

Fossil specimen. Vibrissalana jurassica Robins and Klompmaker, 2019. Naturhistorisches Museum

Wien, Vienna, Austria. Holotype NHMW 2007z0149/0405, a somewhat complete dorsal carapace,

with a broken left side.

Phylogenetic justification. The overall carapace shape of the holotype and sole specimen of *V.*

jurassica fits well within the general diagnosis of Galatheoidea. The ovate, wide, and flattened

dorsal carapace, plus the limited dorsal ornamentation and groove pattern, match previous diagnoses

for Porcellanidae (Ahyong et al., 2010; Schweitzer and Feldmann, 2012; Robins and Klompmaker,

2019).

Minimum age. 145.0 Ma.

Tip maximum age. 152.0 Ma.

Soft maximum age. 191.8 Ma.

Age justification. This fossil was collected from the Ernstbrunn Limestone, exposed in the

Ernstbrunn Quarries, Ernstbrunn, Austria. Although the age of the Ernstbrunn Limestone, based on

microfossil and ammonite biostratigraphy, has been considered to range from the middle Tithonian

(Upper Jurassic) to the Berriasian (Lower Cretaceous) (e.g., Moshammer and Schlagintweit, 1999;

Zeiss, 2001; Schneider et al., 2013), the age of the rocks bearing meiurans has been consistently

treated as middle upper Tithonian (*Richterella richteri* Zone to *Micracanthoceras microcanthum*

Zone, *Simplisphinctes* Subzone) (e.g., Zeiss, 2001; Feldmann and Schweitzer, 2009; Schweitzer and

Feldmann, 2010b; Robins et al., 2013; Fraaije et al., 2019; Robins and Klompmaker, 2019), for

which a tip maximum age and a minimum age for *Vibrissalana jurassica* can be bracketed between

~152.0 and 145.0 Ma.

As described in Wolfe et al. (2019), a soft maximum age is obtained by phylogenetic

bracketing. The oldest crown Brachyura is debatable; *Eocarcinus praecursor* Withers, 1932, and

Eoprosopon klugi Förster, 1986, have both been proposed, but both lack some crown group

characters. Nevertheless, stem lineage positions of these taxa (Scholtz, 2020) allow a calibration of

crown group Meiura with a soft maximum from the base of the Pliensbachian, at 191.8 Ma.

Discussion. *Jurellana tithonia* Schweitzer and Feldmann, 2010b, also from the Ernstbrunn

Limestone, was previously considered the oldest porcellanid, although a recent revision of the taxon

has suggested an affinity with homolodromioid brachyurans (Robins and Klompmaker, 2019). As

such, *Vibrissalana jurassica* stands as the putatively oldest crown porcellanid (~152–145 Mya) and

represents, to date, a better calibration point for the crown group Porcellanidae.

2. Brachyura: Dromiacea: Dromioidea: Dynomenidae (crown)

Fossil specimen. *Graptocarcinus muiri* Stenzel, 1944. Bureau of Economic Geology, University of

Texas at Austin. Holotype BEG00021288.000, a large and complete dorsal carapace.

Phylogenetic justification. The overall ovate carapace outline, being wider than long, with discrete

but low-relief dorsal grooves, fits the diagnosis of Dynomenidae (Schweitzer et al., 2012; Van Bakel

et al., 2012a).

Minimum age. 100.5 Ma.

Tip maximum age. 113.0 Ma.

Soft maximum age. 191.8 Ma.

Age justification. This fossil was collected in the Taniniúl limestone, upper Lower Cretaceous

(Albian), from Choy Cave in Sierra del Abra between Las Palmas and Taninil, at kilometre 550 on

the railroad between Tampico and San Luis Potosí, State of San Luis Potosí, Mexico (Stenzel,

1944).

Soft maximum age as for calibration 1.

Discussion. Six species of the genus *Graptocarcinus* Roemer, 1887, ranging in age from the Lower

Cretaceous (Albian) to the middle Eocene (lower Lutetian) (Schweitzer et al., 2012; Beschin et al.,

2016b), are currently included within the subfamily Graptocarcininae Van Bakel, Guinot, Corral &

Artal, 2012a. Two isolated dactyli from the Barremian "Coulés boueuses", Serre de Bleyton, France,

have been adjudicated to *Graptocarcinus* (Hyzny and Kroh, 2015). Due the fragmentary nature of

the samples, we considered for calibration the earliest occurrence of fossils with more diagnostic

features that would warrant inclusion within crown Dynomenidae.

Several taxonomic studies have highlighted the issues of synonymy among some species

within the genus *Graptocarcinus*, for instance, *Graptocarcinus muiri*, from the Albian of San Luis

Potosí, Mexico, and *Graptocarcinus texanus* Roemer, 1887, from the Albian and Cenomanian of the

US and Europe, with some of them concluding that *G. miuri* is a junior synonym of *G. texanus* (e.g.,

Klompmaker, 2013; Kocova-Veselská et al., 2014, and references therein), whereas other studies

have maintained them as separate, valid species (e.g., Schweitzer et al., 2012; Van Bakel et al.,

2012a; Beschin et al., 2016b, and references therein). Since the earliest confirmed occurrences of

well-preserved fossils of both species are Albian in age, choosing one species over the other would

impact minimally the time calibration bracket for the genus (113–100 Mya).

3. Brachyura: Dromiacea: Dromioidea: Dromiidae (crown)

Fossil specimen. *Dromiopsis gigas* Forir, 1887. Mineral collections of the Université de Liège,

Belgium. Holotype 4936, a large, fragmented dorsal carapace in volume, right side.

Phylogenetic justification. *Dromiopsis* is an extinct genus with over a dozen species known to date

(Schweitzer et al., 2010). The overall circular carapace shape nearly as wide as long, the fronto-

orbital configuration, and the well-developed set of dorsal grooves delimiting the dorsal carapace

regions, guarantee placement within the crown Dromiidae (Schweitzer et al., 2012).

Minimum age. 93.9 Ma.

Tip maximum age. 100.5 Ma.

Soft maximum age. 191.8 Ma.

Age justification. The type material of *Dromiopsis gigas* comes from the Tourtias (Formation?),

Tournai, Belgium, which has been dated as Cenomanian based on ammonite biozonation (Kennedy

et al., 2011).

Discussion. *Costadromia hajzeri* Feldmann and Schweitzer, 2019, from the Campanian (Upper

Cretaceous, ~83–72 Mya) Wenonah Formation in New Jersey, USA, has been recently proposed as

the earliest known sponge crab, and included within Dromiidae sensu lato based mostly on details of

the frontal margin (Feldmann and Schweitzer, 2019). While *C. hajzeri* does seem to fit well within

the total group Dromiidae, the presence of *Dromiopsis gigas* in mid-Cretaceous (Cenomanian,

~100–93 Mya) rocks of Belgium, and its conspicuous similarity to other genera within crown

Dromiidae, indicate that the type material of *D. gigas* provides a better current calibration point for

crown Dromiidae.

Soft maximum age as for calibration 1.

4. Brachyura: Homoloida: Homoloidea: Homolidae (stem)

Fossil specimen. *Doerflesia ornata* Feldmann and Schweitzer, 2009. Naturhistorisches Museum

Wien (NHMW). Holotype 2007z0149/0015, a small dorsal carapace nearly complete.

Phylogenetic justification. The genus *Doerflesia* has several of the diagnostic features of the family

Homolidae, and it shows some resemblance to extant genera (Feldmann and Schweitzer, 2009;

Feldmann et al., 2012). While a homolid affinity is most likely, its position within the crown group

cannot be warranted with the available fossil material (a dorsal carapace of the holotype and sole

specimen), therefore we have chosen to treat it as a stem group member of Homolidae.

Minimum age. 145.0 Ma.

Tip maximum age. 152.0 Ma.

Soft maximum age. 191.8 Ma.

Age justification. This fossil was collected from the Ernstbrunn Limestone, exposed in the

Ernstbrunn Quarries, near the village of Dörfles, Austria.

Soft maximum age as for calibration 1.

5. Brachyura: Raninoida: Palaeocorystoidea: Palaeocorystidae (stem)

Fossil specimen. Joeranina kerri (Luque, Feldmann, Schweitzer, Jaramillo, and Cameron, 2012) (as

Necrocarcinus). Colombian Geological Survey (formerly INGEOMINAS), Museo Geológico José

Royo y Gómez, Bogotá DC, Colombia. Holotype IGM p881128, a nearly complete dorsal carapace

(Luque et al., 2012, pp. 411–413, fig. 4A, B.; Luque et al., 2017, p. 20, fig. 8B).

Phylogenetic justification. Recent phylogenetic studies have shown consistently that crabs of the

extinct superfamily Palaeocorystoidea occupy an intermediate place between the crab-like

Necrocarcinoidea and the frog-like Raninoidea (Karasawa et al., 2014; Luque, 2015a; Schweitzer et

al., 2016a; Luque et al., 2019b). As such, Palaeocorystoidea as a whole is recovered as part of the

stem group of Raninoidea, the latter of which includes all of the extant genera of frog crabs.

Minimum age. 113.0 Ma.

Tip maximum age. 118.0 Ma.

Soft maximum age. 191.8 Ma.

Age justification. The holotype of *Joeranina kerri* was discovered *in-situ* by one of us (J.L.) in grey

shales of the upper portion of the Lower Cretaceous Paja Formation, exposed along the road

between the towns of San Gil and Curití, in the Department of Santander, Colombia. The occurrence

of the gastropod *Turritella* (*Haustator*) *columbiana* Jaworski, 1938, and the ammonite

Acanthohoplites eleganteante Etayo-Serna, 1979, stratigraphically below the horizon yielding the

holotype of *J. kerri*, indicate an upper Aptian age in Colombia (see in Luque et al., 2012; Luque,

2014).

Soft maximum age as for calibration 1.

Discussion. Frog crabs and allies, together constituting the Section Raninoida (Ahyong et al., 2007),

are among the most anatomically disparate brachyuran groups (Schweitzer et al., 2012; Van Bakel et

al., 2012b; Karasawa et al., 2014; Hartzell et al., 2022). Recent phylogenetic studies have shown

that early-branching forms, referred to as the crab-like raninoidans (e.g., Orithopsidae,

Paranecrocarcinidae, Necrocarcinidae, Cenomanocarcinidae) are sister groups to a clade formed by

mostly decarcinized, frog-like raninoidan families such as the extinct Palaeocorystidae and the

extant Raninidae and allied relatives (Luque, 2015a; Schweitzer et al., 2016a; Luque et al., 2019b).

As such, the most recent common ancestor of Palaeocorystidae + Raninidae must be as old or older

than the oldest fossils known within Palaeocorystidae, which correspond to *Joeranina kerri* from the

upper Aptian (Lower Cretaceous, 118–113 Mya) of Colombia, South America, as indicated above.

330   6. Brachyura: Raninoida: Raninoidea: Lyreididae (stem)

Fossil specimen. *Marylyreidus punctatus* (Rathbun, 1935, as *Notopocorystes*). University of Texas.

Holotype Coll. No. 210, a nearly complete dorsal carapace in volume.

Phylogenetic justification. The type material of *Marylyreidus punctatus*, together with additional

specimens known from different localities and preserving dorsal, ventral, cuticular, and pleonal

details, strongly show a crown Raninoidea affinity, with a sternal configuration suggesting a close

proximity to lyreidids (Van Bakel et al., 2012b; Karasawa et al., 2014; Frantescu et al., 2016;

Schweitzer et al., 2018). This is consistent with the phylogenetic position of *Marylyreidus*,

recovered in a parsimony analysis as sister to *Bournelyreidus* van Bakel, Guinot, Artal, Fraaije, and

Jagt, 2012b, both forming an early diverging stem group sister to the remainder of the lyreidid-like

frog crabs (Karasawa et al., 2014).

Minimum age. 100.5 Ma.

Tip maximum age. 104.0 Ma.

Soft maximum age. 191.8 Ma.

Age justification. The type specimen of *Marylyreidus punctatus* was discovered in rocks of the

Lower Cretaceous (Albian), indicated by Rathbun (1935) as belonging to the 'Denton Clay, lower

Comanche Series, Washita Formation', cropping out in Grayson County, two miles north of

Denison, Texas, USA (Rathbun, 1935). Currently, the Washita 'Formation' in Texas is considered at

the Group level, and constituted at its thickest part by seven formations, in ascending order:

Kiamichi, Duck Creek, Fort Worth, Denton, Weno, Paw Paw, and Main Street formations (Scott et

al., 2002; Scott et al., 2016). The entire Washita Group from base to top is thought to have been

deposited between 104–94.4 Ma, with its lower boundary in the lowermost upper Albian, and with

the upper Albian–lower Cenomanian boundary (100.5 Ma) represented in the uppermost Main Street

Limestone (Scott et al., 2000; Scott et al., 2002). Denton Formation has been dated as upper Albian,

based on the occurrence of ammonites from the *Mortoniceras* (*Subschloenbachia*) *rostratum* Zone

(Kennedy et al., 2005). Additional specimens assigned to *M. punctatus* have been recovered from

several localities and stratigraphic intervals in Texas, including the slightly younger but also upper

Albian Paw Paw Formation (Frantescu et al., 2016). As such, relaxing the lowermost and uppermost

potential occurrences of *M. punctatus* within the Washita Group, its tip maximum and minimum

ages can be bracketed within 104.0–100.5 Ma.

Soft maximum age as for calibration 1.

Discussion. Wolfe et al. (2022) recovered a monophyletic Lyreididae nested within a paraphyletic

'Raninidae'. As such, despite *Marylyreidus* being part of the stem group to Lyreididae (or

Lyreininae, if confirmed to be a subfamily within Raninidae), it belongs to the crown Raninoidea.

Besides *M. punctatus*, only a couple other raninoideans are currently known from upper Albian

rocks worldwide, both initially assigned to the genus *Hemioon* Bell, 1863, traditionally considered a

Lyreididae. These are '*H. cunningtoni*' Bell, 1863, the type species of the genus and known from the

upper Albian of England, and '*Hemioon yanini*' Ilyin and Alekseev, 1998, known from the upper

Albian of Crimea. However, both records are problematic. Firstly, '*Hemioon*' has been a

problematic taxon due to the few specimens known and their poor preservation, resulting in some

authors considering it a valid taxon, whereas others have synonymized it with *Raninella* A. Milne-

Edwards, 1862a (see discussions in Bishop and Williams, 2000; Van Bakel et al., 2012b; Karasawa

et al., 2014; Schweitzer et al., 2018 and references therein). Secondly, *Raninella* '*cunningtoni*'

might be a juvenile of *Raninella elongata* Milne-Edwards, 1862b (see Bishop and Williams, 2000;

Van Bakel et al., 2012b; Karasawa et al., 2014 and references therein). Thirdly, *Raninella* is

currently considered a member of the subfamily Ranininae within the Raninidae, in which case this

family would also have an upper Albian record. Unfortunately, besides the conflicting views on

their systematic affinities, the exact age of 'H. cunningtoni' needs further revision (B. van Bakel,

pers. comm. to J.L., February 2023).

Regarding 'Hemioon yanini', its age is better constrained the species above discussed, being

referable to the upper Albian based on its occurrence in the Mortoniceras infiatum zone (Ilyin and

Alekseev, 1998; Ilyin, 2005), but its systematic placement is equally convoluted. Van Bakel et al.

(2012b) included this species within Raninella, whereas Karasawa et al. (2014) included it within

the genus Macroacaena Tucker, 1998, under the lyreidid subfamily Macroacaeninae Karasawa,

Schweitzer, Feldmann, and Luque, 2014 (Schweitzer et al., 2018). Regardless the generic and

specific affinities of 'H. cunningtoni' and 'H. yanini', the presence of disparate raninidoid forms in

the late Albian of USA, UK, and Crimea, indicate that the most recent common ancestor of

Raninodiea and all of its descendants must have originated before the late Albian.

7.   Brachyura: Cyclodorippoida: Cyclodorippoidea: Cymonomidae (crown)

Fossil specimen. Cymonomus primitivus Müller and Collins, 1991. Természcttudományi Múzeum,

Föld-és oslénytar, VIII Muzeum krt. 14-16, H-1088 Budapest, Hungary. Holotype EK-10.1 [M.91-

135], a partial dorsal carapace (Müller and Collins, 1991, p. 61, 63–64, pl. 3, fig. 6)

Phylogenetic justification. The Hungarian specimen can be referred to crown Cyclodorippoidea,

and assignable to Cymonomidae based on diagnostic characteristics of the subquadrate dorsal

carapace, the narrow fronto-orbital margin, and the configuration of the dorsal grooves and regions,

which fit well within the diagnosis of crown Cymonomidae (Müller and Collins, 1991; Tavares,

1993; Ahyong, 2019).

Minimum age. 33.9 Ma.

Tip maximum age. 37.71 Ma.

Soft maximum age. 191.8 Ma.

Age justification. Cymonomus primitivus was collected from 4–5 m thick coral-bearing limestones

of Facies 4 of the Upper Eocene (Priabonian) Szépvölgy Limestone Formation, cropping out at the

Ruprecht quarry, Budapest, Hungary, with the coral fauna indicating a Priabonian age (Müller and

Collins, 1991).

Soft maximum age as for calibration 1.

Discussion. Besides *Cymonomus primitivus*, other presumed cymonomids are known from the

Eocene, e.g., *Spathanomus felicianensis* De Angeli, 2016, and *Caporiondolus bericus* De Angeli,

2016, from the Priabonian of of the Orgiano quarry, Monti Berici, Italy, and *Eonomus californianus*

Nyborg, Garassino, and Slak, 2017, from the early to middle Eocene LLajas Formation, Simi

Valley, California, USA. We selected *C. primitivus* as our vetted calibration point for crown

Cymonomidae (~37–33 Ma), since *Cymonomus* is an extant genus, whereas *Spathanomus*,

Caporiondolus, and *Eonomus* are all extinct genera, and their family-level affinities are considered

by some authors as uncertain (e.g., Schweitzer et al., 2017).

8. Brachyura: Eubrachyura: Potamoidea: Potamidae (crown)

Fossil specimen. Alontecarcinus buratoi De Angeli and Caporiondo, 2019. Museo di Storia

Naturale di Verona. Holotype IGVR 19.38, a complete dorsal carapace.

Phylogenetic justification. The type material of *Alontecarcinus buratoi* shows remarkable

similarities with the dorsal carapaces of extant potamid crabs, including their oval outline and wide

carapace, the well-developed cervical and gastro-cardiac grooves, a front that is entire and directed

downwards, and the convex lateral margins that are smooth and bear a single epibranchial tooth (De

Angeli and Caporiondo, 2019).

Minimum age. 37.71 Ma.

Tip maximum age. 41.2 Ma.

Soft maximum age. 100.9 Ma.

Age justification. This fossil was collected from middle Eocene limestones of the Alonte quarry in
Berici Mounts, Vicenza, Northern Italy (De Angeli and Caporiondo, 2019). The stratigraphic
position of the Alonte quarry Limestones, overlying rocks of middle Eocene age, and underlying
rocks of Priabonian age, together with associations of gastropods and bivalves (e.g., *Ampullina*,
Cerithium, *Campanile*, *Natica*, *Corbis*, *Glycimeris*), echinoids (e.g., *Leiopedina*, *Sismondia*,
Echinolampas, *Schizaster*, *Cidaris*), and calcareous nannofossils, indicate a Bartonian age for the
Alonte quarry Limestones (41.2–37.71 Ma) (Beccaro, 2003; De Angeli and Alberti, 2016; De
Angeli and Caporiondo, 2019).
As this node is within Eubrachyura, a soft maximum age is based on *Cretapsara athanata*
Luque, *in* Luque et al., 2021, which is the oldest crown group Eubrachyura. The position of *C.*
athanata relative to modern eubrachyuran families is not clear, but its membership within the crown
group, based on morphological phylogenetic analysis (Luque et al., 2021), indicates that the
common ancestor of typical eubrachyuran forms must be older than the earliest Late Cretaceous.
The fossil was found in Burmese amber, for which the exact age estimates can vary, depending of
the source. The limited radioisotopic U-Pb information available for Kachin burmite indicates an
age of 98.8 ± 0.6 Ma (Shi et al., 2012), which would suggest a bracketed tip maximum age and
minimum age for *C. athanata* between 99.4 Ma and 98.2 Ma. As radiometric estimates pertain
primarily to the sediments and not necessarily the amber itself (see discussion in Luque et al., 2021).
Therefore, we calibrate a soft maximum close to the Albian–Cenomanian boundary 100.5 Ma ± 0.4
Myr = 100.9 Ma.

9. Brachyura: Eubrachyura: Ocypodoidea: Ocypodidae (crown)
Fossil specimens. *Afruca miocenica* (Artal, 2008, as *Uca*). Museo Geológico del Seminario de
Barcelona (MGSB), Catalonia, Spain. Holotype MGSB 68653, a complete and well-preserved
dorsal carapace, and paratypes 68654a to 68654e, including well-preserved dorsal and ventral males
and females with pleon, maxillipeds, pereopods, and the male major cheliped.

Phylogenetic justification. The type material of *Afruca miocenica* can be confidently assigned to the
genus *Afruca* based on the diagnostic enlarged and flattened dactyli and pollices of the male major
cheliped, and it is closest to the extant *Afruca tangeri* (Eydoux, 1835), which is the only recognized
species of fiddler crabs living today in the Iberic Peninsula and north Atlantic Africa. While *A.*

miocenica might indeed represent its own species, its overall similarities with *A. tangeri* have

invited the question of whether they are synonymous, with the former representing ontogenetic

variations similar to those seen in younger individuals of the later (de Gibert et al., 2013). In either

scenario, the close proximity between *A. miocenica* and *A. tangerii* is remarkable, and the

phylogenetic assignment of the fossils from Catalonia to the extant genus *Afruca*—and thus to the

crown Ocypodidae—are confirmed.

Minimum age. 13.82 Ma.

Tip maximum age. 15.97 Ma.

Soft maximum age. 100.9 Ma.

Age justification. The fossil material of *Afruca miocenica* was collected *in-situ* in terrigenous

yellow claystones from a lower Middle Miocene (Langhian, 15.97–13.82 Ma) locality of the Rubí

(Vallés), Province of Barcelona, Catalonia, Spain (Artal, 2008). These deposits are part of the

Vallès-Penedès basin, and represent marine to coastal mangrove facies deposited during the

maximum transgression of the Langhian in the area (Batllori and García, 1997; de Gibert and

Robles, 2005; Artal, 2008; Garassino et al., 2009), overlying continental Burdigalian facies

(Casanovas-Vilar et al., 2011), and underlying continental Serravalian facies (Batllori and García,

1997; de Gibert and Robles, 2005; Casanovas-Vilar et al., 2016).

Soft maximum age as for calibration 8.

Discussion. The fossil record of fiddler crabs is sparse and fragmentary, with only half a dozen

species recognized from fossilized carapaces (e.g., Artal, 2008; Domínguez Alonso, 2008; de Gibert

et al., 2013; Luque et al., 2017; Luque et al., 2018; Lima et al., 2020). Aside *Afruca miocenina*, the

only other Miocene fiddler crab known is *Uca maracoani* (Latreille, 1802) from Brazil (Brito, 1972;

Luque et al., 2017; Luque et al., 2018; Lima et al., 2020). The fossil material of *U. maracoani* was

collected in rocks of the Pirabas Formation, in the village of Baunilha Grande, Baia de Quatipuru,

Pará State, Brazil (Brito, 1972), and subsequent material from the area has been reported recently

(Lima et al., 2020). However, the exact age and stratigraphic provenance of these specimens is

unclear. Overall, palynological, lithological, petrographic, and geochemical data indicate that the

Pirabas Formation as a whole is constituted by two distinct depositional groups: the oldest and

lowermost one, of uppermost Lower Miocene age, deposited in a shallow marine environment, and

the youngest and uppermost one, of upper Middle Miocene to Upper Miocene, deposited in a coastal

environment near mangroves (e.g., Aguilera et al., 2022; Gomes et al., 2023). In localities such as

Capanema, where the lowermost depositional unit outcrops, absolute Sr/Sr dating obtained from

pectinid bivalve shells have yielded ages of 17.3–16.0 Ma, suggesting an uppermost Lower Miocene

(upper Burdigalian) age (Martinez et al., 2017; Gomes et al., 2023). In localities such as Praia do

Atalaia and Praia do Maçarico, where the uppermost depositional unit outcrops, palynomorphs of

the T16 biozone (Jaramillo et al., 2011) indicate an age of 12.7 to 7.1 Ma, suggesting an uppermost

Middle Miocene to Upper Miocene age (mid Serravalian to lowermost lower Messinian) (Gomes et

al., 2023). The fiddler crab-bearing facies of the Pirabas Formation at the 'Furo de Baunilha Grande'

locality correspond to the latter, younger unit, with a most probable age of uppermost Middle

Miocene to Upper Miocene, roughly corresponding to the Serravalian–Tortonian (13.82–7.246 Ma)

(Gomes et al., 2023). Moreover, these fiddler crab-bearing nodules are not *in-situ*, but found loose

on the Baunilla stream floor (Lima et al., 2020; Orangel Aguilera, pers. comm. to J.L., February

2023), and although previous works have presumably recovered palynological samples associated to

these nodules that would suggest a uppermost Lower Miocene rock age (e.g., Antonioli et al., 2015),

more recent and comprehensive biostratigraphic and paleonvironmental studies across localities

have confirmed the uppermost Middle Miocene to Upper Miocene age of the Baunilha Grande rocks

(Gomes et al., 2023). As such, we opt to use the apparently older and *in-situ* occurrence of *Afruca*

miocenica (Langhian, 15.97–13.82 Ma) as our calibration point for Ocypodidae.

Aside from the two fiddler crab species above mentioned, the only other Ocypodidae with

Miocene fossils known to date is *Ocypode vericoncava* Casadío et al., 2005, a single damaged

dorsal carapace from the upper Miocene (Tortonian? 11.63–7.246 Ma) of Argentina. Despite the

poor preservation of the holotype and sole specimen, the overall shape and fronto-orbital

configuration match those of *Ocypode*. Thus, this occurrence from Argentina, together with those of

Uca from Brazil and *Afruca* in Catalonia, indicate that Ocypodidae was already widespread by the

mid-Miocene, and that the most recent common ancestor of Ocypodidae and its divergence from

other thoracotremes must have occurred in the pre-Neogene, most likely in the Paleogene or even

the Cretaceous (Wolfe et al., 2022).

10. Brachyura: Eubrachyura: Pinnotheroidea: Pinnotheridae (crown)

Fossil specimen. *Pinnixa* sp. Invertebrate Paleontology collections, Florida Museum of Natural

History. Specimen UF 115397, a complete dorsal carapace (Luque et al., 2017, fig. 12Q).

Phylogenetic justification. Based on the overall carapace shape and dorsal patterns, this fossil

specimen can be assigned with confidence to the genus *Pinnixa*. Recent molecular studies, however,

have shown that *Pinnixa* species may not form a monophyletic genus, but be distributed among

three Pinnotheridae subfamilies (e.g., Palacios Theil et al., 2016). Despite this, they all can be

assigned to the crown-group Pinnotheridae, justifying the inclusion of the fossil *Pinnixa* from the

lower Miocene of Panama in the crown group.

Minimum age. 18.7 Ma.

Tip maximum age. 21.68 Ma.

Soft maximum age. 100.9 Ma.

Age justification. The specimen was collected *in-situ* by one of us (J.L.) in marine grey mudstones

from the middle part of the upper member of the Culebra Formation in the Panama Canal Zone,

Lirio section, Culebra Cut (formerly known as Gaillard Cut). The marine rocks of the Culebra

Formation are lower Miocene (uppermost Aquitanian–lower Burdigalian), and not Oligocene as

assumed by Rathbun, 1918 [1919] and repeated as such by subsequent authors (see notes in Luque

et al., 2017). At its base, the marine Culebra Formation overlies in sharp unconformity the terrestrial

upper Oligocene-lower Miocene rocks of the Las Cascadas Formation, and its top, it underlies

conformably the coarser terrestrial rocks of the upper Lower to Middle Miocene Cucaracha

Formation (Montes et al., 2012; Buchs et al., 2019; LeBlanc, 2021). Kirby et al. (2008) reported Sr

isotope dating ages for different parts of the formation, including four ages from *Acropora* corals

and pectinids in the middle member of the Culebra Formation (Emperador Limestone), ranging

between 21.24±0.44 and 20.62±0.58 Ma, and two ages of from a pectinid and an ostreid (19.83±0.39

and 19.12±0.42 Ma, respectively), from the lowermost portion of the Cucaracha Formation, above

the top of the upper member of the Culebra Formation (Kirby et al., 2008, fig. 6, table 2). Montes et

al. (2012) reported a U/Pb age of 19.3 ±0.4 Ma from magmatic zircons in an ash tuff from the lower

half of the upper member of the Culebra Formation, and MacFadden et al. (2014) reported U/Pb and

Ar/Ar ages of 18.81±0.3 and 18.96±0.9, respectively, from detrital zircons in an ash tuff from the

overlying Cucaracha Formation. As such, based on the currently available absolute ages available,

the age of the crab-bearing beds in the middle part of the upper member of the Culebra Formation—

and thus that of *Pinnixa* sp.—can be bracketed with confidence between 21.68 and 18.7 Ma, most

likely around 20–19 Ma (see Farris et al., 2017), in the lowermost Burdigalian, near the Aquitanian–

Burdigalian boundary.

Soft maximum age as for calibration 8.

Discussion. The fossil record of pinnotherids is among the most controversial among

eubrachyurans, in part because of their overall small sizes, their largely fragmentary fossil record,

and the overall carapace outline similarities of some fossils to some species of other eubrachyuran

(e.g., some hexapodids and aphanodactylids). This obscure the systematic affinities of several

fossils, and thus cast doubts of their usefulness for node calibrations of crown Pinnotheridae.

The extinct genus *Viapinnixa* Schweitzer and Feldmann, 2001, comprises four species that

range in age from the lower Paleocene to the lower Eocene (e.g., Philippi, 1887; Vega et al., 2001;

Vega et al., 2007; Vega et al., 2008; Armstrong et al., 2009), which are remarkably similar to

species within Hexapodidae. As such, the genus cannot be assigned with certainty to the

pinnotheroids (Luque et al., 2017), and even less so to the crown group Pinnotheridae. Other fossil

currently placed in the family Pinnotherinae include the spectacularly preserved extinct genus

Pharkidodes Feldmann, Schweitzer, Casadío & Griffin, Feldmann et al., 2011b, from the middle

Miocene of Tierra del Fuego, Argentina, which unfortunately cannot be placed within the crown

group with certainty.

Three fossil species have been dubiously assigned to *Pinnotheres*, i.e., *P.? araucana*

Philippi, 1887, from the 'Tertiary' of Chile, *P.? elatus* Milne-Edwards, 1873, from the upper

Miocene of France, and *P.? promaucanus* Philippi, 1887, from the Miocene of Chile (Schweitzer et

al., 2010). Over half a dozen species have been previously assigned to *Pinnixa*. '*Pinnixa*' *eocenica*

Rathbun, 1926, from the Eocene of Washington, USA, is a junior homonym of *Pinnixa eocenia*

Woods, 1922, and it is currently recognized as a hexapodid: *Palaeopinnixa rathbunae* Schweitzer,

Feldmann, Tucker & Berglund, 2000.

Pinnixa navidadensis Feldmann, Schweitzer and Encinas, 2005, from the middle Miocene of

Chile (Feldmann et al., 2010; Jagt et al., 2015; and *Pinnixa* sp., from the middle Pleistocene of

Florida (Agnew, 2001; Portell and Agnew, 2004), can be assigned to *Pinnixa*, but they are younger

than *Pinnixa* sp. from the lower Miocene Culebra Formation in Panama. *Pinnixa aequipunctata*

Morris and Collins, 1991, and *P. omega* Morris and Collins, 1991, come from the Pliocene Upper

Miri Formation, Brunei. *Pinnixa microgranulosa* Collins et al., 2003, from the Miocene Sandakan

Formation, Sarawak, may also belong to the hexapodid genus *Palaeopinnixa*.

As such, given the reliable systematic placement and the constrained stratigraphic age of

Pinnixa sp. from the lower Miocene Culebra Formation in the Panama Canal Zone (Luque et al.,

2017), this is to date the most reliable calibration datum for the crown group Pinnotheridae, while

the most recent common ancestor of all extant pinnotherids must have originated in the pre-

Neogene, most likely the Paleogene or Late Cretaceous (see Wolfe et al., 2022).

11. Brachyura: Eubrachyura: Grapsoidea: Percnidae (crown)

Fossil specimen. *Percnon santurbanensis* Ceccon and De Angeli, 2019, Museo Civico "D. Dal

Lago" di Valdagno (MCV), Vicenza, northern Italy. Holotype MCV.19/02, a complete dorsal

carapace (Ceccon and De Angeli, 2019, pp. 19–20, fig. 2.2).

Phylogenetic justification. Seven extant species in the genus *Percnon* constitute the monophyletic

family Percnidae Stevcic, 2005, which is nested within Grapsoidea (e.g., Schubart and Cuesta, 2010,

herein). The fossil species *P. santurbanensis* can be assigned to the genus *Percnon*, and thus is a

reliable calibration point for the family.

Minimum age. 27.82 Ma.

Tip maximum age. 33.9 Ma.

Soft maximum age. 100.9 Ma.

Age justification. The holotype was collected from coralline algae and corals of the lower Oligocene

(Rupelian, 33.90–27.82 Ma) of the Berici eastern Lessini mountains, tunnel of Sant'Urbano,

Montecchio Maggiore, Vicenza, northern Italy (Müller and Collins, 1991).

Soft maximum age as for calibration 8.

Discussion. To our knowledge, *P. santurbanensis* is the only known fossil brachyuran attributable

to Percnidae.

12. Brachyura: Eubrachyura: Grapsoidea: Grapsidae (crown)

Fossil specimen. *Pachygrapsus hungaricus* Müller, 1974. Hungarian Natural History Museum,

department of Earth Sciences and Palaeontology. Holotype HNHM 2004.163.1, a nearly complete

dorsal carapace.

Phylogenetic justification. Morphological and molecular phylomitogenomic studies indicate that

Pachygrapsus is phylogenetically well nested within crown Grapsidae, close to *Grapsus* (e.g.,

Karasawa and Kato, 2001; Chen et al., 2019; Lü et al., 2022; Tsang et al., 2022; Wolfe et al., 2022;

Zhang et al., 2022).

Minimum age. 12.6 Ma.

Tip maximum age. 13.82 Ma.

Soft maximum age. 100.9 Ma.

Age justification. The type material of *Pachygrapsus hungaricus* was collected from a patch reef

with *Porites* of the Leitha Limestone Formation, exposed at Tétény-plateau, western part of

the Budapest area, Hungary, dated as Middle Miocene (late Badenian, Decapod Zone 4 sensu

Müller, 1984) (Hyzny, 2016; Hyzny and Dulai, 2021). Several tens of additional specimens referred

to *P. hungaricus* have also been reported from coeval rocks in several localities in the Budapest area

(Gyakorló út, Örs vezér tere, and Rákos), as well as in its vicinity (i.e., Biatorbágy and Diósd) in

Hungary, and from coeval strata in Austria and Poland (Müller, 1984, 1996; Hyzny, 2016; Hyzny

and Dulai, 2021). Other conspecific occurrences are known from slightly older strata as well

(Decapod Zone 3 sensu Müller, 1984). Thus, *P. hungaricus* has been reliably documented from the

entire span of the upper Badenian (13.82–12.6 Ma) (Kovác et al., 2018; M. Hyzny, pers. comm. to

J.L., February 2023).

669   The age of the lower and upper boundaries of the Paratethyan Badenian stage/age have been

670 a matter of ongoing discussion. While some authors have considered that the age of the Badenian

671 lower boundary is positioned at the top of chron C5Cn.2n (16.303 Ma), roughly equivalent to the

672 uppermost part of the global Lower Miocene Burdigalian stage/age (e.g., Piller et al., 2007;

673 Hohenegger et al., 2014; Wagreich et al., 2014), others have considered it to be closer to the base of

674 the global Langhian stage/age (15.97 Ma) (Krijgsman and Piller, 2012; Reichenbacher et al., 2013;

675 Gozhyk et al., 2015; Kovác et al., 2018). Similarly, for several authors, the age of the Badenian

676 upper boundary is positioned at the top of polarity chron C5Ar.2n (12.829), roughly equivalent to

677 the middle part of the global Middle Miocene Serravallian stage/age boundary (e.g.Hohenegger et

678 al., 2014; Wagreich et al., 2014), whereas for others, the Badenian/Sarmatian boundary is slightly

679 younger (12.7–12.6 Ma) (e.g., Piller et al., 2007; Krijgsman and Piller, 2012; Kovác et al., 2018, see

680 also Hyzny and Dulai, 2021 and references therein). This reflects the issues of interpreting minimum

681 and tip maximum ages for taxa based on the literature alone, since the lithostratigraphic boundary

682 ages in different localities might vary in relation with eustatic sea-level changes and local

683 subsidence, and rarely a crab fossil occurrence is tied to absolute dating.

684   Since the goal of our calibration is not to resolve the exact stratigraphic age of a given

685 occurrence but to provide a confidence range that brackets the minimum and tip maximum ages, we

686 opt to consider the minimum age of *P. hungaricus* as 12.6 Ma, and the tip maximum age as 13.82

687 Ma, which roughly delimit the bottom and top of the Late Badenian (Kovác et al., 2018).

688   Soft maximum age as for calibration 8.

690 ***Discussion.*** Fossils of grapsid crabs are rare, in part due to the relatively dynamic and high-energy

691 environments they inhabit. Among grapsids, the genera *Goniopsis* and *Leptograpsus* (monotypic)

692 have no confirmed fossil record. The genera *Geograpsus* and *Grapsus* have fossils known from the

693 Holocene of Hawaii, i.e., *Geograpsus severnsi* Paulay and Starmer, 2011, and the Holocene of the

694 Pacific in Panama, i.e., *G.* aff. *G. grapsus* (Luque et al., 2018). The extant genus *Metopograpsus* has

695 two extinct species currently assigned to it, e.g., *M. badenis* Müller, 2006, from the same middle

696 Miocene (Badenian) locality in Hungary as *Pachygrapsus hungaricus*, and *M. traxleri* Müller, 1998,

697 from the lower Miocene (Karpatian) of Austria (Hyzny, 2016; Hyzny and Dulai, 2021). While any

698 of the fossil metopograpsid species could serve as potential good candidates as vetted calibration

699 points for crown Grapsidae, the fragmentary holotypes missing the fronto-orbital margins make their

700 systematic placement less confident than those fossils referred to *P. hungaricus*.

The extinct genus *Miograpsus* Fleming, 1981, and its sole species *M. papaka* Fleming, 1981,

is represented by a well-preserved female holotype and a ventrally exposed paratype from a "silty

sandstone pebble, not in place" (*ex-situ*) (Fleming, 1981, p. 105), presumably from the lower Hurupi

Formation, lower part of Tongaporutuan Stage (upper Miocene) of New Zealand. Although the

genus *Miograpsus* has been placed within Grapsidae by several authors (e.g., Feldmann, 1993;

Schweitzer et al., 2010), it seems to have closer affinities to genera within 'Varuninae' (now

considered its own family, Varunidae) (e.g., Fleming, 1981; Karasawa and Kato, 2001; Webber et

al., 2010). Since *Miograpsus* is 1) an extinct genus, 2) slightly younger that the fossil record of

crown genera such as *Pachygrapsus* and *Metopograpsus*, and especially 3) may have putative

varunid affinities, we refrain from using this taxon as calibration point for crown Grapsidae.

'*Nautilograpsoides*' *prior* Smirnov, 1929, a fossil species from the lower Miocene of the Caucasus,

is currently envisioned as a juvenile form of the portunid genus *Liocarcinus* (Hyzný et al., 2022, and

references therein).

Pachygrapsus, on the other hand, has a couple of fossil occurrences that can be assigned to

this genus. Fragmentary cheliped remains of *Pachygrapsus* sp. are known from the upper

Pleistocene of Jamaica (Morris, 1993; Luque et al., 2017), and *P. hungaricus* is known from several

tens of specimens from the middle Miocene (Badenian) of Europe (see above). Since *Pachygrapsus*

is an extant genus with a good Miocene fossil record, as represented by the type series and

numerous additional specimens of *P. hungaricus*, we selected the latter as our calibration point for

crown Grapsidae. The presence of *Pachygrapus* and *Metopograpsus* in the Miocene indicates that

the most recent common ancestor of crown Grapsidae and all of its descendants must have a pre-

Miocene origin, and likely rooted into the early Paleogene or Late Cretaceous.

13. Brachyura: Eubrachyura: Grapsoidea: Varunidae (crown)

Fossil specimen. *Brachynotus corallinus* Beschin, Busulini, De Angeli & Tessier, 2007. Museo di

Archeologia e Scienze Naturali "G. Zannato" di Montecchio Maggiore, Vicenza, northern Italy.

Holotype MCZ 1794, a nearly complete dorsal carapace.

Phylogenetic justification. The overall carapace morphology indicates that the holotype of

Brachynotus corallinus can be referred to crown Varunidae, which is fairly distantly related to

Grapsidae, the eponym of the polyphyletic superfamily Grapsoidea (Chen et al., 2019; Lü et al.,

2022; Tsang et al., 2022; Wolfe et al., 2022)

Minimum age. 47.8 Ma.

Tip maximum age. 56.0 Ma.

Soft maximum age. 100.9 Ma.

Age justification. This coral-associated fossil was collected from lower Eocene (Ypresian) marine

calcareous rocks, exposed at the Contrada Gecchelina of Monte di Malo in Vicenza, northern Italy.

The faunal association includes foraminifera, corals, calcareous algae, and molluscs. Foraminifera

like *Nummulites* cf. *partschi*, *N. tauricus*, and *N. nitidus*, indicate a middle–upper Ypresian age

(Beschin et al., 2007), and are stratigraphically correlated with the nearby 'Rossi' quarry (cava

'Rossi') (Beschin et al., 1998).

Discussion. Other Paleogene fossils of crown Varuninae include *Brachynotus oligocenicus* De

Angeli, Garassino and Ceccon, 2010b, a nicely preserved dorsal carapace from the early Oligocene

of Vicenza, Italy (De Angeli et al., 2010b), confirming the presence of the genus and the family

early in the Paleogene.

14. Brachyura: Eubrachyura: Grapsoidea: Gecarcinidae (crown)

Fossil specimen. Cardisoma guanhumi Latreille, *in* Latreille, Le Peletier, Serville and Guérin, 1828.

NCB – Naturalis, Leiden, The Netherlands, specimen NNM RGM 544 482, a large and mostly

complete right chela preserving the dactylus and pollex.

Phylogenetic justification. *Cardisoma guanhumi* is an extant gecarcinid species closely related to

Gecarcinus, the type genus of the family Gecarcinidae. The phylogenetic position of Gecarcinidae is

contentious with respect to other families within Grapsoidea. Recent nuclear and mitochondrial

phylogenetic studies have recovered a non-monophyletic Grapsoidea, with gecarcinids closer to taxa

within the families Sesarmidae and even Varunidae than to any taxon within Grapsidae (e.g., Chen

et al., 2019; Liu et al., 2021; Lü et al., 2022; Tsang et al., 2022; Wolfe et al., 2022).

Minimum age. 0.0117 Ma.

Tip maximum age. 0.129 Ma.

Soft maximum age. 100.9 Ma.

Age justification. This fossil was collected in the Port Morant Formation, parish of St. Thomas, east

Port Morant Harbour, southeast Jamaica (Collins and Donovan, 1997, 2012). The Port Morant

Formation, has been dated as Middle-Upper Pleistocene (132 ± 7 kyr to 125 ± 9 kyr, Sangamonian

Stage) based on electron spin resonance dating on samples of the coral species *Solenastrea bournoni*

and *Siderastrea radians* (Mitchell et al., 2000; Mitchell et al., 2001; James-Williamson and

Mitchell, 2012). Given the age uncertainty of the fossil specimens from the Port Morant Formation

with respect to the coral samples dated, we bracketed the soft maximum and minimum ages for the

fossil occurrences of *C. guanhimu* between the base and the top of the Late Pleistocene (0.129–

0.0117 Ma), although we expect to family as a whole to be much older than Pleistocene, most likely

originating in the Paleogene or possibly late Cretaceous (see in Wolfe et al., 2022).

Soft maximum age as for calibration 8.

Discussion. The fossil record of gecarcinid land crabs is sparse and fragmentary, and largely

represented by isolated claw remains (e.g., Luque, 2017). To date, all of the known fossils

assignable to *Cardisoma* are Quaternary in age, i.e., late Pleistocene (Collins and Donovan, 1997;

Luque, 2017) or late Holocene (Luque, 2017; Luque et al., 2017; Luque et al., 2018), making the

record from the upper Pleistocene of Jamaica the oldest fossil material referable to Gecarcinidae

with confidence.

15. Brachyura: Eubrachyura: Grapsoidea: Sesarmidae (crown)

Fossil specimens. Sesarmidae gen. et sp. indet., in Serrano-Sánchez et al. (2016). Museo de

Paleontología "Eliseo Palacios Aguilera" (Secretaría de Medio Ambiente e Historia Natural),

Calzada de los Hombres Ilustres s/n, Tuxtla Gutiérrez, Chiapas, Mexico, specimens with acronym

IHNFG (Instituto de Historia Natural, Fósil Geográfico) IHNFG-4969, IHNFG-4970, IHNFG-4991,

IHNFG-4992, and IHNFG-5555. A handful of specimens preserved in amber, mostly represented by

complete dorsal and ventral carapaces with limbs attached (Serrano-Sánchez et al., 2016, figs 3, 4.1,

4.3–47; Luque et al., 2017, fig. 13L).

Phylogenetic justification. Despite the unclear generic and specific affinities of the fossil grapsoids

in amber from the Miocene of Chiapas, Mexico, the fossils can be assigned with confidence to

crown Sesarmidae based on the overall morphology of the carapace and limbs (Serrano-Sánchez et

al., 2016; Luque et al., 2017).

Minimum age. 23.0 Ma.

Tip maximum age. 23.03 Ma.

Soft maximum age. 100.9 Ma.

Age justification. These fossils were collected in rocks of the upper La Quinta Formation (Finca

Carmitto Member) in Chiapas, México, dated as Lower Miocene (Aquitanian, 22.8 Ma, in Serrano-

Sánchez et al., 2015) based on the biostratigraphy of corals, molluscs, microfossils, and Strontium

isotopes (Serrano-Sánchez et al., 2015). Bernot et al., 2022 highlighted that the absence of

radiometric ages for La Quinta Formation precludes a more precise constrain of the possible age of

the crab-bearing amber samples, and propose some tentative minimum ages based on previously

reported benthic foraminiferal data from the overlying Mazantic Shale (Solórzano Kraemer, 2007),

while indicating a $^{87}Sr/^{86}Sr$ radiometric date of 23.0 Ma, roughly at the Oligocene–Miocene

boundary, reported in Vega et al. (2009) (see Bernot et al. (2022) supplementary materials and

reference therein). As such, given the uncertainty on the age of the crab-yielding amber deposits, we

constrain the minimum age to the date reported by Serrano-Sánchez et al. (2015), and the tip

maximum age to the base of the Aquitanian and thus the Miocene (23.03 Ma).

Soft maximum age as for calibration 8.

Discussion. Sesarmid crabs trapped in amber are known from the lower Miocene (Aquitanian) of

Simojovel, Chiapas, Mexico (Grimaldi, 1996; Boucot and Poinar Jr, 2010; Serrano-Sánchez et al.,

2016; Luque et al., 2017). '*Sesarma*' *paraensis* Beurlen, 1958, presumably a fossil sesarmid from

the Miocene Pirabas Formation of Pará, Brazil, cannot be confirmed to be a sesarmid as the

specimen is not illustrated in the original paper by Beurlen (1958), but only a schematic line

drawing reconstruction (Beurlen, 1958, pl. IV, fig. 4). As such, as it stands today, this record cannot

be reliably confirmed as a sesarmid. The only other sesarmid fossils known are cheliped remains

from the upper Pleistocene of Jamaica previously described as a new species, *Sesarma primigenium*

Collins, Mitchell and Donovan, 2009, but now considered a junior synonym of the extant species

Sesarma cookei Hartnoll, 1971, to which the fossilized cheliped remains above mentioned belong

(see discussion in Luque et al., 2017, p. 68, note 3).

16. Brachyura: Eubrachyura: Pseudothelphusoidea: Pseudothelphusidae (crown)

Fossil specimen. Pseudothelphusidae gen. & sp. indet. (Luque et al., 2019a). Invertebrate

Paleontology collections, Florida Museum of Natural History. Specimen UF XXXXX, a complete

dorsal and ventral carapace, chelipeds, eyes, and proximal podomeres of pereopods, in volume.

Phylogenetic justification. The detailed preservation of diagnostic features in the fossil specimen,

together with its stratigraphic and paleoenviromental context, e.g., continental leaf-rich level of the

Pedro Miguel Formation, allow its placement with certainty within the freshwater group

Pseutothelphusidae, and thus as part of the crown Pseudothelphusoidea (Luque et al., 2019a; Luque

et al., under study).

Minimum age. 17.93Ma.

Tip maximum age. 18.18 Ma.

Soft maximum age. 100.9 Ma.

Age justification. This fossil pseudothelphusid specimen comes *in-situ* from a leaf-rich level of

tuffaceous mudstones of the Pedro Miguel Formation, Panama Canal, Panama. U/Pb absolute ages

858 from detrital zircons collected *in-situ* in two tuffaceous conglomerates that bracket a leaf-rich

859 horizon reported by Londoño et al. (2018) from which the fossil pseudothelphusid comes, yielded an

860 estimated depositional age of 18.01±0.17 Ma. Previous Ar/Ar dating from the Pedro Miguel

861 Formation in other parts of the Canal by Wegner et al. (2011) has suggested an age of 18.4 ± 1.07

862 Ma, and while details of the exact provenance of such samples within the formation are unclear

863 (Montes et al., 2012; Farris et al., 2017), a corrected datum by MacFadden et al. (2014) suggests an

864 age of 18.90±0.59 Ma.

865  Since the tuffaceous mudstone layer yielding the fossil leaves of Londoño et al. (2018) and

866 the *in-situ* pseudothelphusid are bracketed by the same lower and upper tuffaceous conglomerates,

867 both dated as 18.01±0.17 Ma, we assign the fossil a tip maximum age of 18.18 Ma, and a minimum

868 age of 17.93 Ma, which are closer to the middle Burdigalian (late Early Miocene). As this fossil is

869 clearly part of the crown Pseudothelphusidae (Luque et al., 2019a), the most recent common

870 ancestor of all pseudothelphusids must have originated in the pre-Burdigalian, most likely the

871 Paleogene.

872  Soft maximum age as for calibration 8.

874 ***Discussion.*** There are only two extant superfamilies of freshwater crabs in the Neotropics:

875 Trichodactylidae and Pseudothelphusidae. While the fossil record of Trichodactylidae is represented

876 by numerous cheliped fragments (see below), the fossil record of Pseudothelphusidae is otherwise

877 unknown. The new fossil represents not only the first and thus oldest know fossil of the family, but

878 one of the most complete fossil freshwater crabs known to date (Luque et al., 2019a; Luque et al.,

879 under study).

882 **17. Brachyura: Eubrachyura: Trichodactyloidea: Trichodactylidae (crown)**

883 ***Fossil specimen.*** Trichodactylidae genus and species indet. Museo de Historia Natural de la

884 Universidad Nacional Mayor San Marcos de Lima (MUSM), Peru. Three claw fragments: CTA 47

885 (1 specimen), and CTA 66 (two specimens), representing a small pollex (CTA 47), and a very small

886 dactylus and a small pollex (CTA 66) (Klaus et al., 2017, fig. 3D–E).

888 ***Phylogenetic justification.*** The preservation of diagnostic features on the fossilized dactyli and

889 propodi and the teeth and interteeth on their cutting edges, together with their stratigraphic and

continental limnic paleoenviromental contexts, allow their placement within the family

Trichodactylidae (Klaus et al., 2017; Luque et al., 2017).

Minimum age. 40.94 Ma.

Tip maximum age. 43.44 Ma.

Soft maximum age. 100.9 Ma.

Age justification. These fossils were collected in the lower member of the Pozo Formation, middle

and upper middle Eocene (lower Barrancan, >41.6–40.94 Ma, 43.44 ± 2.5 Ma), Contamana area,

Loreto, Peru. The ages are based on mammalian biochronology and $^{40}Ar/^{39}Ar$ radiometric age

(Antoine et al., 2016; Klaus et al., 2017). The South American land mammal age (SALMA) known

as Barrancan is largely equivalent to the uppermost global Lutetian and most of the Bartonian.

Soft maximum age as for calibration 8.

18. Brachyura: Eubrachyura: Epialtidae + Mithracidae (crown)

Fossil specimen. **'***Micippa***'** *antiqua* Beschin, De Angeli, and Checchi, 2001. Museo Civico "G.

Zannato" di Montecchio Maggiore, Vicenza, northern Italy. Holotype I.G. 286477, a complete

dorsal carapace.

Phylogenetic justification. *Micippa* Leach, 1817, is an extant genus that has been recovered by

several authors as not clustering together with other mithracids (see Windsor and Felder (2014) and

references therein). Windsor and Felder (2014) included *Micippa* and *Stenocionops* Desmarest,

1823, within the family Mithracidae *sensu lato*, and Klompmaker et al. (2015) also included them

within this family in their revision of the fossil record of Mithracidae (see discussion below).

Alternatively, Poore and Ahyong (2023) include *Micippa* within Epialtidae. As such, we consider

'*Micippa***'** *antiqua* as a calibration point for Epialtidae + Mithracidae (crown) (see in Wolfe et al.,

2022). The fossil record of pre-Cenozoic majoid crabs is one of the most problematic for

brachyurans due to the anatomical disparity seen across taxa and its often fragmentary nature, and it

is in need of future thorough revisions.

Minimum age. 27.82 Ma.

Tip maximum age. 33.9 Ma.

Soft maximum age. 100.9 Ma.

Age justification. The holotype of '*Micippa*' *antiqua* was collected associated to shallow-water

corals-bearing limestones from the Castelgomberto Formation ("Formazione di castelgomberto") in

Vicenza, northerm Italy (Beschin et al., 2001). Large foraminifera such as *Nummulites fichteli, N.*

vascus, Operculina complanata, Praerhapydionina delicata, Spirolina cylindracea, Peneroplis

glynnjonesi, and *Asterigerina rotula haeringensis* in the Castelgomberto Formation are indicative of

the lower Oligocene (Rupelian, 33.9–27.82 Ma) biozones SB21–22A (Ungaro, 1978; Cahuzac and

Poignant, 1997; Nebelsick et al., 2013).

Soft maximum age as for calibration 8.

Discussion. Among the extant mithracid genera with a known fossil record, *Micippa* has the oldest

putatively confirmed occurrences, as represented by '*M.*' *antiqua* from the lower Oligocene of Italy.

Dorsal carapaces of two other micippid species, i.e., *M. annamariae* Gatt and De Angeli, 2010, and

M. hungarica (Lorenthey in Lorenthey & Beurlen, 1929), are known from the middle Miocene of

Malta and the middle–upper Miocene of Poland, Hungary, and Austria, respectively. This, together

with some Miocene and several Pliocene and Pleistocene fossils referable to Mithracidae s.s. from

the Caribbean (Klompmaker et al., 2015; Luque et al., 2017), indicate that mithracids were already

present and widespread during the late Paleogene–early Neogene.

'*Stenocionops*' *suwanneeana* Rathbun, 1935, from the upper Eocene fossil of Florida, USA,

is a propodus that cannot be placed with confidence within any extant mithracid genus.

"*Stenocionops*" *primus* Rathbun, 1935, presumably from the Upper Cretaceous (Santonian?) of

Arkansas, USA, is a fragmented cheliped manus of unclear affinities. *Antarctomithrax thomsoni*

Feldmann, 1994, from the Eocene of Antarctica, is an extinct monotypic genus described based on a

partially preserved dorsal carapace that cannot be ascribed to Mithracidae with confidence.

19. Brachyura: Eubrachyura: Portunoidea: Eogeryonidae (stem)

Fossil specimen. *Eogeryon elegius* Ossó, 2016. Museu de Geologia de Barcelona, holotype MGB

69151, a complete dorsal and ventral carapace in volume with cheliped associated.

Phylogenetic justification. In a recent morphological phylogenetic study incorporating fossil and

extant brachyurans, Eogeryonidae has been recovered as a stem Portunoidea, sister to a clade

formed by the extant families Carcinidae, Geryonidae, and Portunidae, and all of them together

forming the total group Portunoidea (Luque et al., 2021).

Minimum age. 93.9 Ma.

Tip maximum age. 97.0 Ma.

Soft maximum age. 100.9 Ma.

Age justification. The type material of *Eogeryon elegius* comes from sandy limestones of the Villa

de Vés Formation, exposed near the village of Condemios de Arriba, Northern Guadalajara

Province, Spain. The presence of the ammonite *Vascoceras gamai* Choffat, 1898, in the Villa de Vés

Formation represents the upper part of the upper Cenomanian (lowermost Upper Cretaceous, 95–93

Ma), and it is found below the *Spathites* (*Jeanrogericeras*) *subconciliatus* Zone, which marks the

end of Cenomanian (Ossó, 2016).

Soft maximum age as for calibration 8.

Discussion. The anatomical similarities of *Eogeryon* to portunoids such as *Geryon* suggest a

potential proximity of Eogeryonidae to Geryonidae (Ossó, 2016). Several small dorsal carapaces of

the eubrachyuran *Romualdocarcinus salesi* Prado and Luque *in* Prado et al., 2018, from the upper

Lower Cretaceous (Aptian–Albian, ~115–110 Ma) of Brazil, share some diagnostic features with

Eogeryon elegius that suggest a plausible eogeryonid affinity, but due to their incompleteness,

especially regarding the lack of ventral and cheliped material associated, a more precise

phylogenetic placement of *R. salesi* is not possible at this time (Prado et al., 2018). *Eogeryon*,

Romualdocarcinus, and other modern-looking crown eubrachyuran fossils, also share several

general features with the remarkably well-preserved *Cretapsara athanata* Luque *in* Luque, Xing et

al., 2021, from the lowermost Cenomanian (~99 Ma) of Burma (Myanmar), but *Cretapsara* differs

from them in having a bilobate rostrum and lacking orbital fissures (Luque et al., 2021).

20. Brachyura: Eubrachyura: Portunoidea: Geryonidae (crown)

Fossil specimen. *Chaceon helmstedtense* (Bachmayer and Mundlos, 1968, as *Coeloma* ?). Mundlos

collection, geological-palaeontological collections of the Natural History Museum at Vienna,

Austria. Holotype 1968/773/2, a complete dorsal carapace in volume with chelipeds and legs

preserved.

Phylogenetic justification. The fossil species *Chaceon helmstedtense* can be assigned to the extant

genus *Chaceon* Manning and Holthuis, 1989, and thus the family Geryonidae, based on diagnostic

features of the dorsal carapace outline, the orbitofrontal and anterolateral margins, the thoracic

sternum, and the chelipeds. Molecular and morphological phylogenetic studies have recovered

Chaceon and Geryonidae as nested within crown Portunoidea (e.g., Karasawa et al., 2008;

Spiridonov et al., 2014; Evans, 2018; Luque et al., 2021; Wolfe et al., 2022).

1002

1003 ***Minimum age***. 27.82 Ma.

1004

1005 ***Tip maximum age.*** 33.9 Ma.

1006

1007 ***Soft maximum age.*** 97.0 Ma.

1008

Age justification. The type locality of *Chaceon helmstedtense* is a marl/clay horizon (Mergel/Ton-

Horizont, crab zone K I) in Helmstedt near Braunschweig, Lower Saxony, Northwestern Germany,

considered lower Oligocene in age (Rupelian, 33.9–27.82 Ma). Associated molluscan fauna in the

marl/clay beds includes "*Pecten corneus, Isocardia multicostata, Cardita* latesulcata, *Ostrea*

ventilabrum, and *Ostrea queteleti*" (Bachmayer and Mundlos, 1968). As crown group members of

Portunoidea cannot be older than their common ancestor shared with the stem group fossil *Eogeryon*

(Ossó, 2016), a soft maximum age is taken from the maximum uncertainty on the age of the Villa de

Vés Formation, at 97.0 Ma.

1017

Discussion. Besides *Chaceon helmstedtense*, another occurrence of *Chaceon* in the Paleogene is *C.*

heimertingensis (Bachmayer and Wagner, 1957), from the uppermost Oligocene (Chattian) of

Austria. *Chaceon peruvianus* (D'Orbigny, 1842, as *Portunus*) an iconic Miocene fossil crab from

southern South America—mainly Argentina, but also Chile, Peru, and Uruguay (e.g., Casadío et al.,

2005; Luque et al., 2017; Perea et al., 2020, and references therein). This species has been reported

from the Estancia 25 de Mayo Formation (previously known as Centinela Formation) in Tierra del

Fuego, Argentina, whose lower portion has been dated as upper Oligocene–Lower Miocene

(Chattian–Aquitanian, 27.28–20.44 Ma) based on $^{87}Sr/^{86}Sr$ radioisotopic dating ages between 25.5–

21.5 Ma from a valve of the oyster *Crassostrea*? *hatcheri* (Ortmann), collected below a tuff bed near

the Oligocene–Miocene (Chattian–Aquitanian) boundary (Casadío et al., 2000; Casadio et al., 2001;

Guerstein et al., 2004), and the palynomorphs *Operculodinium israelianum*, *Spiniferites ramosus*,

Lingulodinium machaerophorum, *Reticulatosphaera actinocoronata*, *Nematosphaeropsis* cf. *N.*

lativittata, *Baumannipolis variapertatus*, *Mutisiapollis viteauensis*, *Nothofagidites* spp., and

Podocarpidites spp. (Guerstein et al., 2004). As such, the most recent common ancestor of

Geryonidae and all of its descendants must have originated in the pre-Oligocene.

1033

1034

1035　## 21. Brachyura: Eubrachyura: Portunoidea: Portunidae: Thalamitinae (crown)

1036　***Fossil specimen.*** *Lessinithalamita gioiae* De Angeli and Ceccon, 2015. Museo Civico di Valdagno,

1037　Italy. Holotype MCV14/15, a fairly complete dorsal carapace.

1038

1039　***Phylogenetic justification***. The extinct, monotypic genus *Lessinithalamita* De Angeli and Ceccon,

1040　2015, shows remarkable affinities with the extant genera *Thalamita* Latreille, 1829, and *Thranita*

1041　Evans, 2018, justifying its inclusion within the subfamily Thalaminitae, which is nested within the

1042　crown group Portunidae (Evans, 2018).

1043

1044　***Minimum age***. 47.8 Ma.

1045

1046　***Tip maximum age.*** 53.3 Ma.

1047

1048　***Soft maximum age.*** 97 Ma.

1049

1050 *Age justification*. The specimen studied comes from compact calcarenites rich in coralline algae

1051 (e.g., *Lithothamnium bolcensis*) and calcareous nannofossils rocks cropping out near Monte Magrè,

1052 Vicenza, Italy, and dated approximately as lower Eocene (middle and upper Ypresian, ~53.3–47.8

1053 Ma) (Beccaro, 2003, in De Angeli and Ceccon, 2012).

1054   Soft maximum age as for calibration 20.

1055

Discussion. Besides *Lessinithalamita gioiae*, there are two other species of Eocene thalamitine crabs

belonging to the extinct genus *Eocharybdis* Beschin, Busulini, De Angeli & Tessier, 2002, i.e., *E.*

rugosa Beschin, Busulini, Tessier & Zorzin, 2016a, from the lower Eocene (Ypresian) of Monte di

Malo, Verona, NE Italy, and *E. cristata* Beschin, Busulini, De Angeli & Tessier, 2002, from the

middle Eocene (Lutetian) Cava "Main" di Arzignano, Vicenza, Italy. While both of these species are

suitable candidates for calibration points for the subfamily Thalamitinae, the completeness of the

type material of *L. gioiae* and its similarities to *Thalamita* and *Thranita*, support a more reliable

affiliation to the crown Thalamitinae.

1064

1065

22. Brachyura: Eubrachyura: Portunoidea: Polybiidae (crown)

1067 ***Fossil specimen.*** *Liocarcinus heintzi* Schweitzer and Feldmann, 2010a. Muséum national d'Histoire

1068 naturelle, Paris. Holotype MNHN R03778, a well-preserved dorsal carapace.

1069

1070 ***Phylogenetic justification.*** *Liocarcinus* Stimpson, 1858, is an extant portunoid genus nested within

1071 crown Polybiidae (Evans, 2018). The overall sub-hexagonal carapace of *L. heintzi*, slightly wider

1072 than long, with four anterolateral spines, the concave posterolateral margins, a convex posterior

1073 margin, the orbits directed forward, and marked dorsal regions, seem to conform with the forms

1074 seen among species of *Liocarcinus* (Schweitzer et al., 2021c).

1075

1076 ***Minimum age***. 27.82 Ma.

1077

1078 ***Tip maximum age.*** 33.9 Ma.

1079

1080 ***Soft maximum age.*** 97 Ma.

1081

Age justification. The studied specimen comes from the carbonatitic Calcaire à Astéries Formation,

cropping out near Monségur, Gironde, Western Aquitaine, France (Schweitzer and Feldmann,

2010a). The Calcaire à Astéries Formation has been dated as lower Oligocene (Rupelian, 33.9–27.82

Ma), based on the recognition of the foraminifera biozones SBZ 21 and 22A, containing *Nummulites*

fichteli, *N. intermedius N. vascus*, *Spiroclypeus*, *Operculina*, *Halkyardia* spp., *Neorotalia* spp.,

Peneroplis, and *Arenagula* and the ostreid *Crassostrea longirostris* (Cahuzac and Londeix, 2012;

Sztrákos and Steurbaut, 2017).

Soft maximum age as for calibration 20.

1090

Discussion. Besides *Liocarcinus heintzi*, there are several other occurrences of polybiid and

polybiid-like genera and species in the Paleogene. *Archaeogeryon corsolini* (Casadío, De Angeli,

Feldmann, Garassino, Hetler, Parras and Schweitzer, 2004, as *Proterocarcinus*), and *A. lophos*

(Feldmann, Casadío, Chirino-Gálvez, and Aguirre-Urreta, 1995), are known from the middle

Oligocene and Paleocene (Danian, 66.0–61.6 Ma) of Argentina, respectively (Feldmann et al., 1995;

Casadío et al., 2004; Casadío et al., 2005; Feldmann et al., 2011b). The latter species, represented by

the holotype and sole specimen, is moderately preserved and shows some dorsal and sternal features

that suggest affinity with portunids in general, but the fronto-orbital margin and part of the

anterolateral margins are eroded, precluding a detailed comparison with polybiids in general.

Moreover, the exact phylogenetic position of *Archaeogeryon* Colosi, 1924, among portunoids, as

well as that of the also extinct genus *Gecchelicarcinus* Beschin, Busulini, De Angeli, and Tessier,

2007, known from two species from the Ypresian of Italy, is unclear. Both taxa could represent

extinct polybiid genera, but they cannot be assigned to the crown group with certainty.

The presumably polybiid monotypic extinct genera *Boschettia* Busulini, Tessier, Beschin,

and De Angeli, 2003, from the lower and middle Eocene (Ypresian to Lutetian, 56.0–41.2 Ma) of

Italy, and *Falsiportunites* Collins and Jakobsen, 2003, from the middle Eocene (?Lutetian, 47.8–

41.2 Ma) of Denmark, show a combination of anatomical features on their fronto-orbital margins,

anterolateral margins, and dorsal carapaces that differ from those seen among polybiids and thus

cannot be accommodated satisfactorily among the crown group either.

Among the extant genera with known fossil records, *Liocarcinus oligocaenicus* (Pauca,

1929, as *Portunus*), and *L. atropatanus* (Aslanova and Dschafarova, 1975) are known from the

lower and upper Oligocene (Rupelian–Chattian, 33.9–23.3 Ma) of Romania and Azerbaijan,

respectively (Schweitzer et al., 2009; Beschin et al., 2016b; Hyzny, 2016), which confirm the

presence of the genus during Oligocene times. *Liocarcinus priscus* Beschin, De Angeli, Checci, and

Zarantonello, 2016b, from the lower Eocene (lower Lutetian, 47.8–41.2 Ma), may represent the

oldest occurrence of the genus and thus a more approximate soft maximum age of calibration for the

polybiid node, but due to its remarkable differences with the type species of the genus *Liocarcinus,*

i.e., *L. holsatus* (Fabricius, 1798), to which *L. heintzi* is anatomically closer, we favor the younger

fossil for a calibration point. Regardless of which of these fossils is considered the oldest reliable

record of the family, the clear occurrence of several Paleogene genera and species attributable to

Polybiidae indicate that the most recent common ancestor of the family and all of its descendants

originated most likely during the early Paleogene (Wolfe et al., 2022), or even the Late Cretaceous.

1123

1124

1125   **23. Brachyura: Eubrachyura: Parthenopoidea: Parthenopidae+Dairoididae (crown)**

1126   ***Fossil specimen.*** *Aragolambrus collinsi* Ferratges, Zamora, and Aurell, 2019. Palaeontological

1127   collection of the Museo de Ciencias Naturales de la Universidad de Zaragoza, Spain. Holotype

1128   MPZ-2019/211, a nearly complete dorsal carapace with chelipeds associated.

1129

Phylogenetic justification. In a recent molecular phylogenetic study, Wolfe et al. (2022) recovered

Dairoides kusei as sister to a monophyletic Parthenopidae, whereas Ferratges et al. (2023, in press),

based on a morphological phylogenetic study of Parthenopoidea, recovered the extinct monotypic

genus *Aragolambrus* Ferratges, Zamora, and Aurell, 2019, in close proximity to the also extinct

genus *Phrynolambrus* Bittner, 1893, and the extant genus *Dairoides* Stebbing, 1920, all three

forming a well-supported monophyletic subfamily Dairoidinae sister to Daldorfiinae and nested

within a paraphyletic Parthenopidae (Ferratges et al., 2023, in press). Future comprehensive

molecular phylogenetic studies with larger taxon coverage across multiple parthenopoid genera will

shed light on the internal relationships among these groups.

1139

1140   ***Minimum age***. 53.02 Ma.

1141

1142   ***Tip maximum age.*** 57.1 Ma.

1143

1144   ***Soft maximum age.*** 100.9 Ma.

1145

Age justification. *Aragolambrus collinsi* comes from the coral-algal Reef Limestones interval

(middle member) of the Serraduy Formation, in the Barranco de Ramals outcrop near Puebla de

Roda and Serraduy, Huesca province, Spain (Ferratges et al., 2019). The Reef Limestones interval

develops over the underlying Alveolina Limestones interval and the overlying marls of the Riguala

Member (Serra-Kiel et al., 1994; Pujalte et al., 2009). Serra-Kiel et al. (1994) assigns an age of

lower Ilerdian to the Alveolina Limestones interval, which is located within the

magnetostratigraphic Chron C24r (57.101–53.9 Ma, Francescone et al., 2019; Ogg, 2020), and

contains the macroforaminifera *Alveolina cucumiformis, A. ellipsoidalis*, and *Nummulites fraasi*,

from the NP9, BB1, and P6a Biozones. For the marls of the Riguala Member, the same authors

assigns an age of middle Ilerdian, roughly located between the top of the magnetostratigraphic

Chron C24r and Chron C24n.1r (~53.9–53.02 Ma, Francescone et al., 2019; Ogg, 2020), and

contains the macroforaminifera *Alveolina moussoulensis*, *A. corbarica*, *Nummulites robustifrons,*

and *N. exilis*, from the BB1–BB2, NP9–NP10, and P6a-P6b Biozones (Serra-Kiel et al., 1994). Due

to the lateral facial discontinuity of the Reef Limestones interval, which is bounded at the bottom by

the Alveolina Limestones and at the top by the Riguala Member, we bracket the tip maximun and

minimum ages of *A. collinsi* between 57.1–53.02 Ma, with a possible age of the fossil itself closer to

53.5 Ma (F. Ferratges, pers. comm. to J. Luque, Feb. 2023).

Soft maximum age as for calibration 8.

1164

Discussion. *Aragolambrus* and *Phrynolambrus* are two of the oldest fossils of crown Parthenopidae

known, sharing with *Dairoides* the presence of two to three anterolateral spines, an ornamented

epistomia with one or more rows of tubercles, and the pterygostome with grooves, which are

synapomorphies that unite them among parthenopoids under the subfamily Dairoidinae (Ferratges et

al., 2023, in press).

The extinct genera *Eogarthambrus* De Angeli, Garassino, and Alberti, De Angeli et al.,

2010a, and *Mesolambrus* Müller and Collins, 1991, both known from the Eocene (middle Ypresian

to Priabonian, ~52.0–33.9 Ma) of Italy, were recovered by Ferratges et al. (2023, in press) outside

the crown Parthenopoidea. The monotypic extinct genus *Braggilambrus* De Angeli and Caporiondo,

2016, from the Ypresian of Italy, seems to fit within the general parthenopoidean body form, but due

to its incompleteness and therefore lack of key diagnostic characters, its affinities at the family and

subfamily levels is still unclear.

The presence of several crown parthenopid forms across the early to late Eocene indicate

that the most recent common ancestor of Parthenopoidea and all of its descendants must have

Paleocene or even Late Cretaceous origins.

1180

1181

1182   **24. Brachyura: Eubrachyura: Calappoidea: Calappidae (crown)**

1183   ***Fossil specimen.*** *Calappa zinsmeisteri* Feldmann and Wilson, 1988. Smithsonian Institution,

1184   National Museum of Natura History (NMHN), USA. Holotype USNM 404877, claws only.

1185

1186   ***Phylogenetic justification.*** *Calappa* Weber, 1795, is the type genus of the family Calappidae De

1187   Haan, 1833, and thus phylogenetically nested within the crown Calappoidea (Lu et al., 2020).

1188

1189   ***Minimum age***. 33.9 Ma.

1190

1191   ***Tip maximum age.*** 41.2 Ma.

1192

1193   ***Soft maximum age.*** 100.9 Ma.

1194

Age justification. The type material of *Calappa zinsmeisteri* comes from the uppermost La Meseta

Formation on Seymour Island, Antarctica, Locality 14, of Feldmann and Wilson (1988). This

locality corresponds to the Telms VI–VII of Sadler (1988), which have been dated as middle to

upper Eocene (Bartonian–Priabonian, ~42.0–34.0 Ma) based on ^{87}Sr/^{86}Sr radiometric dating ages

from *Cucullaea* shells from Telm VII between 46.13 and 34.69 Ma (Dutton et al., 2002), and the

occurrences of the neogastropods *Prosipho lawsi*, *P. lamesetaensis*, *Austroficopsis meridionalis*,

Microfulgur byrdi, *Fusinus*? *suraknisos*, *Eupleura suroabdita*, ?*Adelomelon suropsilos*, *Zemacies*

finlayi, and *Aforia canalomos* (Crame et al., 2014).

Soft maximum age as for calibration 8.

1204

Discussion. Among the extant calappid genera, only *Calappa* and *Mursia* Leach, *in* Desmarest,

1823, have recognized Eocene fossils, to our knowledge. Four species of *Calappa*—including *C.*

zinsmeisteri— are known from the middle and upper Eocene of USA, Antarctica, and presumably

Venezuela, and are constituted largely by cheliped (propodi) remains with diagnostic anatomical

details that can be assigned with certainty to crown Calappidae, and most likely to *Calappa* (Ross et

al., 1964; Feldmann and Wilson, 1988; Luque et al., 2017). One species of *Mursia* Leach, *in*

Desmarest, 1823, i.e., *Mursia aspina* Schweitzer and Feldmann, 2000a, from the Late Eocene of

Washington, USA, is represented by a handful of dorsal carapaces that strongly resemble *Mursia*,

although they lack some of the diagnostic features of the genus such as developed lateral spines.

While we considered the cheliped remains of *Calappa zinsmeisteri* as our calibration point given

their putative affinities to the type genus of the family, *M. aspina*, which is represented by dorsal

carapace material of similar age as *C. zinsmeisteri*, could also be a good candidate to calibrate the

node for crown Calappidae.

Among the extinct calappid genera, *Calappilia* A. Milne-Edwards in de Bouillé, 1873, bears

conspicuous anatomical similitudes with *Calappa*, and has over 20 species known, of which more

that 15 are Eocene in age (e.g., Williams and Child, 1988; Schweitzer et al., 2010; Rumsey et al.,

2016; Luque et al., 2017; Feldmann et al., 2019, and references therein). There are also several

extinct genera known exclusively from the Eocene, such as the monotypic genera

Paracorallomursia Beschin, Busulini, Tessier & Zorzin, 2016a, and *Pseudocorallomursia* Beschin,

Busulini, Tessier & Zorzin, 2016a, from the Ypresian of Italy, *Carinocalappa* Beschin, Busulini,

and Tessier *in* Beschin et al., 2018, from the Priabonian of Italy, and *Tavernolesia* Artal and Onetti,

2017, from the Lutetian of Spain. *Corallomursia* De Angeli and Ceccon, 2014, comprises two

species, both from the Ypresian of Italy (Beschin et al., 2015). All of these extinct genera seem to fit

well within the total group Calappidae, but their precise affinities with crown calappids remain

unclear.

Calappid crabs were already diverse, speciose, and widespread by the Eocene, indicating that

the most recent common ancestor of Calappidae and all of its descendants must have originated in

the earliest Paleogene (Paleocene) or the Cretaceous.

1233

1234

1235 ## 25. Brachyura: Eubrachyura: Cancroidea: Cancridae (crown)

1236 *Fossil specimen.* *Anatolikos undecimspinosus* Schweitzer, Feldmann, González-Barba, and Ćosović,

1237 2006. Museo de Historia Natural, Universidad Autónoma de Baja California Sur, La Paz, Baja

1238 California Sur, México (MHN-UABCS). Holotype MHN-UABCS/Ba12-9, a partially preserved

1239 dorsal left carapace.

1240

Phylogenetic justification. The extant genus *Anatolikos* Schweitzer and Feldmann, 2000b, can be

placed with confidence within the family Cancridae (Schweitzer et al., 2006; Schram and Ng, 2012;

Wolfe et al., 2022). The diagnostic shape and arrangement of the eleven anterolateral spines seen in

the extant genus *Anatolikos* and thus in *A. undecimspinosus*, together with the presence of a nearly

straight rostrum with "five coalesced spines separated by fissures" (Schweitzer and Feldmann,

2000b) are unique traits not shared with other genera within the family (Schweitzer et al., 2006;

Poore and Ahyong, 2023).

1248

1249   *Minimum age*. 41.2 Ma.

1250

1251   *Tip maximum age.* 56.0 Ma.

1252

1253   *Soft maximum age.* 100.9 Ma.

1254

Age justification. The type material of *Anatolikos undecimspinosus* comes from middle Eocene

rocks of the Bateque Formation, cropping out at locality Waypoint 29 of Schweitzer et al. (2006),

northwest of La Paz, near the village of San Ignacio, Baja California Sur, Mexico. The age of the

Bateque Formation has been constrained between the early Eocene and the late middle Eocene

(56.0–41.2 Ma) based on foraminiferal, nannoplankton, and diatom assemblages, more than 150

macroinvertebrates, and ostracods (Mina-Uhink, 1957; Sorensen, 1982; McLean et al., 1987;

Carreño and Cronin, 1993; Morales-Ortega et al., 2016).

Soft maximum age as for calibration 8.

1263

Discussion. Among the cancrids with known fossil record, only the Cancrinae genera *Anatolikos*

and presumably *Cancer* Linnaeus, 1758, have Eocene occurrences. However, *Cancer gabbi*

Rathbun, 1926, from the Eocene of California, is represented by claw fragments only that cannot be

confirmed to belong to the group. *Cancer meticuriensis* Thurmann, 1853, from the Oligocene and

presumably from the lower Eocene of France, is too incomplete, with only the sternum and the

underside of the front legs known, which A. Milne-Edwards (1865) considered to most likely belong

to a portunid. At least six species (*C. aldenardensis*, *C. archiaci*, *C. flandricus*, *C. pratti*, 1850, *C.*

rotnacensis, and *C. villabersiani*) from the Ypresian–Lutetian of Europe, are all *nomina nuda*, many

of which have never been described or figured, and therefore their generic affinities are dubious

(Van Straelen, 1920; Nations, 1975). *Cancer santosi* (Rathbun, 1937, as *Lobocarcinus*?) from the

upper Eocene of Panama, is a poorly preserved right dorsal carapace.

Three extinct monotypic genera with potential Carcininae affinities have also early to middle

Eocene occurrences, i.e., *Santeecarcinus* Blow and Manning, 1996, and *Sarahcarcinus* Blow and

Manning, 1996, both from USA, and *Notocarcinus* Schweitzer and Feldmann, 2000b, from

Argentina. However, given that these three genera are extinct only, we consider them suboptimal

candidates for the calibration of the Cancridae node, given the presence of fossil taxa assigned to

extant genera, e.g., *Anatolikos undecimspinosus*, in rocks of similar ages.

Several other extinct cancroid genera currently placed within the subfamily Lobocarcininae

Beurlen, 1930 (e.g., De Grave et al., 2009; Schweitzer et al., 2010), also have their oldest or only

records in the Eocene. These include *Miocyclus* Müller, 1978, from Bulgaria, *Nicoliscarcinus*

Beschin, Busulini, Tessier & Zorzin, 2016a, and *Ramacarcinus* De Angeli & Ceccon, 2017, from

Italy, and *Tasadia* Müller, *in* Janssen and Muller, 1984, from Belgium. In addition, two out of the

three species of the extinct genus *Ceronnectes* De Angeli and Beschin, 1998, and several species of

Lobocarcinus Reuss, 1857, also occur in the middle Eocene (e.g., Anderson and Feldmann, 1995;

De Angeli and Beschin, 1998; Feldmann et al., 1998; Schweitzer et al., 2010; Beschin et al., 2016a).

1289

1290

1291 **26. Brachyura: Eubrachyura: Dorippoidea (crown)**

1292 ***Fossil specimen.*** *Bartethusa hepatica* Quayle and Collins, 1981. Palaeontological Department,

1293 London's Natural History Museum (formerly British Museum (Natural History)), England, holotype

1294 BM In.61704, a well-preserved dorsal carapace.

1295

Phylogenetic justification. *Bartethusa hepatica* has an overall carapace outline, fronto-orbital

construction, and dorsal carapace region configuration that matches the range of forms seen across

crown dorippoids. While *Bartethusa hepatica* could be allied to either Ethusidae Guinot, 1977 (e.g.,

Quayle and Collins, 1981; Van Bakel et al., 2021) or Dorippidae MacLeay, 1838a (e.g., Luque,

2015b; Schweitzer et al., 2021b), in either case it will still be nested within the crown Dorippoidea

MacLeay, 1838a.

1302

1303 ***Minimum age***. 37.71 Ma.

1304

1305   ***Tip maximum age.*** 41.2 Ma.

1306

1307   ***Soft maximum age.*** 100.9 Ma.

1308

Age justification. The type material of *Bartethusa hepatica* was collected in rocks of the Horizon

A3, Barton Beds, Christchurch Bay, Isle of Wight, England (Quayle and Collins, 1981), which have

been dated as middle Eocene (Bartonian, 41.2–37.71 Ma) based on larger and smaller foraminifera,

calcareous nannofossils, palynomorphs, and paleomagnetism (see Cotton et al., 2021, and references

therein). Tsang et al. (2014) also used *B. hepatica* as their calibration point, but they assigned it an

age of lower Eocene (Ypresian, 56.0–47.8 Ma). However, the available stratigraphic information

indicates that the correct age of the known fossils of *B. hepatica* is middle Eocene (Bartonian, 41.2–

37.71 Ma), based on larger and smaller foraminifera, calcareous nannofossils, palynomorphs, and

paleomagnetism (see Cotton et al., 2021, and references therein).

Soft maximum age as for calibration 8.

1319

Discussion. Dorippoid crabs—extant and fossil—are among the most puzzling of the eubrachyurans

in terms of their phylogenetic affinities, given their unique combination of plesiomorphic and

apomorphic characters and their relatively low representation in previous molecular phylogenies

compared to most other crabs. Based on morphology alone, dorippoids have been typically regarded

as some of the oldest crown eubrachyurans, and thus one of the earliest splitting branches in the

eubrachyuran tree, with their earliest known representatives, e.g., Telamonocarcinidae Larghi, 2004,

and potentially Tepexicarcinidae Luque, 2015b, already present in the Early Cretaceous (Luque,

2015b; Luque et al., 2021; Van Bakel et al., 2021). Although their overall anatomy is strongly

reminiscent of the array of body forms across modern dorippoids in terms of their dorsal carapaces,

thoracic sternum, and shape and size of their pereopods (P2–P5), they cannot be assigned to any of

the two crown families recognized today, i.e., Dorippidae and Ethusidae. Some of these other extinct

families, however, may represent convergent morphology, as dorippoids consistently nest well

within heterotremes in molecular phylogenies (Tsang et al., 2014; Wolfe et al., 2022).

Among the extinct post-Cretaceous dorippoid forms, *Goniochele* Bell, 1858, presumably

belonging to its own family Goniochelidae Schweitzer and Feldmann, 2011, has a confirmed fossil

record restricted to two Eocene species: *G. angulata* Bell, 1858, from the Ypresian of England, and

G. madseni Collins and Jakobsen, 2003, from the Ypresian–Lutetian of Denmark. Rathbun (1918),

based on an isolated dactylus from the Early Miocene (not Oligocene, as Rathbun suggested)

Culebra Formation in the Panama Canal, erected a third species of *Goniochele* that currently cannot

be ascribed to this genus with certainty (see Luque et al., 2017). Another extinct taxon,

Archaeocypoda veronensis Secrétan, 1975, which known from the lower Eocene (upper Ypresian)

of Italy (Pasini et al., 2019), has been presumably assigned to Dorippidae based on the sub-oval

carapace outline, the broad front and continuous supraorbital margin, the smooth anterolateral and

posterolateral margins, the inflated protogastric and branchial regions, the smooth antero- and

posterolateral margins (Casadío et al., 2005; Pasini et al., 2019; Schweitzer et al., 2021b). While

these features match some of the diagnostic characters of dorippids, the possession of at least three

pairs of well-developed pereopods (i.e., P2–P4) dramatically differs from the apomorphic dorippoid

possession of two large pereopods (P2–P3) and two reduced, subchelate, and sub-dorsally carried

pereopods (P4–P5), which is a feature seen across crown dorippoids and even stem dorippoids from

the Cretaceous (e.g., Larghi, 2004; Luque, 2015b; Guinot et al., 2019). As such, while *A. veronensis*

might be related to dorippoids, its systematic affinities remain unclear. Another taxon, the presumed

genus *Titanodorippe* Blow and Manning, 1996, from the middle Eocene of USA, is a single

cheliped propodus that cannot be assigned with confidence to Dorippidae or even Dorippoidea.

Among the extant dorippoid genera, only *Ethusa* Roux, 1830 [in Roux, 1828-1830], has its

oldest record in the uppermost Eocene (Priabonian, 37.71–33.9 Ma) (Müller and Collins, 1991),

while *Dorippe* Weber, 1795, and *Medorippe* Manning and Holthuis, 1991, are known from the

lower-middle Miocene (Burdigalian, 20.44–15.97 Ma) (Studer, 1892; Karasawa, 2000). Given the

close proximity in age between the occurrences of *Bartethusa* and *Ethusa* from the middle and late

Eocene, respectively, the most recent common ancestor of crown Dorippoidea and all of its

descendants must have originated during the early Paleogene or even the Cretaceous, while the

presence of several potential stem dorippoids in the Early and Late Cretaceous worldwide support a

late Mesozoic origin for the superfamily.

1362

1363

1364   ## 27. Brachyura: Eubrachyura: Leucosioidea: Leucosiidae (crown)

1365   ***Fossil specimen.*** *Typilobus alponensis* Beschin, De Angeli, and Zorzin, 2009. Museo di Storia

1366   Naturale di Verona. Holotype IGVR78539, a complete dorsal carapace in volume.

1367

1368 ***Phylogenetic justification.*** The overall oval and domed carapace about as long as wide, with convex

1369 anterolateral and posterolateral margins, a narrow posterior margin, the small and round orbits, and

1370 the presence of a projected bilobed front/rostrum that is mesially depressed, are typical features of

1371 Leucosiidae, an extant family nested within Leucosioidea (Beschin et al., 2009).

1372

1373 ***Minimum age***. 47.8 Ma.

1374

1375 ***Tip maximum age.*** 56.0 Ma.

1376

1377 ***Soft maximum age.*** 100.9 Ma.

1378

1379 ***Age justification***. The holotype and sole specimen of *Typilobus alponensis* comes from lower

1380 Eocene (Ypresian, 56.0–47.8 Ma) rocks cropping out in the valley of Alpone, near Monte Serea, San

1381 Giovanni Ilarione, Verona, Italy (Beschin et al., 2009).

1382   Soft maximum age as for calibration 8.

1383

Discussion. *Typilobus* Stoliczka, 1871, is an extinct genus with presumably over a dozen species

ranging in age from early Eocene (Ypresian) to Miocene and widely distributed across Europe, the

UK, North Africa, Asia, and the Indo-Pacific (Karasawa, 1998; Beschin et al., 2009; Artal and

Hyžný, 2016; Karasawa et al., 2019). However, several of these species are considerably different

from the type species of the genus, *T. granulosus* Stoliczka, 1871, from the Early Miocene of

Pakistan, casting doubts on the monophyletic nature of the genus as currently envisioned.

Regardless of the generic affinities of *Typilobus alponensis*, this species fits well within the overall

body plan seen across crown Leucosiidae (see above under *Phylogenetic justification*), hence our

species selection for the calibration of the node. Other Eocene species assigned to *Typilobus* that

might be related to *T. alponensis* and thus belong to the crown Leucosiidae are *T. belli* Quayle and

Collins, 1981, from the Bartonian of the UK, *T. prevostianus* (Desmarest, 1822, as *Leucosia*), from

the Lutetian of Paris, *T. semseyanus* Lörenthey 1898, from the Lutetian–Priabonian of Hungary and

Italy (Lorenthey and Beurlen, 1929; Beschin et al., 1998), and *T. trispinosus* Lörenthey, 1907, from

the Lutetian–Priabonian of Egypt (Beschin et al., 1998; Feldmann et al., 2011a; Artal and Hyžný,

2016).

The type material of the monotypic genus *Zannatoius* Beschin, Busulini & Tessier *in*

Beschin et al., 2018, from the upper Eocene of Italy, is somewhat damaged and missing the fronto-

orbital margin, but overall it conforms with the leucosiid body plan. Among the extant leucosiid

genera with known fossil record, *Ebalia* Leach, 1817, has its earliest putative occurrence in the

upper Eocene (Priabonian, 37.71–33.9 Ma) of Germany (Förster and Mundlos, 1982; Karasawa et

al., 2019). As such, the presence of crown Leucosiidae in the early and late Eocene suggests that the

family must have originated at least during the early Paleogene.

1406

1407

1408   ## 28. Brachyura: Eubrachyura: Goneplacoidea: Euryplacidae (crown)

1409   ***Fossil specimens.*** *Chirinocarcinus wichmanni* (Feldmann, Casadío, Chirino-Galvez, and Aguirre-

1410   Urreta, 1995, as ?*Glyphithyreus*). Geology Collections at National University of La Pampa, Santa

1411   Rosa, La Pampa, Argentina. Holotype GHUNLPam 7015, a nearly complete dorsal carapace.

1412

1413   ***Phylogenetic justification.*** *Chirinocarcinus wichmanni* has been included within Euryplacidae

1414   based on the relatively broad fronto-orbital margin and large orbits, the well-defined supraorbital

1415   angle, and short anterolateral margins (Feldmann et al., 1995; Karasawa and Kato, 2003; Castro and

1416   Ng, 2010).

1417

1418   ***Minimum age***. 61.6 Ma.

1419

1420   ***Tip maximum age.*** 66.0 Ma.

1421

1422   ***Soft maximum age.*** 100.9 Ma.

1423

Age justification. The type material of *Chirinocarcinus wichmanni* was collected from lowermost

Paleogene (lower Danian) rocks of the Roca Formation at the type section, located to the north of

General Roca, Rio Negro, Neuquén Province, Argentina (Feldmann et al., 1995). The Roca

Formation has been dated as uppermost Cretaceous (Maastrichtian) to lower Paleocene (Danian),

based on foraminifera, calcareous nannoplankton, and palynomorphs (del Rio et al., 2011 and

references therein). The crab-bearing intervals of the Roca Formation are comprised within the

calcareous nannofosil zones NP3 and NP4, with *Chiasmolithus danicus*, *Neochiastozygus modestus*,

Hornibrookina edwardsii, *H. teuriensis*, and *Toweius africanus* from NP3, indicative of the lower

Danian (del Rio et al., 2011).

Soft maximum age as for calibration 8.

1434

Discussion. Goneplacoidea, as traditionally envisioned, is among the most anatomically disparate

and problematic brachyuran groups in terms of the clarity about their internal phylogenetic

affinities, and thus their likely polyphyletic relationships with other eubrachyuran groups (Ng et al.,

2008; Castro et al., 2010). The extant family Euryplacidae, however, seems to be monophyletic

(Castro and Ng, 2010) and nested within the crown Goneplacoidea (Karasawa and Kato, 2003).

Among the euryplacid genera, the monotypic *Chirinocarcinus* Karasawa and Schweitzer,

2004, is the only taxon known from Paleocene fossils. Five other monotypic genera currently

assigned to Euryplacidae are known from the lower Eocene (Ypresian) of Italy and the middle

Eocene (Lutetian) of Spain (Karasawa and Kato, 2003; Beschin and De Angeli, 2011; Beschin et al.,

2016a), as it is the genus *Corallicarcinus* Müller and Collins, 1991, from Eocene fossils from

Hungary and Italy (Müller and Collins, 1991; Karasawa and Kato, 2003; Beschin et al., 2018).

Similarly, the genus *Orbitoplax* Tucker and Feldmann, 1990, also referable to Euryplacidae, is

known from four Eocene species occurring in Mexico and USA (e.g., Rathbun, 1926; Tucker and

Feldmann, 1990; Schweitzer, 2000; Vega et al., 2001).

Among the extant euryplacid genera, *Eucrate* De Haan, 1835, has its oldest putative record

in the Oligocene, as represented by a fairly complete ventral carapace with chelipeds of *E. martini*

Rathbun, 1926 from USA, and *E. puliensis* Hu & Tao, 1996, from Taiwan. ?*Euryplax culebrensis*

Rathbun, 1918 [1919], from the Panama Canal Zone, is of lower Miocene age and not Oligocene,

just like the other fossil decapods from the Culebra Formation in the Panama Canal, as initially

assumed by Rathbun and repeated by several authors (see in Luque et al., 2017 and comments

therein).

The number of extinct and extant euryplacids known from the Paleogene, and the proximity

of *C. wichmanni* to the Cretaceous/Paleogene boundary, indicates that Euryplacidae must have a

pre-Cenozoic origin, most likely in the Cretaceous.

1459

1460

1461   **29. Brachyura: Eubrachyura: Eriphioidea: Eriphiidae (crown)**

1462   ***Fossil specimens.*** *Eriphia verrucosa* (Forskal, 1775), in Betancort et al., 2014. Repository indet.

1463   Specimen number indet. Single left cheliped fragment with palm and pollex.

1464

1465   ***Phylogenetic justification.*** The type species of *Eriphia* Latreille, 1817, is *E. verrucosa* (Forskal,

1466   1775) [*E. spinifrons* (Herbst, 1782)], by subsequent designation (see in Koh and Ng, 2008; Ng et al.,

1467   2008). Although Eriphioidea as previously envisioned is shown to be polyphyletic (see Wolfe et al.,

1468   2022), *Eriphia* is the type genus of the family Eriphiidae and thus the superfamily Eriphioidea. As

1469   such, it is reliably placed within the crown group.

1470

1471   ***Minimum age***. 4.1 Ma.

1472

1473   ***Tip maximum age.*** 9.3 Ma.

1474

1475   ***Soft maximum age.*** 100.9 Ma.

1476

1477   ***Age justification***. The specimen of *E. verrucosa* reported by Betancort et al., 2014 comes from

1478   rocks of the Papagayo cliff, Lanzarote, Canary Islands, Spain, dated as upper Miocene (Messinian?)

1479   to lower Pliocene from K/Ar radiometric dating, ranging between 9.3 and 4.1 Ma (see in Betancort

1480   et al., 2014, and references therein).

1481   Soft maximum age as for calibration 8.

1482

Discussion. The family Eriphiidae, as currently recognized, includes only the extant genera *Eriphia*

Latreille, 1817, and *Eriphides* Rathbun, 1897 (Ng et al., 2008; De Grave et al., 2009). While

Eriphides is monotypic, Eriphia is represented by eight extant and two fossil species (Koh and Ng,

2008; Schweitzer et al., 2010). The extinct *Eriphia cocchii* Ristori, 1886, is known from well-

preserved specimens in volume as well as from isolated dactyli from the lower Pliocene (Zanclean–

Piacentian, ~5,33–3.6 Ma) of Italy (Ristori, 1886; De Angeli et al., 2009). Based solely on an

isolated propodus, Collins and Donovan, 1997, report a presumably new species, *Eriphia*

xaymacaensis Collins and Donovan, 1997, from the upper Pleistocene of Jamaica (Luque et al.,

2017). The latter species, however, might be a junior synonym of *Eriphia gonagra* Fabricius, 1781,

an extant species. Along these lines, among the extant species with putative fossils, Varola (1981)

reported isolated dactyli of the extant species *E. verrucosa* from the mid-Pliocene of Italy, and

Luque et al., 2018, reported a cheliped of *Eriphia* cf. *squamata* Stimpson, 1859, from the

Quaternary of Panama.

As we show here, the fossil record of Eriphiidae is meager and restricted to the uppermost

Neogene and Quaternary, hence why we opt to use a calibration fossil with uncertain specimen data.

As such, the most recent common ancestor of extant eriphiids and all of its descendants must have

originated at least in the early Neogene or, most likely, in the Palaeogene.

1500

1501

1502 **30. Brachyura: Eubrachyura: Trapezioidea: Trapeziidae (crown)**

1503 ***Fossil specimen.*** *Archaeotetra lessinea* De Angeli and Ceccon, 2013. Museo Civico "D. Dal Lago"

1504 of Valdagno (Vicenza) (MCV). Holotype MCV 12/05-I.G.360311, a complete dorsal carapace.

1505

Phylogenetic justification. Phylogenetically, Trapeziidae Miers, 1886 seems to be a monophyletic

family nested within Trapezioidea s.l. (Castro et al., 2004; Lai et al., 2009; Wolfe et al., 2022), but

the monophyly of Trapezioidea as a whole, containing Tetrallidae Castro, Ng, and Ahyong, 2004,

has been questioned (see Wolfe et al., 2022). Despite this, the distinctive carapace outline of

Archaeotetra Schweitzer, 2005, and thus of *Archaeotetra lessinea*, with a wide and slightly bilobed

front, the fronto-orbital margin being about 80–90% as wide as the carapace maximum with, the

orbits placed anterolaterally, the smooth dorsal carapace, and the seemingly spineless, short, and

nearly parallel anterolateral margins, conforms with the body form typical of genera and species in

the family Trapeziidae (Schweitzer, 2005; Karasawa and Schweitzer, 2006; De Angeli and Ceccon,

2013; Poore and Ahyong, 2023).

1516

1517 ***Minimum age***. 47.8 Ma.

1518

1519 ***Tip maximum age.*** 53.3 Ma.

1520

1521 ***Soft maximum age.*** 100.9 Ma.

1522

1523 *Age justification*. The material studied comes from micritic limestones with abundant nullipore

1524 coralline algae and fragments of invertebrates, cropping out in the vicinity of Monte Magré, eastern

1525 margin of Lessini Mounts, Italy. Age as for calibration 21.

1526   Soft maximum age as for calibration 8.

1527

Discussion. Among the few trapeziid crab fossils known, at least four species besides *Archaeotetra*

lessinea have been recovered from Eocene deposits. *Archaeotetra inornata* Schweitzer, 2005, is

known from the lower to middle Eocene of Mexico, *Paratetralia convexa* Beschin, Busulini, De

Angeli, and Tessier, 2007, from the lower Eocene of Italy, and *Eomaldivia pannonica* and *E.*

trispinosa Müller and Collins, 1991, from the upper Eocene of Hungary. This indicates that the most

recent common ancestor of trapeziid crabs and all of its descendants must have a pre-Eocene origin

(Wolfe et al., 2022).

1535

1536

1537 ### 31. Brachyura: Eubrachyura: Eriphioidea: Oziidae (crown)

1538 *Fossil specimen. Ozius collinsi* Karasawa, 1992. Mizunami Fossil Museum (Yamanouchi,

1539 Mizunami, Japan). Holotype MFM39001, a complete dorsal carapace.

1540

1541 *Phylogenetic justification*. The extant genus *Ozius* Milne Edwards, 1834 [in H. Milne Edwards,

1542 1834-1840] is the type genus of the family Oziidae. The overall anatomy of *Ozius collinsi* fits well

1543 the diagnosis of the genus, and thus can be considered within the crown Oziidae s.s.

1544

1545 *Minimum age*. 13.82 Ma.

1546

1547 *Tip maximum age.* 15.97 Ma.

1548

1549 *Soft maximum age.* 100.9 Ma.

1550

Age justification. The type material of *Ozius collinsi* comes from mudstones of the Yoshino

Formation, Katsuta Group, Locality KTT13, in Niida, Tsuyama City, Okayama Prefecture, Japan

(Karasawa, 1992). The Yoshino Formation is considered as uppermost lower Miocene to middle

Miocene (Burdigalian to Langhian), with some estimated ages of 17.9± 2.1 Ma and 18.5–16 Ma

based on the fission track method and palynomorphs (Mori and Yamanoi, 2003; Suzuki et al., 2003,

in Ando et al., 2016), and on mollusc assemblages and planktonic foraminifera biostratigraphic ages,

zones N8 to N10 (Blow, 1969; Yoshimoto, 1979, in Karasawa, 1992 and Taguchi, 2002), which

correspond to the global Burdigalian. Other authors have restricted the age of the oldest fossil

Oziidae from Japan to the Langhian (Schweitzer et al., 2021a).

Soft maximum age as for calibration 8.

1561

Discussion. In some recent mitogenomic studies, oziid taxa such as *Epixathus frontalis* have been

recovered outside the traditional Eriphioidea, highlighting the polyphyletic nature of the superfamily

as traditionally envisioned (Tan et al., 2018; Lü et al., 2022), and further corroborated in Wolfe et al.

(2022). Moreover, previous groupings of genera within the family Oziidae have been shown to be

non-monophyletic (Lai et al., 2014), further adding to the phylogenetic issues highlighted above.

This also adds to the problem that several extinct forms that might superficially resemble oziids

might not necessarily belong to Oziidae s.s., which, together with their incompleteness, preclude a

confident systematic placement. As such, due to the considerably reliable placement of *O. collinsi*

within the genus *Ozius*, and due to its early-mid Miocene age, we consider it as the best calibration

point for the family currently available.

1572

1573

1574   **32. Brachyura: Eubrachyura: Pilumnoidea: Pilumnidae (crown)**

1575 ***Fossil specimen.*** *Glabropilumnus trispinosus* Beschin, Busulini, and Tessier, *in* Beschin et al.,

1576 2016a. Museo di Storia naturale di Verona. Holotype VR 94508, a complete dorsal carapace in

1577 volume.

1578

Phylogenetic justification. The overall carapace shape of *G. trispinosus*, with smooth dorsal surface

and lacking dorsal ridges, and the anterolateral margin with 3 spines, being the posterior one the

smallest of all, match well the diagnosis of *Glabropilumnus* (Poore and Ahyong, 2023).

Glabropilumnus Balss, 1932, is an extant genus represented by six extant species (Ng et al., 2008),

and the only genus in the Family Pilumnidae Samouelle, 1819, and the subfamily Pilumninae

Samouelle, 1819, with a fossil record extending into the Eocene (see below), hence our taxon

selection for calibration.

1586

1587 *Minimum age*. 47.8 Ma.

1588

1589 *Tip maximum age.* 56.0 Ma.

1590

1591 *Soft maximum age.* 100.9 Ma.

1592

1593 *Age justification*. *Glabropilumnus trispinosus* is known from its type locality in the area of Rama di

1594 Bolca (Verona), NE Italy, in calcareous rocks of the 'Calcari Nummulitici' that have been ascribed

1595 to the lower Eocene (Ypresian) based of the presence of *Nummulites partschi*, *Alveolina cremae*, *A.*

1596 *croatica*, *A. decastroi*, *A.* cf. *dainellii*, and *A. distefanoi* (Beschin et al., 2016a, and references

1597 therein).

1598   Soft maximum age as for calibration 8.

1599

1600 *Discussion*. Pilumnid genera and species show a considerable anatomical disparity. Although

1601 previous molecular and mitogenomic phylogenetic studies have suggest that Pilumnidae, in the

1602 broad sense, is monophyletic (Lai et al., 2014; Duan et al., 2022), the monophyly of Pilumnidae as

1603 currently envisioned is questionable (Wolfe et al., 2022). Despite this, *Glabropilumnus* can be

1604 referred to the subfamily Pilumninae in the strict sense, and thus to the crown Pilumnidae (Poore

1605 and Ahyong, 2023).

1606   There are several extinct genera of Eocene crabs from Europe and USA that have been

1607 tentatively assigned to Pilumnidae and, specifically, the subfamily Pilumninae. However, the only

1608 extant pilumnid genus with Eocene representatives known is *Glabropilumnus* Balss, 1932. Besides

1609 *G. trispinosus, G. bizzarinii* Beschin, Busulini, and Tessier, in Beschin et al., 2019, is known from

1610 the upper Eocene (Priabonian) of Italy, together with *G.* cf. *granulatus* De Angeli, Garassino &

1611 Ceccon, 2010b, which was previously known solely from the lower Oligocene of Italy (Beschin et

1612 al., 2019). As such, the most recent common ancestor of Pilumnidae and all of its descendants must

1613 be pre-Eocene.

1614

1615

1616 **33. Brachyura: Eubrachyura: Goneplacoidea: Goneplacidae (crown)**

1617 *Fossil specimens*. *Carcinoplax temikoensis* Feldmann and Maxwell, 1990. New Zealand Geological

1618 Survey, Lower Hutt, New Zealand. Holotype AR 1943, a complete dorsal carapace and a cheliped.

1619

1620   ***Phylogenetic justification.*** *Carcinoplax temikoensis* can be assigned to *Carcinoplax* Milne

1621   Edwards, 1852, based on the transversely rectangular carapace, the forward directed, narrow orbits,

1622   and the presence of three spines on the anterolateral margin, including the one forming the outer

1623   orbital angle (Feldmann and Maxwell, 1990; Ng and Castro, 2020). *Carcinoplax* is nested within

1624   Goneplacidae (Karasawa and Kato, 2003).

1625

1626   ***Minimum age***. 34.6 Ma.

1627

1628   ***Tip maximum age.*** 39.1 Ma.

1629

1630   ***Soft maximum age.*** 100.9 Ma.

1631

Age justification. The type material of *Carcinoplax temikoensis* was collected from the lower part

(unit 2) of the Island Sandstone at Te Miko, locality K30/D7A, exposed along the coast between

Perpendicular Point and Punakaiki, north of Greymouth, Westland, New Zealand, ascribed to the

upper Eocene (Kaiatan to Runangan) based on foraminiferal assemblages (Feldmann and Maxwell,

1990). The Kaiatan (39.1–36.7 Ma) and Runangan (36.7–34.6 Ma) stages in New Zealand roughly

correspond to the global uppermost Bartonian (41.2–37.71 Ma) and Priabonian (37.71–33.9 Ma)

ages (Raine et al., 2015). Given the unclear exact age of the layers bearing the type material of *C.*

temikoensis, for the time being, we bracket its tip maximum and minimum ages between the base of

the Kaiatan and the top of the Runangan (i.e., 39.1–34.6 Ma).

Soft maximum age as for calibration 8.

1642

Discussion. Currently, there are 25 species referred to *Carcinoplax* (Ng and Castro, 2020). As

previously discussed for Euryplacidae (see above), the superfamily Goneplacoidea is an

anatomically disparate and problematic brachyuran groups in terms of the clarity about its internal

phylogenetic affinities (Ng et al., 2008; Castro et al., 2010). Despite this, *Carcinoplax* fits well

within Goneplacidae and thus belongs to the crown Goneplacoidea.

There are over a dozen species of *Carcinoplax* known from fossils, from which only *C.*

temikoensis occur in the Eocene. Extinct, monotypic genera known from the Eocene and presumably

with goneplacid affinities include *Amydrocarcinus* Schweitzer, Feldmann, González-Barba, and

Vega, 2002, from the Bartonian of Mexico, and *Gonioplacoides* Quayle and Collins, 2012, from the

upper Eocene of UK. The presence of extinct and extant goneplacid genera in the Eocene indicate

that the most recent common ancestor of Goneplacidae and all of its descendants must be at least

early Eocene in age, but, as previously seen with most other families already discussed, its origins

might lie in the Paleocene (Wolfe et al., 2022).

1656

1657

1658 ## 34. Brachyura: Eubrachyura: Xanthoidea: Pseudorhombilidae (crown)

1659 ***Fossil specimens.*** *Pseudorhombila patagonica* Glaessner, 1933. Collections of the British Museum.

1660 Holotype In. 28031, a complete dorsal carapace.

1661

Phylogenetic justification. The genus *Pseudorhombila* H. Milne Edwards, 1834 [in 1834-1840], has

been recovered nested within a monophyletic Xanthoidea, but its familial placement has been

included either within Pseudorhombilidae Alcock, 1900 (e.g., Thoma et al., 2014), or within the

subfamily Pseudorhombilinae under the family Panopeidae Ortmann, 1893, and besides other

subfamilies such as Panopeinae, Melybiinae, and an unnamed subfamily (e.g., Mendoza et al.,

2022). In our study (Wolfe et al., 2022), *Pseudorhombila* was recovered in a clade formed by

several genera typically ascribed to Pseudorhombilidae, which in turn was recovered as sister to a

clade formed by several genera ascribed to Panopeidae. As such, here we consider

Pseudorhombilidae as a discrete family.

1671

1672 ***Minimum age***. 5.333 Ma.

1673

1674 ***Tip maximum age.*** 23.03 Ma.

1675

1676 ***Soft maximum age.*** 100.9 Ma.

1677

Age justification. The type specimen of *Pseudorhombila patagonica* was collected from the so-

called Patagonian Beds, in the Santa Cruz Province, Argentina, which have been considered

Miocene in age (Glaessner, 1933, 1969). Unfortunately, it comes from an old collection, and its

precise lithostratigraphic and chronostratigraphic context is unclear, hence the broad minimum and

soft maximum age bracket.

Soft maximum age as for calibration 8.

1684

Discussion. Pseudorhombilidae Alcock, 1900, is a small family of xanthoid crabs with eight extant

genera and over a dozen species known to date (Ng et al., 2008; De Grave et al., 2009). In our study

(Wolfe et al., 2022), species within Pseudorhombilidae form a monophyleic group sister to

Panopeidae. To date, the only known fossil referable to the family is *P. patagonica*, a member of the

extant genus *Pseudorhombila* H. Milne Edwards, 1834 [in 1834-1840], therefore it is our selection

of fossil for calibration.

1691

1692

1693   ### 35. Brachyura: Eubrachyura: Xanthoidea: Panopeidae (crown)

1694   ***Fossil specimens.*** *Panopeus incisus* Beschin, Busulini, De Angeli, and Tessier, 2007. Museo di

1695   Archeologia e Scienze Naturali "G. Zannato" di Montecchio Maggiore, Vicenza, northern Italy.

1696   Holotype MCZ 2009, a nearly complete dorsal carapace.

1697

Phylogenetic justification. *Panopeus incisus* was ascribed to *Panopeus* H. Milne Edwards, 1834 [in

1834-1840], based on the wide, subhexagonal carapace, the broad and mesially notched front, the

sub-ovate orbits bearing two orbital fissures, and the well-developed anterolateral margin with four

spines excluding the outer orbital spine (Schweitzer, 2000; Beschin et al., 2007). *Panopeus* is an

extant genus, the type genus of the family Panopeidae Ortmann, 1893, and together with other

panopeid species forms a clade nested within Xanthoidea (Jennings et al., 2021; Mendoza et al.,

2022Thoma et al., 2014; Wolfe et al., 2022).

1705

1706   ***Minimum age***. 47.8 Ma.

1707

1708   ***Tip maximum age.*** 56.0 Ma.

1709

1710   ***Soft maximum age.*** 100.9 Ma.

1711

1712   ***Age justification***. The type material of *Panopeus incisus* comes from lower Eocene rocks cropping

1713   out at the Contrada Gecchelina of the Monte di Malo in Vicenza, northern Italy (Beschin et al.,

1714   2007, see also Beschin et al., 2016a). Age as for calibration 13.

1715 Soft maximum age as for calibration 8.

1716

Discussion. Panopeidae is a speciose group of xanthoid crabs with an early fossil record extending

back into the Paleogene Period. Among the extant panopeid genera, only *Panopeus*, *Lophopanopeus*

Rathbun, 1898, and *Metopocarcinus* Stimpson, 1860, have putative occurrences in the Eocene

(Beschin et al., 2016b; Pasini et al., 2019), with *Panopeus* being represented by multiple species

(e.g., Schweitzer, 2000; Vega et al., 2008; Gatt and De Angeli, 2010; Beschin et al., 2013; Beschin

et al., 2016a; Beschin et al., 2018, and references therein). While there are numerous extinct

presumed panopeid genera with Eocene representatives, only *Pakicarcinus* Schweitzer, Feldmann

and Gingerich, 2004, and *Glyphithyreus* Reuss, 1859, have pre-Eocene (upper? Paleocene)

occurrences (Collins and Morris, 1978; Charbonnier et al., 2013; Schweitzer et al., 2016b).

The presence of numerous species of extinct and extant panopeid genera in the Eocene and

apparently the late Paleocene, suggest that the origins of the family are rooted into the early

Paleogene.

1729

1730

1731   **36. Brachyura: Eubrachyura: Xanthoidea: Xanthidae (crown)**

1732 ***Fossil specimens.*** *Phlyctenodes edwardsi* Beschin, Busulini, Tessier, and Zorzin, 2016a. Museo di

1733 Storia naturale di Verona. Holotype VR 94284, a complete dorsal carapace.

1734

Phylogenetic justification. The genus *Phlyctenodes* Milne-Edwards, 1862a is ascribed to Xanthidae,

specifically the subfamily Actaeinae Alcock, 1898, based on the ovate and wide carapace with

poorly defined dorsal regions, the large and rimmed sub-circular orbits lacking orbital fissures, the

anterolateral margins moderately convex and bearing several small tubercles or spines, the smooth

posterolateral margins nearly straight to weakly concave, the overall fronto-orbital margin

configuration, a front that is wide and bearing four or more tubercles, and the overall outline of the

frontal and anterolateral margins forming a wide arch (Busulini et al., 2006; Beschin et al., 2016a;

Schweitzer et al., 2021d).The monophyletic status of the family Xanthidae MacLeay, 1838b, has

been corroborated by several molecular and total evidence phylogenetic studies (e.g., Lai et al.,

2011; Tsang et al., 2014; Mendoza et al., 2022; Wolfe et al., 2022).

1745

1746 ***Minimum age***. 47.8 Ma.

1747

1748   ***Tip maximum age.*** 56.0 Ma.

1749

1750   ***Soft maximum age.*** 100.9 Ma.

1751

1752   ***Age justification***. *Phlyctenodes edwardsi* is known from its type locality in the area of Rama di

1753   Bolca (Verona), NE Italy, in calcareous rocks of the 'Calcari Nummulitici' that have been ascribed

1754   to the lower Eocene (Ypresian) (Beschin et al., 2016a). Age as for calibration 32.

1755   Soft maximum age as for calibration 8.

1756

Discussion. *Phlyctenodes* is an extinct genus with most of its species occurring in the Eocene of

Europe (Busulini et al., 2006). We chose *P. edwardsi* as our calibration point given its Ypresian

(56.0–47.8 Ma) age, which is similar to the age assigned to *P. multituberculatus* Beschin, Busulini,

De Angeli, and Tessier, 2007, and *P. nicolisi* Bittner, 1884, from the Ypresian of Italy. *Phlyctenodes*

tuberculosus Milne-Edwards, 1862a, from the middle–upper Eocene (41.2–47.8 Ma) of France and

Italy, was used by Tsang et al. (2014), Wolfe et al. (2019), and Mendoza et al. (2022) as their

calibration points. Beschin et al. (2016a) reported this species also from the Ypresian of Italy, in the

Bolca region from were *P. edwardsi* originated. As such, given the numerous species of

Phlyctenodes now know from the Ypresian, any of those species would represent equally

parsimonious choices of calibration points, and all would indicate that the most recent common

ancestor of xanthid crabs and all of its descendants must have originated in the pre-Eocene.

1768

1769

ACKNOWLEDGEMENTS

Special thanks to Matúš Hyžný, Àlex Ossó, Barry van Bakel, Fernando Ferratges, Carrie

Schweitzer, and Orangel Aguilera, for facilitating us bibliographic references and for valuable

discussion about the systematic affinities and potential ages of several taxa here investigated. We

thank the National Science Foundation (NSF), DEB grants #1856667 and #1856679 (USA) for

supporting this work. This is contribution #X from the Coastlines and Oceans Division of the

Institute of Environment at Florida International University.

1777

REFERENCES

Agnew, J., 2001, Taxonomy, taphonomy, and paleoecology of the Plio-Pleistocene shell beds at 101
Ranch Pit, Okeechobbe County, Florida [Masters of Science thesis: Gainesville, Florida,
USA, 219 p.

Aguilera, O., Martins, M. V. A., Linhares, A. P., Kütter, V. T., and Coletti, G., 2022,
Palaeoenvironment of the Miocene Pirabas Formation mixed carbonate–siliciclastic deposits,
Northern Brazil: Insights from skeletal assemblages: Marine and Petroleum Geology, v. 145,
p. 105855.

Ahyong, S. T., Lai, J. C. Y., Sharkey, D., Colgan, D. J., and Ng, P. K. L., 2007, Phylogenetics of the
brachyuran crabs (Crustacea: Decapoda): The status of Podotremata based on small subunit
nuclear ribosomal RNA: Molecular Phylogenetics and Evolution, v. 45, no. 2, p. 576–586.

Ahyong, S. T., Baba, K., Macpherson, E., and Poore, G. C., 2010, A new classification of the
Galatheoidea (Crustacea: Decapoda: Anomura): Zootaxa, v. 2676, no. 1, p. 57–68.

Ahyong, S. T., 2019, The cymonomid crabs of New Zealand and Australia (Crustacea: Brachyura:
Cyclodoripoida): Records of the Australian Museum, v. 71, no. 2, p. 33-69.

Alcock, A., 1898, Materials for a carcinological fauna of India. No. 3. The Brachyura Cyclometopa.
Part I. The family Xanthidae: Journal of the Asiatic Society of Bengal, v. 67, no. 2, p. 67–
233.

Alcock, A., 1900, Materials for a carcinological fauna of India, No. 6. The Brachyura Catometopa,
or Grapsoidea: Journal of the Asiatic Society of Bengal, v. 69, no. 2–3, p. 279–456.

Anderson, J. L., and Feldmann, R. M., 1995, *Lobocarcinus lumacopius* (Decapoda: Cancridae), a
new species of cancrid crab from the Eocene of Fayum, Egypt: Journal of Paleontology, v.
69, no. 5, p. 922–932.

Ando, Y., Kishimoto, S., and Kawano, S., 2016, Two new species of Thalassina (Decapoda,
Thalassinidae) from the Miocene of Japan: Neues Jahrbuch für Geologie und Paläontologie-
Abhandlungen, p. 107–117.

Antoine, P.-O., Abello, M. A., Adnet, S., Sierra, A. J. A., Baby, P., Billet, G., Boivin, M., Calderón,
Y., Candela, A., and Chabain, J., 2016, A 60-million-year Cenozoic history of western
Amazonian ecosystems in Contamana, eastern Peru: Gondwana Research, v. 31, p. 30-59.

Antonioli, L., de Araújo Távora, V., and Dino, R., 2015, Palynology of carcinolites and limestones
from the Baunilha Grande Ecofacies of the Pirabas Formation (Miocene of Pará state,
northeastern Brazil): Journal of South American Earth Sciences, v. 62, p. 134–147.

Armstrong, A., Nyborg, T., Bishop, G. A., Ossó-Morales, À., and Vega, F. J., 2009, Decapod
crustaceans from the Paleocene of Central Texas, USA: Revsta Mexicana de Ciencias
Geológicas, v. 26, no. 3, p. 745–763.

Artal, P., 2008, *Uca miocenica* (Crustacea, Decapoda), nueva especie del Mioceno de la Prov. de
Barcelona (Cataluña, España): Scripta Musei Geologici Seminarii Barcinonensis
(Palaeontologica), v. 6, p. 3–18.

Artal, P., and Hyžný, M., 2016, An appraisal of *Typilobus* Stoliczka, 1871 (Crustacea, Brachyura,
Leucosioidea), with description of a new family and genus: Zootaxa, v. 4117, no. 3, p. 387-
398.

Artal, P., and Onetti, A., 2017, *Tavernolesia*, new genus (Crustacea, Decapoda, Brachyura), from
the Eocene of the Iberian Peninsula: Batalleria, v. 24, p. 6–12.

Aslanova, S., and Dschafarova, J., 1975, Some finds of Decapoda from the Tertiary deposits of
Azerbaijan: Akademia nauk Azerbaidzhanskoi Soviet Socialist Republic Doklady, v. 31, no.
7, p. 41–46.

Bachmayer, F., and Wagner, R., 1957, *Geryon heimertingensis* n. sp., eine Krabbbe aus dem
Chattian der Erdölbohrung Heimertingen (bayerische Vorlandmolasse): Paläontologische
Zeitschrift, v. 31, no. 1/2, p. 99–102.

Bachmayer, F., and Mundlos, R., 1968, Die tertiären Krebse von Helmstedt bei Braunschweig,
Deutschland: Annalen des Naturhistorischen Museums in Wien, p. 649–692.

Balss, H., 1932, Über einige systematisch interessante Xanthidae (Crustacea, Decapoda, Brachyura)
der Harmsschen Reisen nach dem Sundaarchipel: Zeitschrift für Wissenschaftliche Zoologie,
v. 142, no. 4, p. 510–519.

Batllori, J., and García, J. J., 1997, Malacofauna d'un manglar del Miocè de Bellaterra (depressió del
Vallès-Penedès, Barcelona): Butlletí de la Institució Catalana d'Història Natural, p. 15–21.

Beccaro, L., 2003, Revisioni stratigrafiche nel Paleocene del Veneto occidentale [Dottorato di
Ricerca in Scienze della Terra, Ciclo XVI: Unpublished Ph. D. Thesis, Università degli Studi
di Padova, Padova., 200 p.

Bell, T., 1858, A monograph of the fossil malacostracous Crustacea of Great Britain. Part I.
Crustacea of the London Clay: Monograph of the Palaeontographical Society, v. 10, p. i-viii,
1–44, 11 pls.

Bell, T., 1863, A monograph of the Fossil Malacostracous Crustacea of Great Britain. Part II. Crustacea of the Gault and Greensand: Monograph of the Palaeontographical Society, v. 14, p. vii + 40 pp., pls. 41–11.

Bernot, J. P., Owen, C. L., Wolfe, J. M., Meland, K., Olesen, J., and Crandall, K. A., 2022, Major revisions in pancrustacean phylogeny with recommendations for resolving challenging nodes: bioRxiv, p. 2022.2011. 2017.514186.

Beschin, C., Busulini, A., De Angeli, A., Tessier, G., and Ungaro, S., 1998, Crostacei eocenici di "Cava Rossi" presso Monte di Malo (Vicenz – Italia settentrionale): Studi Trentini di Scienze Naturali – Acta Geologica v. 73, no. 1996, p. 7–34.

Beschin, C., De Angeli, A., and Checchi, A., 2001, Crostacei decapodi associati a coralli della "Formazione di Castelgomberto"(Oligocene)(Vicenza—Italia settentrionale): Studi e Ricerche, Associazione Amici del Museo, Museo Civico ''G. Zannato'', Montecchio Maggiore (Vicenza), v. 2001, p. 13-30.

Beschin, C., Busulini, A., De Angeli, A., and Tessier, G., 2002, Aggiornamento ai crostacei eocenici di cava "Main" di Arzignano (Vicenza-Italia settentrionale) (Crustacea, Decapoda): Studi e Ricerche, Associazione Amici del Museo—Museo Civico "G. Zannato"(Montecchio Maggiore), v. 2002, p. 7–28.

Beschin, C., Busulini, A., De Angeli, A., and Tessier, G., 2007, I decapodi dell'Eocene inferiore di Contrada Gecchelina (Vicenza, Italia settentrionale) (Anomura e Brachyura): Museo di archeologia e scienze naturali G. Zannato, p. 1–84.

Beschin, C., De Angeli, A., and Zorzin, R., 2009, Crostacei fossili del Veneto: una inedita fauna eocenica dei Lessini orientali (Monte Serea di San Giovanni Ilarione, Verona), con descrizione di tre nuove specie: Bollettino del Museo civico di Storia naturale di Verona, v. 33, p. 59–83.

Beschin, C., and De Angeli, A., 2011, *Baldoplax bonizzatoi* gen. nov., sp. nov., (Crustacea, Decapoda, Euryplacidae) dell'Eocene di Ferrara di Monte Baldo (Verona, Italia settentrionale): Studi e Ricerche - Associazione Amici del Museo- Museo Civico "G. Zannato", v. 18, p. 5–10.

Beschin, C., Busulini, A., and Tessier, g., 2013, Crostacei medio-eocenici della "Pietra di Nanto" (Monti Berici, Vicenza- Italia settentrionale): Lavori-Società Veneziana di Scienze Naturali, v. 38, p. 111–146.

Beschin, C., Busulini, A., and Tessier, G., 2015, Nuova segnalazione di crostacei associati a coralli nell'Eocene inferiore dei Lessini orientali (Vestenanova–Verona): Lavori Società veneziana di Scienze naturali, v. 40, p. 47–109.

Beschin, C., Busulini, A., Tessier, G., and Zorzin, R., 2016a, I crostacei associati a coralli nell'Eocene inferiore dell'area di Bolca: Verona e Vicenza, Italia nordorientale, Memorie del Museo civico di storia naturale di Verona, 189 p.:

Beschin, C., De Angeli, A., Checchi, A., and Zarantonello, G., 2016b, Crostacei Decapodi del" tufo a *Lophoranina*"(Luteziano inferiore) della Valle del Chiampo (Vicenza, Italia nordorientale), Montecchio Maggiore (Vicenza), Museo di Archeologia e Scienze Naturali" G. Zannato", 92 p.:

Beschin, C., Busulini, A., Fornaciari, E., Papazzoni, C. A., and Tessier, G., 2018, La fauna di crostacei associati a coralli dell'Eocen Superiore di Campolongo di Val Liona (Monti Berici, Vicenza, Italia Nordorientale): Bolletino del Museo di Storia Naturale di Venezia, v. 69, p. 129–215.

Beschin, C., Busulini, A., Tessier, G., and Zorzin, R., 2019, La fauna di crostacei dell'Eocene superiore di Parona di Verona (Italia nordorientale): nuovi ritrovamenti: Bollettino del Museo di Storia Naturale di Venezia, v. 70, p. 71–142.

Betancort, J. F., Lomoschitz Mora-Figueroa, A., and Meco, J., 2014, Mio-Pliocene crustaceans from the Canary Islands, Spain: Rivista Italiana di paleontologia e Stratigrafia, v. 120, no. 3, p. 337–349.

Beurlen, K., 1958, Contribuição à paleontologia do estado do Para: Crustáceos Decápodos da Formação Pirabas: Boletim do Museu Paraense Emilio Goeldi, (nova série) (Geologia), v. 5, p. 1–48.

Bishop, G. A., and Williams, A. B., 2000, Fossil crabs from tepee buttes, submarine seeps of the late Cretaceous Pierre Shale, South Dakota and Colorado, USA: Journal of Crustacean Biology, v. 20, no. 5, p. 286-300.

Bittner, A., 1893, Decapoden des pannonischen Tertiärs: Sitzungsberichte der kaiserlichen Akademie der Wissenschaften in Wien, v. 102, p. 10–37.

Blow, W. C., and Manning, R. B., 1996, Preliminary descriptions of 25 new decapod crustaceans from the middle Eocene of the Carolinas, U.S.A.: Tulane Studies in Geology and Paleontology, v. 29, no. 1, p. 1–26.

Blow, W. H., Late Middle Eocene to Recent planktonic foraminiferal biostratigraphy, *in*
Proceedings Proceedings of the first international conference on planktonic microfossils
(Geneva, 1967)1969, Volume 1, Leiden, EJ Brill, p. 199–422.

Boucot, A. J., and Poinar Jr, G. O., 2010, Fossil Behavior Compendium, Boca Raton, CRC Press.

Brito, I. M., 1972, Contribuição ao Conhecimento dos Crustáceos Decápodos da Formação Pirabas.
II — O Gênero *Uca* (Brachyura-Ocypodidae): Anais da Academia Brasileira de Ciências, v.
44, no. 1, p. 95–98.

Buchs, D. M., Irving, D., Coombs, H., Miranda, R., Wang, J., Coronado, M., Arrocha, R., Lacerda,
M., Goff, C., and Almengor, E., 2019, Volcanic contribution to emergence of Central
Panama in the Early Miocene: Scientific reports, v. 9, no. 1, p. 1-16.

Busulini, A., Tessier, G., Beschin, C., and De Angeli, A., 2003, *Boschettia giampietroi*, nuovo
genere e specie di Portunidae (Crustacea, Decapoda) dell'Eocene medio della Valle del
Chiampo (Vicenza, Italia settentrionale): Studi e Ricerche–Associazione Amici del Museo–
Museo civico "G. Zannato"–Montecchio Maggiore (Vicenza), p. 13–18.

Busulini, A., Tessier, G., and Beschin, C., 2006, The genus *Phlyctenodes* Milne Edwards, 1862
(Crustacea: Decapoda: Xanthidae) in the Eocene of Europe: Revista mexicana de ciencias
geológicas, v. 23, no. 3, p. 350–360.

Cahuzac, B., and Poignant, A., 1997, Essai de biozonation de l'Oligo-Miocène dans les bassins
européens à l'aide des grands foraminifères néritiques: Bulletin de la Société géologique de
France, v. 168, no. 2, p. 155–169.

Cahuzac, B., and Londeix, L., 2012, Biostratigraphie appliquée à l'étage Aquitanien [Contribution à
la connaissance de l'étage Aquitanien, n° 8] Bulletin de la Société Linnéenne de Bordeaux,
no. 2, p. 213–226.

Carreño, A. L., and Cronin, T. M., 1993, Middle Eocene Ostracoda from Baja California Sur,
Mexico: Journal of Micropalaeontology, v. 12, no. 2, p. 141–153.

Casadio, S., Parras, A., Marenssi, S., and Griffin, M., 2001, Edades 87Sr/86Sr de *Crassostrea*?
hatcheri Ortmann (Bivalvia, Ostreoidea) en el "Patagoniano" de Santa Cruz, Argentina,
Reunión de Comunicaciones de la Asociación Paleontológica Argentina. 29–30 de
noviembre, 2001, Volume 38, Ameghiniana, Suplemento, p. 30R.

Casadío, S., Guerstein, G., Marenssi, S., Santillana, S., Feldmann, R., Parras, A., and Montalvo, C.,
2000, Evidencias para una edad oligocena de la Formación Centinela, suroeste de Santa
Cruz, Argentina: Ameghiniana, v. 37, no. 4, p. 71R.

Casadío, S., De Angeli, A., Feldmann, R. M., Garassino, A., Hetler, J. L., Parras, A., and
Schweitzer, C. E., 2004, New decapod crustaceans (Thalassinidea, Galatheoidea, Brachyura)
from the middle Oligocene of Patagonia, Argentina: Annals of Carnegie Museum, v. 73, p.
25–47.

Casadío, S., Feldmann, R. M., Parras, A., and Schweitzer, C. E., 2005, Miocene fossil Decapoda
(Crustacea: Brachyura) from Patagonia, Argentina, and their paleoecological setting: Annals
of Carnegie Museum, v. 74, no. 3, p. 151–188.

Casanovas-Vilar, I., DeMiguel, D., Galindo, J., Robles, J. M., Garcés, M., and Cabrera, L., 2011,
The continental Burdigalian (Early Miocene) of the Vallès-Penedès Basin (Catalonia, Spain),
in Pérez-García, A., Gascó, F., Gasulla, J. M., and Escaso, F., eds., Viajando a Mundos
Pretéritos: Morella, Ayuntamiento de Morella, p. 93–100.

Casanovas-Vilar, I., Madern, A., Alba, D. M., Cabrera, L., García-Paredes, I., Van den Hoek
Ostende, L. W., DeMiguel, D., Robles, J. M., Furió, M., and van Dam, J., 2016, The
Miocene mammal record of the Vallès-Penedès Basin (Catalonia): Comptes Rendus Palevol,
v. 15, no. 7, p. 791–812.

Castro, P., Ng, P. K., and Ahyong, S. T., 2004, Phylogeny and systematics of the Trapeziidae Miers,
1886 (Crustacea: Brachyura), with the description of a new family: Zootaxa, v. 643, no. 1, p.
1–70.

Castro, P., Guinot, D., and Ng, P. K. L., 2010, A new family for *Sotoplax robertsi* Guinot, 1984,
with a diagnosis and key to the Goneplacoidea MacLeay, 1838 (Crustacea: Decapoda:
Brachyura): Zootaxa, v. 2356, no. 1, p. 36–56.

Castro, P., and Ng, P. K. L., 2010, Revision of the family Euryplacidae Stimpson, 1871 (Crustacea:
Decapoda: Brachyura: Goneplacoidea): Zootaxa, v. 2375, p. 1–130.

Ceccon, L., and De Angeli, A., 2019, Un nuovo costaceo Percnidae dell'Oligocene di Sant'Urbano,
Vicenza, Italia settentrionale: Bolletino del Museo Civico di Storia Naturale di Verona, v.
43, p. 17–22.

Charbonnier, S., Garassino, A., Pasini, G., Métais, G., Merle, D., Bartolini, A., Brohi, I. A., Solangi,
S. H., Lashari, R. A., Welcomme, J.-L., and Marivaux, L., 2013, Early Paleogene decapod
crustaceans from the Sulaiman and Kirthar Ranges, Pakistan: Annales de Paléontologie, v.
99, no. 2, p. 101-117.

Chen, J., Xing, Y., Yao, W., Xu, X., Zhang, C., Zhang, Z., and Liu, Q., 2019, Phylomitogenomics
reconfirm the phylogenetic position of the genus *Metaplax* inferred from the two grapsid
crabs (Decapoda: Brachyura: Grapsoidea): Plos one, v. 14, no. 1, p. e0210763.

Choffat, P. L., 1898, Recueil d´études paléontologiques sur la Faune Crétacique du Portugal, II
Section des Travaux Géologiques du Portugal, Lisbonne, p. 45.

Cohen, K. M., Finney, S. C., Gibbard, P. L., and Fan, J.-X., 2013; updated 2023, The ICS
International Chronostratigraphic Chart: Episodes, v. 36, p. 199–204.

Collins, J. S. H., and Morris, S. F., 1978, New Lower Tertiary crabs from Pakistan: Palaeontology,
v. 21, p. 957-981.

Collins, J. S. H., and Donovan, S. K., 1997, Some new crab records (Crustacea: Decapoda) from the
late Pleistocene Port Morant Formation of southeast Jamaica: Bulletin of the Mizunami
Fossil Museum, v. 24, no. (for 1997), p. 73–77.

Collins, J. S. H., and Jakobsen, S. L., 2003, New crabs (Crustacea, Decapoda) from the Eocene
(Ypresian/Lutetian) Lillebælt Clay Formation of Jutland, Denmark: Bulletin of the
Mizunami Fossil Museum, v. 30, p. 63–96.

Collins, J. S. H., Lee, C., and Noad, J., 2003, Miocene and Pleistocene crabs (Crustacea, Decapoda)
from Sabah and Sarawak: Journal of Systematic Palaeontology, v. 1, no. 3, p. 187–226.

Collins, J. S. H., Mitchell, S. F., and Donovan, S. K., 2009, A new species of land crab, *Sesarma*
Say , 1817(Decapoda, Brachyura), from the Pleistocene of Jamaica: Scripta Geologica, v.
138.

Collins, J. S. H., and Donovan, S. K., 2012, Pleistocene decapod crustaceans of eastern Jamaica:
Caribbean Journal of Science, v. 46, p. 133–142.

Colosi, G., 1924, Una specie fossile de Gerionide (Decapodi brachiuri): Bolettino della Societá dei
Naturalisti in Napoli, Series 2, v. 15 (for 1923), no. 35, p. 248–255.

Cotton, L., Rivero-Cuesta, L., Franceschetti, G., Monechi, S., Iakovleva, A., Alegret, L., Dinares-
Turell, J., Hooker, J., King, C., and Fluegeman, R., 2021, Reassessing the Bartonian unit
stratotype at Alum Bay (Isle of Wight, UK): an integrated approach: Newsletters on
Stratigraphy, v. 54, no. 1, p. 17–42.

Crame, J. A., Beu, A. G., Ineson, J. R., Francis, J. E., Whittle, R. J., and Bowman, V. C., 2014, The
early origin of the Antarctic marine fauna and its evolutionary implications: PloS one, v. 9,
no. 12, p. e114743.

De Angeli, A., and Beschin, C., 1998, *Ceronnectes*, nuovo genere di brachyuro dell'Eocene di
Ungheria e italia: Lavori Società Veneziana di Scienze Naturali, v. 23, p. 87–91.

De Angeli, A., Garassino, A., and Pasini, G., 2009, New reports of anomurans and brachyurans from
the Cenozoic of Tuscany (Italy): Atti della Società italiana di scienze naturali e del museo
civico di storia naturale di Milano, v. 150, no. 2, p. 163–196.

De Angeli, A., Garassino, A., and Alberti, R., 2010a, *Eogarthambrus guinotae* n. gen. and n.
sp.(Decapoda, Brachyura, Parthenopidae) from the Eocene of Vicenza, Italy, Studies on
Brachyura: a homage to Danièle Guinot, Brill, p. 107–116.

De Angeli, A., Garassino, A., and Ceccon, L., 2010b, New report of the coral-associated decapods
from the "Formazione di Castelgomberto" (early Oligocene) (Vicenza, NE Italy): Atti della
Società italiana di Scienze naturali e del Museo civico di Storia naturale in Milano, v. 151,
no. 2, p. 145–177.

De Angeli, A., and Ceccon, L., 2012, *Eouroptychus montemagrensis* n. gen., n. sp. (Crustacea,
Decapoda, Anomura, Chirostylidae) dell'Eocene inferiore (Ypresiano) di Monte Magrè
(Vicenza, Italia settentrionale): Lavori Società veneziana di Scienze naturali, v. 37, p. 19–24.

De Angeli, A., and Ceccon, L., 2013, Tetraliidae and Trapeziidae (Crustacea, Decapoda, Brachyura)
from the Early Eocene of Monte Magrè (Vicenza, NE Italy): Natural History Sciences, v.
154, no. 1, p. 25–40.

De Angeli, A., and Ceccon, L., 2014, Nuovi Brachyura (Decapoda) dell'Eocene inferiore di Monte
Magrè (Vicenza, Italia settentrionale): Lavori Società veneziana di Scienze naturali, v. 39, p.
77–92.

De Angeli, A., and Ceccon, L., 2015, Nuovi crostacei Brachyura dell'Eocene di Monte Magrè
(Vicenza, Italia settentrionale): Lavori Società veneziana di Scienze naturali, v. 40, p. 119–
138.

De Angeli, A., 2016, Nuovi crostacei Cymonomidae (Decapoda: Brachyura) dell'Eocene dei Monti
Berici (Vicenza, Italia settentrionale): Studi Trentini di Scienze Naturali, v. 95, p. 25-32.

De Angeli, A., and Alberti, R., 2016, *Tethyscarpilius bericus* n. gen., n. sp.(Decapoda, Brachyura,
Carpiliidae) dell'Eocene superiore dei Monti Berici (Vicenza, Italia settentrionale): Lavori
Società Veneziana di Scienze Naturali, v. 41, p. 121–128.

De Angeli, A., and Caporiondo, F., 2016, Un nuovo Parthenopidae (Crustacea, Decapoda,
Brachyura) dell'Eocene inferiore dei Monti Lessini orientali (Verona− Italia settentrionale):
Lavori Società veneziana di Scienze naturali, v. 41, p. 137–144.

De Angeli, A., and Ceccon, L., 2017, Contributo ai crostacei decapodi dell'Eocene inferiore dei
Monti Lessini orientali (Italia nordorientale): Natura Vicentina, v. 20, no. 2016, p. 5–38.

De Angeli, A., and Caporiondo, F., 2019, *Alontecarcinus buratoi* n. gen., n. sp. (Decapoda,
Brachyura, Potamonidae) un nuovo crostaceo d'acqua dolce dell'Eocene (Bartoniano) di
Alonte (Monti Berici, Vicenza, Italia settentrionale): Bollettino del Museo Civico di Storia
Naturale di Verona, v. 43, no. 2019 Geologia Paleontologia Preistoria, p. 5–16.

de Bouillé, R., Paléontologie de Biarritz, *in* Proceedings Compte Rendu Travaux Congrès
Scientifique de France, 39e session, Pau1873, p. 11, pl. 14.

de Gibert, J. M., and Robles, J. M., 2005, Firmground ichnofacies recording high-frequency marine
flooding events (Langhian transgression, Vallès-Penedès Basin, Spain): Geologica Acta, v.
3, no. 3, p. 295–305.

de Gibert, J. M., Muñiz, F., Belaústegui, Z., and Hyžný, M., 2013, Fossil and modern fiddler crabs
(*Uca tangeri*: Ocypodidae) and their burrows from SW Spain: ichnologic and biogeographic
implications: Journal of Crustacean Biology, v. 33, no. 4, p. 537–551.

De Grave, S., Pentcheff, N. D., Ahyong, S. T., Chan, T. Y., Crandall, K. A., Dworschak, P. C.,
Felder, D. L., Feldmann, R. M., Fransen, C. H. J. M., Goulding, L. Y. D., Lemaitre, R., Low,
M. E. Y., Martin, J. W., Ng, P. K. L., Schweitzer, C. E., Tan, S. H., Tshudy, D., and Wetzer,
R., 2009, A classification of recent and fossil genera of decapod crustaceans.: The Raffles
Bulletin of Zoology Supplement, v. 21, p. 1–109.

De Haan, W., 1833-1850, Crustacea, *in* Siebold, P. F. v., ed., Fauna Japonica sive Descriptio
Animalium, Quae in Itinere per Japoniam, Jussu et Auspiciis Superiorum, qui Summum in
India Batava Imperium Tenent, Suscepto, Annis 1823–1830 Collegit, Notis, Observationibus
et Adumbrationibus Illustravit: Leiden, Lugduni-Batavorum, p. i-xvii, i-xxxi, ix-xvi, 1–243,
pls. A–J, L–Q, 241–255, table 242.

del Rio, C. J., Concheyro, A., and Martınez, S. A., 2011, The Maastrichtian–Danian at General Roca
(Patagonia, Argentina): a reappraisal of the chronostratigraphy and biostratigraphy of a type
locality: Neues Jahrbuch für Geologie und Paläontologie - Abhandlungen, v. 259, no. 2, p.
129–156.

Desmarest, A. G., 1822, Les crustacés proprement dits, *in* Brongniart, A., and Desmarest, A. G.,
eds., Histoire naturelle des crustacés fossiles: Paris, F.-G. Levrault, p. 67–154.

Desmarest, A. G., 1823, Malacostracés, Malacostraca. (Crust.), *in* Cuvier, F., ed., Dictionnaire des
Sciences Naturelles, dans lequel on trait Méthodiquement des Différens étres de la Nature,

considérés soit en eux-mêmes, d'après l'état actuel de nos connoissances, soit relativement a
l'utilité qu'en peuvent retirer la Médecine, l'Agriculture, le Commerce et les Arts. Suivi
d'une biographie des plus Célèbres Naturalistes. Ouvrage destiné aux médecins, aux
agriculteurs, aux commerçans, aux artistes, aux manufacturiers, et à tous ceux qui ont intérêt
à connoître les productions de la nature, leurs caractères génériques et spécifiques, leur lieu
natal, leurs propiétés et leurs usages, Volume 28: Strasbourg et Paris, F.G. Levrault et Le
Normant, p. 138–425.

Domínguez Alonso, P., 2008, Nuevo cangrejo violinista (género *Uca*, Ocypodidae) en el Plio-
Pleistoceno del litoral pacífico de Honduras: Ameghiniana, v. 45, no. 4, p. 663–676.

Duan, X., Dong, X., Li, J., Lü, J., Guo, B., Xu, K., and Ye, Y., 2022, The complete mitochondrial
genome of *Pilumnopeus makianus* (Brachyura: Pilumnidae), novel gene rearrangements, and
phylogenetic relationships of Brachyura: Genes, v. 13, no. 11, p. 1943.

Dutton, A. L., Lohmann, K. C., and Zinsmeister, W. J., 2002, Stable isotope and minor element
proxies for Eocene climate of Seymour Island, Antarctica: Paleoceanography, v. 17, no. 2, p.
6-1–6-13.

Etayo-Serna, F., 1979, Zonation of the Cretaceous of central Colombia by ammonites, Bogotá,
Colombia, Ministerio de Minas y Energia, Instituto Nacional de Investigaciones Geológico-
Mineras, 186 p.:

Evans, N., 2018, Molecular phylogenetics of swimming crabs (Portunoidea Rafinesque, 1815)
supports a revised family-level classification and suggests a single derived origin of
symbiotic taxa: PeerJ, 6, p. e4260.

Eydoux, F., 1835, Nouvelle espèce de *Gélasime*: Magasin de Zoologie, v. 5, no. 7, p. 29–32.

Fabricius, J., 1781, Species insectorum exhibentes eorum dffirentias specificas, sryonrya auctorum,
loca natalia, rnetarnorphosis adjectis obseruationibus, descriptionibus, Hamburg and
Cologne, Ernest Bohnii, 552 p.:

Fabricius, J., 1798, Supplementatum Entomologiae systematicae, Hafniae, Proft et Storch, 572 p.:

Farris, D. W., Cardona, A., Montes, C., Foster, D., and Jaramillo, C., 2017, Magmatic evolution of
Panama Canal volcanic rocks: A record of arc processes and tectonic change: PloS one, v.
12, no. 5, p. e0176010.

Feldmann, R. M., and Wilson, M. T., 1988, Eocene decapod crustaceans from Antarctica:
Geological Society of America Memoir, v. 169, p. 465–488.

Feldmann, R. M., and Maxwell, P. A., 1990, Late Eocene decapod Crustacea from North Westland,
South Island, New Zealand: Journal of Paleontology, v. 64, no. 5, p. 779–797.

Feldmann, R. M., 1993, Additions to the fossil decapod crustacean fauna of New Zealand: New
Zealand Journal of Geology and Geophysics, v. 36, no. 2, p. 201-211.

Feldmann, R. M., 1994, *Antarctomithrax thomsoni*, a new genus and species of crab (Brachyura;
Majidae) from the La Meseta Formation (Eocene) of Seymour Island, Antarctica: Journal of
Paleontology, v. 68, no. 1, p. 174–176.

Feldmann, R. M., Casadío, S., Chirino-Galvez, L., and Aguirre-Urreta, M., 1995, Fossil decapod
crustaceans from the Jagüel and Roca formations (Maastrichtian–Danian) of the Neuquén
Basin, Argentina: Journal of Paleontology, v. 69, no. S43, p. 1-22.

Feldmann, R. M., Bice, K. L., Hopkins, C. S., Salva, E. W., and Pickford, K., 1998, Decapod
crustaceans from the Eocene Castle Hayne Formation, North Carolina: Paleoceanographic
implications: Memoir (The Paleontological Society), v. 48, p. 1–28.

Feldmann, R. M., Schweitzer, C. E., and Encinas, A., 2005, New decapods from the Navidad
Formation (Miocene) of Chile: Journal of Crustacean Biology, v. 25, p. 427–449.

Feldmann, R. M., and Schweitzer, C. E., 2009, Revision of Jurassic Homoloidea De Haan, 1839,
from the Ernstbrunn and Štramberk limestones, Austria and the Czech Republic: Annalen
des Naturhistorischen Museums in Wien. Serie A für Mineralogie und Petrographie,
Geologie und Paläontologie, Anthropologie und Prähistorie, p. 183-205.

Feldmann, R. M., Schweitzer, C. E., and Encinas, A., 2010, Neogene decapod Crustacea from
Southern Chile: Annals of Carnegie Museum v. 78, no. 4, p. 337–366.

Feldmann, R. M., Schweitzer, C. E., Bennett, O., Franescu, O. D., Resar, N., and Trudeau, A.,
2011a, New Eocene Brachyura (Crustacea: Decapoda) from Egypt: Neues Jahrbuch fur
Geologie und Palaontologie-Abhandlungen, v. 262, no. 3, p. 1–31.

Feldmann, R. M., Schweitzer, C. E., Casadío, S., and Griffin, M., 2011b, New Miocene Decapoda
(Thalassinidea; Brachyura)from Tierra del Fuego, Argentina: Paleobiogeographic
implications: Annals of the Carnegie Museum, v. 79, no. 2, p. 91–123.

Feldmann, R. M., Schweitzer, C. E., and Karasawa, H., 2012, Part R, Revised, Volume 1, Chapter
8M: Systematic Descriptions: Infraorder Brachyura, Section Homoloida: Treatise Online, v.
52, p. 1–12.

Feldmann, R. M., and Schweitzer, C. E., 2019, Earliest known sponge crab (Bacrhyura: Dromiidae)
from the Upper Cretaceous Wenonah Formation, New Jersey, USA: Bulletin of the
Mizunami Fossil Museum, v. 45, p. 1-6.

Feldmann, R. M., Schweitzer, C. E., and Phillips, G., 2019, Paleogene Decapoda (Caridea,
Anomura, Axiidea, Brachyura) from Alabama and Mississippi, USA: Journal of Crustacean
Biology, v. 39, no. 3, p. 279–302.

Ferratges, F. A., Zamora, S., and Aurell, M., 2019, A new genus and species of Parthenopidae
MacLeay, 1838 (Decapoda: Brachyura) from the lower Eocene of Spain: Journal of
Crustacean Biology, v. 39, no. 3, p. 303-311.

Ferratges, F. A., Luque, J., Dominguez, J. L., Ossó, À., Aurell, M., and Zamora, S., 2023, in press,
The oldest dairoidid crab (Decapoda: Brachyura: Parthenopoidea) from the Eocene of Spain:
Papers in Palaeontology.

Fleming, C. A., 1981, A new grapsid crab from the upper Miocene of New Zealand: Journal of the
Royal Society of New Zealand, v. 11, no. 2, p. 103-108.

Forir, H., 1887, Contributions à l'étude du système crétacé de la Belgique. II. Études
complémentaires sur les crustacés: Annales de la Société géologique de Belgique, v. 14, p.
155-175.

Forskal, P., 1775, Descriptiones Animalium, Avium, Amphibiorum, Piscium, Insectorum, Vermium,
quae in itinere orientali observavit Petrus Forskal. Post mortem auct. ed. C. Niebuhr: Hafniae
(= Copenhagen), ex officina Mölleri.

Förster, R., and Mundlos, R., 1982, Krebse aus dem Alttertiär von Helmstedt und Handorf
(Niedersachsen): Palaeontographica (A), v. 179, p. 148–184.

Förster, R., 1986, Der erste Nachweis eines brachyuren Krebses aus dem Lias (Oberes Pliensbach)
Mitteleuropas: Mitteilungen der Bayerischen Staatssammlung für Paläontologie und
Historische Geologie, v. 26, p. 25–31.

Fraaije, R. H., Robins, C., van Bakel, B. W., Jagt, J. W., and Bachmayer, F., 2019, Paguroid
anomurans from the Tithonian Ernstbrunn Limestone, Austria–the most diverse extinct
paguroid assemblage on record: Annalen des Naturhistorischen Museums in Wien. Serie A
für Mineralogie und Petrographie, Geologie und Paläontologie, Anthropologie und
Prähistorie, v. 121, p. 257-290.

Francescone, F., Lauretano, V., Bouligand, C., Moretti, M., Sabatino, N., Schrader, C., Catanzariti,
R., Hilgen, F., Lanci, L., and Turtù, A., 2019, A 9 million-year-long astrochronological

record of the early–middle Eocene corroborated by seafloor spreading rates: Bulletin, v. 131, no. 3–4, p. 499–520.

Frantescu, O., Feldmann, R. M., and Schweitzer, C. E., 2016, Cretaceous fossil Raninoida De Haan, 1839 (Crustacea, Decapoda, Brachyura) from northeast Texas: Journal of Paleontology, v. 90, no. 6, p. 1118–1132.

Gandolfo, M. A., Nixon, K. C., and Crepet, W. L., 2008, Selection of Fossils for Calibration of Molecular Dating Models1: Annals of the Missouri Botanical Garden, v. 95, no. 1, p. 34-42.

Garassino, A., Artal, P., and Pasini, G., 2009, *Upogebia miocenica* n. sp.(Crustacea, Thalassinidea, Upogebiidae) from the Miocene of Catalonia (Spain): Atti della Società italiana di Scienze naturali e del Museo civico di Storia naturale di Milano, v. 150, no. 1, p. 61–68.

Gatt, M., and De Angeli, A., 2010, A new coral-associated decapod assemblage from the Upper Miocene (Messinian) Upper Coralline Limestone of Malta (Central Mediterranean): Palaeontology, v. 53, no. 6, p. 1315–1348.

Glaessner, M. F., 1933, I.—New tertiary crabs in the collection of the British Museum: Journal of Natural History Series 10, v. 12, no. 67, p. 1-28.

Glaessner, M. F., 1969, Decapoda, *in* (ed.), R. C. M., ed., Treatise on Invertebrate Paleontology. Part R. Arthropoda, Volume 4(2): Geological Society of America and University of Kansas Press, Lawrence, p. R399-R651.

Gomes, B. T., Aguilera, O., da Silva-Caminha, S. A. F., D'Apolito, C., Cárdenas, D., Hocking, E. P., and Lemes, K. K. B., 2023, Biostratigraphy and Paleoenvironments of the Pirabas Formation (Neogene, Pará State-Brazil): Marine Micropaleontology, p. 102218.

Gozhyk, P., Semenenko, V., Andreeva-Grigorovich, A., and Maslun, N., 2015, The correlation of the Neogene of Central and Eastern Paratethys segments of Ukraine with the International Stratigraphic Chart based on planktonic microfossils: Geologica Carpathica, v. 66, no. 3, p. 235–244.

Grimaldi, D. A., 1996, Amber: Window to the Past, American Museum of Natural History, New York.

Guerstein, G., Guler, M., and Casadío, S., 2004, Palynostratigraphy and palaeoenvironments across the Oligocene–Miocene boundary within the Centinela Formation, southwestern Argentina: Geological Society, London, Special Publications, v. 230, no. 1, p. 325-343.

Guinot, D., 1977, Propositions pour une nouvelle classification des Crustacés Décapodes
Brachyoures: Comptes rendus hebdomadaires des séances de l'Académie des sciences, v.
285, no. Série D, p. 1049–1052.

Guinot, D., Charbot-Chanona, G., and Vega, F. J., 2019, Archaeochiapasidae n. fam., a new early
Cenomanian brachyuran family from Chiapas, Mexico, new hypothesis on Lecythocaridae
Schweitzer & Feldmann, 2009, and phylogenetic implications (Crustacea, Decapoda,
Brachyura, Eubrachyura): Geodiversitas, v. 41, no. 7, p. 285–322.

Hartnoll, R. G., 1971, *Sesarma cookei* n. sp., a grapsid crab from Jamaica (Decapoda: Brachyura):
Crustaceana, v. 20, p. 257–262.

Hartzell, S. M., Schweitzer, C. E., and Feldmann, R. M., 2022, Extinction and survival of raninoid
crabs (Decapoda: Brachyura: Raninoida) from the Early Cretaceous to the present: Journal of
Crustacean Biology, v. 42, no. 4, p. ruac053.

Hegna, T. A., Luque, J., and Wolfe, J. M., 2020, The fossil record of the Pancrustacea, *in* Poore, G.,
and Thiel, M., eds., The Natural History of the Crustacea: Evolution and Biogeography of
the Crustacea, Volume 8, Volume 8, Oxford University Press, p. 21–52.

Herbst, J. F. W., 1782-1804, Versuch einer Naturgeschichte der Krabben und Krebse, nebst einer
systematischen Beschreibung ihrer verschiedenen Arten, Strals und GA Lange, 515 p.:

Hohenegger, J., Coric, S., and Wagreich, M., 2014, Timing of the middle miocene Badenian stage of
the central Paratethys: Geologica Carpathica, v. 65, no. 1, p. 55–66.

Hyzny, M., and Kroh, A., 2015, Barremian decapod crustaceans from Serre de Bleyton (Drôme, SE
France): Annalen des Naturhistorischen Museums in Wien. Serie A, Fur Mineralogie und
Petrographie, Geologie und Palaontologie, Anthropologie und Prahistorie, v. 117, p. 121.

Hyzny, M., 2016, Diversity and distribution patterns of the Oligocene and Miocene decapod
crustaceans (Crustacea: Malacostraca) of the Western and Central Paratethys: Geologica
Carpathica, v. 67, no. 5, p. 471–494.

Hyzny, M., and Dulai, A., 2021, Badenian decapods of Hungary, Institute de Geosciences,
University of Szeged.

Hyzný, M., Kovalchuk, O., Świdnicka, E., Berezovsky, A., Dumitriu, S., Grădianu, I., Stefaniak, K.,
and Barkaszi, Z., 2022, Revisiting brachyuran crabs (Malacostraca: Decapoda) from
Oligocene and Miocene fish beds of Europe: Geologica Carpathica, v. 73, no. 6, p. 579–597.

Ilyin, I. V., and Alekseev, A. S., 1998, Novye vidy krabov (Decapoda, Brachyura) iz nizhnego mela
ugo-zapadnogo Kryma: Paleontologicheskii Zhurnal, v. 1998/6, p. 46–49.

Ilyin, I. V., 2005, Cretaceous and Paleogene decapod crustaceans of the western part of Northern
Eurasia (in Russian), Moscow, Moscow State University Press, 296 p.:

Jagt, J. W. M., Van Bakel, B. W. M., Guinot, D., Fraaije, R. H., and Artal, P., 2015, Fossil
Brachyura, *in* Castro, P., Davie, P. J. F., Guinot, D., Schram, F. R., and Vaupel Klein, J. C.
v., eds., Decapoda: Brachyura (Part 2). Treatise on Zoology — Anatomy, Taxonomy,
Biology. The Crustacea, Vol. 9C-, Volume 9C-II: Leiden & Boston, Brill, p. 847–920.

James-Williamson, S. A., and Mitchell, S. F., 2012, Revised lithostratigraphy of the Coastal Group
of south-eastern St. Thomas, Jamaica: Caribbean Journal of Earth Science, v. 44, p. 9-17.

Janssen, A. W., and Muller, P., 1984, Miocene Decapoda and Mollusca from Ramsel (province of
Antwerpen, Belgium), with a new crab genus and a new cephalopod species: Scripta
Geologica, v. 75, p. 1–26.

Jaramillo, C. A., Rueda, M., and Torres, V., 2011, A palynological zonation for the Cenozoic of the
Llanos and Llanos Foothills of Colombia: Palynology, v. 35, no. 1, p. 46-84.

Jaworski, E., 1938, Gasterópodos del Cretácico inferior de Colombia: Estudios geológicos y
Paleontológicos sobre la Cordillera Oriental de Colombia, v. Parte 3, p. 109–112.

Jennings, L. A., Blakeslee, A., McCoy, K. A., Behringer, D. C., and Bojko, J., 2021, Systematic
assessment of the Panopeidae and broader Eubrachyura (Decapoda: Brachyura) using
mitochondrial genomics: Arthropod Systematics and Phylogeny, v. 79, p. 569–585.

Karasawa, H., 1992, The crab *Ozius collinsi* sp. nov. (Xanthoidea: Decapoda: Crustacea) from the
Miocene Katsuta Group, southwest Japan: Tertiary Research, v. 14, no. 1, p. 19–24.

Karasawa, H., 1998, *Typilobus kishimotoi*, a new leucosiid crab (Crustacea: Decapoda: Brachyura)
from the Miocene Katsuta Group, Japan: Proceedings of the Biolological Society of
Washington, v. 111, no. 1, p. 97–101.

Karasawa, H., 2000, *Medorippe tanabei*, a new species of Miocene dorippid crab (Crustacea:
Decapoda: Brachyura) from the Katsuta Group, West Honshu, Japan: Proceedings of the
Biological Society of Washington, v. 113, no. 3, p. 810–814.

Karasawa, H., and Kato, H., 2001, The systematic status of the genus *Miosesarma* Karasawa, 1989
with a phylogenetic analysis within the family Grapsidae and a review of fossil records
(Crustacea: Decapoda: Brachyura): Paleontological Research, v. 5, no. 4, p. 259–275.

Karasawa, H., and Kato, H., 2003, The family Goneplacidae MacLeay, 1838 (Crustacea: Decapoda:
Brachyura): Systematics, phylogeny, and fossil records: Paleontological Research, v. 7, no.
2, p. 129–151.

Karasawa, H., and Schweitzer, C. E., 2004, Revision of the genus *Glyphithyreus* Reuss, 1859

(Crustacea, Decapoda, Brachyura, Xanthoidea) and recognition of a new genus:

Paleontological Research, v. 8, no. 3, p. 143-154.

Karasawa, H., and Schweitzer, C. E., 2006, A new classification of the Xanthoidea sensu lato

(Crustacea: Decapoda: Brachyura) based on phylogenetic analysis and traditional

systematics and evaluation of all fossil Xanthoidea sensu lato: Contributions to Zoology, v.

75, no. 1/2, p. 23-73.

Karasawa, H., Schweitzer, C. E., and Feldmann, R. M., 2008, Revision of the Portunoidea

Rafinesque, 1815 (Decapoda: Brachyura) with emphasis on the fossil genera and families:

Journal of Crustacean Biology, v. 28, no. 1, p. 82–127.

Karasawa, H., Schweitzer, C. E., Feldmann, R. M., and Luque, J., 2014, Phylogeny and

classification of the Raninoida (Decapoda: Brachyura): Journal of Crustacean Biology, v. 34,

no. 2, p. 216–272.

Karasawa, H., Schweitzer, C. E., and Feldmann, R. M., 2019, Part R, Revised, Volume 1, Chapter

8T3: Systematic Descriptions: Superfamily Leucosioidea: Treatise Online, v. 115, p. 1–22.

Kennedy, W., Cobban, W., Hancock, J., and Gale, A., 2005, Upper Albian and Lower Cenomanian

ammonites from the Main Street Limestone, Grayson Marl and Del Rio Clay in northeast

Texas: Cretaceous Research, v. 26, no. 3, p. 349-428.

Kennedy, W. J., Amédro, F., Robaszynski, F., and Jagt, J. W., 2011, Ammonite faunas from

condensed Cenomanian-Turonian sections ('Tourtias') in southern Belgium and northern

France: Netherlands Journal of Geosciences, v. 90, no. 2–3, p. 209–238.

Kirby, M. X., Jones, D. S., and MacFadden, B. J., 2008, Lower Miocene srtratigraphy along the

Panama Canal and its bearing on the Central American Peninsula: PloS One, v. e2791, p. 1–

14.

Klaus, S., Magalhães, C., Salas-Gismondi, R., Gross, M., and Antoine, P.-O., 2017, Paleogene and

Neogene brachyurans of the Amazon Basin: A revised first appearance date for primary

freshwater crabs (Crustacea, Brachyura, Trichodactylidae): Crustaceana, v. , no. Special

Issue: Volume in Honour of Michael Türkay, p. 953–967.

Klompmaker, A. A., 2013, Extreme diversity of decapod crustaceans from the mid-Cretaceous (late

Albian) of Spain: Implications for Cretaceous decapod paleoecology: Cretaceous Research,

v. 41, p. 150-185.

Klompmaker, A. A., Portell, R. W., Klier, A. T., Prueter, V., and Tucker, A. L., 2015, Spider crabs
of the Western Atlantic with special reference to fossil and some modern Mithracidae: PeerJ,
v. 3:e1301.

Kocova-Veselská, M., Koci, T., and Kubajko, M., 2014, Dynomenid crabs (Decapoda, Brachyura)
and stalked barnacles (Cirripedia, Scalpelliformes) from upper Cenomanian-lower Turonian
nearshore, shallow-water strata in the Bohemian Cretaceous Basin, Czech Republic: Scripta
Geologica, v. 147, p. 49–81.

Koh, S., and Ng, P. K., 2008, A revision of the shore crabs of the genus Eriphia (Crustacea:
Brachyura: Eriphiidae): Raffles Bulletin of Zoology, v. 56, no. 2, p. 327-355.

Kovác, M., Halásová, E., Hudácková, N., Holcova, K., Hyzný, M., Jamrich, M., and Ruman, A.,
2018, Towards better correlation of the Central Paratethys regional time scale with the
standard geological time scale of the Miocene Epoch: Geologica Carpathica, v. 69, no. 3, p.
283–300.

Krijgsman, W., and Piller, W. E., 2012, Central and Eastern Paratethys, in Gradstein, F. M., Ogg, J.
G., Schmitz, M. D., and Ogg, G., eds., The Geologic Time Scale 2012: Amsterdam, Elsevier,
p. 935–937.

Lai, J. C., Ahyong, S. T., Jeng, M.-S., and Ng, P. K., 2009, Are coral-dwelling crabs monophyletic?
A phylogeny of the Trapezioidea (Crustacea: Decapoda: Brachyura): Invertebrate
Systematics, v. 23, no. 4, p. 402–408.

Lai, J. C. Y., Mendoza, J. C. E., Guinot, D., Clark, P. F., and Ng, P. K. L., 2011, Xanthidae
MacLeay, 1838 (Decapoda: Brachyura: Xanthoidea) systematics: A multi-gene approach
with support from adult and zoeal morphology: Zoologischer Anzeiger - A Journal of
Comparative Zoology, v. 250, no. 4, p. 407–448.

Lai, J. C. Y., Thoma, B. P., Clark, P. F., Felder, D. L., and Ng, P. K., 2014, Phylogeny of eriphioid
crabs (Brachyura, Eriphioidea) inferred from molecular and morphological studies:
Zoologica Scripta, v. 43, no. 1, p. 52–64.

Larghi, C., 2004, Brachyuran decapod Crustacea from the Upper Cretaceous of Lebanon: Journal of
Paleontology, v. 78, no. 3, p. 528–541.

Latreille, P. A., 1802, Histoire Naturelle, générale et particulière des Crustacés et des Insectes :
Ouvrage faisant suite aux Oeuvres de Leclerc de Buffon, et Partie du Cours complet
d'Histoire Naturelle rédigé par C. S. Sonnini, Membre de plusieurs Sociétés Savantes, Paris,
F. Dufart, xii + 467 p.:

Latreille, P. A., 1817, Les crustacés, les arachnides et les insectes, *in* Cuvier, G., ed., Le règne

animal distribué d'après son organization, pour servir de base à l'histoire naturelle des

animaux et d'introduction à l'anatomie comparée, Volume Edition 1, 3: Paris, p. xxix + 653

pp.

Latreille, P. A., 1825-1828, Encyclopédie méthodique, Histoire naturelle: Entomologie, ou histoire

naturelle des crustacés, des arachnides et des insectes, Imprimerie-Libraire, Paris, Chez Mme

veuve Agasse, v. (1, 2), 1–344 [1825]; 1345–1832 [1828] p.:

Latreille, P. A., 1829, Crustacés, arachnides et partie des insectes. (2nd Edition), *in* Cuvier, G., ed.,

Le règne animal distribué d'après son organisation, pour servir de base à l'histoire naturelle

des animaux et d'introduction à l'anatomie comparée. Nouvelle édition, revue et augmentée.,

Volume 4: Paris, Chez Déterville, Libraire; et Chez Crochard, Libraire, p. 1–584.

Leach, W. E., 1817, Monograph on the genera and species of the malacostracous family Leucosidea,

in Leach, W. E., ed., The Zoological Miscellany; being descriptions of new, or interesting

animals, Volume 3: Covent Garden and London, E. Nodder and Son, p. 17–26.

LeBlanc, J., 2021, Stratigraphic Lexicon: The Onshore Cenozoic Sedimentary Formations of The

Republic of Panama: Biosis: Biological Systems, v. 2, no. 1, p. 1–173.

Lima, D., Tavares, M., Lopes, R. T., de Araújo, O. M. O., and Aguilera, O., 2020, *Uca maracoani*

(Crustacea, Decapoda, Ocypodidae) from a Miocene paleomangrove in Brazil: A case of

evolutionary stasis among tropical American fiddler crabs: Journal of South American Earth

Sciences, v. 99, p. 102517.

Linnaeus, C. v., 1758, Systema Naturae per Regna Tria Naturae, Secundum Classes, Ordines,

Genera, Species, cum Characteribus, Differentiis, Synonymis, Locis, Holmiae, Laurentii

Salvii, iii+828 p.:

Liu, Q.-N., Tang, Y.-Y., Yang, T.-T., Li, Y.-T., and Yu, X.-M., 2021, Phylogenetic relationships of

Grapsoidea and insights into the higher phylogeny of Brachyuran: Genomics, v. 113, no. 1,

p. 429-439.

Locatelli, E. R., 2013, Preservation potential of gecarcinid land crabs (Decapoda, Brachyura,

Gecarcinidae) from San Salvador: Palaios, v. 28, no. 12, p. 867–874.

Londoño, L., Royer, D. L., Jaramillo, C., Escobar, J., Foster, D. A., Cárdenas-Rozo, A. L., and

Wood, A., 2018, Early Miocene CO_2 estimates from a Neotropical fossil leaf assemblage

exceed 400 ppm: American Journal of Botany, v. 105, no. 11, p. 1929-1937.

Lorenthey, E., and Beurlen, K., 1929, Die fossilen Decapoden der Länder der Ungarischen Krone:
Geologica Hungarica (Palaeontologica), v. 3, p. 1–421, 412 tables, 416 pls.

Lörenthey , E., 1898, Beiträge zur Decapodenfauna des Ungarischen Tertiärs: Termész Füzetek,
Budapest, v. 21, p. 1–133.

Lörenthey, E., 1907, Palaeontologiai tanulmányok a harmadkorú rákok köréből. V. Adatok Egyptom
eocenkori rákfaunájához: Mathematikai és Természettudományi Közlemények, v. 29, no. 2,
p. 195–242.

Lü, J., Xia, L., Liu, X., Ma, Y., Li, J., Ye, Y., and Guo, B., 2022, The mitochondrial genome of
Grapsus albolineatus (Decapoda: Brachyura: Grapsidae) and phylogenetic associations in
Brachyura: Scientific Reports, v. 12, no. 1, p. 1-12.

Lu, X., Gong, L., Zhang, Y., Chen, J., Liu, L., Jiang, L., Lü, Z., Liu, B., Tong, G., and Wei, X.,
2020, The complete mitochondrial genome of Calappa bilineata: the first representative from
the family Calappidae and its phylogenetic position within Brachyura. Genomics, 112(3),
pp.2516-2523.: Genomics, v. 112, no. 3, p. 2516–2523.

Luque, J., Feldmann, R. M., Schweitzer, C. E., Jaramillo, C., and Cameron, C. B., 2012, The oldest
frog crabs (Decapoda: Brachyura: Raninoida) from the Aptian of northern South America:
Journal of Crustacean Biology, v. 32, no. 3, p. 405–420.

Luque, J., 2014, A new genus and species of raninoidian crab (Decapoda, Brachyura), from the
Lower Cretaceous of Colombia, South America: Scripta Geologica, v. 147, p. 27–34.

Luque, J., 2015a, A puzzling frog crab (Crustacea: Decapoda: Brachyura) from the Early Cretaceous
Santana Group of Brazil: Frog first or crab first?: Journal of Systematic Palaeontology, v. 13,
no. 2, p. 153–166.

Luque, J., 2015b, The oldest higher true crabs (Crustacea: Decapoda: Brachyura): insights from the
Early Cretaceous of the Americas: Palaeontology, v. 58, no. 2, p. 251–263.

Luque, J., 2017, Fossil hermit and land crabs (Anomura and Brachyura) from the Quaternary of
Antigua and Bermuda: Journal of Crustacean Biology, v. 37, no. 2, p. 151–156.

Luque, J., Schweitzer, C. E., Santana, W., Portell, R. W., Vega, F. J., and Klompmaker, A. A., 2017,
Checklist of fossil decapod crustaceans from tropical America, Part I: Anomura and
Brachyura: Nauplius v. 25, p. e2017025.

Luque, J., Christy, J. H., Hendy, A. J. W., Rosenberg, M. S., Portell, R. W., Kerr, K. A., and Palmer,
A. R., 2018, Quaternary intertidal and supratidal crabs (Decapoda, Brachyura) from tropical

America and the systematic affinities of fossil fiddler crabs: Journal of Systematic Palaeontology, v. 16, no. 12, p. 1037–1055.

Luque, J., Allison, W. T., Bracken-Grissom, H. D., Jenkins, K. M., Palmer, A. R., Porter, M. L., and Wolfe, J. M., 2019a, Evolution of crab eye structures and the utility of ommatidia morphology in resolving phylogeny: bioRxiv preprint, p. 786087. https://doi.org/786010.781101/786087.

Luque, J., Feldmann, R. M., Vernygora, O., Schweitzer, C. E., Cameron, C. B., Kerr, K. A., Vega, F. J., Duque, A., Strange, M., Palmer, A. R., and Jaramillo, C., 2019b, Exceptional preservation of mid-Cretaceous marine arthropods and the evolution of novel forms via heterochrony: Science Advances, v. 5, p. eaav3875.

Luque, J., Xing, L., Briggs, D. E. G., Clark, E. G., Duque, A., Hui, J., Mai, H., and McKellar, R. C., 2021, Crab in amber reveals an early colonization of non-marine environments during the Cretaceous Sciences Advances, v. 7, p. eabj5689.

Ma, K. Y., Qin, J., Lin, C.-W., Chan, T.-Y., Ng, P. K. L., Chu, K. H., and Tsang, L. M., 2019, Phylogenomic analyses of brachyuran crabs support early divergence of primary freshwater crabs: Molecular Phylogenetics and Evolution, v. 135, p. 62–66.

MacFadden, B. J., Bloch, J. I., Evans, H., Foster, D. A., Morgan, G., Rincón, A., and Wood, A. R., 2014, Temporal Calibration and Biochronology of the Centenario Fauna, Early Miocene of Panama: The Journal of Geology, v. 122, p. 113–135.

MacLeay, W. S., 1838a, On the brachyurous decapod Crustacea brought from the Cape by Dr. Smith, *in* Smith, A., ed., Illustrations of the Annulosa of South Africa; consisting chiefly of figures and descriptions of the objects of natural history collected during an expedition into the interior of South Africa, in the years 1834, 1835, and 1836; fitted out by "The Cape of Good Hope Association for Exploring Central Africa. . .": London, Smith, Elder and Company, p. 53–71.

MacLeay, W. S., 1838b, On the brachyurous decapod Crustacea brought from the Cape by Dr. Smith, *in* Smith, A., ed., Illustrations of the Annulosa of South Africa; being a portion of the objects of natural history chiefly collected during an expedition into the interior of South Africa, under the directin of Dr. Andrew Smith, in the years 1834, 1835. and 1836; fitted out by "The Cape of Good Hope Association for Exploring Central Africa". London, Smith, Elder, and Co., p. 53–71, pp. 52 plates.

Manning, R. B., and Holthuis, L. B., 1989, Two new genera and nine new species of geryonid crabs
(Crustacea, Decapoda, Geryonidae): Proceedings of the Biological Society of Washington, v.
102, no. 1, p. 50–77.

Manning, R. B., and Holthuis, L. B., 1991, West African Brachyuran Crabs (Crustacea: Decapoda):
Smithsonian Contributions to Zoology, v. 396, p. 1–379.

Martinez, S., Ramos, M. I. F., McArthur, J. M., Del Río, C. J., and Thirlwall, M. F., 2017, Late
Burdigalian (Miocene) age for pectinids (Mollusca-Bivalvia) from the Pirabas Formation
(northern Brazil) derived from Sr-isotope (87 Sr/86 Sr) data: Neues Jahrbuch für Geologie
und Paläontologie-Abhandlungen, v. 284, no. 1, p. 57–64.

McLean, H., Hausback, B., and Knapp, J., 1987, The geology of west-central Baja California Sur,
Mexico: US Geological Survey.

Mendoza, J. C., Chan, K. O., Lai, J. C., Thoma, B. P., Clark, P. F., Guinot, D., Felder, D. L., and
Ng, P. K., 2022, A comprehensive molecular phylogeny of the brachyuran crab superfamily
Xanthoidea provides novel insights into its systematics and evolutionary history: Molecular
Phylogenetics and Evolution, v. 177, p. 107627.

Miers, E. J., 1886, Report on the Brachyura collected by H.M.S. Challenger during the years 1873–
1876 *in* Murray, J., ed., Report of the Scientific Results of the Voyage of H.M.S. Challenger
during the years 1873-1876, (Zoology), Under the Command of Captain George S. Nares,
R.N., F.R.S. and the Late Captain Frank Tourle Thomson, R.N.: Edinburgh, Neil and
Company, p. 1–362.

Milne Edwards, H., 1834-1840, Histoire naturelle des Crustacés, comprenant l'anatomie, la
physiologie et la classification de ces animaux, Roret. Paris, Librairie Encyclopédique de
Roret, v. 2, (1)468, (462)532, (463)638 p.:

Milne Edwards, H., Observations sur les affinités zoologiques et la classification naturelle des
Crustacés, *in* Proceedings Annales des Sciences Naturelles. Zoologie. Série 3, 18: 109-166,
pls 3, 4. (second part in Annales des Sciences naturelles, Zoologie, Série 3, 20: 163–228, pls
6-11 [1853])1852.

Milne-Edwards, A., 1862a, Monographie des Crustacés fossiles de la famille des Cancériens:
Annales des Sciences Naturelles, Zoologie, série 4, v. 18, p. 31–85.

Milne-Edwards, A., 1862b, Sur l'existence de Crustacés de la famille des Raniniens pendant la
période crétacée: Comptes Rendus de l'Academie des Sciences de Paris, v. 55, p. 492-494.

Milne-Edwards, A., 1865, Monographie des Crustacés Fossiles de la Famille des Cancériens:
Annales des Sciences Naturelles, 5e série, v. 3, p. 297–351.

Milne-Edwards, A., 1873, Crustacés fossiles nouveaux des terrains tertiaires de la Gironde: Actes de
la Société Linnéenne de Bordeaux, v. 29, p. 64–66.

Mina-Uhink, F., 1957, Bosquejo geoldgico del territorio de la Baja California: Boletin de la
Asociacidn Mexicana de Geólogos Petroleros, Mexico, D.F., v. 9, no. 3–4, p. 139–269.

Mitchell, S. F., Pickerill, R. K., Blackwell, B. A. B., and Skinner, A. R., 2000, The age of the Port
Morant Formation, south-eastern Jamaica: Caribbean Journal of Earth Science, v. 34, p. 1-4.

Mitchell, S. F., Pickerill, R. K., and Stemann, T. A., 2001, The Port Morant Formation (Upper
Pleistocene, Jamaica): high resolution sedimentology and paleoenvironmental analysis of a
mixed carbonate clastic lagoonal succession: Sedimentary Geology, v. 144, no. 3-4, p. 291-
306.

Montes, C., Cardona, A., McFadden, R., Morón, S., Silva, C., Restrepo-Moreno, S., Ramírez, D.,
Hoyos, N., Wilson, J., and Farris, D., 2012, Evidence for middle Eocene and younger land
emergence in central Panama: Implications for Isthmus closure: Bulletin, v. 124, no. 5-6, p.
780–799.

Morales-Ortega, P., González-Barba, G., Nava-Sánchez, E. H., and Vera-Dimas, D. R., 2016, New
Eocene bivalves from Bateque Formation, Baja California Sur, Mexico: Paleontología
Mexicana, v. 5, no. 1, p. 1–19.

Mori, M., and Yamanoi, T., 2003, Pollen assemblages and palaeovegetation changes of the Bihoku
and Katsuta Groups in Hiroshima Prefecture and Okayama Prefecture, southwest Japan:
Japanese Journal of Palynology, v. 49, no. 1, p. 9–20.

Morris, S. F., and Collins, J. S. H., 1991, Neogene crabs from Brunei, Sabah and Sarawak: Bulletin
of the British Museum (Natural History), (Geology), v. 47, no. 1, p. 1–33.

Morris, S. F., 1993, The fossil arthropods of Jamaica, *in* Wright, R. M., and Robinson, E., eds.,
Biostratigraphy of Jamaica. Geological Society of America Memoirs, Volume 182, p. 115–
124.

Moshammer, B., and Schlagintweit, F., 1999, The Ernstbrunn Limestone (Lower Austria): new data
on biostratigraphy and applied geology: Abhandlungen-Geologischen Bundesanstalt, v. 56,
no. 2, p. 553–566.

Müller, P., 1974, Decapoda (Crustacea) fauna a budapesti miocénből (1). [Les faunes de crustacés

décapodes des calcaires miocènes de Budapest (1)]: Foldtani Kozlony, v. 104, p. 119–132 [in

Hungarian

with French summary].

Müller, P., 1978, Decapoda (Crustacea) fauna a budapesti miocénböl (5): Földtani Közlöny, v. 108,

no. 3, p. 272–312.

Müller, P., 1984, Decapod Crustacea of the Badenian: Institutum Geologicicum Hungaricum,

Geologica Hungarica, Series Palaeontologica, v. 42, p. 3–317.

Müller, P., and Collins, J. S. H., 1991, Late Eocene coral-associated decapods (Crustacea) from

Hungary: Contributions to Tertiary and Quaternary Geology, v. 28, no. 2–3, p. 47–92.

Müller, P., 1996, Middle Miocene decapod Crustacea from southern Poland: Prace Muzeum Ziemi,

v. 43, p. 3–14.

Müller, P., 1998, Decapode Crustacea aus dem Karpat des Korneuburger Beckens (Unter-Miozän,

Niederösterreich): Beiträge zur Paläontologie, v. 23, p. 273-281.

Müller, P., 2006, New decapods from the Miocene of Hungary—with remarks about their

environment: Földtani Közlöny, v. 136, no. 1, p. 37-50.

Nations, J. D., 1975, The genus Cancer (Crustacea: Brachyura): systematics, biogeography and

fossil record: Science Bulletin of the National History Museum of Los Angeles County, v.

23, p. 1–104.

Nebelsick, J. H., Bassi, D., and Lempp, J., 2013, Tracking paleoenvironmental changes in coralline

algal-dominated carbonates of the Lower Oligocene Calcareniti di Castelgomberto formation

(Monti Berici, Italy): Facies, v. 59, no. 1, p. 133-148.

Ng, P. K., and Castro, P., 2020, A revision of *Carcinoplax abyssicola* (Miers, 1885) and seven

related species of *Carcinoplax* H. Milne Edwards, 1852, with the description of two new

species and an updated key to the genus (Crustacea, Decapoda, Brachyura, Goneplacidae):

Zoosystema, v. 42, no. 17, p. 239–284.

Ng, P. K. L., Guinot, D., and Davie, P. J. F., 2008, Systema Brachyurorum: Part I. An Annotated

Checklist of Extant Brachyuran Crabs of the World: The Raffles Bulletin of Zoology, v. 17,

p. 1–286.

Nyborg, T., Garassino, A., and Slak, G., 2017, Eonomus californianus n. gen., n. sp.(Crustacea,

Brachyura, Cymon-omidae) from the Eocene Llajas Formation, California (USA): Neues

Jahrbuch für Geologie und Paläontologie-Abhandlungen, v. 286, p. 189-193.

Ogg, J. G., 2020, Geomagnetic polarity time scale, Geologic time scale 2020, Elsevier, p. 159–192.

Orbigny, A. D. d., 1842, Voyage dans l'Amérique méridional, 1826-1833. Géologie et

Paléontologie, Paris.

Ortmann, A. E., 1893, Die Decapoden-Krebse des Strassburger Museums, mit besonderer

Berücksichtigung der von Herrn Dr. Döderlein bei Japan und bei den Liu-Kiu-Inseln

gesammelten und zur Zeit im Strassburger Museum aufbewahrten Formen. VII. Theil.

Abtheilung: Brachyura (Brachyura genuina Boas) II. Unterabtheilung: Cancroidea, 2.

Section: Cancrinea, 1. Gruppe: Cyclometopa. Zoologische Jahrbücher: Abteilung für

Systematik, Geographie und Biologie der Thiere, v. 7, no. 3, p. 411–495.

Ossó, À., 2016, *Eogeryon elegius* n. gen. and n. sp. (Decapoda: Eubrachyura: Portunoidea), one of

the oldest modern crabs from late Cenomanian of the Iberian Peninsula: Boletín de la

Sociedad Geológica Mexicana, v. 68, no. 2, p. 231–246.

Palacios Theil, E., Cuesta, J. A., and Felder, D. L., 2016, Molecular evidence for non-monophyly of

the pinnotheroid crabs (Crustacea: Brachyura: Pinnotheroidea), warranting taxonomic

reappraisal: Invertebrate Systematics, v. 30, p. 1–27.

Parham, J. F., Donoghue, P. C., Bell, C. J., Calway, T. D., Head, J. J., Holroyd, P. A., Inoue, J. G.,

Irmis, R. B., Joyce, W. G., and Ksepka, D. T., 2012, Best practices for justifying fossil

calibrations: Systematic Biology, v. 61, no. 2, p. 346–359.

Pasini, G., Garassino, A., De Angeli, A., Hyzný, M., Giusberti, L., and Zorzin, R., 2019, Eocene

decapod faunas from the Konservat-Lagerstätten laminites of "Pesciara"(Bolca, Verona) and

Monte Postale (Altissimo, Vicenza) in northeast Italy: a review and update: Neues Jahrbuch

für Geologie und Paläontologie-Abhandlungen, v. 293, p. 233–270.

Pauca, M., 1929, Zwei neue fossile Decapoden aus Oligozanen Clupea crenata Schiefern von

Suslaneski-Muscel und über die Bildung der Menilitschiefern: Bulletin de la Section

Scientifique de l'Académie Roumaine, v. 12, no. 4–5, p. 40–44.

Paulay, G., and Starmer, J., 2011, Evolution, insular restriction, and extinction of oceanic land crabs,

exemplified by the loss of an endemic *Geograpsus* in the Hawaiian Islands: PLoS One, v. 6,

no. 5, p. e19916.

Perea, D., Verde, M., Toriño, P., Montenegro, F., Ubilla, M., and Manzuetti, A., 2020, A Complex

Association of Invertebrates, Vertebrates and Trace Fossils in the Marine Camacho

Formation (Late Miocene of Uruguay): Biostratigraphy and Paleoenvironments:

Ameghiniana, v. 57, no. 3, p. 266–277.

Philippi, R. A., 1887, Die Tertiären und Quartáren Versteinerunge Chiles, Brockhaus, Leipzig, 226
p.:

Piller, W. E., Harzhauser, M., and Mandic, O., 2007, Miocene Central Paratethys stratigraphy–
current status and future directions: Stratigraphy, v. 4, no. 2–3, p. 151–168.

Poore, G. C., and Ahyong, S. T., 2023, Marine Decapod Crustacea: A Guide to Families and Genera
of the World, CSIRO PUBLISHING, 916 p.:

Portell, R. W., and Agnew, J., 2004, Pliocene and Pleistocene decapod crustaceans: Florida Fossil
Invertebrates, v. 4, p. 1–29.

Prado, L. A. C., Luque, J., Barreto, A. M. F., and Palmer, A. R., 2018, New brachyuran crabs from
the Aptian–Albian Romualdo Formation, Santana Group of Brazil: Evidence for a Tethyan
connection to the Araripe Basin: Acta Palaeontologica Polonica, v. 63, no. 2018, p. 737–750.

Pujalte, V., Schmitz, B., Baceta, J. I., Orue-Echebarría, X., Bernaola, G., Dinarès-Turell, J., Payros,
A., Apellaniz, E., and Caballero, F., 2009, Correlation of the Thanetian–Ilerdian turnover of
larger foraminifera and the Paleocene–Eocene thermal maximum: confirming evidence from
the Campo area (Pyrenees, Spain): Geologica Acta, p. 161–175.

Quayle, W., and Collins, J., 2012, A review of the decapod crustaceans from the Tertiary of the Isle
of Wight, Hampshire, UK, with description of three new species: Bulletin of the Mizunami
Fossil Museum, v. 38, p. 33–51.

Quayle, W. J., and Collins, J. S. H., 1981, New Eocene crabs from the Hampshire Basin:
Palaeontology, v. 24, no. 4, p. 733–758.

Raine, J. I., Beu, A. G., Boyes, A. F., Campbell, H. J., Cooper, R. A., Crampton, J. S., Crundwell,
M. P., Hollis, C. J., and Morgans, H. E. G., 2015, Revised calibrationof the New Zealand
geological timescale: NZGT2015/1, GNSScience Report 2012/39, 53 p.:

Rathbun, M. J., 1897, A revision of the nomenclature of the Brachyura: Proceedings of the
Biological Society of Washington, v. 11, p. 153–167.

Rathbun, M. J., 1898, The Brachyura of the biological expedition to the Florida Keys and the
Bahamas in 1893: Bulletin from the Laboratories of Natural History, State University of
Iowa, v. 4, no. 3, p. 250–294.

Rathbun, M. J., 1918 [1919], Decapod crustaceans from the Panama region, *in* Vaughan, T. W., ed.,
Contributions to the geology and paleontology of the Canal Zone, Panama and geologically
related areas in Central America and the West Indies. Bulletin of the United States National
Museum, Volume 103, p. 123–184.

Rathbun, M. J., 1926, The fossil stalk-eyed Crustacea of the Pacific slope of North America, US
Government Printing Office, Bulletin of the United States National Museum.

Rathbun, M. J., 1935, Fossil Crustacea of the Atlantic and Gulf Coastal Plain: Geological Society of
America, v. Special Paper, no. 2, p. i–viii, 1–160.

Rathbun, M. J., 1937, Cretaceous and Tertiary crabs from Panama and Colombia: Journal of
Paleontology, v. 11, no. 1, p. 26–28.

Reichenbacher, B., Krijgsman, W., Lataster, Y., Pipperr, M., Van Baak, C. G., Chang, L., Kälin, D.,
Jost, J., Doppler, G., and Jung, D., 2013, A new magnetostratigraphic framework for the
lower Miocene (Burdigalian/Ottnangian, Karpatian) in the North Alpine Foreland Basin:
Swiss Journal of Geosciences, v. 106, no. 2, p. 309-334.

Reuss, A. E., 1857, Zur Kenntnis fossiler Krabben: Sitzungsberichte der Akademie der
Wissenschaften, v. 27, p. 161–166.

Reuss, A. E., 1859, Zur Kenntnis fossiler Krabben: Denkschriften der Kaiserlichen Akademie der
Wissenschaften. Mathematisch-Naturwissenschaftliche Classe, v. 17, p. 1–90.

Ristori, G., 1886, I crostacei brachiuri e anomuri del pliocene italiano: Bollettino della Società
Geologica Italiana, v. 5, p. 93–128.

Robins, C. M., Feldmann, R. M., and Schweitzer, C. E., 2013, Nine new genera and 24 new species
of the Munidopsidae (Decapoda: Anomura: Galatheoidea) from the Jurassic Ernstbrunn
Limestone of Austria, and notes on fossil munidopsid classification: Annalen des
Naturhistorischen Museums in Wien. Serie A für Mineralogie und Petrographie, Geologie
und Paläontologie, Anthropologie und Prähistorie, p. 167–251.

Robins, C. M., and Klompmaker, A. A., 2019, Extreme diversity and parasitism of Late Jurassic
squat lobsters (Decapoda: Galatheoidea) and the oldest records of porcellanids and
galatheids: Zoological Journal of the Linnean Society, v. 187, no. 4, p. 1131-1154.

Roemer, F. A., 1887, *Graptocarcinus texanus*, ein Brachyure aus der oberen Kreide von Texas:
Leonhardt und Bronn's Neues Jahrbuch für Mineralogie, Geologie, und Paläontologie, v. 1,
p. 173–176.

Ross, A., Lewis, J. E., and Scolaro, R. J., 1964, New Eocene decapods from Florida: Quarterly
Journal of the Florida Academy of Sciences, v. 27, no. 3, p. 187–196.

Roux, P., 1828-1830, Crustacés de la Mediterranée et de son littoral. Décrits et Lithographiés par
Polydore Roux, Conservateur du Cabinet d'histoire naturelle de la Ville de Marseille. , Paris,
Levrault.

Rumsey, B. T., Klompmaker, A. A., and Portell, R. W., 2016, Paleobiogeography of the fossil box
crab Calappilia (Brachyura: Calappidae) with a new species from the Eocene-Oligocene of
Florida, USA: Journal of Crustacean Biology, v. 36, no. 3, p. 329-337.

Sadler, P. M., 1988, Geometry and stratification of uppermost cretaceous and paleogene units on
Seymour Island, Northern Antarctic Peninsula, *in* Feldmann, R. M., and Woodburne, M.,
eds., Geology and Paleontology of Seymour Island, Antarctic Peninsula. Memoirs of the
Geological Society of America, Volume 169: California, Geological Society of America, p.
303–320.

Samouelle, G., 1819, The entomologist's useful compendium; or an introduction to the knowledge
of British insects, comprising the best means of obtaining and preserving them, and a
description of the apparatus generally used; together with the genera of Linné, and the
modern method of arranging the classes Crustacea, Myriapoda, Spiders, Mites and Insects,
from their affinities and structure, according to the views of Dr. Leach, London.

Schneider, S., Harzhauser, M., Kroh, A., Lukeneder, A., and Zuschin, M., 2013, Ernstbrunn
Limestone and Klentnice beds (Kimmeridgian Berriasian; Waschberg-dánice Unit; NE
Austria and SE Czech Republic): state of the art and bibliography: Bulletin of Geosciences,
v. 88, no. 10.

Scholtz, G., and Richter, S., 1995, Phylogenetic systematics of the reptantian Decapoda (Crustacea,
Malacostraca): Zoological Journal of the Linnaean Society, v. 113, p. 289–328.

Scholtz, G., 2014, Evolution of crabs–history and deconstruction of a prime example of
convergence: Contributions to Zoology, v. 83, no. 2, p. 87–105.

Scholtz, G., 2020, *Eocarcinus praecursor* Withers, 1932 (Malacostraca, Decapoda, Meiura) is a
stem group brachyuran: Arthropod Structure & Development.

Schram, F. R., and Dixon, C. J., 2004, Decapod phylogeny: addition of fossil evidence to a robust
morphological cladistic data set: Bulletin of the Mizunami Fossil Museum, v. 31, p. 1–19.

Schram, F. R., and Ng, P. K. L., 2012, What is *Cancer*?: Journal of Crustacean Biology, v. 32, no. 4,
p. 665–672.

Schubart, C. D., and Cuesta, J. A., 2010, Phylogenetic relationships of the Plagusiidae Dana, 1851
(Brachyura), with description of a new genus and recognition of Percnidae Stevcic, 2005, as
an independent family: Crustaceana Monographs, v. 11, p. 279–299.

Schweitzer, C. E., 2000, Tertiary Xanthoidea (Crustacea: Decapoda: Brachyura) from the west coast
of North America: Journal of Crustacean Biology, v. 20, no. 4, p. 715–742.

Schweitzer, C. E., and Feldmann, R. M., 2000a, New species of calappid crabs from western North America and reconsideration of the Calappidae sensu lato: Journal of Paleontology, v. 74, no. 2, p. 230–246.

Schweitzer, C. E., and Feldmann, R. M., 2000b, Reevaluation of the Cancridae Latreille, 1803 (Decapoda: Brachyura) including three new genera and three new species: Contributions to Zoology, v. 69, p. 223–250.

Schweitzer, C. E., Feldmann, R. M., Tucker, A. B., and Berglund, R. E., 2000, Eocene decapod crustaceans from Pulali Point, Washington: Annals of the Carnegie Museum, v. 69, no. 1, p. 23–67.

Schweitzer, C. E., and Feldmann, R. M., 2001, Differentiation of the fossil Hexapodidae Miers, 1886 (Decapoda: Brachyura) from similar forms: Journal of Paleontology, v. 75, p. 330–345.

Schweitzer, C. E., Feldmann, R. M., González–Barba, G., and Vega, F. J., 2002, New crabs from the Eocene and Oligocene of Baja California Sur, Mexico and an assessment of the evolutionary and paleobiogeographic Implications of Mexican fossil decapods: Paleontological Society Memoir, v. 76, p. 1–43.

Schweitzer, C. E., Feldmann, R. M., and Gingerich, P. D., 2004, New Decapoda (Crustacea) from the middle and late Eocene of Pakistan and a revision of *Lobonotus* A. Milne Edwards, 1864: University of Michigan, Contributions from the Museum of Paleontology, v. 31, no. 4, p. 89–118.

Schweitzer, C. E., 2005, The Trapeziidae and Domeciidae (Decapoda: Brachyura: Xanthoidea) in the fossil record and a new Eocene genus from Baja California Sur, Mexico: Journal of Crustacean Biology, v. 25, no. 4, p. 625–636.

Schweitzer, C. E., Feldmann, R. M., González-Barba, G., and Cosovic, V., 2006, New Decapoda (Anomura, Brachyura) from the Eocene Bateque and Tepetate Formations, Baja California Sur, Mexico: Bulletin of the Mizunami Fossil Museum, v. 33, p. 21–45.

Schweitzer, C. E., Feldmann, R. M., and Lazar, I., 2009, Fossil Crustacea (excluding Cirripedia and Ostracoda) in the University of Bucharest Collections, Romania, including two new species: Bulletin of the Mizunami Fossil Museum, v. 35, p. 1–14.

Schweitzer, C. E., and Feldmann, R. M., 2010a, New fossil decapod crustaceans from the Remy Collection, Muséum national d'Histoire naturelle, Paris: Geodiversitas, v. 32, no. 3, p. 399–415.

Schweitzer, C. E., and Feldmann, R. M., 2010b, Earliest known Porcellanidae (Decapoda: Anomura:
Galatheoidea) (Jurassic: Tithonian): Neues Jahrbuch für Geologie und Paläontologie,
Abhandlungen, v. 258, no. 2, p. 243-248.

Schweitzer, C. E., Feldmann, R. M., Garassino, A., Karasawa, H., and Schweigert, G., 2010,
Systematic List of Fossil Decapod Crustacean Species: Crustaceana Monographs, v. 10, p.
222.

Schweitzer, C. E., and Feldmann, R. M., 2011, New fossil Brachyura (Decapoda: Homoloidea,
Dorippoidea, Carpilioidea) from the United Kingdom: Bulletin of the Mizunami Fossil
Museum, v. 37, p. 1–11.

Schweitzer, C. E., and Feldmann, R. M., 2012, Revision of Decapoda deposited in The Muséum
national d'Histoire naturelle, Paris: Bulletin of the Mizunami Fossil Museum, v. 38, p. 15–
27.

Schweitzer, C. E., Feldmann, R. M., and Karasawa, H., 2012, Part R, Revised, Volume 1, Chapter
8M: Systematic Descriptions: Section Dromiacea: Treatise Online, v. 51, p. 1–43, 24 figs.

Schweitzer, C. E., Karasawa, H., Luque, J., and Feldmann, R. M., 2016a, Phylogeny and
classification of Necrocarcinoidea Förster, 1968 (Brachyura: Raninoida) with description of
two new genera: Journal of Crustacean Biology, v. 36, no. 3, p. 338–372.

Schweitzer, C. E., Odumodu, C. F. R., and Feldmann, R. M., 2016b, New Eocene crabs from
Nigeria (Decapoda: Brachyura: Heterotremata): Annals of Carnegie Museum, v. 84, no. 1, p.
59–73.

Schweitzer, C. E., Feldmann, R. M., and Karasawa, H., 2017, Part R, Revised, Volume 1, Chapter
8R: Systematic Descriptions: Section Cyclodorippoida: Treatise Online, v. 97, p. 1–6, 4 figs.

Schweitzer, C. E., Feldmann, R. M., Karasawa, H., and Luque, J., 2018, Part R, Revised, Volume 1,
Chapter 8S: Systematic Descriptions: Section Raninoida: Treatise Online, v. 113, p. 1–42, 20
figs.

Schweitzer, C. E., Feldmann, R. M., and Karasawa, H., 2021a, Part R, Revised, Volume 1, Chapter
8T10: Systematic descriptions: Superfamily Ethioidea, Treatise Online, Volume 132, p. 1–8.

Schweitzer, C. E., Feldmann, R. M., and Karasawa, H., 2021b, Part R, Revised, Volume 1, Chapter
8T17: Systematic descriptions: Superfamily Dorippoidea, Treatise Online, Volume 159, p.
1–8.

Schweitzer, C. E., Feldmann, R. M., and Karasawa, H., 2021c, Part R, Revised, Volume 1, Chapter
2685   8T15: Systematic descriptions: Superfamily Portunoidea, Treatise Online, Volume 151, p. 1–
2686   40.

2687 Schweitzer, C. E., Feldmann, R. M., and Karasawa, H., 2021d, Part R, Revised, Volume 1, Chapter
2688   8T16: Systematic descriptions: Superfamilies Trapezioidea and Xanthoidea, Treatise Online,
2689   Volume 153, p. 1–42.

2690 Scott, R. W., Schlager, W., Fouke, B., and Nederbragt, S. A., 2000, Are mid-Cretaceous eustatic
2691   events recorded in Middle East carbonate platforms?, *in* Alsharhan, A. S., and Scott, R. W.,
2692   eds., Middle East models of Jurassic/Cretaceous Carbonate Systems: SEPM Special
2693   Publication, Volume 69, p. 73–84.

2694 Scott, R. W., Benson, D. G., Morin, R. W., Shaffer, B. L., and Oboh-Ikuenobe, F. E., 2002,
2695   Integrated Albian-Lower Cenomanian chronostratigraphy standard, Trinity River section,
2696   Texas: US Gulf Coast Cretaceous stratigraphy and paleoecology: Gulf Coast Section,
2697   Society of Economic Paleontologists and Mineralogists, Bob F. Perkins Memorial Volume,
2698   v. 23, p. 277–334.

2699 Scott, R. W., Campbel, W., Hojnacki, R., Wang, Y., and Lai, X., 2016, Albian rudist biostratigraphy
2700   (Bivalvia), Comanche shelf to shelf margin, Texas.

2701 Secrétan, S., 1975, Les Crustacés du Monte Bolca. Studi e Ricerche sui Giacimenti Terziari di Bolca
2702   II, Miscellanea Paleontologica: Museo Civico di Storia Naturale - Studi e Ricerche sui
2703   giacimenti terziari di Bolca, Verona, v. 2, p. 315-388, pls. I-XXXVII.

2704 Serra-Kiel, J., Canudo, J., Dinares, J., Molina, E., Ortiz, N., Pascual, J., Samsó, J., and Tosquella, J.,
2705   1994, Cronoestratigrafía de los sedimentos marinos del Terciario inferior de la Cuenca de
2706   Graus-Tremp (Zona Central Surpirenaica): Revista de la Sociedad Geológica de España, v.
2707   7, no. 3–4, p. 273–297.

2708 Serrano-Sánchez, M., Hegna, T. A., Schaaf, P., Pérez, L., Centeno-García, E., and Vega, F. J., 2015,
2709   The aquatic and semiaquatic biota in Miocene amber from the Campo La Granja mine
2710   (Chiapas, Mexico): Paleoenvironmental implications: Journal of South American Earth
2711   Sciences, v. 62, p. 243-256.

2712 Serrano-Sánchez, M. d. L., Guerao, G., Centeno-García, E., and Vega, F. J., 2016, Crabs
2713   (Brachyura: Grapsoidea: Sesarmidae) as inclusions in Lower Miocene amber from Chiapas,
2714   Mexico: Boletín de la Sociedad Geológica Mexicana, v. 68, no. 1, p. 37–43.

Shi, G., Grimaldi, D. A., Harlow, G. E., Wang, J., Wang, J., Yang, M., Lei, W., Li, Q., and Li, X.,
2012, Age constraint on Burmese amber based on U–Pb dating of zircons: Cretaceous
Research, v. 37, p. 155–163.

Solórzano Kraemer, M., 2007, Systematic, palaeoecology, and palaeobiogeography of the insect
fauna from Mexican amber: Palaeontographica Abteilung A, p. 1–133.

Sorensen, P. A., 1982, Sedimentology and sedimentary petrology of a Paleoce Basin near Laguna
San Ignacio, Baja California Sur, Mexico [Master of Arts: University of California, Santa
Barbara, California, United States, 123 p.

Spiridonov, V. A., Neretina, T. V., and Schepetov, D., 2014, Morphological characterization and
molecular phylogeny of Portunoidea Rafinesque, 1815 (Crustacea Brachyura): implications
for understanding evolution of swimming capacity and revision of the family-level
classification: Zoologischer Anzeiger - A Journal of Comparative Zoology, v. 253, no. 5, p.
404–429.

Stebbing, T. R. R., 1920, The Malacostraca of Durban Bay: Durban Museum Novitates, v. 2, no. 6,
p. 263–278.

Stenzel, H. B., 1944, A new Cretaceous crab, *Graptocarcinus muiri*, from Mexico: Journal of
Paleontology, v. 18, no. 6, p. 550–552.

Stevcic, Z., 2005, The reclassification of brachyuran crabs (Crustacea: Decapoda: Brachyura):
Natura Croatica, v. 14, no. 1, p. 1–159.

Stimpson, W., 1858, Prodromus descriptionis animalium evertebratorum, quae in Expeditione ad
Oceanum Pacificum Septentrionalem, a Republica Federata missa, Cadwaladaro Ringgold et
Johanne Rodgers ducibus, observavit et descripsit. Pars VII. Crustacea Anomura:
Proceedings of the Academy of Natural Sciences of Philadelphia, v. 10, no. 7, p. 225–252
[Pages 263-290 on separate].

Stimpson, W., 1859, Notes on North American Crustacea in the Museum of the Smithsonian
Institution No. 1: Annals of the Lyceum of Natural History, New York, v. 7, no. 1962, p. 49–
93.

Stimpson, W., 1860, Notes on North American Crustacea in the Museum of the Smithsonian
Institution No. 2: Annals of the Lyceum of Natural History, New York, v. 7, p. 176–246.

Stoliczka, F., 1871, Observations on fossil crabs from Tertiary deposits in Sind and Kutch: Memoirs
of the Geological Survey of India, Palaeontologica Indica, (7) (14), v. 1, no. 1, p. 1–16.

Studer, T., 1892, Ueber zwei fossile dekapode Krebse aus den Molasseablagerungen des
Belpberges: Abhandlungen der Schweizerischen Paläontologischen Gesellschaft, v. 19, p. 1–
8.

Suzuki, S., Danhara, T., and Tanaka, H., 2003, Fission-track dating of Tertiary tuff samples from the
Kibi Plateau Area, Okayama prefecture, Southwest Japan: Journal of Geography, v. 112, p.
35–49.

Sztrákos, K., and Steurbaut, E., 2017, Révision lithostratigraphique et biostratigraphique de
l'Oligocène d'Aquitaine occidentale (France): Geodiversitas, v. 39, no. 4, p. 741–781.

Taguchi, E., 2002, Stratigraphy, molluscan fauna and paleoenvironment of the Miocene Katsuta
Group in Okayama Prefecture, Southwest Japan: Bulletin of the Mizunami Fossil Museum,
v. 29, p. 95–133.

Tan, M. H., Gan, H. M., Lee, Y. P., Linton, S., Grandjean, F., Bartholomei-Santos, M. L., Miller, A.
D., and Austin, C. M., 2018, ORDER within the chaos: Insights into phylogenetic
relationships within the Anomura (Crustacea: Decapoda) from mitochondrial sequences and
gene order rearrangements: Molecular Phylogenetics and Evolution, v. 127, p. 320-331.

Tang, Y.-Y., Tang, B.-P., Xin, Z.-Z., Li, Y.-T., Zha, X.-H., Zhang, D.-Z., Sun, Y., Liu, Q.-N., and
Ma, Y.-F., 2020, Characterization of the complete mitochondrial genome of *Helice latimera*
and its phylogenetic implications in Brachyura: Genomics, v. 112, no. 6, p. 5180-5187.

Tavares, M., 1993, Crustacea Decapoda: les Cyclodorippidae et Cymonomidae de l'Indo-Ouest-
Pacifique à l'exclusion du genre *Cymonomus*, *in* Crosnier, A., ed., Résultats des Campagnes
MUSORSTOM, Volume 10. Mémoires du Muséum national d'Histoire naturelle, Volume
156, p. 253–313.

Thoma, B. P., Guinot, D., and Felder, D. L., 2014, Evolutionary relationships among American mud
crabs (Crustacea: Decapoda: Brachyura: Xanthoidea) inferred from nuclear and
mitochondrial markers, with comments on adult morphology: Zoological Journal of the
Linnean Society, v. 170, no. 1, p. 86–109.

Thurmann, J., 1853, Lettres écrites du Jura à la Société d'histoire naturelle de Berne: Lettre IX,
Mitteilungen der Naturforschenden Gesellschaft Bern.

Tsang, C. T., Schubart, C. D., Chu, K. H., Ng, P. K., and Tsang, L. M., 2022, Molecular phylogeny
of Thoracotremata crabs (Decapoda, Brachyura): Toward adopting monophyletic
superfamilies, invasion history into terrestrial habitats and multiple origins of symbiosis:
Molecular Phylogenetics and Evolution, v. 177, p. 107596.

Tsang, L. M., Schubart, C. D., Ahyong, S. T., Lai, J. C. Y., Au, E. Y. C., Chan, T.-Y., Ng, P. K. L.,
and Chu, K. H., 2014, Evolutionary history of true crabs (Crustacea: Decapoda: Brachyura)
and the origin of freshwater crabs: Molecular Biology and Evolution, v. 31, no. 5, p. 1173–
1187.

Tucker, A. B., and Feldmann, R. M., 1990, Fossil decapod crustaceans from the lower Tertiary of
the Prince William Sound region, Gulf of Alaska: Journal of Paleontology, v. 64, p. 409–
427.

Tucker, A. B., 1998, Systematics of the Raninidae (Crustacea: Decapoda: Brachyura), with accounts
of three new genera and two new species: Proceedings of the Biological Society of
Washington, v. 111, no. 2, p. 320–371.

Ungaro, S., 1978, L'Oligocene dei Colli Berici: Rivista, Italiana di Paleontologiia e Stratigrafia, v.
84, p. 199–278.

Van Bakel, B. W., Guinot, D., Corral, J. C., and Artal, P., 2012a, Graptocarininae n. subfam., an
extinct subfamily of Dynomenidae Ortmann, 1892 (Crustacea, Brachyura, Podotremata):
Zootaxa, v. 3534, no. 1, p. 40–52.

Van Bakel, B. W., Mychko, E. V., Spiridonov, A., Jagt, J. W., and Fraaije, R. H., 2021, New
Cretaceous crabs (Crustacea, Brachyura) from Moscow Oblast and Dagestan (Russia):
patterns in phylogeny and morphospace of the oldest eubrachyurans (Dorippoidea):
Cretaceous Research, v. 119, p. 104675.

Van Bakel, B. W. M., Guinot, D., Artal, P., Fraaije, R. H. B., and Jagt, J. W. M., 2012b, A revision
of the Palaeocorystoidea and the phylogeny of raninoidian crabs (Crustacea, Decapoda,
Brachyura, Podotremata): Zootaxa, v. 3215, p. 1–216.

Van Straelen, V., 1920, Catalogue des crustacés décapodes des terrains Tertiaires de la Belgique:
Annales de la Société royale zoologique de Belgique, v. 51, p. 111–131.

Varola, A., 1981, Crostacei Decapodi neogenici della penisola salentina (Italia): Thalassia salentina,
v. 11, p. 3–51.

Vega, F. J., Cosma, T., Coutiño, M. A., Feldmann, R. M., Nyborg, T., Schweitzer, C. E., and
Waugh, D. A., 2001, New middle Eocene decapods (Crustacea) from Chiapas, Mexico:
Journal of Paleontology, v. 75, p. 929–946.

Vega, F. J., Nyborg, T., Fraaye, R. H., and Espinosa, B., 2007, Paleocene decapod Crustacea from
the Rancho Nuevo Formation (Parras Basin-Difunta Group), Northeastern México: Journal
of Paleontology, v. 81, no. 6, p. 1432–1441.

Vega, F. J., Nyborg, T., Coutiño, M. A., and Hernández-Monzón, O., 2008, Review and additions to
the Eocene decapod Crustacea from Chiapas, Mexico: Bulletin of the Mizunami Fossil
Museum, v. 34, p. 51–71.

Vega, F. J., Nyborg, T., Coutiño, M. A., Solé, J., and Hernández-Monzón, O., 2009, Neogene
Crustacea from Southeastern Mexico: Bulletin of the Mizunami Fossil Museum, v. 35, p.
51–69.

Wagreich, M., Hohenegger, J., and Ćorić, S., Base and new definition of the Lower Badenian and
the age of the Badenian stratotype (Middle Miocene, Central Paratethys), *in* Proceedings
STRATI 2013: First International Congress on Stratigraphy At the Cutting Edge of
Stratigraphy2014, Springer, p. 615–618.

Wang, Q., Tang, D., Guo, H., Wang, J., Xu, X., and Wang, Z., 2020, Comparative mitochondrial
genomic analysis of *Macrophthalmus pacificus* and insights into the phylogeny of the
Ocypodoidea & Grapsoidea: Genomics, v. 112, no. 1, p. 82-91.

Wang, Z., Wang, Z., Shi, X., Wu, Q., Tao, Y., Guo, H., Ji, C., and Bai, Y., 2018, Complete
mitochondrial genome of *Parasesarma affine* (Brachyura: Sesarmidae): Gene
rearrangements in Sesarmidae and phylogenetic analysis of the Brachyura: International
journal of biological macromolecules, v. 118, p. 31–40.

Watson-Zink, V. M., 2021, Making the grade: physiological adaptations to terrestrial environments
in decapod crabs: Arthropod structure & development, v. 64, p. 101089.

Webber, W., Fenwick, G., Bradford-Grieve, J., Eagar, S., Buckeridge, J., Poore, G., Dawson, E.,
Watling, L., Jones, J., Wells, J., Bruce, N., Ahyong, S., Larsen, K., Chapman, M., Olesen, J.,
Ho, J., Green, J., Shiel, R., Rocha, C., Lorz, A., Bird, G., and Charleston, W., 2010, Phylum
Arthropoda Subphylum Crustacea: shrimps, crabs, lobsters, barnacles, slaters, and kin, *in*
Gordon, D. P., ed., New Zealand Inventory of Biodiversity: Volume Two: Kingdom
Animalia - Chaetognatha, Ecdysozoa, Ichnofossils, : New Zealand, Canterbury University
Press, p. 98–232.

Weber, F., 1795, Nomenclator entomologicus secundum entomologiam systematicam Fabricii,
adjectis speciebus recens detectis et varietatibus, Chilonii and Hamburgi, Viii +171 p.:

Wegner, W., Wörner, G., Harmon, R. S., and Jicha, B. R., 2011, Magmatic history and evolution of
the Central American Land Bridge in Panama since Cretaceous times: Bulletin, v. 123, no. 3-
4, p. 703-724.

Williams, A. B., and Child, C. A., 1988, Comparison of some genera and species of box crabs

(Brachyura: Calappidae), southwestern North Atlantic, with description of a new genus and

species: Fishery Bulletin, Washington, v. 87, no. 1, p. 105–121.

Windsor, A. M., and Felder, D. L., 2014, Molecular phylogenetics and taxonomic reanalysis of the

family Mithracidae MacLeay (Decapoda : Brachyura : Majoidea): Invertebrate Systematics,

v. 28, p. 145–173.

Withers, T. H., 1932, XXXIII.—A Liassic crab, and the origin of the Brachyura: Annals and

Magazine of Natural History, v. 9, no. 51, p. 313–323.

Wolfe, J. M., Daley, A. C., Legg, D. A., and Edgecombe, G. D., 2016, Fossil calibrations for the

arthropod Tree of Life: Earth-Science Reviews, v. 160, p. 43–110.

Wolfe, J. M., Breinholt, J. W., Crandall, K. A., Lemmon, A. R., Moriarty Lemmon, E., Timm, L. E.,

Siddall, M. E., and Bracken-Grissom, H. D., 2019, A phylogenomic framework,

evolutionary timeline and genomic resources for comparative studies of decapod

crustaceans: Proceedings of the Royal Society B, v. 286, p. 20190079.

Wolfe, J. M., Luque, J., and Bracken-Grissom, H., 2021, How to become a crab: A recurring body

plan is a driver of biodiversity: Bioessays, v. 43, no. 5, p. 1–14.

Wolfe, J. M., Ballou, L., Luque, J., Watson-Zink, V. M., Ahyong, S. T., Barido-Sottani, J., Chan, T.-

Y., Chu, K. H., Crandall, K. A., Daniels, S. R., Felder, D. L., Mancke, H., Martin, J. W., Ng,

P. K. L., Ortega-Hernández, J., Palacios Theil, E., Pentcheff, N. D., Robles, R., Thoma, B.

P., Tsang, L. M., Wetzer, R., Windsor, A. M., and Bracken-Grissom, H. D., 2022,

Convergent adaptation of true crabs (Decapoda: Brachyura) to a gradient of terrestrial

environments: bioRxiv, p. 2022-2012.

Woods, H., 1922, Crustacea from the Eocene deposits of Peru, *in* Bosworth, T. O., ed., Geology of

the Tertiary and Quaternary Periods in the northwest part of Peru: London, Macmillan and

Co., p. 114–118.

Yoshimoto, Y., 1979, 38. Tsuyama basin, Okayama Prefecture, *in* Tsuchi, R., ed., Fundamental data

on Japanese Neogene Bio- and Chronostratigraphy, IGCO-114 National Working Group of

Japan, Shizuoka (in Japanese), p. 113–114.

Zeiss, A., 2001, Die Ammonitenfauna der Tithonklippen von Ernstbrunn, Niederösterreich: Neue

Denkschriften des Naturhistorischen Museums in Wien, Neue Serie, v. 6, p. 1–117.

Zhang, Y., Gong, L., Lu, X., Miao, Z., Jiang, L., Liu, B., Liu, L., Li, P., Zhang, X., and Lü, Z., 2022,

Comparative mitochondrial genome analysis of Varunidae and its phylogenetic implications:

Acta Oceanologica Sinica, v. 41, no. 6, p. 119-131.

2874