High-fat diet in early life triggers both reversible and persistent

epigenetic changes in the medaka fish (*Oryzias latipes*)

Authors

Yusuke Inoue[1]†*, Yuta Suzuki[2]†, Yoshimi Kunishima[1]†, Terumi Washio[1], Shinichi

Morishita[2]*, Hiroyuki Takeda[1]*

Affiliations:

1. Department of Biological Sciences, Graduate School of Science, The University of

Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

2. Department of Computational Biology and Medical Sciences, Graduate School of

Frontier Sciences, The University of Tokyo, 5-1-5 Kashiwanoha, Kashiwa, Chiba

277-8561, Japan

†Equal contributions

*Correspondence to; Y.I. (y-inoue@bs.s.u-tokyo.ac.jp), S.M. (moris@cb.k.u-tokyo.ac.jp),

H.T. (htakeda@bs.s.u-tokyo.ac.jp)

Abstract

The nutritional status during early life can have enduring effects on an animal's metabolism, although the mechanisms underlying these long-term effects are still unclear. Epigenetic modifications are considered a prime candidate mechanism for encoding early-life nutritional memories during this critical developmental period. However, the extent to which these epigenetic changes occur and persist over time remains uncertain, in part due to challenges associated with directly stimulating the fetus with specific nutrients in viviparous mammalian systems. In this study, we used medaka as an oviparous vertebrate model to establish an early-life high-fat diet (HFD) model. Larvae were fed with HFD from the hatching stages (one week after fertilization) for six weeks, followed by normal chow (NC) for eight weeks until the adult stage. We examined the changes in the transcriptomic and epigenetic state of the liver over this period. We found that HFD induces fatty liver phenotypes, accompanied by drastic changes in the hepatic transcriptome, chromatin accessibility, and histone modifications, especially in metabolic genes. These changes were largely reversed after the long-term NC, demonstrating the high plasticity of the epigenetic state in hepatocytes. However, we found a certain number of genomic loci showing non-reversible epigenetic changes, especially around genes related to cell signaling, implying persistent changes in the cellular state of the liver triggered by early-life HFD feeding. Our data provide novel insights into the epigenetic mechanism of nutritional programming and a comprehensive atlas of the long-term epigenetic state in an early-life HFD model of non-mammalian vertebrates.

1   **Introduction**

2   Early-life nutrition can have life-long effects on the metabolism of animals. This

3   phenomenon is referred to as "nutritional programming" (Langley-Evans, 2009). It is

4   generally accepted that animals have the ability to adapt to the energy-imbalanced

5   environments imposed during early life and continually shift their metabolic states

6   throughout their lives (Pfennig et al., 2010). Accordingly, if early nutritional

7   programming does not correctly match the postnatal environment, it may cause metabolic

8   disorders such as obesity, type 2 diabetes, and cardiovascular diseases (Barker, 2007;

9   Langley-Evans, 2009). Liver serves as the central organ regulating the metabolic

10 homeostasis of animals across feeding-fasting cycles (Bideyan et al., 2021; Lempradl et

11 al., 2015), adapting its phenotype in response to metabolic states, such as the development

12 of fatty liver (Brunt et al., 2015). Regarding nutritional programming, cohort studies and

13 experimental studies using mammalian models have shown that maternal high-fat diet

14 (HFD) feeding, obesity, and diabetes during gestation and suckling period are associated

15 with an increased risk of non-alcoholic fatty liver disease (NAFLD) in the offspring (de

16 Jesus et al., 2020; Gregorio et al., 2010; Huang et al., 2017; Kruse et al., 2013; M. Li et

17 al., 2015; Oben et al., 2010; Thompson, 2020). Epigenetic modifications are one of the

18 prime candidate mechanisms implicated in long-term nutritional programming of the

19 liver (Costello and Schones, 2018), since they can be maintained across cell divisions

20 over a long period of time (Moazed, 2011). Indeed, in some types of cells such as innate

21 immunity cells (Netea et al., 2016), epidermal stem cells (Larsen et al., 2021; Naik et al.,

22 2017), and mammary epithelial cells (dos Santos et al., 2015), transient external stimuli

23 caused long-term changes in chromatin state even after external stimuli were removed.

24 These persistent epigenetic changes are associated with enhanced transcriptional

responses to recurrent stresses at later stages (Larsen et al., 2021; Naik et al., 2017; Netea et al., 2016). Many studies that examined the effects of maternal nutritional stimuli on livers of offspring have shown long-term changes in epigenetic modifications, especially in DNA methylation (Chen et al., 2022; de Jesus et al., 2020; Maude et al., 2021; Suter et al., 2014). However, in the case of mammalian models, it can sometimes be difficult to determine a causal relationship between maternal nutrition and phenotypes of the offspring, as there are multiple pathways through which an imbalance of maternal nutrition can affect the fetus, such as maternal-fetal nutritional transmission via placenta or milk, transmission of maternal inflammatory cytokines and hormones, placental inflammation, microbiota, and maternal nursing behaviors (M. Li et al., 2015; Wesolowski et al., 2017). Previous studies with adult mice tested whether HFD-induced changes in chromatin state persisted when fatty liver was resolved by long-term feeding with normal chow (NC) (Leung et al., 2016; Siersbæk et al., 2017) or bariatric surgery (Ahrens et al., 2013; Du et al., 2017). However, the results are not consistent across studies, possibly due to the differences in experimental conditions such as age, strains, nutritional compositions of diet, and schedules of HFD feeding.

So far, studies on nutritional programming have been conducted mainly with mammalian models including humans, and knowledge in other non-mammalian vertebrates has been very limited. Medaka (Japanese killifish, *Oryzias latipes*) was recently utilized as a vertebrate model to analyze the effects of nutrition (Asaoka et al., 2014). Indeed, HFD feeding in adult medaka was shown to reproducibly lead to the development of fatty liver, obese phenotypes characterized by enlarged adipose tissues, as well as elevation of serum glucose and triglyceride levels, similar to the results observed in mammals (Matsumoto et al., 2010). Importantly, medaka is oviparous; starts

feeding soon after hatching (7 days post fertilization (dpf)), providing a promising platform with which to apply direct nutritional stimulation from early stages. In addition, medaka has several advantages such as high fecundity and availability of high-quality genome data, making it possible to track epigenetic states throughout life (Ichikawa et al., 2017; Iwamatsu, 2004; Kasahara et al., 2007; Takeda and Shimada, 2010). Furthermore, given the long evolutionary distance between teleosts and mammals, studies with medaka could provide insights into conserved and divergent mechanisms of nutritional adaptation among vertebrates.

As a first step toward understanding the effect of early-life nutritional environment on non-mammalian vertebrates, we examined medaka livers which had experienced HFD feeding at larval stages, and assessed which epigenetic changes were induced and whether they persisted even after shifting to NC. Specifically, we performed RNA-seq, ATAC (Assay for Transposase-Accessible Chromatin)-seq, and ChIP (Chromatin immunoprecipitation)-seq of histone modifications (H3K27ac, H3K27me3, and H3K9me3). Under our experimental conditions, medaka responds well to HFD histologically, transcriptionally, and epigenetically to develop fatty liver. While HFD-induced transcriptomic and epigenetic changes almost entirely returned to normal levels as fatty liver was reversed, we found that some changes persisted at several specific loci. We discuss these non-reversible gene loci in light of nutritional programming.

Results

Fatty liver was induced by early-life HFD feeding in medaka, and was reversed following a switch to NC

To investigate the effects of early-life HFD feeding, we reared medaka fish in the following conditions (Fig. 1A). First, male fish were divided into two groups and were fed either NC or HFD for six weeks from the hatching stage (seven weeks of age, termed NC and HFD fish, respectively). Both groups of fish were then fed NC for eight weeks, the period during which it was expected that the fatty liver condition would reverse (15 weeks of age, termed NC-NC and HFD-NC fish, respectively). To prevent competition between individuals and thus to reduce individual fluctuations as much as possible, we reared fish alone (one fish per one glass or plastic tank) for the duration of the study (Fig. 1B). We found that body weight and liver weight were increased in HFD fish, but the difference in both parameters became smaller between NC-NC and HFD-NC fish (Fig. 1C). We dissected livers from the fish in each group and performed histological analyses to check liver phenotypes. We confirmed that HFD fish showed the typical phenotypes of fatty liver, characterized by lipid droplets visualized by HE staining and neutral lipid accumulation detected by Oil Red O staining (Fig. 1D). However, we did not observe any inflammatory cell aggregates, hepatocyte ballooning, or fibrosis, suggesting that six weeks of HFD feeding induces simple steatosis. After eight weeks of NC, those phenotypes disappeared, and no clear histological differences were detected between NC-NC and HFD-NC livers (Fig. 1D). Electron microscopy observation confirmed the accumulation of lipid droplets in the livers of HFD fish, and their disappearance after 8-weeks of NC (Fig. 1E). Thus, we concluded that early-life HFD feeding effectively induced weight gain and fatty liver, which could both be reversed by subsequent long-

term NC feeding in medaka.

Early-life HFD feeding caused drastic changes in hepatocyte gene expression, which

were largely reversed by subsequent NC

We next investigated the effects of early and transient HFD on hepatic gene

expression by RNA-seq. Since liver generally contains various types of cells other than

hepatocytes (e.g., hepatic stellate cells, endothelial cells, cholangiocytes, erythrocytes,

and immune cells), the analysis of whole-liver tissue would complicate the interpretation

of results. Thus, we first established a transgenic medaka strain that expresses GFP

specifically in hepatocytes. For this, we integrated the Tbait-hs-GFP cassette into the

genomic region upstream of the transcriptional start site of *tdo2* (specifically expressed

in hepatocytes) using the CRISPR-Cas9-mediated genome knockin method (Fig. 2A)

(Watakabe et al., 2018). Immunofluorescent staining confirmed that GFP was detected

specifically in hepatocytes (Supplementary Fig. 1).

For transcriptome analysis, the transgenic fish were reared under the same

conditions as shown in Fig. 1A, then GFP-positive hepatocytes were isolated by

fluorescence-activated cell sorting (FACS) and subjected to RNA-seq (Fig. 2A,

Supplementary Fig. 2). Each liver obtained from the fish in the four groups was

individually analyzed (NC n=8; HFD n=7; NC-NC n=15; HFD-NC n=16). Despite

individual rearing, we still observed large variation in gene expression between

individuals, possibly caused by an accumulation of subtle differences during long-term

rearing and/or inherent features of nutrition-related genes (Abu-Toamih Atamni et al.,

2019). Thus, in the following analyses, we applied a less stringent threshold to identify

differentially expressed genes. At seven weeks of age, 629 genes were up-regulated and

850 genes were down-regulated in HFD fish (Fig. 2B, DESeq2, p-value < 0.01). Gene ontology analysis showed that upregulated genes in HFD hepatocytes were enriched for metabolic pathways, mainly lipid and fatty acid metabolic processes, while downregulated genes were enriched for translation and peptide biosynthetic processes (Fig. 2C). Upregulation of lipid metabolism genes by HFD feeding was frequently reported in mammalian models of HFD feeding (Czech, 2017; Guan et al., 2018; Siersbæk et al., 2017; Soltis et al., 2017). It is unclear why translation-related genes (especially ribosomal protein genes) were downregulated, but it could be a result of endoplasmic reticulum (ER) stress inhibiting protein synthesis, which was also frequently reported in livers of HFD-fed mice (Lebeaupin et al., 2018). On the other hand, only 81 genes were up-regulated and 156 genes were down-regulated in HFD-NC hepatocytes, relative to NC-NC (Fig. 2D, p-value < 0.01). When a more stringent threshold correcting for multiple testing at FDR < 0.1 was applied, only 15 genes were found to be differentially expressed between NC-NC and HFD-NC hepatocytes after 8 weeks of NC (Supplementary Fig. 3A). In addition, the number of genes persistently up-regulated or down-regulated following the eight weeks of NC is small (nine up-regulated and 25 down-regulated, Fig. 2E, F). These data suggest that HFD feeding in early life stages induces drastic hepatic transcriptional changes similar to mammalian models, and that these changes are largely reversed by subsequent long-term NC feeding.

Early-life HFD induced changes in the epigenetic state, which were largely reversed by subsequent NC

Previous studies have shown that some cell types, including innate immunity cells and epidermal stem cells, experience long-term epigenetic changes in response to

transient inflammatory stresses, while transcriptional activity returns to near-normal levels (Larsen et al., 2021; Naik et al., 2017; Netea et al., 2016). To test if such persistent changes in the epigenetic state are induced by early-life HFD in medaka hepatocytes, we profiled genome-wide chromatin accessibility of FACS-isolated hepatocytes by ATAC-seq (Fig. 2A). Each liver obtained from the fish in the four treatment groups was individually analyzed (NC n=7; HFD n=6; NC-NC n=9; HFD-NC n=6). We confirmed that the distribution of ATAC-seq signal exhibits expected patterns—enrichment at promoters and enhancers (Fig. 3A). To identify ATAC-seq peaks that are differently regulated by diet, we initially called peaks for each dataset, merged the resulting peaks, and then selected reliable peaks based on their presence in at least six of the 28 samples. Then we counted the number of reads inside the defined peaks for each dataset and obtained differently accessible regions by DESeq2. Among the 30,484 open chromatin sites identified throughout the genome, 2,421 regions showed changes in chromatin accessibility soon after early-life HFD feeding (998 more accessible regions and 1,423 less accessible regions in HFD hepatocytes, DESeq2, p-value < 0.01, Fig. 3B). Genomic annotation revealed that 70.7% of the differentially accessible peaks were located outside 2 kb on either side of the transcription start sites (TSS), which is a significantly higher proportion than the distribution observed for all peaks (55.4%, p-value < 2.2e-16, Fisher's exact test). This suggests that there is a higher tendency for HFD-induced changes to affect chromatin accessibility of enhancers (Supplementary Fig. 4A). We next annotated all ATAC-seq peaks to their nearby genes (see Methods) and performed gene ontology analysis. Among the 30,484 peaks, 21,227 peaks were linked to at least one gene, and 13,622 genes were linked to at least one peak. Gene ontology analysis showed that genes related to monocarboxylic acid metabolic process were enriched in the set of genes with

increased accessibility following HFD feeding (Fig. 3C), consistent with the results of

the RNA-seq analysis. Motif analyses using HOMER indicated that transcription factor

motifs that regulate lipid metabolism (Nr2f6/Ppar, Srebp) were enriched, corroborating

the observation that HFD feeding promotes lipid metabolic processes (Fig. 3D). Genes

associated with regulation of biological process, cell differentiation, and signaling were

enriched among those with decreased accessibility in HFD hepatocytes (Fig. 3C). Motifs

of several liver-enriched transcription factors (e.g., forkhead box (Fox), GATA, HLF,

C/EBP) and STAT were enriched for these peaks (Fig. 3D). Eight weeks following the

switch to NC, the number of differentially accessible peaks between dietary groups

decreased, with 100 more accessible and 237 less accessible in HFD-NC compared to

NC-NC (p-value < 0.01, Fig. 3E, Supplementary Fig. 3B). Following the eight weeks NC,

the number of peaks that remained persistently more- or less-accessible was limited

(seven for up-regulated and 43 for down-regulated, Fig. 3F), similar to the trend observed

for transcriptional changes. Consequently, the chromatin state of metabolic genes, such

as lipid metabolism genes, which exhibited altered chromatin accessibility after HFD

feeding, was reversed by the subsequent NC (Fig. 3G).

We next performed ChIP-seq to profile the genome-wide distribution of three

histone modifications (H3K27ac, H3K27me3, H3K9me3), which respectively mark

active promoters and enhancers, repressive polycomb domains, and repressive

heterochromatin. We first confirmed by western blot that the global levels of each histone

modification were unchanged between dietary conditions (Supplementary Fig. 5). We

also confirmed that the distribution of each histone modification exhibited expected

patterns; H3K27ac was highly enriched in active gene promoters and enhancers and

highly correlated with ATAC-seq signals, while H3K27me3 and H3K9me3 were enriched

near repressed developmental genes and intergenic regions, and anti-correlated with ATAC-seq signals (Fig. 3A, Supplementary Fig. 6A). To facilitate immunoprecipitation from a sufficient number of cells, three to seven livers were mixed together for each of the following ChIP-seq analyses, and three biological replicates were carried out.

We identified 394 differentially-enriched peaks of H3K27ac soon after early-life HFD feeding (DESeq2, p-value < 0.01, Fig. 4A, Supplementary Fig. 3C). Similar to the ATAC-seq results, differentially enriched peaks were predominantly located on enhancers (80.2% for differential peaks compared to 62.9% for all peaks, p-value = 3.64e-14, Fisher's exact test, Supplementary Fig. 4B). Gene ontology analyses of nearby genes revealed that genes related to fatty acid metabolism were enriched for HFD-upregulated peaks, consistent with the results from the gene expression and chromatin accessibility analyses (Supplementary Fig. 7A). The number of differential peaks decreased to 112 after long-term reversal to NC (Fig. 4B, Supplementary Fig. 3C), and only a limited number of persistent peaks were observed (one for up- and 10 for down-regulated in HFD-NC, Fig. 4C), indicating that most changes in H3K27ac signal also returned to normal levels following the switch to NC (Supplementary Fig. 7B). Comparison of RNA-seq, ATAC-seq, and H3K27ac ChIP-seq results showed that changes in gene expression, chromatin accessibility, and H3K27ac signal were highly correlated with each other soon after HFD feeding, but the correlation weakened after long-term NC (Fig. 4D, supplementary Fig. 6B, supplementary Table 1).

Finally, we focused on the repressive histone modification marks of H3K27me3 and H3K9me3. However, due to their broad distribution pattern, we were unable to identify clear peaks. As an alternative, we divided the entire genome into 4-kb bins (resulting in 183,528 bins in total), counting the reads inside each bin and performing

differential analysis using DESeq2. Surprisingly, we did not detect any regions

differentially enriched for H3K27me3 soon after HFD feeding (Fig. 4E), suggesting that

the distribution pattern of H3K27me3 is not strongly influenced by HFD feeding. On the

other hand, for H3K9me3, we identified 690 bins with significantly decreased ChIP read

density in HFD livers (DESeq2, padj < 0.1, Fig. 4F). Despite moderate enrichment for

regulation of metabolic process in gene ontology analysis of nearby genes

(Supplementary Fig. 8A), changes in H3K9me3 enrichment did not correspond to gene

expression (Supplementary Fig. 8B) or correlate with chromatin accessibility

(Supplementary Fig. 8B). Notably, after switching to NC, H3K9me3 signal mostly

returned to control levels (Fig. 4F, Supplementary Fig. 8C). Taken together, our results

indicate that early-life HFD feeding causes drastic changes in chromatin accessibility and

histone modifications, particularly H3K27ac and H3K9me3. The majority of these

changes were reversed following a switch to NC, except for the particular loci described

below. Representative examples are shown in Fig. 4G.

Persistently altered loci induced by early-life HFD

Although a recovery of fatty liver was observed, differences in the chromatin

state were still found at 337 loci between NC-NC and HFD-NC fish at 15 weeks of age

(Fig. 3E). Gene ontology analysis revealed that these loci were enriched with genes

related to cell surface receptor signaling pathway and cell differentiation (Fig. 5A),

suggesting potential changes at the levels of cellular or tissue properties in the livers that

had experienced HFD, such as cell-cell interaction and differentiation. The 337 loci were

a combination of persistently changed (i.e., difference in chromatin accessibility was

observed at both seven and 15 weeks of age) and transiently changed (i.e., difference in

chromatin accessibility was observed only at 15 weeks of age) ATAC-seq peaks. We

focused on the 50 persistent peaks (Fig. 3F, Fig. 5B, Supplementary Fig. 9A). While

differences at each locus may be sometimes subtle, we reason that some of these loci

could truly maintain altered chromatin accessibility induced by early-life HFD feeding,

if they are accompanied by consistent changes in gene expression and H3K27ac signal.

In this context, at least 16 peaks among the 50 persistent peaks, were accompanied by

consistent changes in gene expression and/or H3K27ac at seven and/or 15 weeks of age

(Fig. 5B, Supplementary Fig. 9). These include peaks showing long-term changes in

expression of nearby genes (*raftlin*, *hhla2b.1*, and *epha5*, Fig. 5C), and peaks showing

persistent epigenetic changes while gene expression returned to normal levels (*cdh18*,

gadd45a, and two unannotated genes, Fig. 5D). Intriguingly, we found that some of the

50 peaks are located close to genes related to cell signaling (e.g., *dscam1*, *epha5*, *ptprfb*,

hhla2b, *cdh18*, *rasgef1bb*, and *raftlin*), and are associated with liver fibrosis and

hepatocellular carcinoma (HCC) (e.g., *epha5*, *raftlin*, *hhla2b1*, and *gadd45a*, Fig. 5B-D).

Additionally, we identified 11 peaks showing persistent changes in H3K27ac signal, and

10 of them showed consistent changes in expression or ATAC-seq signal at either time

point (Supplementary Fig. 7C, 9B). Therefore, despite the nearly reversible chromatin

state triggered by HFD feeding, there are a number of loci showing persistent changes in

chromatin state following a switch to long-term NC. These are candidate genomic loci

responsible for long-term epigenetic memory of early-life HFD feeding (see Discussion).

Discussion

Several studies conducted in mammals have shown that overnutrition during development and growth periods increases susceptibility to obesity, diabetes, and fatty liver disease in later life (de Jesus et al., 2020; Gregorio et al., 2010; Huang et al., 2017; Kruse et al., 2013; M. Li et al., 2015; Oben et al., 2010; Thompson, 2020). Epigenetic modifications have been proposed as the mechanism underlying the long-term memory of early-life overnutrition (Costello and Schones, 2018). However, the extent to which epigenetic changes induced by early-life overnutrition persist and affect later metabolic programming phenotypes remains controversial across studies (Ahrens et al., 2013; Du et al., 2017; Kim et al., 2019; Leung et al., 2016; Siersbæk et al., 2017). To avoid the complexity of mammalian systems (M. Li et al., 2015; Thompson, 2020; Wesolowski et al., 2017), we developed a new system using medaka as an oviparous animal model, and in this study we tested whether HFD feeding during the early growth period can effectively cause fatty liver phenotypes and to what extent induced transcriptional and epigenomic changes persist. To our knowledge, this is the first study to thoroughly analyze the long-term effect of early-life nutritional environment on the epigenetic state in non-mammalian vertebrates.

Our data demonstrate that HFD induces drastic changes in the hepatic transcriptome, chromatin accessibility, and histone modifications especially in metabolic genes, and that subsequent long-term NC feeding returned most of the changes to normal levels. Our results suggest that the hepatic transcriptome and epigenome of medaka are highly adaptable to changes in the feeding environment. However, we also notably found genomic loci which showed persistent epigenetic changes triggered by early-life HFD feeding (discussed later).

Our results are consistent with several recent studies in adult mice investigating

the persistence of HFD-induced epigenetic changes following weight loss by long-term

NC or bariatric surgery (Du et al., 2017; Kim et al., 2019; Siersbæk et al., 2017). For

example, Siersbæk et al. demonstrated that seven weeks of HFD feeding caused fatty liver

accompanied by drastic changes in the hepatic transcriptome and H3K27ac at enhancers,

which returned to levels comparable with control mice after five weeks of NC and

recovery from fatty liver (Siersbæk et al., 2017). However, Siersbæk et al., found that

chromatin accessibility was not affected by HFD feeding, based on DNase-seq analysis.

On the other hand, our data based on ATAC-seq analysis demonstrates that HFD feeding

causes drastic changes in chromatin accessibility. The reason for the discrepancy is

unclear, but one possibility is the difference in the timing of HFD feeding (i.e., Siersbæk

et al. fed HFD to 12 weeks old adult mice, while we started HFD feeding from the

hatching stage in medaka). Similarly, Du et al. surveyed the persistence of HFD-induced

changes in chromatin accessibility long after vertical sleeve gastrectomy by FAIRE-seq,

and found that chromatin accessibility returned to normal levels after weight loss (Du et

al., 2017). In addition, Kim et al. tested the persistence of HFD-induced changes in DNA

methylation following nine weeks of NC, and again showed that the majority of hyper-

and hypomethylated CpGs were reversed after weight loss, although ~5.7% of CpGs (152

CpG in total) showed persistent changes, similar to our results (Kim et al., 2019). In

contrast, Leung et al. demonstrated that HFD-induced changes in chromatin accessibility

persisted eight weeks after a switch to NC; however, this difference can be explained by

the incomplete reversal of obese phenotypes in their study (i.e., heavier body weight and

more lipid accumulation in livers in the HFD-NC mouse group) (Leung et al., 2016).

Taking into account these previous studies, it is reasonable to speculate that HFD-induced

changes in epigenetic state are largely reversible upon recovery from fatty liver phenotypes. The high plasticity of epigenetic states in hepatocytes seems plausible, as hepatocytes need to dynamically regulate metabolic genes according to the feeding/fasting cycle, to maintain whole-body metabolic homeostasis (Bideyan et al., 2021). Indeed, dynamic circadian regulation of enhancer RNA production and H3K27ac accumulation was reported in livers of mice (Fang et al., 2014).

So far, studies of nutritional programming have been mainly conducted in mammals, because of clinical relevance. However, the long-term effects of early-life nutritional intervention have also been widely investigated in teleosts, with the aim of producing high-quality farmed fish at low cost (Panserat et al., 2019). The strategy is based on the concept that the first feeding stage, the transition from endogenous- to exogenous feeding, exhibits a high degree of molecular plasticity for intermediary metabolism, leading long-term metabolic traits that persist throughout life. One example is feeding of an early-life carbohydrate-rich (CH-rich) diet to carnivorous fish (Fang et al., 2014; Geurden et al., 2014, 2007; Gong et al., 2015; Rocha et al., 2016), which generally exhibit poor utilization of carbohydrates due to their weak glucose metabolism such as glucose intolerance, poor carbohydrate digestion, absorption, and utilization (Kamalam et al., 2017). In these studies, a CH-rich diet at first feeding resulted in long-term changes in carbohydrate metabolism in response to a secondary CH-rich diet challenge, such as lower expression of gluconeogenesis genes (zebrafish (Fang et al., 2014) and Siberian sturgeon (Gong et al., 2015)), increased expression of amylase and maltase (rainbow trout (Geurden et al., 2007)), and improved uptake and catabolism of carbohydrates (gilthead seabream (Rocha et al., 2016)). Another example is a plant-based diet at the initial phase of feeding (Balasubramanian et al., 2016; Clarkson et al., 2017;

Geurden et al., 2013; Michl et al., 2017; Vera et al., 2017), which is rich in various

secondary metabolites and poor in long-chain poly-unsaturated fatty acids (PUFAs),

leading to poor growth in some carnivorous and/or marine species (Panserat et al., 2019;

Vera et al., 2017). A first feeding of a plant-based diet to rainbow trout and Atlantic

salmon resulted in improved growth rate and energy utilization during the secondary

plant-based diet challenge, compared to fish fed a marine based diet (Clarkson et al.,

2017; Geurden et al., 2013). Microarray expression analysis in the liver of juvenile

Atlantic salmon challenged with a plant-based diet revealed the upregulation of genes

involved in intermediary metabolism (e.g., glucose metabolism, fatty acid biosynthesis,

and elongation), protein metabolism, and antioxidant defense of cells, suggesting an

improved utilization of a plant-based diet at the molecular level (Vera et al., 2017). These

examples clearly demonstrate that long-term changes can occur in the metabolism of fish.

This somewhat contrasts with our results in medaka liver where the effects of early-life

HFD are mostly reversible. However, it is worth noting that dietary habits are quite

diverse among fish and this is reflected in the manner of nutritional adaptation; narrow

(e.g., carnivorous and/or marine) vs omnivorous feeding freshwater fish (Medaka). To

understand the mechanisms of nutritional programming in fish, it will be intriguing to

apply genome-wide epigenome profiling to the above-mentioned conditioned fish, and

our study provides reference data for those future studies.

Although HFD-induced changes in chromatin state are largely reversible in

medaka, we found a number of ATAC-seq peaks showing non-reversible changes long

after a switch to NC (Fig. 5, Supplementary Fig. 9). At least 16 peaks were accompanied

by consistent changes in gene expression and/or H3K27ac at either seven and/or 15 weeks

of age, suggesting that they underwent persistent epigenetic changes (Fig. 5B,

Supplementary Fig. 9). Some peaks showed consistent changes in expression of nearby genes at both time points (e.g., *raftlin*, *hhla2b.1*, and *epha5*), suggesting that the epigenetic state of chromatin has a direct impact on genes near those peaks (Fig. 5C). On the other hand, a majority of peaks were not associated with changes in gene expression at 15 weeks of age. We speculate that at least some of these peaks exhibit latent epigenetic changes, impacting nearby genes under some specific conditions. The fact that most of those persistent peaks were categorized as enhancers supports this idea (Supplementary Fig. 4), since chromatin accessibility at enhancers does not necessarily reflect gene expression (Klemm et al., 2019; Larsen et al., 2021; Netea et al., 2016). These peaks could be candidates with long-term epigenetic change triggered by early-life HFD feeding.

Interestingly, some of the persistently changed ATAC-seq peaks are located close to genes related to cell signaling and associated with liver fibrosis and hepatocellular carcinoma. For example, *epha5*, a persistently downregulated gene in terms of expression and chromatin accessibility in HFD-NC fish, has been reported to be associated with several types of cancers (S. Li et al., 2015), including hepatocellular carcinoma (Wang et al., 2019). Eph receptors, including Epha5, constitute the largest subfamily of receptor tyrosine kinases and mediate a variety of cellular processes such as cell migration, adhesion, proliferation, and differentiation, in response to their ligands ephrins (Kania and Klein, 2016; Shu et al., 2022). Ephrin/Eph signaling plays pivotal roles in various developmental contexts such as axon guidance, tissue boundary formation, and angiogenesis (Kania and Klein, 2016), but its dysregulation is frequently observed in various types of cancers (Shu et al., 2022). Similarly, *raftlin*, a persistent upregulated gene in this study, tends to be upregulated in patients of HCC compared to healthy controls (Sameh Boshra et al., 2019). Raftlin is one of the major lipid raft proteins and is known

to function in adaptive immunity by mediating B-cell receptor (BCR) and T-cell receptor (TCR) signaling in lymphocytes, as well as in innate immunity by mediating uptake of ligand-bound Toll-like receptors (TLR) 3/4 via endocytosis in dendritic cells and macrophages (Matsumoto and Tatematsu, 2016; Saeki et al., 2003). Although the function of Raftlin in liver, as well as its association with HFD feeding and obesity, has not yet been tested, persistent upregulation of this gene could imply latent activation of innate immunity, which affects the propensity for liver inflammation and fibrosis. In addition, some other persistent genes have been implicated in liver damage and fibrosis (e.g., *dax1*, *gadd45a*, *tsc22d3*, and *miR-122*) (Flamini et al., 2021; Hsu et al., 2012; Tanaka et al., 2017; Yun et al., 2022) and HCC (e.g., *spred2*, *scube1*, *ptprfb*, *tmod1*, *asap3*, *zbtb16*, and *hhla2b1*) (Bera et al., 2014; Feng et al., 2020; Gao et al., 2023; Hui et al., 2015; Luo et al., 2021; Vi et al., 2008; Zhao et al., 2022). Thus, the persistently-affected genes identified in our study could represent latent changes in the cellular state of the liver that may contribute to the increased propensity for fibrosis and HCC development later in life as a result of early-life HFD feeding in medaka. Intriguingly, studies with mice also reported persistent changes in DNA methylation around genes involved in cell signaling and development in adult offspring of HFD-fed dams (Suter et al., 2014; Wankhade et al., 2017). They are associated with a higher propensity to develop liver fibrosis (Wankhade et al., 2017), suggesting that these may represent conserved features of long-term nutritional effects in mice and medaka. To reveal the roles of these persistent genes, further phenotypic analyses focusing on aged medaka, as well as analyses focusing on the interaction of hepatocytes with other types of liver cells (e.g., hepatic stellate cells, endothelial cells, and macrophages) are needed. Finally, further studies with non-mammalian vertebrates with various dietary habits will provide unique and conserved

insights into the mechanism and evolution of nutritional adaptation.

Methods

Animals

Himedaka strain d-rR or *tdo2*:GFP transgenic medaka were used in this study.

All experimental procedures and animal care were performed according to the animal

ethics committee of the University of Tokyo. Fish eggs were collected and incubated at

28.5°C for seven days until hatching. After then, fish were raised in a circular glass tank

with 80 mm diameter and 50 mm depth individually (a single fish per a single tank) for

six weeks to avoid competition between individuals. During this period, fish were fed

either normal chow (Hikari rabo 130, Kyo-rin, approx. 20.0% calories from fat, 61.0%

calories from protein and 19.0% calories from carbohydrate) or high-fat diet (High Fat

Diet 32; CLEA Japan Inc., 56.7% calories from fat, 20.1% calories from protein and

23.2% calories from carbohydrate) twice a day. To prevent high mortality, a small amount

of NC was supplemented to the HFD group fish for the morning feed. All glass cups were

maintained in a large water bath, with a constant temperature of 29.0~29.5°C. After seven

weeks of age, fish were transferred to small plastic tanks with 130 mm width, 90 mm

depth, and 85 mm height, individually (a single fish per single tank) in tap water and fed

NC (Tetramin Super, Tetra, approx. 26.9% calories from fat, 58.7% calories from protein

and 14.4% calories from carbohydrate) twice a day for eight weeks. Throughout fish

rearing, room temperature was maintained at 27°C±1°C, with a strict light-dark cycle

(lights turned on at 8:30 am and turned off at 10:00 pm). In all experiments, fish were

sampled between 1:00 to 3:00 pm after undergoing a period of fasting that began in the

afternoon of the preceding day.

Histology

The fish were anesthetized with ice-cold water, and their livers were dissected

for histological analyses. For hematoxylin and eosin (HE) staining, the liver tissue was

fixed overnight at room temperature with gentle shaking using Bouin's fixative.

Afterward, the tissues were dehydrated with ethanol, cleared using Clear Plus (Falma),

infiltrated with paraffin at 65℃, and embedded in it. 5 µm-thick sections were cut and

stained with HE. For Oil Red O staining, the liver was fixed with 4% paraformaldehyde

in PBS for overnight at 4℃ with gentle shaking. After washing twice with PBS, liver was

transferred to 30% sucrose/PBS and rotated overnight at 4℃, and embedded in O.C.T

compound (Sakura Finetek Japan). 10 µm-thick frozen sections were cut, dried, and

stained with 0.18% Oil Red O/60% isopropanol for 10 min at 37℃.

Transmission electron microscopy

The samples were fixed with 2% paraformaldehyde (PFA) and 2%

glutaraldehyde in 0.1 M phosphate buffer (PB) pH 7.4 at 4℃ overnight. After washing

with 0.1 M PB, liver tissues were postfixed with 2% osmium tetroxide (OsO_4) in 0.1 M

PB at 4℃ for 2h. After dehydration in graded ethanol solutions (50%, 70%, 90%, and

100%), livers were infiltrated with propylene oxide (PO), a 70:30 mixture of PO and resin,

and embedded in 100% resin. Ultra-thin sections (70 nm-thick) were cut, mounted on

copper grids, and stained first with 2% uranyl acetate, then with Lead stain solution

(Sigma). The grids were observed by transmission electron microscopy (JEM-1400Plus;

JEOL Ltd.) at an acceleration voltage of 100 kV. All procedures after fixation were

performed by Tokai Electron Microscopy. Inc. (Aichi, Japan).

Generation of *tdo2*:GFP transgenic medaka

Tdo2:GFP transgenic medaka was generated as described in Watakabe et al.,

2018 (Watakabe et al., 2018). Donor plasmid (Tbait-hs-GFP) and a pDR274 plasmid

containing sgRNA sequence targeting Tbait were kindly provided by Dr. Higashijima at

the National Institute for Basic Biology (NIBB). Cas9 mRNA was synthesized by

digesting the pCS2-Cas9 plasmid by NotI, followed by transcription using an

mMESSAGE mMACHINE SP6 kit (Life Technologies). A sgRNA targeting *tdo2* was

designed using CCTop (Stemmer et al., 2015), using the 1000 bp region upstream of the

transcription start site of *tdo2* as a query. sgRNA was synthesized as follows; first, guide

oligos were annealed by heating to 98℃ for 30 seconds, followed by ramping down to

25℃ at a rate of -0.1℃/sec. Then, 40 ng of pDR274e plasmid and ~0.1 pmol of annealed

oligos were mixed with Ligation High ver. 2 (TOYOBO) and BsaI-HF (NEB) and

incubated at 37℃ for 5 min and 16℃ for 5 min for three cycles, to insert the annealed

oligo to the pDR274e plasmid. After amplification of the template DNA by PCR, sgRNA

was synthesized using NEB's HiScribe T7 Quick High Yield RNA Synthesis Kit (NEB).

For generation of transgenic fish, a cocktail of sgRNAs, donor plasmid, and Cas9 mRNA

(9 ng/µl of sgRNA for digesting Tbait, 18 ng/µl of sgRNA for digesting genome DNA, 9

ng/µl of donor plasmid and 200 ng/µl of Cas9 mRNA) was injected into one-cell stage

medaka embryos.

Immunohistochemistry

Liver was dissected from *tdo2*:GFP medaka and fixed in 4% PFA/PBS for two hours at room temperature. After washing with PBS, the liver was dehydrated with ethanol, cleared using Clear Plus (Falma), infiltrated with paraffin at 65℃, and embedded in it. 5 μm-thick sections were cut and immunostained with anti-GFP antibody (living colors, Clontech, 632592) and secondary anti-rabbit IgG with Alexa fluor 488 (Thermo Fisher). Images were captured by Zeiss LSM700 confocal microscopy.

Isolation of GFP-positive hepatocytes for RNA-seq and ATAC-seq

Livers were dissected from *tdo2*:GFP medaka and minced into small pieces on ice using a razor blade. Then, cells were dissociated into individual cells in 0.25% (w/v) trypsin (Nacalai Tesque)/PBS at 37℃ for 5 min, and dissociation was stopped by adding the same volume of 15% (v/v) FBS/Leiboviz's L-15 (Life Technologies). After gentle pipetting 30 times and filtration using pluriStrainer Mini 40 μm (pluriSelect), cells were centrifuged again and resuspended with 15% FBS/L-15. GFP-positive cells (hepatocytes) were sorted using FACSAria III (BFD Biosciences). Dead cells were stained with propidium iodide (Life Technologies) and removed during cell sorting.

RNA-seq

Total RNA was extracted from 50,000 of the sorted hepatocytes using RNeasy Mini Kit (Qiagen). mRNA was enriched by poly-A capture and mRNA-seq libraries were generated using KAPA Stranded mRNA-seq Kit (KAPA Biosystems). Libraries were sequenced using the Illumina HiSeq 1500 platform.

ATAC-seq

ATAC-seq was performed as described in Buenrostro et al. 2015, with some

modifications (Buenrostro et al., 2015). 50,000 of the sorted hepatocytes were washed

with PBS and resuspended with ice-cold lysis buffer (10 mM Tris-HCl, pH 7.4; 10 mM

NaCl; 3 mM MgCl$_2$; 0.1% (v/v) Igepal CA-630) to extract nuclei. After centrifugation at

500×g, 4°C for 10 min, nuclei pellet was resuspended with the transposition reaction mix

(25 µl of 2x reaction buffer and 2.5 µl of Tn5 transposase from Nextera Sample

Preparation Kit (Illumina), mixed with 22.5 µl nuclease-free water) and incubated at 37°C

for 30 min. Following transposition, DNA was purified using a Qiagen MinElute PCR

Purification Kit (Qiagen) and eluted with 10 µl of Buffer EB. Two sequential PCRs were

performed to enrich small DNA fragments. First, nine-cycles of PCR were performed

using indexed primers from a Nextera Index kit (Illumina), and amplified DNA was size-

selected to a size of <500 bp using AMPure XP beads (Beckman Coulter). Then, seven-

cycles of PCR were performed using the same primers as the first PCR and purified by

AMPure XP beads. Libraries were sequenced using the Illumina HiSeq 1500 platform.

ChIP-seq

Liver chromatin lysate was prepared as described in Schmidt et al. 2009

(Schmidt et al., 2009), with some modifications. First, 6–8 mg of liver (from 2–4 fish)

was cut into small pieces and transferred to 1.4 ml of PBS. Then, cross-linking was

performed by adding formaldehyde to a final concentration of 1% (v/v) and rocking at

room temperature for 8 min. Cross-linking reaction was quenched by adding glycine to a

final concentration of 125 mM. To prevent protein degradation, 20 mM sodium butyrate,

1 mM PMSF and 1×cOmplete[TM] EDTA-free Protease Inhibitor Cocktail (Roche) were

added throughout the above procedures. After washing with PBS twice, tissue was

dounced first with the loose pestle 20 times and later with the tight pestle 50 times (DWK

Life Sciences Inc.) in ice-cold lysis buffer 1 (50 mM HEPES-KOH, pH = 7.6; 140 mM

NaCl; 1 mM EDTA; 10% glycerol; 0.5% NP-40; 0.25% Triton-X100; 20 mM sodium

butyrate; 1 mM PMSF; 1×cOmplete). After incubation on ice for 10 min, lysate was

filtered with pluriStrainer Mini 40 μm (pluriSelect) and washed with the lysis buffer 1

once. The pellet was dissociated with the lysis buffer 2 (10 mM Tris-HCl, pH = 8.0; 200

mM NaCl; 1 mM EDTA; 0.5 mM EGTA; 20 mM sodium butyrate; 1 mM PMSF;

1×cOmplete), rocked at 4°C for 5 min, and centrifuged. Pellets were dissociated with lysis

buffer 3 (50 mM Tris-HCl, pH = 8.0; 10 mM EDTA; 1% SDS; 20 mM sodium butyrate;

1 mM PMSF; 1×cOmplete) and sonicated to sizes ranging from 100–500 bp using

Covaris S220 with the following parameters: Peak Power, 105; Duty Factor, 4.0; cycles

per burst, 200; duration, 12.5 min.

As a positive control for ChIP experiments, zebrafish chromatin lysate derived

from BRF41 fibroblasts was added so that the ratio of medaka and zebrafish DNA was

approximately 10:1. Chromatin lysates were diluted with RIPA buffer (10 mM Tris-HCl,

pH = 8.0; 140 mM NaCl; 1 mM EDTA; 1 mM EGTA; 1% Triton X-100; 0.1% SDS; 0.1%

Na-Deoxycholate; 20 mM sodium butyrate; 1 mM PMSF; 1×cOmplete protease

inhibitor) and rocked with the antibody/protein A Dynabeads complex overnight at 4°C.

After washing three times with RIPA buffer and once with TE buffer (10 mM Tris-HCl,

pH = 8.0; 10 mM EDTA, pH = 8.0), chromatin was de-crosslinked by incubating in lysis

buffer 3 at 65°C overnight. After removing beads, eluted samples were treated with RNase

A for two hours at 37°C, followed by proteinase K for two hours at 55°C. DNA was

purified by phenol/chloroform extraction and ethanol precipitation. Input DNA was

simultaneously treated from the elution step. ChIP-seq libraries were constructed using

KAPA Hyper Prep Kit (KAPA Biosystems). Antibodies used in this ChIP experiment are as follows; H3K27ac, ab4729 (abcam); H3K27me3, C15410069 (diagenode); H3K9me3, ab8898 (abcam). Libraries were sequenced using the Illumina HiSeq 1500 platform.

Immunofluorescence western blotting

Dissected livers were weighed and homogenized in 1×Laemmli sample buffer to the concentration of 1 mg tissue/50 µl buffer. Lysates were separated by 10% polyacrylamide gels and transferred onto PVDF membranes (Immobilon-FL, Millipore). After blocking with 5% skim milk/TBS for 1h at room temperature, membranes were incubated with primary antibodies (rabbit anti-H3K27ac antibody (abcam ab4729), rabbit anti-H3K27me3 antibody (diagenode C15410069) or rabbit anti-H3K9me3 antibody (abcam ab8898) combined with mouse anti-H3 antibody (active motif 39763), 1:2000 dilution each) overnight at 4°C. After several washes, membranes were incubated with secondary antibodies (IRDye 800CW Donkey anti-rabbit IgG (LI-COR) and IRDye 680RD Donkey anti-mouse IgG (LI-COR), 1:10000 dilution) for one hour at room temperature. Visualization of protein bands and quantification of signal intensities were performed using ODYSSEY CLx (LI-COR). The quantity of each histone modification was calculated by dividing the signal intensity of histone modification by the signal intensity of histone H3.

RNA-seq data processing

Sequenced reads were preprocessed to remove low-quality reads/bases using Trimmomatic v0.33 (Bolger et al., 2014) with the following parameters: SLIDINGWINDOW:4:15, LEADING:20, TRAILING:20, MINLEN:20. Trimmed reads

were mapped to the Hd-rR medaka reference genome (Ensembl ASM223467v1.95

(Ichikawa et al., 2017)) by STAR (Dobin et al., 2013). Reads with mapping quality

(MAPQ) larger than or equal to 20 were used for further analyses. Differentially

expressed genes were obtained by DESeq2 (Love et al., 2014), and gene ontology

analyses of differentially expressed genes were performed using PANTHER v17.0

(Thomas et al., 2022), putting all expressed genes in medaka hepatocytes as background.

ATAC-seq data processing

Sequenced reads were preprocessed to remove low-quality reads/bases using

Trimmomatic v0.33 (Bolger et al., 2014) with the following parameters:

SLIDINGWINDOW:4:15, LEADING:20, TRAILING:20, MINLEN:20. Trimmed reads

were mapped to the Hd-rR medaka reference genome (Ensembl ASM223467v1.95

(Ichikawa et al., 2017)) by BWA (Li and Durbin, 2009). Reads with mapping quality

(MAPQ) larger than or equal to 20 were used for further analyses. Peak calling was

performed for each data by MACS2 v2.1.0 (Zhang et al., 2008) with the following

options: --nomodel --extsize 200 --shift -100 -g 600000000 -q 0.01 -B --SPMR. Peaks

common to at least six of the 28 samples (30,484 in total) were used for further analyses.

The number of reads in each peak was counted by annotatePeaks command of HOMER

(Heinz et al., 2010), and differentially accessible peaks were obtained by DESeq2, based

on the read counts inside peaks. Motif analyses of differentially accessible peaks were

performed using HOMER. Annotation of the peaks to nearby genes was performed by

the following definition: Firstly, ATAC-seq peaks were annotated with the genes in which

TSS is located within 2 kb of either end of the ATAC-seq peak. If no TSS was found

within 2 kb of either end of the ATAC-seq peak, then we annotated the ATAC-seq peaks

with the overlapping genes (if any). ATAC-seq peaks can be associated with multiple

genes when several TSSs or genes are closely located (e.g., bidirectional promoters).

Peaks that are located inside 2 kb on either side of the TSS are categorized as promoters,

and the others as enhancers. Gene ontology analyses of genes near differentially

accessible peaks were performed using PANTHER v17.0 (Thomas et al., 2022), putting

all genes annotated to at least one of the ATAC-seq peaks as background (13,622 genes

in total).

ChIP-seq data processing

Sequenced reads were preprocessed to remove low-quality reads/bases using

Trimmomatic v0.33 (Bolger et al., 2014) with the following parameters:

SLIDINGWINDOW:4:15, LEADING:20, TRAILING:20, MINLEN:20. Trimmed reads

were aligned to the concatenated Hd-rR medaka (Ensembl ASM223467v1.95 (Ichikawa

et al., 2017)) and zebrafish (Ensembl GRCz11.95) genomes by BWA (Li and Durbin,

2009). Reads mapped to the medaka genome with mapping quality (MAPQ) larger than

or equal to 20 were used for further analyses. For H3K27ac, peak calling was performed

for each dataset by MACS2 v2.1.0 (Zhang et al., 2008) with the following options: --

broad -g 600000000 -q 0.01 -B --SPMR. Then peaks were merged for all the 12 datasets,

and 35,722 peaks were obtained in total. Differential analyses, gene annotation, and gene

ontology analyses were performed in the same methods as ATAC-seq. For H3K27me3

and H3K9me3, differentially enriched regions were obtained by dividing the medaka

genome into 4-kb bins, counting the number of reads at each bin, and performing DESeq2.

Normalization and visualization of bedgraphs

Counts per million (CPM)-normalized bedgraphs were generated by MACS2

(Zhang et al., 2008) with the following options. ATAC-seq, -g 600000000 -q 0.01 -B --

SPMR --nomodel --extsize 200 --shift -100; H3K27ac ChIP-seq, --broad -g 600000000 -

q 0.01 -B --SPMR; H3K27me3 and H3K9me3 ChIP-seq, --broad -g 600000000 -q 0.01 -

B --SPMR --nomodel --extsize 190. Owing to the variation of peak height in the genome

browser view, possibly due to the difference in signal/noise ratio among samples, we

further performed normalization of ATAC-seq and ChIP-seq track data. First, we

determined the scaling factors for visualization of ChIP-seq/ATAC-seq data by linear

regression procedure as follows. Here, we only used regions with stable signals (i.e., read

count). Started with all 2 kb regions around TSS (for ATAC-seq, and K27ac marks) or

peak regions consistent across the libraries (for K27me3 and K9me3 marks), we excluded

a region if the average signal was within the top 100 or below the global average (average

across all regions and libraries). We also excluded a region if the variation of signals was

within the top 10%. Then, the signals of the remaining regions for each library were

regressed against the averages across libraries. The resulting coefficient (slope) was

identified as the scaling factor for each library. Finally, we divided the track data by the

calculated scaling factors.

Author contributions

Y.I., S.M., and H.T. planned and supervised this research. Y.I. and Y.K. established fish-

rearing conditions. Y.K. performed histological analyses of liver tissues. Y.K. created the

tdo2-GFP transgenic medaka. Y.I. and Y.K. performed RNA-seq library preparation and

sequencing. Y.K. performed ATAC-seq library preparation and sequencing. Y.I. and T.W.

performed ChIP-seq library preparation and sequencing. Y.I. performed western blotting.

Y.I., Y.S., and Y.K. performed data analyses of RNA-seq, ATAC-seq, and ChIP-seq data.

Y.S. prepared normalized track views of ATAC-seq and ChIP-seq data. Y.I. and H.T. wrote

the manuscript.

Acknowledgments

We thank Prof. K. Kamimura, Dr. R. Goto, and Prof. S. Terai for the help in establishing

fish rearing conditions, Dr. Y. Kimura and Prof. S. Higashijima for kindly providing us

with plasmids for CRISPR-Cas9-mediated gene knockin, and Mrs. M. Sakamoto for fish

care. We acknowledge Tokai Electron Microscopy for performing electron micrographs

of liver tissue. Flow cytometry was performed in the IMSUT FACS Core laboratory. We

acknowledge the IMSUT FACS Core laboratory for assistance with flow cytometry

analysis. (FACS). This work was supported by AMED CREST, JST (Grant No:

JP23gm1110007).

Competing financial interests

The authors declare no competing financial interests.

Data access

Sequencing data generated in this study has been submitted to the DDBJ BioProject

database under accession number PRJDB15283.

Figure 1: Early-life HFD feeding induces fatty liver in medaka, which is reversed by

subsequent NC feeding.

(A) Experimental design of early-life HFD feeding in medaka, where male fish were fed

either HFD or NC for six weeks from hatching stage, followed by eight weeks of NC for

both groups. **(B)** All fish were maintained individually to minimize variation of the speed

of growth. **(C)** Body weight and liver weight at seven and 15 weeks of age. **(D)**

Representative images of liver sections stained with H&E and Oil Red O, and **(E)**

transmission electron microscopy (TEM) of liver sections from NC, HFD, NC-NC, and

HFD-NC medaka. Scale bars: 100 μm in (D) and 5 μm in (E).

Figure 2: **Early-life HFD feeding-induced changes in medaka hepatocyte gene expression are largely reversed by subsequent NC feeding.**

(A) A schematic of *tdo2*:GFP transgenic medaka construction using CRISPR-Cas9-mediated gene knockin and sorting of GFP-positive hepatocytes. Scale bar: 100 µm **(B)** RNA-seq results showing differentially expressed genes in hepatocytes of HFD fish compared to NC. X-axis: log2 fold change (HFD/NC) of expression values, y-axis: –log10 p-value. **(C)** Gene ontology analyses of upregulated (upper) and downregulated (lower) genes by HFD for biological processes using PANTHER 17.0. Fold enrichment (orange) and –log10 FDR (bar) are displayed. **(D)** Differentially expressed genes in hepatocytes of HFD-NC fish compared to NC-NC. **(E)** Venn diagram of differentially expressed genes at seven and 15 weeks of age. **(F)** A heatmap of differentially expressed genes at seven weeks of age. Log2-transformed, and Z-transformed, DESeq2 normalized read counts are displayed.

Figure 3: Early-life HFD feeding-induced changes in medaka hepatocyte chromatin

accessibility are largely reversed by subsequent NC feeding.

(A) A representative track view of RNA-seq, ATAC-seq, and ChIP-seq (H3K27ac,

H3K27me3, and H3K9me3) signals for hepatocytes/livers for each of the dietary

conditions. **(B)** Differentially accessible peaks in HFD fish hepatocytes compared to those

in NC fish by ATAC-seq. X-axis: log2 fold change (HFD/NC) of normalized read counts

within peaks, y-axis: –log10 p-value. **(C)** Gene ontology analysis of genes close to peaks

with increased or decreased chromatin accessibility in HFD fish. Fold enrichment

(orange) and –log10 FDR (bar) are displayed. **(D)** Motif analysis of peaks with increased

(upper) and decreased (lower) accessibility, inferred by HOMER v4.11. **(E)** Differentially

accessible peaks in HFD-NC fish hepatocytes compared to NC-NC fish. **(F)** Venn

diagram of differentially accessible peaks at seven and 15 weeks of age. **(G)**

Representative examples of ATAC-seq peaks at the promoters of *fads2*, *srebf1*, and

cyp2r1 genes.

Figure 4: Early-life HFD feeding-induced changes in medaka liver histone modifications are largely reversed subsequent NC feeding.

(A) Differentially-H3K27ac enriched peaks in livers of HFD-fed fish compared to NC by ChIP-seq. X-axis: log2 fold change (HFD/NC) of normalized read counts within peaks, y-axis: –log10 p-value. **(B)** Differentially-H3K27ac-enriched peaks in livers of HFD-NC fish compared to NC-NC. **(C)** Overlaps of differentially-H3K27ac-enriched peaks at seven and 15 weeks of age. **(D)** Overlaps of differentially expressed, differentially accessible, and differentially-H3K27ac-enriched genes, at seven and 15 weeks of age. **(E)** Differentially-H3K27me3-enriched 4-kb bins in livers of HFD fish relative to NC (left) and HFD-NC relative to NC-NC (right). **(F)** Differentially-H3K9me3-enriched 4-kb bins in livers of HFD fish relative to NC (left) and HFD-NC relative to NC-NC (right). **(G)** Representative examples of reversible peaks (i.e., genomic regions where epigenetic states were changed soon after HFD feeding but were returned to normal levels by the following NC feeding). Left, *fads2*; right, *s100a14*.

A

ATAC-seq differently-accessible peaks at 15 weeks of age (337 peaks → 194 genes)

GO biological process complete	#list of genes	#genes in input	Fold enrichment	raw P-value	FDR
cell surface receptor signaling pathway (GO:0007166)	626	24	2.69	1.45E-05	1.12E-01
cellular developmental process (GO:0048869)	994	29	2.04	2.64E-04	2.04E-01
transmembrane receptor protein serine/threonine kinase signaling pathway (GO:0007178)	55	6	7.64	2.09E-04	2.32E-01
cell differentiation (GO:0030154)	988	29	2.06	2.46E-04	2.38E-01
BMP signaling pathway (GO:0030509)	34	5	10.3	2.04E-04	2.64E-01
SMAD protein signal transduction (GO:0060395)	21	4	13.35	3.81E-04	2.68E-01

B ATAC-seq persistent peaks (50)

C

D

Figure 5: Genes showing persistent changes in epigenetic state following long-term NC.

(A) Gene ontology analysis of genes close to differentially accessible ATAC-seq peaks between HFD-NC and NC-NC fish (194 genes from 337 peaks). **(B)** A heatmap of persistent ATAC-seq peaks after reversal to NC (50 peaks in total). Log2-transformed, and Z-transformed, DESeq2 normalized read counts of ATAC-seq at each peak are displayed. DESeq2 results of RNA-seq and H3K27ac ChIP-seq of nearby genes/peaks are displayed on the right (p-value < 0.01). **(C, D)** Representative examples of ATAC-seq peaks showing persistent changes in chromatin accessibility, accompanied by long-term changes in expression of nearby genes (C) or gene expression that returned to normal level after a switch to NC (D).

References

Abu-Toamih Atamni HJ, Kontogianni G, Binenbaum I, Mott R, Himmelbauer H,

Lehrach H, Chatziioannou A, Iraqi FA. 2019. Hepatic gene expression variations in

response to high-fat diet-induced impaired glucose tolerance using RNAseq

analysis in collaborative cross mouse population. *Mammalian Genome* **30**:260–

275. doi:10.1007/s00335-019-09816-1

Ahrens M, Ammerpohl O, von Schönfels W, Kolarova J, Bens S, Itzel T, Teufel A,

Herrmann A, Brosch M, Hinrichsen H, Erhart W, Egberts J, Sipos B, Schreiber S,

Häsler R, Stickel F, Becker T, Krawczak M, Röcken C, Siebert R, Schafmayer C,

Hampe J. 2013. DNA methylation analysis in nonalcoholic fatty liver disease

suggests distinct disease-specific and remodeling signatures after bariatric surgery.

Cell Metab **18**:296–302. doi:10.1016/j.cmet.2013.07.004

Asaoka Y, Terai S, Sakaida I, Nishina H. 2013. The expanding role of fish models in

understanding non-alcoholic fatty liver disease. *DMM Disease Models and*

Mechanisms. **6**:905–914. doi:10.1242/dmm.016022

Balasubramanian MN, Panserat S, Dupont-Nivet M, Quillet E, Montfort J, le Cam A,

Medale F, Kaushik SJ, Geurden I. 2016. Molecular pathways associated with the

nutritional programming of plant-based diet acceptance in rainbow trout following

an early feeding exposure. *BMC Genomics* **17**:449. doi:10.1186/s12864-016-2804-

1

Barker DJP. 2007. The origins of the developmental origins theory. *Journal of Internal*

Medicine. **261**:412–417. doi:10.1111/j.1365-2796.2007.01809.x

Bera R, Chiou CY, Yu MC, Peng JM, He CR, Hsu CY, Huang HL, Ho UY, Lin SM, Lin

YJ, Hsieh SY. 2014. Functional genomics identified a novel protein tyrosine

phosphatase receptor type f-mediated growth inhibition in hepatocarcinogenesis. *Hepatology* **59**:2238–2250. doi:10.1002/hep.27030

Bideyan L, Nagari R, Tontonoz P. 2021. Hepatic transcriptional responses to fasting and feeding. *Genes & Development* **35**:635–657. doi:10.1101/gad.348340

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: A flexible trimmer for Illumina Sequence Data. *Bioinformatics* **30**:2114–2120.

Brunt EM, Wong VWS, Nobili V, Day CP, Sookoian S, Maher JJ, Bugianesi E, Sirlin CB, Neuschwander-Tetri BA, Rinella ME. 2015. Nonalcoholic fatty liver disease. *Nat Rev Dis Primers* **1**:1–22. doi:10.1038/nrdp.2015.80

Buenrostro JD, Wu B, Chang HY, Greenleaf WJ. 2015. ATAC-seq: A method for assaying chromatin accessibility genome-wide. *Curr Protoc Mol Biol* **2015**:21.29.1–21.29.9. doi:10.1002/0471142727.mb2129s109

Chen B, Du YR, Zhu H, Sun ML, Wang C, Cheng Y, Pang H, Ding G, Gao J, Tan Y, Tong X, Lv P, Zhou F, Zhan Q, Xu ZM, Wang L, Luo D, Ye Y, Jin Li, Zhang S, Zhu Y, Lin X, Wu Y, Jin Luyang, Zhou Y, Yan C, Sheng J, Flatt PR, Xu GL, Huang H. 2022. Maternal inheritance of glucose intolerance via oocyte TET3 insufficiency. *Nature* **605**:761–766. doi:10.1038/s41586-022-04756-4

Clarkson M, Migaud H, Metochis C, Vera LM, Leeming D, Tocher DR, Taylor JF. 2017. Early nutritional intervention can improve utilisation of vegetable-based diets in diploid and triploid Atlantic salmon (Salmo salar L.). *British Journal of Nutrition* **118**:17–29. doi:10.1017/S0007114517001842

Costello KR, Schones DE. 2018. Chromatin modifications in metabolic disease: Potential mediators of long-term disease risk. *Wiley Interdiscip Rev Syst Biol Med*. **10**:e1416. doi:10.1002/wsbm.1416

Czech MP. 2017. Insulin action and resistance in obesity and type 2 diabetes. *Nat Med.*
23:804–814. doi:10.1038/nm.4350

de Jesus DF, Orime K, Kaminska D, Kimura T, Basile G, Wang CH, Haertle L, Riemens
R, Brown NK, Hu J, Männistö V, Silva AM, Dirice E, Tseng YH, Haaf T, Pihlajamäki
J, Kulkarni RN. 2020. Parental metabolic syndrome epigenetically reprograms
offspring hepatic lipid metabolism in mice. *Journal of Clinical Investigation*
130:2391–2404. doi:10.1172/JCI127502

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M,
Gingeras TR. 2013. STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics*
29:15–21. doi:10.1093/bioinformatics/bts635

dos Santos CO, Dolzhenko E, Hodges E, Smith AD, Hannon GJ. 2015. An Epigenetic
Memory of Pregnancy in the Mouse Mammary Gland. *Cell Rep* **11**:1102–1109.
doi:10.1016/j.celrep.2015.04.015

Du J, Tian J, Ding L, Trac C, Xia B, Sun S, Schones DE, Huang W. 2017. Vertical sleeve
gastrectomy reverses diet-induced gene-regulatory changes impacting lipid
metabolism. *Sci Rep* **7**:5274. doi:10.1038/s41598-017-05349-2

Fang B, Everett LJ, Jager J, Briggs E, Armour SM, Feng D, Roy A, Gerhart-Hines Z, Sun
Z, Lazar MA. 2014. Circadian enhancers coordinate multiple phases of rhythmic
gene transcription in vivo. *Cell* **159**:1140–1152. doi:10.1016/j.cell.2014.10.022

Fang L, Liang XF, Zhou Y, Guo XZ, He Y, Yi TL, Liu LW, Yuan XC, Tao YX. 2014.
Programming effects of high-carbohydrate feeding of larvae on adult glucose
metabolism in zebrafish, Danio rerio. *British Journal of Nutrition* **111**:808–818.
doi:10.1017/S0007114513003243

Feng L, Zhang Y, Yang Q, Guo L, Yang F. 2020. MicroRNA-885 regulates the growth

and epithelial mesenchymal transition of human liver cancer cells by suppressing tropomodulin 1 expression. *Arch Biochem Biophys* **693**:108588. doi:10.1016/j.abb.2020.108588

Flamini S, Sergeev P, Viana de Barros Z, Mello T, Biagioli M, Paglialunga M, Fiorucci C, Prikazchikova T, Pagano S, Gagliardi A, Riccardi C, Zatsepin T, Migliorati G, Bereshchenko O, Bruscoli S. 2021. Glucocorticoid-induced leucine zipper regulates liver fibrosis by suppressing CCL2-mediated leukocyte recruitment. *Cell Death Dis* **12**:421. doi:10.1038/s41419-021-03704-w

Gao T, Yang X, Fujisawa M, Ohara T, Wang T, Tomonobu N, Sakaguchi M, Yoshimura T, Matsukawa A. 2023. SPRED2: A Novel Regulator of Epithelial-Mesenchymal Transition and Stemness in Hepatocellular Carcinoma Cells. *Int J Mol Sci* **24**:4996. doi:10.3390/ijms24054996

Geurden I, Aramendi M, Zambonino-Infante J, Panserat S. 2007. Early feeding of carnivorous rainbow trout (Oncorhynchus mykiss) with a hyperglucidic diet during a short period: effect on dietary glucose utilization in juveniles. *Am J Physiol Regul Integr Comp Physiol* **292**:2275–2283. doi:10.1152/ajpregu.00444.2006

Geurden I, Borchert P, Balasubramanian MN, Schrama JW, Dupont-Nivet M, Quillet E, Kaushik SJ, Panserat S, Médale F. 2013. The positive impact of the early-feeding of a plant-based diet on its future acceptance and utilisation in rainbow trout. *PLoS One* **8**:e83162. doi:10.1371/journal.pone.0083162

Geurden I, Mennigen J, Plagnes-Juan E, Veron V, Cerezo T, Mazurais D, Zambonino-Infante J, Gatesoupe J, Skiba-Cassy S, Panserat S. 2014. High or low dietary carbohydrate: Protein ratios during firstfeeding affect glucose metabolism and intestinal microbiota in juvenile rainbow trout. *Journal of Experimental Biology*

217:3396–3406. doi:10.1242/jeb.106062

Gong G, Xue M, Wang J, Wu X feng, Zheng Y hua, Han F, Liang X fang, Su X ou. 2015. The regulation of gluconeogenesis in the Siberian sturgeon (Acipenser baerii) affected later in life by a short-term high-glucose programming during early life. *Aquaculture* **436**:127–136. doi:10.1016/j.aquaculture.2014.10.044

Gregorio BM, Souza-Mello V, Carvalho JJ, Mandarim-De-Lacerda CA, Aguila MB. 2010. Maternal high-fat intake predisposes nonalcoholic fatty liver disease in C57BL/6 offspring. *Am J Obstet Gynecol* **203**:495.e1–495.e8. doi:10.1016/j.ajog.2010.06.042

Guan D, Xiong Y, Borck PC, Jang C, Doulias PT, Papazyan R, Fang B, Jiang C, Zhang Y, Briggs ER, Hu W, Steger D, Ischiropoulos H, Rabinowitz JD, Lazar MA. 2018. Diet-Induced Circadian Enhancer Remodeling Synchronizes Opposing Hepatic Lipid Metabolic Processes. *Cell* **174**:831–842.e12. doi:10.1016/j.cell.2018.06.031

Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H, Glass CK. 2010. Simple Combinations of Lineage-Determining Transcription Factors Prime cis-Regulatory Elements Required for Macrophage and B Cell Identities. *Mol Cell* **38**:576–589. doi:10.1016/j.molcel.2010.05.004

Hsu SH, Wang B, Kota J, Yu J, Costinean S, Kutay H, Yu L, Bai S, Perle K La, Chivukula RR, Mao H, Wei M, Clark KR, Mendell JR, Caligiuri MA, Jacob ST, Mendell JT, Ghoshal K. 2012. Essential metabolic, anti-inflammatory, and anti-tumorigenic functions of miR-122 in liver. *Journal of Clinical Investigation* **122**:2871–2883. doi:10.1172/JCI63539

Huang Y, Ye T, Liu C, Fang F, Chen Y, Dong Y. 2017. Maternal high-fat diet during pregnancy and lactation affects hepatic lipid metabolism in early life of offspring rat. *J Biosci* **42**:311–319. doi:10.1007/s12038-017-9675-8

Hui AWH, Lau HW, Cao CY, Zhou JW, Lai PBS, Tsui SKW. 2015. Downregulation of

PLZF in human hepatocellular carcinoma and its clinical significance. *Oncol Rep*

33:397–402. doi:10.3892/or.2014.3578

Ichikawa K, Tomioka S, Suzuki Y, Nakamura R, Doi K, Yoshimura J, Kumagai M, Inoue

Y, Uchida Y, Irie N, Takeda H, Morishita S. 2017. Centromere evolution and CpG

methylation during vertebrate speciation. *Nat Commun* **8**:1833. doi:10.1038/s41467-

017-01982-7

Iwamatsu T. 2004. Stages of normal development in the medaka Oryzias latipes. *Mech*

Dev **121**:605–618. doi:10.1016/j.mod.2004.03.012

Kamalam BS, Medale F, Panserat S. 2017. Utilisation of dietary carbohydrates in farmed

fish: New insights on influencing factors, biological limitations and future strategies.

Aquaculture **467**:3–27. doi:10.1016/j.aquaculture.2016.02.007

Kania A, Klein R. 2016. Mechanisms of ephrin-Eph signalling in development,

physiology   and   disease.   *Nat   Rev   Mol   Cell   Biol*   **17**:240–256.

doi:10.1038/nrm.2015.16

Kasahara M, Naruse K, Sasaki S, Nakatani Y, Qu W, Ahsan B, Yamada T, Nagayasu Y,

Doi K, Kasai Y, Jindo T, Kobayashi D, Shimada A, Toyoda A, Kuroki Y, Fujiyama

A, Sasaki T, Shimizu A, Asakawa S, Shimizu N, Hashimoto S-I, Yang J, Lee Y,

Matsushima K, Sugano S, Sakaizumi M, Narita T, Ohishi K, Haga S, Ohta F,

Nomoto H, Nogata K, Morishita T, Endo T, Shin-I T, Takeda H, Morishita S, Kohara

Y. 2007. The medaka draft genome and insights into vertebrate genome evolution.

Nature **447**:714–719. doi:10.1038/nature05846

Kim H, Worsley O, Yang E, Purbojati RW, Liang AL, Tan W, Moses DID, Hartono S, Fan

V, Lim TKH, Schuster SC, Foo RS, Chow PKH, Pettersson S. 2019. Persistent

changes in liver methylation and microbiome composition following reversal of diet-induced non-alcoholic-fatty liver disease. *Cellular and Molecular Life Sciences* **76**:4341–4354. doi:10.1007/s00018-019-03114-4

Klemm SL, Shipony Z, Greenleaf WJ. 2019. Chromatin accessibility and the regulatory epigenome. *Nat Rev Genet* **20**:207–220. doi:10.1038/s41576-018-0089-8

Kruse M, Seki Y, Vuguin PM, Du XQ, Fiallo A, Glenn AS, Singer S, Breuhahn K, Katz EB, Charron MJ. 2013. High-fat intake during pregnancy and lactation exacerbates high-fat diet-induced complications in male offspring in mice. *Endocrinology* **154**:3565–3576. doi:10.1210/en.2012-1877

Langley-Evans SC. 2009. Nutritional programming of disease: Unravelling the mechanism. *Journal of Anatomy.* **215**:36–51. doi:10.1111/j.1469-7580.2008.00977.x

Larsen SB, Cowley CJ, Sajjath SM, Barrows D, Yang Y, Carroll TS, Fuchs E. 2021. Establishment, maintenance, and recall of inflammatory memory. *Cell Stem Cell* **28**:1758–1774.e8. doi:10.1016/j.stem.2021.07.001

Lebeaupin C, Vallée D, Hazari Y, Hetz C, Chevet E, Bailly-Maitre B. 2018. Endoplasmic reticulum stress signalling and the pathogenesis of non-alcoholic fatty liver disease. *J Hepatol* **69**:927–947. doi:10.1016/j.jhep.2018.06.008

Lempradl A, Pospisilik JA, Penninger JM. 2015. Exploring the emerging complexity in transcriptional regulation of energy homeostasis. *Nat Rev Genet* **16**:665–681. doi:10.1038/nrg3941

Leung A, Trac C, Du J, Natarajan R, Schones DE. 2016. Persistent chromatin modifications induced by high fat diet. *Journal of Biological Chemistry* **291**:10446–10455. doi:10.1074/jbc.M115.711028

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**:1754–1760. doi:10.1093/bioinformatics/btp324

Li M, Reynolds CM, Segovia SA, Gray C, Vickers MH. 2015. Developmental programming of nonalcoholic fatty liver disease: The effect of early life nutrition on susceptibility and disease severity in later life. *Biomed Res Int* **2015**:437107. doi:10.1155/2015/437107

Li S, Zhu Y, Ma C, Qiu Z, Zhang X, Kang Z, Wu Z, Wang H, Xu X, Zhang H, Ren G, Tang J, Li X, Guan M. 2015. Downregulation of EphA5 by promoter methylation in human prostate cancer. *BMC Cancer* **15**:18. doi:10.1186/s12885-015-1025-3

López-Pastor AR, Infante-Menéndez J, Escribano Ó, Gómez-Hernández A. 2020. miRNA Dysregulation in the Development of Non-Alcoholic Fatty Liver Disease and the Related Disorders Type 2 Diabetes Mellitus and Cardiovascular Disease. *Front Med (Lausanne)* **7**:527059. doi:10.3389/fmed.2020.527059

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**:550. doi:10.1186/s13059-014-0550-8

Luo M, Lin Y, Liang R, Li Y, Ge L. 2021. Clinical significance of the hhla2 protein in hepatocellular carcinoma and the tumor microenvironment. *J Inflamm Res* **14**:4217–4228. doi:10.2147/JIR.S324336

Matsumoto M, Tatematsu M. 2016. Cell type-specific role of Raftlin in the regulation of endosomal TLR signaling. *Inflamm Cell Signal* **4**:e1326. doi:10.14800/ics.1326

Matsumoto T, Terai S, Oishi T, Kuwashiro S, Fujisawa K, Yamamoto N, Fujita Y, Hamamoto Y, Furutani-Seiki M, Nishina H, Sakaida I. 2010. Medaka as a model for human nonalcoholic steatohepatitis. *DMM Disease Models and Mechanisms* **3**:431–

440. doi:10.1242/dmm.002311

Maude H, Sanchez-Cabanillas C, Cebola I. 2021. Epigenetics of Hepatic Insulin Resistance. *Front Endocrinol (Lausanne)*. **12**:681356. doi:10.3389/fendo.2021.681356

Michl SC, Weis B, Hutchings JA, Schulz C. 2017. Plastic responses by wild brown trout (Salmo trutta) to plant-based diets. *Aquaculture* **476**:19–28. doi:10.1016/j.aquaculture.2017.04.006

Moazed D. 2011. Mechanisms for the inheritance of chromatin states. *Cell* **146**:510–518. doi:10.1016/j.cell.2011.07.013

Naik S, Larsen SB, Gomez NC, Alaverdyan K, Sendoel A, Yuan S, Polak L, Kulukian A, Chai S, Fuchs E. 2017. Inflammatory memory sensitizes skin epithelial stem cells to tissue damage. *Nature* **550**:475–480. doi:10.1038/nature24271

Netea MG, Joosten LAB, Latz E, Mills KHG, Natoli G, Stunnenberg HG, O'Neill LAJ, Xavier RJ. 2016. Trained immunity: A program of innate immune memory in health and disease. *Science* **352**:aaf1098. doi:10.1126/science.aaf1098

Oben JA, Mouralidarane A, Samuelsson AM, Matthews PJ, Morgan ML, Mckee C, Soeda J, Fernandez-Twinn DS, Martin-Gronert MS, Ozanne SE, Sigala B, Novelli M, Poston L, Taylor PD. 2010. Maternal obesity during pregnancy and lactation programs the development of offspring non-alcoholic fatty liver disease in mice. *J Hepatol* **52**:913–920. doi:10.1016/j.jhep.2009.12.042

Panserat S, Marandel L, Seiliez I, Skiba-Cassy S. 2019. New Insights on Intermediary Metabolism for a Better Understanding of Nutrition in Teleosts. *Annu Rev Anim Biosci* **7**:195–220. doi:10.1146/annurev-animal-020518

Pfennig DW, Wund MA, Snell-rood EC, Cruickshank T, Schlichting CD, Moczek AP.

2010. Phenotypic plasticity's impacts on diversification and speciation. *Trends Ecol Evol* **25**:459–467. doi:10.1016/j.tree.2010.05.006

Pirola CJ, Gianotti TF, Castaño GO, Mallardi P, Martino JS, Ledesma MMGL, Flichman D, Mirshahi F, Sanyal AJ, Sookoian S. 2015. Circulating microRNA signature in non-alcoholic fatty liver disease: From serum non-coding RNAs to liver histology and disease pathogenesis. *Gut* **64**:800–812. doi:10.1136/gutjnl-2014-306996

Rocha F, Dias J, Geurden I, Dinis MT, Panserat S, Engrola S. 2016. Dietary glucose stimulus at larval stage modifies the carbohydrate metabolic pathway in gilthead seabream (Sparus aurata) juveniles: An in vivo approach using 14C-starch. *Comp Biochem Physiol A Mol Integr Physiol* **201**:189–199. doi:10.1016/j.cbpa.2016.07.016

Saeki K, Miura Y, Aki D, Kurosaki T, Yoshimura A. 2003. The B cell-specific major raft protein, Raftlin, is necessary for the integrity of lipid raft and BCR signal transduction. *EMBO Journal* **22**:3015–3026. doi:10.1093/emboj/cdg293

Sameh Boshra M, Shawky El Meteini M, El-Sayed Shafei A, Matboly M, Said Ali H. 2019. EVALUATION OF RAFTLIN MRNA IN HEPATOCELLULAR CARCINOMA. *Ain Shams Med J* **70**:349–359. doi:10.21608/asmj.2019.111197

Schmidt D, Wilson MD, Spyrou C, Brown GD, Hadfield J, Odom DT. 2009. ChIP-seq: Using high-throughput sequencing to discover protein-DNA interactions. *Methods* **48**:240–248. doi:10.1016/j.ymeth.2009.03.001

Shu X, Wang Q, Wu Q. 2022. The Eph/Ephrin System in Hepatocellular Carcinoma: Functional Roles and Potential Therapeutic Targets. *Oncologie* **24**:427–439. doi:10.32604/oncologie.2022.023248

Siersbæk M, Varticovski L, Yang S, Baek S, Nielsen R, Mandrup S, Hager GL, Chung

JH, GrØntved L. 2017. High fat diet-induced changes of mouse hepatic transcription and enhancer activity can be reversed by subsequent weight loss. *Sci Rep* **7**:40220. doi:10.1038/srep40220

Soltis AR, Kennedy NJ, Xin X, Zhou F, Ficarro SB, Yap YS, Matthews BJ, Lauffenburger DA, White FM, Marto JA, Davis RJ, Fraenkel E. 2017. Hepatic Dysfunction Caused by Consumption of a High-Fat Diet. *Cell Rep* **21**:3317–3328. doi:10.1016/j.celrep.2017.11.059

Stemmer M, Thumberger T, del Sol Keyer M, Wittbrodt J, Mateo JL. 2015. CCTop: An intuitive, flexible and reliable CRISPR/Cas9 target prediction tool. *PLoS One* **10**:e0124633. doi:10.1371/journal.pone.0124633

Suter MA, Ma J, Vuguin PM, Hartil K, Fiallo A, Harris RA, Charron MJ, Aagaard KM. 2014. In utero exposure to a maternal high-fat diet alters the epigenetic histone code in a murine model. *Am J Obstet Gynecol* **210**:463.e1–463.e11. doi:10.1016/j.ajog.2014.01.045

Takeda H, Shimada A. 2010. The art of medaka genetics and genomics: What makes them so unique? *Annu Rev Genet*. **44**:217–241. doi:10.1146/annurev-genet-051710-151001

Tanaka N, Takahashi S, Hu X, Lu Y, Fujimori N, Golla S, Fang ZZ, Aoyama T, Krausz KW, Gonzalez FJ. 2017. Growth arrest and DNA damage-inducible 45α protects against nonalcoholic steatohepatitis induced by methionine- and choline-deficient diet. *Biochim Biophys Acta Mol Basis Dis* **1863**:3170–3182. doi:10.1016/j.bbadis.2017.08.017

Thomas PD, Ebert D, Muruganujan A, Mushayahama T, Albou LP, Mi H. 2022. PANTHER: Making genome-scale phylogenetics accessible to all. *Protein Science*

31:8–22. doi:10.1002/pro.4218

Thompson MD. 2020. Developmental Programming of NAFLD by Parental Obesity. *Hepatol Commun*. **4**:1392–1403. doi:10.1002/hep4.1578

Tsai WC, Hsu S da, Hsu CS, Lai TC, Chen SJ, Shen R, Huang Y, Chen HC, Lee CH, Tsai TF, Hsu MT, Wu JC, Huang H da, Shiao MS, Hsiao M, Tsou AP. 2012. MicroRNA-122 plays a critical role in liver homeostasis and hepatocarcinogenesis. *Journal of Clinical Investigation* **122**:2884–2897. doi:10.1172/JCI63455

Vera LM, Metochis C, Taylor JF, Clarkson M, Skjærven KH, Migaud H, Tocher DR. 2017. Early nutritional programming affects liver transcriptome in diploid and triploid Atlantic salmon, Salmo salar. *BMC Genomics* **18**:886. doi:10.1186/s12864-017-4264-7

Vi LH, Bharti S, Inoue H, Vass WC, Campa F, Nie Z, De Gramont A, Ward Y, Randazzo PA. 2008. ASAP3 is a focal adhesion-associated Arf GAP that functions in cell migration and invasion. *Journal of Biological Chemistry* **283**:14915–14926. doi:10.1074/jbc.M709717200

Wang X, Zhang M, Ping F, Liu H, Sun J, Wang Y, Shen A, Ding J, Geng M. 2019. Identification and Therapeutic Intervention of Coactivated Anaplastic Lymphoma Kinase, Fibroblast Growth Factor Receptor 2, and Ephrin Type-A Receptor 5 Kinases in Hepatocellular Carcinoma. *Hepatology* **69**:573–586. doi:10.1002/hep.29792/suppinfo

Wankhade UD, Zhong Y, Kang P, Alfaro M, Chintapalli S v., Thakali KM, Shankar K. 2017. Enhanced offspring predisposition to steatohepatitis with maternal high-fat diet is associated with epigenetic and microbiome alterations. *PLoS One* **12**:e0175675. doi:10.1371/journal.pone.0175675

Watakabe I, Hashimoto H, Kimura Y, Yokoi S, Naruse K, Higashijima SI. 2018. Highly

efficient generation of knock-in transgenic medaka by CRISPR/Cas9-mediated

genome engineering. *Zoological Lett* **4**:3. doi:10.1186/s40851-017-0086-3

Wesolowski SR, Kasmi KCE, Jonscher KR, Friedman JE. 2017. Developmental origins

of NAFLD: A womb with a clue. *Nat Rev Gastroenterol Hepatol*. **14**:81-96.

doi:10.1038/nrgastro.2016.160

Yun HJ, Suh YJ, Kim Y Bin, Kang EJ, Choi JH, Choi YK, Lee IB, Choi DH, Seo YJ, Noh

JR, Choi HS, Kim YH, Lee CH. 2022. Hepatocyte DAX1 Deletion Exacerbates

Inflammatory Liver Injury by Inducing the Recruitment of CD4+ and CD8+ T Cells

through NF-κB p65 Signaling Pathway in Mice. *Int J Mol Sci* **23**:14009.

doi:10.3390/ijms232214009

Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nussbaum C, Myers

RM, Brown M, Li W, Shirley XS. 2008. Model-based analysis of ChIP-Seq (MACS).

Genome Biol **9**:R137. doi:10.1186/gb-2008-9-9-r137

Zhao J, Li R, Li J, Chen Z, Lin Z, Zhang B, Deng L, Chen G, Wang Y. 2022. CAFs-

derived SCUBE1 promotes malignancy and stemness through the Shh/Gli1 pathway

in hepatocellular carcinoma. *J Transl Med* **20**:520. doi:10.1186/s12967-022-03689-

w

2 **Supplementary figure 1: Confirmation of hepatocyte-specific expression of GFP in**

3 ***tdo2*:GFP transgenic medaka.**

4 Immunofluorescent images of a liver section stained with GFP antibody are shown. Note

5 that GFP signal was detected in hepatocytes, but not in red blood cells, endothelial cells

6 in the liver, or cells in the intestine. Scale bar: 50 µm (upper), 20 µm (lower).

2 **Supplementary figure 2: Sorting of GFP-positive hepatocytes from livers of**

3 *tdo2*:**GFP medaka.**

4 FACS-sorting strategy of GFP-positive hepatocytes from liver cell suspensions of

5 *tdo2*:GFP transgenic medaka. GFP-positive, propidium-iodide (PI)-negative, singlet cells

6 were sorted.

Supplementary figure 3: Differentially expressed genes/peaks at FDR < 0.1, corrected for multiple testing.

Volcano plots showing **(A)** differentially expressed genes by RNA-seq, **(B)** differentially accessible peaks by ATAC-seq, and **(C)** differentially H3K27ac-enriched peaks by ChIP-seq. X-axis: log2 fold change of normalized read counts within genes/peaks (upper, HFD/NC; lower, HFD-NC/NC-NC), y-axis: –log10(FDR).

Supplementary figure 4: Categorization of peaks into promoters and enhancers.

(A) ATAC-seq peaks (30,484 peaks in total) were categorized as promoters if peaks were located inside 2 kb on either side of the transcription start sites (TSS), while others were categorized as enhancers. **(B)** H3K27ac ChIP-seq peaks (35,722 peaks in total) were categorized as promoters if peaks were located inside 2 kb on either side of the TSS, while others were categorized as enhancers. Enrichment of peaks with differential signal at enhancers was tested by Fisher's exact test.

H3K27ac

H3K27me3

H3K9me3

Supplementary figure 5: Western blotting of H3K27ac, H3K27me3, and H3K9me3

histone modifications in livers.

Left: Raw images of western blotting of each histone modification and histone H3. One

lane indicates liver lysate of a single fish, concentration of which was adjusted by liver

weight. Right: plots of signal intensities of each histone modification, normalized by

signal intensities of histone H3. Note that there is little difference in the global levels of

each histone modification in medaka liver between dietary conditions.

Supplementary figure 6: Distribution pattern of histone modifications around

ATAC-seq peaks.

(A) Distribution pattern of H3K27ac (orange), H3K27me3 (blue), and H3K9me3 (green)

around all 30,484 ATAC-seq peaks, sorted by ATAC-seq signal intensity. For ATAC-seq,

signal intensities within a 1 kb window from the centers of ATAC-seq peaks are displayed.

For histone modifications, signal intensities within an 8 kb window from the centers of

ATAC-seq peaks are displayed. Note the positive correlation of ATAC-seq signal with

H3K27ac, and the negative correlation with H3K27me3 and H3K9me3 signal. **(B)**

Distribution patterns of each histone modification around differentially accessible peaks.

For peaks differentially accessible between HFD and NC fish, a moderate positive

correlation was observed for H3K27ac. However, the correlation was not observed for

differentially accessible peaks between HFD-NC and NC-NC fish. For H3K27me3 and

H3K9me3, little changes were seen for differentially accessible peaks at any time point.

Supplementary figure 7: Differentially enriched H3K27ac peaks by HFD feeding.

(A) Gene ontology analyses of genes close to H3K27ac peaks with different enrichment between HFD and NC group fish, inferred by PANTHER v17.0. **(B)** Heatmaps of H3K27ac peaks differentially enriched at seven and 15 weeks of age. Signal intensities in ±2.5 kb regions from the centers of H3K27ac peaks are displayed. **(C)** A heatmap of persistent H3K27ac ChIP-seq peaks after NC (11 peaks in total). Log2-transformed, and Z-transformed, DESeq2 normalized read counts at each peak were displayed. DESeq2 results of RNA-seq and ATAC-seq of nearby genes/peaks were displayed on the right.

Supplementary figure 8: H3K9me3 domains downregulated by HFD feeding.

(A) Gene ontology analyses of genes close to H3K9me3 bins differentially enriched between HFD and NC group fish. **(B)** Venn diagram of genes close to H3K9me3-downregulated bins and genes differentially expressed after HFD feeding. **(C)** A heatmap of H3K9me3 levels for the downregulated bins by HFD. Log2-transformed, and Z-transformed, DESeq2 normalized read counts of H3K9me3 ChIP-seq at each bin are displayed.

1 **Supplementary figure 9: Track views of persistent peaks.**

2 **(A)** Track views of the 50 persistent ATAC-seq peaks. DESeq2 results of RNA-seq and

3 H3K27ac ChIP-seq of nearby genes/peaks are displayed on the upper left. **(B)** Track

4 views of the 11 persistent H3K27ac ChIP-seq peaks. DESeq2 results of RNA-seq and

5 ATAC-seq of nearby genes/peaks are displayed on the upper left.

Supplementary Figure 10: Rescaling procedure of read counts for visualization of ChIP-seq and ATAC-seq.

The top left panels describe the regression process. Each point in these panels represents read count data from a single region. For each library, the signals on the selected regions (y-axis) were regressed on the averages of all libraries (x-axis). In this process, the regions whose average signal was within the top 100 or below the global average (average across all regions and libraries) were excluded (red dotted lines). Then, the raw counts (left bottom panel) were divided by the corresponding regression coefficients (blue dotted lines) to obtain the rescaled counts (the right bottom panel).