Climate change promoted allopatric divergence and explains the

current disjunct geographic distribution of the *Ophrys aveyronensis*

species complex (Orchidaceae)

Anaïs Gibert[1], Roselyne Buscail[2], Michel Baguette[3,4], Christelle Fraïsse[5], Camille

Roux[5], Bertrand Schatz[6] & Joris A. M. Bertrand[1]

[1] Laboratoire Génome et Développement des Plantes (LGDP), UMR 5096, Université de

Perpignan Via Domitia (UPVD) - Centre National de la Recherche Scientifique (CNRS) -

Institut de Recherche pour le Développement (IRD), EMR 269 MANGO, F-66860, Perpignan.

[2] Centre de Formation et de Recherche sur les Environnements Méditerranéens (CEFREM),

Université de Perpignan Via Domitia (UPVD) - Centre National de la Recherche Scientifique

(CNRS), F-66860, Perpignan.

[3] Institut Systématique, Évolution, Biodiversité (ISEB), UMR 7205, Museum National

d'Histoire Naturelle (MNHN) - Centre National de la Recherche Scientifique (CNRS) -

Sorbonne Université - École Pratique des Hautes Études (EPHE) - Université des Antilles, F-

75005, Paris.

[4] Station d'Écologie Théorique et Expérimentale (SETE), UMR 5321, Centre National de la

Recherche Scientifique (CNRS) - Université Toulouse III, F-09200, Moulis

[5] Evo-Eco-Paleo, UMR 8198, Centre National de la Recherche Scientifique (CNRS) - Université

de Lille, F-59000, Lille

[6] Centre d'Écologie Fonctionnelle et Évolutive (CEFE) - Université de Montpellier - Centre

National de la Recherche Scientifique (CNRS) - École Pratique des Hautes Études (EPHE) -

Institut de Recherche pour le Développement (IRD), F-34293, Montpellier.

Correspondence: anais.gibert@gmail.com; joris.bertrand@univ-perp.fr

Abstract

Species with disjunct geographic distributions provide natural opportunities to investigate

recent or incipient allopatric divergence. Although not rare, many of the cases observed

result from successful colonization from a historical to a new range through dispersal or

human-induced introduction which make the underlying eco-evolutionary processes

sometimes difficult to decipher. The *Ophrys aveyronensis* species complex presents a

disjunct geographic distribution with two ranges currently separated by 600 km on both

sides of the Pyrenees Mountains. To uncover the causes of such intriguing biogeographic

pattern, we combined population genomics and Ecological Niche Modelling approaches.

Population genomic data show that all the populations studied display similar patterns of

genetic diversity and dramatic decrease in effective size. We found significant genetic

differentiation between the two ranges of the *O. aveyronensis* species complex. Our results

indicate a recent divergence (*i.e.* ~3500 generations ago). Ecological Niche Modelling results

further support that the disjunct geographic distribution of the *O. aveyronensis* species

complex is consistent with a range split of a broad ancestral range, contraction and shifts in

opposite directions in response to climate warming since the Last Glacial Maximum (LGM,

about 21000 years ago). The congruence of both the results of population genomics and

ENM approaches indicate that allopatric speciation is involved in the *Ophrys* adaptive

radiation. The *O. aveyronensis* complex thus provides a promising candidate to further

investigate putative incipient ecological divergence following an initial phase of geographic

isolation.

Keywords Population genomics**,** RAD sequencing, Species Distribution Modelling (SDM),

Biogeography, Speciation, Orchids

Introduction

Allopatric speciation (also referred as geographic speciation, or vicariant speciation) occurs when populations of a given species become reproductively isolated from each other because of geographic separation. Aside from being conceptually intuitive, allopatric speciation has been evidenced by theoretical models, laboratory experiments and empirical studies to the point that it is generally regarded as the "default mode" of speciation (see Coyne & Orr, 2004). In allopatric speciation, spatial and reproductive isolation are often assumed to result from geologic-caused topographic changes that form the geographic barrier. In continental systems however, allopatric divergence is prone to experience only transient geographic and reproductive isolation involving one of several phases of secondary contact (e.g. through climate change-induced oscillations during the Quaternary, Hewitt, 2000). Thus, allopatry is a spatio-temporally dynamic process. Aside, not only the fore-mentioned barriers but any kind of fragmentation of species distribution range, including climate change and/or human activities may also initiate and be responsible for evolutionary divergence.

The putative complexity of the speciation process makes difficult to elucidate its ecological and evolutionary causes as current spatial patterns of distribution only provide a snapshot from the speciation continuum. In this regard, species with disjunct geographic distributions are likely to correspond to species whose populations became recently allopatric, *i.e.* with insufficient time to accumulate enough divergence to be considered as distinct species yet. Although such natural example may help to understand the consequences of physical barriers in early stages of evolutionary divergence, disjunct geographic distributions do not necessary result from the fragmentation of an ancestral

range. Indeed, dispersal and successful colonization of a new range from the historical one

may also lead to the same pattern.

Several animal species including terrestrial mollusks (e.g. Reich et al., 2015 and

references therein) and plants (e.g. Beatty & Provan, 2013; 2014; Beatty et al., 2015;

Fagúndez & Díaz-Tapia, 2023) occur in the North-East of the Iberian Peninsula and in Ireland

(a so called 'Lusitanian' distribution) and when they occur elsewhere, display noticeable

genetic similarities between these two regions (see Grindon & Davison, 2013 for another

example in terrestrial mollusks, Davison et al., 2001 and Mascheretti et al., 2003 for

examples in mammals). To explain these disjunct geographic distributions, several authors

have considered that all these species may have survived in ice-free pocket (located in

southwest Ireland) and in southern European refugia (such as the Iberian Peninsula) during

the Last Glacial Maximum (LGM, 21000 years *Before Present*) or should have colonized

Ireland thanks to land-bridges, that disappeared ever since. However, molecular data have

more recently supported that most of the Irish animal populations have been introduced in

the Mesolithic times (more than 8000 years ago) through human activities whereas plants

could have been more prone to accomplish post-glacial recolonization of this region. In such

examples, the combination of genetic data with sufficient resolution and/or Environmental

Niche Modelling (ENM) have been crucial to compare the different scenarios and better

understand the biogeography of both species and regions.

The species of the genus *Ophrys* have long been known for their unusual pollination

syndrome (called 'sexual swindling') that involves insect-like flowers mimicking their female

pollinator to lure conspecific males and ensure pollination but are now also increasingly

considered as a promising biological model to investigate adaptive radiations. The degree of

plant-pollinator specialization (Joffard et al., 2019) required to ensure the success of the

Ophrys pollination strategy was shown to be the driver of the impressive rate of

diversification of the genus within the Mediterranean basin, and to a lesser extent in other

parts of Western Europe (see Baguette et al., 2020, for a recent review). If *Ophrys* are

notorious for their relatively high number of likely cases of speciation with gene flow, the

number of island endemic taxa prove that allopatric speciation is also an important

mechanism to explain *Ophrys* species richness observable nowadays (see Bertrand et al.,

2021a).

For continental taxa, clear evidences of allopatric speciation in *Ophrys* are missing.

Nevertheless, one species, *Ophrys aveyronensis*, displays an intriguing case of disjunct

geographic distribution. This species was initially described as endemic to a very restricted

area in the Grands Causses region (Southern France). Observations of morphologically very

similar populations were reported later from Northern Spain and described as a distinct

species (*Ophrys vitorica*, Kreutz, 2007). *Ophrys aveyronensis* is now rather considered as a

complex consisting of two subspecies (*O. aveyronensis* subsp. *aveyronensis* and *Ophrys*

aveyronensis subsp. *vitorica*) that share a mutual insect pollinator (*i.e.* the solitary bee

Andrena hattorfiana, Paulus & Gack, 1999; Paulus, 2017; Benito Ayuso, 2019 (but see Gibert

et al., 2022 reporting subtle differences in morphometry and coloration). This taxon is thus

present in two rather restricted ranges separated by a gap of about 600 km (and Pyrenees

Mountains), in the Grands Causse region (France) and in Northern Spain.

The formation of the Pyrenees starting during the Eocene (from 45 Myrs ago) cannot

explain by itself the split of the ancestral range of the species as all phylogenetic hypotheses

published so far conclude that the *Ophrys* genus emerged no more than 5-6 Myrs ago

(Breitkopf et al., 2015, see also Inda et al., 2012). This does not rule out 'natural' allopatry as

factor to explain current disjunct geographic distribution but suggests that it would have

occurred more recently. According to this "vicariance hypothesis", the current disjunct

geographic distribution may correspond to two relict pockets of an ancestral range that got

fragmented following a scenario according to which populations from Northern and

Southern ranges diverged in allopatry. In this case, the splitting time between northern and

southern populations is expected to be relatively old and current levels of genetic diversity

are not necessary expected to vary among populations. Alternatively, we also envisage

recent successful colonization from one historical range to the other. This event could have

occurred 'naturally'. The notoriously small *Ophrys* seeds are likely to be wind-dispersed over

hundreds of kilometers and may have found in the newly colonized area suitable conditions

for their germination and development. In this case, it may be difficult to provide a dating of

splitting time, but the levels of genetic diversity are expected to be lower in the newly

colonized area, as the recently founded *Ophrys* population is only a subset of the genetic

pool of the origin population. The colonization of one part of the range from the other may

also have occurred 'non-naturally' (namely, because of anthropogenic activities). In all cases,

the environmental conditions are expected to be highly similar in the two regions.

In this study, we used a population genomic approach (*i.e.* through a double digest

RAD-seq like protocol called 'nGBS') to first investigate patterns of genetic diversity and

differentiation among *Ophrys aveyronensis* populations from both part of its disjunct

geographic range. We then couple demographic inference analyses and Ecological Niche

Modelling approaches to test whether the current disjunct geographic distribution of this

species complex is rather consistent with either allopatry or recent introduction.

Materials and Methods

Sampling and study sites

We used a minimally destructive sampling method to collect a total of 86 plant tissue

samples from six localities that span the entire geographic range of the *O. aveyronensis*

species complex: three in Southern France (Grands Causses region): 1-Guilhaumard, 2-

Lapanouse-de-Cernon and 3-Saint-Affrique, and three in Northern Spain: 4-Valgañón, 5-

Larraona and 6-Bercedo, in June 2019, Fig. 1, see also Gibert et al., 2022). At each sampling

site, we collected 6 to 10, 0.6 mm diameter leaf punches from 13 to 15 individuals which

were stored in 100% ethanol until DNA extraction (see Table 1 and Supplementary Table S1).

As *Ophrys aveyronensis* is a nationally protected plant species in France, sampling was done

under permit 'Arrêté préfectoral n°2019-s-16' issued by the 'Direction Régionale de

l'Environnement de l'Aménagement et du Logement (DREAL)' from the 'Région Occitanie',

on 07-May-2019). *Ophrys aveyronensis* is not legally protected in Spain.

DNA extraction, library preparation and genotyping

Genomic DNA extraction as well as library preparation and genotyping were subcontracted

to LGC Genomics GmbH (Berlin, Germany). A fractional genome sequencing strategy called

normalized Genotyping-by-Sequencing (nGBS) was used to subsample the *Ophrys*

aveyronensis genome for which we estimated a relatively high size of ~5-6 Gbp based on

flow cytometry techniques (Bertrand, unpublished data). The nGBS protocol makes use of a

combination of two restriction enzymes (PstI and ApeKI, in our case) to produce a

reproducible set of fragments across samples/individuals. It also comprises a normalization

step whose aim is to avoid further sequencing of highly repetitive regions. Illumina

technology was then used to aim at obtaining a minimum of 1.5 million reads per sample

(with 2 x 150 bp read length) for each of the 86 individually barcoded libraries.

Population genomic dataset processing

We used different scripts included in *Stacks* v.2.60 (Catchen et al., 2011; 2013) to build loci

from Illumina reads, *de novo* (*i.e.* without aligning reads to a reference genome). We first

used *process_radtags* to demultiplex and clean reads, then *denovo_map.pl* to build loci

within individuals, create a catalog and match all samples against it and finally *populations* to

further filter the SNPs obtained at the population level and compute basic population

genetic statistics. Before running the pipeline on the complete dataset, we first optimized

several key parameters: *-m* (the minimum number of identical raw reads required to form a

putative allele), *-M* (the number of mismatches allowed between alleles to form a locus) and

-n (the number of mismatches allowed between loci during construction of the catalog) by

running it on a subset of 12 individuals representative of the whole data set (*i.e.* geographic

origin and coverage). As recommended by several authors, we varied M and n from 1 to 9

(fixing $M = n$) while keeping $m = 3$ (see Paris et al., 2017). The combination *-m* 3, *-M* 5 and *-n*

5 was found to be the most suitable to maximize the number of SNPs, assembled and

polymorphic loci in our case and was used to run *Stacks* on the whole data set (see

Supplementary Table S2). Then, we ran the *populations* script included in *Stacks*, from which

only loci that were genotyped in at least 80% of the individuals per population (*-r* 80) and

found in all six sampling sites were kept (*-p* 6), while loci exhibiting estimates of observed

heterozygosity greater than 70% (*--max-obs-het* 0.7) were filtered out to reduce the risk of

including remaining paralogs. We also discarded sites whose minor allele frequency was

lower than 5% (*--min-maf* 0.05). The genotype matrix of the remaining 9301 SNPs was then

exported in several formats for downstream analyses.

Genetic diversity and population differentiation

To get an overview of the overall genetic diversity and differentiation among individuals and

populations, we first performed a Principal Component Analysis (PCA) based on a matrix of

86 individuals (as rows) and 9301 SNPs (as columns) coded as a *genlight* object with the R-

package *adegenet* (Jombart, 2008; Jombart & Ahmed, 2011). The number of private alleles

and the average nucleotide diversity (π) were directly available from the output of the

populations script from *Stacks*. We used Genodive v.3.04 (Meirmans, 2020) to compute

expected and observed heterozygosity (H_E and H_O, respectively) as well as to evaluate the

deviation from panmixia by computing G_{IS}. Allelic richness (A_R) was computed with the R-

package *hierfstat* (Weir & Goudet, 2017). Overall and pairwise genetic differentiation was

also assessed based on G-statistics (G_{ST}, G'_{ST}, G''_{ST}) as well as Jost's D also implemented in

Genodive. An Analysis of Molecular Variance (AMOVA) allowed to assess the proportion of

genetic variance at different hierarchical levels, especially between sampling sites among

subspecies/country (F_{SC}) and between subspecies/country (F_{CT}). The statistical significance of

the obtained values was estimated based on 10000 permutations.

To further investigate population structure and characterize putative

migration/admixture event, we used sNMF (Frichot et al., 2014) as implemented in the R-

package *LEA* (Frichot & François, 2015) to estimate individual ancestry coefficients based on

sparse non-negative matrix factorization algorithms. The number of ancestral populations

(genetic clusters) was varied from $K = 1$ to 10, and analyses were run with 10 replicates at

each value of K. We followed the cross-entropy criterion to determine the optimal number

of clusters and the ancestry proportions matrices (Q-matrices) obtained were plotted with

the R-package *pophelper* (Francis, 2017).

Demographic and evolutionary history reconstruction

We reconstructed the past demographic history of each of the six populations, separately,

using STAIRWAYPLOT2 (Liu & Fu, 2015; 2020). This program relies on the Site Frequency

Spectrum (SFS) to infer the temporal dynamics of effective population size (N_e) and has the

advantage of not requiring whole-genome sequence data or reference genome information.

First, we used the *easySFS.py* script (https://github.com/isacovercast/easySFS) to compute

the folded SFS of each population. To maximize the number of segregating sites, the

population was downsampled to 24, 24, 22, 24, 24 and 22 individuals for 1-Guilhaumard, 2-

Lapanouse, 3-St-Affrique, 4-Valgañón, 5-Larraona and 6-Bercedo haploid genomes,

respectively. We then ran STAIRWAYPLOT2 that fits a flexible multi-epoch demographic

model that estimates N_e at different time periods using the SFS to infer estimated N_e

fluctuation coinciding with coalescent events. We assumed a generation time of 5 years for

Ophrys and a mutation rate of 7 x 10^{-9} per site per generation for angiosperms autosomal

markers (following Krasovec et al., 2018) and performed 200 bootstrap replicates to

estimate 95% confidence intervals.

We then used the Approximate Bayesian Computation (ABC) framework

implemented in DILS (Fraïsse et al., 2021) to infer the best demo-genomic scenario and

associated parameters for each pair of *O. a.* subsp. *aveyronensis* and *O. a.* subsp. *vitorica*

populations. The two-population model of DILS compares different scenarios in a

hierarchical manner. First, current isolation models, *i.e.* Strict Isolation (SI) versus Ancient

Migration (AM) are compared to ongoing migration models, *i.e.* Isolation Migration (IM)

versus Secondary Contact (SC). Then, for the best demographic model, DILS compares

genomic models to test for background selection by assuming variation in effective

population size (N_e) across sites (*i.e.* hetero versus homo Ne models are compared). For

migration models (IM and SC), DILS also tests for selection against migrants by assuming

variation in migration rates $N.m$ across sites (*i.e.* hetero versus homo $N.m$ models are

compared). Depending on the best selected model, DILS provides estimates and confident

intervals for parameters such as the time of split (T_{split}), the migration rate ($N.m$), current and

ancestral effective population sizes ($Ne_{current}$ and Ne_{past}) and whenever relevant, the time of

ancient migration (T_{AM}) or the time of secondary contact (T_{SC}). We formatted the input file

with a custom Perl script and used the online facility (http://dils.univ-lyon1.fr/) to run

computations with the following settings and priors: *population_growth*: variable,

modeBarrier: bimodal (*i.e.* there is a class of species barrier loci with $N.m=0$ and a class of

loci migrating at the background rate $N.m$) for model type to test), *max_N_tolerated*: 0.25,

LMin: 10, *nMin*: 8, *mu*= 7.10^{-9}, *rho_over_theta*: 0.2, *N_min* = 0, *N_max*: 200000, *Tsplit_min*:

0, *Tsplit_max*: 100000, *M_min*: 1 and *M_max*: 40. As the first step of DILS comparing the

models 'with' *versus* 'without' current migration did not result in strong statistical support

for either model, we estimated the parameters of the strict isolation model based on our

knowledge of current biogeography. This parameter estimation was carried out using the

same prior distribution as described above. Finally, to quantify the fit of the estimated model

to the data we conducted a goodness-of-fit test using 2000 simulations performed under the

SI model based on the estimated parameter values.

Past and current Species Distribution Modelling

We used an Ecological Niche Modelling (or Environmental Niche Modelling, ENM or Species

Distribution Modelling, SDM) framework similar to the one we followed and described in

detail in Salvado et al., (2022) to infer the spatio-temporal dynamics of suitable habitat for

the *O. aveyronensis* species complex since the Last Glacial Maximum (LGM, 21 kyrs BP). We

then compared the results with those obtained from population genomic analyses. To model

and spatially project the current bioclimatic niche, we first downloaded 19 bioclimatic layers

at a resolution of 2.5 arc-minutes (~5 km^2) available from WorldClim 2 (Fick & Hijmans,

2017). Current climate data correspond to time averaged variables over the period 1970-

2000. To spatially project the inferred niche onto past bioclimatic conditions data, we then

downloaded paleoclimate data either from WorldClim v1.4 (Hijmans et al., 2005) and from

PaleoClim (www.paleoclim.org).

Paleoclimate data from WorldClim are available for both the LGM (21000 kyrs ago)

and the Mid-Holocene (about 6000 years ago), downscaled at a resolution of 2.5 arc-minutes

for at least three General Circulation Models (GCMs) proposed by the National Center for

Atmospheric Research (NCAR) though the Community Climate System Model (CCSM4), the

Japan Agency for Marine-Earth Science and Technology, the Center for Climate System

Research of the University of Tokyo, and the National Institute for Environmental Studies

(MIROC-ESM) and the Max Plant Institute for Meteorology Earth System Model (MPI-ESM-

P). Only CCSM data are available from PaleoClim but over different time periods over the

Pleistocene (at a resolution of 2.5 arc-minutes): the LGM (as from CHELSA, Karger et al.,

2021), the Heinrich Stadial1 (17.0-14.7 kyrs BP), the Bølling-Allerød (14.7-12.9 kyrs BP) the

Younger Dryas Stadial (12.9-11.7 kyrs BP) and the Holocene: early-Holocene, Greenlandian

(11.7-8.326 kyrs BP), mid-Holocene, Northgrippian (8.326-4.2 ka BP) and late-Holocene,

Meghalayan (4.2-0.3 kyrs BP) (as from Fordham et al., 2017).

The *O. aveyronensis* occurrences dataset consists of a set of field observations as well

as from records downloaded from the gbif (www.gbif.org) and the iNaturalist

(www.inaturalist.org) databases before manual curation. The Ecological Niche Modelling

analyses were conducted thanks to the R-package *ENMwizard v.0.3.7* (Heming et al., 2018,

see also Bagley et al. 2020) which is a convenient wrapper of several tools. We first spatially

filtered occurrences to keep only those that were at least 5 km away from each other using

the R-package *spThin* (Aiello-Lammens et al., 2015). The calibration area for the models was

created as a buffer of 0.5° around the minimum convex polygon encompassing all

occurrences. From the 19 bioclimatic variables, we selected the less correlated ones (Person

correlation coefficient < 0.75) thanks to the R-package *caret* (Kuhn, 2019) and kept 7

variables: bio2 (the mean diurnal range which is the mean of monthly maximum

temperature minus the mean of monthly minimum temperature), bio3 (the isothermality,

i.e. bio2 divided by the temperature annual range), bio8 (the mean temperature of the

wettest quarter), bio9 (the mean temperature of the driest quarter), bio10 (the mean

temperature of the warmest quarter), bio15 (the precipitation seasonality, *i.e.* its coefficient

of variation) and bio17 (the precipitation amount of the driest quarter) for further analyses.

We used the maximum entropy method (implemented in MaxEnt ver. 3.4.1, Phillips

et al., 2006; Philips & Dudík, 2008; Phillips et al., 2017) to calibrate models and evaluated

models' performance thanks to the package *ENMeval* (Muscarella et al., 2014) as

implemented in *ENMwizard*. We evaluated models using a geographic partition scheme of

type "block" and optimized two parameters of MaxEnt: the Regularization Multipliers (RM)

and the Feature Classes (FCs). RM was varied from 0.5 to 4.5, incremented by 0.5 whereas a

suite of 15 FCs (L, for Linear, P, for Product, Q, for Quadratic and H for Hinge) or combination

of them were evaluated: L, P, Q, H, LP, LQ, LH, PQ, PH, QH, LPQ, LPH, LQH, PQH, LPQH,

resulting in a total of 135 models. Model selection was done by computing the corrected

Akaike Information Criterion ("LowAIC"). Model accuracy was also evaluated by calculating

omission rates calculated when a 10[th] percentile threshold is applied (x10ptp). The final

model was projected on current and paleoclimatic contexts.

Results

Population genomic dataset

We obtained a total of 150 694 350 of read pairs across the 86 individuals (1 382 768 - 7 370 366 of raw reads per ind., mean 3 504 519, SD = 1 282011.52) of which 290 431 210 barcoded reads (*i.e.* 96.36%) were retained, 10 819 014 (3.6%) were removed because RAD cutsite was not found in the sequence and 138 476 reads (< 1%) were removed because of low quality. We genotyped a total of 859 035 *loci* (composed of 135 255 870 sites) including 368 712 variant sites (SNPs) with *Stacks*. Average read depth ranged from 24.1X to 81.6X (mean 48.7X, SD = 12.4X) based on the combination of parameters we used. After filtering the data with *populations*, we finally kept 4302 *loci* from which we retained 9 301 variant sites (SNPs).

Genetic diversity and population differentiation

The PCA biplot shows that individuals are arranged by subspecies/country of origin along PC1 (which explains 8.50% of the total genetic variance) and by sampling site along PC2 (which explains 3.88% of the total genetic variance) even though the sampling sites of 1-Guilhaumard and 2-Lapanouse-de-Cernon slightly overlap (Fig. 2A). Based on PC2, intra-population genetic variation seems minimal for 1-Guilhaumard and 2-Lapanouse-de-Cernon and maximal for 3-Saint-Affrique and 5-Larraona. Basic population genetics statistics are indicated in Table 1. All populations displayed weak but significant deviation from panmixia with a mean G_{IS} = 0.060 (p < 0.01, 95% CI: 0.056-0.064) and population G_{IS} values ranging from 0.006 in 3-Saint-Affrique to 0.081 in 4-Valgañón (all p < 0.01). Average nucleotide diversity (π) varies from 0.269 in 3-Saint-Affrique to 0.290 in 4-Valgañón and 5-Larraona, and allelic richness (A_R) from 16696.70 (in 3-Saint-Affrique) to 17297.27 (in 5-Larraona). Based on

the filtering procedure we followed, the number of private alleles was relatively low but

consistently higher for *O. a.* subsp. *vitorica* (*i.e.* from 9 to 23) compared to *O. a.* subsp.

aveyronensis (*i.e.* from 0 to 2).

Overall *G*-statistics show that the whole set of populations is genetically structured:

G_{ST} = 0.091 (*p* < 0.01, 95% CI: 0.089-0.093); G'_{ST} = 0.134 (*p* < 0.01, 95% CI: 0.105-0.011); G''_{ST}

= 0.150 (*p* < 0.01, 95% CI: 0.146-0.153) and Jost's *D* = 0.047 (*p* < 0.01, 95% CI: 0.046-0.048).

The AMOVA confirms that although a major part of the genetic variance is found within

individual (81.8%, F_{IT} = 0.182, 95% CI: 0.177-0.186), significant proportions of the total

genetic variance are found among populations within subspecies/country (6.8%, F_{SC} = 0.072,

95% CI: 0.070-0.074) and between subspecies/country (6.1%, F_{CT} = 0.061, F_{CT} = 0.058-0.064).

Pairwise differentiation between sampling sites varies from F_{ST} = 0.049 to 0.082 within

subspecies/country and reaches F_{ST} = 0.111 to 0.147 between subspecies/country of origin

(all *p* < 0.01; see Supplementary Appendix S3 for details).

The clustering analysis carried out with sNMF suggests an optimal number of genetic

clusters of *K*=5 based on the cross-entropy criterion (Fig. 2B). We plotted the *Q*-matrices

(that contain individual admixture coefficients) corresponding to the best replicate runs for

K=2 (to investigate whether the program could distinguish the two subspecies) and *K*=5 (Fig.

2C). For *K*=2, the individuals of *O. a.* subsp. *aveyronensis* and *O. a.* subsp. *vitorica* were

unambiguously assigned to two distinct clusters. For the most likely combination of *K*=5, we

observe one cluster per sampling site except for the geographically close localities of 1-

Guilhaumard and 2-Lapanouse-de-Cernon that were grouped together in a unique cluster. In

the locality of 3-Saint-Affrique 4 out of 13 individuals (19-Oav-039, 19-Oav-040, 19-Oav-049

and 19-Oav-051) display >50% of ancestry proportion that are associated to the cluster

associated with 1-Guilhaumard and 2-Lapanouse-de-Cernon.

Inference of past demographic history

Reconstruction of historical dynamics of effective population size using STAIRWAYPLOT2

shows congruent trends consisting of progressive >95% decline in estimated N_e for all 6

populations, since the LGM, or at least, over the last 25000 years (see Fig. 3). Current

effective population sizes were estimated to vary between 302 (2-Lapanouse-de-Cernon)

and 642 individuals (3-St-Affrique) with this method. Despite the current geographical

discontinuity between *O. a.* subsp. *aveyronensis* and *O. a.* subsp. *vitorica* populations, we

attempted to statistically compare speciation patterns with and without ongoing migration

using DILS. For each of the 9 trans-Pyrenean pairs tested, the Posterior Probabilities

associated with the models with ongoing migration vary between 0.45 and 0.69, which

represent cases of inferential ambiguities that do not allow strong conclusions to be drawn

from model comparisons with DILS. This can happen when the real history is an intermediate

model, or when the separation time between *O. a.* subsp. *aveyronensis* and *O. a.* subsp.

vitorica is recent compared to the current population sizes. Thus, based on biogeographical

knowledge of the species, we estimate demographic parameters assuming a strict isolation

model after the subdivision of the common ancestor into two daughter populations. The 9

parameter estimates made under the strict isolation model indicate similar scenarios: the

subdivision of an ancestral population of about 120 000 individuals (from 103916 to 129860

among the 9 analyses) into two daughter populations about 3060 generations ago (from

2050 to 6884 generations among the 9 analyses, which convert into 6150 to 20652 years

assuming a generation time of 3 years, 10250 to 33420 years assuming a generation time of

5 years and 14350 to 48188 years assuming a generation time of 7 years). This subdivision is

indeed considered recent as it is about six times smaller than current population sizes. This

suggests that most coalescence events are taking place more in the ancestral population and

not in the respective populations. Similar to what was inferred with STAIWAYPLOT2, our

analyses also point to a strong demographic decline. At the time of subdivision, we infer

population sizes for the *O. a.* subsp. *aveyronensis* lineages of about 13% of ancestral size,

and 15% of ancestral size for the *O. a.* subsp. *vitorica* lineages. On average, the sizes of the

current *O. a.* subsp. *aveyronensis* populations correspond to 15% of the ancestral

population, while the current *O. a.* subsp. *vitorica* sizes correspond to 28% of the ancestral

size. The goodness-of-fit test indicates that this estimated scenario reproduces the observed

data set with high fidelity (see Supplementary Appendix S4).

Current and past Species Distribution Modelling

ENMeval analyses identified RM = 1 and a mixture of Product (P), Quadratic (Q) and Hinge

(H) feature classes as the best-performing parameters for calibrating the final ENM. These

parameters yielded a single "best" candidate model with the lowest AICc score (925.02) and

an AICc weight of ~0.23. This model had mean omission rates of 0.165 (for the 10[th]

percentile) and a mean test AUC of 0.834 (see Supplementary Table S5, for details). With a

contribution of 41.65%, bio15 (the precipitation seasonality, *i.e.*, its coefficient of variation)

was found to be the bioclimatic variable the more prone to explain the *O. aveyronensis*

species complex distribution. It was followed by bio17 (the precipitation amount of the

driest quarter) with a contribution of 23.34% and bio10 (the mean temperature of the

warmest quarter), with a contribution of 11.31%, bio2 (the mean diurnal range) with 8.14%,

bio3 (the isothermality), with 5.32% and to a lesser extent bio8 (the mean temperature of

the wettest quarter) and bio9 (the mean temperature of the driest quarter) with 3.21% and

3.01%, respectively.

The spatial projection of the current bioclimatic inferred for the *O. aveyronensis*

species complex corresponds to its actual geographic distribution but show that some

additional areas may be climatically suitable for this orchid (Fig. 5). Spatial projections

considering paleoclimatic conditions show that the extent of suitable habitat for the *O.*

aveyronensis species complex may have reached its apex during the Younger Dryas cold

period (12.9-11.7 kyrs BP) before starting to decrease across the Holocene. The LGM

conditions (21 kyrs BP) were not suitable for the establishment of viable populations in the

studied area but important suitable habitats may have existed during the Heinrich Stadial 1

event (17-14.7 kyrs BP) and the Bøllering-Allerød warming (14.7 – 12.9 kyrs BP). Altogether,

these results may be consistent with the existence of a unique ancestral area, centered on

South-Western France that could have experienced a split and a combination of reduction

and shift in two distinct directions during the Holocene. The results available from

WorldClim data do not contradict these conclusions but show significant differences across

GCMs and even for a given GCM but downscaled based on different interpolation

algorithms: namely, WorldClim instead of CHELSA (see Supplementary Appendix S5).

Discussion

Ophrys orchids exhibit among the higher rate of speciation reported so far (Breitkopf et al.,

2015) and are being increasingly mentioned as biological model of interest to investigate

ecological speciation and adaptive radiations (e.g. Nosil et al., 2017; Baguette et al., 2020;

Nürk et al., 2020). The selection pressures exerted by pollinators and the high degree of

specificity necessary to ensure the success of the sexual swindling strategy certainly explains

the overwhelming importance of ecological and sometimes sympatric speciation in *Ophrys*

(see Baguette et al., 2020). This does not preclude for a significant role of allopatric

speciation in *Ophrys*, even in non-insular systems (see Bertrand et al., 2021a). Our study, is

to our knowledge, the first to document a scenario of allopatric speciation in a continental

system event if divergence is probably too recent to accurately date the actual cessation of

gene flow. The *Ophrys aveyronensis* species complex therefore provide a promising

opportunity to investigate the relative contribution of biogeographic and ecological factors

in shaping evolutionary divergence in rapidly and intensely diversifying groups of organisms.

Patterns of genetic diversity and differentiation based on population genomics

Although distinguishing *Ophrys aveyronensis* subsp. *aveyronensis* and *O. a.* subsp. *vitorica* is

difficult but still overall possible based on phenotypic data (Gibert et al., 2022), the French

and Iberian counterparts of the *O. aveyronensis* complex can be unambiguously teased apart

based on population genomics data. Both the PCA, *F*-statistics (and/or their analogues) and

the clustering analysis confirm that the two subspecies form distinct genetic entities,

themselves exhibiting a genetic substructure consistent with geography (sampling sites).

Altogether, these results suggest that each sampled population of *O. aveyronensis* is more

genetically differentiated from populations from the other subspecies than from populations

of its own subspecies. This pattern *a priori* rules out a recent colonization and expansion

from one range to the other from a unique source population. Patterns of genetic diversity

also support this conclusion. We found that all the populations sampled displayed relatively

similar levels of allelic richness, heterozygosity, departure from panmixia and number of

private alleles. There is therefore no evidence for any of these six populations to appear as a

genetic subsample of the genetic diversity of a putative source population elsewhere.

Finally, the inference of the temporal dynamics of effective population sizes suggests that

each sampled population may be considered as kind of replicate of each other. They all

displayed consistent decreases in effective population size over the last tens of thousands of

years suggesting that they shared a common demographic history.

It has also been proposed that *Ophrys aveyronensis* may have a hybrid origin and

Delforge (2016) hypothesized that the Iberian *O. a.* subsp. *vitorica* may result from a hybrid

swarm (without mentioning the putative parental species). Our results do not support the

later hypothesis that consider that the two subspecies would have emerged from distinct

hybridization events. However, the demographic scenarios we tested remain relatively

simple and the Bayesian Posterior Probabilities we obtained with DILS show that the

approach implemented in this program could not strongly support the Ongoing Migration

over a Strict Isolation model. Events that are known to deceives demographic inferences

such as 'ghost introgression' from an unsampled lineage have a non-negligible probability in

Ophrys. Investigating whether the *Ophrys aveyronensis* complex has or not a hybrid origin

and at least whether the two taxa have been introgressed by other *Ophrys* taxa will thus

require the genomic analyses of additional species in a future study, including more

informative data such as transcriptomes.

On the causes of the disjunct geographic distribution of the *O. aveyronensis* complex

The Ecological Niche Modelling results suggest that the LGM conditions were not suitable for

the establishment of viable populations of the *O. aveyronensis* species complex. This

suggests that either this species complex diversified from an ancestral *Ophrys* lineage in the

study area after this period, or that the populations of the complex occupied a relatively

remote glacial refugia during this period. Over the last 21000 years, the extent of the habitat

suitable for *O. aveyronensis* was rather large until the Holocene (about 12000 years ago)

(Fig. 5) where its area started to dramatically decrease while keeping an area of relatively

high suitability centered on the Southwest of France. The split and, to a lesser extent, the

shift Northward and Southward may have occurred during the Late Holocene (starting about

4000 years ago) (Fig. 5). This timing is on the same order of magnitude than the time of split

we inferred based on population genomic data and is consistent with a scenario of allopatric

divergence (without being able to fully exclude transient gene flow through climate-induced

oscillations) from an ancestral population with relatively high effective size.

Evidence for ecological speciation in addition to vicariance?

Selection pressures exerted by insect pollinators are likely to be the main driver of ecological

and putatively sympatric speciation in *Ophrys* orchids (see Baguette et al., 2020). In this way,

populations of a given *Ophrys* species being pollinated by distinct insect pollinator species

may be considered as engaged in a speciation process. So far, field observations support that

O. aveyronensis from the Grands Causses and their Iberian counterparts are pollinated by

the same pollinator species: the solitary bee *Andrena hattorfiana* (see Benito Ayuso, 2019).

In a recent study on this system, however, we showed that *O. aveyronensis* from the Grands

Causses and from the Iberian Peninsula present subtle differences at a set of morphological

and coloration traits that may be involved in pollination (Gibert et al., 2022). Although

additional work is required to address whether this variation may be adaptive and involved

in evolutionary divergence, these results call for an assessment of i) which phenotypic traits

may be under divergent selection across the *O. aveyronensis* species complex geographic

distribution and ii) an investigation of putative genomic regions associated with phenotypic

trait values, genetic structure or ecological features, something that remains largely

underexplored in *Ophrys* (but see Sedeek et al., 2014, for a noticeable exception).

507 **Taxonomic and conservation implications**

508 Considering that populations of the *O. aveyronensis* species complex are pollinated by the

509 same insect pollinator species across their range and have similar environmental niches, we

510 have currently no evidence to even consider the existence of different ecotypes in this

511 species complex. However, the disjunct geographic distribution in two subranges, separated

512 by ~600 km and the Pyrenees Mountains suggest that populations from the Grands Causses

513 and from the Iberian Peninsula may be considered as distinct evolutionary (and

514 conservation) units. Our population genomic analyses indicate that both sets of populations

515 are genetically differentiated and do not show any evidence of ongoing gene flow.

516 Accordingly, the proposition to consider them as two subspecies: *Ophrys aveyronensis*

517 subsp. *aveyronensis* and *Ophrys aveyronensis* subsp. *vitorica*, (Paulus, 2017) seems

518 appropriate.

519 Recent studies elaborated from some empirical evidence that delineating the so-

520 called 'micro-species' in *Ophrys* is practically cumbersome (Bateman et al., 2021; Bateman

521 and Rudall, 2023), especially in the section *Sphegodes* to which the *O. aveyronensis* species

522 complex belongs. Some genomic data such as whole plastomes that are useful to delineate

523 the main *Ophrys lineages*, present sequence similarity degree of more than 99.5% between

524 'microspecies' and therefore do not present enough variation to allow to discriminate them

525 (Bertrand et al., 2019; Bertrand et al., 2021b; Bateman et al., 2021). Here, however, our

526 results demonstrate that an approach consisting of sampling populations (*i.e.* a sufficient

527 number of individuals at a given location) and appropriate genetic markers (e.g. SNPs

528 genotyped on populations based on RAD-seq protocols) give signals that are consistent with

529 taxonomy. We therefore claim that such approaches of population genomics within an

530 integrative taxonomy framework (relying on flower morphology, pollinator identity,

ecological features...) is by far the most fruitful to better understand the process of

evolutionary divergence and the most relevant to improve at the same time our knowledge

on taxonomy, systematics, ecology and evolution of *Ophrys* orchids.

Author contributions

JB, BS, AG and MB designed the study. AG, JB, BS and RB gathered the data.

AG and JB performed the analyses with contribution from CF and CR and wrote the first

version of the manuscript. All authors contributed substantially to the revisions.

Data availability statement

Sequencing data have been submitted to the European Nucleotide Archive (ENA;

https:/www.ebi.ac.uk/ena/browser/home) under Study with primary accession

n°PRJEB61037 (and secondary accession number ERP146119) and samples accession

n°ERS14864169 (SAMEA112857509) to n°ERS14864254 (SAMEA112857594).

Acknowledgements

This work was primarily supported by an inter-LabEX TULIP-CEMEB initiative (n°197310) to

Joris Bertrand and Bertrand Schatz. This research was also funded by an Agence Nationale

pour la Recherche Jeune Chercheur Jeune Chercheuse (ANR JCJC) grant to Joris Bertrand,

grant number ANR-21-CE02-0022-01, and is set within the framework of the "Laboratoires

d'Excellences (LABEX)" TULIP [ANR-10- LABX-41]. Lastly, this research was also funded by the

Observatoire de REcherche Méditerranéen de l'Environnement (SO Ocove & SO PolliMed,

OSU OREME) to Bertrand Schatz. We thank Jean-Luc Roux and Daniel Vizcaïno for the

information provided about sampling sites in Spain. We thank Christian Moliné for his

contribution to fieldwork.

Conflict of interest

The authors declare no conflict of interest

References

Aiello-Lammens, M.E., Boria, R.A., Radosavljevic, A., Vilela, B., & Anderson, R.P. (2015)

spThin: an R package for spatial thinning of species occurrence records for use in

ecological   niche models. *Ecography*, 38, 541-545. doi: doi.org/10.1111/ecog.01132

Bagley, J.C., Heming, N.M., Gutiérrez, E. E., Devisetty, U.K., Mock, K. E., Eckert, A.J., &

Strauss,   S.H. (2020) Genotyping-by-sequencing and ecological niche modeling

illuminate   phylogeography, admixture, and Pleistocene range dynamics in quaking aspen

(*Populus tremuloides*) *Ecology and Evolution*, 10, 4609-4629. doi: 10.1002/ece3.6214

Baguette, M., Bertrand, J.A.M, Stevens, V.M., & Schatz, B. (2020) Why are there so many

bee- orchid species? Adaptive radiation by intra-specific competition for mnesci

pollinators.   *Biol Rev.*, 95, 1630-1663. doi: 10.1111/brv.12633

Bateman, R.M., Rudall, P. J., Murphy, A.R.M., Cowan, R.S., Devey, D.S. & Peréz-Escobar, O.A.

(2021) Whole plastomes are not enough: phylogenomic and morphometric

exploration at multiple demographic levels of the bee orchid clade *Ophrys* sect.

Sphegodes. Journal of Experimental Botany, 72, 654-681. doi: 10.1093/jxb/eraa467

Bateman, R.M. & Rudall, P.J. (2023) Morphological continua make poor species: genus-wide

morphometric survey of the European Bee Orchids (*Ophrys* L.). *Biology*, 12, 136. doi:

10.3390/ biology12010136

Beatty, G.E., & Provan, J. (2013) Post-glacial dispersal, rather than *in situ* glacial survival, best

explains the disjunct distribution of the Lusitanian plant species *Daboecia cantabrica*

(Ericacea). *Journal of Biogeography*, 40, 335-344. doi: 10.1111/j.1365-

2699.2012.02789.x

Beatty, G.E., & Provan, J. (2014) Phylogeographical analysis of two cold-tolerant plants with

disjunct Lusitanian distributions does not support *in situ* survival during the last

glaciation. *Journal of Biogeography*, 41, 2185-2193. doi: 10.1111/jbi.12371

Beatty, G.E., Lennon, J.J., O'Sullivan, C.J., & Provan, J. (2015) The not-so-Irish spurge:

Euphorbia hyberna (Euphorbiaceae) and the Littletonian plant 'steeplechase'.

Biological Journal of the Linnean Society, 114, 249-259. doi: 10.1111/bij.12435

Benito Ayuso, J. (2019) Estudios sobre polinización en el généro *Ophrys* (Orchidaceae), I,

Flora   Montiberica, 74, 32-37.

Bertrand, J. A. M., Gibert, A., Llauro, C., & Panaud, O. (2019) Characterization of the

complete plastome of *Ophrys aveyronensis*, a Euro-Mediterranan orchid with an

intriguing geographic distribution. *Mitochondrial DNA Part B*, 4, 3256-3257. doi:

10.1080/23802359.2019.1670748

Bertrand, J. A. M., Gibert, A., Llauro, C., & Panaud, O. (2021) Whole plastid genome-based

phylogenomics supports an inner placement of the *O. insectifera* group rather than a

basal position in the rapidly diversifying *Ophrys* genus (Orchidaceae). *Biology Letters*,

168, 452-457. doi: 10.1080/23818107.2021.1893216

Bertrand, J. A. M., Baguette, M., Joffard, N., & Schatz, B. (2021) Challenges inherent in the

systematics and taxonomy of genera that have recently experienced explosive

radiation: the case of orchids of the genus *Ophrys*. In Grancolas, P. & Maurel, M.-C.

Systematics and the Exploration of Life (pp 113-134). Hoboken, New Jersey: Wiley.

Breitkopf, H., Onstein, R.E., Cafasso, D., Schlüter, P.M., & Cozzolino, S. (2015) Multiple shifts

to different pollinators fuelled rapid diversification in sexually deceptive *Ophrys*

orchids. *New Phytologist*, 207, 377-389. doi: 10.1111/nph.13219

Catchen, J., Amores, A., Cresko, W., & Postlewait, J. (2011) Stacks: building and genotyping

loci de novo from short-read sequences. *G3: genes, Genomes, Genetics*, 1, 171-182.

doi: 10.1534/g3.111.000240

Catchen, J., Hohenlohe, P.A., Bassham, S., Amores, A., & Cresko, W.A. (2013) Stacks: an

analysis tool set for population genomics. *Molecular Ecology*, 22, 3124-3140. doi:

/10.1111/mec.12354

Coyne, J.A., & Orr, H.A. (2004). *Speciation*. Sunderland, Mass: Sinauer Associates.

Claessens, J., & Kleynen, J. (2016) *Orchidées d'Europe : fleurs et pollinisation*. Biotope, Mèze,

France : Biotope.

Davison, A., Birks, J.D.S., Brookes, R.C., Messenger, J.E., & Griffiths, H.I. (2001) Mitochondrial

phylogeography and population history of pine martens *Martes martes* compared

with   polecats *Mustela putorius*. *Molecular Ecology*, 10, 2479-2488. doi: 10.1046/j.1365-

294x.2001.01381.x.

Delforge, P. (2016) *Orchidées d'Europe, d'Afrique du Nord et du Proche Orient*. Lausanne,

Switzerland: Delachaux & Niestlé.

Fagúndez, J. & Díaz-Tapia, P. (2023) Comparative phylogeography of a restricted and a

widespread heather: genetic evidence of multiple independent introductions of Erica

mackayana into Ireland from northern Spain. *Botanical Journal of the Linnean Society*.

doi: 10.1093/botlinnean/boac071

Fick, S.E., & Hijmans, R.J. (2017) WorldClim 2: new 1km spatial resolution climate surfaces

for global land areas. *International Journal of Climatology*, 37, 4302-4315. doi:

10.1002/joc.5086

Fordham, D.A., Saltré, F., Haythorne, S., Wigley, T. M. L., Otto-Bliesner, B. L., Chan, K. C., &

Brook, B.W. (2017) PaleoView: a tool for generating continuous climate projections

spanning the last 21 000 years at regional and global scales. *Ecography*, 40, 1348-

1358. Doi: doi.org/10.1111/ecog.03031

Fraïsse, C., Popovic, I., Mazoyer, C., Spataro, B., Delmotte, S., Romiguier, J., Loire E., Simon,

A., Galtier, N., Duret, L., Bierne, N., Vekemans, X., & Roux C. (2021) DILS :

Demographic inferences with linked selection using ABC. *Molecular Ecology*

Resources, 2629-2644. doi: doi.org/10.1111/1755-0998.13323

Francis, R.M. (2017) POPHELPER: an R package and web app to analyse and visualize

population structure. *Molecular Ecology Resources*, 17, 27-32. doi: 10.1111/1755-

0998.12509

Frichot, E., Mathieu, F., Trouillon T., Bouchard, G., & François O. (2014) Fast and efficient

estimation of individual ancestry coefficients. *Genetics*, 196, 973-983. doi:

10.1534/genetics.113.160572

Frichot, E. & François O. (2015) LEA: An R package for landscape and ecological association

studies. *Methods in Ecology and Evolution*, 6, 925-929. doi: 10.1111/2041-

210X.12382

Gibert, A., Louty, F., Buscail, R., Baguette, M., Schatz, B. &, Bertrand, J.A.M. (2022) Extracting

quantitative information from images taken in the wild: a case study of two vicariants

of the *Ophrys aveyronensis* species complex. *Diversity*, **14**, 400. doi:

10.3390/d14050400

Grindon, A. J., & Davison, A. (2013) Irish *Cepea nemoralis* land snails have a cryptic Franco-

Iberian origin that is most easily explained by the movements of Mesolithic humans.

PLoS ONE, 8, e65792. doi: doi.org/10.1371/journal.pone.0065792

Heming, N. M., Dambros, C., & Gutiérrez, E. E. (2018). ENMwizard: AIC model averaging and

other advanced techniques in Ecological Niche Modeling made easy. R package

version   0.1.7. Retrieved from https:// github.com/HemingNM/ENMwizard

Hewitt, G. (2000) The genetic legacy of the Quaternary ice ages. *Nature*, 405, 907-913. doi:

10.1038/35016000/

Hijmans, R. J., Cameron, S.E., Parra, J. -L., Jones, P. J., & Jarvis A. (2005) Very high resolution

interpolated climate surfaces for global land areas. *International Journal of*

Climatology, 25, 1965-1978. doi: 10.1002/joc.1276

Inda, L.A., Pimentel, M., & Chase, M.W. (2012) Phylogenetics of tribe Orchideae

(Orchidaceae: Orchidoideae) based on combined DNA matrices: inferences regarding timing

of   diversification and evolution of pollination syndromes. *Annals of Botany.*, 110, 71-90.

doi: 10.1093/aob/mcs083

Joffard, N., Massol, F., Grenié, M., Montgellard, C. & Schatz, B. (2019) Effect of pollination

strategy, phylogeny and distribution niches of Euro-Mediterranean orchids. *Journal of*

Ecology, 107, 478-490. doi: 10.1111/1365-2745.13013

Jombart, T. (2008) adegenet: a R package for the multivariate analysis of genetic markers.

Bioinformatics, 24, 1403-1405. doi: 10.1093/bioinformatics/btn129

Jombart, T., & Ahmed I. (2011) adegenet 1.3-1: new tools for the analysis of genome-

wideSNP   data. *Bioinformatics*. doi: 10.1093/bioinformatics/btr521

Karger, D. N., Nobis, M. P., Normand, S., Graham, C. H., & Zimmermann, N. E. (2021):

CHELSA-   TraCE21k v1. 0. Downscaled transient temperature and precipitation data

since the   last glacial maximum. *Climate of the Past Discussions*, 1-27. doi: 10.5194/cp-

2021-30

Krasovec, M., Chester, M., Ridout, K., & Filatov, D.A. (2018) The mutation rate and the age of

the sex chromosomes in *Silene latifolia*. *Current Biology*, 28, 1832-1838. doi:

10.1016/j.cub.2018.04.069

Kreutz, C.A. (2007) Beitrag zur Taxonomie und Nomenklatur europäischer,

mediterraner, nordafrikanischer und vorderasiatischer Orchideen. *Ber. Arbeitskrs.*

Heim. Orchid. 24, 77-141.

Kuhn, M. (2019) Caret: classification and regression training. https://CRAN. R-

project.org/package=caret.

Liu, X., & Fu, Y.-X. (2015) Exploring population changes using SNP frequency spectra. *Nature*

Genetics, 47, 555-559. doi: 10.1038/ng.3254.

Liu, X., & Fu, Y.-X. (2020) Stairway Plot 2: demographic inference with folded SNP frequency

spectra. *Genome Biology*, 21, 280. doi: 10.1186/s13059-020-02196-9

Mascheretti, S., Rogatcheva M. B., Gündüz, I., Fredga, K., & Searle, J.B. (2003) How did

pygmy shrews colonize Ireland? Clues from a phylogenetic analysis of mitochondrial

cytochrome *b* sequences. *Proceedings of the Royal Society B*, 270, 1593-1599. doi:

doi.org/10.1098/rspb.2003.2406

Meirmans, P. G. (2020) GENODIVE version 3.0: Easy-to-use software for the analysis of

genetic   data of diploids and polyploids. *Molecular Ecology Resources*, 20, 1126-1131.

doi: 10.1111/1755-0998.13145

Muscarella, R., Galante, P.J., Soley-Guardia, M., Boria, R.A., Kass, J.M., Uriarte, M., &

Anderson, R.P. (2014) ENMeval: an R package for conducting spatially

independent evaluations and estimating optimal model complexity for Maxent

ecological niche models. *Methods in Ecology and Evolution*, 5, 1198-1205. doi:

10.1111/2041-210X.12261

Nosil, P., Feder, J.L., Flaxman, S.M. & Gompert Z. (2017) Tipping points in the dynamics of

speciation. *Nature Ecology and Evolution*, 1, 0001. doi: 10.1038/s41559-016-0001

Nürk, M.N., Linder, H.P., Onstein, R.E., Larcombe, M.J., Hughes, C.H., Piñeiro Fernández, L.,

Schlüter, P.M., Valente, L.M., Beierkuhnlein, C., Cutts, V., Donoghue, M.J., Edwards,

E.J., Field, R., Flantua, S.G.A., Higgins, S.II, Liede-Schumann, S. & Pirie, M.D. (2020)

Diversification in evolutionary arenas – assessment and synthesis. *Ecology and*

Evolution, **10**, 6163-6182. doi:/10.1002/ece3.6313

Paris, J., Stevens, J.R., & Catchen J.M. (2017) Lost in parameter space: a road map for

STACKS.   *Methods in Ecology and Evolution*, 8, 1360-1373. doi: 10.1111/2041-

210X.12775

Paulus, H.F. & Gack, C. (1999) Bestäubungsbiologische Untersuchungen an der Gattung

Ophrys in der Provence (SO Frankreich), Ligurien und Toscana (NW Italien)

(Orchidaceae und Insecta, Apoidea). *Journal Europäischer Orchideen*. 31, 347-422.

Paulus, H.F. (2017) Bestäubungs-biologie *Ophrys* in Nordspanien. *Journal Europäischer*

Orchideen. 49, 427-471.

Phillips, S.J., Anderson, R.P., & Schapire, R.E. (2006) Maximum entropy modelling of species

geographic   distributions.   *Ecological   Modelling*,   190,   231-259.   doi:

10.1016/j.ecolmodel.2005.03.026

Phillips, S. J., & Dudík, M. (2008). Modelling of species distributions with Maxent: New

extensions and a comprehensive evaluation. *Ecography*, 31, 161–175. doi:

doi.org/10.1111/j.0906-7590.2008.5203.x

Philipps, S.J., Anderson, R.P., Dudík, M., Shapire, R.E., & Blair, M.E. (2017) Opening the black

box: an open-source release of Maxent. *Ecography*, 40, 887-893. doi:

doi.org/10.1111/ecog.03049

Reich, I., Gormally, M., Allcock, A.L., McDonnel, R., Castillejo, J., Iglesias, J., ... Smith, C.J.

(2015) Genetic study reveals close link between Irish and Northern Spanish

specimens   of the protected Lusitanian slug *Geomalacus maculus*. *Biological Journal of the*

Linnean   *Society*, 116, 156-168. doi: 10.1111/bij.12568

Rochette, N.C., & Catchen J.M. (2017) Deriving genotypes from RAD-seq short-read data

using   Stacks. *Nature Protocols*, 12, 2640-2659. doi:10.1038/nprot.2017.123

Salvado, P., Aymerich Boixader, P., Vila Bonfill, A., Martin, M., Quélennec, C., Lewin, J.-M.,

Hinoux, V., & Bertrand, J.A.M. (2022) Little hope for the polyploid endemic Pyrenean

Larkspur (*Delphinium montanum*): evidences form population genomics and

Ecological   Niche Modeling. *Ecology and Evolution*, 12, e8711. doi: 10.1002/ece3.8711

Sedeek, K.E.M., Scopece, G., Staedler, Y.M., Schönenberger, J., Cozzolino, S., Schiestl, F. P., &

Schlüter, P.M. (2014) Genic rather than genome-wide differences between sexually

deceptive *Ophrys* orchids with different pollinators. *Molecular Ecology*, **23**, 6192-

6205. doi: 10.1111/mec.12992

Weir, B.S. & Goudet, J. (2017) A unified characterization of population structure and

relatedness.   *Genetics*,   206,   2085-2103.   doi:

https://doi.org/10.1534/genetics.116.198424

Table 1. Sampling site, sample size (*n*), geographic coordinates and elevation, population

summary statistics: number of private alleles, expected and observed heterozygosity (H_E and

H_O), inbreeding coefficient (G_{IS}) and nucleotide diversity (π)

Sampling site	*n*	Latitude (°)	Longitude (°)	Elevation (m)	Allelic richness (A_R)	Private alleles	H_O	H_E	G_{IS}	π
1-Guilhaumard	15	43.85	3.19	726	17043.95	0	0.262	0.282	0.071	0.282
2-Lapanouse-de-Cernon	15	43.99	3.09	651	16833.53	2	0.260	0.276	0.059	0.276
3-Saint-Affrique	13	43.98	2.93	585	16669.70	1	0.268	0.269	0.006	0.269
4-Valgañón	15	42.32	-3.08	1023	17167.45	24	0.268	0.291	0.081	0.290
5-Larraona	15	42.78	-2.28	812	17297.27	9	0.269	0.291	0.076	0.290
6-Bercedo	13	43.09	-3.44	774	17102.52	13	0.264	0.283	0.066	0.281
	86									

744 Table 2. Summary of the main demographic parameters inferred with DILS for each *Ophrys*

745 *aveyronensis* subsp. *aveyronensis* and <u>*Ophrys aveyronensis*</u> subsp. *vitorica* population pairs

746 (all details are provided in Table S4). N_e 1, N_e 2, N_e anc. correspond to effective population

747 sizes estimated for population 1, population 2 and the ancestral population, t_{split} to the time

748 of split between population 1 and population 2 in number of generations, respectively.

Population 1	Population 2	N_e 1	N_e 2	N_e anc.	t_{split}
Guilhaumard	Valgañón	19479	10389	129860	3837
Guilhaumard	Larraona	37410	44684	103916	2568
Guilhaumard	Bercedo	7302	26772	121691	2050
Lapanouse	Valgañón	20495	89214	120560	2476
Lapanouse	Larraona	29891	9197	114966	2853
Lapanouse	Bercedo	15855	7318	121961	3023
St-Affrique	Valgañón	16668	31253	104177	6884
St-Affrique	Larraona	4830	13282	120745	4432
St-Affrique	Bercedo	2540	66049	127017	2107

Figure 1 Map displaying the 6 sites sampled: 3 for *Ophrys aveyronensis* subsp. *aveyronensis*:

1-Guilhaumard, 2-Lapanouse-de-Cernon, 3-St-Affrique (in blue) and 3 for *O. a.* subsp.

vitorica: 4- Valgañón, 5-Larraona, 6-Bercedo (in red). The approximate known geographic

distribution is highlighted for both taxa with the same color code.

Figure 2 A) Principal Component Analysis (PCA) displaying the two first axes (PC1 and PC2)

representing 8.50% and 3.88% of the total genetic variance. PCA was computed based on 86

individuals genotyped at 9,301 SNP and colors depict sampling localities. B) Values of the

cross-entropy criterion for a number of clusters ranging from $K = 1$ to 10 (10 sNMF runs

each). The optimal number of K was found to be 5. C) Barplots of ancestry coefficients

obtained from sNMF for 86 individuals for $K = 2$ and $K = 5$.

Figure 3 Demographic history of the 6 studied populations of *O. a.* subsp. *aveyronensis*

(shades of blue) and *O. a.* subsp. *vitorica* (shades of red) depicting the mean effective

population size N_e (and associated 95 confidence intervals) over time as inferred from

STAIRWAYPLOTv2, assuming a mutation rate of 7×10^{-9} and a generation time of 5 years.

Figure 4 Schematic overview of the retained demographic scenario and range of associated

parameters inferred for each pair of populations (*O. aveyronensis* subsp. *aveyronensis*/ *O. a.*

subsp. *vitorica*). N_a: ancestral effective population size, N_1: (current) effective population size

for the *O. a.* subsp. *aveyronensis* population (given in number of individuals), N_2: Current)

effective population size for the *O. a.* subsp. *vitorica*, *founders1*: effective population size of

O. a. subsp. *aveyronensis* before T_{dem1} and *founders2*: effective size of *O. a.* subsp. *vitorica*

before T_{dem2} (each of them given in proportion of N_a). T_{split}: time of split between the

populations, T_{dem1}: time of *O. a.* subsp. *aveyronensis* effective population size change and

T_{dem2}: time of *O. a.* subsp. *vitorica* effective population size change (all in number of

generations).

Figure 5 Spatial projections of the bioclimatic niches inferred for the *Ophrys aveyronensis*

species complex depicting climate suitability at different time periods from the Last Glacial

Maximum (LGM, about 21 000 kyrs ago) and with projections for the Heinrich Stadial1 (17.0-

14.7 kyrs BP), the Bølling-Allerød (14.7-12.9 kyrs BP) the Younger Dryas Stadial (12.9-11.7

kyrs BP) and the Holocene: early-Holocene, Greenlandian (11.7-8.326 kyrs BP), mid-

Holocene, Northgrippian (8.326-4.2 kyrs BP) and late-Holocene, Meghalayan (4.2-0.3 kyrs

BP) and currently.

3-St-Affrique 2-Lapanouse
1-Guilhaumard

6-Bercedo

5-Larraona

4-Valgañón

0   75   150 km