Holocene climate change promoted allopatric divergence and

explains the current disjunct geographic distribution of the *Ophrys*

***aveyronensis* species complex (Orchidaceae)**

Anaïs Gibert[1], Roselyne Buscail[2], Michel Baguette[3,4], Christelle Fraïsse[5], Camille

Roux[5], Bertrand Schatz[6] & Joris A. M. Bertrand[1]

[1] Laboratoire Génome et Développement des Plantes (LGDP), UMR 5096, Université de

Perpignan Via Domitia (UPVD) - Centre National de la Recherche Scientifique (CNRS) -

Institut de Recherche pour le Développement (IRD), EMR 269 MANGO, F-66860 Perpignan.

[2] Centre de Formation et de Recherche sur les Environnements Méditerranéens (CEFREM),

Université de Perpignan Via Domitia (UPVD) - Centre National de la Recherche Scientifique

(CNRS), F-66860 Perpignan.

[3] Institut Systématique, Évolution, Biodiversité (ISEB), UMR 7205, Museum National

d'Histoire Naturelle (MNHN) - Centre National de la Recherche Scientifique (CNRS) -

Sorbonne Université - École Pratique des Hautes Études (EPHE) - Université des Antilles, F-

75005 Paris.

[4] Station d'Écologie Théorique et Expérimentale (SETE), UMR 5321, Centre National de la

Recherche Scientifique (CNRS) - Université Toulouse III, F-09200 Moulis.

[5] CNRS, Univ. Lille, UMR 8198 – Evo-Eco-Paleo, F-59000 Lille.

[6] Centre d'Écologie Fonctionnelle et Évolutive (CEFE) - Université de Montpellier - Centre

National de la Recherche Scientifique (CNRS) - École Pratique des Hautes Études (EPHE) -

Institut de Recherche pour le Développement (IRD), F-34293 Montpellier.

Correspondence: anais.gibert@gmail.com; joris.bertrand@univ-perp.fr

Abstract

Species with disjunct geographic distributions provide natural opportunities to investigate recent or incipient allopatric divergence. Although not rare, many of the cases observed result from successful colonization from a historical to a new range through dispersal or human-induced introduction which make the underlying eco-evolutionary processes sometimes difficult to decipher. The *Ophrys aveyronensis* species complex presents a disjunct geographic distribution with two ranges currently separated by 600 km on both sides of the Pyrenees Mountains. To uncover the causes of such intriguing biogeographic pattern, we combined population genomics and Ecological Niche Modelling approaches. Population genomic data show that all the populations studied display similar patterns of genetic diversity and dramatic decrease in effective size. We found significant genetic differentiation between the two ranges of the *O. aveyronensis* species complex. Our results support a very recent divergence (*ca.* 1500 generations ago). Ecological Niche Modelling results further indicate that the disjunct geographic distribution of the *O. aveyronensis* species complex is consistent with a range split of a broad ancestral range, contraction and shifts in opposite directions in response to climate warming during the Holocene. The congruence of both the results of population genomics and ENM approaches demonstrates that continental allopatric divergence is involved in the *Ophrys* adaptive radiation. The *O.*

45   *aveyronensis* complex is a promising candidate to study the onset of reproductive isolation

46   immediately following an initial stage of geographic separation.

48   **Keywords** Population genomics**,** RAD sequencing, Species Distribution Modelling (SDM),

49   Biogeography, Speciation, Orchids

Introduction

Allopatric speciation (also referred as geographic speciation, or vicariant speciation) occurs when populations of a given species become reproductively isolated from each other because of geographic separation. Aside from being conceptually intuitive, allopatric speciation has been evidenced by theoretical models, laboratory experiments and empirical studies to the point that it is generally regarded as the "default mode" of speciation (see Coyne & Orr, 2004). In allopatric speciation, spatial and reproductive isolation are often assumed to result from geologic-caused topographic changes that form the geographic barrier. In continental systems however, allopatric divergence is prone to experience only transient geographic and reproductive isolation involving one of several phases of secondary contact (e.g. through climate change-induced oscillations during the Quaternary, Hewitt, 2000). Thus, allopatric divergence is a spatio-temporally dynamic process. Aside, not only the fore-mentioned barriers but any kind of fragmentation of species distribution range, including human activities may also initiate and be responsible for evolutionary divergence.

The putative complexity of the speciation process makes difficult to elucidate its ecological and evolutionary causes as current spatial patterns of distribution only provide a snapshot from the speciation continuum (Nosil et al., 2017; Stankowski & Ravinet, 2021). In this regard, species with disjunct geographic distributions are likely to correspond to species whose populations became recently allopatric, *i.e.* with insufficient time to accumulate enough divergence to be considered as distinct species yet. Although such natural example may help to understand the consequences of physical barriers in early stages of evolutionary divergence, disjunct geographic distributions do not necessary result from the fragmentation of an ancestral range. Indeed, dispersal and successful colonization of a new range from the historical one may also lead to the same pattern.

For instance, several animal species including terrestrial mammals and mollusks (e.g.

Davison et al., 2001; Mascheretti et al., 2003; Grindon & Davison, 2013; Reich et al., 2015

and references therein) and plants (e.g. Beatty & Provan, 2013; 2014; Beatty et al., 2015;

Fagúndez & Díaz-Tapia, 2023) occur in Northern Iberia and in Ireland (a so called 'Lusitanian'

distribution) and display noticeable genetic similarities between these two regions. To

explain these disjunct geographic distributions, several authors have considered that all

these species may have survived in ice-free pocket (located in southwest Ireland) and in

southern European refugia (such as the Iberian Peninsula) during the Last Glacial Maximum

(LGM, 21000 years *Before Present*) or may have colonized Ireland thanks to land-bridges,

that disappeared ever since. However, molecular data have more recently supported that

most of the Irish animal populations have been introduced in the Mesolithic times (more

than 8000 years ago) through human activities whereas plants could have been more prone

to accomplish post-glacial recolonization of this region. In such examples, the combination of

genetic data with sufficient resolution and/or Environmental Niche Modelling (ENM) have

been crucial to compare the different scenarios and better understand the biogeography of

both species and regions.

The species of the genus *Ophrys* have long been known for their unusual pollination

syndrome (called 'sexual swindling') that involves insect-like flowers mimicking their female

pollinator to lure conspecific males and ensure pollination but are now also increasingly

considered as a promising biological model to investigate adaptive radiations. The degree of

plant-pollinator specialization (Joffard et al., 2019) required to ensure the success of the

Ophrys pollination strategy was shown to be the driver of the impressive rate of

diversification of the genus within the Mediterranean basin, and to a lesser extent in other

parts of Western Europe (see Baguette et al., 2020, for a recent review). If *Ophrys* are

notorious for their relatively high number of likely cases of speciation with gene flow, the

number of island endemic taxa suggests that geographic isolation induced by insularity is

also a key factor to explain *Ophrys* species richness observable nowadays (see Bertrand et

al., 2021a).

On continents, assessing the role of allopatry in *Ophrys* diversification is challenging,

in particular because clear evidences of taxa displaying subranges without being connected

by gene flow are lacking. Nevertheless, one species, *Ophrys aveyronensis*, displays an

intriguing case of disjunct geographic distribution. This species was initially described as

endemic to a very restricted area in the Grands Causses region (Southern France).

Observations of morphologically very similar populations were reported later from Northern

Spain and described as a distinct species (*Ophrys vitorica*, Kreutz, 2007). *Ophrys aveyronensis*

is now rather considered as a species complex consisting of two subspecies (*O. aveyronensis*

subsp. *aveyronensis* and *Ophrys aveyronensis* subsp. *vitorica*) that share, at least a mutual

insect pollinator (*i.e.* the solitary bee *Andrena hattorfiana*, Paulus & Gack, 1999; Paulus,

2017; Benito Ayuso, 2019 (but see Gibert et al., 2022 reporting subtle differences in

morphometry and coloration)). This taxon is thus present in two rather restricted ranges

separated by a gap of about 600 km (and Pyrenees Mountains), in the Grands Causse region

(France) and in Northern Spain.

The formation of the Pyrenees starting during the Eocene (from 45 Myrs ago) cannot

explain by itself the split of the ancestral range of the species as all phylogenetic hypotheses

published so far conclude that the *Ophrys* genus emerged no more than 5-6 Myrs ago and

that the *Sphegodes* clade, to which the *O. aveyronensis* complex belongs may have

diversified over the last million year (Breitkopf et al., 2015, see also Inda et al., 2012). This

does not rule out allopatry as factor to explain current disjunct geographic distribution but

suggests that geographic separation would have occurred more recently. According to this

'vicariance hypothesis', the current disjunct geographic distribution may correspond to two

relict pockets of an ancestral range that got fragmented following a scenario according to

which populations from Northern and Southern ranges diverge in allopatry. In this case, the

degree of genetic differentiation between Northern and Southern populations is expected to

be significant and positively correlated with the duration of reproductive isolation induced

by geographic separation. Namely, genetic differentiation is expected to be relatively higher

between Northern and Southern ranges than between populations within subranges.

Alternatively, we also envisage a 'biological introduction hypothesis': a colonization from

one historical range to the other as a result of human activities, or not. The notoriously small

Ophrys seeds are likely to be wind-dispersed over hundreds of kilometers (*i.e.* thus being

able to travel across most physical barriers including mountains and vast open water areas)

and may have found in the newly colonized area, suitable conditions for their germination,

development and reproduction. Although 'orchid dusty seed rains' may allow to generate

founder populations with non negligeable genetic diversity, such newly established

populations typically consist of a subset of the genetic pool of the source population (*i.e.* a

relatively little divergent population with lower effective population size and genetic

diversity).

In this study, we used a population genomic approach (*i.e.* through a double digest

RAD-seq like protocol called 'nGBS') to first investigate patterns of genetic diversity and

differentiation among *Ophrys aveyronensis* populations from both part of its disjunct

geographic range. We then couple demographic inference analyses and Ecological Niche

Modelling approaches to test whether the current disjunct geographic distribution of this

species complex is rather consistent with either allopatry or recent introduction. Specifically,

we aimed at determining how recent the divergence between Northern and Southern

populations of the *O. aveyronensis* complex may be and whether or not this disjunct

geographic distribution may have originated with a colonization event from an ancestral

range to a new one.

Materials and Methods

Sampling and study sites

We used a minimally destructive sampling method to collect a total of 86 plant tissue

samples from six localities that span the entire geographic range of the *O. aveyronensis*

species complex: three in Southern France (Grands Causses region): 1-Guilhaumard, 2-

Lapanouse-de-Cernon and 3-Saint-Affrique, and three in Northern Spain: 4-Valgañón, 5-

Larraona and 6-Bercedo, in June 2019, Fig. 1, see also Gibert et al., 2022). At each sampling

site, we collected 6 to 10, 0.6 mm diameter leaf punches from 13 to 15 individuals which

were stored in 100% ethanol until DNA extraction (see Table 1 and Supplementary Table S1).

As *Ophrys aveyronensis* is a nationally protected plant species in France, sampling was done

under permit 'Arrêté préfectoral n°2019-s-16' issued by the 'Direction Régionale de

l'Environnement de l'Aménagement et du Logement (DREAL)' from the 'Région Occitanie',

on 07-May-2019). *Ophrys aveyronensis* is not legally protected in Spain.

DNA extraction, library preparation and genotyping

Genomic DNA extraction as well as library preparation and genotyping were subcontracted

to LGC Genomics GmbH (Berlin, Germany). A fractional genome sequencing strategy called

normalized Genotyping-by-Sequencing (nGBS) was used to subsample the *Ophrys*

aveyronensis genome for which we estimated a relatively high size of ~5-6 Gbp based on

flow cytometry techniques (Bertrand, unpublished data). The nGBS protocol makes use of a

combination of two restriction enzymes (PstI and ApeKI, in our case) to produce a

reproducible set of fragments across samples/individuals. It also comprises a normalization

step whose aim is to avoid further sequencing of highly repetitive regions. Illumina

technology was then used to aim at obtaining a minimum of 1.5 million reads per sample

(with 2 x 150 bp read length) for each of the 86 individually barcoded libraries.

Population genomic dataset processing

We used different scripts included in *Stacks* v.2.60 (Catchen et al., 2011; 2013) to build loci

from Illumina reads, *de novo* (*i.e.* without aligning reads to a reference genome). We first

used *process_radtags* to demultiplex and clean reads, then *denovo_map.pl* to build loci

within individuals, create a catalog and match all samples against it and finally *populations* to

further filter the SNPs obtained at the population level and compute basic population

genetic statistics. Before running the pipeline on the complete dataset, we first optimized

several key parameters: -*m* (the minimum number of identical raw reads required to form a

putative allele), -*M* (the number of mismatches allowed between alleles to form a locus) and

-*n* (the number of mismatches allowed between loci during construction of the catalog) by

running it on a subset of 12 individuals representative of the whole dataset (*i.e.* geographic

origin and coverage). As recommended by several authors, we varied M and n from 1 to 9

(fixing $M = n$) while keeping $m = 3$ (see Paris et al., 2017). The combination -*m* 3, -*M* 5 and -*n*

5 was found to be the most suitable to maximize the number of SNPs, assembled and

polymorphic loci in our case and was used to run *Stacks* on the whole data set (see

Supplementary Table S2). Then, we ran the *populations* script included in *Stacks*, from which

only SNPs that were genotyped in at least 80% of the individuals per population (-*r* 80) and

found in all six sampling sites were kept (-*p* 6), while SNPs exhibiting estimates of observed

heterozygosity greater than 70% (--*max-obs-het* 0.7) were filtered out to reduce the risk of

including remaining paralogs. We also discarded sites whose minor allele frequency was

lower than 5% (--*min-maf* 0.05). The genotype matrix of the remaining 9301 SNPs was then

exported in several formats for downstream analyses.

The following population genomic analyses aimed to i) infer patterns of genetic

diversity and divergence between subspecies/countries and ii) infer the demographic

scenario (and associated parameters) likely to best explain the current biogeographic

pattern observed in the *O. aveyronensis* species complex.

Genetic diversity, population differentiation and evolutionary divergence

To get an overview of the overall genetic diversity and differentiation among individuals and

populations, we first performed a Principal Component Analysis (PCA) based on a matrix of

86 individuals (as rows) and 9301 SNPs (as columns) coded as a *genlight* object with the R-

package *adegenet* (Jombart, 2008; Jombart & Ahmed, 2011). The number of private alleles

and the average nucleotide diversity (π) were directly available from the output of the

populations script from *Stacks*. We used Genodive v.3.04 (Meirmans, 2020) to compute

expected and observed heterozygosity (H_E and H_O, respectively) as well as to evaluate the

deviation from panmixia by computing G_{IS}. Allelic richness (A_R) was computed with the R-

package *hierfstat* (Weir & Goudet, 2017). Overall and pairwise genetic differentiation was

also assessed based on *G*-statistics (G_{ST}, G'_{ST}, G''_{ST}) as well as Jost's *D* also implemented in

Genodive. An Analysis of Molecular Variance (AMOVA) allowed to assess the proportion of

genetic variance at different hierarchical levels, especially between sampling sites among

subspecies/country (F_{SC}) and between subspecies/country (F_{CT}). The statistical significance of

the obtained values was estimated based on 10000 permutations.

To further investigate population structure and characterize putative

migration/admixture event, we used sNMF (Frichot et al., 2014) as implemented in the R-

package *LEA* (Frichot & François, 2015) to estimate individual ancestry coefficients based on

sparse non-negative matrix factorization algorithms. The number of ancestral populations

(genetic clusters) was varied from K = 1 to 10, and analyses were run with 10 replicates at

each value of K. We followed the cross-entropy criterion to determine the optimal number

of clusters and the ancestry proportions matrices (Q-matrices) obtained were plotted with

the R-package *pophelper* (Francis, 2017).

Finally, we performed Maximum-Likelihood phylogenetic tree reconstruction with IQ-

Tree v.2.2.6 (Minh *et al.*, 2020) based on the concatenated SNP matrix with 1000 replicates

(-B 1000) of Ultrafast Bootstrap Approximation (UFBoot) to assess nodes support. The

substitution model that best fitted the data was selected with *ModelFinder Plus*

(Kalyaanamoorthy *et al.*, 2017) and Ascertainment bias correction (-MFP+ASC) as

recommended for SNP data.

Demographic and evolutionary history reconstruction

We reconstructed the past demographic history of each of the six populations, separately,

using STAIRWAYPLOT2 (Liu & Fu, 2015; 2020). This program relies on the Site Frequency

Spectrum (SFS) to infer the temporal dynamics of effective population size (N_e) and has the

advantage of not requiring whole-genome sequence data or reference genome information.

First, we used the *easySFS.py* script (https://github.com/isacovercast/easySFS) to compute

the folded SFS of each population. To maximize the number of segregating sites, the

population was downsampled to 24, 24, 22, 24, 24 and 22 individuals for 1-Guilhaumard, 2-

Lapanouse, 3-St-Affrique, 4-Valgañón, 5-Larraona and 6-Bercedo haploid genomes,

respectively. We then ran STAIRWAYPLOT2 that fits a flexible multi-epoch demographic

model that estimates N_e at different time periods using the SFS to infer estimated N_e

fluctuation coinciding with coalescent events. We assumed a generation time of 5 years for

Ophrys and a mutation rate of 7×10^{-9} per site per generation for angiosperms autosomal

markers (following Krasovec et al., 2018) and performed 200 bootstrap replicates to

estimate 95% confidence intervals.

We then used the Approximate Bayesian Computation (ABC) framework

implemented in DILS (Fraïsse et al., 2021) to infer the best demo-genomic scenario and

associated parameters of the evolutionary history between *O. a.* subsp. *aveyronensis* and *O.*

a. subsp. *vitorica*. We considered two types of datasets i) a pooled dataset grouping the 3

populations *O. a.* subsp. *aveyronensis* together and the 3 populations of *O. a.* subsp. *vitorica*

together and ii) paired datasets consisting of all possible *O. a.* subsp. *aveyronensis* and *O. a.*

subsp. *vitorica* population pairs (3^2 = 9 pairwise comparisons). The two-population model of

DILS compares different scenarios in a hierarchical manner. First, current isolation models,

i.e. Strict Isolation (SI) versus Ancient Migration (AM) are compared to ongoing migration

models, *i.e.* Isolation Migration (IM) versus Secondary Contact (SC). Then, for the best

demographic model, DILS compares genomic models to test for background selection by

assuming variation in effective population size (N_e) across sites (*i.e.* hetero versus homo Ne

models are compared). For migration models (IM and SC), DILS also tests for selection

against migrants by assuming variation in migration rates $N.m$ across sites (*i.e.* hetero versus

homo $N.m$ models are compared). Depending on the best selected model, DILS provides

estimates and confident intervals for parameters such as the time of split (T_{split}), the

migration rate ($N.m$), current and ancestral effective population sizes ($Ne_{current}$ and Ne_{past})

and whenever relevant, the time of ancient migration (T_{AM}) or the time of secondary contact

(T_{SC}). We formatted the input file with a custom Perl script and used the online facility

(https://www.france-bioinformatique.fr/cluster-ifb-core/) to run computations with the

following settings and priors: *population_growth*: constant, *modeBarrier*: bimodal (*i.e.* there

is a class of species barrier loci with $N.m$ = 0 and a class of loci migrating at the background

rate $N.m$) for model type to test), *max_N_tolerated:* 0.1*, LMin* = 30, *nMin* = 10 (20 for the

pooled dataset), *mu* = 7.10^{-9}, *rho_over_theta*: 0.1, *N*_min = 100, *N_max*: 200000, *Tsplit_min*:

100, *Tsplit_max*: 25000, *M_min*: 0.4 and *M_max*: 4. To quantify the fit of the best model to

the data we conducted a goodness-of-fit test using 2000 simulations performed under the

best model based on the estimated parameter values.

Past and current Species Distribution Modelling

We used an Ecological Niche Modelling (or Environmental Niche Modelling, ENM or Species

Distribution Modelling, SDM) framework similar to the one we followed and described in

detail in Salvado et al., (2022) to infer the spatio-temporal dynamics of suitable habitat for

the *O. aveyronensis* species complex since the Last Glacial Maximum (LGM, 21 kyrs BP). We

then compared the results with those obtained from population genomic analyses. To model

and spatially project the current bioclimatic niche, we first downloaded 19 bioclimatic layers

at a resolution of 2.5 arc-minutes (~5 km^2) available from WorldClim 2 (Fick & Hijmans,

2017). Current climate data correspond to time averaged variables over the period 1970-

2000. To spatially project the inferred niche onto past bioclimatic conditions data, we then

downloaded paleoclimate data either from WorldClim v1.4 (Hijmans et al., 2005) and from

PaleoClim (www.paleoclim.org).

Paleoclimate data from WorldClim are available for both the LGM (21 kyrs ago) and

the Mid-Holocene (about 6 kyrs ago), downscaled at a resolution of 2.5 arc-minutes for at

least three commonly used General Circulation Models (GCMs) proposed by the National

Center for Atmospheric Research (NCAR) though the Community Climate System Model

(CCSM4), the Japan Agency for Marine-Earth Science and Technology, the Center for Climate

System Research of the University of Tokyo, and the National Institute for Environmental

Studies (MIROC-ESM) and the Max Plant Institute for Meteorology Earth System Model

(MPI-ESM-P). Only CCSM data are available from PaleoClim but over different time periods

over the Pleistocene (at a resolution of 2.5 arc-minutes): the LGM (as from CHELSA, Karger et

al., 2021), the Heinrich Stadial1 (17.0-14.7 kyrs BP), the Bølling-Allerød (14.7-12.9 kyrs BP)

the Younger Dryas Stadial (12.9-11.7 kyrs BP) and the Holocene: early-Holocene,

Greenlandian (11.7-8.326 kyrs BP), mid-Holocene, Northgrippian (8.326-4.2 ka BP) and late-

Holocene, Meghalayan (4.2-0.3 kyrs BP) (as from Fordham et al., 2017).

The *O. aveyronensis* occurrences dataset consists of a set of field observations as well

as from records downloaded from the gbif (www.gbif.org) and the iNaturalist

(www.inaturalist.org) databases before manual curation. The Ecological Niche Modelling

analyses were conducted thanks to the R-package *ENMwizard v.0.3.7* (Heming et al., 2018,

see also Bagley et al. 2020) which is a convenient wrapper of several tools. We first spatially

filtered occurrences to keep only those that were at least 5 km away from each other using

the R-package *spThin* (Aiello-Lammens et al., 2015). The calibration area for the models was

created as a buffer of 0.5° around the minimum convex polygon encompassing all

occurrences. From the 19 bioclimatic variables, we selected the less correlated ones (Person

correlation coefficient < 0.75) thanks to the R-package *caret* (Kuhn, 2019) and kept 7

variables: bio2, bio3, bio8, bio9, bio10 and bio17 (see details in Supplementary Appendix S5)

for further analyses.

We used the maximum entropy method (implemented in MaxEnt ver. 3.4.1, Phillips

et al., 2006; Philips & Dudík, 2008; Phillips et al., 2017) to calibrate models and evaluated

models' performance thanks to the package *ENMeval* (Muscarella et al., 2014) as

implemented in *ENMwizard*. We evaluated models using a geographic partition scheme of

type "block" and optimized two parameters of MaxEnt: the Regularization Multipliers (RM)

and the Feature Classes (FCs). RM was varied from 0.5 to 4.5, incremented by 0.5 whereas a

suite of 15 FCs (L, for Linear, P, for Product, Q, for Quadratic and H for Hinge) or combination

of them were evaluated: L, P, Q, H, LP, LQ, LH, PQ, PH, QH, LPQ, LPH, LQH, PQH, LPQH,

resulting in a total of 135 models. Model selection was done by computing the corrected

Akaike Information Criterion ("LowAIC"). Model accuracy was also evaluated by calculating

omission rates calculated when a 10^{th} percentile threshold is applied (x10ptp). The final

model was projected on current and paleoclimatic contexts.

Results

Population genomic dataset

We obtained a total of 150 694 350 of read pairs across the 86 individuals (1 382 768 - 7 370

366 of raw reads per ind., mean 3 504 519, SD = 1 282011.52) of which 290 431 210

barcoded reads (*i.e.* 96.36%) were retained, 10 819 014 (3.6%) were removed because RAD

cutsite was not found in the sequence and 138 476 reads (< 1%) were removed because of

low quality. We genotyped a total of 859 035 *loci* (composed of 135 255 870 sites) including

368 712 variant sites (SNPs) with *Stacks*. Average read depth ranged from 24.1X to 81.6X

(mean 48.7X, SD = 12.4X) based on the combination of parameters we used. After filtering

the data with *populations*, we finally kept 4302 *loci* from which we retained 9301 variant

sites (SNPs).

Genetic diversity, population differentiation and divergence

The PCA biplot shows that individuals are arranged by subspecies/country of origin along

PC1 (which explains 8.50% of the total genetic variance) and by sampling site along PC2

(which explains 3.88% of the total genetic variance) even though the sampling sites of 1-

Guilhaumard and 2-Lapanouse-de-Cernon slightly overlap (Fig. 2A). Based on PC2, intra-

population genetic variation seems minimal for 1-Guilhaumard and 2-Lapanouse-de-Cernon

and maximal for 3-Saint-Affrique and 5-Larraona. Basic population genetics statistics are

indicated in Table 1. All populations displayed weak but significant deviation from panmixia

with a mean G_{IS} = 0.060 (p < 0.01, 95% CI: 0.056-0.064) and population G_{IS} values ranging

from 0.006 in 3-Saint-Affrique to 0.081 in 4-Valgañón (all p < 0.01). Average nucleotide

diversity (π) varies from 0.269 in 3-Saint-Affrique to 0.290 in 4-Valgañón and 5-Larraona, and

allelic richness (A_R) from 16696.70 (in 3-Saint-Affrique) to 17297.27 (in 5-Larraona). Based on

the filtering procedure we followed, the number of private alleles was relatively low but

consistently higher for *O. a.* subsp. *vitorica* (*i.e.* from 9 to 23) compared to *O. a.* subsp.

aveyronensis (*i.e.* from 0 to 2).

Overall G-statistics show that the whole set of populations is genetically structured:

G_{ST} = 0.091 (p < 0.01, 95% CI: 0.089-0.093); G'_{ST} = 0.134 (p < 0.01, 95% CI: 0.105-0.011); G''_{ST}

= 0.150 (p < 0.01, 95% CI: 0.146-0.153) and Jost's D = 0.047 (p < 0.01, 95% CI: 0.046-0.048).

The AMOVA confirms that although a major part of the genetic variance is found within

individual (81.8%, F_{IT} = 0.182, 95% CI: 0.177-0.186), significant proportions of the total

genetic variance are found among populations within subspecies/country (6.8%, F_{SC} = 0.072,

95% CI: 0.070-0.074) and between subspecies/country (6.1%, F_{CT} = 0.061, F_{CT} = 0.058-0.064).

Pairwise differentiation between sampling sites varies from F_{ST} = 0.049 to 0.082 within

subspecies/country and reaches F_{ST} = 0.111 to 0.147 between subspecies/country of origin

(all p < 0.01; see Supplementary Appendix S3 for details).

The clustering analysis carried out with sNMF suggests an optimal number of genetic

clusters of K=5 based on the cross-entropy criterion (Fig. 2B). We plotted the Q-matrices

(that contain individual admixture coefficients) corresponding to the best replicate runs for

K=2 (to investigate whether the program could distinguish the two subspecies) and K=5 (Fig.

2C). For K=2, the individuals of $O. a.$ subsp. $aveyronensis$ and $O. a.$ subsp. $vitorica$ were

unambiguously assigned to two distinct clusters. For the most likely combination of K=5, we

observe one cluster per sampling site except for the geographically close localities of 1-

Guilhaumard and 2-Lapanouse-de-Cernon that were grouped together in a unique cluster. In

the locality of 3-Saint-Affrique 4 out of 13 individuals (19-Oav-039, 19-Oav-040, 19-Oav-049

and 19-Oav-051) display >50% of ancestry proportion that are associated to the cluster

associated with 1-Guilhaumard and 2-Lapanouse-de-Cernon.

The phylogenetic tree inferred from the SNP matrix corroborates these results. $O. a.$

subsp. $aveyronensis$ and $O. a.$ subsp. $vitorica$ appeared as two well supported reciprocally

monophyletic groups. Sub-clades corresponding to sampling sites also confirm a

phylogenetic sub-clustering consistent with geography in each country for each subspecies

(with the exception of a group consisting of a mixture of individuals of all sites in France).

Inference of past demographic history

Reconstruction of historical dynamics of effective population size using STAIRWAYPLOT2

shows congruent trends consisting of progressive >95% decline in estimated N_e for all 6

populations, since the LGM, or at least, over the last 25000 years (see Fig. 3). Current

effective population sizes were estimated to vary between 302 (2-Lapanouse-de-Cernon)

and 642 individuals (3-St-Affrique) with this method. Despite the current geographical

discontinuity between *O. a.* subsp. *aveyronensis* and *O. a.* subsp. *vitorica* populations, we

attempted to statistically compare demographic scenarios with and without ongoing

migration using DILS. The Posterior Probabilities associated with the models of ongoing

migration (*i.e.* IM and SC) relative to models without (*i.e.* SI and AM) was 0.59 based on the

pooled dataset and ranged from 0.55 to 0.66 based on the population pairs analyses. These

probabilities are low and lead to ambiguous model choices. Therefore, we cannot draw

strong conclusions from model comparisons with DILS (see Table 2 and Appendix S4). This

can happen when the real history is an intermediate model, or because the separation time

between *O. a.* subsp. *aveyronensis* and *O. a.* subsp. *vitorica* is recent compared to the

current population sizes. The median of the parameters estimated based on posterior

distributions made under the most probable model (*i.e. Isolation with continuous Migration*,

IM) indicate very low migration rates ($N.m$ = 0.60 for the pooled dataset, and it ranges from

0.56 to 0.92 based on population pairs). With the pooled dataset, we inferred a subdivision

of the ancestral population of about 75 000 individuals based on two pools (90 000 to 103

200 based on population pairs) into two daughter populations about 1500 generations ago

(650 to 3880 based on population pairs), which convert into 4500 years assuming a

generation time of 3 years, 7500 years assuming a generation time of 5 years and 10500

years assuming a generation time of 7 years, an globally less based on population pairs

estimates. This subdivision is indeed considered recent as it is about five times smaller than

current population sizes. This suggests that most coalescence events are taking place in the

ancestral population and not in the respective daughter populations. In other words, the

amount of shared polymorphism is still relatively high because of insufficient time for drift to

differentially fix/eliminate alleles within the two subranges (*i.e.* Incomplete Lineage Sorting).

Similar to what was inferred with STAIWAYPLOT2, our analyses also point to a decrease of

effective sizes compared to the ancestral population (*i.e.* current N_e have values

corresponding to about 10% of $N_{anc.}$. The goodness-of-fit test indicates that the best

estimated scenario reproduces the observed data set with high fidelity (see Supplementary

Appendix S4).

Current and past Species Distribution Modelling

ENMeval analyses identified RM = 1 and a mixture of Product (P), Quadratic (Q) and Hinge

(H) feature classes as the best-performing parameters for calibrating the final ENM. These

parameters yielded a single "best" candidate model with the lowest AICc score (925.02) and

an AICc weight of ~0.23. This model had mean omission rates of 0.165 (for the 10^{th}

percentile) and a mean test AUC of 0.834 (see variable contributions following this model in

Table 3 and Supplementary Appendix S5, for additional details). The spatial projection of the

current bioclimatic inferred for the *O. aveyronensis* species complex corresponds to its

actual geographic distribution but show that some additional areas may be climatically

suitable for this orchid (Fig. 5). Spatial projections considering paleoclimatic conditions show

that the extent of suitable habitat for the *O. aveyronensis* species complex may have

reached its apex during the Younger Dryas cold period (12.9-11.7 kyrs BP) before starting to

decrease across the Holocene. The LGM conditions (21 kyrs BP) were not suitable for the

establishment of viable populations in the studied area but important suitable habitats may

have existed during the Heinrich Stadial 1 event (17-14.7 kyrs BP) and the Bøllering-Allerød

warming (14.7 − 12.9 kyrs BP). Altogether, these results are consistent with the possible

existence of a unique ancestral area, centered on South-Western France that could have

experienced a split and a combination of reduction and shift in two distinct directions during

the Holocene. The results available from WorldClim data do not contradict these conclusions

but show significant differences across GCMs and even for a given GCM but downscaled

based on different interpolation algorithms: namely, WorldClim instead of CHELSA (see

Supplementary Appendix S5).

Discussion

Some *Ophrys* orchid lineages exhibit among the highest rate of speciation reported so far

(Breitkopf et al., 2015) and are being increasingly mentioned as biological model of interest

to investigate ecological speciation and adaptive radiations (e.g. Nosil et al., 2017; Baguette

et al., 2020; Nürk et al., 2020). The selection pressures exerted by pollinators and the high

degree of specificity necessary to ensure the success of the sexual swindling strategy is likely

to explain the overwhelming importance of ecological and sometimes sympatric speciation

in *Ophrys* (see Baguette et al., 2020). This does not preclude for a significant role of allopatry

in *Ophrys* diversification, even in non-insular systems (see Bertrand et al., 2021a). Our study,

is to our knowledge, the first to thoroughly characterize a scenario of very recent allopatric

divergence in a continental *Ophrys* species. The *Ophrys aveyronensis* species complex

therefore provides a promising opportunity to investigate the relative contribution of

biogeographic and ecological factors in shaping early stages of evolutionary divergence in

rapidly diversifying groups of organisms.

Patterns of genetic diversity and differentiation based on population genomics

Although Northern and Southern individuals of *Ophrys aveyronensis* only present subtle

differences based on phenotypic data (Gibert et al., 2022), the French and Iberian

counterparts of this species complex can be unambiguously teased apart based on

population genomics data. Both the PCA, the *F*-statistics (and/or their analogues), the

clustering analysis and the phylogenetic tree confirm that the two taxa form distinct genetic

entities, themselves exhibiting a lower genetic substructure between sampling sites within

subranges. Altogether, these results show that each sampled population of *O. aveyronensis*

is more genetically differentiated from populations from the other region than from

populations of its own region. This pattern *a priori* rules out a recent colonization and

expansion from one range to the other from a unique source population. Patterns of genetic

diversity corroborate this conclusion. We found that all the populations sampled displayed

similar levels of allelic richness, heterozygosity, departure from panmixia and number of

private alleles. There is therefore no evidence for any of these six populations to appear as a

subsample of the genetic diversity of a putative source population elsewhere. Finally, the

inference of the temporal dynamics of effective population sizes suggests that each sampled

population may be considered as kind of replicate of each other. All the populations

displayed consistent decreases in effective population size over the last tens of thousands of

years which suggests that they shared the same demographic history. In summary,

population genomics results are fully consistent with a scenario of range split, and

subsequent contractions of two subranges experiencing restricted gene flow.

Delforge (2016) hypothesized that the Iberian populations may result from a hybrid

swarm (without mentioning the putative parental species) undergoing spatial and

demographic expansion. None of our results support the hypothesis according to which

Northern and Southern populations of *Ophrys aveyronensis* may result from parallel

convergent evolution from distinct regional hybrid swarms. However, the demographic

scenarios we tested remain relatively simple and the Bayesian Posterior Probabilities we

obtained with DILS show that the approach implemented in this program could not strongly

support Migration over a Isolation models. Events that are known to deceives demographic

inferences such as 'ghost introgression' from an unsampled lineage have a non-negligible

probability in *Ophrys*. Investigating whether the *Ophrys aveyronensis* complex has or not a

hybrid origin and at least whether the two taxa have been introgressed, and to which extent,

by other *Ophrys* taxa will thus require the genomic analyses of additional species in a future

study, including data with higher marker density (transcriptomes or whole-genome

resequencing data).

On the causes of the disjunct geographic distribution of the *O. aveyronensis* complex

The Ecological Niche Modelling results show that the LGM conditions were not suitable for

the establishment of viable populations of the *O. aveyronensis* species complex. This

suggests that either this species complex diversified from an ancestral *Ophrys* lineage in the

study area after this period, or that the populations of the complex occupied a relatively

remote glacial refugia during this period (the ENM results suggest that the Italian peninsula

could have provided such refugia, see Fig 5). Over the last 21000 years, the extent of the

habitat suitable for *O. aveyronensis* was rather large until the Holocene (about 12000 years

ago) (Fig. 5) where its area started to dramatically decrease while keeping an area of

relatively high suitability centered on the Southwest of France. The split and, to a lesser

extent, the shift Northward and Southward may have occurred during the Late Holocene

(starting about 4000 years ago) (Fig. 5). This timing is on the same order of magnitude than

the time of split we inferred based on population genomic data and is consistent with a

scenario of allopatric divergence (without being able to fully exclude transient gene flow

through climate-induced oscillations all along the Holocene) from an ancestral population

with relatively high effective size.

The Pyrenees mountains have the particularity to act as a biogeographic boundary

for many terrestrial organisms while being at the crossroads between temperate and

Mediterranean biomes (see Pironon et al., 2022 and references therein). Although orchid

propagules have the capacity to efficiently spread across long distances, several taxa occur

on either the Southern or the Northern slopes of the Pyrenees but have ranges that do not

extend latitudinally on both sides, or only weakly so (*i.e.* the so-called Northern or Southern

peripheral species in Pironon et al., 2022). The Pyrenees may thus alter the dispersal of at

least some orchid populations (e.g. *Dactylorhiza insularis*, *Gymnadenia gabasiana*, *Neotinea*

conica, *Ophrys castellana*, *O. riojana* and *O. subinsectifera*, *Orchis langei* and *O. simia*)

because of various biotic or abiotic factors. This mostly granitic and relatively high mountain

channel forms a vast area of unsuitable habitat for *Ophrys* orchids that generally require

calcareous soils and lowland or midland environments. Although data on their geographic

distribution remain incomplete, we may also speculate that the absence of *Ophrys* specific

pollinator species and/or of fungi to which orchid depend for their germination and their

development may also have limited the possibility of establishment for intermediate orchid

populations between the two current subranges. For *O. aveyronensis*, our results are

consistent with an overall reduction of gene flow between both sides of the Pyrenees

through the formation and the contraction of the two subranges during the Holocene until

the contemporary arrest of gene exchange that seems to be too recent to be determined

based on current data.

Evidence for ecological speciation in addition to vicariance?

Selection pressures exerted by insect pollinators are likely to be the main driver of ecological

and putatively sympatric speciation in *Ophrys* orchids (see Baguette et al., 2020). In this way,

populations of a given *Ophrys* species being pollinated by distinct insect pollinator species

may be considered as engaged in a speciation process. So far, field observations support that

O. aveyronensis from the Grands Causses and their Iberian counterparts are pollinated by

the same pollinator species: the solitary bee *Andrena hattorfiana* (see Benito Ayuso, 2019).

In a recent study on this system however, we showed that *O. aveyronensis* from the Grands

Causses and from the Iberian Peninsula present subtle differences at a set of morphological

and coloration traits that may be involved in pollination (Gibert et al., 2022). These subtle

differences could be emerging as a result of local adaptation to a "secondary" pollinator

species. Although additional work is required to address whether this variation may be

actually adaptive and involved in evolutionary divergence, these results call for i) an

assessment of which phenotypic traits may be under divergent selection across the *O.*

aveyronensis species complex geographic distribution and which ones may evolve through

drift only and ii) an investigation of putative genomic regions associated with phenotypic

trait values, genetic structure or ecological features, something that remains largely

underexplored in *Ophrys* (but see Sedeek et al., 2014, for a noticeable exception).

Carrying out cross-pollination experiments between individuals of the two subranges

may also help to highlight the insurgence of post-zygotic reproductive isolation within the

Ophrys aveyronensis species complex. However, such work may be impeded by the difficulty

to grow *Ophrys* under controlled conditions and because of the protection status of *Ophrys*

aveyronensis in France. Current knowledge suggest that such reproductive barriers may be

weak or absent at the *Ophrys* genus scale (Soliva and Widmer, 2003; Scopece et al., 2007,

Gervasi et al., 2017). To our best knowledge, only two noticeable exceptions to this pattern

involve the phylogenetically distant *Ophrys incubacea* and *Ophrys iricolor* (Cortis et al., 2009)

or species with different levels of ploidy: the diploid *Ophrys exaltata* and the tetraploid

Ophrys lupercalis (Vereecken et al., 2010).

Taxonomic and conservation implications

Considering that populations of the *O. aveyronensis* species complex are pollinated by the

same insect pollinator species across their range and have similar environmental niches, we

have currently no evidence to even consider the existence of different ecotypes in this

species complex. However, the disjunct geographic distribution in two subranges, separated

by ~600 km and the Pyrenees Mountains suggest that populations from the Grands Causses

and from the Iberian Peninsula may be considered as distinct evolutionary (and

conservation) units. Our population genomic analyses indicate that both sets of populations

are genetically differentiated and do not show evidence of ongoing gene flow. Accordingly,

the proposition to consider them as two subspecies: *Ophrys aveyronensis* subsp.

aveyronensis and *Ophrys aveyronensis* subsp. *vitorica*, (Paulus, 2017) seems appropriate.

Recent studies elaborated from some empirical evidence that delineating the so-

called 'micro-species' in *Ophrys* is practically cumbersome (Bateman et al., 2021; Bateman

and Rudall, 2023), especially in the section *Sphegodes* to which the *O. aveyronensis* species

complex belongs. Some genomic data such as whole plastomes that are useful to delineate

the main *Ophrys lineages*, present sequence similarity degree of more than 99.5% between

'microspecies' and therefore do not present enough variation to allow to discriminate them

(Bertrand et al., 2019; Bertrand et al., 2021b; Bateman et al., 2021). Here, however, our

results demonstrate that an approach consisting of sampling populations (*i.e.* a sufficient

number of individuals at a given location) and appropriate genetic markers (e.g. SNPs

genotyped on populations based on RAD-seq protocols) give signals that are consistent with

taxonomy. We therefore claim that such approaches of population genomics within an

integrative taxonomy framework (relying on flower morphology, pollinator identity,

ecological features…) is by far the most fruitful to better understand the process of

evolutionary divergence and the most relevant to improve at the same time our knowledge

on taxonomy, systematics, ecology and evolution of *Ophrys* orchids.

***Ophrys aveyronensis* as a promising biological model to study speciation**

Allopatric divergence is a very common route toward speciation (Hewitt et al., 2000), but

most of empirical cases document it retrospectively, between nascent species that exhibit at

the same time obvious geographic separation and substantial levels of both phenotypic and

genetic differentiation (Dool et al., 2021). By contrast studies reporting cases of incipient

allopatric divergence are rare, especially in continental systems. The allopatric divergence in

the *Ophrys aveyronensis* species complex we report here is extremely recent compared to

other species complex harboring disjunct geographic distribution in terrestrial (e.g.

González-Serna et al., 2018; Tomasello et al., 2020; Wang et al., 2022; Cao et al., 2022;

Balmori-de la Puente et al., 2022) or marine systems (Schmid et al., 2023; Marcionetti et al.,

2023). It is likely to have arisen because of climate warming that dissected an ancestral

geographic distribution and subsequently constrained the spatial dynamics of its subranges

and does not *a priori* imply a human-induced biological introduction. This system therefore

may serve as a natural laboratory to assess early stages of reproductive isolation insurgence,

immediately after geographic separation.

Author contributions

JB, BS, AG and MB designed the study. AG, JB, BS and RB gathered the data.

AG and JB performed the analyses with contribution from CF and CR and wrote the first

version of the manuscript. All authors contributed substantially to the revisions.

Data availability statement

Acknowledgements

This work was primarily supported by an inter-LabEX TULIP-CEMEB initiative (n°197310) to

Joris Bertrand and Bertrand Schatz. This research was also funded by an Agence Nationale

pour la Recherche Jeune Chercheur Jeune Chercheuse (ANR JCJC) grant to Joris Bertrand,

grant number ANR-21-CE02-0022-01, and is set within the framework of the "Laboratoires

d'Excellences (LABEX)" TULIP [ANR-10- LABX-41]. Lastly, this research was also funded by the

Observatoire de REcherche Méditerranéen de l'Environnement (SO Ocove & SO PolliMed,

OSU OREME) to Bertrand Schatz. We thank Jean-Luc Roux and Daniel Vizcaïno for the

information provided about sampling sites in Spain. We thank Christian Moliné for his

contribution to fieldwork, Moaine El Baidouri and Pascaline Salvado for their support with

bioinformatic analyses.

Conflict of interest

The authors declare no conflict of interest.

References

Aiello-Lammens, M.E., Boria, R.A., Radosavljevic, A., Vilela, B., & Anderson, R.P. (2015)

spThin: an R package for spatial thinning of species occurrence records for use in

ecological niche models. *Ecography*, 38, 541-545. doi: doi.org/10.1111/ecog.01132

Bagley, J.C., Heming, N.M., Gutiérrez, E. E., Devisetty, U.K., Mock, K. E., Eckert, A.J., & Strauss,

S.H. (2020) Genotyping-by-sequencing and ecological niche modeling illuminate

phylogeography, admixture, and Pleistocene range dynamics in quaking aspen

(*Populus tremuloides*) *Ecology and Evolution*, 10, 4609-4629. doi: 10.1002/ece3.6214

Baguette, M., Bertrand, J.A.M, Stevens, V.M., & Schatz, B. (2020) Why are there so many

bee-orchid species? Adaptive radiation by intra-specific competition for mnesic

pollinators. *Biol Rev.*, 95, 1630-1663. doi: 10.1111/brv.12633

Balmori-de la Puente, A., Ventura, J., Miñarro, M., Somoano, A., Hey, J. & Castesana, J.

(2022) Divergence time estimation using ddRAD data and an isolation-with-migration

model applied to water vole populations of *Arvicola*, *Scientific Reports*, 12, 4065. doi:

10.1038/s41598-022-07877-y

Bateman, R.M., Rudall, P. J., Murphy, A.R.M., Cowan, R.S., Devey, D.S. & Peréz-Escobar, O.A.

(2021) Whole plastomes are not enough: phylogenomic and morphometric

exploration at multiple demographic levels of the bee orchid clade *Ophrys* sect.

Sphegodes. Journal of Experimental Botany, 72, 654-681. doi: 10.1093/jxb/eraa467

Bateman, R.M. & Rudall, P.J. (2023) Morphological continua make poor species: genus-wide

morphometric survey of the European Bee Orchids (*Ophrys* L.). *Biology*, 12, 136. doi:

10.3390/ biology12010136

Beatty, G.E., & Provan, J. (2013) Post-glacial dispersal, rather than *in situ* glacial survival, best

explains the disjunct distribution of the Lusitanian plant species *Daboecia cantabrica*

(Ericacea). *Journal of Biogeography*, 40, 335-344. doi: 10.1111/j.1365-

2699.2012.02789.x

Beatty, G.E., & Provan, J. (2014) Phylogeographical analysis of two cold-tolerant plants with

disjunct Lusitanian distributions does not support *in situ* survival during the last

glaciation. *Journal of Biogeography*, 41, 2185-2193. doi: 10.1111/jbi.12371

Beatty, G.E., Lennon, J.J., O'Sullivan, C.J., & Provan, J. (2015) The not-so-Irish spurge:

Euphorbia hyberna (Euphorbiaceae) and the Littletonian plant 'steeplechase'.

Biological Journal of the Linnean Society, 114, 249-259. doi: 10.1111/bij.12435

Benito Ayuso, J. (2019) Estudios sobre polinización en el généro *Ophrys* (Orchidaceae), I,

Flora Montiberica, 74, 32-37.

Bertrand, J. A. M., Gibert, A., Llauro, C., & Panaud, O. (2019) Characterization of the

complete plastome of *Ophrys aveyronensis*, a Euro-Mediterranan orchid with an

intriguing geographic distribution. *Mitochondrial DNA Part B*, 4, 3256-3257. doi:

10.1080/23802359.2019.1670748

Bertrand, J. A. M., Gibert, A., Llauro, C., & Panaud, O. (2021) Whole plastid genome-based

phylogenomics supports an inner placement of the *O. insectifera* group rather than a

basal position in the rapidly diversifying *Ophrys* genus (Orchidaceae). *Biology Letters*,

168, 452-457. doi: 10.1080/23818107.2021.1893216

Bertrand, J. A. M., Baguette, M., Joffard, N., & Schatz, B. (2021) Challenges inherent in the

systematics and taxonomy of genera that have recently experienced explosive

radiation: the case of orchids of the genus *Ophrys*. In Grancolas, P. & Maurel, M.-C.

Systematics and the Exploration of Life (pp 113-134). Hoboken, New Jersey: Wiley.

Breitkopf, H., Onstein, R.E., Cafasso, D., Schlüter, P.M., & Cozzolino, S. (2015) Multiple shifts

to different pollinators fuelled rapid diversification in sexually deceptive *Ophrys*

orchids. *New Phytologist*, 207, 377-389. doi: 10.1111/nph.13219

Cao, Y., Almeida-Silva, F., Zhang, W.-P., Ding, Y.-M., Bai, D., Bai, W.-N., Zhang, B.W., Van de

Peer, Y.- & Zhang, D.Y. (2023) Genomic Insights into Adaptation to Karst Limestone

and Incipient Speciation in East Asian *Platycarya* spp. (Juglandaceae). *Molecular*

Biology and Evolution, 40, msad121. doi: 10.1093/molbev/msad121

Catchen, J., Amores, A., Cresko, W., & Postlewait, J. (2011) Stacks: building and genotyping

loci de novo from short-read sequences. *G3: genes, Genomes, Genetics*, 1, 171-182.

doi: 10.1534/g3.111.000240

Catchen, J., Hohenlohe, P.A., Bassham, S., Amores, A., & Cresko, W.A. (2013) Stacks: an

analysis tool set for population genomics. *Molecular Ecology*, 22, 3124-3140. doi:

/10.1111/mec.12354

Cortis, P., Vereecken, N.J., Schiestl, F.P., Barone Lumaga, M.R., Scrugli, A. & Cozzolino, S.

(2009) Pollinator convergence and the nature of species' boundaries in sympatric

Sardinian *Ophrys* (Orchidaceae). *Annals of Botany*, 104, 497-506. doi:

/10.1093/aob/mcn219

Coyne, J.A., & Orr, H.A. (2004). *Speciation*. Sunderland, Mass: Sinauer Associates.

Claessens, J., & Kleynen, J. (2016) *Orchidées d'Europe : fleurs et pollinisation*. Biotope, Mèze,

France: Biotope.

Davison, A., Birks, J.D.S., Brookes, R.C., Messenger, J.E., & Griffiths, H.I. (2001) Mitochondrial

phylogeography and population history of pine martens *Martes martes* compared

with polecats *Mustela putorius*. *Molecular Ecology*, 10, 2479-2488. doi:

10.1046/j.1365-294x.2001.01381.x.

Delforge, P. (2016) *Orchidées d'Europe, d'Afrique du Nord et du Proche Orient*. Lausanne,

Switzerland: Delachaux & Niestlé.

Dool, S.E., Picker, M.D. & Eberhard, M.J.B. (2021) Limited dispersal and local adaptation

promote allopatric speciation in a biodiversity hotspot. *Molecular Ecology*, 31, 279-

295. doi: 10.1111/mec.16219

Fagúndez, J. & Díaz-Tapia, P. (2023) Comparative phylogeography of a restricted and a

widespread heather: genetic evidence of multiple independent introductions of *Erica*

mackayana into Ireland from northern Spain. *Botanical Journal of the Linnean*

Society. doi: 10.1093/botlinnean/boac071

Fick, S.E., & Hijmans, R.J. (2017) WorldClim 2: new 1km spatial resolution climate surfaces

for global land areas. *International Journal of Climatology*, 37, 4302-4315. doi:

10.1002/joc.5086

Fordham, D.A., Saltré, F., Haythorne, S., Wigley, T. M. L., Otto-Bliesner, B. L., Chan, K. C., &

Brook, B.W. (2017) PaleoView: a tool for generating continuous climate projections

spanning the last 21 000 years at regional and global scales. *Ecography*, 40, 1348-

1358. Doi: doi.org/10.1111/ecog.03031

Fraïsse, C., Popovic, I., Mazoyer, C., Spataro, B., Delmotte, S., Romiguier, J., Loire E., Simon,

A., Galtier, N., Duret, L., Bierne, N., Vekemans, X., & Roux C. (2021) DILS :

Demographic inferences with linked selection using ABC. *Molecular Ecology*

Resources, 2629-2644. doi: doi.org/10.1111/1755-0998.13323

Francis, R.M. (2017) POPHELPER: an R package and web app to analyse and visualize

population structure. *Molecular Ecology Resources*, 17, 27-32. doi: 10.1111/1755-

0998.12509

Frichot, E., Mathieu, F., Trouillon T., Bouchard, G., & François O. (2014) Fast and efficient

estimation of individual ancestry coefficients. *Genetics*, 196, 973-983. doi:

10.1534/genetics.113.160572

Frichot, E. & François O. (2015) LEA: An R package for landscape and ecological association

studies. *Methods in Ecology and Evolution*, 6, 925-929. doi: 10.1111/2041-

210X.12382

Gervasi, D.L., Selosse, M.-A., Sauve, M., Francke, W., Vereecken, N.J., Cozzolino, S. & Schiestl,

F.P. (2017) Floral scent and species divergence in a pair of sexually deceptive orchids.

Ecology and Evolution, 7, 6023-6034. doi: 10.1002/ece3.3147

Gibert, A., Louty, F., Buscail, R., Baguette, M., Schatz, B. &, Bertrand, J.A.M. (2022) Extracting

quantitative information from images taken in the wild: a case study of two vicariants

of the *Ophrys aveyronensis* species complex. *Diversity*, **14**, 400. doi:

10.3390/d14050400

Grindon, A. J., & Davison, A. (2013) Irish *Cepea nemoralis* land snails have a cryptic Franco-

Iberian origin that is most easily explained by the movements of Mesolithic humans.

PLoS ONE, 8, e65792. doi: doi.org/10.1371/journal.pone.0065792

González-Serna, M., Cordero, P.J. & Ortego, J. (2018) Using high-throughput sequencing to

investigate the factors structuring genomic variation of a Mediterranean grasshopper

of great conservation concern. *Scientific Reports*, 8, 13436. doi: 10.1038/s41598-018-

31775-x

Heming, N. M., Dambros, C., & Gutiérrez, E. E. (2018). ENMwizard: AIC model averaging and

other advanced techniques in Ecological Niche Modeling made easy. R package

version 0.1.7. Retrieved from https:// github.com/HemingNM/ENMwizard

Hewitt, G. (2000) The genetic legacy of the Quaternary ice ages. *Nature*, 405, 907-913. doi:

10.1038/35016000/

Hijmans, R. J., Cameron, S.E., Parra, J. -L., Jones, P. J., & Jarvis A. (2005) Very high resolution

interpolated climate surfaces for global land areas. *International Journal of*

Climatology, 25, 1965-1978. doi: 10.1002/joc.1276

Inda, L.A., Pimentel, M., & Chase, M.W. (2012) Phylogenetics of tribe Orchideae

(Orchidaceae: Orchidoideae) based on combined DNA matrices: inferences regarding

timing of diversification and evolution of pollination syndromes. *Annals of Botany.*,

110, 71-90. doi: 10.1093/aob/mcs083

Joffard, N., Massol, F., Grenié, M., Montgellard, C. & Schatz, B. (2019) Effect of pollination

strategy, phylogeny and distribution niches of Euro-Mediterranean orchids. *Journal of*

Ecology, 107, 478-490. doi: 10.1111/1365-2745.13013

Jombart, T. (2008) adegenet: a R package for the multivariate analysis of genetic markers.

Bioinformatics, 24, 1403-1405. doi: 10.1093/bioinformatics/btn129

Jombart, T., & Ahmed I. (2011) adegenet 1.3-1: new tools for the analysis of genome-wide

SNP data. *Bioinformatics*. doi: 10.1093/bioinformatics/btr521

Kalyaanamoorthy, S., Minh, B.Q., Wong, T.F.K., von Haeseler, A. & Jermiin, L.S. (2017)

ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature*

Methods, 14, 587-589. doi: 10.1038/nmeth.4285

Karger, D. N., Nobis, M. P., Normand, S., Graham, C. H., & Zimmermann, N. E. (2021):

CHELSA-TraCE21k v1. 0. Downscaled transient temperature and precipitation data

since the last glacial maximum. *Climate of the Past Discussions*, 1-27. doi: 10.5194/cp-

2021-30

Krasovec, M., Chester, M., Ridout, K., & Filatov, D.A. (2018) The mutation rate and the age of

the sex chromosomes in *Silene latifolia*. *Current Biology*, 28, 1832-1838. doi:

10.1016/j.cub.2018.04.069

Kreutz, C.A. (2007) Beitrag zur Taxonomie und Nomenklatur europäischer, mediterraner,

nordafrikanischer und vorderasiatischer Orchideen. *Ber. Arbeitskrs. Heim. Orchid.* 24,

77-141.

Kuhn, M. (2019) Caret: classification and regression training. https://CRAN. R-

project.org/package=caret.

Liu, X., & Fu, Y.-X. (2015) Exploring population changes using SNP frequency spectra. *Nature*

Genetics, 47, 555-559. doi: 10.1038/ng.3254.

Liu, X., & Fu, Y.-X. (2020) Stairway Plot 2: demographic inference with folded SNP frequency

spectra. *Genome Biology*, 21, 280. doi: 10.1186/s13059-020-02196-9

Marcionetti, A., Bertrand, J. A.M, Cortesi, F., Donati, G.F.A., Heim, S., Huyghe, F., Kochzius,

M., Pelissier, L. & Salamin N. (2023) Recurrent gene flow events shaped the

diversification of the clownfish skunk complex. *bioRxiv*. doi:

10.1101/2023.10.24.562491

Mascheretti, S., Rogatcheva M. B., Gündüz, I., Fredga, K., & Searle, J.B. (2003) How did

pygmy shrews colonize Ireland? Clues from a phylogenetic analysis of mitochondrial

cytochrome *b* sequences. *Proceedings of the Royal Society B*, 270, 1593-1599. doi:

doi.org/10.1098/rspb.2003.2406

Meirmans, P. G. (2020) GENODIVE version 3.0: Easy-to-use software for the analysis of

genetic data of diploids and polyploids. *Molecular Ecology Resources*, 20, 1126-1131.

doi: 10.1111/1755-0998.13145

Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, A., von Haeseler, A. &

Lanfear, R. (2020) IQ-TREE 2: New methods and efficient methods for phylogenetic

inference in the genomic era. *Molecular Biology and Evolution*, 37, 1530-1534. Doi:

10.1093/molbev/msaa015

Muscarella, R., Galante, P.J., Soley-Guardia, M., Boria, R.A., Kass, J.M., Uriarte, M., &

Anderson, R.P. (2014) ENMeval: an R package for conducting spatially independent

evaluations and estimating optimal model complexity for Maxent ecological niche

models. *Methods in Ecology and Evolution*, 5, 1198-1205. doi: 10.1111/2041-

210X.12261

Nosil, P., Feder, J.L., Flaxman, S.M. & Gompert Z. (2017) Tipping points in the dynamics of

speciation. *Nature Ecology and Evolution*, 1, 0001. doi: 10.1038/s41559-016-0001

Nürk, M.N., Linder, H.P., Onstein, R.E., Larcombe, M.J., Hughes, C.H., Piñeiro Fernández, L.,

Schlüter, P.M., Valente, L.M., Beierkuhnlein, C., Cutts, V., Donoghue, M.J., Edwards,

E.J., Field, R., Flantua, S.G.A., Higgins, S.II, Liede-Schumann, S. & Pirie, M.D. (2020)

Diversification in evolutionary arenas – assessment and synthesis. *Ecology and*

Evolution, 10, 6163-6182. doi:/10.1002/ece3.6313

Paris, J., Stevens, J.R., & Catchen J.M. (2017) Lost in parameter space: a road map for

STACKS. *Methods in Ecology and Evolution*, 8, 1360-1373. doi: 10.1111/2041-

210X.12775

Paulus, H.F. & Gack, C. (1999) Bestäubungsbiologische Untersuchungen an der Gattung

Ophrys in der Provence (SO Frankreich), Ligurien und Toscana (NW Italien)

(Orchidaceae und Insecta, Apoidea). *Journal Europäischer Orchideen*. 31, 347-422.

Paulus, H.F. (2017) Bestäubungs-biologie *Ophrys* in Nordspanien. *Journal Europäischer*

Orchideen. 49, 427-471.

Pironon, S., Gómez, D., Font, X. & García, M.B. (2022) Living at the limit in the Pyrenees:

Peripheral and endemic plants are rare but underrepresented in protection lists.

Diversity and distribution, 28, 930-943. doi: 10.1111/ddi.13487

Phillips, S.J., Anderson, R.P., & Schapire, R.E. (2006) Maximum entropy modelling of species

geographic   distributions.   *Ecological   Modelling*,   190,   231-259.   doi:

10.1016/j.ecolmodel.2005.03.026

Phillips, S.J., & Dudík, M. (2008). Modelling of species distributions with Maxent: New

extensions   and   a   comprehensive   evaluation.   *Ecography*,   31,   161–175.   doi:

doi.org/10.1111/j.0906-7590.2008.5203.x

Philipps, S.J., Anderson, R.P., Dudík, M., Shapire, R.E., & Blair, M.E. (2017) Opening the black

box: an   open-source   release   of   Maxent.   *Ecography*,   40,   887-893.   doi:

doi.org/10.1111/ecog.03049

Reich, I., Gormally, M., Allcock, A.L., McDonnel, R., Castillejo, J., Iglesias, J., Quinteiro J. &

Smith, C.J. (2015) Genetic study reveals close link between Irish and Northern Spanish

specimens of the protected Lusitanian slug *Geomalacus maculus*. *Biological Journal of*

the Linnean Society, 116, 156-168. doi: 10.1111/bij.12568

Rochette, N.C., & Catchen J.M. (2017) Deriving genotypes from RAD-seq short-read data

using Stacks. *Nature Protocols*, 12, 2640-2659. doi:10.1038/nprot.2017.123

Salvado, P., Aymerich Boixader, P., Vila Bonfill, A., Martin, M., Quélennec, C., Lewin, J.-M.,

Hinoux, V., & Bertrand, J.A.M. (2022) Little hope for the polyploid endemic Pyrenean

Larkspur   (*Delphinium   montanum*):   evidences   form   population   genomics   and

Ecological Niche Modeling. *Ecology and Evolution*, 12, e8711. doi: 10.1002/ece3.8711

Schmid, S., Bachmann Salvy, M., Garcia Jimenez, A., Bertrand, J.A.M., Cortesi F., Heim, S.,

Huyghe, F., Kochzius, M., Pelissier, L. & Salamin, N. Gene flow throughout the

835 evolutionary history of a polymorphic and generalist clownfish. *bioRxiv*. Doi:

836 10.1101/2023/10.19.563036

837 Scopece, G., Musacchio, A., Widmer, A. & Cozzolino, S. (2007) Patterns of reproductive

838 isolation in Mediterranean deceptive orchids. *Evolution*, 61, 2623-2642. doi:

839 doi.org/10.1111/j.1558-5646.2007.00231.x

840 Sedeek, K.E.M., Scopece, G., Staedler, Y.M., Schönenberger, J., Cozzolino, S., Schiestl, F. P., &

841 Schlüter, P.M. (2014) Genic rather than genome-wide differences between sexually

842 deceptive *Ophrys* orchids with different pollinators. *Molecular Ecology*, 23, 6192-

843 6205. doi: 10.1111/mec.12992

844 Soliva, M. & Widmer, A. (2003) Gene flow across species boundaries in sympatric sexually

845 deceptive *Ophrys* (Orchidaceae) species. *Evolution*, 57, 2252-2261. doi:

846 10.1111/j.0014-3820.2003.tb00237.x.

847 Stankowski, S., Westram, A.M., Zagrodzka, Z.B., Eyres, I., Broquet, T., Johannesson, K., &

848 Butlin, R.K. (2020). The evolution of strong reproductive isolation between sympatric

849 intertidal snails. *Philosophical Transactions of the Royal Society B: Biological Sciences*,

850 375, 20190545. doi:10.1098/rstb.2019.0545

851 Stankowski, S. & Ravinet, M. (2021) Defining the speciation continuum. *Evolution*, 75, 1256-

852 1273. doi:10.1111/evo.14215

853 Tomasello, S., Karbstein, K., Hodač, L., Paetzold, C. & Hörandl, E. (2020) Phylogenomics

854 unravels Quaternary vicariance and allopatric speciation patterns in temperate-

855 montane plant species: A case study on the *Ranunculus auricomus* species complex.

856 *Molecular Ecology*, 29, 2031, 2049. doi: 10.1111/mec.15458

Vereecken, N.J., Cozzolino, S. & Schiestl, F.P. (2010) Hybrid floral scent novelty drives

pollinator shift in sexually deceptive orchids. *BMC Evolutionary Biology*, 10, 103. doi:

10.1186/1471-2148-10-103

Wang, R.-H., Yang, Z.-P., Zhang, Z.-C., Comes, H.P., Qi, Z.-C., Li, P. & Fu, C.-X. (2022) Plio-

Pleistocene climatic change drives allopatric speciation and population divergence

within the *Scrophularia incisa* complex (Scrophulariaceae) of desert and steppe

subshrubs in Northwest China. *Frontiers in Plant Science*, 13, 985372. doi:

10.3389/fpls.2022.985372.

Weir, B.S. & Goudet, J. (2017) A unified characterization of population structure and

relatedness. *Genetics*, 206, 2085-2103. doi:

https://doi.org/10.1534/genetics.116.198424

Table 1. Sampling site, sample size (n), geographic coordinates and elevation, population

summary statistics: number of private alleles, expected and observed heterozygosity (H_E and

H_O), inbreeding coefficient (G_{IS}) and nucleotide diversity (π)

Sampling site	n	Latitude (°)	Longitude (°)	Elevation (m)	Allelic richness (A_R)	Private alleles	H_O	H_E	G_{IS}	π
1-Guilhaumard	15	43.85	3.19	726	17043.95	0	0.262	0.282	0.071	0.282
2-Lapanouse-de-Cernon	15	43.99	3.09	651	16833.53	2	0.260	0.276	0.059	0.276
3-Saint-Affrique	13	43.98	2.93	585	16669.70	1	0.268	0.269	0.006	0.269
4-Valgañón	15	42.32	-3.08	1023	17167.45	24	0.268	0.291	0.081	0.290
5-Larraona	15	42.78	-2.28	812	17297.27	9	0.269	0.291	0.076	0.290
6-Bercedo	13	43.09	-3.44	774	17102.52	13	0.264	0.283	0.066	0.281

	86											

Table 2. Summary of the main demographic parameters inferred with DILS i) for pooled dataset (*Ophrys aveyronensis* subsp. *aveyronensis* and <u>*Ophrys aveyronensis*</u> subsp. *vitorica*) and ii) for each *Ophrys aveyronensis* subsp. *aveyronensis* and *Ophrys aveyronensis* subsp. *vitorica* population pairs. N_e 1, N_e 2, N_e anc. correspond to effective population sizes estimated for population 1, population 2 and the ancestral population, t_{split} to the time of split between population 1 and population 2 in number of generations, respectively and M_{12}/M_{21} the migration rates (as N_e.m) from population 1 to population 2 and from population 2 to population 1. Parameters have been estimated based on a model using neural network.

Population 1	Population 2	Ne 1	Ne 2	Ne anc.	tsplit	M_{12}/M_{21}
O. a. *aveyronensis* pool (France)	O. a. *vitorica* pool (Spain)	6474 (5737-7276)	8425 (7433-9693)	74803 (66810-83607)	1453 (1271-1642)	0.63/0.56 (0.53-0.73/0.44-0.70)
1-Guilhaumard	4-Valgañón	4934 (4290-5662)	4105 (3622-4589)	98474 (92517-105003)	1226 (1079-1383)	0.90/0.74 (0.73-1.07/0.60-0.88
1-Guilhaumard	5-Larraona	2403 (2089-2751)	2968 (2569-3429)	103206 (97489-109542)	651 (573-724)	0.89/0.64 (0.72-1.06/0.52-0.76)
1-Guilhaumard	6-Bercedo	9824 (8553-11446)	10477 (7866-13078)	97808 (93302-103547)	3029 (2672-3410)	0.73 /0.44 (0.35-0.53/(0.60-0.86)
2-Lapanouse	4-Valgañón	2729 (2506-3035)	2640 (2311-2953)	99913 (94450-105541)	803 (728-872)	0.46/0.67 (0.38-0.55/0.54-0.79)
2-Lapanouse	5-Larraona	4061 (3556-4554)	6049 (5291-6972)	93345 (89784-97211)	1416 (1255-1564)	0.42/1.01 (0.34-0.50/0.71-1.28)
2-Lapanouse	6-Bercedo	5812 (4988-6822)	7719 (6431-9262)	97785 (90460-105961)	2069 (1731-2410)	0.84/0.74 (0.67-1.00/0.58-0.93)
3-St-Affrique	4-Valgañón	9995 (8557-11455)	13510 (11837-15468)	94633 (89865-100052)	3879 (3236-4549)	0.89/0.81 (0.71-1.05/0.62-0.99)
3-St-Affrique	5-Larraona	7270 (6386-8230)	10997 (9603-12667)	90012 (86305-93995)	2891 (2494-3300)	0.82/0.79 (0.66-0.97/0.56-0.98)

| 3-St-Affrique | 6-Bercedo | 9562 (8393-10964) | 10997 (9632-12459) | 99195 (92748-104706) | 3682 (3184-4204) | 0.95 /0.89 (0.77-1.14-0.69-1.08) |

883 **Table 3.** List of bioclimatic variables available from WorldClim. From these 19 variables (bio1

884 to bio19) the ones used for ENM analyses (Person correlation coefficient < 0.75) are

885 indicated in bold with their relative contribution in the selected ENM model.

Variable name	Denomination	Relative contribution (in %)
bio1	Annual Mean Temperature	
bio2	**Mean Diurnal Range (Mean of monthly (max temp - min temp))**	**8.14**
bio3	**Isothermality (bio2/bio7) (×100)**	**5.32**
bio4	Temperature Seasonality (standard deviation ×100)	
bio5	Max Temperature of Warmest Month	
bio6	Min Temperature of Coldest Month	
bio7	Temperature Annual Range (bio5-bio6)	
bio8	**Mean Temperature of Wettest Quarter**	**3.21**
bio9	**Mean Temperature of Driest Quarter**	**3.01**
bio10	**Mean Temperature of Warmest Quarter**	**11.31**
bio11	Mean Temperature of Coldest Quarter	
bio12	Annual Precipitation	
bio13	Precipitation of Wettest Month	
bio14	Precipitation of Driest Month	
bio15	**Precipitation Seasonality (Coefficient of Variation)**	**41.65**
bio16	Precipitation of Wettest Quarter	
bio17	**Precipitation of Driest Quarter**	**23.34**
bio18	Precipitation of Warmest Quarter	
bio19	Precipitation of Coldest Quarter	

Figure 1 Map displaying the 6 sites sampled: 3 for *Ophrys aveyronensis* subsp. *aveyronensis*:

1-Guilhaumard, 2-Lapanouse-de-Cernon, 3-St-Affrique (in blue) and 3 for *O. a.* subsp.

vitorica: 4- Valgañón, 5-Larraona, 6-Bercedo (in red). The approximate known geographic

distribution is highlighted for both taxa with the same color code.

Figure 2 A) Principal Component Analysis (PCA) displaying the two first axes (PC1 and PC2)

representing 8.50% and 3.88% of the total genetic variance. PCA was computed based on 86

individuals genotyped at 9301 SNP and colors depict sampling localities. B) Maximum

Likelihood phylogenetic tree (rooted at midpoint) obtained from the concatenation of the

SNPs (node support for main nodes are given as UltraFast Boostrap Approximation based on

1000 replicates). C) Barplots of ancestry coefficients obtained from sNMF for 86 individuals

for $K = 2$ and $K = 5$.

Figure 3 Demographic history of the 6 studied populations of *O. a.* subsp. *aveyronensis*

(shades of blue) and *O. a.* subsp. *vitorica* (shades of red) depicting the mean effective

population size N_e (and associated 95 confidence intervals) over time as inferred from

STAIRWAYPLOTv2, assuming a mutation rate of 7×10^{-9} and a generation time of 5 years.

Figure 4 Schematic overview of the best demographic scenario (Isolation with Migration, IM)

for pooled dataset (*i.e. O. aveyronensis* subsp. *aveyronensis/ O. a.* subsp. *vitorica*). The

median values(and 2.5-97.5% boundaries of Highest Posterior Distribution probability) are

shown for the parameters inferred with DILS. N_a: ancestral effective population size, N_1:

(current) effective population size for the *O. a.* subsp. *aveyronensis* population (given in

number of individuals), N_2: (current) effective population size for the *O. a.* subsp. *vitorica*,

M_{12}: migration rate from *O. a.* subsp. *aveyronensis* to *O. a.* subsp. *vitorica* and M_{21}: migration

rate from *O. a.* subsp. *vitorica* to *O. a.* subsp. *aveyronensis*. Alternative demographic

scenarios tested are also illustrated: Strict Isolation (SI), Ancestral Migration (AM, involving a

cessation of gene flow after an initial stage of migration after the split) and Secondary

Contact (SC, involving a secondary contact after an initial stage of isolation after the split).

Figure 5 Spatial projections of the bioclimatic niches inferred for the *Ophrys aveyronensis*

species complex depicting climate suitability at different time periods from the Last Glacial

Maximum (LGM, about 21 000 kyrs ago) and with projections for the Heinrich Stadial1 (17.0-

14.7 kyrs BP), the Bølling-Allerød (14.7-12.9 kyrs BP) the Younger Dryas Stadial (12.9-11.7

kyrs BP) and the Holocene: early-Holocene, Greenlandian (11.7-8.326 kyrs BP), mid-

Holocene, Northgrippian (8.326-4.2 kyrs BP) and late-Holocene, Meghalayan (4.2-0.3 kyrs

BP) and currently.

IM

t_{split}
1453 gen.
(1271-1642)

$N_{anc.}$
74803 ind.
(66810-83607)

M_{12}
0.63 *N.m*
(0.53-0.73)

M_{21}
0.56 *N.m*
(0.44-0.70)

N_1
6474 ind. (5737-7276)

N_2
8425 ind. (7433-9693)

SI

AM

SC