Zooarchaeological investigation of the Hoabinhian exploitation of reptiles and amphibians in Thailand and Cambodia with a focus on the Yellow-Headed tortoise (*Indotestudo elongata* (Blyth, 1854))

Corentin Bochaton[1,2*], Sirikanya Chantasri[3], Melada Maneechote[4], Julien Claude[2,5], Christophe Griggo[6], Wilailuck Naksri[7], Hubert Forestier[8], Heng Sophady[9], Prasit Auertrakulvit[4], Jutinach Bowonsachoti[3], and Valéry Zeitoun[1]

[1]UMR 7207. CR2P-CNRS-MNHN-SU, Sorbonne Université, campus Jussieu, T. 46-56, 5ème étage, case 104, 4, place Jussieu, 75 252 Paris Cedex 05, France.
[2]Institut des sciences de l'évolution de Montpellier, UMR 5554 CNRS/UM/IRD/EPHE, Université de Montpellier, Montpellier, France.
[3]12th Regional Office of Fine Art Department, Nakhon Si Thammarat, Thailand.
[4]Department of Archaeology, Faculty of Archaeology, Silpakorn University, Bangkok, Thailand.
[5]Faculty of Science, University of Chulalongkorn, Bangkok, Thailand.
[6]Laboratoire Environnements, DYnamiques et Territoires de Montagne, UMR 5204 – Université Grenoble Alpes, Le Bourget-du-Lac, France.
[7]Northeastern Research Institute of Petrified Wood and Mineral Resources, Nakhon Ratchasima Rajabhat University, Nakhon Ratchasima, Thailand.
[8]Muséum national d'Histoire naturelle, Musée de l'Homme, UMR 7194 HNHP, Paris, France.
[9]Ministry of Culture and Fine Arts of Cambodia, Phnom Penh, Cambodia.
*Corresponding author: corentin.bochaton@cnrs.fr

ABSTRACT

Although non-marine turtles are nearly ubiquitous in the Southeast Asian archaeological record, their zooarchaeological study has been very poorly undertaken in that tropical region of the world. This lack of study makes the understanding of past human subsistence strategies very complex especially regarding the prehistoric hunter gatherer populations which may have massively exploited inland chelonian taxa. In order to try to start a new dynamic regarding the study of the past human-turtle interactions in Southeast Asia we propose here an in-depth zooarchaeological analysis of the turtle bone remains recovered

from four Hoabinhian Hunter-gatherer archaeological assemblages located in Thailand and Cambodia, and dated from the Late Pleistocene to the first half of the Holocene. Our study is focused on the bone remains attributed to the Yellow-Headed Tortoise (*Indotestudo elongata*) as they account for the majority of the turtle archaeological assemblages identified in the target area. For this species, we developed osteo-metric equations enabling the estimation of the carapace size of the archaeological individuals of this species. This allowed us to study the size structure of the archaeological populations in the different sites and to reveal the human exploitation strategies of these animals. We found a strong taphonomic homogeneity between the studied assemblages suggesting similarities of the subsistence behaviors in the different sites despite their very different environmental settings. We thus hypothesize putative cultural similarities across time and space. In addition, we also provide a baseline for future zooarchaeological studies as well as a methodological frame for the detailed studies of archaeological turtle bones in continental Southeast Asia.

Keywords: Reptiles, Hunter-gatherer, Southeast Asia, Turtle, Zooarchaeology

                                              Introduction

The Hoabinhian has been a major topic in prehistoric research in Mainland Southeast Asia for
nearly 90 years. Since its first definition by the French archaeologist Madeleine Colani in the early 1930s
(Collectif 1932), the Hoabinhian has no doubt been one of the most debated topics of the field. A wide
range of subjects pertaining to the Hoabinhian populations have been discussed, such as their
distribution in space and time, their definition, the technological and functional characteristics of their
lithic industries, their economic organization, and their environmental context (Forestier *et al.* 2021). The
chronology of the Hoabinhian has been largely extended since the earliest known Hoabinhian sites.
Indeed, the occupation of the Xiaodong Rock shelter in Southwest China has been dated to about 43, 000
BP (Ji *et al.* 2016) and is considered to be the earliest occurrence of the Hoabinhian culture while the site
of Huai Hin in Northwest Thailand, dated to about 3700 BP is considered to be the last occurrence with a
lithic production associated with ceramic sherds (Zeitoun *et al.* 2008; Forestier *et al.* 2013). More than a
hundred Hoabinhian sites have been reported in Southeast Asia (Moser 2001; Chung 2008; Zeitoun *et al.*
2008; White 2011; Forestier *et al.* 2017) leading inevitably to the question of the variability of Hoabinhian
lithic assemblages with unexpected operational sequences on pebble matrix (Forestier *et al.* 2022, 2023).
Despite this, the Hoabinhian people remain, however, quite poorly understood from cultural and
material point of views. In fact, the homogeneity and lack of diversity of their lithic material culture,
probably related to their putative heavy use of objects made of perishable vegetal material (Forestier
2003) does not allow to characterize the precise uses of the sites by past populations, and it is difficult to
expect a cultural stasis over 30,000 years among different hunter-gatherer groups in such an extensive
region presenting diverse environmental, ecological and geographic settings.
White (2011) proposed that the emergence of cultural diversity in mainland Southeast Asia
began precisely in the late Late Pleistocene which is also suggested by burial practices (Imdirakphol *et al.*
2017). Forestier et al. (2013) argued that more analyses were still needed to evaluate the whole corpus of
Southeast Asian lithic industries to describe putative "cultural variations" and since then some patterns
started to emerge (Forestier *et al.* 2017, 2021). However, while a technological definition of the
Hoabinhian culture is in progress, the lack of detailed zooarchaeological analyses makes our knowledge
of the Hoabinhian paleoecology and subsistence strategies variability blurry. This is also an issue for the
characterization of the archaeological deposits as faunal data are of major importance to characterize the
use and occupation periods of the archaeological sites. The economic aspects of the Hoabinhian culture
were previously addressed by several authors (Gorman 1969, 1970, 1971; Glover 1977; Yen 1977; Vu
1994). Yet, the pioneering prehistoric zooarchaeological studies relying on occurrence data (Gorman
1971) offer only a poor understanding of the choices made by the hunters by focusing on qualitative data
and the diversity of the exploited animals rather than on quantitative information reflecting the intensity
of the exploitation of each species. These first works also lack the detailed taphonomic and taxonomic
analyses needed to describe past human behaviors and bone accumulation processes in more depth.
Because of these lacks, we currently have only a vague idea of the spatial and chronological variability of
the subsistence strategies of the Hoabinhian people. These problems impacting southeast Asian
zooarchaeology have previously been reviewed in depth by Conrad (2015). Among the identified issues is
the lack of detailed study of each animal group but especially of the non-mammal taxa including reptiles
and non-marine turtles that are often accounting for a large part of the animal bone assemblages found
in the archaeological record.
This limit is not proper to continental Southeast Asia but is more impactful here than in many
other areas. Zooarchaeological studies fully focused on non-marine turtles have been conducted in non-
tropical areas as Europe (Blasco 2008; Nabais & Zilhão 2019; Nabais *et al.* 2019), Near East (Speth &
Tchernov 2002; Blasco *et al.* 2016; Biton *et al.* 2017), South Africa (Avery *et al.* 2004; Thompson &

Henshilwood 2014), and Northern America (Rhodin 1992). Such studies remained, however, limited in tropical areas including in South-East Asia despite the fact that turtle bones are way more common in tropical settings than in the temperate regions where there is a strong tradition of prehistoric studies. In Southeast Asia, this problem is partly rooted in the facts that there is a general lack of detailed anatomical data allowing for the identification of the taxa on the basis of isolated plate remains, and that appropriate methodological framework were never developed to analyze this material. Several works have been conducted regarding the osteology of Southeast Asian turtles most of which focusing on the Geoemydidae family in order to address questions related to the phylogeny and the paleo-biodiversity of the group (Naksri 2007, 2013; Naksri *et al.* 2013; Garbin *et al.* 2018). However, only few works have been interested in the study of the isolated elements found in the archaeological record (Pritchard *et al.* 2009; Claude *et al.* 2019). Despite these limitations several zooarchaeological studies of Hoabinhian archaeological deposits in mainland Southeast Asia have started to characterize the exploitation of the non-marine turtles by these prehistoric populations. In the few existing studies, turtle remains are often left unidentified: Ban Rai Rockshelter (Treerayapiwat 2005); Banyan Valley Cave (Higham 1977); Gua Gunung Runtuh (Zuraina 1994); Gua Harimau (Bulbeck 2003); Gue Kechil (Dunn 1964; Medway 1969); Gua Ngaum (Bulbeck 2003); Gua Peraling (Adi 2000); Gua Teluk Kelawar (Bulbeck 2003); Tham Lod Rockshelter (Amphansri 2011); Lang Kamnan Cave (Shoocongdej 1996); Khao Toh Chong Rockshelter (Van Vlack 2014); Moh Khiew II Rockshelter (Auetrakulvit 2004); Spirit Cave (Higham 1977); Tham Phaa Can (Higham 2002); Thung Nong Nien Rockshelter (Auetrakulvit 2004). In some studies they are identified but not quantified by species for instance in the Lang Rongrien Rock Shelter assemblage (Anderson 1990; Mudar & Anderson 2007). In the rare studies in which turtle bones are identified and quantified, the most abundant species is often by far the Yellow-Headed tortoise (*Indotestudo elongata* (Blyth, 1854)) -Doi Pha Kan Rockshelter (Frère *et al.* 2018); Laang Spean Cave (Frère *et al.* 2018); Spirit Cave (Conrad *et al.* 2016)- at least in its modern distribution area as Geoemydidae turtles are the best represented in the Malaysian sites Gua Sagu (Rabett 2012), and Gua Tenggek (Rabett 2012). There is also a site in Vietnam (Hiem Cave) where the tortoise *Manouria* is the most abundant turtle taxa but the small size of the faunal assemblage does not allow drawing conclusions from this observation (Masojc *et al.* 2023). However, even in the quantified above mentioned studies, the study of the turtle bone remains is still superficial. For instance, a detailed analysis of the taphonomy of the bone assemblages is not conducted and the population of turtle exploited is not characterized further than from the aspect of its species composition.

To start proposing solutions to these issues and also to provide the first detailed data regarding the prehistoric exploitation of Southeast Asian turtles we conducted an in-depth zooarchaeological analysis of the turtle bone remains recovered from four hunter-gatherer archaeological assemblages. These sites are located in Thailand and Cambodia and dated from the Late Pleistocene to the first half of the Holocene. They are the Doi Pha Kan rockshelter, the Moh Khiew Cave, and the Khao Tha Phlai Cave located in Thailand, and the Laang Spean Cave located in Cambodia. In addition, in order to be able to characterize with more precision the exploitation strategies of the non-marine turtles by the archaeological human populations, we developed osteo-metric equations enabling to estimate the carapace size of the archaeological individuals of *Indotestudo elongata*. We choose to focus our methodological approach on this species as most of the rich assemblages of turtle bones collected in the four considered sites correspond to this species (Frère *et al.* 2018; C. B., J. C. preliminary observations). This analytical tool allows for the study of the size structures of the archaeological populations in the different sites, and the characterization the choices made by the hunters. Altogether these data provide the first characterization of the exploitation of non-marine turtle by Pleistocene and Holocene hunter-gatherer populations of continental Southeast Asia.

Material and Methods

Main characteristics of *Indotestudo elongata*, the Yellow-Headed tortoise

The genus *Indotestudo* currently includes three species: the Forsten's Tortoise *Indotestudo*
forstenii (Schlegel & Müller, 1845), the Travancore Tortoise *Indotestudo travancorica* (Boulenger, 1907),
and the Yellow-Headed tortoise *Indotestudo elongata* (Blyth, 1854), the two later being sister taxa
(Iverson *et al.* 2001). These species are currently distributed in India (*I. travancorica*), Sulawesi (*I.*
forstenii), and northern India and continental Southeast Asia (*I. elongata*) (Rhodin *et al.* 2021). This last
species is the only *Indotestudo* species present in continental Southeast Asia. It is nowadays present in
most areas of Thailand, Cambodia, Vietnam, and Laos (Ihlow *et al.* 2016; Rhodin *et al.* 2021). It is also
occurring in northwestern Malaysia but is absent from most of this country and from the insular Sunda
(Ihlow et al., 2016). *I. elongata* is a medium size tortoise whose adult Straight Carapace Length (SCL)
reaches about 300 mm (Taylor 1970). The largest specimen ever recorded is a male of about 380 mm SCL
(Rhodin *et al.* 2021). It is possible that the sexual dimorphism in this species vary from one population to
another although males are generally larger than females (Ihlow *et al.* 2016). The size of the hatchlings
ranges from 50 to 55 mm SCL (Ihlow & Handschuh 2011). The specie is present in a wide variety of
environments including many forest types (Ihlow et al., 2016). *I. elongata* is active during daytime mostly
in the early morning and late afternoon. It present seasonal activity patterns in order to avoid the highest
temperatures during the dry season during which it aestivate spending most of its time hiding in former
burrows of other animal species including porcupines (Van Dijk 1998; Som & Cottet 2016; Ihlow *et al.*
2016). In their active period the individuals spend most of their time in open areas during the rainy
season then move in more closed environment (mostly semi-evergreen and pine forests) when the
climate becomes dryer (Van Dijk 1998; Som & Cottet 2016). Depending of their sex, the individuals reach
sexual maturity between 175 mm and 240 mm SCL at an age of 6-8 years old (Van Dijk 1998; Eberling
2011; Sriprateep *et al.* 2013). Reproduction takes place during the rainy season.

Presentation of the studied sites and assemblages

The Doi Pha Kan rockshelter

The site of Doi Pha Kan is a rockshelter located in northern Thailand (E 99° 46' 37.2'' ; N 18° 26'
57.0''). The site is studied since 2011 by P. Auetrakulvit and V. Zeitoun and its excavation is still in
progress. The site is mostly known for its three Hoabinhian sepultures dated between 11,170 ± 40 and
12,930 ± 50 BP (Imdirakphol *et al.* 2017; Zeitoun *et al.* 2019). Two of these sepultures contained turtle
shells elements in anatomical connection interpreted as funeral offerings. The site also provided a rich
archaeological assemblage corresponding to a Hoabinhian occupation older than the sepultures. The
stratigraphy of the site being homogeneous from a sedimentary point of view, its archaeological material
has been so far considered as a single assemblage.

Samples of the lithic material and of the animal bone assemblages collected on the site have
already been the object of studies (Celiberti *et al.* 2018; Frère *et al.* 2018) but most of the material is still
under study. In this paper we will present zooarchaeological data collected on the herpetofaunal taxa
bone remains recovered on the site until 2019. To collect this dataset the whole archaeofaunal material
has been observed to extract and then study the reptile and amphibian bone remains. The material
screened this way corresponds to the material previously studied by S. Frère (Frère *et al.* 2018) with the
addition of the material collected following this first study. In the first study, S. Frère analyzed 4256
animal remains among which 2541 were attributed to vertebrate species. So far, no complete study of
this faunal assemblage has been completed and no Minimal Number of Individual (MNI) data have ever
been published. In these conditions, the overall weight of the herpetofauna we collected in respect to the
full sample cannot be assessed with precision. However, the existing data indicates it could account for a
very significant part of the full assemblage as it corresponds to 17.1% of the total bone weight and 51% of

the vertebrate total Number of Indentified Skeletal Parts (NISP) analyzed in the previous study (Frère *et*
al. 2018).

The Moh Khiew Cave

The site of Moh Khiew Cave is a 30m long archaeological rock-shelter located in southern
Thailand in the Krabi Province (E 98° 55' 49.27''; N 08° 09' 36.32''). It was first excavated by S. Pookajorn
between 1991 and 1994 (Pookajorn 2001) before being the object of a later excavation in 2008 by P.
Auetrakulvit in order to clarify its stratigraphy (Auetrakulvit *et al.* 2012). The stratigraphy of the site is
composed of several archaeological layers dated from the Holocene to the Late Pleistocene, most of
which corresponding to Hoabinhian occupations. Five sepultures were also discovered, one during the
first excavations, and four in 2008.

Regarding the zooarchaeological data, the complete assemblages of the first excavations was
studied by P. Auetrakulvit (Auetrakulvit 2004). In this assemblage, MNI data indicate that herpetofaunal
species account for 24.9% of the assemblage. This proportion dramatically increases to more than 70% of
the material if the NISP is considered with nearly 60% for the non-marine turtles bone alone.
Unfortunately, the turtle remains were not identified further at the moment of this first study and we
were not able to locate this material in order to study it again in the framework of the present analysis.
We however had access to the material collected during the 2008 excavation but only to previously
studied herpetofaunal bone samples that were extracted from the complete bone samples by several
master students. These bones were recovered from the four different layers identified during the 2008
excavation of Moh Khiew Cave (Auetrakulvit et al., 2012). The first layer (Layer 1) is composed of the
upper first 90 cm of the sequence. It is a disturbed layer that has not been dated and which corresponds
to the levels 1 and 2 identified by S. Pookajorn in the first excavations. The second layers include the
sediment collected between 90 and 170 cm of depth. It has been the object of three radiocarbon dates
between 7520+-420 BP. and 8660+-480 BP. This layer corresponds to the level 3 identified in the previous
excavations. The third layer correspond to a depth between 170 and 210 cm and has been dated with
two radiocarbon dates of 8730+-480 BP. and 9270+-510 BP. It was also identified as corresponding to the
level 3 previously described by S. Pookajorn. The last layer corresponds to the remaining of the
stratigraphy. It is a layer of scree and was not dated but associated to the level 4 described in the
previous excavations.

The Khao Tha Phlai cave

The site of Khao Tha Phlai is a cave located in southern Thailand in the province of Chumphon (E
10°36'12.39"; N 99° 5'49.08"). It has been excavated by the 12[th] Regional Office of Fine Arts Department,
Nakhon Si Thammarat since 2014, and has so far been the object of two excavation campaigns with two
test-pits conducted in the deposit. A first test-pit of 9 m² (TP1) in 2014, and a second of 20 m² (TP2) in
2021-2022.

The archaeofaunal material collected in the TP1 has been the object of a preliminary study by S.
Jeawkok in the framework of a Bachelor thesis that remained unpublished. This first study conducted on
6945 bone remains indicated that the reptiles represent around 25% of the NISP of the complete
assemblage, the remaining bones being nearly all attributed to large mammals. In the present study we
considered the herpetofaunal material previously extracted by S. Jeawkok in the faunal samples collected
in the TP1. We also consulted the full archaeofaunal sample recovered from the TP2 in order to extract
the reptile and amphibian bones from it. We considered separately the samples collected in the two test-
pits and subdivided the samples in two assemblages corresponding to Metal Ages (between 75 and 130
cm of depth in the TP2 and between 65 and 180 cm in TP1), and Neolithic periods (between 130 and 320
cm of depth in the TP2 and between 180 and 320 in TP1). These layers have been dated on the basis of
the typology of the archaeological artifacts they have provided.

The Laang Spean cave

The site of Laang Spean is a large cave of more than 1000 m² located in northwest Cambodia, in
the Battambang province (E 102° 51' 00.0''; N 12°51' 00.0''). The site has been the object of a first
excavation between 1965 and 1968 by R. Mourer and C. Mourer-Chauviré (Mourer-Chauviré *et al.* 1970;
Mourer-Chauviré & Mourer 1970). The archaeofaunal material collected during these excavations has
been the object of a preliminary study but has never been studied in depth at the exception of the
rhinoceros remains (Guerin & Mourer-Chauviré 1969), and no zooarchaeological study have been
conducted. Following this first exploration, the site has been the object of a new detailed archaeological
excavation by H. Forestier between 2009 and 2019 aiming to document in more detail the Hoabinhian
occupation previously identified (Forestier *et al.* 2015; Sophady *et al.* 2016). These excavations conducted
on a surface of over 40 m² led to the discovery of an important undisturbed Hoabinhian layer dated
between 5018 ± 29 cal. BP and 10 042 ± 43 cal. BP as well as several Neolithic burials dug in the
Hoabinhian level (Zeitoun *et al.* 2012, 2021). These sepultures have been dated from 3335 ± 30 to 2960 ±
30 BP (Sophady 2016). Regarding the subdivision of the material, the archaeological remains collected in
the squares lacking traces of Neolithic perturbations (see Forestier et al., 2015) have been associated to
the Hoabinhian occupation. There is no stratigraphic evidence suggesting a subdivision of this Hoabinhian
assemblage obviously representing several occupations over a time span of more than 5000 years.
Regarding the other squares, the squares in which sepultures were found are grouped together under the
term "Sepulture layer" and the first 120 cm of the disturbed squares are considered as being a "Neolithic
layer". These subdivisions are however very artificial and probably correspond to a mix of Neolithic and
Hoabinhian material has it has been evidenced on the lithic material these contexts provided (H.
Forestier, com. pers.).

A sample of the complete faunal assemblage collected in the Hoabinhian squares has been the
object of a first zooarchaeological study by S. Frère (Frère *et al.* 2018). In this first study, among the 5885
vertebrate remains identified, turtle account for 44% of the NISP, monitor lizards for 1%, and large
mammals for 37%. Unfortunately no MNI data were reported and the fact that most of the small
fragments were not attributed to a least a size class of animal make these results difficult to interpret.
The study presented in this paper corresponds to the herpetofaunal material collected in the complete
assemblage of bones recovered during all the excavations since 2009. This material has been extracted by
C. B. upon the consultation of all the bone samples collected on the site. The zooarchaeological study of
the other groups of vertebrate for the complete Hoabinhian assemblage of Laang Spean is currently in
progress and the final results are not available at the moment.

Quantification of the zooarchaeological data

The basic units of quantification considered are the Number of Indentified Skeletal Parts (NISP)
and the Weight of the Remains (WR). The fragmentation of each bone has been recorded by describing
the Percentage of Completion (PC) of the anatomical elements represented by the fragment. The
laterality of the bones has been registered when possible for the best represented and easiest to identify
anatomical elements (i. e. peripheral plates, and all paired elements of the plastron). A Minimal Number
of Elements (MNE) has been compute for each anatomical part in order to study the anatomical
distribution of the remains in the different archaeological contexts. To do so, we have added the PCs of a
given element and divided the result by 100. The results were rounded to the superior unit to obtain the
MNEs. The Minimal Number of Individual (MNI) is defined using the anatomical element with the highest
MNE. The anatomical distributions are represented by the Percentage of Representation (PR) of Dodson
and Wexlar (1979) using the MNE of each anatomical elements and the MNI of the considered
assemblage. All the archaeological bones have been weighted individually. In order to avoid a potential
impact of taxonomic identification bias on the anatomical distribution of the remains we considered all

the turtle/tortoises taxa in these analyses and not only the bone fragments attributed to *Indotestudo*. As
tortoises are by far the best represent taxa in the different assemblage most of the unidentified turtle
bones likely represents *Indotestudo*. The positions of the peripheral plates have been identified but only
for *Indotestudo* remains. The peripheral plates for which it was not possible to give a position have been
posteriorly assigned to the different ranks following the distribution of those for which a position was
determined. Regarding the size estimations (see below), the mean of the obtained size estimation is
considered in the case several measurements were recorded on a single bone. Chi² tests were performed
on the Microsoft Excel software 2007 version.

Size estimation of archaeological Indotestudo

In order to reliably estimate the body size of the archaeological individuals of *Indotestudo* sp. we
build size estimation equations on the models of what was previously done for Southeast Asian monitor
lizards (Bochaton *et al.* 2019), and recently on the size and weight of species of tortoises (Esker *et al.*
2019; Codron *et al.* 2022). These approaches are more powerful than considering isolated measurement
(e. g. Klein and Cruz-Uribe 1983) to describe the size of subfossil animal populations because: 1) they
enable to take account on several measurements from different anatomical parts to reconstruct the body
size structure of a past population, and 2) it converts measurements taken on the skeleton into a variable
used to describe the size of modern individuals which make easier the comparisons needed to address
biological questions. To build the equations we defined a set of 86 measurements (See Supplementary
Table 1; Appendix 1) that we recorded on a sample of 34 museum specimens of *Indotestudo* sp. from the
Florida Museum of Natural History (UF), and the Comparative Anatomy collection of the Muséum
national d'Histoire naturelle (MNHN-ZA-AC) (see all details in Supplementary Table 1). In order to have
enough specimens to produce relevant and reliable predictive equations we pooled altogether the three
species currently includes in the genus *Indotestudo*: *I. forstenii* (Schlegel & Müller, 1845): 8 specimens, *I.
travancorica* (Boulenger, 1907): 14 specimens, and *I. elongata* (Blyth, 1854): 12 specimens. Differences of
body proportions among species were controlled before applying this strategy to avoid including bias
related to interspecific differences in the size estimations.

The measurements recorded were distributed on all the bones of the plastron and the carapace
as well as on the long bones. Vertebrae and skull elements were not taken into account as their
occurrences were too rare in the archaeological record. In addition to these measurements, in order to
be able to choose a variable accounting for the "body size" of each individual, we took three
measurements considered as "size variables" on the complete carapace of the modern specimens: the
Carapace Straight Length (CSL), the Shell Height (SH), and the Plastron Length (PL). All the measurements
collected on modern and archaeological specimens were recorded using a digital dial calliper [IP 67
(Mitutoyo Corporation, Japan)]. All the measurements recorded are included in the Supplementary Table
1. All statistical analyses were performed using the basic library Stats of the open-source software R (R
Core Team 2020). Each size estimation equation produced is the result of a linear regression between a
given log transformed measurement recorded on a bone/plate and the log transformed "size variable" of
the specimens. The variables are log transformed in order to make linear the simple allometric
relationship between the used variables (Huxley 1932; Gould 1966). Consequently, the obtained CSL
estimation has to be log reversed using an exponential function to be obtained in the same unit as the
used measurements. Obtained equations are of the form:

Log ("size variable") = (Beta1)* log (osteological measurement in mm) + (Beta0)

From this initial set of equations we choose to discard all the equation that were not significant
(p.value above 0.01) and/or with a low coefficient of determination (R2) (below 0.85) in order to keep
only the best equations to estimate the size of the archaeological individuals.

Specific identification of the *I. elongata* archaeological bone sample

The archaeological bones attributed to *I. elongata* have been identified on the basis of a direct comparison with pictures of the skeletal specimens of this species used to build the SCL estimation equations. The fact that the family of the Testudinidae is only represented by very few species in Southeast Asia (*Geochelone platynota*, *Indotestudo elongata*, *Manouria emys*, and *Manouria impressa*) (Das 2010) presenting different sizes, morphologies, and distribution areas allowed to attribute with relative ease nearly all of the studied Testudinidae remains to *I. elongata*. Among the *Indotestudo* genus, as the only currently available qualitative diagnostic criteria for *I. elongata* is located on the nuchal plate (presence of a long and narrow cervical scute), we based most of our identification obtained from other plates/bones on the exact similarity between the archaeological bone remains and the morphologies present on the modern specimens of different ages we observed. An overview of the carapace morphologies of juvenile and adult *I. elongata* is provided here (Fig. 1). The remains attributed to this species also present the morphological traits common to most Testudinidae: a carapace lacking lateral keels, a costal pattern of odd costal plates with short distal end and long medial end, and even costal plates with long distal end and short medial end, octagonal and squared neural plates, peripheral plates without musk ducts, a costo-marginal sulcus superimposed to the costo-peripheral suture, a pygal plate not intersected by the posterior sulcus of the fifth vertebral scute, thickened epiplastra, and thin and vertical inguinal and axillary buttresses. Among Testudinidae the genus *Indotestudo* is characterized by the fact that the humeropectoral sulcus is crossing the entoplastron (Auffenberg 1974). The establishment of robust and quantified diagnostic criteria for the identification of isolated bones of Southeast Asian turtles has still to be performed. As a comment, we signal that the characteristic nuchal scute morphology of *I. elongata* was present on all the nuchal plate (N=109) attributed to this species at the exception of a single remain from Laang Spean cave. It has been previously signaled that the nuchal scute could be absent on some specimens (Ihlow *et al.* 2016) and our data indicate a frequency of such feature inferior to 1%.

Figure 1 - A) Drawing of the carapace of an adult specimen of Indotestudo elongata (CUMZ-R-
TT181); B) Drawing of the plastron of an adult specimen of Indotestudo elongata (CUMZ-R-TT181);
C) Drawing of the carapace of a juvenile specimen of Indotestudo elongata lacking peripherals (UF-
34760); D) Drawing of the plastron of a juvenile specimen of Indotestudo elongata showing central
fontanel (UF-34760). **Abbreviations:** A. s.: Abdominal scute, An. s.: Anal scute, G. s.: Gular scute,
Co.: Costal plate, F. s.: Femoral scute, Ep.: Epiplastron, En.: Entoplastron, F.: plastral fontanel, H. s.:
Humeral scute, Hyo.: Hyoplastron, Hyp. : Hypoplastron, M. s.: Marginal scute, Ne.: Neural plate, Nu.
p.: Nuchal plate, C. s.: Cervical scute, P. s.: Pectoral scute, Pe.: Peripheral plate, Pl. s.: Pleural scute,
Py.: Pygal plate, S.p. 1: Supra-pygal 1, S.p. 2: Supra-pygal 2, S. s.: Supra-caudal scute, V. s.: Vertebral
scute, Xi.: Xiphiplastron.

368   **Results**

369 **Size predictive equations**

370   In order to choose the variable we will use as our "size" variable we tested the correlations
371 between the three "body size" measurements taken on the complete carapaces of our *Indotestudo* spp.
372 modern specimens. Our results indicate that the Straight Carapace Length (SCL) is strongly correlated to
373 the Plastron Length (PL) (R^2=0.97), and that the Shell Height is more weakly correlated to the two other
374 measurements (R^2=0.93 and 0.92). This could be related to sex specific or interspecific differences
375 regarding the height of the carapace among the considered specimens but these hypotheses are not
376 possible to test considering the small size of our sample. As a result and because the carapace length is
377 the most used variable to describe the size of a turtle we choose the SCL as our size scalar but the PL
378 could have been equally considered.

379   From our complete modern sample, a set of 86 equations corresponding to the 86 initial
380 recorded measurements has been produced. This set has been refined to keep only the equations
381 providing the most reliable (significant linear relationships) and precise (high R^2 linear relationships) SCL
382 estimations. This final set of equations include 52 equations from 52 measurements distributed on
383 epiplastron, entoplastron, hyoplastron, hypoplastron, xiphiplastron, nuchal plate, neural plates (ranks 1,
384 2, 3, 4, 6, and 7), peripheral plates (ranks 1, 2, 3, 8, 9, and 10), 2nd supra-pygal plate, pygal plate,
385 humerus, radius, ulna, femur, tibia, and fibula (Fig. 2; Table. 1).

386
387    **Table 1** - Equations retained for the prediction of the Straight Carapace Length (SCL) of the
388    archaeological Indotestudo specimens. Are indicated: the measurement used (to be recorded on
389    the archaeological specimen, the slope X ("Beta1" to integrate in the equation indicated in the
390    Material and Method section), and the intercept Y ("Beta0" to integrate in the equation indicated in
391    the Material and Method section). Is also indicated the coefficient of determination (R2) of each
392    relation, the p.values, and the degree of freedom of each linear regression.

Measurement Used	Y	X	R²	P. value	Freedom degree
Greatest Length Nuchal plate (GL)	1.44	1.05	0.89	2.48E-14	27
Greatest Heigth Peripheral plate 1 (GH)	2.30	0.91	0.89	2.77E-14	27
Greatest Width Peripheral plate 1 (GW)	1.95	1.01	0.90	3.43E-15	27
Greatest Heigth Peripheral plate 2 (GH)	1.68	1.11	0.92	4.10E-16	26
Greatest Width Peripheral plate 3 (GW)	1.96	1.06	0.88	5.70E-14	27
Greatest Heigth Peripheral plate 3 (GH)	2.42	0.84	0.93	2.20E-16	27
Greatest Width Peripheral plate 8 (GW)	2.45	0.89	0.86	5.22E-13	27
Greatest Heigth Peripheral plate 8 (GH)	2.19	0.91	0.94	2.20E-16	27
Greatest Width Peripheral plate 9 (GW)	2.58	0.86	0.89	1.47E-14	27
Greatest Heigth Peripheral plate 9 (GH)	2.27	0.89	0.90	7.29E-15	27
Greatest Width Peripheral plate 10 (GW)	2.26	0.95	0.89	3.04E-14	27
Greatest Heigth Peripheral plate 10 (GH)	2.69	0.77	0.90	3.55E-15	27
Greatest Length Neural plate 2 (GL)	2.54	0.91	0.87	2.82E-13	27
Greatest Width Neural plate 2 (GW)	1.55	1.14	0.92	3.36E-16	27
Greatest Length Neural plate 3 (GL)	2.20	1.07	0.90	4.73E-15	27
Greatest Length Neural plate 4 (GL)	2.52	0.93	0.85	1.09E-12	27
Greatest Width Neural plate 4 (GW)	1.82	1.05	0.92	2.20E-16	27
Greatest Width Neural plate 6 (GW)	2.10	0.98	0.91	6.00E-16	27
Greatest Width Neural plate 7 (GW)	2.23	0.96	0.86	3.95E-13	27
Greatest Width Supra-pygal plate 2 (GW)	2.19	0.86	0.88	1.61E-13	26
Greatest Heigth Pygal plate (GH)	2.98	0.68	0.93	2.20E-16	26
Greatest Length Epiplastron (GL)	1.48	1.06	0.95	2.20E-16	27
Greatest Length Entoplastron (GL)	1.54	1.09	0.86	3.09E-13	27
Greatest Width Entoplastron (GW)	1.76	1.02	0.92	4.25E-16	27
Greatest medial Length Hyoplatron (GmL)	1.81	0.96	0.87	1.14E-13	27
Greatest Width Hyoplastron (GW)	0.02	1.29	0.94	2.20E-16	25
Greatest medial Length Hypoplastron (GmL)	0.44	1.25	0.90	7.32E-15	27
Greatest Width Hypoplastron (GW)	-0.09	1.32	0.95	2.20E-16	25
Greatest Length Xiphiplastron (GL)	1.53	1.02	0.88	6.69E-14	27
Greatest Width Xiphiplastron (GW)	1.09	1.15	0.92	2.20E-16	27
Greatest Length Humerus (GL)	1.16	1.08	0.95	2.20E-16	24
Greatest proximal antero-posterior Width Humerus (GpapW)	2.74	0.99	0.91	3.32E-14	24
Greatest proximal dorso-ventral Width Humerus (GpdvW)	2.59	1.00	0.94	8.45E-16	24
Greatest Length Ulna (GL)	1.52	1.12	0.96	2.20E-16	23
Greatest distal antero-posterior Width Ulna (GdapW)	3.60	0.93	0.90	4.69E-13	26
Greatest proximal latero-medial Width Ulna (GplmW)	3.13	1.02	0.87	1.22E-11	23
Greatest Length Radius (GL)	1.34	1.17	0.94	1.82E-15	23
Greatest proximal latero-medial Width Radius (GplmW)	3.18	1.10	0.92	3.64E-14	23
Greatest proximal antero-posterior Width Radius (GpapW)	3.96	0.89	0.89	1.20E-12	23
Greatest distal antero-posterior Width Radius (GdapW)	3.91	0.99	0.86	2.08E-11	23
Greatest Length Femur (GL)	1.41	1.03	0.96	2.20E-16	23
Greatest proximal antero-posterior Width Femur (GpapW)	3.15	0.86	0.93	1.89E-14	23
Greatest proximal dorso-ventral Width Femur (GpdvW)	2.91	0.90	0.97	2.20E-16	23
Greatest distal antero-posterior Width Femur (GdapW)	3.46	0.77	0.86	2.91E-11	23
Greatest Length Fibula (GL)	1.57	1.06	0.94	1.81E-14	21
Greatest distal latero-medial Width Fibula (GdlmW)	3.59	0.94	0.89	1.06E-11	21
Greatest distal antero-posterior Width Fibula (GdapW)	3.92	0.96	0.90	9.30E-12	21
Greatest Length Tibia (GL)	1.39	1.09	0.95	1.25E-15	22
Greatest proximal latero-medial Width Tibia (GplmW)	3.06	1.01	0.92	2.16E-13	22
Greatest proximal antero-posterior Width Tibia (GpapW)	3.20	1.02	0.94	3.03E-15	22
Greatest distal latero-medial Width Tibia (GdlmW)	3.45	1.00	0.92	1.99E-13	22
Greatest distal antero-posterior Width Tibia (GdapW)	3.60	0.97	0.91	9.04E-13	22

395
Figure 2 - Measurements corresponding to the 52 equations retained to predict the SCL of our
archaeological sample of Indotestudo elongata bone remains. Measurements names: GB: Greatest
Width (on the latero-medial axis), GdapW: Greatest distal antero-posterior Width, GddvW: Greatest
distal dorso-ventral Width, GdlvW: Greatest distal latero-ventral Width, GH: Greatest Height (on the
dorso-ventral axis), GL: Greatest Length (on the antero-posterior axis), GmL: Greatest medial
Length(on the antero-posterior axis), GpapW: Greatest proximal antero-posterior Width, GpdvW:
Greatest proximal dorso-ventral Width, GplvW: Greatest proximal latero-ventral Width.

403
404 **Zooarchaeological and taphonomic analyses of the herpetofaunal assemblages**

Figure 3 - Examples of the turtle bone remains studied. A) Plastron of an adult *I. elongata* in anatomical connection from the Hoabinhian layer of Laang Spean Cave (ventral view); B) Plastron of a juvenile *I. elongata* in anatomical connection from the Hoabinhian layer of Laang Spean Cave (ventral view); C) Nuchal plate of *I. elongata* from the Hoabinhian layer of Laang Spean Cave (dorsal view); D) Pygal plate of a juvenile *I. elongata* from the Neolithic layer of Laang Spean Cave (posterior view); E) Pygal plate of an adult *I. elongata* from the Neolithic layer of Laang Spean Cave (posterior view); F) Left peripheral plates (1st to 3rd) of *I. elongata* in anatomical connection from the layer 2 of Moh Khiew Cave presenting burning traces limited to the internal side of the carapace; G) Left peripheral plates (8th to 11th) of a Geoemydidae in anatomical connection whose ventral part has been cut-down from the site of Doi Pha Kan.

Doi Pha Kan Rockshelter

<u>Composition of the herpetofaunal assemblages</u>

The herpetofaunal assemblage of Doi Pha Kan consist of 8414 bone remains weighting a total of
6875 gr. and representing a least 115 individuals. Most of these bones correspond to non-marine turtles
(Fig. 3; Table. 2) in term of WR (74%), NISP (56%), and MNI (47%) with the second most represented
group being Monitor lizards (23% of the WR, 38% of the NISP, and 38% of the MNI) followed by rare
snakes, amphibians, and small lizards bone remains.

Table 2 - Weight of the Remains (WR), Number of Identified Skeletal Parts (NISP), and Minimal
Number of Individual (MNI) corresponding to the different taxa identified in the complete Doi Pha
Kan Rock-shelter herpetofaunal assemblage.

	NISP	WR	NMI
Turtle/Tortoise	4762	5113.02	55
Monitor lizard	3203	1610.39	44
Snake	375	141	3
Amphibian	67	10.57	10
Small lizards	7	0.82	3
Total	**8414**	**6875.8**	**115**

428
429

Regarding the bones attributed to turtles and tortoises (Table. 3), 66.8% of them (50.8% of the
WR) could not be attributed to a given family. Among the fragments that were identified to at least the
family level, 71 % of them were attributed to *Indotestudo elongata* (71% of the WR and 76% of the MNI),
28% (28% of the WR and 18% of the MNI) to Geoemydidae, and less than 1% (0.6% of the WR and 5% of
the MNI) to Trionychidae. Note that for Trionychids remains from all localities, attribution can be done
for all the plates because of their ornamentation, they are therefore not underrepresented because they
cannot be easily identified.

Table 3 - Number of Identified Skeletal Parts (NISP), Weight of the Remains (WR), and Minimal
Number of Individual (MNI) corresponding to the different turtle/tortoise taxa identified in the
complete Doi Pha Kan Rock-shelter assemblage.

	NISP	WR	NMI
Indotestudo elongata	1122	1787.17	42
Geoemydidae	447	711.34	10
Trionychidae	12	14.13	3
Turtle ind.	3181	2600.38	
Total	**4762**	**5113.02**	**55**

441
442

Taphonomy of the turtle/tortoise bone assemblage
Among the 4762 bone fragments attributed to turtle/tortoises, 303 are complete elements
(6.3%), and 249 are nearly complete (at least 90% of the bone is preserved) while 1389 (29%) are small
fragments representing less than 5% of the complete anatomical part. The average percentage of
completion of the bones is 32%. The overall PR is 28%. The best represented bones (Fig. 4 – A) are the
stylopods (humerus and femur with PR>75%) followed by the tibia, the radius, the scapula, the coracoid,
and the epiplastron (PR>50%). Then follow the rest of the long bones, and most of the plates that are the
easiest to identify. The peripheral plates of the bridge are less represented (PR=10%) than all the other
peripheral plates (PR=39%). The skull, vertebrae and all small elements of the hands and foot are nearly
absent. The largest plates (hyoplastron and hypoplastron) are the most fragmented with completion
means of less than 26%. The peripheral plates corresponding to the bridge are also more heavily

fragmented (completion mean of 59%) compare to the other peripheral plates (completions means 67-89%). Burning (black) and carbonization (grey/white) traces were observed on 690 bones (14% of the total NISP). Such traces were present indiscriminately on every anatomical element of the carapace and skeleton. Cut marks were observed on only five bones, six peripheral plates, and one hypoplastron. A series of peripheral plates attributed to a Geoemydidae turtle bears clear traces of a clean cut aiming to cut the ventral part of the carapace (Fig. 3-G). Lastly, 120 bones have been observed as corresponding to preserved anatomical connections on the field between unfused plates. This indicates that at least a part of the assemblage was undisturbed prior to the excavation.

Figure 4 - Anatomical distributions of the turtle/tortoise remains collected in the sites of Doi Pha Kan rockshelter (A), and Moh Khiew Cave (B-E). The percentage of representation (PR) is considered here to provide a graphical visualization of the different values observed for the different anatomical elements.

 Size of *Indotestudo elongata* archaeological individuals

469   The measurements recorded on the *I. elongata* archaeological material of Doi Pha Kan enabled
470 for the reconstruction of 201 SCL estimations included between 64 and 292 mm with a mean of 182 mm
471 (Figure. 5-A) and corresponding to at least 42 individuals. The distribution of these sizes was not
472 unimodal (Hartigans' dip test, p.val>0.05) but could be bi-modal with a population of small individuals
473 around 120 mm, and a second one between 140 and 260 mm.
474

Figure 5 - Histograms of the *Indotestudo elongata* size reconstructions (Standard Carapace Length)
obtained from the different herpetofaunal bone assemblages studied: A) Doi Pha Kan Rock-shelter
(NMI=42); B) Moh Khiew Cave (NMI=59); C) Laang Spean Cave (NMI=75); D) Khao Ta Phlai Cave
(NMI=26). The black bars represent the minimal size of the sexually mature specimens based on
modern data collected on modern *I. elongata* populations.

Moh Khiew cave
Composition of the herpetofaunal assemblages

The herpetofaunal assemblage of Moh Khiew consists in 9 108 bone remains weighting 8351 gr.
Those bones are mostly distributed in the layers 2 (51% of the NISP, and 52% of the WR), 1 (26% of the
NISP, and 24% of the WR), and 3 (17% of the NISP, and 18% of the WR) (Table. 4). The complete
assemblage includes bone fragment from a least 152 individuals.
The full sample mostly correspond to non-marine turtle remains (63% of the NISP, 74% of the WR, and
52% of the MNI), followed by Monitor lizards (25% of the NISP, 16% of the WR, and 24% of the MNI), and
snakes (8% of the NISP, 8% of the WR, and 4% of the MNI). The small size lizards (excluding snakes),
amphibians and crocodile remains altogether represent less than 5% of the assemblage in term of NISP
and WR. The distribution of these groups among the layers is however far from homogenous. Indeed,

turtle bone remains represent more than 75% of the NISP, 85% of the WR, and 60% of the MNI in the layers 2 to 4 but only 20% of the NISP, 35% of the WR, and 21% of the MNI in the layer 1. At the opposite, Monitor lizards represent 46% of the NISP, 31% of the WR, and 35% of the MNI in the layer 1 but less than 20% of NISP, 13% of the WR, and 25% of the MNI in the other layers. Snakes are also better represented in the layer 1 (23% of the NISP) than in the subsequent levels (less than 2.6% of the NISP). Chi² tests performed on the NISP indicate that the faunal composition of the layer 1 significantly differs (P.val< 0.01) from the layers 2 and 3, the effective of the layer 4 being too low to conduct a statistical test.

Table 4 - Number of Identified Skeletal Parts (NISP), Weight of the remains (WR), and Minimal Number of Individuals (MNI) identified in the complete herpetofaunal assemblage of the different layers of the 2008 excavation of Moh Khiew Cave.

	Layer 1			Layer 2			Layer 3			Layer 4			Total		
	NISP	WR	NMI	NISP	WR	NMI	NISP	WR	NMI	NISP	WR	NMI	NISP	WR	NMI
Turtle/Tortoise	503	702.8	6	3614	3789	53	1184	1288	15	439	481.3	5	**5740**	**6261**	**79**
Monitor lizard	1111	641.4	17	872	463.8	12	298	190.1	5	43	25.4	2	**2324**	**1321**	**36**
Snake	560	612.2	3	117	52.8	1	37	29.15	1	7	6.4	1	**721**	**700.6**	**6**
Amphibian	144	38.45	14	34	6.5	4	22	3.92	3	0	0	0	**200**	**48.87**	**21**
Small lizards	89	8.27	7	4	0.3	1	28	3.1	1	**0**	**0**	**0**	**121**	**11.67**	**9**
Crocodile	2	8.5	1	0	0	0	0	0	0	0	0	0	**2**	**8.5**	**1**
Total	**2409**	**2012**	**48**	**4641**	**4312**	**71**	**1569**	**1514**	**25**	**489**	**513.1**	**8**	**9108**	**8351**	**152**

Regarding the identification of non-marine turtle taxa (Table 5), between 66 and 79% of the bone fragment have not been associated with at least a family rank identification. Regarding the identified families, Testudinidae (*Indotestudo elongata*) represent a stable major part of the identified bone remains (between 68 and 63% of the NISP, 70-56% of the WR, and 86-50% of the MNI depending of the layer). The other fragments have been attributed to Geoemydidae turtles (36-29% of the NISP, 36-28% of the WR, and 33-12% of the MNI), and Trionychidae (0-2% of the NR and WR). The composition of the turtle assemblages in term of families seems then to be fairly stable across the layers as indicated by results of Chi² tests that did not indicated significant differences in the distribution of the NISP from the different testable layers (P.val> 0.01).

Table 5 - Number of Identified Skeletal Parts (NISP), Weight of the remains (WR), and Minimal Number of Individuals (MNI) in the non-marine turtle assemblage from the different layers of the 2008 excavation of Moh Khiew Cave.

	Layer 1			Layer 2			Layer 3			Layer 4			Total		
	NISP	WR	NMI	NISP	WR	NMI	NISP	WR	NMI	NISP	WR	NMI	NISP	WR	NMI
Indotestudo elongata	109	239.9	4	853	1274	43	193	323.9	10	60	105	2	**1215**	**1943**	**59**
Geoemydidae	61	107.3	2	386	512.9	6	111	253.9	3	26	62.2	1	**584**	**936**	**12**
Trionychidae	1	0.7	0 (1)	16	37.8	1	0	0	1	2	2.5	1	**19**	**41**	**3**
Turtle ind.	332	354.9	0	2359	1964	3	880	709.9	1	351	311.6	1	**3922**	**3340**	**5**
Total	**503**	**703**	**6**	**3614**	**3789**	**53**	**1184**	**1288**	**15**	**439**	**481**	**5**	**5740**	**6261**	**79**

Taphonomy of the turtle/tortoise bone assemblage

The fragmentation of the 5740 bone fragments analyzed increase with the depth. In the layer 1, 8.9% of the bones are complete, and 12.9% nearly complete, in layer 2 these percentages falls to 7.5 and 12.1%, then to 5.6 and 9.6% in layer 3, and finally to 3.9 and 7.7% in layer 4. The average percentage of completeness of the bones is also slightly higher in layers 1 and 2 (36 and 37%) than in the layers 3 and 4 (34 and 28%). These differences are only significant between the layers 1 and 4 (Chi² test; p. value<0.01).The anatomical distribution of the remains present strong variations between the layers (Fig.

4) but the sizes of the assemblages are also very dissimilar with the bone samples of the layers 1 and 4
containing only 503 and 439 remains and those of the layers 2 and 3 containing 3614 and 1184 bone
fragments. Distributions in small samples could be more strongly impacted by random effects than larger
assemblages and a strict comparison of the four layers might not make sense at all. Some general trends
can, however, still be noted just as the overall PR which is between 27 and 24% in all the layers at the
exception of the first in which it is slightly higher (36%). Nearly all the anatomical parts are present in
every layer but the skulls, vertebrae, and extremities are nearly absent. The stylopods (humerus and
femur) are the best represented bones in the richest layers. They are also well represented in layers 1
and 2 but are outnumbered by some specific carapace plates. Girdles and zeugopods are also present but
in smaller number. Regarding the carapace and the plastron, no clear pattern emerges at the exception
of the nearly systematic lower representation of the peripheral plates of the bridge (PR=25-21%) in
respect to the other peripheral plates (mean PR=52-29%). This could be explained by an identification
bias related to the lower mean completion rate of peripheral plates of the bridge in respect to the others.
Burning (black) and carbonization (grey/white) traces were observed on 325 bones (5.7% of the total
NISP). Such traces were present indiscriminately on every anatomical element and every side of the
carapace and skeleton parts. They were recorded on the internal side of several peripheral plates that
were still in anatomic connection at the moment of the excavation (Fig. 3-F). Such observations were,
however, not repeated on the rest of the material. Cut marks were observed on only three bones: one
peripheral plate, one nuchal plate, and one xiphiplastron. Among the full assemblage 61 fragments of
carapaces were still in anatomical connection at the moment of the excavation. These elements were
distributed mostly in the layers 2 and 3 but also in the lower part of the layer 1 at a depth of 70-80 cm.
549
Size of _Indotestudo elongata_ individuals
The measurements recorded on the _I. elongata_ archaeological material of Moh Khiew Cave
enabled for the reconstruction of 201 SCL estimations included between 98 and 310 mm and with a mean
of 193 mm (Figure. 5-B) and corresponding to at least 59 individuals. The mean size of the tortoises is
similar in all layers and no statistically significant differences was noted (student T-test; p.val>0.01). In
layer 1 (N=25) the mean size was 198 mm, 194 mm in layer 2 (N=169), 187 mm in layer 3 (N=52), and
finally 195 mm in layer 4 (N=19). The largest observed specimen was in the layer 1. The global
distribution of these sizes was not unimodal (Hartigans' dip test, p.val>0.01) and mixture models indicate
it is most likely bimodal with a best represent group of individuals around 220 mm and a second group of
smaller specimens around 150 mm.
560
Khao Ta Phlai Cave

Taxonomic composition
A total of 3763 bone remains of herpetofauna weighting 7239 gr. and representing at least 43
individuals were analyzed from the two excavated test-pits of the site of Khao Ta Phlai (Table. 6). Most of
these bones correspond to turtle or tortoises in term of WR (87% in TP1 and 90% in TP2), NISP (81% in
TP1 and 86% in TP2), and MNI (65% in TP1 and 70% in TP2). The second most represented herpetofaunal
group is the Monitor lizards (12 and 9% of the WR, 16 and 12% of the NISP, and 23 and 22% of the MNI)
followed by some rare remains of snakes as well as few amphibian bones in the TP2 only.
569
570   **Table 6** - Number of Identified Skeletal Parts (NISP), Weight of the Remains (WR), Number of
571   Remains (NR), and Minimal Number of Individual (MNI) corresponding to the different
572   herpetofaunal taxa identified in the herpetofaunal bone assemblages collected in the two test-pits
573   of the site of Khao Ta Phlai.

	TP1			TP2		
	NISP	**WR**	**NMI**	**NISP**	**WR**	**NMI**
Amphibians				3	0.51	1
Snakes	38	33.4	2	37	69.3	2
Turtle/Tortoise	1335	2509.7	19	1825	3901	18
Monitor lizards	274	347.7	6	251	377.9	6
Total	**1647**	**2891**	**27**	**2116**	**4349**	**27**

Although the distribution of the material between the two TPs is somewhat homogenous, the repartition of the bones across the two main periods documented (metal ages and Neolithic) is quite different. Indeed, in the TP2, most of the material (95% of the NISP) is located in the Neolithic layers while in TP1 the bone are more evenly distributed (55% of the NISP in the Metal Ages layers and 44% of the NISP in the Neolithic levels).

The distribution of the main herpetofaunal taxa across the layers does not present strong variations (Tab. 7) as only the very poor metal age layer of the TP2 significantly differ from the other levels (Chi²; p.value <0.01). Turtle bones are always the best represented (between 71 and 87% of the NISP) but Monitor lizards seems a bit better represented in Metal Ages layers from TP1 and 2 (18 and 20% compared to Neolithic layers (14 and 11% of the NISP). These tendencies are, however, not statistically significant and difficult to interpret in the absence of a study of the complete faunal assemblages.

Table 7 - Number of Identified Skeletal Parts (NISP), Weight of the Remains (WR), and Minimal Number of Individual (MNI) corresponding to the different herpetofaunal taxa identified in the herpetofaunal bone assemblages of the two periods identified in the two test-pits of the site of Khao Ta Phlai.

	TP1-Metal Age			**TP1-Neolithic**			**TP2-Metal Age**			**TP2-Neolithic**		
	NISP	**WR**	**NMI**	**NISP**	**WR**	**NMI**	**NISP**	**WR**	**NMI**	**NISP**	**WR**	**NMI**
Amphibians										3	0.51	1
Snakes	33	22.4	2	5	11	2	9	13.4	2	28	55.9	2
Turtle/Tortoise	720	1011.8	14	615	1497.8	9	70	131.2	2	1755	3769.8	21
Monitor lizards	166	146.2	1	108	201.5	1	20	36.2	1	231	341.7	3
Total	**919**	**1180**	**17**	**728**	**1710**	**12**	**99**	**180.8**	**5**	**2017**	**4168**	**27**

The identification rate of non-marine turtles bones was lower in the TP1 (41% of the WR and 25% of the NR) than in TP2 (57% of the WR and 35% of the NR). Regarding the bones attributed to a given family (Tab. 8), TP1 provided nearly as much Testudinidae as Geoemydidae in terms of WR and NISP while *Indotestudo elongata* is much more represented than the latter in the TP 2 (61% of the WR and 69% of the NISP). These differences are statistically significant (Chi² test; p.value<0.01). Trionychidae are present in the two TPs. If the chronological phases are considered (Tab. 9), Geoemydidae and Trionychidae are significantly better represented in the upper layers of the TP2 and TP1 corresponding to the metal ages. These two layers do not significantly differ in term of family composition (Chi² test; p.value>0.01) but significantly differ from the two Neolithic layers (Chi² test; p.value<0.01). This trend to a more important exploitation of freshwater turtles during the metal ages in regard to the Neolithic period is for now difficult to interpret considering the possible issues of chronological associations between the layers of the two TPs and the possibility of a spatial variation in the distribution of the remains inside the site. Indeed freshwater turtles are also better represented in the Neolithic layer of the TP1 compare to TP2.

Table 8 - Number of Identified Skeletal Parts (NISP), Weight of the Remains (WR), and Minimal Number of Individual (MNI) corresponding to the different turtle/tortoise taxa identified in the bone assemblages collected in the two test-pits of the site of Khao Ta Phlai.

	TP1			TP2		
	NISP	**WR**	**NMI**	**NISP**	**WR**	**NMI**
Indotestudo elongata	175	435.5	8	441	1357	14
Geoemydidae	150	581.9	5	191	838.6	3
Trionychidae	10	14.1	1	9	32.2	1
Turtle ind.	1000	1478		1184	1673	
Total	**1335**	**2510**	**14**	**1825**	**3901**	**18**

Table 9 - Number of Identified Skeletal Parts (NISP), Weight of the Remains (WR), and Minimal Number of Individual (MNI) corresponding to the different turtle/tortoise taxa identified in the bone assemblages collected in the different chronological phases of the two test-pits of the site of Khao Ta Phlai.

	TP1-MA			TP1-NE			TP2-MA			TP2-NE		
	NISP	**WR**	**NMI**	**NISP**	**WR**	**NMI**	**NISP**	**WR**	**NMI**	**NISP**	**WR**	**NMI**
Indotestudo elongata	86	149	7	89	286.5	6	14	40.2	1	427	1316.6	12
Geoemydidae	81	243.2	2	69	338.7	3	13	28.7	1	178	809.9	4
Trionychidae	10	14.1	1				3	5.2	1	6	27	1
Turtle ind.	543	605		457	872.63		70	57.1		1144	1616.3	
Total	**720**	**1011**	**10**	**615**	**1498**	**9**	**100**	**131.2**	**3**	**1755**	**3770**	**17**

Taphonomy of the turtle/tortoise bone assemblage

Among the 3 160 bone fragments attributed to turtle/tortoises in the material of Khao Ta Phlai, 221 were complete elements (6.9%), and 370 were nearly complete (at least 90% of the complete bone) while 760 (24%) were small fragments representing less than 5% of the complete anatomical part. The average percentage of completion of the bones is 29%. This value is similar in the Neolithic layers of TP1 and TP2 (29%). It is slightly lower in the metal ages layer of the TP1 (25%), and higher in the same period layers from TP2 (39%) but the small size of this assemblage does not allow to consider this result as significant.

Figure 6 - Anatomical distributions of the non-marine turtle remains collected in the different test-pits and layers of the site of Khao Ta Phlai: (A) Test-pit 1 – Metal Ages, (B) Test-pit 1– Neolithic, (C), Test-pit 2– Neolithic. The percentage of representation (PR) is considered here to provide a graphical visualization of the different values observed for the different anatomical elements.

The anatomical distributions of the turtle bone elements present strong variations among the different layers (Fig. 6). The distribution observed in the Neolithic layers of the TP2 is fairly homogenous (mean PR=33%) with a representation of all the anatomical part at the exception of the smallest elements (phalanges, carpal and tarsal articulations, and vertebrae), and the skull. The most robust anatomical parts are the best represented (peripheral plates, epiplastron, entoplastron, and nuchal plate -PR>38%-) and the most fragile the least represented (zeugopods, and most girdles elements - PR<5%-). The only exception to that pattern is the peripheral plates of the bridge (PR=15%) which are least represented than the other elements of the carapace and other peripherals (mean PR=73%). The distribution pattern is quite less homogenous in TP1 where the mean PR is lower (24% in the Neolithic layers and 15% in the metal ages layers) but the Neolithic layers follow the same general pattern as the one of the TP2 with a lower global representation due to the strong presence of a single peripheral plate rank. In this last layer, another difference with TP2 is that the stylopods are also better represented than the elements from the carapace. The anatomical distribution of the bones collected in the metal ages layers of TP1 is very different with a very strong representation of the stylopods (mean PR=76%) compare to the most robust elements of the carapace (PR=29-15%). Otherwise, the same general observations applies with a lack of skull and extremities elements, a better representation of the most robust elements of the carapace, and a lower representation of the peripheral plates of the bridge (PR=5%) in regard of other peripheral plates (Mean PR=18%).

Most of the material (72% of the NISP) was covered by a veil of calcite which made very difficult the observation of the surface alterations of the bones. Despite this important limitation, nine bones

were recorded as presenting traces of dissolution under the effect of flowing water, and 182 as bearing
traces of burning and carbonization. During the study we also recorded 22 associations of bones from the
same context being in anatomical connection in the two TPs below 120 cm of depth in the TP 2 and 135
cm in the TP 1. This indicates that the material from the deepest layers was not strongly disturbed since
its deposition. Nine combination of bone in anatomical connection linked together by concretion were
also found in the same layers.
660
<u>Size of *Indotestudo elongata* individuals</u>
The measurements recorded on the *I. elongata* archaeological material of the Khao Ta Phlai site
have enabled for the reconstruction of 219 SCL estimations included between 108 and 252 mm and with
a mean of 184 mm (Figure. 5-D) and corresponding to at least 26 individuals. Most of the size estimations
are from the Neolithic layers of the TP2 (n=158) and TP1 (n=31) while only 29 estimations were obtained
from the metal ages layers (mean size = 178 mm). The strong disparities of the distributions of the size
estimations between the archaeological contexts do not allow for an individual comparison of the
different layers. The Neolithic layers provided mean SCL values of 185 and 186 mm. The global
distribution of these sizes was not unimodal (Hartigans' dip test, p.val>0.01) and mixture models indicate
it is most likely bimodal with a group of individuals around 210 mm and a second group of smaller
specimens around 165 mm.
672
673 *Laang Spean Cave*
674
<u>Composition of the herpetofaunal assemblage</u>
The complete herpetofaunal assemblage of Laang Spean consists of 9533 bone fragments
weighting 18 804 gr. and representing at least 115 individuals (Tab. 10). Most of them come from the
Hoabinhian layer accounting for 76% of the NISP, 78% of the WR, and 62% of the MNI. The "Neolithic"
and "sepulture" assemblages are of similar sizes and account respectively for 10.5 and 13.2% of the NR.
The material corresponds nearly exclusively to non-marine turtle remains which account for 92% of the
NISP, 95% of the WR, and 70% of the MNI. Monitor lizards represent 5% of the NISP of the assemblage,
snakes are rare (2% of the NISP), and the occurrence of amphibians and smaller lizards is insignificant
(below 1%). The distribution of the taxa is not statistically different across the sub-assemblages (Chi² test;
p.value>0.01).
684
685   **Table 10** - Number of Identified Skeletal Parts (NISP), Weight of the remains (WR), and Minimal
686   Number of Individuals (MNI) studied in the complete herpetofaunal assemblage from the different
687   layers of the Laang Spean Cave.

	Hoabinhian layer			Neolithic layer			Sepulture layers			Total		
	NISP	WR	NMI	NISP	WR	NMI	NISP	WR	NMI	NISP	WR	NMI
Turtle/Tortoise	6650	13893	56	904	1854	12	1195	2057	12	**8749**	**17804**	**80**
Monitor lizard	406	497	3	30	48	2	39	56	3	**475**	**601**	**8**
Snake	158	311	3	16	37	1	19	30	2	**193**	**378**	**6**
Amphibian	45	9	7	33	6	5	6	1	2	**84**	**16**	**14**
Small lizards	13	2	3	16	2	3	3	1	1	**32**	**5**	**7**
Total	**7272**	**14712**	**72**	**999**	**1947**	**23**	**1262**	**2145**	**20**	**9533**	**18804**	**115**

688
689
Regarding the identifications of the turtle and tortoise bone fragments (Tab. 11), only 33% of the
NISP and 52% of the WR have been attributed at least to a family. *Indotestudo elongata* bone remains
account for most of the identified turtle/tortoise bones with 86% of the NISP, 86% of the WR, and 87% of

the MNI. Geoemydidae are rare with only 14% of the NISP, 13% of the WR, and 9% of the MNI, while the

occurrence of Trionychidae is anecdotic (less than 1% of the NR and WR, and 3.5% of the MNI).

Table 11 - Number of Identified Skeletal Parts (NISP), Weight of the remains (WR), and Minimal
Number of Individuals (MNI) identified in the turtle/tortoise bones assemblage from the different
layers of Laang Spean Cave.

	Hoabinhian layer			Neolithic layer			Sepulture layers			Total		
	NISP	WR	NMI	NISP	WR	NMI	NISP	WR	NMI	NISP	WR	NMI
Indotestudo elongata	2046	6706	51	204	649	12	260	692	12	2510	8047	75
Geoemydidae	325	955	4	34	115	2	47	162	2	406	1232	8
Trionychidae	7	20.5	1	2	7	1	5	16	1	14	43.5	3
Turtle ind.	4272	6212		664	1084		883	1187		5819	8483	0
Total	**6650**	**13894**	**56**	**904**	**1855**	**15**	**1195**	**2057**	**15**	**8749**	**17806**	**86**

<u>Taphonomy of the turtle/tortoise bone assemblage</u>

Regarding the taphonomy of the turtle bones collected in the different squares, the mean

completion rate is slightly lower in the "sepulture" assemblage (32%) than in the "Neolithic" and

"Hoabinhian" assemblages (39.2 and 36.7%). The general fragmentation pattern is otherwise similar in all

layers. The complete bones constitute between 9.3% of the assemblages for the Hoabinhian assemblage,

and 7.2-6.2% for the "Neolithic" and "Sepulture" assemblages while nearly complete elements account

for 14.2% of the "Hoabinhian", 13.8% of the "Neolithic", and 11% of the "Sepulture" assemblages. Most

anatomical parts are represented in the different assemblages but the extremities, vertebrae, and skull

remains are very rare with a PR below 4% in all assemblages (Fig. 7). There is also a global tendency to

the lower representation of the peripheral plates of the bridge (11%-31%) in respect to the other

peripheral plates (58-70%) although this trend is more strongly marked in the Hoabinhian layer (11% vs.

70%). The peripheral plates of the bridge are also systematically more fragmented (59-67% of mean

completion) than the others (83-87% of mean completion). Outside of these common trends, significant

strong differences emerge between the "Hoabinhian" assemblages and the two other layers. Indeed,

although the general PR is similar in the different assemblages (41% for the Hoabinhian assemblage, 37%

for the Neolithic assemblage, and 34% for the Sepulture assemblage) the carapace elements are

dramatically better represented in the Hoabinhian bones (Fig. 7-A) compare to the two other

assemblages (Fig. 7-B, C) (Chi² test; p.value<0.01). Indeed, although the PR of the carapace and plastron

bones are more or less similar in the Hoabinhian assemblage (57.8% vs. 67%) the plastron elements are

mostly missing in the other assemblages (54% vs. 20.8% in the Neolithic assemblage and 53% vs. 15.1% in

the Sepulture assemblage). The stylopods are also better represented in the Hoabinhian squares (84%)

compare to the other assemblages (50.4% and 28%).

Figure 7 - Anatomical distributions of the turtle remains collected in the different layers of the site of Lang Spean. The percentage of representation (PR) is considered here to provide a graphical visualization of the different values observed for the different anatomical elements.

The observation of traces on the bones is made very challenging by the fact that 61% of them are covered by a veil of calcite. Interestingly, this calcite deposit was more frequent in the Hoabinhian squares where it covered 65% of the bones but was scarcer in the Neolithic and sepulture squares where it covered respectively 56% and 38% of the bones. This is probably related to the position of the remains in the cave, more or less close to the walls, which influenced their exposition to water flows during the rainy season. There is also a possibility that the calcite veil might be more frequent on the oldest remains. The presence of water flow in the site is also indicated by the occurrence of 61 bones having been polished by water flows. Porcupine gnawing traces were observed on only 13 elements distributed in several areas and layers of the site and digestion traces on only one. This clearly indicates a minor impact of animal species on the integrity of the archaeological assemblage. Putative burning traces were observed 10% of the remains. These traces were better represented in the Hoabinhian squares where they were present on 12% of the bones while they only occur on 2.8 and 3.8% of the bones recovered in the Neolithic and Sepulture squares. The characterization of burning traces was made very difficult by the fact that the material present a strong variability of surface color probably related to post-depositional chemical alteration. The occurrence of these traces might thus have been underestimated given the fact that we choose to record them only when their nature was undisputable. No cut mark was observed.

Among the full assemblage 327 fragments of carapaces (3.7% of the NR) were still in anatomical connection at the moment of the excavation. These elements are mostly from the Hoabinhian squares (N=292) where they account for 4.3% of the turtle remains. Elements in anatomical connection are scarcer in the other assemblages with only 35 occurrences (1.7% of the turtle NR). This indicates that the Hoabinhian squares have been indeed less disturbed than the "Neolithic" and "Sepulture" squares.

 <u>Size of *Indotestudo elongata* individuals</u>

The measurements recorded on the *I. elongata* archaeological material of Laang Spean Cave
enabled for the reconstruction of 688 SCL estimations included between 68 and 345 mm and with a mean
of 201 mm (Figure. 5-C) and corresponding to at least 75 individuals. Most of the data (N=564) are from
Hoabinhian layers while the Neolithic squares only provided 124 SCL data but no significant difference
emerged from the comparison of these two assemblages (Student t-test, p.val>0.05). The global
distribution (all squares) of these sizes is unimodal (Hartigans' dip test, p.val>0.05) with a peak of
specimens around 200 mm SCL. In this site, small specimens below 170 mm represent only 16% of the
population, and specimens below 140 mm only 4.7%.

756   **Discussion**

Taxonomic composition of the herpetofaunal assemblages

In all the assemblages the distribution of the herpetofaunal groups in the four investigated sites
present strong similarities. Non-marine turtles are nearly systematically the best represented
herpetofaunal group (between 59 and 91% of the NISP) before Monitor lizards (between 6 and 25% of
the NISP), snakes (below 3.5% of the NISP), and amphibians. The only exception to this trend is the layer
1 of Moh Khiew cave in which Monitor Lizards (25% of the NISP), and snakes (23% of the NISP) bone
remains are more numerous than turtle skeletal elements (20% of the NISP). This layer is disturbed and
not dated so interpreting this observation is impossible for now. However, the other sites follow a clear
pattern showing that hunter-gatherers groups have preferentially exploited turtles over other reptile and
amphibian taxa. This follow the general regional pattern previously observed in similar zooarchaeological
assemblages (Conrad 2015). Regarding the proportion of the turtle/tortoise families in the assemblages,
Testudinidae (*Indotestudo elongata*) is always the most represented group as it account for between 52
and 89% of the turtle bones NISPs. The proportions of Geoemydidae turtles vary a lot between 48 and
11% of the same NISPs. As most of the species from this group are aquatic freshwater turtles this
variability could be explained by the accessibility of streams, rivers, and lakes by the inhabitants of the
sites. In most sites the proportion of Geoemydidae is around 30% of the NISP but these turtles are much
less prevalent in Laang Spean cave, and the best represented in the TP1 of the site of Khao Ta Phlai. The
published data regarding the faunal assemblage from Laang Spean indicate a weak contribution of
freshwater taxa (mussels and fish) to the overall diet (Forestier *et al.* 2015; Frère *et al.* 2018) which is in
accordance with our observation of the scarcity of freshwater turtles in the site. Regarding the
prevalence of Geoemydidae species in the TP1 of Khao Ta Phlai, this could indicate a stronger reliance on
freshwater resources than in the other sites. However, considering the chronology of the two test-pits of
the site is not yet fully resolved and that general importance of aquatic resources in this assemblage still
need to be estimated, this fact cannot be related to a cultural/chronological trend for now. From a
general point of view, the data regarding the herpetofaunal assemblages, however, point to strong
similarities between assemblages of different ages and from very different environmental settings. This
should be considered at the light of the studies regarding the mammal bone assemblages of the same
sites to test the hypothesis of a putative homogeneity of the Hoabinhian subsistence strategies in
continental Southeast Asia.

Taphonomy of the turtle assemblages

The fragmentation rate of the bones is fairly homogenous among the sites. The average
percentage of completeness of the bones is between 37 and 28%. The material from three first layers of
Moh Khiew Cave and Laang Spean are the less fragmented (average percentage of completeness above
33%). The layer 4 of Moh Khiew cave provided the most fragmented material (average percentage of
completeness of 28%). The presence of large limestone blocks in this layer may be indicative of crumbling

that may have altered the faunal material. Regarding the anatomical distribution of the turtle remains the sites present strong differences with mean PR between 41% (Laang Spean) and 15% (Khao Ta Phlai metal ages layer from TP1). This means that the anatomical representation of the bone remains is more or less biased toward some elements. Two main cases occur in the assemblages: sites in which stylopds are the best represented parts (Khao Ta Phlai metal ages layer from TP1, Doi Pha Kan, layers 2 to 4 of Moh Khiew cave), and sites in which the most robust parts of the carapace are the best represented elements. The mean PR is systematically higher in the assemblages where the stylopds are the most numerous. This indicates that these assemblages are the least altered by post-depositional phenomena. Indeed, a natural alteration would rather lead to the situation observed in the other assemblages in which the elements that are the most often founds are the most robust and thus would have the highest survival rates. However this is not sufficient to explain an overrepresentation of long bones that are supposed to preserve less well than carapaces elements. Considering that all the sediment of the studied deposit has been screened, a major recovery bias is unlikely although some of the smallest elements might have been missed. A post-depositional sorting of the material could also be ruled out as we shown no evidence of differential fragmentation and no abundant trace of water circulation in the different deposits studied. The most likely hypothesis is thus that human inhabitants of some of the sites discarded or transported for further use some of the carapaces of the consumed animals and left on place the smallest elements among which the largest and toughest (humerus and femurs) have been recovered and identified. This behavior would in any case not be systematic as the anatomical distributions indicate that complete individuals have been brought on the sites. The absence of the head of the specimens could be either related to an identification bias or a removal of these parts outside of the site. Humerus and femur put apart, the anatomical distributions of turtle bones follow a global pattern where the most robust anatomical elements are better represented than the more fragile ones. The only exception to this trend concern the peripheral plates of the bridge which are always less represented than the other peripheral. This is very likely to be related to an identification bias itself related to the nearly complete absence of complete pieces of such element in the material. This is undoubtedly linked to the separation of the carapace from the plastron by the inhabitants of the sites who have broken the bones in the area that links both part of the shell, a step that is mandatory to access to most of the meat content of the animal.

The observations of the surface traces on the bones indicate a nearly complete lack of predation and digestion traces which combined with the general weak fragmentation of the material allow to completely ruling out a putative role of non-human predators in the constitution of the studied assemblages. This is not surprising as although some predators, including Monitor lizards, are known to hunt juvenile tortoise individuals, adults' specimens probably have few non-human predators although some modern specimen bear traces of predation attempts (Ihlow *et al.* 2016) and that predation on other Southeast Asian tortoises species have been reported (Platt *et al.* 2021). Large felids (Emmons 1989), and eagles (Gil-Sánchez *et al.* 2022) are known to be able to hunt adults tortoises but such predator would undoubtedly have left predation traces on the subfossil bone assemblages we studied. Some very rare bones bearing porcupine traces indicate that these animals had a minor impact on some of the assemblages but not enough to impact the zooarchaeological interpretations. However, although it seems fairly evident that the animals present in the sites have been hunted to be consumed as there is no trace of bone industry in the assemblages, finding direct traces of culinary preparation on the bones is very challenging. Indeed, in Khao Ta Phlai and Laang Spean, some remains (72 and 37%) were covered by a veil of calcite making impossible to observe the surface of the bones. In addition, very few cut marks have been characterized on the bones of the different sites. Many burned bones were observed in all the sites but linking these to a cooking technique is very questionable. Indeed, these traces do not seem to be located on specific part of the bones (i. e. external side of the carapace) and appear randomly on

every area of every anatomical part. It is likely that these traces are related to post-depositional events
unrelated to the cooking of the animals. The frequency of fire traces combined with the strong
fragmentation of the large vertebrate remains in most sites (C. Griggo; C. Bochaton pers. obs) could
indicate the use of the bones as combustible (Villa *et al.* 2002). Such a use is unlikely for the turtle
skeletons considering their small size and the very good preservation stage of their remains but a
proximity to the fire places (Bennett 1999) could explain the random occurrence of fire traces on their
bone elements.

Size of Indotestudo elongata archaeological specimens
The size of *I. elongata* individuals observed in the four archaeological deposits (Fig. 5) shows
common patterns but also some differences. The distributions of the estimated sizes are bimodal in all
sites except Laang Spean. In all the sites, most of the estimations correspond to adult size specimens
above 170 mm SCL reaching maximums of 270-345 mm SCL. These specimens all fall in the size range of
the modern representatives of the species. Yet, all sites present a variable proportion of smaller, likely
immature individuals. The representation of this second group is the lowest in Laang Spean (16% of the
total number of estimations) but is important enough in the other sites to make their distributions
bimodal with 35%-33% in Doi Pha Kan and Khao Ta Phlai, and 24% in Moh Khiew cave. Specimens below
140mm SCL are rare in all sites as they account for less than 10% of the estimations in all sites but
represent more than 15% of the Doi Pha Kan population.
Interpreting the size distribution of the archaeological tortoises is a difficult task as it first
requires an idea of what the size structure of a wild population would look like and basic biological data
(season of birth, activity pattern, growth speed…) regarding modern and past *I. elongata* populations.
However, these data are currently mostly missing which make very challenging a detailed interpretation
of the collected archaeological data. The recovery of size distribution data in a natural modern population
is always challenging as it could be influenced by many factors (i. e. climate, environment, seasonality,
behaviors, and sizes of the individuals) that could bias the observations by making one or several size
classes more difficult to observe than the others. In addition, the history and specific conditions of a wild
population itself could have a dramatic impact on its size structure. In these conditions, defining a
modern comparison point to the documented archaeological populations is difficult. To our knowledge,
the only data collected on *I. elongata* concern the population of the Ban Kok Village (Khon Kaen Province,
Thailand). This study shows that the pre-adult individuals have a low survivability rate as their population
was mostly composed of newly born and old adult individuals (Sriprateep *et al.* 2013). The authors
suggest that this strongly biased structure could be related to an absence of predation on the large
individuals, and partly to several phenomena having a stronger impact on small specimens (e. g.
predation, trampling of domestic bovids) but admit the main cause is still unknown while not discussing a
potential poaching of the smaller individuals. Similarly, another publication about *I. travancorica*
indicates a lack of juvenile specimen in the population but highlight that it could be related to a seasonal
activity specific pattern. Indeed, juvenile specimens were much more commonly found at the beginning
of the rainy season than during the dry season when their study has been conducted (Ramesh 2008).
Other published distributions from other tortoises' species also indicate a strong representation of adult
size individuals of different ages having completed their growth but also a much more balanced
distribution of juvenile specimens of all sizes (Hailey & Coulson 1999; Znari *et al.* 2005; Rouag *et al.* 2007).
In all these distributions the juvenile specimens are way scarcer than adult ones which make sense as
adult class specimens correspond to individuals of very different ages having reached their final size. The
only case in which this situation would be reversed is a population in which adult individuals would be
subject to a strong predation pressure superior to the pressures imposed on the smaller individuals.

The site of Laang Spean presents a unimodal size distribution in which juvenile specimens are mostly
excluded. In that sense, this distribution is very different from that of a natural population and indicates a
strong selection on adult specimens of moderate to large size. This is clearly indicative of a very selective
hunting strategy that may have been enabled by the abundance of resources in the vicinity of the site.
Such a selection, although visible in other deposits is less marked as juvenile specimens composed more
important parts of the assemblages, especially in Doi Pha Kan. In these sites, it is impossible to estimate
whether or not the proportions of juvenile specimens present in the assemblages are similar to those of
the exploited natural populations and thus to estimate the exact intensity of the selection toward large
size individuals. In any case, it is the sign of an opportunistic foraging as such a combination of juvenile
specimens has been observed on modern hunter-gatherer population actively collecting tortoises this
way (Mena *et al.* 2000). But this might also be influenced by the hunting method in the case a direct
selection by the hunter is not made, for instance with the use of trapping that was also hypothesized in
Doi Pha Kan for the hunting of monitor lizards (Bochaton *et al.* 2019). This technique is also the most
used to hunt tortoise in the Amazon as it is the most efficient method before active searching (Santos *et*
al. 2020). This implies no selection on the specimens in the wild although the type of trap used (i. e. size
of the ground hole) might induce some size bias. The use of traps could thus explain the strong
representation of smaller individuals present in the archaeological assemblages and indicate a very
opportunistic strategy indicative of either a poor selection by the hunter and/or a relative scarcity of the
tortoises in the environments making harder the collect of large individuals. The hunting season could
also be an explanation to the stronger or weaker presence of juvenile specimens in the assemblages.
Indeed, during the dry season tortoises are less active and harder to find which could led the hunter to be
less selective especially in the case of a use of non-selective hunting methods allowing to find these
animals. Theobald (1868) mentions the hunts of tortoises by Burmese hunters in the dry season by
clearing grass lands and forests with fire in order to destroy their shelters and locate them. At the
opposite, smaller tortoises are more active in the rainy season during which dogs are more used to track
them (Blythe 1854; Theobald 1868). Ultimately, both seasonal hypotheses could explain the occurrence
of small individual using different explanations (hunting method vs. activity season). Only the use of non-
traditional approaches such as skeletochronology (Ehret 2007) could help to clarify this question by
estimating the season of death of the tortoise individuals as well as the occupation seasonality of the
different sites given the absence of other seasonality makers in the materials.

Tortoise populations are vulnerable to intensive exploitation focused on the larger mature
individuals. As such, and their exploitation has been considered as a marker of small scale hunting and
thus of small human groups (Stiner *et al.* 2000). In the studied sites, the strong focus on a single turtle
species (*I. elongata*) and the focus placed on large individuals would undoubtedly damage the natural
populations and lead to a size reduction of the individuals (Close & Seigel 1997). Such exploitation would
be sustainable only if it was not intensive meaning that not many individuals were collected in order to
feed a putatively small size human group. It is difficult to address the question of the overall importance
of the tortoises in the diet of the studied Southeast Asian hunter-gatherer groups as we still lack a
complete and quantified study of the mammal fauna of the sites as well as strong data regarding the use
of the studied sites. It is, however, clear that the investigated prehistoric population have exploited
tortoises which composed a significant part of their meat diet. This is not surprising as turtle species are
supposed to represent an important biomass in the ecosystems (Iverson 1982) and are also fairly easy to
collect. This behavior have subsisted until nowadays in continental Southeast Asia (Hansel 2004) although
all population does not choose to exploit reptile species (Tungittiplakornl & Dearden 2002).

928   **Conclusion and perspective**

The present work has been built as a stepping stone to provide the basic data and study tools enabling the study of tortoise's assemblages from continental Southeast Asia. As an effect, the true potential of this work will only be reached through the use of its analytical protocol for future studies and comparison with additional assemblages. We were however able to reach several conclusions as we demonstrated a putative strong similarities between the exploitation of the herpetofaunal taxa in the different sites as well as in the taphonomy of the non-marine turtle assemblages in different chronological and environmental settings. These data thus open many interesting questions regarding the trends of hunter-gatherer subsistence strategies in continental Southeast Asia during the Pleistocene and through the Holocene. However, much work remains to be done to reach a satisfactory zooarchaeological documentation level regarding these prehistoric human groups. As we demonstrate it in the introduction of this paper, many of the previously excavated Hoabinhian archaeological deposits of continental Southeast Asia, including sites which are known to have provided rich assemblages of non-marine turtle bone remains (i. e. Lang Rongrien), have not benefited from quantified zooarchaeological analyses. The complete study of these sites will be important to provide additional relevant comparison points to the present study. The non-herpetofaunal taxa of the sites included in this study should also be investigated to estimate the relative part of the reptile and amphibian exploitation in the global diet of these hunter-gatherer populations. Such studies should however be carried out in combination with the elaboration of appropriate study protocols regarding the estimation of the size/weight of the exploited individuals of large mammal species. Much needed is also the elaboration of identification methods, should they be morphological or molecular, designed for Southeast Asian species to complement the existing works (Pritchard *et al.* 2009; Bochaton *et al.* 2019). Only at the cost of such investment the zooarchaeology of Southeast Asia will be on par with the rich literature existing on the material productions of prehistoric groups but this will also require the local development of a strong research community interested in that discipline that is still lacking at the present.

Acknowledgments

The authors are grateful to all the excavation teams who collected the studied remains as well to all the technical staff who helped us in the study of the different assemblages. We are especially grateful to Silpakorn University who host C. B. during most of the studies as well as to the Fine Art Departement of Thailand. We also thank S. Coleman and the Florida Museum of Natural History for welcoming us in their herpetology collections.

Data, scripts, code, and supplementary information availability

Supplementary information is available online: XXXXDOI of the webpage hosting the data
https://doi.org/10.5802/fake3.doi

Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article.

Funding

969 This work has been done thanks to the support of several funding agencies: the FYSSEN foundation, the
970 DIM-MAP funded by the Region île-de-France, the French Ministry of Europe and Foreign Affairs, and the
971 IRN PalBioDivASE 0846 funded by the CNRS. It was also supported by the Mission Préhistorique Franco-
972 Cambodgienne and the Mission Paléolithique Franco-Thaïe of the French Ministry of Europe and Foreign
973 Affairs (MEAE, Paris).
974

References

ADI B.T.H. 2000. — Archaeology of Ulu Kelantan PhD thesis. Canberra, *Australian National University*.
297 p.
AMPHANSRI A. 2011. — Temporal and Spatial Analysis of Faunal Remains from Tham Lod Rockshelter
Site, Pang Mapha District, Mae Hong Son Province M. A. Thesis. Bangkok, Thailand, *Silpakorn
University*.
ANDERSON D.D. 1990. — *Lang Rongrien Rockshelter: A Pleistocene, Early Holocene Archaeological Site
from Krabi, Southwestern Thailand*. Philadelphia, University of Pennsylvania Press. 108 p.
AUETRAKULVIT P. 2004. — Faunes du pléistocène final à l'holocène de Thaïlande : approche
archéozoologique PhD thesis. Marseille, *Aix Marseille 1*. 433 p.
AUETRAKULVIT P., FORESTIER H., KHAOKHIEW C. & ZEITOUN V. 2012. — New excavation at MohKhiew site
(Southern Thailand), *in* BONATZ D., REINECKE A. & TJOA-BONATZ M.L. (eds.), *Crossing Borders in
Southeast Asian Archaeology*. p. 62–74.
AUFFENBERG W. 1974. — Checklist of Fossil Land Tortoises (Testudinidae). *Bulletin of the Florida State
Museum Biological Sciences* 18 (3): 122–247
AVERY G., KANDEL A.W., KLEIN R.G., CONARD N.J. & CRUZ-URIBE K. 2004. — Tortoises as food and
taphonomic elements in palaeo « landscapes »*In:* XXIVe Rencontres internationales d'archéologie et
d'histoire d'Antibes, Antibes, France. p. 147–161.
BENNETT J.L. 1999. — Thermal Alteration of Buried Bone. *Journal of Archaeological Science* 26 (1): 1–
8. https://doi.org/10.1006/jasc.1998.0283
BITON R., SHARON G., ORON M., STEINER T. & RABINOVICH R. 2017. — Freshwater turtle or tortoise? The
exploitation of testudines at the Mousterian site of Nahal Mahanayeem Outlet, Hula Valley, Israel.
Journal of Archaeological Science: Reports 14: 409–419.
https://doi.org/10.1016/j.jasrep.2017.05.058
BLASCO R. 2008. — Human consumption of tortoises at Level IV of Bolomor Cave (Valencia, Spain).
Journal of Archaeological Science 35 (10): 2839–2848. https://doi.org/10.1016/j.jas.2008.05.013
BLASCO R., ROSELL J., SMITH K.T., MAUL L.C., SAÑUDO P., BARKAI R. & GOPHER A. 2016. — Tortoises as a
dietary supplement: A view from the Middle Pleistocene site of Qesem Cave, Israel. *Quaternary
Science Reviews* 133: 165–182. https://doi.org/10.1016/j.quascirev.2015.12.006
BLYTHE E. 1854. — Notices and descriptions of various reptiles, new or little-known. Part I. *The journal
of the Asiatic Society of Bengal* 22 (7): 639–655
BOCHATON C., HANOT P., FRÈRE S., CLAUDE J., NAKSRI W., AUETRAKULVIT P. & ZEITOUN V. 2019. — Size and
weight estimations of subfossil monitor lizards (Varanus sp. Merrem 1820) with an application to the
Hoabinhian assemblage of Doi Pha Kan (Late Pleistocene, Lampang province, Thailand). *Annales de
Paléontologie* 4 (105): 295–304. https://doi.org/10.1016/j.annpal.2019.05.003
BOCHATON C., IVANOV M. & CLAUDE J. 2019. — Osteological criteria for the specific identification of
Monitor lizards (*Varanus* Merrem, 1820) remains in subfossil deposits of Sundaland and continental
Southeast Asia. *Amphibia-Reptilia* 40 (2): 219–232. https://doi.org/10.1163/15685381-20181101
BULBECK F. 2003. — Hunter-Gatherer Occupation of the Malay Peninsula from the Ice Age to the Iron
Age, *Under the Canopy: The Archaeology of Tropical Rain Forests*. Brunswick, Rutgers University
Press. p. 119–160.

CELIBERTI V., FORESTIER H., AUETRAKULVIT P. & ZEITOUN V. 2018. — Doi Pha Kan, Lampang Province, North
of Thailand: A New Lithic Assemblage in the Hoabinhian Chrono-Cultural Context, *in* TAN N. (ed.),
Advancing Southeast Asian Archaeology. p. 91–99, 316–319.
CHUNG T.N. 2008. — Research on Hoa Binh culture in Viet Nam, Laos and Cambodia. *Vietnam*
Archaeology 3: 19–32
CLAUDE J., AUETRAKULVIT P., NAKSRI W., BOCHATON C., ZEITOUN V. & TONG H. 2019. — The recent fossil
turtle record of the central plain of Thailand reveals local extinctions. *Annales de Paléontologie* 4
(105): 305–315. https://doi.org/10.1016/j.annpal.2019.04.005
CLOSE L.M. & SEIGEL R. 1997. — Differences in body size among populations of red-eared sliders
(Trachemys scripta elegans) subjected to different levels of harvesting. *Chelonian Conservation and*
Biology 2 (4): 563–566
CODRON D., HOLT S., WILSON B. & HORWITZ L.K. 2022. — Skeletal allometries in the leopard tortoise
(Stigmochelys pardalis): Predicting chelonian body size and mass distributions in archaeozoological
assemblages, Quaternary Environments and Archaeology of the Northern Cape (South Africa).
Quaternary International 614: 59–72. https://doi.org/10.1016/j.quaint.2021.07.021
COLLECTIF 1932. — *Premier congrès des préhistoriens d'Extrême-Orient*. Hanoi, Imprimerie d'Extrême-
Orient. 154 p.
CONRAD C. 2015. — Archaeozoology in Mainland Southeast Asia: Changing Methodology and
Pleistocene to Holocene Forager Subsistence Patterns in Thailand and Peninsular Malaysia. *Open*
Quaternary 1 (1). https://doi.org/10.5334/oq.af
CONRAD C., HIGHAM C., EDA M. & MARWICK B. 2016. — Palaeoecology and Forager Subsistence
Strategies during the Pleistocene – Holocene Transition: A Reinvestigation of the Zooarchaeological
Assemblage from Spirit Cave, Mae Hong Son Province, Thailand. *Asian Perspectives* 55 (1): 2–27.
https://doi.org/10.1353/asi.2016.0013
DAS I. 2010. — *A field guide to the reptiles of South-East Asia*. London, New Delhi, New York, Sydney,
Bloomsbury Publishing. 376 p.
DODSON P. & WEXLAR D. 1979. — Taphonomic Investigations of Owl Pellets. *Paleobiology* 5 (3): 275–
284
DUNN F.L. 1964. — Excavations at Gua Kechil, Pahang. *Journal of the Malaysian Branch of the Royal*
Asiatic Society 37 (2 (206)): 87–124
EBERLING G. 2011. — Haltung und Nachzucht von *Indotestudo elongata* (Blythe, 1853). *Draco* 2: 61–72
EHRET D. 2007. — Skeletochronology: A Method for Determing the Individual Age and Growth of
Modern and Fossil Tortoises (Reptilia: Testudines). *Bulletin of the Florida Museum of Natural History*
47 (2): 49–72
EMMONS L.H. 1989. — Jaguar Predation on Chelonians. *Journal of Herpetology* 23 (3): 311–314.
https://doi.org/10.2307/1564460
ESKER D.A., FORMAN S.L. & BUTLER D.K. 2019. — Reconstructing the mass and thermal ecology of North
American Pleistocene tortoises. *Paleobiology* 45 (2): 363–377. https://doi.org/10.1017/pab.2019.6
FORESTIER H. 2003. — Des outils nés de la forêt : l'importance du végétal en Asie du Sud-Est dans
l'imagination et l'invention technique aux périodes préhistoriques, *in* FROMENT A., GUFFROY J., &
PEUPLEMENTS ANCIENS ET ACTUELS DES FORÊTS TROPICALES : SÉMINAIRE-ATELIER, ORLÉANS (FRA), 1998/10/15-
16 (eds.), *Peuplements anciens et actuels des forêts tropicales : actes du séminaire-atelier*. Paris, IRD.
p. 315–337. (Colloques et Séminaires).
FORESTIER H., GRIGGO C. & SOPHADY H. 2023. — L'Hoabinhien ou le paradigme égaré de la modernité
européenne en Extrême-Orient : l'exemple de la Préhistoire du Cambodge*In:* 29e Congrès
préhistorique de France, Toulouse. Société préhistorique française. p. 59–76.
FORESTIER H., SOPHADY H. & CELIBERTI V. 2017. — Le techno-complexe hoabinhien en Asie du Sud-est
continentale : L'histoire d'un galet qui cache la forêt. *Journal of Lithic Studies* 4 (2): 305–349.
https://doi.org/10.2218/jls.v4i2.2545
FORESTIER H., SOPHADY H., PUAUD S., CELIBERTI V., FRÈRE S., ZEITOUN V., MOURER-CHAUVIRÉ C., MOURER R.,
THAN H. & BILLAULT L. 2015. — The Hoabinhian from Laang Spean Cave in its stratigraphic,

chronological, typo-technological and environmental context (Cambodia, Battambang province). *Journal of Archaeological Science: Reports* 3: 194–206. https://doi.org/10.1016/j.jasrep.2015.06.008

FORESTIER H., ZEITOUN V., WINAYALAI C. & MÉTAIS C. 2013. — The open-air site of Huai Hin (Northwestern Thailand): Chronological perspectives for the Hoabinhian. *Comptes Rendus Palevol* 12 (1): 45–55. https://doi.org/10.1016/j.crpv.2012.09.003

FORESTIER H., ZHOU Y., AUETRAKULVIT P., KHAOKHIEW C., LI Y., JI X. & ZEITOUN V. 2021. — Hoabinhian variability in Mainland Southeast Asia revisited: The lithic assemblage of Moh Khiew Cave, Southwestern Thailand. *Archaeological Research in Asia* 25: 100236. https://doi.org/10.1016/j.ara.2020.100236

FORESTIER H., ZHOU Y., VIALLET C., AUETRAKULVIT P., LI Y. & SOPHADY H. 2022. — Reduction Sequences During the Hoabinhian Technocomplex in Cambodia and Thailand: A New Knapping Strategy in Southeast Asia from the Terminal Upper Pleistocene to mid Holocene. *Lithic Technology* 47 (2): 147–170. https://doi.org/10.1080/01977261.2021.1981654

FRÈRE S., AUETRAKULVIT P., ZEITOUN V., SOPHADY H. & FORESTIER H. 2018. — Doi Pha Kan (Thailand), Ban Tha Si (Thailand) and Laang Spean (Cambodia) Late Paleolithic animal bone assemblages. A new perception of meat supply strategies for Early Holocene Mainland South-east Asia?, *in* TAN N.H. (ed.), *Advancing Southeast Asian Archaeology*. p. 100–108, 320–323.

GARBIN R.C., ASCARRUNZ E. & JOYCE W.G. 2018. — Polymorphic characters in the reconstruction of the phylogeny of geoemydid turtles. *Zoological Journal of the Linnean Society* 184 (3): 896–918. https://doi.org/10.1093/zoolinnean/zlx106

GIL-SÁNCHEZ J.M., RODRÍGUEZ-CARO R.C., MOLEÓN M., MARTÍNEZ-PASTOR M.C., LEÓN-ORTEGA M., EGUÍA S., GRACIÁ E., BOTELLA F., SÁNCHEZ-ZAPATA J.A., MARTÍNEZ-FERNÁNDEZ J., ESTEVE-SELMA M.A. & GIMÉNEZ A. 2022. — Predation impact on threatened spur-thighed tortoises by golden eagles when main prey is scarce. *Scientific Reports* 12 (1): 17843. https://doi.org/10.1038/s41598-022-22288-9

GLOVER I.C. 1977. — The Hoabinhian hunter-gatherers or early agriculture list in Southeast Asia, *in* MEGAW J.V.S. (ed.), *Hunter Gatherers and First Farmer beyond Europe*. Leicester, UK, Leicester University Press. p. 145–166.

GORMAN C. 1971. — The Hoabinhian and After: Subsistence Patterns in Southeast Asia during the Late Pleistocene and Early Recent Periods. *World Archaeology* 2 (3): 300–320

GORMAN C.F. 1970. — Excavations at Spirit Cave, North Thailand: SOME INTERIM INTERPRETATIONS. *Asian Perspectives* 13: 79–107

GORMAN C.F. 1969. — Hoabinhian: A Pebble-Tool Complex with Early Plant Associations in Southeast Asia. *Science* 163 (3868): 671–673. https://doi.org/10.1126/science.163.3868.671

GOULD S.J. 1966. — Allometry and size in ontogeny and phylogeny. *Biological Reviews* 41 (4): 587–640

GUERIN C. & MOURER-CHAUVIRÉ C. 1969. — Le Rhinoceros sondaicus Desmarest du gisement néolithique de Loang Spean, province de Battambang, Cambodge. *Ann. Fac. Sc. Phnom Penh, 1970* 2: 261–274

HAILEY A. & COULSON I. 1999. — The growth pattern of the African tortoise Geochelone pardalis and other chelonians. *Canadian Journal of Zoology* 77: 181–193. https://doi.org/10.1139/cjz-77-2-181

HANSEL T. 2004. — Observations on subsistence hunting along the Phu Yai Mountain Range, Xanakham District, Vientiane Province, Lao PDR. *Natural History Bulletin of the Siam Society* 52: 195–200

HIGHAM C. 2002. — *Early Cultures of Mainland Southeast Asia*.Art Media Resources. 380 p.

HIGHAM C.F.W. 1977. — Economic Change in Prehistoric Thailand, *in* REED C. (ed.), *Origins of Agriculture*. The Hauge, De Gruyter Mouton. p. 385–412. https://doi.org/10.1515/9783110813487.385

HUXLEY J.S. 1932. — *Problems of relative growth*. London, Methuen & co. 319 p.

IHLOW F., DAWSON J.E., HARTMANN T. & SOM S. 2016. — *Indotestudo elongata* (Blyth 1854) – Elongated Tortoise, Yellow-headed Tortoise, Yellow Tortoise. *Chelonian Conservation and Biology* 5: 1–14. https://doi.org/10.3854/crm.5.096.elongata.v1.2016

IHLOW F. & HANDSCHUH M. 2011. — Auswilderung von Indotestudo elongata im Kulen Promtep
Wildlife Sanctuary im Norden Kambodschas. *Marginata* 8 (4): 16–23

IMDIRAKPHOL S., ZAZZO A., AUETRAKULVIT P., TIAMTINKRIT C., PIERRET A., FORESTIER H. & ZEITOUN V. 2017. —
The perforated stones of the Doi Pha Kan burials (Northern Thailand): A Mesolithic singularity?
Comptes Rendus Palevol 16 (3): 351–361. https://doi.org/10.1016/j.crpv.2016.12.003

IVERSON J.B., SPINKS P.Q., SHAFFER B.H., MCCORD W.P. & DAS I. 2001. — Phylogenetic relationships
among the asian tortoises of the genus *Indotestudo* (Reptilia: Testudines:testudinidae). *Hamadryad*
26 (2): 283–286

IVERSON J.B. 1982. — Biomass in turtle populations: A neglected subject. *Oecologia* 55 (1): 69–76.
https://doi.org/10.1007/BF00386720

JI X., KUMAN K., CLARKE R.J., FORESTIER H., LI Y., MA J., QIU K., LI H. & WU Y. 2016. — The oldest
Hoabinhian technocomplex in Asia (43.5 ka) at Xiaodong rockshelter, Yunnan Province, southwest
China, Peking Man and related studies. *Quaternary International* 400: 166–174.
https://doi.org/10.1016/j.quaint.2015.09.080

KLEIN R.G. & CRUZ-URIBE K. 1983. — Stone Age Population Numbers and Average Tortoise Size at
Byneskranskop Cave 1 and DieKelders Cave 1, Southern Cape Province, South Africa. *The South
African Archaeological Bulletin* 38 (137): 26–30

MASOJC M., LE H.D., GRALAK T., MICHALEC G., APOLINARSKA K., BADURA M., CENDROWSKA M., GALAS A.,
KRUPA-KURZYNOWSKA J., MIAZGA B., OSPINSKA M., ROZOK Z. & VIET N. 2023. — The Early Holocene
Hoabinhian (8300-8000 cal BC) occupation from Hiem Cave, Vietnam. *Comptes Rendus Palevol* 22
(5): 59–76

MEDWAY, LORD 1969. — Excavations At Gua Kechil, Pahang III. Animal Remains. *Journal of the
Malaysian Branch of the Royal Asiatic Society* 42 (2 (216)): 197–205

MENA P., STALLINGS J.R., REGALADO J. & CUEVA R. 2000. — The Sustainability of Current Hunting Practices
by the Huaorani. *In Hunting for Sustainability in Tropical Forests*: 57–78

MOSER J. 2001. — *Hoabinhian: Geographie und Chronologie eines steinzeitlichen Technokomplexes in
Südostasien*. Germany, Linden Soft. 194 p. (AVA - Forschungen; 6).

MOURER-CHAUVIRÉ C. & MOURER R. 1970. — The Prehistoric Industry of Laang Spean, Province of
Battambang, Cambodia. *Archaeology and Physical Anthropology in Oceania* 5: 128–146.
https://doi.org/10.1002/j.1834-4453.1970.tb00110.x

MOURER-CHAUVIRÉ C., MOURER R. & THOMMERET Y. 1970. — Premières datations absolues de l'habitat
préhistorique de la grotte de Laang Spean, province of Battambang (Cambodge). *C. R. Acad. Sci.
Paris, Ser. D* 270: 471–473

MUDAR K. & ANDERSON D.D. 2007. — New Evidence for Southeast Asian Pleistocene Foraging
Economies: Faunal Remains from the Early Levels of Lang Rongrien Rockshelter, Krabi, Thailand.
Asian Perspectives 46 (2): 298–334

NABAIS M., BONETA I. & SOARES R. 2019. — Chelonian use in Portugal: Evidence from Castelo Velho de
Safara. *Journal of Archaeological Science: Reports* 28: 102054.
https://doi.org/10.1016/j.jasrep.2019.102054

NABAIS M. & ZILHÃO J. 2019. — The consumption of tortoise among Last Interglacial Iberian
Neanderthals, Neanderthals: Ecology and Evolution. *Quaternary Science Reviews* 217: 225–246.
https://doi.org/10.1016/j.quascirev.2019.03.024

NAKSRI W. 2013. — Origin and changes in continental turtle diversity from the Miocene to Holocene
in Thailand PhD thesis. Maha Sarakham, Thailand, *Mahasarakham University*.

NAKSRI W. 2007. — Appraisal of the evolution of testudinoid turtle diversity from the Oligocene and
Neogene of Thailand Master thesis. Maha Sarakham, Thailand, *Mahasarakham University*. 230 p.

NAKSRI W., TONG H., LAUPRASERT K., SUTEETHORN V. & CLAUDE J. 2013. — A new species of *Cuora*
(Testudines: Geoemydidae) from the Miocene of Thailand and its evolutionary significance.
Geological Magazine 150 (5): 908–922. https://doi.org/10.1017/S0016756812001082

PLATT S.G., AUNG S.H.N., SOE M.M., LWIN T., PLATT K., WALDE A.D. & RAINWATER T.R. 2021. — Predation
on Translocated Burmese Star Tortoise (Geochelone platynota) by Asiatic Jackals (Canis aureus) and

Wild Pigs (Sus scrofa) at a Wildlife Sanctuary in Myanmar. *Chelonian Conservation and Biology* 20 (1): 133–138. https://doi.org/10.2744/CCB-1461.1

POOKAJORN S. 2001. — New perspectives for Palaeolithic research in Thailand, *Origine des peuplements et chronologie des cultures paléolithiques dans le sud-est asiatique*. Paris, France, Semenanjung. p. 167–188.

PRITCHARD P.C.H., RABETT R.J. & PIPER P.J. 2009. — Distinguishing species of Geoemydid and Trionychid turtles from shell fragments: evidence from the Pleistocene at Niah Caves, Sarawak. *International Journal of Osteoarchaeology* 19 (4): 531–550. https://doi.org/10.1002/oa.1038

R CORE TEAM 2020. — R: A language and environment for statistical computing

RABETT R.J. 2012. — Tropical Subsistence Strategies at the End of the Last Glacial, *Human Adaptation in the Asian Palaeolithic: Hominin Dispersal and Behaviour during the Late Quaternary*. Cambridge, UK, Cambridge University Press. p. 208–265.

RAMESH M. 2008. — Relative Abundance and Morphometrics of the Travancore Tortoise, Indotestudo travancorica, in the Indira Gandhi Wildlife Sanctuary, Southern Western Ghats, India. *Chelonian Conservation and Biology* 7 (1): 108–113

RHODIN A.G.J. 1992. — Chelonian Zooarchaeology of Eastern New England: Turtle bone remains from Cedar Swamp and other prehistoric sites. *Bulletin of the Massachussetts Archaeological Society* 53 (1): 21–30

RHODIN A.G.J., IVERSON J., BOUR R., FRITZ U., GEORGES A., SHAFFER B.H. & VAN DIJK P.P. 2021. — *Turtles of the World: Annotated Checklist and Atlas of Taxonomy, Synonymy, Distribution, and Conservation Status (9th Ed.)*. United States of America, Chelonian Research Foundation and Turtle Conservancy. 482 p. (Chelonian Research Monograph; 8).

ROUAG R., BENYACOUB S., LUISELLI L., EL MOUDEN E.H., TIAR G. & FERRAH C. 2007. — Population structure and demography of an Algerian population of the Moorish tortoise, Testudo graeca. *Animal Biology* 57: 267–279. https://doi.org/10.1163/157075607781753065

SANTOS A., MAYOR P., LOUREIRO L., GILMORE M., PEREZ PEÑA P., BOWLER M., LEMOS L., SVENSSON M., NEKARIS K.A., NIJMAN V., VALSECCHI J. & MORCATTY T. 2020. — Widespread Use of Traditional Techniques by Local People for Hunting the Yellow-Footed Tortoise (Chelonoidis denticulatus) Across the Amazon. *Journal of Ethnobiology* 40: 268. https://doi.org/10.2993/0278-0771-40.2.268

SHOOCONGDEJ R. 1996. — Forager mobility organization in seasonal tropical environments: A view from Lang Kamnan Cave, western Thailand. Thesis.

SOM S. & COTTET M. 2016. — Rescue and relocation programme of turtles and tortoises and elongated tortoise monitoring programme in the Nam Theun 2 Reservoir (Laos). *Hydroécologie Appliquée* 19: 383–406. https://doi.org/10.1051/hydro/2015007

SOPHADY H. 2016. — Archeo-stratigraphy of Laang Spean prehistoric site (Battambang province): a contribution to Cambodian prehistory PhD thesis. Paris, France, *Muséum national d'Histoire naturelle*. 474 p.

SOPHADY H., FORESTIER H., ZEITOUN V., PUAUD S., FRÈRE S., CELIBERTI V., WESTAWAY K., MOURER R., MOURER-CHAUVIRÉ C., THAN H., BILLAULT L. & TECH S. 2016. — Laang Spean cave (Battambang province): A tale of occupation in Cambodia from the Late Upper Pleistocene to Holocene, Southeast Asia: human evolution, dispersals and adaptation. *Quaternary International* 416: 162–176. https://doi.org/10.1016/j.quaint.2015.07.049

SPETH J.D. & TCHERNOV E. 2002. — Middle Paleolithic Tortoise Use at Kebara Cave (Israel). *Journal of Archaeological Science* 29 (5): 471–483. https://doi.org/10.1006/jasc.2001.0740

SRIPRATEEP K., ARANYAVALAI V., AOWPHOL A. & THIRAKHUPT K. 2013. — Population Structure and Reproduction of the Elongated Tortoise Indotestudo elongata (Blyth, 1853) at Ban Kok Village, Northeastern Thailand. *Tropical Natural History* 13 (1): 21–37

STINER M.C., MUNRO N.D. & SUROVELL T.A. 2000. — The Tortoise and the Hare: Small-Game Use, the Broad-Spectrum Revolution, and Paleolithic Demography. *Current Anthropology* 41 (1): 39–79. https://doi.org/10.1086/300102

TAYLOR E.H. 1970. — The turtles and crocodiles of Thailand and adjacent waters; with a synoptic
herpetological bibliography. *The University of Kansas Science Bulletin* 49 (3): 87–179
THEOBALD W. 1868. — Catalogue of the Reptiles of British Birma, embracing the Provinces of Pegu,
Martaban, and Tenasserim; with descriptions of new or little-known species. *Journal of the Linnean*
Society of London, Zoology 10 (41): 4–67. https://doi.org/10.1111/j.1096-3642.1868.tb02007.x
THOMPSON J.C. & HENSHILWOOD C.S. 2014. — Tortoise taphonomy and tortoise butchery patterns at
Blombos Cave, South Africa. *Journal of Archaeological Science* 41: 214–229.
https://doi.org/10.1016/j.jas.2013.08.017
TREERAYAPIWAT C. 2005. — Patterns of Habitation and Burial Activity in the Ban Rai Rock Shelter,
Northwestern Thailand. *Asian Perspectives* 44 (1): 231–245. https://doi.org/10.1353/asi.2005.0001
TUNGITTIPLAKORNL W. & DEARDEN P. 2002. — Hunting and wildlife use in some Hmong communities in
northern Thailand. *Natural History Bulletin of the Siam Society* 50: 57–73
VAN DIJK P.P. 1998. — The Natural History of the Elongated Tortoise Indotestudo Elongata (Blyth,
1853) (Reptilia: Testudines) in a Hill Forest Mosaic in Western Thailand with Notes on Sympatric
Turtle Species Ph.D. dissertation. Galway, Ireland, *National University of Ireland*. 255 p.
VAN VLACK H. 2014. — Forager Subsistence Regimes in the Thai-Malay Peninsula: An Environmental
Archaeological Case Study of Khao Toh Chong Rockshelter, Krabi Master thesis. 4484. San Jose, *San*
José State University. 188 p.
VILLA P., BON F. & CASTEL J.-C. 2002. — Fuel, fire and fireplaces in the Palaeolithic of western Europe.
Review of Archaeology 23: 33–42
VU T.L. 1994. — The Hoabinhian animal complex. *Vietnam Soc. Sci.* 5 (43): 70–74
WHITE J.C. 2011. — Emergence of cultural diversity in mainland Southeast Asia: A view from
prehistory, *in* ENFIELD N.J. (ed.), *Dynamics of Human Diversity: The Case of Mainland Southeast Asia*.
Canberra, Australian National University. p. 9–46.
YEN E. 1977. — Hoabinhian horticulture? The evidence and the questions from Northwest Thailand,
in ALLEN J.G. & JONES R. (eds.), *Sunda and Sahul. Prehistoric Studies in Southeast Asia, Melanesia and*
Australia. London, Academic Press. p. 567–599.
ZEITOUN V., AUETRAKULVIT P., ZAZZO A., PIERRET A., FRÈRE S. & FORESTIER H. 2019. — Discovery of an
outstanding Hoabinhian site from the Late Pleistocene at Doi Pha Kan (Lampang province, northern
Thailand). *Archaeological Research in Asia* 18: 1–16. https://doi.org/10.1016/j.ara.2019.01.002
ZEITOUN V., FORESTIER H., SOPHADY H., PUAUD S. & BILLAULT L. 2012. — Direct dating of a Neolithic burial
in the Laang Spean cave (Battambang Province, Cambodia): First regional chrono-cultural
implications. *Comptes Rendus Palevol* 11 (7): 529–537. https://doi.org/10.1016/j.crpv.2012.06.006
ZEITOUN V., FORESTIER H. & SUPAPORN N. 2008. — *Préhistoires au sud du Triangle d'Or* 252 p.
ZEITOUN V., HENG S. & FORESTIER H. 2021. — The funeral cave of Laang Spean*In:* SEAMEO SPAFA
International Conference on Southeast Asian Archaeology and Fine Arts, 13-17 December 2021,
Bangkok, Thailand. SEAMEO SPAFA. p. 1–14. https://doi.org/10.26721/spafa.pqcnu8815a-14
ZNARI M., GERMANO D.J. & MACÉ J.-C. 2005. — Growth and population structure of the Moorish
Tortoise (Testudo graeca graeca) in Westcentral Morocco: possible effects of over-collecting for the
tourist trade. *Journal of Arid Environments* 62 (1): 55–74.
https://doi.org/10.1016/j.jaridenv.2004.11.013
ZURAINA M. 1994. — *The Excavation of Gua Gunung Runtuh and the Discovery of the Perak Man in*
Malaysia. Kuala Lumpur, Department of Museums and Antiquity Malaysia. 174 p.
1262

1263   **Appendix**

1264   Appendix 1: All measurements recorded on the modern turtle skeletons.

1265

A

M. s. 1   C. s.

Pe.

Nu. p.

Pl. s. 1   *Co.*

Ne. 1

V. s. 3

S. p. 1

Dorsal view

3 cm

Ne.

S. p. 1

Co.

S. p. 2

S. s.   *Py.*   *Pe.*

Posterior view

C

1 cm

Posterior view

B

G. s.

Ep.

H. s.   *En.*

P. s.

Hyo.

A. s.

Hyp.

F. s.

Xi.

An. s.

Ventral view

Ep.

En.

Hyo.

Hyp.

Xi.

Dorsal view

D

F.

1 cm

Ventral view

Nuchal plate
(dorsal view)

Peripheral plate
(lateral view)

Pygal plate
(posterior view)

Neural plate
(dorsal view)

Entoplastron
(ventral view)

Supra-pygal plate 2
(posterior view)

Epiplastron
(ventral view)

Hyoplastron
(ventral view)

Hypoplastron
(ventral view)

Xiphiplastron
(ventral view)

Humerus
(Posterior view)   (Dorsal view)

Radius

Ulna

Femur
(Posterior view)   (Dorsal view)

Tibia
(Posterior view)   (Lateral view)

Fibula
(Lateral view)(Posterior view)

A **Doi Pha Kan Rock Shelter**

Carapace   Plastron   Skeleton

PR
0%
1-10%
10-20%
20-30%
30-40%
40-50%
50-60%
60-70%
70-80%
80-90%
90-100%

NMI= 55

B **Moh Khiew Cave Layer 1**

Carapace   Plastron   Skeleton

PR
0%
1-10%
10-20%
20-30%
30-40%
40-50%
50-60%
60-70%
70-80%
80-90%
90-100%

NMI= 6

C **Moh Khiew Cave Layer 2**

Carapace   Plastron   Skeleton

PR
0%
1-10%
10-20%
20-30%
30-40%
40-50%
50-60%
60-70%
70-80%
80-90%
90-100%

NMI= 53

D **Moh Khiew Cave Layer 3**

Carapace   Plastron   Skeleton

PR
0%
1-10%
10-20%
20-30%
30-40%
40-50%
50-60%
60-70%
70-80%
80-90%
90-100%

NMI= 15

E **Moh Khiew Cave Layer 4**

Carapace   Plastron   Skeleton

PR
0%
1-10%
10-20%
20-30%
30-40%
40-50%
50-60%
60-70%
70-80%
80-90%
90-100%

NMI= 5

Laang Spean Cave Hoabinhian level

Carapace   Plastron   Skeleton

PR

0%
1-10%
10-20%
20-30%
30-40%
40-50%
50-60%
60-70%
70-80%
80-90%
90-100%

NMI= 56

Laang Spean Cave "Neolithic" layers

Carapace   Plastron   Skeleton

PR

0%
1-10%
10-20%
20-30%
30-40%
40-50%
50-60%
60-70%
70-80%
80-90%
90-100%

NMI= 12

Laang Spean Cave "Sepulture" layers

Carapace   Plastron   Skeleton

PR

0%
1-10%
10-20%
20-30%
30-40%
40-50%
50-60%
60-70%
70-80%
80-90%
90-100%

NMI= 12

Kao Tah Plai Test Pit 1 - Metal Ages

Carapace Plastron Skeleton

PR
0%
1-10%
10-20%
20-30%
30-40%
40-50%
50-60%
60-70%
70-80%
80-90%
90-100%

NMI= 14

Kao Tah Plai Test Pit 1 - Neolithic

Carapace Plastron Skeleton

PR
0%
1-10%
10-20%
20-30%
30-40%
40-50%
50-60%
60-70%
70-80%
80-90%
90-100%

NMI= 9

Kao Tah Plai Test Pit 2 - Neolithic

Carapace Plastron Skeleton

PR
0%
1-10%
10-20%
20-30%
30-40%
40-50%
50-60%
60-70%
70-80%
80-90%
90-100%

NMI= 21