1   **Small ADP-ribosylation factor-like GTPase TITAN5 (TTN5/ARL2) is linked with the**
2   **dynamic regulation of IRON-REGULATED TRANSPORTER1**
3

4   Inga Mohr[1], Monique Eutebach[1], Naima Schommen[1], Regina Gratz[1], Kalina Angrand[2], Lara
5   Genders[1], Tzvetina Brumbarova[1], Petra Bauer[1,3] and Rumen Ivanov[1]
6
7   [1]Institute of Botany, Heinrich Heine University, Universitätsstr. 1, 40225 Düsseldorf, Germany.
8   [2]former address: Department of Biosciences-Plant Biology, Saarland University, 66123
9   Saarbrücken, Germany.
10 [3]Cluster of Excellence on Plant Sciences (CEPLAS), Heinrich-Heine University, 40225
11 Düsseldorf, Germany.

12
13 **Author contributions:** P.B. and R.I. conceived the project; I.M., T.B., P.B. and R.I. designed
14 experiments, supervised the research and analyzed data; I.M., M.E., N.S., R.G., and K.A.
15 performed experiments; L.G. contributed key materials; I.M. wrote the manuscript; I.M., P.B.
16 and R.I. edited the manuscript; P.B. acquired funding. P.B. and R.I. agreed to serve as the
17 author responsible for contact and communication.
18
19
Highlights
• The small GTPase TITAN 5 (TTN5/ARL2) is linked to iron homeostasis
• TTN5 interacts with the large intracellular loop and variable region of IRON-REGULATED
TRANSPORTER 1 (IRT1) and has a positive effect on root iron (Fe) reduction
• TTN5 interacts with peripheral membrane proteins of the IRT1 interactome, EHB1, PATL2
and SNX1 suggesting a coordinating role in IRT1 regulation
• TTN5 and IRT1 colocalize at the plasma membrane and the endomembrane system
related to vesicle transport
28
29 **Running title**
30 ARF-like GTPase TTN5 and iron acquisition
31
32 **One-sentence summary**
33 ADP-ribosylation factor-like small GTPase TITAN 5 (TTN5/ARL2) links vesicle transport and
34 iron homeostasis by binding the intracellular regulatory protein interaction region of IRON-
35 REGULATED TRANSPORTER 1 (IRT1).
36
37 **Keywords**
38 ARF-like; ARLC1; ARL2; EHB1; endomembrane; GTPase; HALLIMASCH; iron;
39 PATL2; plasma membrane; SEC14; SNX1; vesicle
40
Abstract

The endomembrane system is the central sorting machinery for membrane proteins where
signals can be integrated to control plant nutritional responses. IRON-REGULATED
TRANSPORTER1 (IRT1) is tightly controlled through trafficking between the endomembrane
system and the plasma membrane to balance the demands for iron (Fe) and toxic effects that

may arise from excess of this micronutrient in roots. Small GTPases of the Ras superfamily
regulate signaling inputs in endomembrane trafficking, but their functions and activities are
poorly known in plants. One plant endomembrane system-associated regulatory component is
the small ARF-like GTPase TITAN 5/ARL2/ARLC1/HALLIMASCH (hereafter termed TTN5).
TTN5 is required during early embryo development but many open questions remain with
regard to its physiological function during the later life cycle or in response to environmental
cues. Here, we demonstrate that TTN5 interacts with IRT1vr, the large cytoplasmic variable
region and protein-regulatory platform of IRT1. TTN5 also interacts with peripheral membrane
proteins that are components of the IRT1 regulation machinery and can be involved in protein
trafficking, like SNX1, the C2 domain protein EHB1 and the SEC14-GOLD domain-containing
PATL2. We show that heterozygous *ttn5-1*$^{+/-}$ plants have a Fe reduction phenotype.
Fluorescent fusion proteins of TTN5 and IRT1 colocalize at the plasma membrane and in
endosomes/multivesicular bodies, where IRT1 sorting and cycling between the plasma
membrane and the vacuole are coordinated. Hence, this work links IRT1 and iron nutritional
regulation with a small GTPase. This opens up the possibility that iron nutrient levels are
controlled *via* the cellular regulation of the endomembrane system.
62

63 **Abbreviations**

AD	activation domain
Arabidopsis	*Arabidopsis thaliana*
ARF-like / ARL	ADP-ribosylation factor-like
BD	binding domain
EE	early endosomes
EHB1	ENHANCED BENDING 1
FRET-APB	Förster Resonance Energy Transfer-Acceptor Photobleaching
GAP	GTPase-activating protein
GEF	guanine nucleotide exchange factor
ICQ	intensity correlation quotient
MVB	multivesicular body
PATL2	PATELLIN 2
ROI	region of interest
SNX1	SORTING NEXIN 1
TGN	*trans*-Golgi network
TTN5	TITAN 5
Y2H	yeast two-hybrid
ZIP	ZRT/IRT-LIKE PROTEIN

Introduction
Plants adjust to environmental changes, such as variations of soil nutrient availability.
Cellular membranes are very dynamic in plants, and this is crucial for sessile organisms. Plants
can scan their environment for nutrients and control the activities of plasma membrane
transporter proteins by vesicular trafficking. Many open questions still exist with regard to the
environmental responsiveness and the complexity of the plant cell endomembrane system.
One essential element, iron (Fe), is abundant in the soil. But since Fe is mostly present in
insoluble ferric Fe complexes, it must be mobilized into a bioavailable form. Plants like

Arabidopsis thaliana (Arabidopsis) are capable of mobilizing ferric Fe^{3+} by reducing it *via* plasma membrane FERRIC REDUCTION OXIDASE 2 (FRO2) (Robinson et al., 1999). The reactive ferrous Fe^{2+} is taken up by IRON-REGULATED TRANSPORTER 1 (IRT1) into root epidermis cells (Eide et al., 1996; Vert et al., 2002).

IRT1 is one of the founding members of the evolutionarily conserved ZIP (ZRT, IRT-LIKE PROTEIN) divalent metal ion transporter family (Eide et al., 1996; Guerinot, 2000). Divalent metal ions are potentially cytotoxic. The activity of IRT1, like that of other ZIP proteins, is controlled at the level of protein abundance at the plasma membrane (Hu, 2021). In this respect, IRT1 serves as a model system for studying the control of a ZIP transporter. Importantly, IRT1 protein activity is controlled through transport from the *trans*-Golgi network/early endosomes (TGN/EE) to the plasma membrane, degradation by the lytic pathway and recycling back to the plasma membrane (Barberon et al., 2011; Shin et al., 2013; Barberon et al., 2014; Foot et al., 2017; Zhang et al., 2023) and (Ivanov et al. 2014). The large variable cytosolic region between transmembrane domains three and four (vr) is a characteristic regulatory feature of ZIP proteins (Guerinot, 2000; Gaither and Eide, 2001). In IRT1, this regulatory variable region, IRT1vr, is crucial for IRT1 control and target of metal ion binding and post-translational modification (Grossoehme et al., 2006; Kerkeb et al., 2008; Dubeaux et al., 2018; Cointry and Vert, 2019). Clearly, Fe acquisition is affected by regulation of the vesicular transport mechanism, but how this is achieved and which proteins are involved are open questions that still remain largely elusive.

Cycling of plasma membrane proteins is an important function of endomembrane trafficking (Valencia et al., 2016; Ivanov and Vert, 2021). One protein family involved in this process in mammalian cells and yeast is the Ras superfamily of small GTPases. Small GTPases are important in vesicle-related processes as they function as molecular switches due to their GTPase activities to transmit signals. Small GTPases have usually low intrinsic nucleotide exchange and hydrolysis activity, and therefore, require the support of guanine nucleotide exchange factors (GEFs) and GTPase-activating proteins (GAPs). GEFs recruit the inactive, GDP-bound, GTPase to their site of action and lead to nucleotide exchange by GDP release. GTP binding leads to a conformational change of two regions referred to as switch I and II. The active, GTP-loaded, GTPases then exert their function, e.g. the recruiting and subsequent assembly of coat proteins by ARF1 or SAR1, until hydrolysis of the GTP by GAPs. Most of the known interactions occur in the active conformation of the GTPases (Sztul et al., 2019; Nielsen, 2020; Adarska et al., 2021). Although the ARF-GTPase family is well studied in mammals, there are only few well-established examples of an ARF-GTPase functioning in plasma membrane protein trafficking in plants, like that of ARF1-GNOM-mediated recycling of auxin carrier PIN FORMED 1 (PIN1), with GNOM being an ARF-GEF (Steinmann et al., 1999; Geldner et al., 2003). The presence of 21 ARF and ARF-like (ARL) GTPases encoded by the Arabidopsis genome raises the question whether other ARF members may act in endomembrane trafficking and what are their functions in the diverse physiological and nutrient assimilation pathways besides auxin signaling and cell division.

TITAN5 (TTN5), also known as HALLIMASCH (HAL)/ARL2/ARLC1, is an ARF-like protein. It was identified in two independent screens for abnormal-embryo mutants arrested soon after egg cell division, indicating a central cellular function (Mayer et al., 1999; McElver et al., 2000), in agreement with its ubiquitous expression in plants (Mohr et al., 2023). Physiological functions of this small GTPase, relevant during other phases of the plant life cycle, remain elusive until today. TTN5 is more closely related in sequence to human ADP-ribosylation factor-like2

(HsARL2) than any Arabidopsis protein. HsARL2 is associated with very diverse roles in cells,
ranging from microtubule development, also identified for yeast and Caenorhabditis homologs
(Bhamidipati et al., 2000; Fleming et al., 2000; Radcliffe et al., 2000; Antoshechkin and Han,
2002; Tzafrir et al., 2002; Mori and Toda, 2013), adenine nucleotide transport in mitochondria
(Sharer et al., 2002) and control of phosphodiesterase activity in cilia (Ismail et al., 2011; Fansa
and Wittinghofer, 2016). TTN5 orthologs are therefore not only involved in a diverse set of
functions and signaling cascades but they also bind very different protein partners for that.
TTN5 is expressed in the root epidermis (Mohr et al., 2023), like *IRT1* (Vert et al., 2002). TTN5
is present at the plasma membrane and in the endomembrane compartment. Its unusually high
intrinsic GDP-GTP exchange activity, that causes TTN5 to be present predominantly in a GTP-
binding form, seems a prerequisite for its dynamic cellular localization (Mohr et al., 2023). TTN5
can be considered atypical as it does not require a GEF to be in its active GTP-bound form and
has a slow GTPase activity relying on a GAP (Mohr et al., 2023).

A yeast two-hybrid screen identified novel IRT1vr regulators (Khan et al., 2019; Hornbergs
et al., 2022). We report here, that with this same strategy, we retrieved the HsARL2 homolog
and small putative GTPase TTN5. We validated protein interactions of TTN5 and IRT1vr as
well as the IRT1vr interactome and found a physiological Fe reductase phenotype linked with
root Fe acquisition. Colocalization analysis indicated that the active GTP-bound TTN5 form
promoted the presence of IRT1 in intracellular vesicles. Hence, the dynamic small GTPase
TTN5 acts in a nutritional-environmental context in the vesicular trafficking system that serves
to control the presence of a root iron transporter at the plasma membrane.

Results

TTN5 interacts with the variable region of IRT1

To identify new cytoplasmic proteins regulating IRT1 we had developed a strategy for
identifying IRT1vr candidate interaction partners of IRT1vr (residues 145-192), and used
IRT1vr as bait against a cDNA expression library prepared from Fe-deficient Arabidopsis roots
(Khan et al., 2019). We report here that 25 of the colonies (representing 25 %) obtained in the
yeast two-hybrid (Y2H) screen carried a fragment of the coding sequence of the gene
AT2G18390, encoding the ARL-type small GTPase TTN5. We verified the interaction in a
targeted Y2H test (Figure 1A). Small GTPases act as molecular switches in cells, like it is the
case for TTN5 (Figure 1B) (Mohr et al., 2023). We further verified the protein interaction in plant
cells using Bimolecular Fluorescence Complementation (BiFC) as a reconstitution of YFP in
positively transformed cells that expressed the control marker mRFP (Figure 1C). We
generated mutants of conserved amino acids in the GTP-binding pocket, Thr30-to-Asn (T30N,
previously found to be dominant-negative and GDP-bound) and Gln70-to-Leu (Q70L,
characterized as dominant-active and GTP-bound) (Figure 1B) (Mohr et al., 2023). Both, the
dominant-negative cYFP-TTN5^{T30N} and active cYFP-TTN5^{Q70L} forms interacted with IRT1vr
(Figure 1D-E). The BiFC interactions were specific as neither protein partner was able to bind
negative controls in this assay (Figure 1F-G). We further elucidated whether IRT1vr might
discriminate between active and inactive TTN5 variant forms using a quantitative analysis of
the interaction by Förster Resonance Energy Transfer-Acceptor Photobleaching (FRET-APB).
This approach is based on close-proximity-dependent energy transfer of an excited GFP-
tagged TTN5 donor to a mCherry acceptor, here IRT1vr-mCherry. Energy transfer was
quantified as FRET efficiency (Figure 1H-J). FRET efficiencies between all GFP-TTN5 forms
and IRT1vr-mCherry were significantly higher compared to the donor-only sample, confirming

protein interactions between all TTN5 forms and IRT1vr. On the other hand, it still remains to be shown that full-length IRT1 can interact with TTN5 at the membrane, e.g. by *in planta* pull-down experiments of respective tagged proteins, an assay that was previously successful to find hints for full-length IRT1-protein interactions with other peripheral membrane proteins (Khan et al., 2019; Hornbergs et al., 2022).

In sum, we can conclude that TTN5 is a novel interactor of the cytoplasmic loop IRT1vr. TTN5 is a link in the coordination of IRT1 regulation with vesicle trafficking. Surprisingly, the GTP/GDP-binding conformations did not influence significantly the protein interaction in the case of TTN5 with IRT1vr.

TTN5 affects Fe homeostasis

Following up on the IRT1-TTN5 interaction, we assumed that TTN5 might affect the regulation of the Fe deficiency response. Homozygous TTN5 knock-outs are embryo-lethal (Mayer et al., 1999; McElver et al., 2000), and heterozygous plants (*ttn5-1$^{+/-}$*) showed no obvious developmental defects in comparison with wild type sibling plants (Figure 2A). Root length increased under Fe deficient conditions but did not differ between wild type and *ttn5-1$^{+/-}$* siblings, and chlorophyll content was not affected (Figure 2B-C). *TTN5* expression was nearly 50 % reduced in *ttn5-1$^{+/-}$* compared to wild type siblings, consistent with single allele presence. Very interestingly, we detected a 15 % decrease of *TTN5* gene expression between Fe sufficiency and deficiency in wild type and *ttn5-1$^{+/-}$*, indicating that TTN5 might be perhaps less needed under low Fe supply (Figure 2D). We examined the heterozygous seedlings at the molecular and physiological level. No difference of gene expression levels of *FRO2* and *IRT1* occurred in *ttn5-1$^{+/-}$* roots compared with wild type (Figure 2E-F). This indicated that root Fe deficiency responses might not be compromised in *ttn5-1$^{+/-}$*. However, we then noted that in two out of three experiments (each including the indicated biological replicates), Fe reductase activity was altered. It was up-regulated under Fe-deficient versus -sufficient conditions in wild type and *ttn5-1$^{+/-}$* roots, but upon Fe deficiency, Fe reductase activity was reduced in *ttn5-1$^{+/-}$* compared with wild type siblings (Figure 2G).

Taken together, heterozygous *ttn5-1$^{+/-}$* seedlings have an Fe reductase phenotype that indicates a positive effect of TTN5 at the physiological level of Fe acquisition. This was interesting, because two other interactors of IRT1vr had an opposite effect on Fe reductase activity. The C2 domain protein EHB1/CAR6 had a negative effect and also the SEC14 protein PATELLIN 2 (PATL2) (Khan et al., 2019; Hornbergs et al., 2022).

TTN5 colocalizes with IRT1 in endocytosed plasma membrane material

To investigate how TTN5 may interact with IRT1, we compared localization of IRT1-mCitrine (proIRT1::IRT1-mCitrine/*irt1-1*) (Dubeaux et al., 2018) with YFP-TTN5 (pro35S::YFP-TTN5) in the root. The YFP-TTN5 fusion protein was found to be localized at the plasma membrane and in the endomembrane system, where IRT1 is present, in a similar manner as immunolocalization signals of a complementing HA$_3$-TTN5 protein (Mohr et al., 2023). 6-day old transgenic Arabidopsis plants of the TTN5 and IRT1 reporter fusion lines were grown side-by-side (Supplemental Figure S1). As previously described, IRT1-mCitrine was present at the plasma membrane of root epidermis cells in our growth system (Supplemental Figure S1A) (Dubeaux et al., 2018; Hornbergs et al., 2022). The same was the case for YFP-TTN5 (Supplemental Figure S1B), corresponding to the previously identified pattern (Mohr et al.,

2023). Virtual cross sections of both constructs highlighted the polar expression of IRT1-
mCitrine (Supplemental Figure S1C) compared with an equally distributed YFP-TTN5 in the
epidermis (Supplemental Figure S1D). Next to plasma membrane localization both, IRT1-
mCitrine and YFP-TTN5 were expressed in vesicle-like structures in root cells (Supplemental
Figure S1E-F). These findings suggested that IRT1 and TTN5 may locate to similar structures
inside root cells, supporting an interaction at these places.
To confirm the obtained Arabidopsis data and to investigate potential colocalization of TTN5
with IRT1, we used the transient expression system of *Nicotiana benthamiana*. We created a
constitutively expressing GFP-tagged IRT1 with GFP inserted inside the cytoplasmic loop
(hereafter termed IRT1-GFP). This IRT1 fusion protein was present at the plasma membrane.
Fluorescent spots were also found in close proximity to the plasma membrane (Supplemental
Figure S2A), consistent with previous reports (Barberon et al., 2011; Ivanov et al., 2014). To
further evaluate its usefulness for localization studies, we compared IRT1-GFP signals with
already described C-terminally-tagged IRT1-mCherry signals in transformed *N. benthamiana*
cells (Ivanov et al., 2014) before and after wortmannin treatment. Wortmannin is a fungal
metabolite with an inhibitory effect on phosphatidylinositol-3-kinase (PI3K) function, leading to
swollen multivesicular bodies (MVBs) and therefore is commonly used to investigate
endocytosis (Cui et al., 2016). The two constructs showed good colocalization and, following
wortmannin treatment, could be observed among others in swollen structures, suggesting
partial presence in the MVBs (Supplemental Figure S2B-C), indicating that the new IRT1-GFP
construct was suited for IRT1 localization studies. IRT1-GFP complemented an early rosette
growth deficit of *irt1-1* (in pro35S::IRT1-GFP/*irt1-1* plants), indicating functionality of the IRT1-
GFP construct for this trait (Supplemental Figure S2D).
We detected colocalization of IRT1-GFP and the three different mCherry-TTN5 constructs at
the plasma membrane (Figure 3A-C). We found YFP-TTN5 localization in vesicle-like
structures and close proximity of YFP-TTN5 to the plasma membrane, as previously reported
(Mohr et al., 2023). YFP-TTN5^{T30N} is less mobile than TTN5 and TTN5^{Q70L} (Mohr et al., 2023).
Further investigation revealed a significant colocalization difference between the inactive and
active variant of TTN5 with IRT1 (Supplemental Figure S2E). While Pearson coefficients of
IRT1-GFP with mCherry-TTN5 and mCherry-TTN5^{Q70L} were 0.77 and 0.80 respectively, it was
0.90 for IRT1-GFP and mCherry-TTN5^{T30N}, demonstrating higher general colocalization
between IRT1-GFP and the low-mobile low-activity TTN5 form (Supplemental Figure S2E;
similar results for overlap coefficients). All three mCherry-TTN5 constructs were also localized
in IRT1-GFP-positive structures close to the plasma membrane, suggesting the two proteins
might colocalize in plasma membrane-derived vesicles (Figure 3D-F). Surprisingly, the
Pearson coefficients changed when these spotted structures were analyzed (Supplemental
Figure S2F; similar results for overlap coefficients). While the values for mCherry-TTN5 (0.80)
and mCherry-TTN5^{Q70L} (0.78) with IRT1-GFP remain relatively the same, we obtained with 0.78
now a similar coefficient with mCherry-TTN5^{T30N}. Similar results were achieved while checking
the overlaps of mCherry-TTN5 structures with IRT1-GFP, with overlaps of 36-41 %
(Supplemental Figure S2G). In contrast, only 54 % of IRT1-GFP-positive structures colocalized
with mCherry-TTN5^{T30N}, whereas 61 % and up to 83 % colocalized with mCherry-TTN5 and
mCherry-TTN5^{Q70L}, respectively. From this, we hypothesized that IRT1 along with dominant
negative TTN5 is more likely to be localized at the plasma membrane while the active variant
may play a role in intracellular cycling. This was consistent with the previous observation that
YFP- TTN5^{T30N} was moving slower in the cell than the other two TTN5 fusion proteins, indicating

that it might have led to a slow-down of IRT1 removal from the plasma membrane. In an attempt to identify the intracellular structures, we treated the cells with wortmannin (Figure 3G). Structures positive for both IRT1-GFP and mCherry-TTN5 became doughnut-shaped suggesting IRT1-TTN5 colocalization in MVBs (Figure 3H-J). Presence of IRT1 in these late endosomal compartments has been observed previously, as part of the pathway to IRT1 vacuolar degradation (Dubeaux et al., 2018), same for YFP-TTN5 (Mohr et al., 2023). Thus, our findings emphasize the involvement of TTN5 in the endomembrane trafficking of the Fe transporter IRT1.

TTN5 interacts with IRT1 regulators EHB1, SNX1 and PATL2

We previously found that IRT1 associates and colocalizes with certain peripheral membrane proteins. To further undermine that TTN5 associates with IRT1, we investigated the possibility that TTN5 also binds these other proteins of the IRT1 interactome in close proximity.

At first, we tested IRT1-interacting C2-domain protein ENHANCED BENDING 1 (EHB1) (Khan et al., 2019) that belongs to the ten-member calcium-dependent C2-DOMAIN ABSCISIC ACID-RELATED (CAR) protein family (EHB1 is also known as CAR6) (Rodriguez et al., 2014). CAR proteins tether signaling and transport proteins involved in drought, defense or iron nutrition to the membrane (Cui et al., 2023). They can also oligomerize and cause membrane deformations, making them candidate proteins involved in vesicle budding (Chen et al., 2023). CAR proteins have a plant-specific insertion of an extra region within the C2 domain known as signature (sig) domain, that is involved in binding proteins (Rodriguez et al., 2014; Khan et al., 2019). EHB1 interacts *via* its sig domain with IRT1vr and it has an inhibitory effect on Fe acquisition (Khan et al., 2019). Interestingly, while EHB1Δsig did not interact with TTN5 in BiFC, the intact EHB1 protein and the EHB1 sig domain alone interacted with TTN5 in BiFC (Figure 4A-C, see also Figure 1F). FRET-APB experiments were used for validation taking EHB1-GFP as the donor sample and all three TTN5 variants as the acceptor with a C-terminal mCherry-tag. EHB1-GFP with TTN5-mCherry yielded a significantly higher FRET efficiency compared to the donor-only sample (Figure 4F). Surprisingly, the FRET pairs EHB1-GFP with TTN5^{T30N}-mCherry and TTN5^{Q70L}-mCherry resulted in an even higher FRET-efficiency compared to TTN5 (Figure 4F).

SORTING NEXIN 1 (SNX1) is a positive regulator of IRT1 recycling at the TGN (Ivanov et al., 2014). TTN5 was able to bind SNX1 (Figure 4G). The active TTN5 GTPase variant interacted with SNX1 (Figure 4I), while the inactive TTN5 mutant failed (Figure 4H). Sorting nexins are classified by their characteristic domains. SNX1 represents the class I SNX proteins, characterized by the presence of PHOX-homology (PX) and Bin-Amphiphysin-Rvs (BAR) domains (Heucken and Ivanov, 2018). Such a domain organization is typical for proteins that function in membrane and phosphoinositide binding by the PX domain, whereas the BAR domain is involved in membrane curvature sensing and the formation of endosomal tubules (Peter et al., 2004; Frost et al., 2009). Interestingly, ARF-GAPs which are needed for the GTP hydrolysis, contain a BAR domain as well (Memon, 2004). We found evidence that TTN5 binds to both the SNX1 PX and BAR domains using BiFC (Figure 4J-K).

Lastly, plasma membrane-localized SEC14 protein PATELLIN 2 (PATL2) binds IRT1vr (Hornbergs et al., 2022). The SEC14 domain is a lipid-binding pocket that serves to present or transfer lipophilic substances at membranes (Montag et al., 2023). Thereby, PATL2 is a lipid transfer protein that may connect oxidative stress prevention and Fe acquisition, possibly by

transferring and exchanging phospholipids and α-tocopherol at the membrane. PATL2 may
also initiate vesicle formation (Hornbergs et al., 2022). We tested interaction ability of cYFP-
TTN5 with nYFP-PATL2 in BiFC experiments. YFP signals indicating complementation and
interaction were visible at the plasma membrane, fitting to the localization studies of both
PATL2 and TTN5 (Figure 4L). We were not able to observe complementation for the
combination of PATL2 and any of the two GTPase variants TTN5^{T30N} (Figure 4M) and TTN5^{Q70L}
(Figure 4N). This was very interesting, as it indicates that the TTN5 switching mechanism may
be required for a potential interaction between these two proteins. To further identify the region
of interaction, we tested already published deletion mutants of PATL2 (Montag et al., 2020).
PATL2 can be divided into two protein halves, consisting of the N-terminal part and the
conserved C-terminal CRAL-TRIO-N-terminal extension (CTN), SEC14 and Golgi dynamics
(GOLD) domains of PATL2 (Montag et al., 2020). The N-terminal part is needed for the
interaction with IRT1vr (Montag et al., 2020). We used the PATL2 deletion mutants nYFP-
PATL2ΔC lacking the three consecutive CTN, SEC14-GOLD domain and nYFP-PATL2ΔN
lacking the IRT1vr-interacting N-terminal part. nYFP-PATL2ΔC together with cYFP-TTN5
complemented YFP (Figure 4O), whereas no signal was obtained by nYFP-PATL2ΔN in
combination with cYFP-TTN5 (Figure 4P), similar as in the case of IRT1vr (Hornbergs et al.,
2022). Interestingly, the fluorescent complementation signal between cYFP-TTN5 and nYFP-
PATL2ΔC was no longer present at the plasma membrane consistent with the observation that
the PATL2 plasma membrane localization is dependent on C-terminal PATL2 domain region
(Montag et al., 2020).
We also analyzed *EHB1, SNX1* and *PATL2* gene expression. All genes were expressed in
similar levels in both wild type and *ttn5-1*$^{+/-}$ (Supplemental Figure S3A-C) indicating that there
was no feedback at the level of regulation of genes encoding the TTN5 interactors.
In summary, we could show that TTN5 is not only interacting with IRT1vr but also with three
peripheral membrane proteins that can be in the vicinity of IRT1 at membrane sites in root cells.
This suggests formation of a protein complex at the IRT1vr regulatory platform in which TTN5
may have a coordinating role due to its switch mechanism.
328

Discussion

Here, we show that the small GTPase TTN5 links Fe homeostasis with the endomembrane
trafficking system (Figure 5). We demonstrate that TTN5 interacted with the large intracellular
loop and variable region of IRT1 and colocalized with IRT1 in plant cells at the plasma
membrane and in vesicle structures. TTN5 had a positive effect on root iron reduction. *TTN5*
gene expression was found to be slightly lower upon Fe deficiency than sufficiency. TTN5
interacted with peripheral membrane proteins of the IRT1 interactome, EHB1, PATL2 and
SNX1. Hence, our findings shed new light on a post-embryonic physiological ARL GTPase
function in plant nutrition. Binding of TTN5 directly to the cargo IRT1 *via* IRT1vr indicates
specificity of TTN5 function beyond housekeeping roles. The concert with Ca^{2+}-dependent and
oxidative stress-related proteins, EHB1 and PATL2, as well as protein sorting by SNX1 indicate
that TTN5 links dynamic plasma membrane and vesicle trafficking responses with Fe regulation
to control an iron transporter. Nucleotide exchange and GTPase activities of TTN5 were
important for several of the mentioned activities since respective mutants of TTN5 showed
partly different behavior from TTN5.
344

TTN5 is part of the endomembrane system, targeting IRT1vr

The interaction between TTN5 and IRT1vr is supported by multiple findings. First, the interaction was detected in a Y2H screen, that yielded only few interaction partners, which could all be validated. Second, the interaction was verified by using independent methods (targeted Y2H, BiFC and FRET-APB). Third, TTN5 and IRT1 reporter proteins colocalized in cells, and their localization overlapped in intracellular membrane-associated locations in cells of the root epidermis, in which both genes and proteins are expressed and present according to transcriptome studies and promoter reporter lines. Further, TTN5 was found to interact with proteins of the IRT1vr environment (Figure 5A-B). These findings altogether indicate that in plant roots, TTN5 and IRT1 indeed can be present to interact in root epidermis cells in places related with Fe acquisition. Since TTN5 is involved in vesicle trafficking, like other members of the ARF family of small GTPases (Memon, 2004; Sztul et al., 2019) it has diverse localization patterns within the cell. The previous TTN5 localization (Mohr et al., 2023) was confirmed upon colocalization with IRT1 in this study. Hence, IRT1 localization dependent on endomembrane trafficking may be linked with the presence of TTN5 in this process.

A very interesting finding was that the dominant-negative mCherry-TTN5^{T30N} variant tended to colocalize more at the plasma membrane with IRT1-GFP while intracellular vesicle-like structures of IRT1-GFP overlapped more with the active mCherry-TTN5 variant. This is consistent with our colocalization data with the Golgi, MVB and plasma membrane markers (Mohr et al., 2023) and reinforces our suggestion that TTN5 has functions in vesicle trafficking. Since TTN5^{T30N} is predicted to strongly bind a GEF, we can suppose that TTN5^{T30N} presence at the plasma membrane might be linked with a TTN5-interacting GEF at this site. The presence of mCherry-TTN5 together with IRT1-GFP in wortmannin-induced swollen structures, probably indicating MVBs, is supporting the idea of a role in transporter degradation. MVBs are known to be the last step before vacuolar degradation (Kolb et al., 2015; Arora and Van Damme, 2021). This could indicate that TTN5 presence is a critical point of IRT1 sorting fate. We therefore propose a role of TTN5 in targeting intracellular IRT1 protein (Figure 5B). In this process, TTN5 is present at the plasma membrane in a resting state. In the move of GTPase activation, it could be recruited to the place of action and active TTN5 would play a role in IRT1 cycling. TTN5 should predominantly bind GTP and be in an active state in the cell (Mohr et al., 2023). Further, there is support of a role of TTN5 in IRT1 recycling by the binding of TTN5 and SNX1. SNX1 is involved in the recycling of IRT1 at these sites (Ivanov et al. 2014). Hence, TTN5 can be in a functional complex with SNX1 to control the recycling of the cargo protein IRT1.

The observed roles of TTN5 fit well with the general understanding of ARF-signaling in plasma membrane protein trafficking. The auxin transporters PIN1 and PIN2 and the boron transporter BOR1, like IRT1, are polar localized in the plasma membrane and undergo a constant degradation-recycling process (Benková et al., 2003; Takano et al., 2010; Barberon et al., 2014). Apart from vesicle formation, as described for ARF1 or SAR1, a role in this process was shown for other regulatory mechanisms of ARF-signaling through the GEFs and GAPs (Nielsen, 2020). GNOM and BIG5, two ARF-GEFs, and ARF1-regulators, are described to be involved in the proper localization of the plasma membrane transporter PIN1, functioning at different regulatory steps (Geldner et al., 2003; Anders et al., 2008; Kleine-Vehn et al., 2008; Tanaka et al., 2009). In addition, another ARF GTPase ARF1A1C may not only partially colocalize with GNOM, but modified activity also resulted in impaired localization of PIN1 to the plasma membrane and also altered vacuolar targeting of PIN2 (Tanaka et al., 2014). BOR1

plasma membrane localization depends on GNOM as well (Yoshinari et al., 2021), showing a good connection between ARF-GEFs and plasma membrane protein localization.

A limitation in our study was that we were not able to show TTN5 interaction with full-length IRT1 by coimmunoprecipitation. This might have been due to a transient interaction. Alternatively, fluorescent protein-based interactions approaches are often limited by technical issues, e.g. a spatial distance of the fluorescent proteins. Perhaps the locations of the tags can be problematic for certain interactions like that of TTN5 and IRT1.

Taken together, we could not only identify TTN5 as part of the endomembrane system but also uncover another link between ARF signaling and plasma membrane protein trafficking by its interaction with IRT1vr. Future identification of potential TTN5-GEF and -GAP proteins will be of great interest to clarify the role of TTN5 in endomembrane trafficking with particular reference to IRT1 regulation. For future studies, stable coexpression of both proteins under the control of the endogenous promoter will be suitable to investigate the effect of TTN5 on IRT1.

TTN5 might play a general role in Fe nutrition

TTN5 does not only have an influence on IRT1 but rather potentially possesses a general role in Fe nutrition. A partial lack of TTN5 results in decreased Fe reductase activity. It was already investigated that IRT1 can interact with FRO2 and AHA2 and potentially form a complex for a highly effective Fe uptake machinery (Martín-Barranco et al., 2020). Reduced Fe reductase activity could be due to an indirect influence of TTN5 on the complex formation or it could be directly driven by TTN5. Interestingly, both *ehb1* and *patl2* seedlings have a higher Fe reductase activity under Fe deficiency compared to wild type suggesting an opposing role of TTN5 on the Fe deficiency response. Gene expression analysis revealed no great impact on Fe-related genes in *ttn5-1*$^{+/-}$ roots, implying an effect of TTN5 mostly on posttranslational level.

It is a possibility that TTN5 has very different effects depending on its localization in the cell. For example, the effect on Fe reduction could be different from its effect on colocalization with IRT1 in MVBs. Different types of interaction partners were reported for the human homolog ARL2 with connection to distinct cellular functions in tubulin folding and microtubule dynamics (Bhamidipati et al., 2000) or in mitochondria (Sharer et al., 2002). Based on these distinct locations and the associated functions of ARL2, it was proposed that ARL2 may act at a high level in signaling (Francis et al., 2016). This can be also the case in Arabidopsis. For example, TTN5 may bridge TTN5 microtubule function (Mayer et al., 1999) with our here-found role in IRT1 regulation.

TTN5 interacts with EHB1, SNX1 and PATL2 and may coordinate IRT1 regulation

It is already known that IRT1 regulation is a process of high complexity which is not only influenced by Fe-sufficient or -deficient conditions, but also by other metals such as zinc and manganese or Ca^{2+}- and ROS-signaling among others (Dubeaux et al., 2018; Brumbarova and Ivanov, 2019; Khan et al., 2019; Gratz et al., 2021).

Interaction with the IRT1 regulators EHB1, PATL2 and SNX1 suggests that TTN5 could not only be linked with Fe nutrition but also the associated Ca^{2+}- and ROS-signaling events, previously studied for EHB1 and PATL2 (Khan et al. 2019; Hornbergs et al. 2022). We identified interaction with EHB1 by two independent methods (BiFC and FRET-APB) and with PATL2 and SNX1 by BiFC. The interactions are further supported by deletion mutant approaches. The interacting sites of PATL2 and EHB1 with TTN5 were similar to those for IRT1vr, supporting

again the possibility that these domains and motifs are at the basis of a protein complex forming
at the IRT1vr regulatory platform.
Fe deficiency leads to an increase of cytosolic Ca^{2+} (Gratz et al., 2019; Matthus et al., 2019).
Physiological data indicate that CBL1/9-dependent CIPK23 activation is promoting an induction
of the Fe reductase activity (Tian et al., 2016). Contrary to this, CBL1/9 are also responsible
for potential IRT1 endocytosis by CIPK23-dependent phosphorylation and the transporter
activity is inhibited by Ca^{2+}-induced recruiting of EHB1 to the plasma membrane (Dubeaux et
al., 2018; Khan et al., 2019), suggesting opposing effects of Ca^{2+} in the Fe deficiency response.
EHB1 is additionally interesting as a member of the CAR family. Several CAR proteins can
initiate Ca^{2+}-dependent membrane curvature, which may provide a direct link to vesicle
trafficking. Interestingly, Ca^{2+} is bound by EHB1 *via* its C2 domain, which is also a common
feature of class III ARF-GAPs (Vernoud et al., 2003; Knauer et al., 2011). Indeed, EHB1 is
directly linked to ARF signaling in an antagonistic interaction with ARF-GAP AGD12 in
gravitropic bending (Dümmer et al., 2016; Rath et al., 2020). Further research needs to
investigate whether TTN5 can interact with the C2-domain containing ARF-GAPs AGD11-
AGD13 as well. Additionally, crystallization experiments revealed a potential Ca^{2+}-binding site
between HsARF6 and its GAP ASAP3. The GTP-hydrolysis was stimulated in the presence of
Ca^{2+} proposing a linkage between Ca^{2+}- and ARF-signaling (Ismail et al., 2010). We therefore
consider the possibility of an interaction between TTN5 and EHB1 in a Ca^{2+}-dependent manner.
PATL2 may interact with IRT1 to prevent lipid peroxidation stress (Hornbergs et al., 2022).
Prolonged Fe deficiency leads to an increase in hydrogen peroxide concentration as well as an
activity increase of catalase CAT2 (Le et al., 2015; von der Mark et al., 2020; Gratz et al., 2021).
Some catalases are present in the PATL2 interactome in roots (Hornbergs et al., 2022). Future
studies may address whether TTN5 is connected with oxidative stress.
Next to EHB1 and PATL2, we identified SNX1 as a positive regulator for IRT1 recycling to
the plasma membrane in the sorting endosomes (Ivanov et al., 2014). In addition to IRT1,
endosomal recycling of the auxin efflux carrier PIN2 is also dependent on SNX1 (Jaillais et al.,
2006). SNX1 is therefore part of the recycling of specific plasma membrane proteins.
Interestingly, deficiency of BLOS1 leads to accumulation of PIN1 and PIN2 at the plasma
membrane (Cui et al., 2010). BLOS1 is one of the homologs to the mammalian biogenesis of
lysosome-related organelles complex 1 (BLOC-1), which is responsible for vesicle transport
from endosomes to lysosomes (Li et al., 2007; Raposo et al., 2007). BLOS1 could therefore
perform a similar function to BLOC-1 in Arabidopsis. SNX1 can interact with both BLOS1 and
BLOS2 and could therefore also play a role in vesicle-mediated transport through these
interactions (Cui et al., 2010; Heucken and Ivanov, 2018). SNX1 possesses a BAR domain
which is also typical for class I ARF-GAPs (Vernoud et al., 2003; Heucken and Ivanov, 2018).
A connection between the SNX1 BAR domain and an ARF6-GEF was identified in mouse by
their interaction. This interaction was associated with a positive effect of ARF6 function (Fukaya
et al., 2014). Colocalization analysis revealed overlapping expression in endosomes, which fits
known SNX1 localization in Arabidopsis (Fukaya et al., 2014; Ivanov et al., 2014), this could be
a possible interconnection between ARF-signaling and SNX1. Here, we identified an interaction
between TTN5 and SNX1, which could be the critical link in the decision between SNX1-
dependent recycling and degradation of IRT1. SNX1 could be needed for a potential TTN5-
GEF and subsequent TTN5 recruiting to membranes. Based on the opposing roles of EHB1
and SNX1 in IRT1 regulation and their here-described connection to TTN5, we suggest a
coordinating role of TTN5 in transporting IRT1 between plasma membrane and endosomes by

interacting with its regulators such as EHB1, PATL2 and SNX1. The findings that some ARF-
GAPs have a C2 or BAR domain, and that SNX1 interacts with an ARF-GEF strengthen the
interest in the identification of TTN5-GEFs and -GAPs.

Further research has to focus on physiological aspects in either combined knockout mutants
or tagged-protein lines to arrange and decipher the IRT1 regulatory network. An intriguing open
question is whether a large IRT1vr-EHB1-PATL2-TTN5-SNX1 complex is assembled. If so, the
next question is how, when and in which order the individual or all components of the IRT1vr
complex are assembled. Another question is which are the structural requirements for proper
protein conformations and membrane interfaces (Figure 5A, B).

Conclusion and perspectives

491
In this study, we identified the ARF-like GTPase TTN5 as an interactor of the iron transporter
IRT1 and three of its known regulators, SNX1, PATL2 and EHB1, suggesting that they form a
complex at the IRT1vr site. One role of TTN5 and perhaps the entire identified IRT1vr complex
including TTN5 is likely to contribute to endocytosis and recycling of IRT1. The proposed model
of complex formation to affect IRT1 localization at the plasma membrane can be studied further
in the future. TTN5 manipulation can show whether indeed TTN5 affects IRT1 endocytosis and
degradation or its recycling. Furthermore, certain physiological effects may facilitate that a
complex forms composed of TTN5 with PATL2, EHB1 and SNX1, also in combination with
IRT1. An intriguing question is whether TTN5 binds specifically with IRT1 or also targets other
transmembrane cargo proteins. A challenging task is to obtain proofs of TTN5 protein
complexes of this kind in roots and determine the requirements and order of events for complex
assembly. Finally, due to the very low intrinsic GTP hydrolysis activity of TTN5, identification of
TTN5 GAPs will shed light on TTN5 function in Fe homeostasis and potential other cellular
signaling events. Likewise, it will be interesting to identify a membrane-localized GEF of TTN5.

Material & Methods

Yeast two-hybrid (Y2H)

507
The yeast two-hybrid screen for identification of IRT1vr interactors was performed previously
and is described in Khan et al. (2019). Targeted Y2H interactions were tested using the GAL4
transcription factor system with histidine as selection marker. *TTN5* coding sequence was
amplified using TITAN5 nter B1 and TITAN5 stop B2 primers (for primer sequences, see
Supplemental Table S1). Obtained fragment was cloned into pDONR207 *via* Gateway BP
reaction with following LR reactions into pGBKT7-GW (binding domain, BD) or pACT2-GW
(activation domain, AD). Both vectors were a kind gift from Dr. Yves Jacob. The yeast strain
AH109 was cotransformed with the corresponding AD- and BD-tagged expression vectors. The
combination pACT2:SNX1 and pGBKT7:SNX1 (Pourcher et al., 2010) was used as a positive
control. Negative controls were cotransformations with the respective non-recombined pACT2-
GW or pGBKT7-GW. Drop test was performed with SD -Leu, -Trp-cultures at an OD$_{600}$ of 0.1
and three additional 1:10 dilution steps (10^1-10^4) on SD -Leu, -Trp- and SD -Leu, -Trp, -His +
0,5 mM 3-amino-1,2,4-triazole-plates. One of each plate was incubated either at 30°C or at
room temperature, wrapped in aluminum foil for up to two weeks.

Arabidopsis plant material and growth conditions

522
The Arabidopsis *ttn5-1* mutant was previously described (McElver et al., 2000).
Heterozygous seedlings were selected by PCR on gDNA using the primer TTN5 intron1 fwd
and pDAP101 LB1 (Supplemental Table S1). Arabidopsis seeds were sterilized with sodium

hypochlorite solution (6 % Sodium hypochlorite and 0.1 % Triton X-100) and stored 24 hours
at 4°C for stratification. Seedlings were grown upright on Hoagland media [1.5 mM Ca(NO$_3$)$_2$,
0.5 mM KH$_2$PO$_4$, 1.25 mM KNO$_3$, 0.75 mM MgSO$_4$, 1.5 µM CuSO$_4$, 50 µM H$_3$BO$_3$, 50 µM KCl,
10 µM MnSO$_4$, 0.075 µM (NH$_4$)$_6$Mo$_7$O$_{24}$, 2 µM ZnSO$_4$ and 1 % sucrose, pH 5.8, supplemented
with 1.4 % Plant agar (Duchefa)] with sufficient (50 µM FeNaEDTA, +Fe) or deficient (0 µM
FeNaEDTA, -Fe) Fe supply in growth chambers (CLF Plant Climatics) under long day condition
(16 hours light at 21°C, 8 hours darkness at 19°C). Seedlings were grown in a two-week growth
system with plants growing 14 days on Fe-sufficient media and then transferred to either fresh
Fe-sufficient or Fe-deficient media for additional three days.
Nicotiana benthamiana plants were grown on soil for 2-4 weeks in a greenhouse facility
under long day conditions (16 hours light, 8 hours darkness).

Root length

538   For root length comparison plants were growing for 10 days in Fe-sufficient or -deficient
539 conditions photographed including a scale in the picture. Root length were measured from root
540 tip to hypocotyl using the JMicro Vision program (Roduit, N. JMicroVision: Image analysis
541 toolbox for measuring and quantifying components of high-definition images. Version 1.3.4.
542 https://jmicrovision.github.io) (+Fe: WT n = 19; *ttn5-1* n = 28; -Fe: WT n = 16; *ttn5-1* n = 32).

Chlorophyll content

544   For chlorophyll content comparison plants were grown in the two-week system in Fe-
545 sufficient or -deficient conditions. Chlorophyll content was measured by SPAD values using
546 SPAD-502plus (Konica Minolta Optics, 2012) with up to ten biological replicates (+Fe: n = 10;
547 -Fe: n = 9). Chlorophyll content was determined in three individual experiments.

Fe reductase activity assay

Fe reductase activity assay was performed as described in (Gratz et al., 2019). For Fe
reductase activity plants were grown in the two-week system in Fe-sufficient or -deficient
conditions. Two plants per replicate were washed in 100 mM Ca(NO$_3$)$_2$, then incubated for
1 hour in the dark in 1.5 ml Fe reductase solution (300 mM ferrozine and 100 mM FeNaEDTA).
The Fe reductase activity-dependent color change was measured at 562 nm using Infinite
200® PRO (Tecan) plate reader. Activity calculation was done using the ferrozine extinction
coefficient ε = 28.6 mM^{-1}*cm^{-1} and was normalized to root weight. The assay was performed
with a minimum of three replicates (n ≥ 3), each consisting of a pool of two plants. Fe reductase
activity was determined in three individual experiments.

Rosette size

559   For rosette size measurements plants were grown in parallel for one week in Fe-sufficient
560 conditions on Hoagland plates before transferring them to soil. 14 days after sowing rosette
561 sizes was measured using ImageJ (Schneider et al., 2012). The size was determined between
562 tips of opposite leaves of twelves plants each (n = 12).

Gene expression analysis by RT-qPCR

For gene expression analysis, plants were grown in the two-week system. Total RNA was
isolated from roots using the RNeasy Mini Kit (Qiagen) and cDNA synthesis was performed
with the RevertAid RT Reverse Transcription Kit (ThermoFisher Scientific). The SFX96
Touch^{TM}Real-Time PCR Detection System (Bio-Rad) was used for RT-qPCR performance.
Data processing was done with Bio-Rad SFX Manager^{TM} (version3.1) software. Mass standard
curve analysis was used for determination of the absolute gene expression and the elongation

factor *EF1Bα* expression served as a reference for normalization. RT-qPCR was performed in
three biological replicates (n = 3) and two technical replicates each. All primer pairs used in this
study are listed in Supplemental Table S1.

Nicotiana benthamiana leaf infiltration

N. benthamiana leaf infiltration was performed with the Agrobacterium (*Rhizobium*
radiobacter) strain C58 (GV3101) carrying the respective constructs for confocal microscopy.
Agrobacteria cultures were grown overnight at 28°C, centrifuged for 5 min at 4°C at 5000*g*,
resuspended in infiltration solution (5 % sucrose, a pinch of glucose, 0.01 % Silwet Gold,
150 µM acetosyringone) and incubated for 1 hour at room temperature. Bacterial suspension
was set to an OD_{600}=0.4 and infiltrated into the abaxial side of *N. benthamiana* leaves.
Expression of pABind and pMDC7 (Curtis and Grossniklaus, 2003; Bleckmann et al., 2010)
constructs was induced with a β-estradiol solution (20 µM β-estradiol, 0.1 % Tween 20) 16
hours before imaging.

Subcellular localization of fluorescent protein fusions

The creation of TTN5 dominant negative (T30N) and dominant active (Q70L) constructs by
introduced point mutations and generation of pro35S::YFP-TTN5 is described in (Mohr et al.,
2023). mCherry-tagged constructs were created by overlap-extension PCR. Fluorescent
protein was amplified from pABind:mCherry (Bleckmann et al., 2010) using primer pair
mCherry B1 and mCh R ns BIND. *TTN5*, *TTN5^{T30N}* and *TTN5^{Q70L}* CDS were amplified with a
20 bp overlap of mCherry using mCh to TTN5 and TITAN5 stop B2. In a third PCR mCherry-
tagged TTN5 constructs were generated with the mCherry B1 and TITAN5 stop B2 primer
pair. BP and LR reaction were performed as described above. As destination vector the XVE-
driven β-estradiol inducible promoter driven vector pMDC7 (Curtis and Grossniklaus, 2003)
was chosen. IRT1-GFP construct, with the GFP at position 173 in the variable, cytoplasmic
loop, was cloned by overlap-extension PCR. *GFP* was amplified from pABind:GFP
(Bleckmann et al., 2010) using primer pair IG173 F and GI173 R. IRT1 halves were amplified
using IRT1 B1 and IG173 R and GI173 R and IRT1 stop B2. In a last PCR IRT1-GFP was
amplified with primer pair IRT1 B1 and IRT1 stop B2. BP and LR reaction were performed as
described above with pMDC32 (Curtis and Grossniklaus, 2003) as destination vector. Cloning
of IRT1-mCherry construct is described in Ivanov et al., (2014) and proIRT1::IRT1-
mCitrine/*irt1-1* in Dubeaux et al., (2018). Agrobacteria were transformed with the obtained
constructs and tested by *N. benthamiana* leaf infiltration after two days of expression.
Localization studies were carried out by laser-scanning confocal microscopy (LSM 780,
Zeiss) with a 40x C-Apochromat water immersion objective. YFP constructs were excited at
488 nm and detected at 491-560 nm. mCherry or FM4-64 fluorescence was excited at 561 nm
and detected at 570-633 nm.

Chemical treatments

607   Wortmannin (10 µM, Sigma-Aldrich) and plasma membrane dye FM4-64 (165 µM,
608   ThermoFisher Scientific) were infiltrated into *N. benthamiana* leaves. FM4-64 was detected
609   after five min incubation and Wortmannin was incubated for 25 min before checking the
610   treatment effect.

JACoP based colocalization analysis

612   Colocalization analysis was performed with the ImageJ (Schneider et al., 2012) Plugin Just
613   Another Colocalization Plugin (JACoP) (Bolte and Cordelières, 2006). A comparison of

Pearson's and Overlap coefficients was done. Object-based analysis was performed for spotted-structures, adapted by (Ivanov et al., 2014). Percentage of colocalization for both channels was calculated based on distance between geometrical centers of signals. Analysis was conducted in three replicates each (n = 3).

Bimolecular Fluorescence Complementation (BiFC)

The interaction studies of TTN5 and the respective GTPase dominant active (TTN5^{Q70L}) or dominant negative (TTN5^{T30N}) variants *in planta* were performed by using BiFC. *TTN5*, *TTN5^{Q70L}* and *TTN5^{T30N}* coding sequences with stop codon were amplified using the primers TITAN5 n-ter B1 and TITAN5 stop B4 (Supplemental Table S1).

IRT1vr coding sequence with stop codon was amplified using the primers I1L1B3 and I1LB2. Amplifying *EHB1* the primer pair EHB1 n-ter B3 and EHB1 stop B2 was used. *SNX1* coding sequence without stop codon for C-terminal fusion was amplified using the primer SNX1 B3 and SNX1 ns B2. Cloning of SNX1 deletion mutants SNX1PX and SNX1BAR was done using the primers SNX1 B3 and PXSNX1 rev and BARSNX1 fwd and SNX1 ns B2 respectively. The used PATL2 constructs are described in (Hornbergs et al., 2022).

The amplified PCR products were cloned *via* Gateway BP reaction (Life Technologies) into pDONR221-P1P4 (Invitrogen) or pDONR221-P3P2 (Invitrogen). The obtained constructs were used in an LR reaction (Life Technologies) for cloning into pBiFCt-2in1 vectors (Grefen and Blatt, 2012). Agrobacteria were transformed with correct constructs and used for *N. benthamiana* leaf infiltration. After 48 hours, the mRFP expression control signal and YFP signals were detected by fluorescent microscopy (LSM 780, Zeiss) with a 40x C-Apochromat water immersion objective. YFP constructs were detected at 491-560 nm after exciting at 488 nm and mCherry fluorescence was excited at 561 nm and emission detected at 570-633 nm.

The BiFC constructs were tested in three independent replicates with three infiltrated leaves each. The vectors pBiFC-2in1-NN and pBiFC-2in1-CN were kindly provided by Dr. Christopher Grefen, Tübingen, Germany.

Förster-Resonance-Energy Transfer Acceptor Photo Bleaching (FRET-APB)

To verify protein-protein interactions of TTN5 and the corresponding GTPase dominant active (TTN5^{Q70L}) and dominant negative (TTN5^{T30N}) variants the FRET-APB was used. The coding sequences without stop codon were amplified with the primers TITAN5 B1 and TITAN5 ns B2, and cloned into the pDONR207 (Invitrogen, BP reaction, Life Technologies). Cloning of pDONR207:IRT1vr and pDONR207:EHB1 is described in Khan et al. (2019). The obtained constructs were used for LR reactions (Life Technologies) with the pABind-GFP, pABind-mCherry and pABind-FRET for C-terminal tagging (Bleckmann et al., 2010). N-terminal GFP-TTN5 constructs were cloned by overlap extension PCR simultaneously with mCherry-TTN5 constructs (check paragraph 'Subcellular localization of fluorescent protein fusions'). In the first step the primer pairs GFP B1 with GFP R ns BIND and GFP to TTN5 with TITAN5 stop B2 were used with pABind-FRET (Bleckmann et al., 2010) as template for the fluorescent protein coding sequence. The final construct was obtained with the primer pair GFP B1 with TITAN5 stop B2. Constructs were cloned into pDONR207 for further cloning into pMDC7 *via* LR reactions. Agrobacteria were transformed and used for *N. benthamiana* leaf infiltration. FRET-APB was performed using laser-scanning confocal microscopy (LSM 780, Zeiss) with a 40x C-Apochromat water immersion objective. GFP was excited at 488 nm and detected at 491-560 nm and mCherry fluorescence was excited at 561 nm and detected at 570-633 nm. GFP

and mCherry channels were recorded for five scans. The mCherry signal was then bleached
with 70 iterations of maximum laser intensity in a specific region of interest (ROI) which was
set to the nucleus. Both channels were detected for additional 20 post-bleaching scans. FRET-
APB measurements were performed with a minimum of 10 repetitions (n ≥ 10).

Statistical analysis
One-way ANOVA was used for statistical analysis and performed in OriginPro 2019. Fisher
LSD was chosen as post-hoc test with $p < 0.05$.

ACCESSION NUMBERS
Sequence data from this article can be found in the TAIR and GenBank data libraries under
accession numbers: *FRO2* (TAIR: AT1G01580), *EHB1* (TAIR: AT1G70800), *IRT1* (TAIR:
AT4G19690), *PATL2* (TAIR: AT1G22530), *TTN5* (TAIR: AT2G18390), and *SNX1* (TAIR:
AT5G06140).

Acknowledgements

We thank Gintaute Matthäi and Elke Wieneke for excellent technical assistance. We thank
Ksenia Trofimov for microscopic help and advice, and Rubek Merina Basgaran for help with
interaction studies. We are thankful for the excellent assistance from Stefanie Weidtkamp-
Peters and Sebastian Hänsch, members of the Center for Advanced Imaging (CAi) at the
Heinrich Heine University.
This work was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research
Foundation, SFB 1208 project B05 to P.B.). Funding for instrumentation: Zeiss LSM780: DFG-
INST 208/551-1 FUGG.

Figure Legends

Figure 1. TTN5 interacts with IRT1vr in yeast and in plant cells.

(A), Targeted Y2H assay between BD-IRT1vr (IRT1-variable region) and AD-TTN5. Yeast
cotransformations were spotted in 10-fold dilution series ($A_{600} = 10^1\text{-}10^4$) on double selective
(LW; control) and triple selective +0.5 mM 3-AT (LWH, selection for protein interactions) SD-
plates. SNX1/SNX1, positive control (Pourcher et al., 2010), BD-IRT1vr/AD-empty and BD-
empty/AD-TTN5 respective negative controls. (B), Schematic representation of the switching
mechanism of small GTPases based on TTN5. TTN5 can switch between an inactive GDP-
loaded form to an active GTP-loaded one. Intrinsic GDP to GTP nucleotide exchange is
unusually high for TTN5 and help by a guanidine exchange factor (GEF) may not be needed,
but is not excluded, while GTP hydrolysis has to be catalyzed by a GTPase-activating protein
(GAP) (Mohr et al., 2023). GTP-loaded TTN5 can interact with effector proteins for signal
transition. (C-G), Validation of TTN5-IRT1vr interaction by bifunctional complementation (BiFC)
of split YFP. Genes of interest were fused either to the C- or N-terminal half of split-YFP.
Potential interaction is indicated by a complemented YFP signal. mRFP serves as a
transformation control. (C-E), nYFP-IRT1vr showed YFP complementation in combination with
(C) cYFP-TTN5, (D) dominant-negative (cYFP-TTN5^{T30N}) and dominant-active (cYFP-
TTN5^{Q70L}) variants. The complementation takes place in nucleus and cytoplasm predominantly.
(F-G), The combinations (F) cYFP-TTN5/nYFP-EHB1Δsig and (G) cYFP-EHB1Δsig/nYFP-
IRT1vr served as negative controls. Each combination was tested a minimum of three times

with comparable results. Scale bar 50 µm. (H-J), Confirmed interaction via FRET-APB.
Significant increase of FRET-efficiency between IRT1vr-mCherry and (H) GFP-TTN5, (I)
dominant-negative TTN5^{T30N} or (J) dominant-active TTN5^{Q70L} indicate short protein distances.
GFP-fusions are donor-only samples and serve as a negative control. GFP-mCherry-tagged
constructs show intra-molecular FRET as a respective positive control. Analysis was done in
the nucleus. Each combination was tested a minimum of three times (n = 3) with comparable
results. One-way ANOVA with Fisher-LSD post-hoc test was performed. Different letters
indicate statistical significance (p < 0.05, n ≥ 10).
711

Figure 2. Heterozygous *ttn5-1*$^{+/-}$ seedlings can have an Fe reductase phenotype.

Heterozygous *ttn5-1*$^{+/-}$ and their wild type sibling seedlings were grown on +/- Fe Hoagland
plates for ten days (A-B) or in the two-week system (C-G). All plants were individually
genotyped as wild type or *ttn5-1*$^{+/-}$. (A), *ttn5-1*$^{+/-}$ seedlings appeared phenotypically as wild type
(WT) siblings. Scale bar 1 cm. (B), Quantified root length of (A). Root length increased under
Fe deficient conditions (-Fe, light grey boxes) compared to sufficient conditions (+Fe, dark grey
boxes), similar in *ttn5-1*$^{+/-}$ and wild type siblings (+Fe: WT n = 19; *ttn5-1* n = 28; -Fe: WT n =
16; *ttn5-1* n = 32). (C), Chlorophyll contents, measured by SPAD values, were similar under
both Fe conditions (-Fe, light grey boxes; +Fe, dark grey boxes) and comparable between the
plants (+Fe: n = 10; -Fe: n = 9). (D-F), Gene expression data by RT-qPCR of *TTN5, FRO2* and
IRT1 in *ttn5-1*$^{+/-}$ roots compared to wild type siblings in Fe-sufficient (+Fe, black bars) or -
deficient (-Fe, grey bars) conditions. (D), *TTN5* expression was reduced under Fe deficient
conditions in wild type seedlings. Same took place in *ttn5-1*$^{+/-}$ seedlings under Fe deficiency.
(E-F) *FRO2* and *IRT1* expression was induced upon Fe deficiency in wild type and *ttn5-1*$^{+/-}$
seedlings in comparable amounts. Data was obtained from three biological replicates (n = 3).
(G), Ferric reductase activity was measured as color change due to formation of a Fe^{2+}-
ferrozine complex, detected at an OD of 562 nm. The increase in iron reductase activity in
response to Fe deficiency was higher in wild type than in *ttn5-1*$^{+/-}$ sibling seedlings. The assay
was performed with three (wild type) and four (*ttn5-1*$^{+/-}$) biological replicates (WT n = 3; *ttn5-1*$^{+/-}$
n = 4). The experiment was repeated in total three times. In two cases, Fe reductase activity
was reduced in *ttn5-1*$^{+/-}$ compared with wild type siblings, and once it was not different. One-
way ANOVA with Fisher-LSD as post-hoc test was performed. Different letters indicate
statistical significance (p < 0.05).
735

Figure 3. TTN5 co-localizes with IRT1 at the plasma membrane and in multi-vesicular
bodies (MVBs).

(A, D, H) mCherry-TTN5 and its (B, E, I) dominant-negative (mCherry-TTN5^{T30N}) and (C, F, J)
dominant-active (mCherry-TTN5^{Q70L}) variants were co-localized with IRT1-GFP in
N. benthamiana leaf epidermal cells. (A-C), mCherry-TTN5 and mCherry-TTN5 variants were
present at the plasma membrane and in distinct spots together with IRT1. (D-F), Internalized
IRT1-GFP was present in mCherry-TTN5 and mCherry-TTN5 variant-containing vesicle-like
structures. (G), Schematic illustration of the inhibiting function of the fungal metabolite
wortmannin on cellular trafficking routes between the vacuole and the *trans*-Golgi network via
MVBs, leading to swelling of MVBs. (H-J), Cells were treated with wortmannin. Swollen MVBs

are visible with the GFP- and mCherry-channel, respectively. Colocalization is indicated with
filled arrowheads. Scale bar 10 µm.
748

Figure 4. TTN5 interacts with IRT1 regulators EHB1, SNX1 and PATL2.

(A-E, G-P) Bimolecular fluorescence complementation (BiFC) of split YFP experiments with
TTN5 and known IRT1 regulators EHB1 (Khan et al., 2019), SNX1 (Ivanov et al., 2014) and
PATL2 (Hornbergs et al., 2022). mRFP expression serves as a transformation control. (A-E),
TTN5 interacts with EHB1 via the sig domain. (B, D, E), TTN5 and EHB1 showed
complementation whereas inactive and active variants were not able to complement split-YFP.
(A, C), TTN5 was tested together with EHB1sig (only the sig, signature domain) and with EHB1
harboring a deletion of the CAR sig domain (sig domain defined in (Rodriguez et al., 2014;
Khan et al. 2019), EHB1Δsig). Deletion of the sig domain resulted in loss of complementation.
(F), Interaction between TTN5 and TTN5 variants (GFP-tagged, donor) with EHB1 (mCherry-
tagged, acceptor) was confirmed by FRET-APB. EHB1-GFP as donor-only sample (negative
control), EHB1-GFP-mCherry for intra-molecular FRET (positive control). Each combination
was tested a minimum of three times (n = 3) with comparable results. In this assay, TTN5
variants interacted with EHB1. (G-K), BiFC showing that TTN5 interacts with SNX1. (G, I),
TTN5 and TTN5^{Q70L} showed complemented YFP signal together with SNX1. (H), No YFP
fluorescence was detectable for TTN5^{T30N} with SNX1. (J, K), An interaction took place between
TTN5 and the (J) PX and (K) the BAR domain of SNX1. (L-P), BiFC showing that TTN5
interacted with PATL2. (L), PATL2 showed complemented YFP signal with N-terminally-tagged
TTN5. (M-N), TTN5^{T30N} and TTN5^{Q70L} did not complement YFP with PATL2. (O-P), Deletion
mutants of PATL2 showed dependency of the N-terminus for complementation with TTN5. (O)
YFP signal was visible for PATL2 N-terminus (PATL2ΔC). (P) On the other hand, the C-
terminus alone (PATL2ΔN) had no detectable YFP. Every construct was tested a minimum of
three times (n ≥ 3) with similar results. Scale bar 50 µm. One-way ANOVA with Fisher-LSD
post-hoc test was performed. Different letters indicate statistical significance ($p < 0.05$).
773

Figure 5. Summary model: Action of TTN5 within the IRT1-regulatory network.

A schematic overview of the identified TTN5 interactors within the IRT1-regulatory network (A)
at the plasma membrane (PM) or (B) in the endomembrane compartments of the *trans*-Golgi
network (TGN) or multivesicular body (MVB). IRT1 undergoes a constant cycling due to
endocytosis from the plasma membrane and either degradation via the lytic pathway in the
vacuole or recycling back to the plasma membrane. (A), We identified TTN5 interaction with
the large cytosolic, variable region of IRT1. TTN5 and IRT1 co-localize at the plasma
membrane (circled 1). PATL2 interacts with the large cytosolic, variable region of IRT1 (IRT1vr)
via its N-terminus and is involved in α-tocopherol recruitment to the membrane, to prevent lipid
peroxidation induced by IRT1-related iron import (Hornbergs et al., 2022). TTN5 also interacts
with the N-terminus of PATL2. A potential TTN5-PATL-IRT1 complex formation is suggested
(circled 2). EHB1 interacts with the large cytosolic, variable region of IRT1 via its signature (sig)
domain and inhibits IRT1 function at the plasma membrane in a Ca^{2+}-dependent manner (Khan
et al. 2019). TTN5 also interacts with EHB1 via the sig domain. A potential TTN5-EHB1-IRT1
complex formation is suggested (circled 3). A possible four complex of IRT1, TTN5, PATL2 and
EHB1 (circled 4). (B), We found that IRT1 co-localized with SNX1 (Ivanov et al., 2014) and

TTN5 in the TGN. IRT1 and TTN5 also co-localized in MVBs. The protein symbols are indicated
in the figure legend. The question mark indicates that a potential protein complex may form.
792

Supplemental Figure S1: Localization of fluorescently labeled IRT1 and TTN5 in
Arabidopsis roots.
IRT1-mCitrine expression (A, C, E) of proIRT1::IRT1-mCit/*irt1-1* seedlings previously described
in Dubeaux et al. (2018) in comparison with YFP-TTN5 expression of pro35S::YFP-TTN5 (B,
D, F) in Arabidopsis roots. (A-B), IRT1-mCitrine was expressed at the plasma membrane of
epidermal cells (A) (empty white arrowhead). YFP-TTN5 expression was localized at the
plasma membrane of epidermal cells and other cell types (B). Dotted white lines indicate the
position of the longitudinal sections shown in C and D. (C-D), Longitudinal sections highlighted
IRT1-mCitrine (C) expression only in epidermal cells compared to nearly ubiquitously
expression of YFP-TTN5 (D). (E-F), Next to plasma membrane localization IRT1-mCitrine (E)
and YFP-TTN5 (F) were present in intracellular spots (filled white arrowheads). Scale bar
10 µm.
805

Supplemental Figure S2. IRT1 colocalization analysis at the plasma membrane and in
intracellular spots.
(A), Plasma membrane dye FM4-64 confirmed IRT1-GFP expression at the plasma membrane.
IRT1 is expressed in a spotted structure, indicating regions of higher transporter accumulation.
(B-C), IRT1-GFP colocalized with published IRT1-mCherry (Ivanov et al., 2014) at the plasma
membrane (PM) (B) and in intracellular spots (C), confirming correct localization of the IRT1-
GFP construct. Scale bar 50 µm. (D), Complementation of rosette growth defect of IRT1-GFP.
Rosette sizes of *irt1-1* and pro35S::IRT1-GFP/*irt1-1* were measured 14 days after sowing.
IRT1-GFP complemented the growth defect of *irt1-1*. (E-G), JACoP-based colocalization
analysis (Bolte and Cordelières, 2006) performed with ImageJ (Schneider et al., 2012). (E-F),
Comparison of Pearson's and Overlap coefficients for IRT1-GFP with the three mCherry-TTN5
variants at the plasma membrane and in vesicle-like structures was done. (G), Object-based
analysis was performed for spotted-structures based on distance between geometrical centers
of signals. Analyses were conducted in three replicates each (n = 3).
One-way ANOVA with Fisher-LSD post-hoc test was performed. Different letters indicate
statistical significance (p < 0.05).
822

Supplemental Figure S3. TTN5 effect on its interactors on transcriptional level.
(A-C), Gene expression data by RT-qPCR of TTN5 interactors EHB1, SNX1 and PATL2 in *ttn5-*
1$^{+/-}$ roots compared to wild type. Plants were grown in the two-week system in Fe-sufficient
(+Fe, black bars) or -deficient (-Fe, grey bars) conditions. (A), *EHB1* expression was repressed
under Fe-deficient conditions in wild type seedlings. Equal expression levels were present in
ttn5-1 seedlings under both Fe supply conditions. (B), *SNX1* expression was not altered
between Fe-sufficient or -deficient conditions in wild type and *ttn5-1* seedlings. (C), Next to
SNX1, *PATL2* expression did not change upon Fe supply in both wild type and *ttn5-1* seedlings.
Data was obtained from three biological replicates (n = 3). One-way ANOVA with Fisher-LSD
post-hoc test was performed. Different letters indicate statistical significance (p < 0.05).

References

Adarska P, Wong-Dilworth L, Bottanelli F (2021) ARF GTPases and Their Ubiquitous Role in Intracellular Trafficking Beyond the Golgi. Frontiers in cell and developmental biology **9:** 679046-679046

Anders N, Nielsen M, Keicher J, Stierhof YD, Furutani M, Tasaka M, Skriver K, Jürgens G (2008) Membrane association of the Arabidopsis ARF exchange factor GNOM involves interaction of conserved domains. Plant Cell **20:** 142-151

Antoshechkin I, Han M (2002) The *C. elegans* evl-20 Gene Is a Homolog of the Small GTPase ARL2 and Regulates Cytoskeleton Dynamics during Cytokinesis and Morphogenesis. Developmental Cell **2:** 579-591

Arora D, Van Damme D (2021) Motif-based endomembrane trafficking. Plant Physiology **186:** 221-238

Barberon M, Dubeaux G, Kolb C, Isono E, Zelazny E, Vert G (2014) Polarization of IRON-REGULATED TRANSPORTER 1 (IRT1) to the plant-soil interface plays crucial role in metal homeostasis. Proceedings of the National Academy of Sciences **111:** 8293-8298

Barberon M, Zelazny E, Robert S, Conéjéro G, Curie C, Friml J, Vert G (2011) Monoubiquitin-dependent endocytosis of the IRON-REGULATED TRANSPORTER 1 (IRT1) transporter controls iron uptake in plants. Proceedings of the National Academy of Sciences **108:** E450-E458

Benková E, Michniewicz M, Sauer M, Teichmann T, Seifertová D, Jürgens G, Friml J (2003) Local, Efflux-Dependent Auxin Gradients as a Common Module for Plant Organ Formation. Cell **115:** 591-602

Bhamidipati A, Lewis SA, Cowan NJ (2000) ADP ribosylation factor-like protein 2 (Arl2) regulates the interaction of tubulin-folding cofactor D with native tubulin. Journal of Cell Biology **149:** 1087-1096

Bleckmann A, Weidtkamp-Peters S, Seidel CAM, Simon R (2010) Stem cell signaling in Arabidopsis requires CRN to localize CLV2 to the plasma membrane. Plant physiology **152:** 166-176

Bolte S, Cordelières FP (2006) A guided tour into subcellular colocalization analysis in light microscopy. J Microsc **224:** 213-232

Brumbarova T, Ivanov R (2019) The Nutrient Response Transcriptional Regulome of Arabidopsis. iScience **19:** 358-368

Chen W, Zhou H, Xu F, Yu M, Coego A, Rodriguez L, Lu Y, Xie Q, Fu Q, Chen J, Xu G, Wu D, Li X, Li X, Jaillais Y, Rodriguez PL, Zhu S, Yu F (2023) CAR modulates plasma membrane nano-organization and immune signaling downstream of RALF1-FERONIA signaling pathway. New Phytol **237:** 2148-2162

Cointry V, Vert G (2019) The bifunctional transporter-receptor IRT1 at the heart of metal sensing and signalling. New Phytologist **223:** 1173-1178

Cui M, Gupta SK, Bauer P (2023) Review: Role of the plant-specific calcium-binding C2-DOMAIN ABSCISIC ACID-RELATED (CAR) protein family in environmental signaling. . OSF Preprints

Cui Y, Li X, Chen Q, He X, Yang Q, Zhang A, Yu X, Chen H, Liu N, Xie Q, Yang W, Zuo J, Palme K, Li W (2010) BLOS1, a putative BLOC-1 subunit, interacts with SNX1 and modulates root growth in Arabidopsis. Journal of Cell Science **123:** 3727-3733

Cui Y, Shen J, Gao C, Zhuang X, Wang J, Jiang L (2016) Biogenesis of Plant Prevacuolar Multivesicular Bodies. Molecular Plant **9:** 774-786

Curtis MD, Grossniklaus U (2003) A gateway cloning vector set for high-throughput functional analysis of genes in planta. Plant physiology **133:** 462-469

Dubeaux G, Neveu J, Zelazny E, Vert G (2018) Metal Sensing by the IRT1 Transporter-Receptor Orchestrates Its Own Degradation and Plant Metal Nutrition. Molecular Cell **69:** 953-964.e955

Dümmer M, Michalski C, Essen LO, Rath M, Galland P, Forreiter C (2016) EHB1 and
AGD12, two calcium-dependent proteins affect gravitropism antagonistically in
Arabidopsis thaliana. J Plant Physiol **206:** 114-124
Eide D, Broderius M, Fett J, Guerinot ML (1996) A novel iron-regulated metal transporter
from plants identified by functional expression in yeast. Proceedings of the National
Academy of Sciences **93:** 5624-5628
Eide D, Broderius M, Fett J, Guerinot ML (1996) A novel iron-regulated metal transporter
from plants identified by functional expression in yeast. Proceedings of the National
Academy of Sciences of the United States of America **93:** 5624-5628
Fansa EK, Wittinghofer A (2016) Sorting of lipidated cargo by the Arl2/Arl3 system. Small
GTPases **7:** 222-230
Fleming JA, Vega LR, Solomon F (2000) Function of tubulin binding proteins in vivo. Genetics
156: 69-80
Foot N, Henshall T, Kumar S (2017) Ubiquitination and the Regulation of Membrane Proteins.
Physiological Reviews **97:** 253-281
Francis JW, Turn RE, Newman LE, Schiavon C, Kahn RA (2016) Higher order signaling:
ARL2 as regulator of both mitochondrial fusion and microtubule dynamics allows
integration of 2 essential cell functions. Small GTPases **7:** 188-196
Frost A, Unger VM, De Camilli P (2009) The BAR Domain Superfamily: Membrane-Molding
Macromolecules. Cell **137:** 191-196
Fukaya M, Fukushima D, Hara Y, Sakagami H (2014) EFA6A, a guanine nucleotide exchange
factor for Arf6, interacts with sorting nexin-1 and regulates neurite outgrowth. Journal of
Neurochemistry **129:** 21-36
Gaither LA, Eide DJ (2001) Eukaryotic zinc transporters and their regulation. Biometals **14:**
251-270
Geldner N, Anders N, Wolters H, Keicher J, Kornberger W, Muller P, Delbarre A, Ueda T,
Nakano A, Jürgens G (2003) The Arabidopsis GNOM ARF-GEF mediates endosomal
recycling, auxin transport, and auxin-dependent plant growth. Cell **112:** 219-230
Gratz R, Manishankar P, Ivanov R, Köster P, Mohr I, Trofimov K, Steinhorst L, Meiser J,
Mai HJ, Drerup M, Arendt S, Holtkamp M, Karst U, Kudla J, Bauer P, Brumbarova
T (2019) CIPK11-Dependent Phosphorylation Modulates FIT Activity to Promote
Arabidopsis Iron Acquisition in Response to Calcium Signaling. Dev Cell **48:** 726-
740.e710
Gratz R, von der Mark C, Ivanov R, Brumbarova T (2021) Fe acquisition at the crossroad of
calcium and reactive oxygen species signaling. Curr Opin Plant Biol **63:** 102048
Grefen C, Blatt MR (2012) A 2in1 cloning system enables ratiometric bimolecular fluorescence
complementation (rBiFC). Biotechniques **53:** 311-314
Grossoehme NE, Akilesh S, Guerinot ML, Wilcox DE (2006) Metal-Binding
Thermodynamics of the Histidine-Rich Sequence from the Metal-Transport Protein
IRT1 of *Arabidopsis thaliana*. Inorganic Chemistry **45:** 8500-8508
Guerinot ML (2000) The ZIP family of metal transporters. Biochimica et Biophysica Acta (BBA)
- Biomembranes **1465:** 190-198
Heucken N, Ivanov R (2018) The retromer, sorting nexins and the plant endomembrane
protein trafficking. J Cell Sci **131**
Hornbergs J, Montag K, Loschwitz J, Mohr I, Poschmann G, Schnake A, Gratz R,
Brumbarova T, Eutebach M, Angrand K, Fink-Straube C, Stühler K, Zeier J,
Hartmann L, Strodel B, Ivanov R, Bauer P (2022) SEC14-GOLD protein PATELLIN2
binds IRON-REGULATED TRANSPORTER1 linking root iron uptake to vitamin E. Plant
Physiol
Hu J (2021) Toward unzipping the ZIP metal transporters: structure, evolution, and implications
on drug discovery against cancer. The FEBS Journal **288:** 5805-5825

Ismail SA, Chen YX, Rusinova A, Chandra A, Bierbaum M, Gremer L, Triola G, Waldmann H, Bastiaens PI, Wittinghofer A (2011) Arl2-GTP and Arl3-GTP regulate a GDI-like transport system for farnesylated cargo. Nat Chem Biol **7:** 942-949

Ismail SA, Vetter IR, Sot B, Wittinghofer A (2010) The Structure of an Arf-ArfGAP Complex Reveals a Ca^{2+} Regulatory Mechanism. Cell **141:** 812-821

Ivanov R, Brumbarova T, Blum A, Jantke A-M, Fink-Straube C, Bauer P (2014) SORTING NEXIN1 is required for modulating the trafficking and stability of the Arabidopsis IRON-REGULATED TRANSPORTER1. The Plant cell **26:** 1294-1307

Ivanov R, Vert G (2021) Endocytosis in plants: Peculiarities and roles in the regulated trafficking of plant metal transporters. Biology of the Cell **113:** 1-13

Jaillais Y, Fobis-Loisy I, Miège C, Rollin C, Gaude T (2006) AtSNX1 defines an endosome for auxin-carrier trafficking in Arabidopsis. Nature **443:** 106-109

Kerkeb L, Mukherjee I, Chatterjee I, Lahner B, Salt DE, Connolly EL (2008) Iron-Induced Turnover of the Arabidopsis IRON-REGULATED TRANSPORTER1 Metal Transporter Requires Lysine Residues   Plant Physiology **146:** 1964-1973

Khan I, Gratz R, Denezhkin P, Schott-Verdugo SN, Angrand K, Genders L, Basgaran RM, Fink-Straube C, Brumbarova T, Gohlke H, Bauer P, Ivanov R (2019) Calcium-Promoted Interaction between the C2-Domain Protein EHB1 and Metal Transporter IRT1 Inhibits Arabidopsis Iron Acquisition. Plant Physiology **180:** 1564-1581

Kleine-Vehn J, Dhonukshe P, Sauer M, Brewer PB, Wiśniewska J, Paciorek T, Benková E, Friml J (2008) ARF GEF-dependent transcytosis and polar delivery of PIN auxin carriers in Arabidopsis. Curr Biol **18:** 526-531

Knauer T, Dümmer M, Landgraf F, Forreiter C (2011) A negative effector of blue light-induced and gravitropic bending in Arabidopsis. Plant physiology **156:** 439-447

Kolb C, Nagel M-K, Kalinowska K, Hagmann J, Ichikawa M, Anzenberger F, Alkofer A, Sato MH, Braun P, Isono E (2015) FYVE1 is essential for vacuole biogenesis and intracellular trafficking in Arabidopsis. Plant physiology **167:** 1361-1373

Le CTT, Brumbarova T, Ivanov R, Stoof C, Weber E, Mohrbacher J, Fink-Straube C, Bauer P (2015) ZINC FINGER OF ARABIDOPSIS THALIANA12 (ZAT12) Interacts with FER-LIKE IRON DEFICIENCY-INDUCED TRANSCRIPTION FACTOR (FIT) Linking Iron Deficiency and Oxidative Stress Responses Plant Physiology **170:** 540-557

Li W, Feng Y, Hao C, Guo X, Cui Y, He M, He X (2007) The BLOC Interactomes Form a Network in Endosomal Transport. Journal of Genetics and Genomics **34:** 669-682

Martín-Barranco A, Spielmann J, Dubeaux G, Vert G, Zelazny E (2020) Dynamic Control of the High-Affinity Iron Uptake Complex in Root Epidermal Cells1. Plant Physiology **184:** 1236-1250

Matthus E, Wilkins KA, Davies JM (2019) Iron availability modulates the *Arabidopsis thaliana* root calcium signature evoked by exogenous ATP. Plant Signaling & Behavior **14:** 1640563

Mayer U, Herzog U, Berger F, Inzé D, Jürgens G (1999) Mutations in the PILZ group genes disrupt the microtubule cytoskeleton and uncouple cell cycle progression from cell division in Arabidopsis embryo and endosperm. European Journal of Cell Biology **78:** 100-108

McElver J, Patton D, Rumbaugh M, Liu C-m, Yang LJ, Meinke D (2000) The TITAN5 Gene of Arabidopsis Encodes a Protein Related to the ADP Ribosylation Factor Family of GTP Binding Proteins. The Plant Cell **12:** 1379-1392

Memon AR (2004) The role of ADP-ribosylation factor and SAR1 in vesicular trafficking in plants. Biochim Biophys Acta **1664:** 9-30

Mohr I, Mirzaiebadizi A, Sanyal SK, Chuenban P, Ahmadian MR, Ivanov R, Bauer P (2023) Characterization of the small Arabidopsis thaliana GTPase and ADP-ribosylation factor-like 2 protein TITAN 5. BIORXIV/2023/538563

Montag K, Hornbergs J, Ivanov R, Bauer P (2020) Phylogenetic analysis of plant multi-
domain SEC14-like phosphatidylinositol transfer proteins and structure–function
properties of PATELLIN2. Plant Molecular Biology **104:** 665-678
Montag K, Ivanov R, Bauer P (2023) Role of SEC14-like phosphatidylinositol transfer proteins
in membrane identity and dynamics. OSF Preprints
Mori R, Toda T (2013) The dual role of fission yeast Tbc1/cofactor C orchestrates microtubule
homeostasis in tubulin folding and acts as a GAP for GTPase Alp41/Arl2. Mol Biol Cell
24: 1713-1724, s1711-1718
Nielsen E (2020) The Small GTPase Superfamily in Plants: A Conserved Regulatory Module
with Novel Functions. Annual Review of Plant Biology **71:** 247-272
Peter BJ, Kent HM, Mills IG, Vallis Y, Butler PJ, Evans PR, McMahon HT (2004) BAR
domains as sensors of membrane curvature: the amphiphysin BAR structure. Science
303: 495-499
Pourcher M, Santambrogio M, Thazar N, Thierry A-M, Fobis-Loisy I, Miège C, Jaillais Y,
Gaude T (2010) Analyses of sorting nexins reveal distinct retromer-subcomplex
functions in development and protein sorting in *Arabidopsis thaliana*. The Plant cell **22:**
3980-3991
Radcliffe PA, Vardy L, Toda T (2000) A conserved small GTP-binding protein Alp41 is
essential for the cofactor-dependent biogenesis of microtubules in fission yeast. FEBS
Letters **468:** 84-88
Raposo G, Marks MS, Cutler DF (2007) Lysosome-related organelles: driving post-Golgi
compartments into specialisation. Current Opinion in Cell Biology **19:** 394-401
Rath M, Dümmer M, Galland P, Forreiter C (2020) A gravitropic stimulus alters the distribution
of EHB1, a negative effector of root gravitropism in Arabidopsis. Plant Direct **4:** e00215
Robinson NJ, Procter CM, Connolly EL, Guerinot ML (1999) A ferric-chelate reductase for
iron uptake from soils. Nature **397:** 694-697
Rodriguez L, Gonzalez-Guzman M, Diaz M, Rodrigues A, Izquierdo-Garcia AC, Peirats-
Llobet M, Fernandez MA, Antoni R, Fernandez D, Marquez JA, Mulet JM, Albert A,
Rodriguez PL (2014) C2-domain abscisic acid-related proteins mediate the interaction
of PYR/PYL/RCAR abscisic acid receptors with the plasma membrane and regulate
abscisic acid sensitivity in Arabidopsis. Plant Cell **26:** 4802-4820
Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image
analysis. Nature Methods **9:** 671-675
Sharer JD, Shern JF, Van Valkenburgh H, Wallace DC, Kahn RA (2002) ARL2 and BART
Enter Mitochondria and Bind the Adenine Nucleotide Transporter. Molecular Biology of
the Cell **13:** 71-83
Shin L-J, Lo J-C, Chen G-H, Callis J, Fu H, Yeh K-C (2013) IRT1 degradation factor1, a ring
E3 ubiquitin ligase, regulates the degradation of iron-regulated transporter1 in
Arabidopsis. The Plant cell **25:** 3039-3051
Steinmann T, Geldner N, Grebe M, Mangold S, Jackson CL, Paris S, Gälweiler L, Palme
K, Jürgens G (1999) Coordinated polar localization of auxin efflux carrier PIN1 by
GNOM ARF GEF. Science **286:** 316-318
Sztul E, Chen P-W, Casanova JE, Cherfils J, Dacks JB, Lambright DG, Lee F-JS,
Randazzo PA, Santy LC, Schürmann A, Wilhelmi I, Yohe ME, Kahn RA (2019) ARF
GTPases and their GEFs and GAPs: concepts and challenges. Molecular biology of the
cell **30:** 1249-1271
Takano J, Tanaka M, Toyoda A, Miwa K, Kasai K, Fuji K, Onouchi H, Naito S, Fujiwara T
(2010) Polar localization and degradation of Arabidopsis boron transporters through
distinct trafficking pathways. Proc Natl Acad Sci U S A **107:** 5220-5225
Tanaka H, Kitakura S, De Rycke R, De Groodt R, Friml J (2009) Fluorescence imaging-
based screen identifies ARF GEF component of early endosomal trafficking. Curr Biol
19: 391-397

Tanaka H, Nodzyński T, Kitakura S, Feraru MI, Sasabe M, Ishikawa T, Kleine-Vehn J,
Kakimoto T, Friml J (2014) BEX1/ARF1A1C is Required for BFA-Sensitive Recycling
of PIN Auxin Transporters and Auxin-Mediated Development in Arabidopsis. Plant and
Cell Physiology **55:** 737-749

Tian Q, Zhang X, Yang A, Wang T, Zhang W-H (2016) CIPK23 is involved in iron acquisition
of Arabidopsis by affecting ferric chelate reductase activity. Plant Science **246:** 70-79

Tzafrir I, McElver JA, Liu Cm C-m, Yang LJ, Wu JQ, Martinez A, Patton DA, Meinke DW
(2002) Diversity of TITAN functions in Arabidopsis seed development. Plant physiology
128: 38-51

Valencia JP, Goodman K, Otegui MS (2016) Endocytosis and Endosomal Trafficking in
Plants. Annual Review of Plant Biology **67:** 309-335

Vernoud V, Horton AC, Yang Z, Nielsen E (2003) Analysis of the small GTPase gene
superfamily of Arabidopsis. Plant Physiol **131:** 1191-1208

Vert G, Grotz N, Dédaldéchamp F, Gaymard F, Guerinot ML, Briat J-F, Curie C (2002)
IRT1, an Arabidopsis Transporter Essential for Iron Uptake from the Soil and for Plant
Growth. The Plant Cell **14:** 1223-1233

von der Mark C, Ivanov R, Eutebach M, Maurino VG, Bauer P, Brumbarova T (2020)
Reactive oxygen species coordinate the transcriptional responses to iron availability in
Arabidopsis. Journal of Experimental Botany **72:** 2181-2195

Yoshinari A, Toda Y, Takano J (2021) GNOM-dependent endocytosis maintains polar
localisation of the borate exporter BOR1 in Arabidopsis. Biology of the Cell **113:** 264-
269

Zhang C, Tong C, Cao L, Zheng P, Tang X, Wang L, Miao M, Liu Y, Cao S (2023) Regulatory
module WRKY33-ATL31-IRT1 mediates cadmium tolerance in Arabidopsis. Plant, Cell
& Environment **46:** 1653-1670

1065

Figure 1

Figure 1. TTN5 interacts with IRT1vr in yeast and in plant cells.

(A), Targeted Y2H assay between BD-IRT1vr (IRT1-variable region) and AD-TTN5. Yeast cotransformations were spotted in 10-fold dilution series ($A_{600} = 10^1$-10^4) on double selective (LW; control) and triple selective +0.5 mM 3-AT (LWH, selection for protein interactions) SD-plates. SNX1/SNX1, positive control (Pourcher et al., 2010), BD-IRT1vr/AD-empty and BD-empty/AD-TTN5 respective negative controls. (B), Schematic representation of the switching mechanism of small GTPases based on TTN5. TTN5 can switch between an inactive GDP-loaded form to an active GTP-loaded one. Intrinsic GDP to GTP nucleotide exchange is unusually high for TTN5 and help by a guanidine exchange factor (GEF) may not be needed, but is not excluded, while GTP hydrolysis has to be catalyzed by a GTPase-activating protein (GAP) (Mohr et al., 2023). GTP-loaded TTN5 can interact with effector proteins for signal transition. (C-G), Validation of TTN5-IRT1vr interaction by bifunctional complementation (BiFC) of split YFP. Genes of interest were fused either to the C- or N-terminal half of split-YFP. Potential interaction is indicated by a complemented YFP signal. mRFP serves as a transformation control. (C-E), nYFP-IRT1vr showed YFP complementation in combination with (C) cYFP-TTN5, (D) dominant-negative (cYFP-TTN5^{T30N}) and dominant-active (cYFP-TTN5^{Q70L}) variants. The complementation takes place in nucleus and cytoplasm predominantly. (F-G), The combinations (F) cYFP-TTN5/nYFP-EHB1Δsig and (G) cYFP-EHB1Δsig/nYFP-IRT1vr served as negative controls. Each combination was tested a minimum of three times with comparable results. Scale bar 50 µm. (H-J), Confirmed interaction via FRET-APB. Significant increase of FRET-efficiency between IRT1vr-mCherry and (H) GFP-TTN5, (I) dominant-negative TTN5^{T30N} or (J) dominant-active TTN5^{Q70L} indicate short protein distances. GFP-fusions are donor-only samples and serve as a negative control. GFP-mCherry-tagged constructs show intra-molecular FRET as a respective positive control. Analysis was done in the nucleus. Each combination was tested a minimum of three times (n = 3) with comparable results. One-way ANOVA with Fisher-LSD post-hoc test was performed. Different letters indicate statistical significance ($p < 0.05$, $n \geq 10$).

Figure 2

Figure 2. Heterozygous *ttn5-1*[+/-] seedlings have an Fe reductase phenotype.

Heterozygous *ttn5-1*[+/-] and their wild type sibling seedlings were grown on +/- Fe Hoagland plates for ten days (A-B) or in the two-week system (C-G). All plants were individually genotyped as wild type or *ttn5-1*[+/-].

(A), *ttn5-1*[+/-] seedlings appeared phenotypically as wild type (WT) siblings. Scale bar 1 cm. (B), Quantified root length of (A). Root length increased under Fe deficient conditions (-Fe, light grey boxes) compared to sufficient conditions (+Fe, dark grey boxes), similar in *ttn5-1*[+/-] and wild type siblings (+Fe: WT n = 19; *ttn5-1* n = 28; -Fe: WT n = 16; *ttn5-1* n = 32). (C), Chlorophyll contents, measured by SPAD values, were similar under both Fe conditions (-Fe, light grey boxes; +Fe, dark grey boxes) and comparable between the plants (+Fe: n = 10; -Fe: n = 9). (D-F), Gene expression data by RT-qPCR of *TTN5, FRO2* and *IRT1* in *ttn5-1*[+/-] roots compared to wild type siblings in Fe-sufficient (+Fe, black bars) or -deficient (-Fe, grey bars) conditions. (D), *TTN5* expression was reduced under Fe deficient conditions in wild type seedlings. Same took place in *ttn5-1*[+/-] seedlings under Fe deficiency. (E-F) *FRO2* and *IRT1* expression was induced upon Fe deficiency in wild type and *ttn5-1*[+/-] seedlings in comparable amounts. Data was obtained from three biological replicates (n = 3). (G), Ferric reductase activity was measured as color change due to formation of a Fe^{2+}-ferrozine complex, detected at an OD of 562 nm. The increase in iron reductase activity in response to Fe deficiency was higher in wild type than in *ttn5-1*[+/-]. The assay was performed with three (wild type) and four (*ttn5-1*[+/-]) biological replicates (WT n = 3; *ttn5-1*[+/-] n = 4). The experiment was repeated in total three times. In two cases, Fe reductase activity was reduced in *ttn5-1*[+/-] compared with wild type siblings, and once it was not different.

One-way ANOVA with Fisher-LSD as post-hoc test was performed. Different letters indicate statistical significance ($p < 0.05$).

Figure 3

Figure 3. TTN5 co-localizes with IRT1 at the plasma membrane and in multi-vesicular bodies (MVBs).

(A, D, H) mCherry-TTN5 and its (B, E, I) dominant-negative (mCherry-TTN5^{T30N}) and (C, F, J) dominant-active (mCherry-TTN5^{Q70L}) variants were co-localized with IRT1-GFP in *N. benthamiana* leaf epidermal cells. (A-C), mCherry-TTN5 and mCherry-TTN5 variants were present at the plasma membrane and in distinct spots together with IRT1. (D-F), Internalized IRT1-GFP was present in mCherry-TTN5 and mCherry-TTN5 variant-containing vesicle-like structures. (G), Schematic illustration of the inhibiting function of the fungal metabolite wortmannin on cellular trafficking routes between the vacuole and the *trans*-Golgi network via MVBs, leading to swelling of MVBs. (H-J), Cells were treated with wortmannin. Swollen MVBs are visible with the GFP- and mCherry-channel, respectively. Colocalization is indicated with filled arrowheads. Scale bar 10 µm.

Figure 4

Figure 4. TTN5 interacts with IRT1 regulators EHB1, SNX1 and PATL2.

(A-E, G-P) Bimolecular fluorescence complementation (BiFC) of split YFP experiments with TTN5 and known IRT1 regulators EHB1 (Khan et al., 2019), SNX1 (Ivanov et al., 2014) and PATL2 (Hornbergs et al., 2022). mRFP expression serves as a transformation control. (A-E), TTN5 interacts with EHB1 via the sig domain. (B, D, E), TTN5 and EHB1 showed complementation whereas inactive and active variants were not able to complement split-YFP. (A, C), TTN5 was tested together with EHB1sig (only the sig, signature domain) and with EHB1 harboring a deletion of the CAR sig domain (sig domain defined in (Rodriguez et al., 2014; Khan et al. 2019), EHB1Δsig). Deletion of the sig domain resulted in loss of complementation. (F), Interaction between TTN5 and TTN5 variants (GFP-tagged, donor) with EHB1 (mCherry-tagged, acceptor) was confirmed by FRET-APB. EHB1-GFP as donor-only sample (negative control), EHB1-GFP-mCherry for intra-molecular FRET (positive control). Each combination was tested a minimum of three times (n = 3) with comparable results. In this assay, TTN5 variants interacted with EHB1. (G-K), BiFC showing that TTN5 interacts with SNX1. (G, I), TTN5 and TTN5^{Q70L} showed complemented YFP signal together with SNX1. (H), No YFP fluorescence was detectable for TTN5^{T30N} with SNX1. (J, K), An interaction took place between TTN5 and the (J) PX and (K) the BAR domain of SNX1. (L-P), BiFC showing that TTN5 interacted with PATL2. (L), PATL2 showed complemented YFP signal with N-terminally-tagged TTN5. (M-N), TTN5^{T30N} and TTN5^{Q70L} did not complement YFP with PATL2. (O-P), Deletion mutants of PATL2 showed dependency of the N-terminus for complementation with TTN5. (O) YFP signal was visible for PATL2 N-terminus (PATL2ΔC). (P) On the other hand, the C-terminus alone (PATL2ΔN) had no detectable YFP. Every construct was tested a minimum of three times (n ≥ 3) with similar results. Scale bar 50 µm.

One-way ANOVA with Fisher-LSD post-hoc test was performed. Different letters indicate statistical significance ($p < 0.05$).

Figure 5

Figure 5. Summary model: Action of TTN5 within the IRT1-regulatory network.
A schematic overview of the identified TTN5 interactors within the IRT1-regulatory network (A) at the plasma membrane (PM) or (B) in the endomembrane compartments of the *trans*-Golgi network (TGN) or multivesicular body (MVB). IRT1 undergoes a constant cycling due to endocytosis from the plasma membrane and either degradation via the lytic pathway in the vacuole or recycling back to the plasma membrane. (A), We identified TTN5 interaction with the large cytosolic, variable region of IRT1. TTN5 and IRT1 co-localize at the plasma membrane (circled 1). PATL2 interacts with the large cytosolic, variable region of IRT1 (IRT1vr) via its N-terminus and is involved in α-tocopherol recruitment to the membrane, to prevent lipid peroxidation induced by IRT1-related iron import (Hornbergs et al. 2022). TTN5 also interacts with the N-terminus of PATL2. A potential TTN5-PATL-IRT1 complex formation is suggested (circled 2). EHB1 interacts with the large cytosolic, variable region of IRT1 via its signature (sig) domain and inhibits IRT1 function at the plasma membrane in a Ca^{2+}-dependent manner (Khan et al. 2019). TTN5 also interacts with EHB1 via the sig domain. A potential TTN5-EHB1-IRT1 complex formation is suggested (circled 3). A possible four complex of IRT1, TTN5, PATL2 and EHB1 (circled 4). (B) We found that IRT1 co-localized with SNX1 (Ivanov et al. 2014) and TTN5 in the TGN. IRT1 and TTN5 also co-localized in MVBs. The protein symbols are explained in the legend. The question mark indicates that the potential protein complex (pot. complex) may form.

Supplemental Figure 1

IRT1-mCitrine   YFP-TTN5

Supplemental Figure S1: Localization of fluorescently labeled IRT1 and TTN5 in Arabidopsis roots.

(A, C, E) IRT1-mCitrine localization in proIRT1::IRT1-mCit/*irt1-1* seedling roots (line previously described in Dubeaux et al. 2018) in comparison with (B, D, F) YFP-TTN5 localization in pro35S::YFP-TTN5/WT seedling roots (line previously described in Mohr et al. 2023). (A, B) Longitudinal sections through the stele. (A), IRT1-mCitrine was present at the plasma membrane of epidermal cells (empty white arrowhead). (B) YFP-TTN5 was localized at the plasma membrane of epidermal cells and other cell types, as well as in the stele (empty magenta arrowhead). Dotted white lines indicate the position of the cross sections shown in C and D. (C-D), Cross sections highlighting (C) IRT1-mCitrine in epidermal cells compared to (D) nearly ubiquitously localized YFP-TTN5. (E-F) Longitudinal section through the epidermis regions. Next to plasma membrane, (E) IRT1-mCitrine and (F) YFP-TTN5 were present in intracellular spots (filled white arrowheads). YFP-TTN5 was also detected in the nucleus (filled magenta arrowhead). Scale bar 10 μm.

Supplemental Figure 2

Channel		Colocalization spots	
GFP/YFP	mRFP/mCh	GFP/mRFP	mRFP/GFP
IRT1-GFP	mCh-TTN5	64.10±25.58	41.13±28.78
IRT1-GFP	mCh-TTN5T30N	54.44±30.81	36.65±8.25
IRT1-GFP	mCh-TTN5Q70L	82.59±12.71	36.34±9.66

Supplemental Figure S2. IRT1 co-localization analysis at the plasma membrane and in intracellular spots.

(A), Plasma membrane dye FM4-64 indicated IRT1-GFP localization at the plasma membrane. IRT1-GFP is present in a spotted structure, indicating regions of higher transporter accumulation. (B-C), IRT1-GFP co-localized with IRT1-mCherry (Ivanov et al., 2014) (B) at the plasma membrane (PM) and (C) in intracellular spots, confirming correct localization of IRT1-GFP. Scale bar 50 µm. (D), Complementation of rosette growth defect of *irt1-1* by IRT1-GFP. Rosette sizes of *irt1-1* and pro35S::IRT1-GFP/*irt1-1* were measured 14 days after sowing. IRT1-GFP complemented the growth defect of *irt1-1*. (E-G), JACoP-based colocalization analysis of mCherry-TTN5 and variants with IRT1-GFP (Bolte and Cordelières, 2006) performed with ImageJ (Schneider et al., 2012). (E-F), Comparison of (from left to right) Pearson's and Overlap coefficients for IRT1-GFP with mCherry-TTN5, mCherry-TTN5^{T30N} and mCherry-TTN5^{Q70L} (E) at the plasma membrane and (F) in vesicle-like structures. (G), Object-based analysis was performed for spotted-structures based on distance between geometrical centers of signals. Analyses were conducted in three replicates each (n=3). One-way ANOVA with Fisher-LSD post-hoc test was performed. Different letters indicate statistical significance (p < 0.05).

Supplemental Figure 3

Supplemental Figure S3. Gene expression of *EHB1*, *SNX1* and *PATL2* is not affected in *ttn5-1*[+/-] plants.

Gene expression data by RT-qPCR of TTN5 interactors *EHB1*, *SNX1* and *PATL2* in *ttn5-1*[+/-] roots compared to wild type sibling roots. Plants were grown in the two-week system in Fe- sufficient (+Fe, black bars) or -deficient (-Fe, grey bars) conditions. See also Figure 2D-F. (A), *EHB1* expression was down-regulated under Fe deficient conditions in wild type seedlings, while expression levels were not differing in *ttn5-1* seedlings under both Fe supply conditions. (B), *SNX1* expression was not altered between Fe-sufficient or -deficient conditions in wild type and *ttn5-1* seedlings. (C), *PATL2* expression did not change upon Fe supply in both wild-type and *ttn5-1* seedlings. Data were obtained from three biological replicates (n = 3). One-way ANOVA with Fisher-LSD post-hoc test was performed. Different letters indicate statistical significance ($p < 0.05$).