GluN2C/D-containing NMDA receptors enhance temporal summation and increase sound-

evoked and spontaneous firing in the inferior colliculus

Abbreviated title: GluN2C/D-containing NMDARs in the IC

Audrey C. Drotos[1], Rachel L. Zarb[1], Victoria Booth[3,4], Michael T. Roberts[1,2,]*

1. Kresge Hearing Research Institute, Department of Otolaryngology – Head and Neck Surgery,

University of Michigan, Ann Arbor, Michigan 48109

2. Department of Molecular and Integrative Physiology, University of Michigan, Ann Arbor,

Michigan 48109

3. Department of Mathematics, University of Michigan, Ann Arbor, Michigan 48109

4. Department of Anesthesiology, University of Michigan, Ann Arbor, Michigan 48109

*Corresponding author: microb@umich.edu

Number of pages: 51

Number of figures: 9

Number of tables: 2

Number of words in abstract: 234

Number of words in introduction: 599

Number of words in discussion: 1398

Conflict of Interest Statement: The authors declare no competing financial interests.

Acknowledgements: The authors would like to thank Yoani Herrera, Marina Silveira, Pierre

Apostolides, Marie Walicki, Will Birdsong, and Kelly Sovacool for helpful discussions and advice.

This work was supported by a Rackham Graduate Student Research Grant (ACD), the National

Science Foundation Graduate Research Fellowship Program under Grant No. DGE 1256260

(ACD), the National Institutes of Health Grant R01 DC018284 (MTR), and the National Institutes

of Health Grant F31 DC021344 (ACD).

Author contributions: ACD and MTR conceived of the study and designed experiments. ACD,

VB, and RLZ conducted experiments. ACD analyzed data, prepared figures, and prepared the

initial draft of the manuscript. ACD and MTR revised the manuscript, and all authors approved

the submitted version.

Significance statement

NMDA receptors are critical components of most glutamatergic circuits in the brain, and the

diversity of NMDA receptor subtypes yields receptors with a variety of functions. We found that

many neurons in the auditory midbrain express GluN2C and/or GluN2D NMDA receptor

subunits, which are less sensitive to Mg^{2+} block than the more commonly expressed GluN2A/B

subunits. We showed that GluN2C/D-containing receptors conducted current at resting

membrane potential and enhanced temporal summation of synaptic inputs. In a model, we

show that GluN2C/D-containing receptors provide additive gain for input-output functions

driven by trains of synaptic inputs. In line with this, we found that blocking GluN2C/D-

containing NMDA receptors in vivo decreased both spontaneous firing rates and firing evoked

by amplitude-modulated sounds.

Abstract

Along the ascending auditory pathway, there is a broad shift from temporal coding,

which is common in the lower auditory brainstem, to rate coding, which predominates in

auditory cortex. This temporal-to-rate transition is particularly prominent in the inferior

colliculus (IC), the midbrain hub of the auditory system, but the mechanisms that govern how

individual IC neurons integrate information across time remain largely unknown. Here, we

report the widespread expression of *Glun2c* and *Glun2d* mRNA in IC neurons. GluN2C/D-

containing NMDA receptors are relatively insensitive to voltage-dependent Mg^{2+} block, and thus

can conduct current at resting membrane potential. Using in situ hybridization and

pharmacology, we show that VIP neurons in the IC express GluN2D-containing NMDA receptors

that are activatable by commissural inputs from the contralateral IC. In addition, GluN2C/D-

containing receptors have much slower kinetics than other NMDA receptors, and we found that

GluN2D-containing receptors facilitate temporal summation of synaptic inputs in VIP neurons.

In a model neuron, we show that a GluN2C/D-like conductance interacts with the passive

membrane properties of the neuron to alter temporal and rate coding of stimulus trains.

Consistent with this, we show in vivo that blocking GluN2C/D-containing receptors decreases

both the spontaneous firing rate and the overall firing rate elicited by amplitude-modulated

(AM) sounds in many IC neurons. These results suggest that GluN2C/D-containing NMDA

receptors influence rate coding for auditory stimuli in the IC by facilitating the temporal

integration of synaptic inputs.

Introduction

The inferior colliculus (IC) in the midbrain plays a critical role in auditory processing,

functioning as a hub of integration in the auditory pathway (Adams, 1979; Oliver and Huerta,

1992) and as a site where phase-locked ascending inputs are transformed into a firing rate code

(Hewitt and Meddis, 1994; Joris et al., 2004). However, the cellular and circuit mechanisms that

govern how IC neurons integrate synaptic inputs and convert temporal to rate codes remain

largely unknown. In other brain regions, the slow kinetics of NMDA receptors (NMDARs) can

play a critical role in temporal integration of synaptic inputs (D'Angelo et al., 1995; Lo et al.,

2013). Intriguingly, previous studies have found that many IC neurons express NMDARs that are

activated at resting membrane potential (Kelly and Zhang, 2002; Ma et al., 2002; Wu, 2004;

Oberle et al., 2022). In line with this previous work, we recently found in adult mice that

optogenetic stimulation of commissural inputs elicited excitatory postsynaptic potentials

(EPSPs) in IC VIP neurons that had a significant contribution from NMDARs at resting membrane

potential (Goyer et al., 2019).

These results suggest that NMDARs that conduct current at resting membrane potential

are a common feature of IC circuits, but this is unusual. At most glutamatergic synapses,

activation of AMPARs is required to depolarize the membrane potential and remove voltage-

dependent Mg^{2+} block of NMDARs, allowing NMDARs to contribute to the response. One

potential explanation for this phenomenon is that NMDARs in IC neurons may be less sensitive

to Mg^{2+} block due to their subunit composition (Paoletti et al., 2013). NMDARs are

heterotetramers comprised of two obligatory GluN1 subunits and a combination of GluN2(A-D)

and/or GluN3(A-B) subunits (Traynelis et al., 2010). A residue in the GluN2 subunit confers Mg^{2+}

sensitivity (Monyer et al., 1992; Siegler Retchless et al., 2012), with GluN2C/D subunit-

containing receptors being much less sensitive to Mg^{2+} block than GluN2A/B subunit-containing

receptors. Inclusion of GluN2C/D subunits in NMDARs thus gives rise to receptors with

significant currents at resting membrane potential, as has been demonstrated in the mouse

barrel cortex (Binshtok et al., 2006).

In addition to their differences in Mg^{2+} sensitivity, NMDARs containing GluN2C or

GluN2D subunits have slower kinetics than NMDARs with GluN2A or GluN2B subunits (Paoletti

et al., 2013). These slow kinetics can expand the time window for temporal summation of

synaptic inputs, as seen in neocortex, where GluN2C/D-containing receptors enable pyramidal

neurons to integrate intracortical inputs over broader time windows than GluN2A/B-containing

receptors (Kumar and Huguenard, 2003).

Here, we tested whether GluN2C/D-containing NMDARs mediate NMDAR currents in

the IC and the role of these receptors in synaptic integration and sound processing. We found

that IC VIP neurons, a class of excitatory principal neurons (Goyer et al., 2019), express

NMDARs that activate at resting potential. We show that VIP neurons express *Glun2d* mRNA

and that many other IC neurons express *Glun2c* and/or *Glun2d* mRNA, pointing to a prominent

role for GluN2C/D-containing NMDARs in the IC. NMDAR-mediated responses in VIP neurons

were sensitive to GluN2C/D-specific pharmacology and were elicited by commissural inputs

from the contralateral IC. Additionally, GluN2C/D-specific antagonists decreased temporal

summation of commissural inputs in VIP neurons. In a model neuron, addition of an NMDAR-

like conductance lacking Mg^{2+} sensitivity altered temporal and rate coding. In line with this,

during in vivo recordings from awake mice, local infusion of a GluN2C/D-specific antagonist

decreased spontaneous and sound-evoked firing rates in many IC neurons. Overall, these

results provide a molecular mechanism for the presence of NMDAR currents at resting potential

in many IC neurons and show that the widespread expression of GluN2C/D subunits in the IC

influences sound processing.

Materials and Methods

Animals

All experiments were approved by the University of Michigan Institutional Animal Care

and Use Committee and were in accordance with NIH guidelines for the care and use of

laboratory animals. Animals were kept on a 12-hour day/night cycle with ad libitum access to

food and water. To visualize VIP neurons, VIP-IRES-Cre mice (*Vip^TM1(cre)Zjh*/J, Jackson Laboratory,

stock #010908) (Taniguchi et al., 2011) were crossed with Ai14 reporter mice (B6.Cg-

Gt(ROSA)26Sor^TM 14(CAG–tdTomato)Hze/J, Jackson Laboratory, stock #007914) (Madisen et al., 2010)

so that the fluorescent protein tdTomato was expressed in VIP neurons (Beebe et al., 2022).

Since mice on the C57BL/6J background undergo age-related hearing loss after 3 months of age

(Zheng et al., 1999), all experiments were performed on mice between postnatal days (P)30 –

66.

Brain slice preparation

Whole-cell patch-clamp recordings were performed in acutely prepared IC slices from

VIP-IRES-Cre x Ai14 mice. Male (n = 18) and female (n = 15) mice aged P30 – P66 were used.

Mice were deeply anesthetized with isoflurane and then rapidly decapitated. The brain was

removed and the IC was dissected in 34 °C artificial cerebrospinal (ACSF) solution containing (in

mM): 125 NaCl, 12.5 glucose, 25 $NaHCO_3$, 3 KCl, 1.25 NaH_2PO_4, 1.5 $CaCl_2$, 1 $MgSO_4$, 3 sodium

pyruvate and 0.40 L-ascorbic acid (Acros Organics), bubbled with 5% CO_2 in 95% O_2. 200 μm

coronal slices containing the IC were cut using a vibrating microtome (VT1200S, Leica

Biosystems). Slices were incubated at 34 °C for 30 min in ACSF bubbled with 5% CO_2 in 95% O_2

and then placed at room temperature for at least 30 min before initiating recordings.

Recordings were targeted to tdTomato-expressing VIP neurons in the central nucleus of the IC

using a Nikon FN1 or Olympus BX51 microscope. All chemicals were obtained from Thermo

Fisher Scientific unless otherwise noted.

Voltage-clamp electrophysiological recordings

Slices were placed in a recording chamber and continuously perfused at a rate of ~2

ml/min with 34 °C ACSF bubbled in 5% CO_2/95% O_2. Whole-cell voltage-clamp recordings were

performed with an Axopatch 200A patch clamp amplifier (Axon Instruments). For each

recording, series resistance compensation was performed using 80% prediction and 80%

correction, and whole cell capacitance was compensated. Series resistance for all neurons

included in this study was <15 MΩ. Data were low-pass filtered at 10 kHz, sampled at 50 kHz

with a National Instruments PCIe-6343 data acquisition board, and acquired using custom

software written in Igor Pro (Sutter Instrument). Recording pipettes were pulled from

borosilicate glass capillaries (outer diameter 1.5 mm, inner diameter 0.86 mm, Sutter

Instrument) with a P-1000 microelectrode puller (Sutter Instrument) and filled with an internal

solution containing (in mM): 115 CsOH, 115 D-gluconic acid, 7.76 CsCl, 0.5 EGTA, 10 HEPES, 10

Na_2 phosphocreatine, 4 MgATP, 0.3 NaGTP, supplemented with 0.1% biocytin (w/v), pH

adjusted to 7.4 with CsOH and osmolality to 290 mmol/kg with sucrose. Voltage clamp

recordings were not corrected for the liquid junction potential. All electrophysiological

recordings were targeted to neurons in the central nucleus of the IC, but it is possible a small

number of recordings were performed in the dorsal cortex of the IC.

To apply glutamate puffs to brain slices, we pulled puffer pipettes from borosilicate

glass (outer diameter 1.5 mm, inner diameter 0.86 mm, Sutter Instrument) to a resistance of

$3.5 - 5.0$ MΩ using a P-1000 microelectrode puller and filled them with 300 μM glutamate

(Sigma, # G5889) dissolved in a vehicle solution containing (in mM): 125 NaCl, 3 KCl, 12.5

glucose and 3 HEPES. The solution was balanced to a pH of 7.40 with NaOH. Puffer pipettes

were connected to a pressure ejection system built based on the OpenSpritzer design (Forman

et al., 2017). The tips of puffer pipettes were placed near the soma of the recorded cell, and 10

ms puffs were presented either 10 ms or 30 ms apart (30 ms for drug conditions) with 5 puffs

presented per condition. Puffs containing only the vehicle solution did not elicit any response in

the neurons tested (data not shown).

To examine whether NMDAR activation in VIP neurons at resting membrane potential is

mediated by NMDARs containing GluN2C/D subunits, we performed whole-cell patch-clamp

recordings targeted to VIP neurons as described above. To isolate the contribution of NMDARs,

recordings were performed in the presence of 10 μM NBQX, an AMPAR antagonist. Glutamate

was puffed onto VIP neurons during bath application of 1.5 μM PPDA (Hello Bio, #HB0531), a

GluN2C/D specific antagonist (Lozovaya et al., 2004; Li et al., 2020; Jing et al., 2022), or 20 μM

CIQ (Hello Bio, HB0197), a GluN2C/D specific positive allosteric modulator (Mullasseril et al.,

2010; Feng et al., 2014; Zhang et al., 2014; Nouhi et al., 2018; Liu et al., 2021). The

concentrations of PPDA and CIQ used throughout this study are based on previous studies that

used PPDA (Lozovaya et al., 2004; Li et al., 2020; Jing et al., 2022) or CIQ (Feng et al., 2014,

2014; Nouhi et al., 2018; Liu et al., 2021). All drugs were washed in for 10 minutes before

testing how the drugs affected responses to glutamate puffs.

CIQ and PPDA were prepared in 0.04% and 0.003% DMSO (volume/volume),

respectively. To test for vehicle effects, we performed control experiments where puff-elicited

EPSCs were compared between control and DMSO (vehicle) conditions. No differences in puff-

elicited EPSCs were found after DMSO wash-in (Linear mixed model tests: amplitude, β = -5.96,

p = 0.42, CI [-20.2, 8.28], halfwidth, β =7.50, p = 0.36, CI [-8.36, 23.37], rise time, β = 13.87, p =

0.53, CI [-30.16, 57.91], decay tau, β = -115.62, p = 0.20, CI [-281.84, 50.60]).

Mg^{2+}-free voltage-clamp recordings

To determine what proportion of NMDARs on VIP neurons contain GluN2C/D subunits

vs GluN2A/B subunits, we performed whole-cell patch-clamp recordings targeted to VIP

neurons as described above, except immediately prior to patching brain slices were transferred

to a Mg^{2+}-free ACSF solution containing (in mM): 125 NaCl, 12.5 glucose, 25 $NaHCO_3$, 3 KCl, 1.25

NaH_2PO_4, 2.5 $CaCl_2$, 0 $MgSO_4$, 3 sodium pyruvate and 0.40 L-ascorbic acid (Acros Organics),

bubbled with 5% CO_2 in 95% O_2. To isolate the contribution of NMDARs, recordings were

performed in the presence of 10 μM NBQX. To determine whether GluN2A/B-containing

NMDARs were present on VIP neurons, we first performed control glutamate puffs and then

puffs after bath application of 1.5 μM PPDA, followed by puffs after bath application of 100 μM

D-AP5, a broad NMDAR antagonist. All drugs were washed in for 10 minutes before testing how

the drugs affected the response to glutamate puffs.

Intracranial virus injections

Mice used for intracranial injections of recombinant adeno-associated viruses (AAVs)

were between ages P26 – 39. Mice were anesthetized with 1-3% isoflurane (Piramal Critical

Care, # NDC 66794-017-25) and body temperature was maintained using a homeothermic

heating pad. To minimize postoperative pain, an injection of carprofen (5 mg/kg, CarproJect,

Henry Schein Animal Health) was administered subcutaneously. The scalp was shaved using

scissors and an incision was made along the midline of the scalp to expose the skull. The

injection site was identified using previously validated coordinates for the IC (all coordinates are

relative to the skull surface at lambda; in μm: 900 caudal, 1000-1250 lateral, 1850 deep, with

two penetrations, one for each lateral coordinate) and a craniotomy was drilled using a

micromotor drill (K.1050, Foredom Electric Co.) with a 0.5 mm burr (Fine Science Tools). Glass

injection pipettes were pulled from capillary glass (Drummond Scientific Company) using a P-

1000 microelectrode puller (Sutter Instruments) and cut on a diagonal for a beveled opening of

~20 μm. Pipettes were first backloaded with mineral oil and then front filled with

AAV1.Syn.Chronos-GFP.WPRE.bGH (Addgene, #59170-AAV1, titer: 1.4e13, 2.2e13),

AAV5.Syn.Chronos-GFP.WPRE.bGH (Addgene, #59170-AAV5, titer: 5.3e12), or

AAV9.Syn.Chronos-GFP-WPRE.bGH (University of North Carolina Vector Core, Addgene plasmid

#59170, titer: 4.5e12) (Klapoetke et al., 2014). The virus was injected using a NanoJect III

nanoliter injector (Drummond Scientific Company) connected to an MP-285 micromanipulator

(Sutter Instrument). IC injections were made in two penetrations along the medial-lateral axis

that were 250 μm apart, and viral deposits were made at 250 μm intervals along the ventral-

dorsal axis for a total of 4 deposits at the more medial site and 3 deposits at the more lateral

site. 20 nL of virus was deposited per injection, for a total of 150 nL virus injected per IC. After

the injections were completed, the scalp was closed either by suturing with Ethilon 6-0 (0.7

metric) nylon sutures (Ethicon USA, LLC) or applying Vetbond (3M, #1469SB) on top of the

closed incision. For postoperative analgesia, 0.5 mL of 2% Lidocaine hydrochloride jelly (Akorn

Inc) was placed on top of the sutures. Mice were observed for 1 hr for indications of pain or

distress and then returned to the vivarium once they were ambulatory. Mice were monitored

daily until sutures fell out and the wound was completely healed. Sutures remaining on the 10th

post-operative day were manually removed.

Current-clamp recordings

Slices were placed in a recording chamber and continuously perfused at a rate of ~2

ml/min with 34 °C oxygenated ACSF. Whole-cell current-clamp recordings were performed with

a BVC-700A patch clamp amplifier (Dagan Corporation). Data were low-pass filtered at 10 kHz,

sampled at 50 kHz with a National Instruments PCIe-6343 data acquisition board, and acquired

using custom software written in Igor Pro. Recording pipettes were pulled from borosilicate

glass pipettes (outer diameter 1.5 mm, inner diameter 0.86 mm, Sutter Instrument) with a P-

1000 microelectrode puller (Sutter Instrument) and filled with an internal solution containing

(in mM): 115 K-gluconate, 7.73 KCl, 0.5 EGTA, 10 HEPES, 10 Na_2phosphocreatine, 4 MgATP, 0.3

NaGTP, supplemented with 0.1% biocytin (w/v), pH adjusted to 7.3 with KOH and osmolality to

290 mmol/kg with sucrose. All membrane potentials for current-clamp recordings were

corrected for an 11 mV liquid junction potential.

Optogenetics

Current-clamp recordings were conducted 2 – 4 weeks after virus injections to allow

time for Chronos expression. Recordings were conducted as described above, except that brain

slices were prepared and incubated under red light to limit Chronos activation. To verify the

virus injection location in IC injections, fluorescence was visualized under the microscope

during the recording session. Recordings were targeted to neurons contralateral to the

injection site.

Chronos was activated using 2-10 ms pulses of 470 nm light emitted by a blue LED

coupled to the epi-fluorescence port of the microscope and delivered to the slice through a

0.80 NA 40x water immersion objective. Blue light flashes illuminated the entire field of the

0.80 NA 40x objective, corresponding to optical power densities of 6 to 48 mW/mm^2. Optical

power was set using the minimum stimulation needed to elicit an EPSP from the recorded

neuron. Recording sweeps with light flashes were repeated 10-30 times with 10 ms between

light flashes.

RNAScope in situ hybridization

Fluorescent in situ hybridization was performed using the RNAscope Multiplex

Fluorescent V2 Assay (Advanced Cell Diagnostics, catalog # 320850). Our methods were

identical to those previously described (Beebe et al., 2022) and followed manufacturer

recommendations (Wang et al., 2012). Briefly, one male (P47) and two female (P45) mice were

deeply anesthetized with isoflurane and brains were rapidly removed and frozen on dry ice.

Brains were placed in a -80 °C freezer until the day of slicing. Prior to slicing, brains were

equilibrated at -20 °C for 1 hour. Brains were sliced into 15 µM sections using a cryostat at -20

°C and sections were mounted on Superfrost Plus slides (Fisher Scientific, catalog # 22037246).

For each mouse, five representative sections were chosen spanning the caudal—rostral axis.

Sections were fixed using 10% neutral buffered formalin (Sigma-Aldrich, catalog # HT501128)

and dehydrated using repeated washes in 50%-100% ethanol. Slides were dried using a Kim

wipe and a hydrophobic barrier was drawn around each section. Slices were next incubated in

hydrogen peroxide for 10 minutes at room temperature, followed by application of Protease IV

for 30 minutes. Probes for *tdTomato*, *Glun2c*, and *Glun2d* (all experimental slices except

controls; Advanced Cell Diagnostics catalog #: 317041-C3, 445581-C2, 425951, respectively),

along with positive and negative controls (one control slice each), were applied to slices and

incubated for 2 hours at 40 °C. The probes were amplified three times for 30 minutes each time

at 40 °C using AMP 1-3. The signal was developed using the HRP for each channel and then opal

dyes diluted at 1:1000 were assigned for each channel: *tdTomato* was assigned to Opal 690

(Akoya Bioscience, catalog # FP1497001KT), *Glun2c* was assigned to Opal 570 (Akoya

Bioscience, catalog # FP1488001KT), and *Glun2d* was assigned to Opal 520 (Akoya Bioscience,

catalog # FP1487001KT). Slices were counterstained with DAPI and coverslipped with ProLong

Gold antifade mountant (Fisher Scientific, catalog # P36934). Slices were imaged within 1 week

of performing the assay using the 0.75 NA 20X objective on a Leica TCS SP8 laser scanning

confocal microscope (Leica Microsystems) at 2 μm depth intervals. Emission wavelengths were

adjusted for each channel as follows: DAPI (405 nm laser, 410-441 nm), *Glun2d* (488 nm laser,

491-528 nm), *Glun2c* (552 nm laser, 565-583 nm), *tdTomato* (638 nm laser, 690-727 nm).

Co-localization of *Glun2c* and *Glun2d* in *tdTomato*-positive neurons was quantified

manually using Neurolucida 360 (MBF Bioscience). One side of the IC (left or right) was selected

randomly for quantification in each slice. *tdTomato*-positive neurons were first identified by

placing a marker on top of the cell, and then *Glun2c* and *Glun2d* fluorescence was quantified

separately for each marked cell. Cells were considered positive for *Glun2c* or *Glun2d* when one

or more puncta co-localized with the *tdTomato* fluorescence. All *tdTomato*-positive neurons

counted co-labeled with DAPI.

Subdivisions of the IC were determined for each IC slice used in the above analysis by

comparison to a reference series of sections from a control C57BL/6J mouse aged P49 that

were immunolabeled for GAD67 and GlyT2. This pattern of labeling is routinely used to identify

IC subdivisions (Choy Buentello et al., 2015; Beebe and Schofield, 2021), including in our

previous studies (Silveira et al., 2020; Anair et al., 2022).

Analysis of electrophysiological recordings

Amplitude, halfwidth, rise time, and decay time constant measurements were made

using custom algorithms in Igor Pro 8. Voltage-clamp data were low-pass filtered at 3 kHz prior

to analysis, except for the commissural optogenetic experiment (**Figure 5**), where data were

low-pass filtered at 1 kHz and the following function was fit to each response, which was then

analyzed:

310
$$f(x) = y_0 + amp\left(1 - e^{\frac{-(x-x_0)}{\tau_{rise}}}\right) * e^{\frac{-(x-x_0)}{\tau_{decay}}}$$

For the temporal summation experiment (**Figure 6**), individual sweeps from each condition

were averaged and measurements were made from the average waveform.

Headbar implantation

Mice used for in vivo experiments were either CBA/CaJ (N = 1 female, age at recording =

P51), CBA/C57 (N = 7; 4 females, age range at recording = P81-127), or VIPxAi14 (N = 2

females, ages 50, 53 at recording). Mice were anesthetized with 1-3% isoflurane (Piramal

Critical Care, # NDC 66794-017-25) and body temperature was maintained using a

homeothermic heating pad. To minimize postoperative pain, injections of carprofen (5 mg/kg,

CarproJect, Henry Schein Animal Health) and buprenorphine (0.1 mg/kg) were administered

subcutaneously. The scalp was shaved using scissors and an incision was made along the

midline of the scalp to expose the skull. The inferior colliculus was identified using previously

validated coordinates for either the IC (all coordinates are relative to lambda; in μm: 900

caudal, 1125 lateral) and marked using a surgical marker. The skull was numbed with 0.2 mL

bupivacaine and then the periosteum was removed using a scalpel and the skull was scored.

After the skull was prepped, a thin layer of Metabond (C&B Metabond from Parkell, cat #s

S398, S399, S371) was applied using a fine paintbrush and allowed to dry for 10 minutes. A thin

layer of dental acrylic (Henry Schein, Inc., # 1259208 and # 1250143) was then applied on top of

the Metabond, and then dental acrylic was applied to a custom-designed headbar which was

placed on the skull. Layers of dental acrylic were added to the headbar to secure it to the skull,

and then the opening above the IC was filled with silicon elastomer (Smooth-On Inc. Body

Double FAST) and secured with a thin strip of dental acrylic. Mice were observed for 1 hr for

indications of pain or distress and then returned to the vivarium once they were ambulatory.

For postoperative analgesia, an additional injection of buprenorphine (0.1 mg/kg) was given the

day following the surgery, and mice were monitored daily for 7 days.

In vivo electrophysiological recordings with drug delivery

Mice were allowed to recover for 3 days following headbar implantation and then began

habituation to the head fixation setup, which consisted of 1 day of handling within the cage and

3-5 days of head fixation in the sound booth for increasing time duration (10 minutes, 20

minutes, 40 minutes, 60 minutes, 80 minutes). After the habituation period, a craniotomy was

surgically drilled above the left IC using the procedures outlined here under "intracranial virus

injections". Mice were allowed to recover for 1 hour, and then placed in the head fixation

setup. For drug delivery, drug delivery pipettes were pulled identically to those used for virus

injection in the "intracranial virus injections" section. Drug delivery pipettes were affixed at a

~30 degree angle to recording pipettes (pulled identically to as described in "current clamp

recordings") using melted beeswax (Fisher Scientific, # S25192A) such that the tips of both

pipettes were 50-200 μm apart. The drug delivery pipette was backfilled with 20 μM DQP

(Tocris, #4491) diluted in the in vivo internal solution (in mM, 1.5 $CaCl_2$, 135 NaCl, 1 $MgCl_2$, 5.4

KCl, 5 HEPES) and connected to FEP tubing which was attached to a 10 μL Hamilton syringe in a

syringe pump (World Precision Instruments Micro4 MicroSyringe Pump Controller). The

recording pipette was backfilled with in vivo internal solution and advanced into the brain

slowly using an MP-285 micromanipulator (Sutter Instruments). Cells were identified by an

increase in resistance and often the presence of spikes in the test pulse. Loose seals (11-30 MΩ)

were made to cells using application of a small amount of negative pressure.

Sounds were presented to awake mice using a calibrated, free-field electrostatic

speaker (Tucker-Davis Technologies, ES1) positioned ~10 cm in front of the right ear

(contralateral to the IC recording site) at a ~45 ° angle relative to the midline of the mouse.

Sounds consisted of 1 s white noise bursts (4-64 kHz, 70 dB SPL) that were amplitude

modulated at 100% modulation depths using modulation frequencies ranging from 16 – 512 Hz

at 4 steps/octave. Each stimulus was repeated 8 times in a pseudorandomized order. After

collection of the control data, DQP or vehicle solution was infused into the brain at a rate of 0.1

µL/min while the sound presentation was repeated. Mice were recorded from no more than

once per day and for fewer than 7 sessions. After recording sessions were finished, the location

of the recordings was verified post hoc by dissection.

Model neuron

The model neuron used here was based, with minimal changes, on the fast-spiking

interneuron model described previously by Wang and Buzsáki (Wang and Buzsáki, 1996).

Simulated currents interacted with passive membrane properties according to the equation

$C_m \frac{dV}{dt} = -I_{Na} - I_K - I_L - I_{syn}$ where C_m = 1 µF/cm^2. Leak current was varied in the model to

approximate a range of membrane resistances that might be found in the IC, and the leak

reversal potential, E_L, was set to -65 mV.

Action potentials were modeled using Hodgkin-Huxley-type, voltage-dependent Na$^+$ and

K$^+$ conductances (Hodgkin and Huxley, 1952). The transient sodium current is described as

$I_{Na} = g_{Na} m^3{}_\infty h(V - E_{Na})$ where the activation variable m is assumed to be fast, and is thus

substituted by its steady state function $m_\infty = \frac{\alpha_m}{\alpha_m + \beta_m}$ where $\alpha_m(V) = -0.1 \frac{V+35}{e^{-0.1(V+35)}-1}$ and

$\beta_m(V) = 4e^{-\frac{V+60}{18}}$. The inactivation variable h obeys first order kinetics, where $\frac{dh}{dt} =$

$\varphi(\alpha_h(1-h) - \beta_h h)$, $\alpha_h(V) = 0.07 e^{-\frac{V+58}{20}}$, and $\beta_h(V) = \frac{1}{e^{-0.1(v+28)}+1}$. The sodium conductance

was set to g_{Na} = 35 mS/cm^2 with a reversal potential E_{Na} = 55 mV.

The delayed rectifier potassium channel was modeled as $I_K = g_K n^4(V - E_K)$, where

the activation variable n is given by $\frac{dn}{dt} = \varphi(\alpha_n(1-n) - \beta_n n)$. Here, $\alpha_n(V) = -0.01 \frac{V+34}{e^{-0.1(V+34)}-1}$

and $\beta_n(V) = 0.125e^{-\frac{V+44}{80}}$, and we set the potassium conductance to $g_K = 9$ mS/cm^2 with a

reversal potential E_K = -90 mV.

Simulations using the model neuron were performed in MATLAB Version 2022a-2024a

(MathWorks) using custom MATLAB scripts adapted from functions provided in (Börgers,

Christoph, 2017). All code used for the model simulations can be found at:

https://github.com/audreydrotos/NMDA_modeling .

Statistical analyses

Data analysis was performed using custom algorithms and statistical packages in Igor

Pro 8, MATLAB R2021a-2024a (MathWorks), and R 4.1.0 (The R Project for Statistical

Computing). Statistical hypothesis testing was performed using two approaches. Comparisons

between control and drug condition data for in vitro experiments were made using linear mixed

model (LMM) analyses, which were conducted in R using the "lme4" and "lmerTest" packages

(Bates et al., 2015; Kuznetsova et al., 2017). Note that in cases where two groups were being

compared, the *p*-values from the LMM are identical to those obtained from a paired-samples *t*-

test (Shravan Vasishth et al., 2022). For comparisons between control and vehicle or drug

conditions in vivo, we used a Wilcoxon Signed Rank test, which was performed using the

'signrank' function in MATLAB. Effects were considered significant when $p < 0.05$.

For in vitro electrophysiology data involving repeated measures from individual cells,

effect sizes and confidence intervals are shown using Gardner-Altman estimation plots (two

groups) or Cumming estimation plots (three groups). These plots were made using custom

MATLAB scripts modeled after the DABEST package (Ho et al., 2019). Both types of estimation

plots use parallel coordinate plots to show the responses of individual cells for each condition,

and these are accompanied by histograms showing the range and 95% confidence intervals of

the paired mean differences expected between each drug condition and the control based on a

bootstrap analysis. Our approach to generating these plots has been detailed previously

(Rivera-Perez et al., 2021) and was influenced by the estimation statistics approach (Bernard,

2019; Calin-Jageman and Cumming, 2019). While estimation plots are provided to give

additional insight into the data, all determinations of statistical significance were based on the

standard null-hypothesis statistical tests detailed in the preceding paragraph.

Results

NMDAR currents in VIP neurons are sensitive to GluN2C/D-specific pharmacology

Several previous studies have noted that NMDARs often contribute to subthreshold

postsynaptic responses in IC neurons (Kelly and Zhang, 2002; Ma et al., 2002; Wu, 2004; Oberle

et al., 2022) . Similarly, we recently found that NMDARs contributed to subthreshold EPSPs

elicited by commissural inputs to IC VIP neurons (Goyer et al., 2019). NMDAR currents at resting

potential are unusual, as most NMDARs in the adult brain have a voltage-dependent Mg^{2+} block

which prevents ions from flowing through the channel pore at or near resting potential

(Traynelis et al., 2010; Paoletti et al., 2013). However, sensitivity to Mg^{2+} block depends on the

subunit composition of the receptor. While receptors containing GluN2A/B subunits are highly

sensitive to Mg^{2+} block, receptors containing GluN2C/D subunits are less sensitive, allowing

them to conduct current even at hyperpolarized potentials (Siegler Retchless et al., 2012).

Based on this, we hypothesized that VIP neurons express NMDARs containing GluN2C and/or

GluN2D subunits.

To test this hypothesis, we voltage-clamped VIP neurons at -70 mV and used a puffer

pipette containing 1 mM glutamate to elicit EPSCs (**Figure 1A**). We found that 10 ms glutamate

puffs reliably elicited EPSCs in VIP neurons (**Figure 1B**). We next performed puffs in the

presence of 1.5 µM PPDA, a GluN2C/D-selective antagonist (Lozovaya et al., 2004; Li et al.,

2020; Jing et al., 2022), and then in the presence of 20 µM CIQ, a GluN2C/D-selective positive

allosteric modulator (Mullasseril et al., 2010; Feng et al., 2014; Zhang et al., 2014; Nouhi et al.,

2018; Liu et al., 2021). PPDA significantly decreased the amplitude (LMM: $p = 0.038$) of the

elicited EPSC. Our experimental design did not favor large CIQ effects since we used a

saturating glutamate concentration (1 mM) in our puff solution. At the population level, CIQ did

not alter the amplitude ($p = 0.088$), halfwidth ($p = 0.15$), rise time ($p = 0.49$), or decay tau ($p = $

0.10) of the EPSC (**Figure 1C**). The sensitivity of NMDAR currents in VIP neurons to PPDA and, in

some cases to CIQ, suggests that at least a portion of the NMDARs expressed in VIP neurons

contain GluN2C/D subunits.

Many IC neurons in adult mice express *Glun2c* and/or *Glun2d* mRNA, with VIP neurons

predominately expressing *Glun2d*

GluN2C and GluN2D subunits are similar in their sensitivity to Mg^{2+} block, but expression

levels of *Glun2c* and *Glun2d* mRNA change dramatically and in different directions during

development (Paoletti et al., 2013). GluN2D-containing NMDARs (along with GluN2B-containing

NMDARs) are primarily found early in development (Wenzel et al., 1996) and are largely

replaced with GluN2A and GluN2C subunits by adulthood (Paoletti et al., 2013), with some

exceptions in small populations of interneurons, such as those found in hippocampus (Monyer

et al., 1992). In many cases, the developmental switch in NMDAR subunit composition is

thought to be driven by sensory experience (Traynelis et al., 2010). Expression of GluN2

subunits is also spatially regulated: while GluN2A and GluN2B-containing NMDARs are typically

found at the synapse, GluN2C and GluN2D-containing NMDARs are often found at extrasynaptic

sites (Paoletti et al., 2013). However, growing evidence points to GluN2C/D expression at

synapses, such as in striatum (Logan et al., 2007) and substantia nigra (Brothwell et al., 2008).

Based on these precedents, we hypothesized that IC VIP neurons in adult animals

express GluN2C subunits. To test this hypothesis, we used the RNAScope assay to perform

fluorescence in situ hybridization on IC brain slices from three VIP-IRES-Cre x Ai14 mice aged

P45-47 (**Figure 2A,B**). Using probes for *tdTomato*, *Glun2c*, and *Glun2d* mRNA, we were

surprised to find that 91.4% of VIP neurons expressed *Glun2d* and only 8.1% expressed *Glun2c*.

More specifically, 84.3% of VIP neurons expressed only *Glun2d*, 1.0% of VIP neurons expressed

only *Glun2c*, and 7.1% of VIP neurons expressed both. The remaining 7.7% of VIP neurons did

not express *Glun2c* or *Glun2d* (**Table 1**, **Figure 3A**).

We also investigated whether NMDAR subunit expression differed among VIP neurons

present in the three major IC subdivisions: central nucleus, dorsal cortex, and lateral cortex. We

found that the percentage of VIP neurons expressing various NMDAR subunits was similar

across all IC subdivisions (**Table 2**, **Figure 3B**). These results suggest that VIP neurons

preferentially express GluN2D rather than GluN2C subunits, regardless of their location in the

IC.

Importantly, our results also show that *Glun2c* and *Glun2d* mRNA is widely expressed in

the IC (**Figure 2A**). This suggests that *Glun2c* and *Glun2d*-containing NMDARs are a prominent

feature in the IC and likely have a significant and widespread impact on synaptic computations.

VIP neurons also express GluN2A/B-containing NMDARs

We next investigated whether IC VIP neurons also express GluN2A or GluN2B-containing

receptors. In the adult brain, GluN2A-containing NMDARs are more commonly found at

synapses and confer more typical properties of the NMDA receptor, including sensitivity to

Mg^{2+} block at rest (Siegler Retchless et al., 2012). To determine whether VIP neurons express

GluN2A/B-containing NMDARs in addition to GluN2C/D-containing NMDARs, we targeted VIP

neurons for whole-cell recordings and puffed glutamate in the presence of AMPAR antagonist

NBQX (10 µM, **Figure 4A**). To examine the contributions of all NMDARs to the resulting current,

we removed Mg^{2+} from the ACSF and replaced it with equimolar Ca^{2+} to maintain the overall

concentration of divalent ions. We found that puffs of glutamate elicited much larger currents

in the Mg^{2+}-free condition than we observed in the earlier puff experiments (compare **Figures**

1C, 4C).

Next, we bath applied 1.5 µM PPDA, a GluN2C/D-specific antagonist, and as expected,

we found that this significantly decreased the amplitude (LMM: $p = $ 9.26e-4), halfwidth ($p = $

8.6e-4), rise time ($p = $ 8.45e-3), and decay time constant ($p = 0.028$) of the response, again

suggesting that a portion of the NMDAR current in VIP neurons is mediated by GluN2C/D-

containing NMDARs (**Figure 4C**). Next, we bath applied 100 µM D-AP5, an NMDAR antagonist

that blocks NMDARs regardless of GluN2 subunit identity and found that this completely

abolished the response in all neurons (amplitude, $p = 1.71e-4$). Overall, 5 out of 6 VIP neurons

had both PPDA sensitive and Insensitive currents, indicating activation of both GluN2A/B and

GluN2C/D currents. On average in these cells, 82.0% ± 26.2% of the current was sensitive to

PPDA (mean ± standard deviation, **Figure 4D**). Two neurons had current that was completely

abolished by PPDA indicating exclusive activation of GluN2C/D-containing receptors. These

results suggest that individual VIP neurons can express NMDARs that differ by subunit

composition and are thus specialized for different roles in the cell.

Commissural inputs to VIP neurons activate GluN2C/D-containing NMDARs

We previously found that inputs from the contralateral IC via the IC commissure activate

NMDARs on VIP neurons at resting potential (Goyer et al., 2019). To test whether these

responses were also mediated by GluN2C/D-containing NMDAR receptors, we repeated these

experiments in the presence of GluN2C/D-specific drugs. Commissural projections were labeled

with the excitatory opsin Chronos by injecting AAV-Synapsin-Chronos-GFP (serotype 1, 5, or 9)

into the right IC of VIP-IRES-Cre x Ai14 mice. 1.5 to 4 weeks later, we targeted whole-cell

recordings to fluorescently labeled VIP neurons in the IC contralateral to the injection site

(**Figure 5A**). We found that even in the presence of AMPAR antagonist NBQX (10 µM), flashes of

blue light elicited small EPSCs (-2 to -18 pA, **Figure 5B**). The GluN2C/D-specific positive allosteric

modulator CIQ did not alter the light-evoked response, again possibly because a saturating

concentration of glutamate was used and/or because CIQ is less effective at tri-heteromeric

NMDARs that contain a GluN2A/B subunit in addition to a GluN2C/D subunit (Mullasseril et al.,

2010) (**Figure 5C,** LMM: amplitude p = 0.86, halfwidth p = 0.064, rise time p = 0.26, decay tau

p = 0.17), but the GluN2C/D-specific antagonist PPDA completely abolished the response in all

cells tested (amplitude, LMM: p = 5.13e-3). These results support the hypothesis that

commissural inputs to VIP neurons activate NMDARs containing GluN2C/D subunits.

Activation of GluN2C/D-containing NMDARs facilitates temporal summation

Since NMDARs with GluN2C/D subunits have slower kinetics and conduct more current

at resting membrane potential than NMDARs with GluN2A/B subunits (Paoletti et al., 2013),

expression of GluN2C/D-containing NMDARs could have a particularly strong effect on the

duration of excitation elicited by glutamatergic transmission. We therefore hypothesized that

GluN2C/D-containing NMDARs in the IC facilitate temporal integration by widening the time

window for integration of synaptic inputs. To test this hypothesis, we used optogenetics to

stimulate commissural inputs to VIP neurons (**Figure 6A**). Trains of five light pulses at 30 Hz

elicited temporal summation in all the cells tested, with the second through fifth EPSPs starting

at a more depolarized potential than the first EPSP (**Figure 6B**). To test whether GluN2C/D-

containing receptors facilitate this temporal summation, we next elicited trains of input in the

presence of the GluN2C/D-containing NMDAR antagonist PPDA (1.5 μM). In one cell, the light

train elicited temporal summation great enough to elicit an action potential in the control

condition, and this was impaired in the PPDA condition, where the cell only reached action

potential threshold in one trial (**Figure 6C,D**). In trials that did not reach action potential

threshold, we assessed temporal summation by comparing the peak amplitudes of each of the

five EPSPs elicited by the train stimuli in the control and PPDA conditions (**Figure 6B**). The

amplitude of the peaks significantly increased during the train (LMM: *p* = 9.15e-6, *n* = 5, **Figure**

6E) indicating temporal summation occurred during the train. This change in amplitude was not

correlated with small changes in the resting membrane potential over the course of the

recording (r^2 = 0.001). In addition, PPDA significantly decreased the area under the curve of the

train response (LMM: *p* = 0.038, **Figure 6F**), indicating a decrease in temporal summation when

GluN2C/D-containing receptors were blocked. These results show that GluN2C/D-containing

NMDARs strongly enhance the time window for synaptic integration in VIP neurons.

GluN2C/D-containing NMDA receptors provide additive gain to rate coding in a neuron model

Our in vitro data suggest that GluN2C/D-containing receptors facilitate synaptic

integration in IC neurons, and we hypothesized that this boosts gain in the IC through an

increase in temporal summation, which could in turn promote rate coding for auditory stimuli.

To test this hypothesis, we created a neuron model based on a previously published Hodgkin-

Huxley model of fast-spiking hippocampal interneurons (Wang and Buzsáki, 1996). We based

our model on the Wang and Buzsáki model because it provides a well-known and widely used

model for neurons that exhibit sustained firing over a large range of synaptic input frequencies,

similar to what we observe for VIP neurons in vitro (Goyer et al., 2019). In our adapted model,

we included two synaptic conductances: one with kinetics similar to AMPA receptors on VIP

neurons, and one with kinetics similar to a GluN1-GluN2D-GluN2D NMDA receptor (Yi et al.,

2019). We set the maximal NMDA conductance to 25% of the maximal AMPA conductance and

set the leak conductance to 0.7 mS/cm^2 to mimic the average membrane resistance of a VIP

neuron (~200 MΩ), assuming a 15 μm diameter soma (Goyer et al., 2019). We next simulated

trains of synaptic inputs at frequencies ranging from 16 – 512 Hz, activating either the AMPA

conductance alone or both the AMPA and NMDA conductances.

We found that activation of the AMPA-only synapse by a slow train of stimuli (16 Hz)

elicited a series of phase-locked EPSPs in the model neuron (**Figure 7A**, left panel). When the

NMDA conductance was added, the slow kinetics of the NMDA receptor caused the EPSPs to

temporally sum across stimulus presentations, providing an additive gain that pushed the

neuron over action potential threshold and elicited phase-locked action potentials (**Figure 7A**,

right panel). When the input frequency was increased to 64 Hz, the neuron fired phase-locked

action potentials in the AMPA-only condition (**Figure 7B**, left panel), and addition of the NMDA

conductance again increased the overall firing rate (**Figure 7B**, right panel). To better examine

how the NMDA conductance shapes the response to inputs at different stimulus frequencies,

we simulated the response using input frequencies from 16 – 512 Hz (4 steps/octave) in both

the AMPA-only and AMPA plus NMDA conditions. We found that addition of the NMDA

conductance caused an overall increase in firing rate in response to all input frequencies tested

(**Figure 7C**, orange line). Addition of the NMDA conductance also altered phase-locking to the

response (**Figure 7D**): in the AMPA-only condition, the neuron exhibited near perfect phase

locking as soon as the input frequency was high enough to elicit spiking (blue line). When the

NMDA conductance was added (orange line), phase locking was high at lower input frequencies

but quickly became poorer as the input frequency increased, an effect due in part to the slow

kinetics of the NMDA receptor which decreased the ability of the fast, temporally precise AMPA

conductance to facilitate phase-locking in the model neuron. In line with this, addition of the

NMDA conductance shifted the phase of the input response at which the neuron fired (**Figure**

7E). These results suggest that GluN2C/D-type NMDA receptors provide additive gain to neuron

input-output functions, in the form of an upward shift along the *y*-axis of firing vs stimulus

curves (e.g., **Figure 7C**) (Silver, 2010). Our model also indicates that, as stimulus frequencies

increase, this type of additive gain increases firing rate while simultaneously decreasing the

precision of spike timing.

Membrane resistance alters modulation of rate coding by GluN2C/D-containing NMDARs

IC neurons are heterogeneous in their intrinsic physiology (reviewed in Drotos and

Roberts, 2024), and these differences may alter how AMPA and NMDA receptors contribute to

temporal and rate coding for stimuli. To examine how membrane resistance influences

temporal integration, we simulated rate and temporal modulation transfer functions (MTFs) to

input trains with frequencies between 16 – 512 Hz, as above, in model neurons where the leak

conductance (g_L) was set to 0.282, 0.50, 0.75, 1.99, and 5.3 mS/cm^2 (corresponding to

membrane resistances of 502, 283, 189, 71, and 27 MΩ, respectively, assuming 15 µm cell

diameter). At the three lowest values for g_L, neurons exhibited high-pass rate MTFs with a

relatively smooth increase in firing rate as input frequency increased (**Figure 8A-C**, left panel).

These cells also exhibited temporally precise firing at slow input frequencies, but the temporal

precision of firing fell apart at higher input frequencies, and overall, the firing rate provided a

more consistent indicator of input frequency. In addition, in these cells, the NMDA conductance

contributed additive gain to the rate MTFs that shifted the input-output functions upwards

along the y-axis, towards higher firing rates.

As the leak conductance increased to 1.99, and 5.3 mS/cm^2, we observed two major

shifts: first, neurons increasingly exhibited better phase-locking and poorer rate coding of the

input frequency, and second, addition of the NMDA conductance began to decrease, rather

than increase, the gain of the rate MTF (**Figure 8D,E**). In our model, this was due to the

additional conductance causing depolarization block, which shifted the response profile of

these cells to an onset response. Note that rate MTFs in **Figure 8D,E** decreased to one spike per

stimulus train, not zero, at higher input frequencies, and vector strengths were only calculated

when input trains elicited >4 spikes per train. While it is likely that adding low-voltage-activated

K$^+$ conductances and/or other voltage-gated conductances to the model could convert onset

firing neurons to phasic firing neurons, our main goal here was to mimic VIP neurons and other

IC neurons with sustained firing, where such conductances are not prominent (Goyer et al.,

2019). Overall, our modeling results highlight that the intrinsic physiology of a neuron

influences both how that neuron uses temporal and rate codes for stimuli and how NMDA

receptors shape these codes.

GluN2C/D-containing receptors enhance gain for encoding AM stimuli

The results from our optogenetics experiments and neuron model suggest that NMDA

receptors shape temporal and rate coding of synaptic stimuli. In the IC, temporal and rate codes

are particularly important for encoding amplitude-modulated (AM) sounds, as IC neurons can

encode the modulation frequency of AM stimuli with changes in firing rate and/or by phase-

locking their firing to the modulation waveform (Krishna and Semple, 2000). Based on our

model results, we hypothesized that GluN2C/D-containing NMDA receptors contribute to rate

coding in the IC by broadening the time window for integrating phase-locked synaptic inputs

and enhancing firing rates across different modulation frequencies. In support of this

hypothesis, Zhang and Kelly showed that blocking NMDARs decreased rate coding in IC neurons

while leaving temporal coding of AM stimuli intact (Zhang and Kelly, 2001), similar to what we

observed in our model neuron (**Figures 7,8**). To examine how GluN2C/D-containing NMDARs

influence rate coding for AM stimuli in the mouse IC, we made juxtacellular (loose-patch)

current-clamp recordings from IC neurons in awake, headfixed mice while playing 1 s

broadband noise bursts (4-64 kHz) that were amplitude modulated with modulation

frequencies of 16 – 512 Hz and 100% modulation depths (**Figure 9A**).

In line with previous studies, we found that IC neurons encode AM stimuli via changes in

firing rate (rate coding) and by phase locking their firing to the modulation waveform (temporal

coding) (example cell response, **Figure 9B**). We next found that local infusion of 20 µM DQP, a

GluN2C/D-selective antagonist, significantly decreased the firing rate during AM stimuli (signed-

rank test, *p* = 0.042, median difference = -1.11 Hz) (**Figure 9C,F**) but had no effect on the

average phase-locked responses (vector strength, signed-rank test, *p* = 0.19, median difference

= -0.008) (**Figure 9C,G**) or the phase of the modulation frequency at which the neuron fired

(**Figure 9E**). Overall, blocking GluN2C/D-containing receptors did not alter the shape of rate or

temporal modulation transfer functions, nor did it significantly alter the relative change in firing

rates (max – min) across the range of AM stimuli (dynamic range, *p* = 0.067, median difference

= -0.13 Hz, **Figure 9H**). However, blocking GluN2C/D-containing receptors did significantly

decrease the spontaneous firing rate during periods between stimulus presentations (p = 0.024,

median difference = -3.03 Hz, **Figure 9I**). Together, these results suggest that GluN2C/D-

containing receptors influence rate coding in IC neurons by providing a tonic, additive boost in

gain to both spontaneous and sound-evoked synaptic inputs.

In addition, we ran control experiments to evaluate whether changes in firing could be

due to infusion of solution into the IC or drift in the firing rate over the course of each

recording. We found that infusion of vehicle solution did not alter the firing rate during AM

stimuli (p = 0.74, median difference = -0.062 Hz), the vector strength of the response (p = 0.063,

median difference = -0.018), the dynamic range in response to AM stimuli (p = 0.086, median

difference = -0.12 Hz), or the spontaneous firing rate between sound presentations (p = 0.20,

median difference = 0.76 Hz) (vehicle panels, **Figure 9D-I**).

Discussion

In this study, we found that GluN2C/D-containing NMDARs shape synaptic integration

and auditory processing in the IC. By combining optogenetics with whole-cell recordings, we

found that NMDARs in VIP neurons are activated at resting potential by commissural

projections. By using puffs of glutamate and whole-cell recordings combined with GluN2C/D-

specific pharmacology, we showed that GluN2C/D-containing NMDARs activate at resting

potential in VIP neurons. We demonstrated that 91% of VIP neurons express mRNA for the

GluN2D subunit. We also found that *Glun2c* and *Glun2d* mRNA is prevalent throughout the IC,

including in many non-VIP neurons, suggesting that GluN2C/D-containing NMDARs exert

widespread influence over excitatory postsynaptic responses in the IC. Additionally, we

demonstrated that VIP neurons can also express GluN2A/B-containing receptors, suggesting

that different types of NMDARs may play distinct roles in these cells. Finally, we showed that

GluN2C/D-containing receptors facilitate temporal summation in VIP neurons in vitro, provide

additive gain to neuron input-output functions in silico, and enhance the spontaneous and

sound-evoked firing rates of IC neurons in vivo. Thus, NMDAR diversity shapes synaptic

integration and thus auditory processing in the IC.

VIP neurons express NMDARs with GluN2D subunits

One of the most intriguing findings from our study was that the NMDAR currents

elicited at resting potential in VIP neurons were predominately attributable to GluN2D-

containing NMDARs rather than GluN2C. This result was unexpected for two reasons: first, in

most brain regions, expression of *Glun2d* mRNA is developmentally regulated and disappears in

adulthood (Monyer et al., 1994; Wenzel et al., 1996). Second, in adult animals, GluN2D-

containing NMDARs are conventionally thought to be restricted to extrasynaptic locations, for

example, in dorsal horn neurons (Momiyama, 2000) and Golgi cells of the cerebellum (Misra et

al., 2000; Brickley et al., 2003), but there are some exceptions, such as in spinal cord laminar

cells (Hildebrand et al., 2014). In addition, NMDARs are highly motile and can rapidly diffuse

between extrasynaptic and synaptic sites (Tovar and Westbrook, 2002), as occurs during

learning. During induction of NMDAR-LTP in the medial perforant path of the dentate gyrus,

GluN2D-containing NMDARs are trafficked to the synapse where they contribute to synaptic

transmission (Harney et al., 2008). Based on this, we hypothesize that GluN2D-containing

receptors diffuse to synaptic locations on VIP neurons during learning. An alternative

hypothesis is that GluN2D-containing NMDARs are natively found in the synapse and do not

undergo a GluN2D-to-GluN2A/C switch during development.

Additionally, while we show here that many NMDARs on VIP neurons contain the

GluN2D subunit, the full subunit composition of the receptor remains unknown. NMDARs are

tetrameric receptors made up of four subunits, including two obligatory GluN1 subunits and a

combination of GluN2A-D and/or GluN3 subunits (Traynelis et al., 2010). Our in situ

hybridization results showed that 91% of VIP neurons express *Glun2d* mRNA. This could give

rise to NMDARs that are diheteromeric with two GluN1 subunits and two GluN2D subunits.

However, GluN1-GluN2D receptor assemblies have much slower kinetics than we observed in

our study (decay τ = 10-100 ms), with decay time constants on the order of 2 seconds (Vicini et

al., 1998). An alternative hypothesis is that GluN2D-containing receptors in VIP neurons are

triheteromeric GluN1-GluN2B-GluN2D complexes. These receptors display properties that are

intermediate to those of GluN1-GluN2B and GluN1-GluN2D receptors, including slightly faster

kinetics than GluN1-GluN2D assemblies (Yi et al., 2019) and a reduced sensitivity to Mg^{2+} block

compared to GluN1-GluN2B assemblies (Huang and Gibb, 2014). In addition, receptors

containing GluN2B tend to be more motile than NMDARs with GluN2A subunits (Groc et al.,

2006). Thus, this proposed GluN1-GluN2B-GluN2D assembly may also help explain how

GluN2D-containing receptors contribute to synaptic transmission in mature VIP neurons. This

conclusion is also supported by our CIQ experiments: we found a small effect of CIQ in the puff

experiments and no effect of CIQ in the optogenetic activation experiments. The ability of CIQ

to potentiate GluN2C/D-containing receptors is diminished in triheteromeric receptor

assemblies compared to diheteromeric ones (Mullasseril et al., 2010), suggesting that the

receptors studied here may be triheteromeric.

Distribution of NMDARs on VIP neurons is synapse-specific

Previous work from our lab showed that NMDAR activation at resting potential on VIP

neurons occurs with commissural inputs but not inputs from the DCN (Goyer et al., 2019). This

suggests that the distribution of NMDARs on VIP neuron synapses is pathway-dependent, with

commissural inputs targeting synapses that are enriched in GluN2D-containing receptors while

DCN inputs target synapses predominated by AMPARs and possibly GluN2A/B-containing

NMDARs. Heterogeneity in NMDAR composition at IC neuron synapses is also supported by the

results from our in vitro and in vivo experiments, where some cells exhibited greater magnitude

changes in response to blockade of GluN2C/D-containing NMDARs as compared to other cells.

Such differences in receptor distribution could underlie functional differences between these

synapses, as has been seen in other brain regions such as neocortex (Kumar and Huguenard,

2003). For example, commissural inputs may be integrated over longer time scales than those

from the DCN, and a neuron may require more DCN inputs within a smaller time window to

generate a postsynaptic spike.

NMDAR subtype distribution might also affect synaptic plasticity. GluN2A/B-containing

NMDARs have been well-studied for their role in the initiation of long-term potentiation (LTP)

(Paoletti et al., 2013). In the IC, previous studies show that electrical stimulation of the lateral

lemniscus can induce LTP in IC neurons (Hosomi et al., 1995), and NMDARs are required for this

process (Zhang and Wu, 2000; Wu et al., 2002). Our glutamate puff experiments showed that

VIP neurons also express GluN2A/B-containing NMDARs, indicating that at least some VIP

neuron synapses may exhibit AMPAR-dependent LTP (Malenka and Nicoll, 1993; Rebola et al.,

2010; Hunt and Castillo, 2012). It will be important for future studies to determine the

contributions of GluN2D trafficking and GluN2A/B expression to synaptic plasticity in the IC.

GluN2D-containing receptors may alter rate coding for stimuli in the IC

Auditory structures primarily encode changes in sound amplitude envelopes in two

ways: using temporal codes, where neurons phase-lock their firing to the modulation

waveform, and using rate codes, where a neuron's firing rate changes based on the frequency

of the amplitude modulation (AM). While early auditory structures such as the cochlear nucleus

primarily use temporal codes for AM stimuli (Rhode and Greenberg, 1994), neurons in auditory

cortex primarily use rate codes (Yin et al., 2011). Since many IC neurons only phase lock to

lower AM modulation frequencies (Rees and Møller, 1983; Rees and Palmer, 1989; Krishna and

Semple, 2000) but exhibit strong and diverse dependencies of firing rate on changes in AM

modulation frequency (Krishna and Semple, 2000; Nelson and Carney, 2007; Geis and Borst,

2009; Kim et al., 2020), the IC has been proposed as a critical site for the temporal-to-rate code

transition that occurs between brainstem and cortex.

Our model and in vivo data highlight multiple important considerations for this

transition. First, our model VIP neuron (**Figure 7**) and in vivo data (**Figure 9**) show that NMDA

receptors contribute additive gain for rate coding of AM stimuli, providing a boost that pushes

the neuron closer to spike threshold and increases the overall firing rate across modulation

frequencies. This provides evidence for a mechanism that may promote rate coding in IC

neurons by enhancing the dynamic range over which neurons encode AM modulation

frequency using changes in firing rate. Second, when we investigated how adding an NMDA

conductance altered temporal and rate coding in our model neuron, we found that the coding

shifts were highly dependent on the input resistance of the neuron. Cells modeled with a lower

leak conductance (**Figure8A-C**) and thus higher input resistance had clear rate coding for AM

stimuli but exhibited poorer phase-locking. Addition of the NMDA conductance to the higher

input resistance models provided additive gain which increased firing output, similar to what

was seen in our in vivo experiments (**Figure 9C,F**). However, neurons modeled with a higher

leak conductance and thus lower input resistance tended to exhibit stronger phase locking and

poorer rate coding. In addition, adding the NMDA conductance to these models pushed the cell

into depolarization block at higher input frequencies, thus turning the model neuron into a cell

with an onset response profile.

764   Overall, our results show that GluN2C/D-containing NMDA receptors interact with

765   AMPA receptors and passive membrane properties to shape rate and temporal coding in many

766   IC neurons, helping give rise to the diverse ways that IC neurons encode AM stimuli and other

767   temporally modulated sounds.

Figure 1. VIP neurons express GluN2C/D-containing NMDARs

A, Puffs of glutamate onto VIP neurons activated NMDARs. Pharmacology was used to examine whether GluN2C/D-containing receptors contributed to elicited currents. *B,* Brief (10 ms) puffs of glutamate onto a VIP neuron elicited a small inward current (left) which was completely abolished by the GluN2C/D antagonist PPDA (middle) and enhanced by the GluN2C/D positive allosteric modulator CIQ (right). *C,* PPDA decreased the amplitude (LMM: β = -30.46, 95% CI [-56.32, -4.61], *p* = 0.038, *n* = 8) of the response to the glutamate puff. CIQ trended towards enhancing the amplitude (β = 24.33, 95% CI [-1.53, 50.18], *p* = 0.088, *n* = 8) of the response, but did not alter the halfwidth (β = 16.13, 95% CI [-7.89, 37.16], *p* = 0.15, *n* = 8), rise time (β = 5.56, 95% CI [-10.24, 21.36], *p* = 0.49, *n* = 8) or decay τ (β = 13.23, 95% CI [-1.29, 27.74], *p* = 0.10, *n* = 8) of the response to the glutamate puff. Vertical black lines on the paired mean difference plots indicate 95% bootstrap confidence intervals, and gray histograms show the expected distribution of paired mean differences based on bootstrap analysis.

Figure 2. *Glun2c and Glun2d* mRNA is widely expressed throughout the IC, and most VIP

neurons express *Glun2d*

A, 40x confocal images of IC brain slices following RNAscope in situ hybridization showing

expression of *tdTomato* mRNA ***(i),*** which is expressed by VIP neurons in VIP-IRES-Cre x Ai14

mice, *Glun2c* mRNA ***(ii)***, and *Glun2d* mRNA ***(iii)***. 'Cb' indicates portions of the cerebellum that

were captured in the images. ***B,*** 63x confocal images with white arrows indicating *tdTomato-*

positive neurons ***(i)***. 8.1% of *tdTomato* positive neurons expressed *Glun2c* mRNA ***(ii)*** and 91.4%

expressed *Glun2d* mRNA ***(iii)***. *Glun2c* and *Glun2d* mRNA was also commonly observed in

tdTomato-negative neurons, suggesting that these NMDAR subunits are expressed in multiple

IC neuron types. A merge of these images is shown in ***iv***.

Figure 3. 91.4% of VIP neurons express *Glun2d* mRNA

A, Quantification of mRNA expressed in *tdTomato*-positive (VIP) neurons in the IC. 1.0% of VIP

neurons expressed only *Glun2c* mRNA, while 84.3% of VIP neurons expressed only *Glun2d*

mRNA. 7.1% of VIP neurons expressed both *Glun2c* and *Glun2d* mRNA. 7.7% of VIP neurons did

not express *Glun2c* or *Glun2d* mRNA. Individual data points represent the mean for individual

mice. ***B,*** The central nucleus, lateral cortex, and dorsal cortex of the IC did not differ in their

expression profiles of tdTomato-positive neurons expressing only *Glun2c* (0.8%, 0.0%, 3.8%,

respectively), only *Glun2d* (83.7%, 85.0%, 86.5%), both *Glun2c* and *Glun2d* (7.9%, 6.2%, 3.8%),

and neither *Glun2c*/*Glun2d* (7.6%, 8.8%, 5.8%). Individual data points represent the means for

individual mice.

Figure 4. VIP neurons express GluN2A/B-containing NMDARs in addition to GluN2C/D-

containing NMDARs

A, Puffs of glutamate onto VIP neurons activated NMDARs. Pharmacology was used to examine

the proportion of GluN2A/B and GluN2C/D containing receptors present on VIP neurons. ***B,*** An

EPSC elicited by a brief (10 ms) puff of glutamate onto a VIP neuron was abolished by

application of the GluN2C/D antagonist PPDA. ***C,*** PPDA decreased the amplitude (β = -155.49,

95% CI [-220.62, -90.36], *p* = 9.26e-4, *n* = 6) of the response to the glutamate puff. D-AP5

completely abolished the response in all cells tested (amplitude, LMM: β = -194.83, 95% CI [-

259.96, -129.70], *p* = 1.71e-4, *n* = 6). Effects on EPSC kinetics for cells with residual current after

PPDA application are shown in ***C,*** right panels. Vertical lines on the paired mean difference plots

indicate 95% bootstrap confidence intervals, and gray histograms show the expected

distribution of paired mean differences based on bootstrap analysis. ***D,*** Percent of EPSC

amplitude that was sensitive to PPDA application in each of the 6 cells from **C**. The distribution

was created using the 'ggdist' and 'stat_halfeye' functions from the 'ggplot' package for R,

which uses a kernel density estimate to generate a curve that reflects the distribution of the

data (Kay, 2024).

Figure 5. Commissural inputs to VIP neurons activate GluN2C/D-containing NMDARs at resting potential.

A, Viral injections in the right IC drove Chronos expression in commissural projections to the left IC. Optogenetic circuit mapping and pharmacology were used to examine postsynaptic responses in VIP neurons in the left IC. **B,** An EPSC elicited by a brief (10 ms) flash of light (left) was enhanced by the GluN2C/D-specific positive allosteric modulator CIQ (middle) and diminished by the GluN2C/D-specific antagonist PPDA (right). **C,** CIQ did not significantly alter the amplitude (LMM: β = 0.39, 95% CI [4.63, -3.86], *p* = 0.86, *n* = 7), halfwidth (β = 2.95, 95% CI [0.22, 5.69, *p* = 0.064], *p* = 0.42, *n* = 7), rise time (β = 0.56, 95% CI [-0.38, 1.50], *p* = 0.26, *n* = 7), or decay τ (β = 6.13, 95% CI [-2.02, 14.27], *p* = 0.16, *n* = 7) of EPSCs. PPDA abolished the response in all cells tested (amplitude, LMM: β = -7.44, 95% CI [-11.68, -3.20], *p* = 5.13e-3, *n* = 7). Vertical lines on the paired mean difference plots indicate 95% bootstrap confidence intervals, and gray histograms show the expected distribution of paired mean differences based on bootstrap analysis.

Figure 6. GluN2C/D-containing NMDARs facilitate temporal summation in VIP neurons.

A, Viral injections in the right IC drove Chronos expression in commissural projections to the left

IC. Trains of optogenetic stimuli combined with pharmacology were used to examine the

contributions of GluN2C/D-containing receptors to temporal summation. *B,* EPSP summation

elicited by a 30 Hz train of light pulses (left) was decreased by application of the GluN2C/D-

specific antagonist PPDA (right, comparison inset). *C,* Example of a cell in which summation of

optogenetically elicited EPSPs reached action potential threshold, eliciting trains of action

potentials. PPDA eliminated spiking in all but one sweep in this cell. *D,* Example traces from *C*

scaled to demonstrate the differences in temporal summation between control and PPDA

conditions. *E,* The peak amplitude of each EPSP was calculated from the baseline (see inset).

The amplitude of the response increased with increasing peak number (LMM: β = 1.06, 95% CI

[0.66, 1.47], *p* = 9.15e-6, *n* = 5), although the interaction between peak and drug condition was

not statistically significant (β = -0.48, 95% CI [-1.06, 0.095], p = 0.11, n = 5). **F,** PPDA significantly

decreased the area under the curve for the train of EPSPs (LMM: β = -116.57, 95% CI [-198.93, -

34.21], p = 0.038, n = 5), which can be seen in both the non-normalized (left) and normalized

(right) plots. Vertical lines on the paired mean difference plots indicate 95% bootstrap

confidence intervals, and gray histograms show the expected distribution of paired mean

differences based on bootstrap analysis.

Figure 7. GluN2C/D-like NMDA conductance provides additive gain for rate coding in model IC

neuron

A, Model neuron response to a 1 second, 16 Hz input train when the synapse contains an AMPA

conductance (left panel) or both AMPA and NMDA conductances (right panel). To mimic

GluN2C/D-containing NMDA receptors, the NMDA conductance was not voltage dependent. **B,**

Model neuron response to a 1 second, 64 Hz input train when the synapse contains an AMPA

conductance (left panel) or both AMPA and NMDA conductances (right panel). **C,** Rate

modulation transfer functions for modulation frequencies ranging between 16 – 512 Hz at 4

steps/octave. Addition of the NMDA conductance (AMPA + NMDA, orange line) increased the

firing rate at all modulation frequencies tested compared to the AMPA conductance alone

condition (blue line). **D,** Temporal modulation transfer functions in response to the same

modulation frequencies shown in **C**. The model neuron exhibited strong phase locking over the

mid-range of input frequencies in the AMPA only condition (blue line), and poorer phase

locking in the AMPA plus NMDA condition (orange line). Note that the spike in phase locking at

215 Hz for the AMPA plus NMDA condition reflects instance where the sustained firing rate of

the model neuron happened to align well with the input frequency. **E,** Distribution of firing

phases for the AMPA conductance alone (blue dots) and the AMPA plus NMDA conductance

(orange dots) conditions. Each dot represents a single modulation frequency. Concentric circles

on the polar plot indicate modulation frequency.

Figure 8. GluN2C/D-like NMDA conductance interacts with membrane resistance and AMPA

conductance to shift temporal and rate coding for trains of inputs in the neuron model

Model neuron rate and temporal modulation transfer functions for leak conductances of *A,*

0.282, *B,* 0.50, *C,* 0.75, *D,* 1.99, and *E,* 5.30 mS/cm^2. For a spherical neuron with a 15 µm

diameter, these leak conductances correspond to 502, 283, 189, 71, and 27 MΩ membrane

resistances, respectively. Blue lines indicate responses of the model neuron in the AMPA

conductance alone condition, while orange lines indicate responses when the model synapse

contained both AMPA and NMDA conductances.

Figure 9. GluN2C/D-containing NMDARs enhance sound-evoked and spontaneous firing rates in vivo

A, Schematic of in vivo recording setup. Adapted from (Petrucco, 2020). In vivo juxtacellular recordings were performed in the IC of awake, headfixed mice using glass electrodes, and DQP

or vehicle solution was infused using a piggyback pipette connected to a syringe pump. **B,**

Juxtacellular recordings show both rate and temporal coding of the modulation frequency for

AM noise bursts. In this example cell, the firing rate of the neuron increased with increasing

modulation frequency. Second panel, the response of the same cell after infusion of the

GluN2C/D-specific antagonist DQP. **C,** Blocking GluN2C/D-containing NMDA receptors via

infusion of DQP decreased the firing rate of the example neuron from **B** at all modulation

frequencies tested (blue line, left panel). Application of DQP had minimal or no effect on the

vector strength of the response at any modulation frequency (right panel). **D,** Infusion of

vehicle solution into the IC did not alter the firing rate (left panel) or vector strength (right

panel) of the response to AM noise bursts. **E,** Infusion of DQP (left panel) or vehicle (right panel)

did not alter the phases at which firing was phase locked in one DQP example neuron and one

vehicle example neuron (orange is pre-infusion, blue is post-infusion). Each dot represents a

single modulation frequency. Concentric circles on the polar plot indicate modulation

frequency. **F,** DQP significantly decreased the number of spikes elicited during the sound

stimulus (left panel, Z = 56, p = 0.042, median difference = -1.11 Hz, while vehicle infusion did

not change the spike counts (right panel, Z = 21, p = 0.74, median difference = -0.062 Hz). **G,**

DQP infusion (left panel) did not change the average vector strength of the response to AM

stimuli (Z = 41, p = 0.19, median difference = -0.008), and vehicle infusion (right panel) also did

not affect vector strength (Z = 15, p = 0.063, median difference = -0.018). **H,** Infusion of DQP

(left panel) did not change the dynamic range of the firing rates over which IC neurons encoded

AM modulation frequency (Z = 54, p = 0.067, median difference = -0.13 Hz), and this was also

not altered by infusion of vehicle solution (right panel) (Z = 30.5, p = 0.086, median difference =

-0.11 Hz). **I,** DQP decreased the overall spontaneous firing rate of IC neurons between stimulus

presentations (left panel, Z = 58, p = 0.024, median difference = -3.03 Hz), while vehicle infusion

had no effect on spontaneous firing rate (right panel, Z = 8, p = 0.20, median difference = 0.76

Hz).

Table 1. Percentage of *tdTomato+* neurons expressing *Glun2d* and/or *Glun2c* mRNA.

Animal	Slice (n)	No. of tdT+	No. of Glun2d+	No. of Glun2c+	No. of tdT+ Glun2d+ Glun2c+	% tdT+ Glun2d+	% tdT+ Glun2c+	% tdT+ Glun2d+ GluN2c+
Mouse 1	A caudal	42	36	3	3	85.7%	7.1%	7.1%
	B caudal	52	49	4	4	94.2%	7.7%	7.7%
	C caudal	48	47	2	2	97.9%	4.2%	4.2%
	D rostral	3	3	0	0	100.0%	0.0%	0.0%
	E rostral	4	4	0	0	100.0%	0.0%	0.0%
	Total	**149**	**139**	**9**	**9**	**93.3%**	**6.0%**	**6.0%**
Mouse 2	D caudal	71	68	9	9	95.8%	12.7%	12.7%
	B medial	13	10	2	1	76.9%	15.4%	7.7%
	C medial	26	26	4	4	100.0%	15.4%	15.4%
	A rostral	3	3	0	0	100.0%	0.0%	0.0%
	Total	**113**	**107**	**15**	**14**	**94.7%**	**13.3%**	**12.4%**
Mouse 3	B caudal	94	83	10	7	88.3%	10.6%	7.4%
	D caudal	85	76	5	5	89.4%	5.9%	5.9%
	A medial	70	62	2	1	88.6%	2.9%	1.4%
	C rostral	10	9	1	1	90.0%	10.0%	10.0%
	Total	**259**	**230**	**18**	**14**	**88.8%**	**6.9%**	**5.4%**
Grand total		**521**	**476**	**42**	**37**	**91.4%**	**8.1%**	**7.1%**

Table 1.

The number and percentage of *tdTomato* positive cells that expressed *Glun2c* and *Glun2d*

mRNA using an RNAScope in situ hybridization assay. Data is from a series of coronal IC sections

collected from three VIP-IRES-Cre x Ai14 mice aged P45-47.

Table 2. Percentage of neurons that expressed *Glun2d* and *Glun2c* in IC central nucleus, dorsal

cortex, and lateral cortex

Location	Mouse	No. of *tdT+*	No. of *Glun2d+*	No. of *Glun2c+*	No. of *tdT+ Glun2d+ Glun2c+*	*% tdT+ Glun2d+*	*% tdT+ Glun2c+*	*% tdT+ Glun2d+ Glun2c+*
Central nucleus	1	89	84	5	5	94.4%	5.6%	5.6%
	2	68	67	12	12	98.5%	17.6%	17.6%
	3	199	175	14	11	87.9%	7.0%	5.5%
	Total	**356**	**326**	**31**	**28**	**91.6%**	**8.7%**	**7.9%**
Dorsal cortex	1	47	42	4	4	89.4%	8.5%	8.5%
	2	28	25	1	1	89.3%	3.6%	3.6%
	3	38	36	2	2	94.7%	5.3%	5.3%
	Total	**113**	**103**	**7**	**7**	**91.2%**	**6.2%**	**6.2%**
Lateral cortex	1	13	13	0	0	100.0%	0.0%	0.0%
	2	17	15	2	1	88.2%	11.8%	5.9%
	3	22	19	2	1	86.4%	9.1%	4.5%
	Total	**52**	**47**	**4**	**2**	**90.4%**	**7.7%**	**3.8%**
Grand total		**521**	**476**	**42**	**37**	**91.4%**	**8.1%**	**7.1%**

Table 2.

The number and percentage of *tdTomato* positive cells in the three major IC subdivisions

(central nucleus, lateral cortex, and dorsal cortex) that expressed *Glun2c* and *Glun2d* mRNA

using an RNAScope in situ hybridization assay. Data is from a series of coronal IC sections

collected from three VIP-IRES-Cre x Ai14 mice aged P45-47.

References

Adams JC (1979) Ascending projections to the inferior colliculus. J Comp Neurol 183:519–538.

Anair JD, Silveira MA, Mirjalili P, Beebe NL, Schofield BR, Roberts MT (2022) Inhibitory NPY
Neurons Provide a Large and Heterotopic Commissural Projection in the Inferior
Colliculus. Front Neural Circuits 16:871924.

Bates D, Mächler M, Bolker B, Walker S (2015) Fitting Linear Mixed-Effects Models Using lme4. J
Stat Softw 67:1–48.

Beebe NL, Schofield BR (2021) Cholinergic boutons are closely associated with excitatory cells
and four subtypes of inhibitory cells in the inferior colliculus. J Chem Neuroanat
116:101998.

Beebe NL, Silveira MA, Goyer D, Noftz WA, Roberts MT, Schofield BR (2022) Neurotransmitter
phenotype and axonal projection patterns of VIP-expressing neurons in the inferior
colliculus. J Chem Neuroanat 126:102189.

Bernard C (2019) Changing the Way We Report, Interpret, and Discuss Our Results to Rebuild
Trust in Our Research. eNeuro 6:ENEURO.0259-19.2019.

Binshtok AM, Fleidervish IA, Sprengel R, Gutnick MJ (2006) NMDA receptors in layer 4 spiny
stellate cells of the mouse barrel cortex contain the NR2C subunit. J Neurosci 26:708–
715.

Börgers, Christoph (2017) An Introduction to Modeling Neuronal Dynamics. In: An Introduction
to Modeling Neuronal Dynamics, 1st ed., pp 153–192 Texts in Applied Mathematics.
Springer International Publishing.

Brickley SG, Misra C, Mok MHS, Mishina M, Cull-Candy SG (2003) NR2B and NR2D subunits
coassemble in cerebellar Golgi cells to form a distinct NMDA receptor subtype restricted
to extrasynaptic sites. J Neurosci Off J Soc Neurosci 23:4958–4966.

Brothwell SLC, Barber JL, Monaghan DT, Jane DE, Gibb AJ, Jones S (2008) NR2B- and NR2D-
containing synaptic NMDA receptors in developing rat substantia nigra pars compacta
dopaminergic neurones. J Physiol 586:739–750.

Calin-Jageman RJ, Cumming G (2019) Estimation for Better Inference in Neuroscience. eNeuro
6:ENEURO.0205-19.2019.

Choy Buentello D, Bishop DC, Oliver DL (2015) Differential distribution of GABA and glycine
terminals in the inferior colliculus of rat and mouse. J Comp Neurol 523:2683–2697.

D'Angelo E, De Filippi G, Rossi P, Taglietti V (1995) Synaptic excitation of individual rat
cerebellar granule cells in situ: evidence for the role of NMDA receptors. J Physiol 484 (
Pt 2):397–413.

Drotos AC, Roberts MT (2024) Identifying neuron types and circuit mechanisms in the auditory
midbrain. Hear Res 442:108938.

Feng Z-J, Zhang X, Chergui K (2014) Allosteric modulation of NMDA receptors alters
neurotransmission in the striatum of a mouse model of Parkinson's disease. Exp Neurol
255:154–160.

Forman CJ, Tomes H, Mbobo B, Burman RJ, Jacobs M, Baden T, Raimondo JV (2017)
Openspritzer: an open hardware pressure ejection system for reliably delivering picolitre
volumes. Sci Rep 7:2188.

Geis H-R, Borst JGG (2009) Intracellular responses of neurons in the mouse inferior colliculus to
sinusoidal amplitude-modulated tones. J Neurophysiol 101:2002–2016.

Goyer D, Silveira MA, George AP, Beebe NL, Edelbrock RM, Malinski PT, Schofield BR, Roberts
MT (2019) A novel class of inferior colliculus principal neurons labeled in vasoactive
intestinal peptide-Cre mice. eLife 8:32.

Groc L, Heine M, Cousins SL, Stephenson FA, Lounis B, Cognet L, Choquet D (2006) NMDA
receptor surface mobility depends on NR2A-2B subunits. Proc Natl Acad Sci U S A
103:18769–18774.

Harney SC, Jane DE, Anwyl R (2008) Extrasynaptic NR2D-containing NMDARs are recruited to
the synapse during LTP of NMDAR-EPSCs. J Neurosci Off J Soc Neurosci 28:11685–11694.

Hewitt MJ, Meddis R (1994) A computer model of amplitude-modulation sensitivity of single
units in the inferior colliculus. J Acoust Soc Am 95:2145–2159.

Hildebrand ME, Pitcher GM, Harding EK, Li H, Beggs S, Salter MW (2014) GluN2B and GluN2D
NMDARs dominate synaptic responses in the adult spinal cord. Sci Rep 4:4094.

Ho J, Tumkaya T, Aryal S, Choi H, Claridge-Chang A (2019) Moving beyond P values: data
analysis with estimation graphics. Nat Methods 16:565–566.

Hodgkin AL, Huxley AF (1952) A quantitative description of membrane current and its
application to conduction and excitation in nerve. J Physiol 117:500–544.

Hosomi H, Hirai H, Okadaa Y, Amatsu M (1995) Long-term potentiation of neurotransmission in
the inferior colliculus of the rat. Neurosci Lett 195:175–178.

Huang Z, Gibb AJ (2014) Mg2+ block properties of triheteromeric GluN1-GluN2B-GluN2D NMDA
receptors on neonatal rat substantia nigra pars compacta dopaminergic neurones. J
Physiol 592:2059–2078.

Hunt DL, Castillo PE (2012) Synaptic plasticity of NMDA receptors: mechanisms and functional
implications. Curr Opin Neurobiol 22:496–508.

Jing P-B, Chen X-H, Lu H-J, Gao Y-J, Wu X-B (2022) Enhanced function of NR2C/2D-containing
NMDA receptor in the nucleus accumbens contributes to peripheral nerve injury-
induced neuropathic pain and depression in mice. Mol Pain 18:17448069211053256.

Joris PX, Schreiner CE, Rees A (2004) Neural processing of amplitude-modulated sounds. Physiol
Rev 84:541–577.

Kay M (2024) ggdist: Visualizations of Distributions and Uncertainty in the Grammar of
Graphics. IEEE Trans Vis Comput Graph 30:414–424.

Kelly JB, Zhang H (2002) Contribution of AMPA and NMDA receptors to excitatory responses in
the inferior colliculus. Hear Res 168:35–42.

Kim DO, Carney LH, Kuwada S (2020) Amplitude modulation transfer functions reveal opposing
populations within both the inferior colliculus and medial geniculate body. J
Neurophysiol Available at: http://dx.doi.org/10.1152/jn.00279.2020.

Klapoetke NC et al. (2014) Independent optical excitation of distinct neural populations. Nat
Methods 11:338–346.

Krishna BS, Semple MN (2000) Auditory Temporal Processing: Responses to Sinusoidally
Amplitude-Modulated Tones in the Inferior Colliculus. J Neurophysiol 84:255–273.

Kumar SS, Huguenard JR (2003) Pathway-specific differences in subunit composition of synaptic
NMDA receptors on pyramidal neurons in neocortex. J Neurosci Off J Soc Neurosci
23:10074–10083.

Kuznetsova A, Brockhoff PB, Christensen RHB (2017) lmerTest Package: Tests in Linear Mixed
Effects Models. J Stat Softw 82:1–26.

Li Y-H, Li Y, Zheng L, Wang J (2020) Postsynaptic GluN2B-containing NMDA receptors contribute
to long-term depression induction in medial vestibular nucleus neurons of juvenile rats.
Neurosci Lett 715:134674.

Liu J, Shelkar GP, Sarode LP, Gawande DY, Zhao F, Clausen RP, Ugale RR, Dravid SM (2021)
Facilitation of GluN2C-containing NMDA receptors in the external globus pallidus
increases firing of fast spiking neurons and improves motor function in a
hemiparkinsonian mouse model. Neurobiol Dis 150:105254.

Lo F-S, Akkentli F, Tsytsarev V, Erzurumlu RS (2013) Functional significance of cortical NMDA
receptors in somatosensory information processing. J Neurophysiol 110:2627–2636.

Logan SM, Partridge JG, Matta JA, Buonanno A, Vicini S (2007) Long-lasting NMDA receptor-
mediated EPSCs in mouse striatal medium spiny neurons. J Neurophysiol 98:2693–2704.

Lozovaya NA, Grebenyuk SE, Tsintsadze TS, Feng B, Monaghan DT, Krishtal OA (2004)
Extrasynaptic NR2B and NR2D subunits of NMDA receptors shape "superslow"
afterburst EPSC in rat hippocampus. J Physiol 558:451–463.

Ma CL, Kelly JB, Wu SH (2002) AMPA and NMDA receptors mediate synaptic excitation in the
rat's inferior colliculus. Hear Res 168:25–34.

Madisen L, Zwingman TA, Sunkin SM, Oh SW, Zariwala HA, Gu H, Ng LL, Palmiter RD, Hawrylycz
MJ, Jones AR, Lein ES, Zeng H (2010) A robust and high-throughput Cre reporting and
characterization system for the whole mouse brain. Nat Neurosci 13:133–140.

Malenka RC, Nicoll RA (1993) NMDA-receptor-dependent synaptic plasticity: multiple forms and
mechanisms. Trends Neurosci 16:521–527.

Misra C, Brickley SG, Farrant M, Cull-Candy SG (2000) Identification of subunits contributing to
synaptic and extrasynaptic NMDA receptors in Golgi cells of the rat cerebellum. J Physiol
524 Pt 1:147–162.

Momiyama A (2000) Distinct synaptic and extrasynaptic NMDA receptors identified in dorsal
horn neurones of the adult rat spinal cord. J Physiol 523 Pt 3:621–628.

Monyer H, Burnashev N, Laurie DJ, Sakmann B, Seeburg PH (1994) Developmental and regional
expression in the rat brain and functional properties of four NMDA receptors. Neuron
12:529–540.

Monyer H, Sprengel R, Schoepfer R, Herb A, Higuchi M, Lomeli H, Burnashev N, Sakmann B,
Seeburg PH (1992) Heteromeric NMDA receptors: molecular and functional distinction
of subtypes. Science 256:1217–1221.

Mullasseril P, Hansen KB, Vance KM, Ogden KK, Yuan H, Kurtkaya NL, Santangelo R, Orr AG, Le
P, Vellano KM, Liotta DC, Traynelis SF (2010) A subunit-selective potentiator of NR2C-
and NR2D-containing NMDA receptors. Nat Commun 1:90.

Nelson PC, Carney LH (2007) Neural Rate and Timing Cues for Detection and Discrimination of
Amplitude-Modulated Tones in the Awake Rabbit Inferior Colliculus. J Neurophysiol
97:522–539.

Nouhi M, Zhang X, Yao N, Chergui K (2018) CIQ, a positive allosteric modulator of GluN2C/D-
containing N-methyl-d-aspartate receptors, rescues striatal synaptic plasticity deficit in a
mouse model of Parkinson's disease. CNS Neurosci Ther 24:144–153.

Oberle HM, Ford AN, Dileepkumar D, Czarny J, Apostolides PF (2022) Synaptic mechanisms of
top-down control in the non-lemniscal inferior colliculus. eLife 10:e72730.

Oliver DL, Huerta MF (1992) Inferior and Superior Colliculi. In: The Mammalian Auditory
Pathway: Neuroanatomy (Webster DB, Popper AN, Fay RR, eds), pp 168–221. New York,
NY: Springer New York.

Paoletti P, Bellone C, Zhou Q (2013) NMDA receptor subunit diversity: impact on receptor
properties, synaptic plasticity and disease. Nat Rev Neurosci 14:383–400.

Petrucco L (2020) Mouse head schema. Available at: https://zenodo.org/records/3925903
[Accessed April 15, 2024].

Rebola N, Srikumar BN, Mulle C (2010) Activity-dependent synaptic plasticity of NMDA
receptors. J Physiol 588:93–99.

Rees A, Møller AR (1983) Responses of neurons in the inferior colliculus of the rat to AM and
FM tones. Hear Res 10:301–330.

Rees A, Palmer AR (1989) Neuronal responses to amplitude-modulated and pure-tone stimuli in
the guinea pig inferior colliculus, and their modification by broadband noise. J Acoust
Soc Am 85:1978–1994.

Rhode WS, Greenberg S (1994) Encoding of amplitude modulation in the cochlear nucleus of
the cat. J Neurophysiol 71:1797–1825.

Rivera-Perez LM, Kwapiszewski JT, Roberts MT (2021) α3β4 ∗ Nicotinic Acetylcholine Receptors
Strongly Modulate the Excitability of VIP Neurons in the Mouse Inferior Colliculus. Front
Neural Circuits 15:709387.

Shravan Vasishth, Daniel Shad, Audrey Bürki, Reinhold Kliegl (2022) From the t-test to the linear
(mixed) model. In: Linear Mixed Models in Linguistics and Psychology: A Comprehensive
Introduction Available at: https://vasishth.github.io/Freq_CogSci/.

Siegler Retchless B, Gao W, Johnson JW (2012) A single GluN2 subunit residue controls NMDA
receptor channel properties via intersubunit interaction. Nat Neurosci 15:406–413, S1-2.

Silveira MA, Anair JD, Beebe NL, Mirjalili P, Schofield BR, Roberts MT (2020) Neuropeptide Y
Expression Defines a Novel Class of GABAergic Projection Neuron in the Inferior
Colliculus. J Neurosci 40:4685–4699.

Silver RA (2010) Neuronal arithmetic. Nat Rev Neurosci 11:474–489.

Taniguchi H, He M, Wu P, Kim S, Paik R, Sugino K, Kvitsani D, Fu Y, Lu J, Lin Y, Miyoshi G, Shima
Y, Fishell G, Nelson SB, Huang ZJ (2011) A Resource of Cre Driver Lines for Genetic
Targeting of GABAergic Neurons in Cerebral Cortex. Neuron 71:995–1013.

Tovar KR, Westbrook GL (2002) Mobile NMDA receptors at hippocampal synapses. Neuron
34:255–264.

Traynelis SF, Wollmuth LP, McBain CJ, Menniti FS, Vance KM, Ogden KK, Hansen KB, Yuan H,
Myers SJ, Dingledine R (2010) Glutamate Receptor Ion Channels: Structure, Regulation,
and Function Sibley D, ed. Pharmacol Rev 62:405–496.

Vicini S, Wang JF, Li JH, Zhu WJ, Wang YH, Luo JH, Wolfe BB, Grayson DR (1998) Functional and
pharmacological differences between recombinant N-methyl-D-aspartate receptors. J
Neurophysiol 79:555–566.

Wang F, Flanagan J, Su N, Wang L-C, Bui S, Nielson A, Wu X, Vo H-T, Ma X-J, Luo Y (2012)
RNAscope: a novel in situ RNA analysis platform for formalin-fixed, paraffin-embedded
tissues. J Mol Diagn JMD 14:22–29.

Wang XJ, Buzsáki G (1996) Gamma oscillation by synaptic inhibition in a hippocampal
interneuronal network model. J Neurosci Off J Soc Neurosci 16:6402–6413.

Wenzel A, Villa M, Mohler H, Benke D (1996) Developmental and regional expression of NMDA
receptor subtypes containing the NR2D subunit in rat brain. J Neurochem 66:1240–
1248.

Wu SH (2004) Contribution of AMPA, NMDA, and GABAA Receptors to Temporal Pattern of
Postsynaptic Responses in the Inferior Colliculus of the Rat. J Neurosci 24:4625–4634.

Wu SH, Ma CL, Sivaramakrishnan S, Oliver DL (2002) Synaptic modification in neurons of the
central nucleus of the inferior colliculus. Hear Res 168:43–54.

Yi F, Bhattacharya S, Thompson CM, Traynelis SF, Hansen KB (2019) Functional and
pharmacological properties of triheteromeric GluN1/2B/2D NMDA receptors. J Physiol
597:5495–5514.

Yin P, Johnson JS, O'Connor KN, Sutter ML (2011) Coding of Amplitude Modulation in Primary
Auditory Cortex. J Neurophysiol 105:582–600.

Zhang H, Kelly JB (2001) AMPA and NMDA Receptors Regulate Responses of Neurons in the
Rat's Inferior Colliculus. J Neurophysiol 86:871–880.

Zhang X, Feng Z-J, Chergui K (2014) Allosteric modulation of GluN2C/GluN2D-containing NMDA
receptors bidirectionally modulates dopamine release: implication for Parkinson's
disease. Br J Pharmacol 171:3938–3945.

Zhang Y, Wu SH (2000) Long-term potentiation in the inferior colliculus studied in rat brain slice.
Hear Res 147:92–103.

Zheng QY, Johnson KR, Erway LC (1999) Assessment of hearing in 80 inbred strains of mice by
ABR threshold analyses. Hear Res 130:94–107.

1135