BEING A TREE CROP INCREASES THE ODDS OF EXPERIENCING YIELD DECLINES IRRESPECTIVE OF POLLINATOR DEPENDENCE

Marcelo A. Aizen[1,2*,] Gabriela Gleiser[1,3], Thomas Kitzberger[1], and Rubén Milla[4]

[1]Instituto de Investigaciones en Biodiversidad y Medioambiente (INIBIOMA), Universidad Nacional del Comahue-CONICET, Pasaje Gutiérrez 1415, 8400 San Carlos de Bariloche, Río Negro, Argentina.

[2]Wissenschaftskolleg zu Berlin, Wallotstrasse 19, 14193 Berlin, Germany

[3]Centro de Investigaciones sobre Desertificación (CSIC-UV-GV), Ctra Moncada-Náquera km4.5, 46113 Moncada, Valencia, Spain

[4]Departamento de Biología, Geología, Física y Química Inorgánica, Universidad Rey Juan Carlos, Tulipán s/n, 28933 Móstoles, Spain

*Author for correspondence (email: maizen@comahue-conicet.gob.ar)

ORCID ID's

Marcelo A. Aizen < 0000-0001-9079-9749>

Gabriela Gleiser <0000-0002-4178-6703>

Thomas Kitzberger <0000-0002-9754-4121>

Rubén Milla <0000-0001-8912-4373>

ABSTRACT

Crop yields, *i.e.*, harvestable production per unit of cropland area, are in decline for a number of

crops and regions, but the drivers of this process are poorly known. Global decreases in

pollinator abundance and diversity have been proposed as a major driver of yield declines in

crops that depend on animals, mostly bees, to produce fruits and seeds. Alternatively, widespread

tree mortality has been directly and indirectly related to global climate change, which could also

explain yield decreases in tree crops. As tree crops are expected to be more dependent on

pollinators than other crop types, disentangling the relative influence of growth form and

pollinator dependence is relevant to identify the ultimate factors driving yield declines. Yield

decline, defined here as a negative average annual yearly change in yield from 1961 to 2020, was

measured in 4270 time series, involving 136 crops and 163 countries and territories. About

one-fourth of all time series showed declines in crop yield, a characteristic associated with both

high pollinator dependence and a tree growth form. Because pollinator dependence and plant

growth form were partially correlated, we disentangled the effect of each of these two predictors

using a series of generalized linear mixed models that evaluated direct and indirect associations.

Our analyses revealed a stronger association of yield decline with growth form than with

pollinator dependence, a relationship that persisted after partialling out the effect of pollinator

dependence. In particular, yield declines were more common among tree than herbaceous and

shrub crops in all major regions but in Africa, a continent showing a high incidence of yield

declines irrespective of growth form. These results suggest that pollinator decline is not the main

reason behind crop productivity loss, but that other factors such as climate change could be

already affecting crop yield.

Keywords: agriculture, climate change, growth form, pollination crisis, pollinator decline,

pollinator dependence, tree crops, tree mortality, yield decline.

INTRODUCTION

Plant breeding has played a crucial role in improving agricultural productivity through techniques such as hybridization and polyploidization, artificial selection, and genetic engineering. Along with the expansion of agriculture and the intensification in the use of external subsidies (Matson and Vitousek 2006; Aizen et al. 2022), these methods have helped feed a growing human population (Borlaug 1983; Moose and Mumm 2008; Tester and Langridge 2010). However, the sustained increase in crop yields (*i.e.*, a positive change in harvestable production per unit of cropland area) of many crop species in different parts of the world is now decelerating, indicating that the increase in productivity might be reaching a ceiling (Bennett et al. 2012; Ray et al. 2012; Grassini et al. 2013). This may be in part because crop improvements are limited by various plant trade-offs that impose upper boundaries on yield growth (Huot et al. 2014; Fernandez et al. 2021; Garibaldi et al. 2021). Even though resource allocation patterns in crop plants may be driven to extremes by human selection, crops have rarely explored the phenotypic space existing beyond the boundaries set by the evolution of wild plants (Milla et al. 2018; Garibaldi et al. 2021; Cunha et al. 2023).

In addition to intrinsic constraints, environmental degradation associated with global change may cause declines in crop yield (*i.e.*, a negative change in harvestable production per unit of cropland area) over the last decades. In particular, the interaction of different plant traits with a plant's changing abiotic and biotic environment may also contribute to diminishing yield growth, and result in yield decline (Bennett et al. 2012; Ray et al. 2012). One of the most influential factors affecting the yield of many crops is the availability of efficient pollinators, which have declined in many regions during the last decades due to a combination of habitat destruction, land-use change, intensive pesticide use, pathogen transmission, and climate change (Winfree et al. 2009; Potts et al. 2010; Zattara and Aizen 2021). Additionally, climate change has

resulted in extreme temperature fluctuations, droughts, and flooding events (Coumou and

Rahmstorf 2012; Stott 2016; Ummenhofer and Meehl 2017), which have increased plant stress

(Greenwood et al. 2017; Onyekachi et al. 2019), pest susceptibility (Jaime et al. 2019; IPPC

Secretariat 2021), and phenological mismatches (Beard et al. 2019). Two plant-related factors

that can predict crop yield decline in relation to dwindling pollination services and climate

change are pollinator dependence and growth form, respectively. Pollinator dependence can

determine a crop's susceptibility to changes in pollinator availability (Aizen et al. 2022), whereas

a plant's growth form and other correlated functional traits can determine susceptibility to

extreme weather events and plasticity to respond to global warming (Greenwood et al. 2017;

Alecrim et al. 2023).

Dependency on pollinators varies greatly among crops (Aizen et al. 2022). On one

extreme, some crops grown for their vegetative organs (*e.g.*, potato, carrot, tea, etc.) or their

wind-pollinated seeds or fruits (*e.g.*, wheat, maize, olive, etc.) do not rely on pollinators to

produce the parts we consume. On the other extreme, some seed and fruit crops have a high

degree of pollinator dependence (*e.g.*, cacao, watermelon, vanilla, etc.), to the point that their

yield would be reduced close to zero in the absence of pollinators (Klein et al. 2007). However,

more than half of all cultivated crops fall somewhere between these two extremes, which means

that their yield can be improved to different degrees in the presence of pollinators (Aizen et al.

2009, 2022). In any event, the presence of a diverse group of pollinators, sometimes including

rare but highly effective pollinators, is relevant for increasing the yield of most pollinator-

dependent crops, particularly of those with high dependency (Garibaldi et al. 2013; Sáez et al.

2022). For instance, the yield of several crops tends to decrease with increasing distance from the

field edge of the cultivated field in association with a decline in the abundance and diversity of

wild pollinators that thrive in field margins (Garibaldi et al. 2011b). In addition, the yield of

economically important tropical crops such as coffee, which benefit from the pollination

provided by diverse pollinator assemblages, has declined in different countries (Aizen et al.

2020). A recent meta-analysis revealed that relying solely on honey bees (*Apis mellifera*), the

most important managed pollinator globally, is insufficient to significantly reduce the pollination

deficit of most cross-pollinated crops (Sáez et al. 2022). Based on these findings, it is reasonable

to expect that the degree of a crop's dependence on pollinators will influence the yield response

to declines in pollinator populations. Specifically, we predict that as the dependency on

pollinators increases, the occurrence of negative trends in crop yield over the last few decades is

expected to increase as well.

The growth form of plants can determine their susceptibility to global warming. Trees are

expected to show much less adaptive plasticity than herbs to a rapidly changing climate because

of their longer lifespans and slower growth rates, with shrubs characterized by intermediate life-

history attributes. For instance, spring-flowering forest herbs are advancing their phenologies

faster than trees, thus taking advantage of the longer growing season (Alecrim et al. 2023). On

the other hand, trees have deeper root systems and greater leaf biomass than shrubs and herbs,

which allows them to tap water and nutrients from deeper in the soil and capture more sunlight

for photosynthesis. However, these traits could also make trees more vulnerable to changes in

temperature and precipitation because, in addition to their more stringent hydraulic limitations

(Choat et al. 2012), they require more resources to support their growth and metabolism. In

particular, secondary growth in trees and shrubs requires a significant amount of energy and

resources to produce the lignin that makes up the tree's woody tissue (Novaes et al. 2010), which

could increase their susceptibility to stress factors, such as drought and pests (Cailleret et al.

2017). Also, increasing occurrences of wildfires can cause direct damage to the cambium layer

responsible for wood formation (Dickinson et al. 2004), whereas extreme temperatures and

increasing incidence of droughts and frosts can cause xylem embolism and cavitation (Martínez-

Vilalta and Pockman 2002; Savi et al. 2015), all leading to growth abnormalities, partial crown

dieback, and increased mortality (Barigah et al. 2013; Cailleret et al. 2017; Greenwood et al.

2017). On the other hand, herbaceous plants can experience reduced reproduction and high

mortality in relation to extreme temperatures; however, they are probably better adapted to

overcome unsuitable climatic episodes as they can survive as dormant seeds or underground

structures (Keeley et al. 1981; Gardarin and Colbach 2015; Jongen et al. 2015). In an agricultural

context, long-lasting yield declines in tree crops can be triggered by sporadic but increasingly

frequent heat waves, frost events, or pest outbreaks. On the other hand, extreme climatic events

or pest outbreaks do not have long-lasting consequences in herbaceous crops, most of them

annuals, as, unlike long-lived crops, they are sown anew every year and breeding can provide

adaptations within a shorter time frame. Given this background, we expect tree crops to exhibit a

higher incidence of long-term yield declines than herb crops, whereas shrub crops would fall in

between.

The effects of pollinator dependence and growth form on yield decline cannot be studied

independently because these two factors are expected to be associated for two reasons. First,

most crops harvested for their vegetative parts, which are thus pollinator-independent, are herbs

(Klein et al. 2007). Second, among crops cultivated for their fruits and seeds, pollinator

dependence is expected to increase from herbs to trees because the incidence of self-

incompatibility, which implies mandatory cross-pollination for successful fertilization, is higher

in long-lived plants (Ramírez 2022; Cunha and Aizen 2023). Thus, the effect of pollinator

dependence on the probability of yield decline can be easily confounded with the effect of

growth form and vice-versa when both factors are considered separately.

Here we investigated the relationship between pollinator dependence and growth form

across crops and then analyzed whether either pollinator dependence or growth form is

associated with the probability of crop yield decline over the last six decades. We examined the

effects of pollinator dependence and growth form separately and jointly after accounting for any

confounding effect associated with total cultivated area per crop, which might also relate to yield

decline (Aizen et al. 2009). Although our assessment is global, whether a crop's yield declines or

not was estimated at the country scale (the smallest spatial scale available for long-term data),

rather than at the regional or global scale. This relatively small scale was chosen because

declines in yield in some countries may be compensated by increases in others, thus hiding any

effect that spatially heterogeneous either abiotic, biotic, economic or political factors may have

on yield decline. Compilation and analysis of country-level data also allow for the detection of

geographical areas where yield decline may be more prevalent, and to determine whether

potential associations between the probability of yield decline and either pollinator dependence

or growth form vary among regions or can be regarded as a truly global trend.

MATERIALS AND METHODS

The database

We obtained yearly data (1961-2020) on yield and cultivated area at the country level from the

United Nations Food and Agriculture Organization database (FAOSTAT 2021) for a total of 136

crops and crop items (*i.e.*, aggregations of different species or subdivisions in terms of different

harvested parts; see Aizen et al. 2019 for details) for which there is available information on

pollinator dependence (Klein et al. 2007; updated and expanded in Aizen et al. 2022) and there

are uninterrupted yield records since 1961 for at least one country. Data were retrieved for 163

countries and territories that maintained their physical integrity since 1961, resulting in a total of

4280 complete 60-yr time series (crop x country combinations). We discarded 10 series that

showed unrealistic differences in yield (> two orders of magnitude) between any two years.

For classifying crops into pollinator-dependence categories, we considered initially the

five categories established by Klein et al. (2007) based on the expected reduction in crop yield in

the absence of animal pollination: none (0% reduction), little (>0 to <10%), modest ($\geq$10 to

<40%), high ($\geq$40 to <90%) and essential ($\geq$90%). However, given the highly unbalanced

number of crops in each category (see Table S1), particularly when pollinator dependence was

crossed with growth form, we reduced the number of categories to three by maintaining the

category "none" and merging the categories "little" and "modest" into the category "modest" and

the categories "high" and "essential" into the category "high". Among pollinator-independent

crops (category "none"), we further distinguished among those cultivated for their vegetative

parts (*i.e.*, leaves, stems, bark, roots, tubers, etc.) *vs.* reproductive parts (*i.e.*, either fruits or

seeds). By definition, all pollinator-dependent crops (our categories "moderate" and "high")

were cultivated for either their fruits or seeds. Finally, crops were classified into one of three

commonly used growth-form categories (*i.e.*, herbs, shrubs, and trees) using existing databases

(Milla 2020; Gleiser et al. 2021). As a result, the two main focal factors, pollinator dependence

and growth form, had the same number of categories and degrees of freedom, which make them

comparable in terms of inferential testing and associated statistical power (Cottingham et al.

2005). To explore for potential geographical differences in yield decline associated with either

pollinator dependence or growth form, countries were grouped into one of the five following

geographical regions defined by the United Nations Geoscheme

(https://unstats.un.org/unsd/methodology/m49/): Africa, the Americas (including South, Central,

and North America), Asia, Europe, and Oceania (including Australia and New Zealand).

Data analyses

For each time series, we estimated the yield growth rate (*i.e.*, the average annual yearly change in

yield) as the slope of the least-squares linear regression of (log) yield *vs*. year (Fig. S1). This

simple statistical method provides similar point estimates as other regression methods that

consider the autocorrelation nature of time series (Altinay 2003). Given the goal of our study, an

average growth rate <0 over the period 1961-2020 was considered evidence of yield decline,

independent of the absolute value of the growth rate and without imposing any extra arbitrary

criteria or value thresholds. This simple dichotomous classification provided clear evidence of

whether the yield a given crop in a given country was declining or not for the vast majority of

trends (see Results).

In terms of statistical analyses, we first evaluated the association between pollinator

dependence and growth form by means of a chi-square test. Then, we ran a step-up, hypothesis-

driven series of sequential generalized linear mixed-effects models (GLMMs) to disentangle the

influence of pollinator dependence and growth form on yield decline, considering the response

variable as dichotomous based on the sign of the yield growth rate (negative slope=declining

yield, coded as 1; positive slope = non-declining yield, coded as 0). The first model (model

GLMM_0) was a pure random model to characterize variation among crops (independent of

country) and countries (independent of crop) in yield decline, and thus only included crop and

country as random crossed factors. This model was aimed at obtaining "raw" descriptive

estimates of yield decline by crop and country. In addition to the random factors crop and

country, the second and third models included pollinator dependence (GLMM_1a) and growth

form (GLMM_1b), respectively, while the fourth model (GLMM_2) included both factors

together to assess the effect of pollinator dependence independent of growth form, and vice

versa. GLMM_1 and GLMM_2 also included the geographical region as another fixed factor to

account for regional variation in yield decline and, more specifically, to test for differences in

yield decline between regions. A last model (GLMM_3) assessed whether the effect of pollinator

dependence depended on growth form (i.e., the "pollinator dependence x growth form"

interaction), and whether the influence of either of these two crop factors on yield decline

depended on geographical region (i.e., the "pollinator dependence x region" and "growth form x

region" interactions). All the mixed models *sensu stricto* (*i.e.*, GLMM_1a, GLMM_1b,

GLMM_2, and GLMM_3) included the (log10) cumulative total harvested area (in square

kilometers) over the period 1961-2020 for each crop (summed across countries) and for each

country (summed across crops) as covariates to account for variation among crops and among

countries in the probability of yield decline that could relate to the agricultural relevance of the

crop and country, respectively. The log-transformation of these two variables was necessary

because the raw data encompass about six orders of magnitude with an extremely right-skewed

distribution. Using the same random structure and area covariates as in the former models but

considering only data for pollinator-independent crops, we tested for differences in the

probability of yield decline between crops cultivated for their reproductive *vs*. vegetative organs.

All models above considered a binomial (0,1) distribution and a *logit* link function and

were implemented with the statistical software R version 4.0.2 (R_Core_Team 2020), using the

glmmTMB function of the glmmTMB package (Brooks et al. 2017) and rechecked for

consistency using the glmer function of the lme4 package (Bates et al. 2015). Because both

functions produced highly similar results (only differing at the second or third decimal in their

parameter estimates), we only report here results from the models run with glmmTMB. The

extent of multicollinearity among all fixed factors and covariables was assessed using the

adjusted Generalized Variance Inflation Factor (GVIF, adjusted GVIF^[1/(2*df)], where df

stands for degrees of freedom). GVIFs were estimated using the vif function from the car

package (Fox and Weisberg 2019), where adjusted GVIF values <3 indicate no serious

multicollinearity problems. Effects were evaluated statistically using Wald's type II tests because

we were firstly interested in assessing the overall average effects and secondarily the effects of

the interactions. Finally, pairwise means were compared using Tukey's test for multiple

comparisons by running the contrast function of R's lsmeans package (Lenth 2016).

Ignorance of phylogenetic non-independence due to shared evolutionary history among

crops could bias results and inflate type-I errors. However, no need for a phylogenetic correction,

with the consequent loss of statistical power, is required when phylogenetic regression models do

not outperform equivalent non-phylogenetic regression models and there is no evidence of

phylogenetic structure in the residuals of the non-phylogenetic models (Revell 2010). To

evaluate these two conditions, we first constructed a phylogeny for the 136 crops following the

protocol of Gleiser et al. (2021) and Milla (2020). In brief, obtaining the crop phylogeny

involved checking accepted scientific species names based on the Plant List (The_Plant_List

2013) and pruning the GBOTB.extended megaphylogeny (Jin and Qian 2019) according to the

unique scientific names assigned to each of the 136 crops. In the case of crops including a

taxonomically diverse group of species, the whole group was assigned to the most representative

species within the group, provided that the vast majority of groups are composed of a single

species that is agriculturally relevant plus other minor crops (Milla and Osborne 2021). Then, we

compared all the above non-phylogenetic models with homologous generalized linear mixed

linear models with the same fixed and random structures, but that also included a correlation

matrix based on the phylogenetic distances between crops (Revell 2010). These models were run

using the phylo_glmm function written by Li and Bolker (2019) in R language (R_Core_Team

2020). Rather than imposing a given phylogenetic correlation structure on the random effects,

this function models trait evolution following a flexible Brownian motion process that in practice

is implemented as a sequence of independent errors (Li and Bolker 2019). Corresponding models

with and without phylogeny were compared using Akaike Information Criterion (AIC). Also,

residuals of phylogeny-ignorant models were averaged by crop species and these averaged

residuals were analyzed for evidence of a phylogenetic signal using Blomberg's K (Blomberg et

al. 2003) estimated by using the R's package phytools (Revell 2012).

RESULTS

The incidence and distribution of yield decline

We found that 22.58% (n=964) of the 4270 times-series compiled and analyzed were

characterized by negative growth rates in yield over the period 1961-2020. Consistent with the

proposal that a negative growth rate can be taken as evidence of long-term yield decline, the

frequency distribution of the year of maximum yield across the 964 time-series exhibiting

negative growth rates peaked in 1961, the first year of the 1961-2020 time series. About 90% of

all these time series had a reduction of >5%, and 82.5% a reduction of >10% in yield over the

whole period. In contrast, the frequency distribution of the year of maximum yield across the

3306 time-series with positive growth rates peaked in 2020, the last year of the 1961-2020 time

series (Fig. 1).

The expected mean probability of yield decline estimated by the random model

(GLMM_0) was 0.234 (95%CI=[0.215, 0.255]). However, there was high variability in the

incidence of yield decline among crops and countries. The expected probability of yield decline

varied from <0.1 in widespread crops such as maize, wheat, and rape, to >0.4 for assorted crops

such as dates, cherries, walnuts, and cauliflowers (Fig. S2). The number of crops per country

ranged from one in small island states like Nauru and Tuvalu to 76 in Turkey, with a median of

23 crops (Table S2). Variation among countries in the probability of yield decline was even more

diverse than among crops, ranging from <0.08 in countries such as Hungary, Turkey, Myanmar,

Turkey, and China, to >0.50 in countries such as Ecuador, Mauritius, Zimbabwe, and Trinidad

and Tobago (Fig. S3). Particularly, there seems to be a high concentration of countries with a

high probability of yield decline in Central and Southern Africa, Oceania, and to a lesser extent,

the Pacific rim of South America (Fig. 2). Regional differences in the probability of decline were

strong (Table 1), with countries in Africa and Oceania, followed by countries in the Americas,

depicting the highest expected probability of yield decline (Fig. S4).

The importance of pollinator dependence and crop growth form

Pollinator dependence and plant growth form were related traits. Pollinator dependence and

growth form were associated across the whole set of 136 crops (X^2=26.3, df=4, P<0.001).

Specifically, the yield of about 64.55% of all herbaceous crops did not depend on pollinators,

whereas only 12.66% of crops with that growth form depended highly on pollinators. On the

other hand, only 24.39% of all tree crops were pollinator independent, whereas 48.78% were

highly dependent on pollinators (Fig. 3). Shrub crops showed intermediate values, with 37.5 and

18.75% of these crops exhibiting no or high pollinator dependence, respectively. Considering

crops cultivated exclusively for their reproductive organs (105 crops) did not alter this

association (X^2=18.4, df=4, P=0.001), with 18.18, 25.0, and 52.62% of herb, shrub, and tree

crops depending highly on pollinators, respectively.

High pollinator dependence and the tree growth form were both plant characteristics

associated with an enhanced probability of yield decline when considered separately (models

GLMM_1a and GLMM_1b; Fig. 4 and Table 1). Particularly, these two traits increased the

probability of yield decline by ~40 and ~60%, in comparison with pollinator-independent and

herbaceous crops, respectively (Fig. 4). In the absence of the other predictive focal factor,

pollinator dependence and growth form accounted for 5.6 and 12.9% of the among-crop variance

in yield decline, respectively (Fig. 5). However, the effect of pollinator dependence almost

vanished, whereas the effect of plant growth form persisted, when both factors were included in

the same model (GLMM_2; Figs. 4-5 and Table 1). These statistical results were robust to

different data manipulations, such as excluding the modal years 1961 and 2020 (Fig. 1) in the

estimation of yield decline or, as in Deguines et al. (2014), considering pollinator dependence as

a numerical rather than as a categorical variable (Table S3).

We found no evidence of an interaction between pollinator dependence and growth form,

but there was evidence of regional differences in the effect of pollinator dependence on yield

decline (GLMM_3; Table 1). This interaction effect was mostly attributed to Asia where,

contrary to expectation, the estimated probability of yield decline was somewhat lower among

pollinator-dependent crops (Fig. 6). There was also an indication of regional differences in the

effect of growth form on the probability of yield decline (GLMM_3; Table 1). Results show that

a decline in the yield of tree crops was strongest in Europe and Oceania followed by Asia and the

Americas. Africa was the only region where yield decline was not clearly associated with tree

crops (Fig. 7).

Accounting for potentially confounding factors

We found a non-linear strong decrease in the probability of yield decline with increasing total

cultivation area per crop and country (Table 1, Fig. S5). However, the relations between the focal

variables, pollinator dependence and plant growth form, and the probability of decline reported

were independent of these effects, as these area variables were included as fixed-factor effects in

our models. Also, data analyses revealed no evidence for differences in the estimated probability

of yield decline between crops cultivated for their reproductive *vs.* vegetative parts (least-square

means [-1SE/+1SE]=0.18 [-0.023/+0.026] *vs.* 0.14 [-0.021/+0.025]; X^2=1.89, P=0.17). Despite

the association between pollinator dependence and plant growth form, there was no evidence that

multicollinearity could introduce biases in the assessment of any of the factors and covariates

included in our analyses (all adjusted GVIFs <1.1). Last, non-phylogenetic models provided a

better fit to the data than phylogenetically-explicit models based on AIC criteria (ΔAICs> 3),

whereas there was no evidence of phylogenetic structure in the crop-averaged residuals as

revealed by the low magnitude of Blomberg's K signal (Table 1).

DISCUSSION

Crop yield declines are widespread, with approximately one-quarter of all crop × country yield

trends in our dataset exhibiting signs of decline. However, the probability of yield decline was

highly heterogeneous among crops and countries. Part of that heterogeneity was explained by

crops´ dependence on pollinators when this factor was considered in isolation. But this

association mostly disappeared when accounting for the stronger effect of plant growth form.

Therefore, given the association between high pollinator dependence and the tree growth form in

crops, the effect of pollinator dependence on yield declines seems to be a side effect of a more

direct association between the probability of yield decline and plant growth form. In particular,

we reveal that tree crops are more likely to experience yield declines.

The association between growth form and pollinator dependence across crops we report

here (Fig. 3) has relevant conceptual and practical implications. Despite thousands of years of

plant domestication, patterns of covariation between vegetative and reproductive traits in crop

plants still reflect the contrasting life-history strategies that characterize wild plants at large. Both

wild flowering plants and crops show associations between growth form, lifespan, flower size

and numbers, outcrossing, and, as a consequence, the extent of pollinator dependence. Trees, in

particular, are long-lived, produce lots of relatively small flowers, and show high frequencies of

self-incompatibility that enforce outcrossing, which results in high pollinator dependence

(Friedman 2020; Cunha and Aizen 2023; Lanuza et al. 2023). Because somatic mutations may be

passed on to seeds due to a lack of a segregated germline in plants, mutational load accumulation

with increasing perenniality is probably the ultimate driver behind these associations,

(Klekowski 1988; Schoen and Schultz 2019). Thus, selection for complete pollinator

independence in pollinator-dependent tree crops may be unfeasible (Sáez et al. 2020). More

generally, the above patterns of covariation may limit the phenotypic space that can be explored

via either artificial selection or genetic engineering (Milla et al. 2018; Garibaldi et al. 2021;

Cunha et al. 2023).

Pollinators are declining worldwide as a part of the ongoing global biodiversity crisis

(Potts et al. 2010; Zattara and Aizen 2021). This trend threatens not only the yield of hundreds of

crops but also the reproduction of thousands of wild plant species (Rodger et al. 2021). In

particular, diverse pollinator assemblages play a crucial role in maintaining high yields of many

common, nutritionally important, and economically valuable crops, such as coffee, stone-fruit

crops, and cucurbits (Garibaldi et al. 2013). Although dozens of studies have shown that spatial

and temporal local declines in wild pollinator populations negatively affect the yield of many

pollinator-dependent populations, evidence that pollinator decline has impacted crop yields

globally has remained elusive (Aizen et al. 2008, 2022). For instance, global data indicate that

yield growth rates and their stability seem to decrease with increasing pollinator dependence

(Garibaldi et al. 2011a). However, these findings cannot be taken as unequivocal evidence of the

impact of pollinator decline despite widespread pollination limitation (Ashman et al. 2004), as

they may be more strongly and proximately influenced by mate than pollinator availability

(Harder and Aizen 2010; Sáez et al. 2022) or by other correlated factors as shown here. In

addition, previous analyses of global data have failed to find a deceleration in yield growth with

increasing pollinator dependence (Aizen et al. 2022). In our study, we did not find evidence to

support the proposal that a crop's pollinator dependence is a proximal driver of yield decline

after accounting for the association between growth form and pollinator dependence. Asia was

the only region where there was evidence that the probability of yield decline changes with

pollinator dependence irrespective of growth form. However, the observed pattern was contrary

to expectations, suggesting that pollination management, including human hand pollination

(Wurz et al. 2021), might have counteracted potential impacts of pollinator decline in that region.

Therefore, even though evidence implies that pollinator decline is occurring at small as well as

continental spatial scales, this phenomenon does not seem to have affected crop yield globally.

This lack of evidence does not rule out the possibility that pollinator decline may be affecting the

yield of particular crops in some areas, but it implies that pollinator dependence cannot be

considered the primary driver of yield decline in most pollinator-dependent crops. Breeding of

new less pollinator-dependent varieties of typically highly pollinator-dependent crops, such as

almonds (Sáez et al. 2020), and more efficient management of crop pollination (Mueller et al.

2012; Röös et al. 2018) might be offsetting the effect of increasing pollination deficits due to

dwindling pollinator populations.

Beyond logging and habitat destruction, tree mortality rates have increased in forests

around the world over the past few decades, likely as a result of climate change and associated

stressors, such as higher occurrences of insect outbreaks, wildfires, heat waves, and frosts (Allen

et al. 2010; BGCI 2021). Sustained declines in fruit and seed production, the most common

harvest of tree crops, precede tree death (Pesendorfer et al. 2019). Thus, decreasing tree crop

yields over time could signal the impacts of climate change. In addition, interannual yield

variation is higher in woody than in herbaceous crops, independent of pollinator dependence and

harvest organ (Gleiser et al. 2021). Although comparative studies on the susceptibility of

different plant growth forms to climate change are lacking, a recent study has revealed that

herbaceous plants have been more tolerant to frost than woody plants over evolutionary time

(Klimeš et al. 2022). This differential susceptibility could be due to frost-caused xylem

embolism and cavitation, which could trigger wood dieback (Martínez-Vilalta and Pockman

2002; Mayr et al. 2003). Because one of the consequences of climate change is an increasing

incidence of spring-frost damage due to advances in spring phenology (Lamichhane 2021), this

could be one of the contributing factors behind the differential yield decline of tree crops

compared to that of herbaceous crops. Even though frosts can ruin the harvest of an annual

herbaceous crop, re-sowing during the same or the year following the crop's failure would

restore prior yields. Also, the short lifespan and high relevance of several herbaceous plants as

basic staple crops make their genetic manipulation more short-term amenable and profitable than

most long-lived perennial plants (McCown 2000), resulting in the production of new crop

varieties that can adapt rapidly to a changing climate (Henry 2020). Interestingly, the two crops

showing the lowest probability of yield decline were maize and wheat (Fig. S2), two of the most

genetically engineered crops (Takeda and Matsuoka 2008).

Our findings suggest that the association between pollinator dependence and the

probability of yield decline is largely explained by its partial association with growth form.

Specifically, we found a higher frequency of high pollinator dependence among tree crops,

confounding the effects of pollinator dependence and growth form. However, this association

was not as complete as to prevent evaluation of the independent effect of each of these two

factors on the probability of yield decline. In addition to showing that the relationship between

growth form and the probability of yield decline is relatively strong, we found that the increase

in the probability of yield decline among tree crops was consistent across continents, except in

Africa. Africa was the continent with the highest probability of crop yield declines, which might

relate to a combination of events, including an increasing occurrence of dry spells, and political,

social, and economic factors that affect the agricultural sector as a whole (*e.g.*, absence of

significant irrigation infrastructure), irrespective of plant growth form (Ray et al. 2012). The total

cultivated area per country and crop was also a potential confounding factor. The most widely

cultivated crops were both herbaceous and pollinator independent, which showed the lowest

probabilities of yield declines. However, the reported association between plant growth form and

the probability of yield decline accounts for this potential confounding effect. In addition, the

fact that most tree crops are cultivated for either their seeds or fruits may also confound the

effect of the type of organ harvested (vegetative *vs.* reproductive) with the effect of plant growth

form. Yet, we did not find evidence that the type of organ harvested influences the probability of

decline to any significant extent. Finally, we can discard any effects of unmeasured

phylogenetically-conserved factors on the probability of yield decline, given the worse goodness

of fit of the phylogenetically-explicit models and the lack of phylogenetic signal in model

residuals. Therefore, the growth form of a crop plant seems to connect more proximately with

the likelihood of exhibiting a long-term decline in yield than any other of the factors studied.

Although we are interpreting the association between growth form and the probability of

yield decline in the context of climate change, it is essential to explore other explanations for this

relationship that are independent of the environmental context. One possible explanation is that

potentially decreasing market prices may have discouraged the cultivation of tree crops and the

proper management of existing cultivated fields. However, the area cultivated with fruit and seed

crops has been increasing steadily for decades (Aizen et al. 2022), with market prices for those

crops that are several times higher than those of cereals and most other crops cultivated for their

vegetative parts (Gallai et al. 2009). Another potential explanation could be the replacement of

slow-growing tree crops by fast-growing and fast-cash herbaceous crops, which may have left

remaining fields of at least some tree crops unattended, or restricted those crops to less

productive marginal areas. For example, the rapid expansion of soybean cultivation in several

countries of the Americas in the last decades has impacted the diversity of cultivated crops.

However, this replacement cannot be considered a global phenomenon, and in some regions such

as Europe and countries elsewhere, the replacement seems to have been in the opposite direction

(Aizen et al. 2019). In conclusion, the proximate factors explaining the relationship between

growth form and yield declines need to be investigated more in depth. However, climate change

seems to be a plausible overarching phenomenon behind the reported association.

CONCLUDING REMARKS

The deceleration in yield growth is a complex phenomenon with multiple causes, including

limitations in crop improvement and diminishing yield returns to increasing external subsidies

such as irrigation, fertilizers, and pesticides (Ray et al. 2012). However, yield decline, rather than

just yield growth deceleration, is also likely to reflect the consequences of widespread

environmental degradation and not just the reach of human management skills. This is

particularly so when negative growth rates are also related to biological crop traits like growth

form, as reported here. Our study also found that yield decline is widespread but exhibits high

geographic variability. While further research is needed to understand why yield decline is more

severe in some countries and regions, we found that plant growth form, rather than a crop's

pollinator dependence, is a more proximate factor explaining variation in yield decline at

regional scales. This highlights the importance of not considering any single factor in isolation

but contrasting the explanatory power of each factor with other correlated factors in

observational studies. For example, we might have reached a misleading conclusion about the

relationship between pollinator dependence and yield decline if we did not consider that

pollinator dependence is associated with growth form. In particular, our study revealed a

differential incidence of yield decline among tree crops compared to crops with other growth

forms, paralleling reports of widespread mass tree mortality associated with climate change.

While climate change can provide a general explanation for this association, more research is

needed to understand the physiological mechanisms behind it for proper crop management and

breeding. Beyond highlighting the need for this type of follow-up research, the reported

association between plant growth form and yield decline adds to the evidence of the potential

hazards of climate change for food security.

Acknowledgments. The authors acknowledge C.L. Morales, L.D. Harder, and A. Sáez for

helpful discussions, and I. Bartomeus, N. Deguines, and an anonymous reviewer for useful

comments and suggestions on a previous version of this contribution.

Authors contribution. M.A.A, G.G., T.K, and R.M. conceived the study. M.A.A. and G.G.

compiled and analyzed the data. R.M. provided feedback on the analyses. M.A.A. wrote the

paper with input from all authors.

Data and script accessibility. Data and scripts for this publication are available on the Zenodo

Repository (Aizen et al. 2023): https://doi.org/10.5281/zenodo.8206104.

Supplementary material. Tables S1-S3 and Figures S1-S5 are available in the Appendix.

Conflict of interest and disclosure. The authors of this contribution declare that they have no

financial conflict of interest with the content of this article.

Funding. This work was supported by grants from the National Fund for Scientific and

Technological Research of Argentina (FONCYT) [PICT 2015-2333, PICT 2018-2145, the

Ministry of Science and Innovation of Spain (MICINN) [PID2021-122296NB-I00], and the

European Commission (Call HORIZON-CL6-2023-BIODIV-01) [COUSIN - 101135314]. This

work was also supported by a sabbatical fellowship from the Wissenschaftskolleg zu Berlin.

REFERENCES

Aizen, M. A., S. Aguiar, J. C. Biesmeijer, L. A. Garibaldi, D. W. Inouye, C. Jung, D. J. Martins,

R. Medel, C. L. Morales, H. Ngo, A. Pauw, R. J. Paxton, A. Sáez, and C. L. Seymour. 2019.

Global agricultural productivity is threatened by increasing pollinator dependence without a

parallel increase in crop diversification. Global Change Biology 25:3516–3527.

https://doi.org/10.1111/gcb.14736.

Aizen, M. A., M. P. Arbetman, N. P. Chacoff, V. R. Chalcoff, P. Feinsinger, L. A. Garibaldi, L.

D. Harder, C. L. Morales, A. Sáez, and A. J. Vanbergen. 2020. Invasive bees and their

impact on agriculture. Advances in Ecological Research 63:49–92.

https://doi.org/10.1016/bs.aecr.2020.08.001.

Aizen, M. A., L. A. Garibaldi, S. A. Cunningham, and A. M. Klein. 2008. Long-term global

trends in crop yield and production reveal no current pollination shortage but increasing

pollinator dependency. Current Biology 18:1572–5.

https://doi.org/10.1016/j.cub.2008.08.066.

Aizen, M. A., L. A. Garibaldi, S. A. Cunningham, and A. M. Klein. 2009. How much does

agriculture depend on pollinators? Lessons from long-term trends in crop production.

Annals of Botany 103:1579–1588. https://doi.org/10.1093/aob/mcp076.

Aizen, M. A., L. A. Garibaldi, and L. D. Harder. 2022. Myth and reality of a global crisis for

agricultural pollination. Ecología Austral 32:698–715.

https://doi.org/10.25260/ea.22.32.2.1.1875.

Alecrim, E. F., R. D. Sargent, and J. R. K. Forrest. 2023. Higher-latitude spring-flowering herbs

advance their phenology more than trees with warming temperatures. Journal of Ecology

111:156–169. https://doi.org/10.1111/1365-2745.14023.

Allen, C. D., A. K. Macalady, H. Chenchouni, D. Bachelet, N. McDowell, M. Vennetier, T.

Kitzberger, A. Rigling, D. D. Breshears, E. H. (Ted. Hogg, P. Gonzalez, R. Fensham, Z.

Zhang, J. Castro, N. Demidova, J. H. Lim, G. Allard, S. W. Running, A. Semerci, and N.

Cobb. 2010. A global overview of drought and heat-induced tree mortality reveals emerging

climate change risks for forests. Forest Ecology and Management 259:660–684.

https://doi.org/10.1016/j.foreco.2009.09.001.

Altinay, G. 2003. Estimating growth rate in the presence of serially correlated errors. Applied

Economics Letters 10:967–970. https://doi.org/10.1080/1350485032000165485.

Ashman, T.-L., T. M. Knight, J. A. Steets, P. Amarasekare, M. Burd, D. R. Campbell, M. R.

Dudash, M. O. Johnston, S. J. Mazer, R. J. Mitchell, M. T. Morgan, and W. G. Wilson.

2004. Pollen limitation of plant reproduction: Ecological and evolutionary causes and

consequences. Ecology 85:2408–2421. https://doi.org/10.1890/03-8024.

Barigah, T. S., O. Charrie, M. Douris, M. Bonhomme, S. Herbette, T. Améglio, R. Fichot, F.

Brignolas, and H. Cochard. 2013. Water stress-induced xylem hydraulic failure is a causal

factor of tree mortality in beech and poplar. Annals of Botany 112:1431–1437.

https://doi.org/10.1093/aob/mct204.

Bates, D., M. Mächler, B. M. Bolker, and S. C. Walker. 2015. Fitting linear mixed-effects

models using lme4. Journal of Statistical Software 67:1–48.

https://doi.org/10.18637/jss.v067.i01.

Beard, K. H., K. C. Kelsey, A. J. Leffler, and J. M. Welker. 2019. The missing angle: Ecosystem

consequences of phenological mismatch. Trends in Ecology and Evolution 34:885–888.

https://doi.org/10.1016/j.tree.2019.07.019.

Bennett, A. J., G. D. Bending, D. Chandler, S. Hilton, and P. Mills. 2012. Meeting the demand

for crop production: The challenge of yield decline in crops grown in short rotations.

Biological Reviews 87:52–71. https://doi.org/10.1111/j.1469-185X.2011.00184.x.

BGCI. 2021. State of the World`s Trees. Botanic Gardens Conservation International, Richmond,

UK.

Blomberg, S. P., T. Garland, and A. R. Ives. 2003. Testing for phylogenetic signal in

comparative data: Behavioral traits are more labile. Evolution 57:717–745.

https://doi.org/10.1111/j.0014-3820.2003.tb00285.x.

Borlaug, N. E. 1983. Contributions of conventional plant breeding to food production. Science

219:689–693. https://doi.org/10.1126/science.219.4585.689.

Brooks, M. E., K. Kristensen, K. J. van Benthem, A. Magnusson, C. W. Berg, A. Nielsen, H. J.

Skaug, M. Mächler, and B. M. Bolker. 2017. glmmTMB balances speed and flexibility

among packages for zero-inflated generalized linear mixed modeling. R Journal 9:378–400.

https://doi.org/10.32614/rj-2017-066.

Cailleret, M., S. Jansen, E. M. R. Robert, L. Desoto, T. Aakala, J. A. Antos, B. Beikircher, C.

Bigler, H. Bugmann, M. Caccianiga, V. Čada, J. J. Camarero, P. Cherubini, H. Cochard, M.

R. Coyea, K. Čufar, A. J. Das, H. Davi, S. Delzon, M. Dorman, G. Gea-Izquierdo, S.

Gillner, L. J. Haavik, H. Hartmann, A. M. Hereş, K. R. Hultine, P. Janda, J. M. Kane, V. I.

Kharuk, T. Kitzberger, T. Klein, K. Kramer, F. Lens, T. Levanic, J. C. Linares Calderon, F.

Lloret, R. Lobo-Do-Vale, F. Lombardi, R. López Rodríguez, H. Mäkinen, S. Mayr, I.

Mészáros, J. M. Metsaranta, F. Minunno, W. Oberhuber, A. Papadopoulos, M. Peltoniemi,

A. M. Petritan, B. Rohner, G. Sangüesa-Barreda, D. Sarris, J. M. Smith, A. B. Stan, F.

Sterck, D. B. Stojanović, M. L. Suarez, M. Svoboda, R. Tognetti, J. M. Torres-Ruiz, V.

Trotsiuk, R. Villalba, F. Vodde, A. R. Westwood, P. H. Wyckoff, N. Zafirov, and J.

Martínez-Vilalta. 2017. A synthesis of radial growth patterns preceding tree mortality.

Global Change Biology 23:1675–1690. https://doi.org/10.1111/gcb.13535.

Choat, B., S. Jansen, T. J. Brodribb, H. Cochard, S. Delzon, R. Bhaskar, S. J. Bucci, T. S. Feild,

S. M. Gleason, U. G. Hacke, A. L. Jacobsen, F. Lens, H. Maherali, J. Martínez-Vilalta, S.

Mayr, M. Mencuccini, P. J. Mitchell, A. Nardini, J. Pittermann, R. B. Pratt, J. S. Sperry, M.

Westoby, I. J. Wright, and A. E. Zanne. 2012. Global convergence in the vulnerability of

forests to drought. Nature 491:752–755. https://doi.org/10.1038/nature11688.

Cottingham, K. L., J. T. Lennon, and B. L. Brown. 2005. Knowing when to draw the line:

Designing more informative ecological experiments. Frontiers in Ecology and the

Environment 3:145–152. https://doi.org/10.1890/1540-

9295(2005)003[0145:KWTDTL]2.0.CO;2.

Coumou, D., and S. Rahmstorf. 2012. A decade of weather extremes. Nature Climate Change

2:491–496. https://doi.org/10.1038/nclimate1452.

Cunha, N. L., and M. A. Aizen. 2023. Pollen production per flower increases with floral display

size across animal-pollinated flowering plants. American Journal of Botany 110:e16180.

https://doi.org/10.1002/ajb2.16180.

Cunha, N. L. da, N. P. Chacoff, A. Sáez, R. Schmucki, L. Galetto, M. Devoto, J. Carrasco, M. P.

Mazzei, S. E. Castillo, T. P. Palacios, J. L. Vesprini, K. Agostini, A. M. Saraiva, B. A.

Woodcock, J. Ollerton, and M. A. Aizen. 2023. Soybean dependence on biotic pollination

decreases with latitude. Agriculture, Ecosystems & Environment 347:108376.

https://doi.org/https://doi.org/10.1016/j.agee.2023.108376.

Deguines, N., C. Jono, M. Baude, M. Henry, R. Julliard, and C. Fontaine. 2014. Large-scale

trade-off between agricultural intensification and crop pollination services. Frontiers in

Ecology and the Environment 12:212–217. https://doi.org/10.1890/130054.

Dickinson, M. B., J. Jolliff, and A. S. Bova. 2004. Vascular cambium necrosis in forest fires:

Using hyperbolic temperature regimes to estimate parameters of a tissue-response model.

Australian Journal of Botany 52:757–763. https://doi.org/10.1071/BT03111.

FAOSTAT. 2021. Data available at http://www.fao.org/faostat/en. Last accessed December-27-

2021.

Fernandez, A. R., A. Sáez, C. Quintero, G. Gleiser, and M. A. Aizen. 2021. Intentional and

unintentional selection during plant domestication: Herbivore damage, plant defensive traits

and nutritional quality of fruit and seed crops. New Phytologist 231:1586–1598.

https://doi.org/10.1111/nph.17452.

Fox, J., and S. Weisberg. 2019. CAR - An R Companion to Applied Regression. Thousand Oaks

CA: Sage.

Friedman, J. 2020. The evolution of annual and perennial plant life histories: Ecological

correlates and genetic mechanisms. Annual Review of Ecology, Evolution, and Systematics

51:461–481. https://doi.org/10.1146/annurev-ecolsys-110218-024638.

Gallai, N., J.-M. M. Salles, J. Settele, and B. E. Vaissière. 2009. Economic valuation of the

vulnerability of world agriculture confronted with pollinator decline. Ecological Economics

68:810–821. https://doi.org/10.1016/j.ecolecon.2008.06.014.

Gardarin, A., and N. Colbach. 2015. How much of seed dormancy in weeds can be related to

seed traits? Weed Research 55:14–25. https://doi.org/10.1111/wre.12121.

Garibaldi, L. A., M. A. Aizen, A. M. Klein, S. A. Cunningham, and L. D. Harder. 2011a. Global

growth and stability of agricultural yield decrease with pollinator dependence. Proceedings

of the National Academy of Sciences 108:5909–5914.

https://doi.org/10.1073/pnas.1012431108.

Garibaldi, L. A., M. A. Aizen, A. Sáez, G. Gleiser, M. M. Strelin, and L. D. Harder. 2021. The

influences of progenitor filtering, domestication selection and the boundaries of nature on

the domestication of grain crops. Functional Ecology 35:1998–2011.

https://doi.org/10.1111/1365-2435.13819.

Garibaldi, L. A., I. Steffan-Dewenter, C. Kremen, J. M. Morales, R. Bommarco, S. A.

Cunningham, L. G. Carvalheiro, N. P. Chacoff, J. H. Dudenhöffer, S. S. Greenleaf, A.

Holzschuh, R. Isaacs, K. Krewenka, Y. Mandelik, M. M. Mayfield, L. A. Morandin, S. G.

Potts, T. H. Ricketts, H. Szentgyörgyi, B. F. Viana, C. Westphal, R. Winfree, and A. M.

Klein. 2011b. Stability of pollination services decreases with isolation from natural areas

despite honey bee visits. Ecology letters 14:1062–1072. https://doi.org/10.1111/j.1461-

0248.2011.01669.x.

Garibaldi, L. A., I. Steffan-Dewenter, R. Winfree, M. A. Aizen, R. Bommarco, S. A.

Cunningham, C. Kremen, L. G. Carvalheiro, L. D. Harder, O. Afik, I. Bartomeus, F.

Benjamin, V. Boreux, D. Cariveau, N. P. Chacoff, J. H. Dudenhöffer, B. M. Freitas, J.

Ghazoul, S. Greenleaf, J. Hipólito, A. Holzschuh, B. Howlett, R. Isaacs, S. K. Javorek, C.

M. Kennedy, K. M. Krewenka, S. Krishnan, Y. Mandelik, M. M. Mayfield, I. Motzke, T.

Munyuli, B. A. Nault, M. Otieno, J. Petersen, G. Pisanty, S. G. Potts, R. Rader, T. H.

Ricketts, M. Rundlöf, C. L. Seymour, C. Schüepp, H. Szentgyörgyi, H. Taki, T. Tscharntke,

C. H. Vergara, B. F. Viana, T. C. Wanger, C. Westphal, N. Williams, and A. M. Klein.

2013. Wild pollinators enhance fruit set of crops regardless of honey bee abundance.

Science 339:1608–1611. https://doi.org/10.1126/science.1230200.

Gleiser, G., N. L. Da Cunha, A. Sáez, and M. A. Aizen. 2021. Ecological correlates of crop yield

growth and interannual yield variation at a global scale. Web Ecology 21:15–43.

https://doi.org/10.5194/we-21-15-2021.

Grassini, P., K. M. Eskridge, and K. G. Cassman. 2013. Distinguishing between yield advances

and yield plateaus in historical crop production trends. Nature Communications 4:2918.

https://doi.org/10.1038/ncomms3918.

Greenwood, S., P. Ruiz-Benito, J. Martínez-Vilalta, F. Lloret, T. Kitzberger, C. D. Allen, R.

Fensham, D. C. Laughlin, J. Kattge, G. Bönisch, N. J. B. Kraft, and A. S. Jump. 2017. Tree

mortality across biomes is promoted by drought intensity, lower wood density and higher

specific leaf area. Ecology Letters 20:539–553. https://doi.org/10.1111/ele.12748.

Harder, L. D., and M. A. Aizen. 2010. Floral adaptation and diversification under pollen

limitation. Philosophical transactions of the Royal Society of London. Series B, Biological

Sciences 365:529–543. https://doi.org/10.1098/rstb.2009.0226.

Henry, R. J. 2020. Innovations in plant genetics adapting agriculture to climate change. Current

Opinion in Plant Biology 56:168–173. https://doi.org/10.1016/j.pbi.2019.11.004.

Huot, B., J. Yao, B. L. Montgomery, and S. Y. He. 2014. Growth-defense tradeoffs in plants: A

balancing act to optimize fitness. Molecular Plant 7:1267–1287.

https://doi.org/10.1093/mp/ssu049.

IPPC Secretariat. 2021. Scientific Review of the Impact of Climate Change on Plant Pests. FAO

on behalf of the IPPC Secretariat. Rome, Italy. Retrieved from

https://policycommons.net/artifacts/2194918/scientific-review-of-the-impact-of-climate-

change-on-plant-pests/2950895/ on 14 Mar 2023. CID: 20.500.12592/snqmv2.

https://doi.org/10.4060/cb4769en.

Jaime, L., E. Batllori, J. Margalef-Marrase, M. Á. Pérez Navarro, and F. Lloret. 2019. Scots pine

(*Pinus sylvestris* L.) mortality is explained by the climatic suitability of both host tree and

bark beetle populations. Forest Ecology and Management 448:119–129.

https://doi.org/10.1016/j.foreco.2019.05.070.

Jin, Y., and H. Qian. 2019. V.PhyloMaker: An R package that can generate very large

phylogenies for vascular plants. Ecography 42:1–7. https://doi.org/10.1111/ecog.04434.

Jongen, M., C. Hellmann, and S. Unger. 2015. Species-specific adaptations explain resilience of

herbaceous understorey to increased precipitation variability in a Mediterranean oak

woodland. Ecology and Evolution 5:4246–4262. https://doi.org/10.1002/ece3.1662.

Keeley, S. C., J. E. Keeley, S. M. Hutchinson, and A. W. Johnson. 1981. Postfire succession of

the herbaceous flora in southern California chaparral. Ecology 62:1608–1621.

https://doi.org/10.2307/1941516.

Klein, A.-M., B. E. Vaissière, J. H. Cane, I. Steffan-Dewenter, S. A. Cunningham, C. Kremen,

and T. Tscharntke. 2007. Importance of pollinators in changing landscapes for world crops.

Proceedings of the Royal Society B: Biological Sciences 274:303–313.

https://doi.org/10.1098/rspb.2006.3721.

Klekowski, E. J. 1988. Genetic load and its causes in long-lived plants. Trees 2:195–203.

https://doi.org/10.1007/BF00202374.

Klimeš, A., I. Šímová, A. Zizka, A. Antonelli, and T. Herben. 2022. The ecological drivers of

growth form evolution in flowering plants. Journal of Ecology 110:1525–1536.

https://doi.org/10.1111/1365-2745.13888.

Lamichhane, J. R. 2021. Rising risks of late-spring frosts in a changing climate. Nature Climate

Change 11:554–555. https://doi.org/10.1038/s41558-021-01090-x.

Lanuza, J. B., R. Rader, J. Stavert, L. K. Kendall, M. E. Saunders, and I. Bartomeus. 2023.

Covariation among reproductive traits in flowering plants shapes their interactions with

pollinators. Functional Ecology 7:2072–2084. https://doi.org/10.1111/1365-2435.14340.

Lenth, R. V. 2016. Least-Squares Means: The R Package lsmeans. Journal of Statistical Software

69:1–33. https://doi.org/10.18637/jss.v069.i01.

Li, M., and B. Bolker. 2019. wzmli/phyloglmm: First release of phylogenetic comparative

analysis in lme4- verse. Zenodo. https://doi.org/10.5281/zenodo.2639887.

Martínez-Vilalta, J., and W. T. Pockman. 2002. The vulnerability to freezing-induced xylem

cavitation of *Larrea tridentata* (Zygophyllaceae) in the Chihuahuan desert. American

Journal of Botany 89:1916–1924. https://doi.org/10.3732/ajb.89.12.1916.

Matson, P. A., and P. M. Vitousek. 2006. Agricultural intensification: Will land spared from

farming be land spared for nature? Conservation Biology 20:709–710.

https://doi.org/10.1111/j.1523-1739.2006.00442.x.

Mayr, S., A. Gruber, and H. Bauer. 2003. Repeated freeze–thaw cycles induce embolism in

drought stressed conifers (Norway spruce, stone pine). Planta 217:436–441.

McCown, B. H. 2000. Recalcitrance of woody and herbaceous perennial plants: Dealing with

genetic predeterminism. In Vitro Cellular & Developmental Biology - Plant 36:149–154.

Milla, R. 2020. Crop Origins and Phylo Food: A database and a phylogenetic tree to stimulate

comparative analyses on the origins of food crops. Global Ecology and Biogeography

29:606–614. https://doi.org/10.1111/geb.13057.

Milla, R., J. M. Bastida, M. M. Turcotte, G. Jones, C. Violle, C. P. Osborne, J. Chacón-Labella,

Ê. E. Sosinski, J. Kattge, D. C. Laughlin, E. Forey, V. Minden, J. H. C. Cornelissen, B.

Amiaud, K. Kramer, G. Boenisch, T. He, V. D. Pillar, and C. Byun. 2018. Phylogenetic

patterns and phenotypic profiles of the species of plants and mammals farmed for food.

Nature Ecology and Evolution 2:1808–1817. https://doi.org/10.1038/s41559-018-0690-4.

Milla, R., and C. P. Osborne. 2021. Crop origins explain variation in global agricultural

relevance. Nature Plants 7:598–607. https://doi.org/10.1038/s41477-021-00905-1.

Moose, S. P., and R. H. Mumm. 2008. Molecular plant breeding as the foundation for 21st

century crop improvement. Plant Physiology 147:969–977.

https://doi.org/10.1104/pp.108.118232.

Mueller, N. D., J. S. Gerber, M. Johnston, D. K. Ray, N. Ramankutty, and J. A. Foley. 2012.

Closing yield gaps through nutrient and water management. Nature 490:254–257.

https://doi.org/10.1038/nature11420.

Novaes, E., M. Kirst, V. Chiang, H. Winter-Sederoff, and R. Sederoff. 2010. Lignin and

biomass: A negative correlation for wood formation and lignin content in trees. Plant

Physiology 154:555–561. https://doi.org/10.1104/pp.110.161281.

Onyekachi, O. G., O. O. Boniface, N. F. Gemlack, and N. Nicholas. 2019. The effect of climate

change on abiotic plant stress: A review. Pages 71–83 *in* A. Oliveira, editor. Abiotic and

Biotic Stress in Plants. IntechOpen. https://doi.org/10.5772/intechopen.82681.

Pesendorfer, M. B., M. Bogdziewicz, W. D. Koenig, M. Ledwoń, and M. Żywiec. 2019.

Declining fruit production before death in a widely distributed tree species, *Sorbus*

aucuparia L. Annals of Forest Science 76:11. https://doi.org/10.1007/s13595-018-0791-x.

Potts, S. G., J. C. Biesmeijer, C. Kremen, P. Neumann, O. Schweiger, and W. E. Kunin. 2010.

Global pollinator declines: Trends, impacts and drivers. Trends in Ecology and Evolution

25:345–353. https://doi.org/10.1016/j.tree.2010.01.007.

R_Core_Team. 2020. R: A language and environment for statistical computing. Available at

http://www.R-project.org/. R foundation for statistical computing, Vienna, Austria.

Ramírez, N. 2022. Breeding systems of four plant communities in the Venezuelan central plains.

Plant Systematics and Evolution 308:17. https://doi.org/10.1007/s00606-021-01794-9.

Ray, D. K., N. Ramankutty, N. D. Mueller, P. C. West, and J. A. Foley. 2012. Recent patterns of

crop yield growth and stagnation. Nature Communications 3:1293.

https://doi.org/10.1038/ncomms2296.

Revell, L. J. 2010. Phylogenetic signal and linear regression on species data. Methods in Ecology

and Evolution 1:319–329. https://doi.org/10.1111/j.2041-210X.2010.00044.x.

Revell, L. J. 2012. phytools: An R package for phylogenetic comparative biology (and other

things). Methods in Ecology and Evolution 3:217–223. https://doi.org/10.1111/j.2041-

210X.2011.00169.x.

Rodger, J. G., J. M. Bennett, M. Razanajatovo, T. M. Knight, M. van Kleunen, T. L. Ashman, J.

A. Steets, C. Hui, G. Arceo-Gómez, M. Burd, L. A. Burkle, J. H. Burns, W. Durka, L.

Freitas, J. E. Kemp, J. Li, A. Pauw, J. C. Vamosi, M. Wolowski, J. Xia, and A. G. Ellis.

2021. Widespread vulnerability of flowering plant seed production to pollinator declines.

Science Advances 7:eabd3524. https://doi.org/10.1126/sciadv.abd3524.

Röös, E., A. Mie, M. Wivstad, E. Salomon, B. Johansson, S. Gunnarsson, A. Wallenbeck, R.

Hoffmann, U. Nilsson, C. Sundberg, and C. A. Watson. 2018. Risks and opportunities of

increasing yields in organic farming. A review. Agronomy for Sustainable Development

38:14. https://doi.org/10.1007/s13593-018-0489-3.

Sáez, A., R. Aguilar, L. Ashworth, G. Gleiser, C. L. Morales, A. Traveset, and M. A. Aizen.

2022. Managed honeybees decrease pollination limitation in self-compatible but not in self-

incompatible crops. Proceedings of the Royal Society B: Biological Sciences

289:20220086. https://doi.org/10.1098/rspb.2022.0086.

Sáez, A., M. A. Aizen, S. Medici, M. Viel, E. Villalobos, and P. Negri. 2020. Bees increase crop

yield in an alleged pollinator-independent almond variety. Scientific Reports 10:3177.

https://doi.org/10.1038/s41598-020-59995-0.

Savi, T., S. Bertuzzi, S. Branca, M. Tretiach, and A. Nardini. 2015. Drought-induced xylem

cavitation and hydraulic deterioration: Risk factors for urban trees under climate change?

New Phytologist 205:1106–1116. https://doi.org/10.1111/nph.13112.

Schoen, D. J., and S. T. Schultz. 2019. Somatic mutation and evolution in plants. Annual Review

of Ecology, Evolution, and Systematics 50:49–73. https://doi.org/10.1146/annurev-ecolsys-

110218-024955.

Stott, P. 2016. How climate change affects extreme weather events: Research can increasingly

755 determine the contribution of climate change to extreme events such as droughts. Science

756 352:1517–1518. https://doi.org/10.1126/science.aaf7271.

757 Takeda, S., and M. Matsuoka. 2008. Genetic approaches to crop improvement: Responding to

758 environmental and population changes. Nature Reviews Genetics 9:444–457.

759 https://doi.org/10.1038/nrg2342.

760 Tester, M., and P. Langridge. 2010. Breeding technologies to increase crop production in a

761 changing world. Science 327:818–822. https://doi.org/10.1126/science.1183700.

762 The_Plant_List. 2013. Version 2. Available at http://www.theplantlist.org.

763 Ummenhofer, C. C., and G. A. Meehl. 2017. Extreme weather and climate events with ecological

764 relevance: A review. https://doi.org/10.1098/rstb.2016.0135.

765 Winfree, R., R. Aguilar, D. P. Vazquez, G. LeBuhn, and M. A. Aizen. 2009. A meta-analysis of

766 bees´responses to anthropogenic disturbance. Ecology 90:2068–2076.

767 https://doi.org/10.1890/08-1245.1.

768 Wurz, A., I. Grass, and T. Tscharntke. 2021, August 14. Hand pollination of global crops – A

769 systematic review. Urban & Fischer. https://doi.org/10.1016/j.baae.2021.08.008.

770 Zattara, E. E., and M. A. Aizen. 2021. Worldwide occurrence records suggest a global decline in

771 bee species richness. One Earth 4:114–123. https://doi.org/10.1016/j.oneear.2020.12.005.

Fig. 1. Relative frequency of year of maximum yield for 4270 crop x country yield trends (1961-2020) categorized as trends showing positive or negative average yield growths. The dashed lines indicate the median year of maximum yield for each yield-growth category.

Fig. 2. World map of yield decline. The map depicts the probability of yield decline (*i.e.*, the estimated proportion of crops that showed negative average growth rates in each country from 1961-2020) according to model GLMM_0 (see also Fig. S3).

Fig. 3. Relative frequency of crops with none, modest, and high dependence on pollinators across growth-form categories (*i.e.*, herb, shrub, and tree crops). In parentheses, the number of crops in each growth-form category.

Fig. 4. Probability of yield decline in relation to pollinator dependence and growth form. The figure depicts the mean estimates (+/- 1 SE) of the proportion of crops in each pollinator dependence and growth-form category with negative average growth rates during the period 1961-2020. The left two panels (GLMM_1a and GLMM_1b) show the results of the effect of each factor when the other factor was not included in the model (*i.e.*, model GLMM_1a tested the effect of pollinator dependence and model GLMM_1b tested the effect of growth form), whereas the right two panels the results of each factor after accounting for the confounding effect of the other factor (model GLMM_2). Means with the same letter do not provide evidence of statistical differences at the level of α = 0.05 according to a pairwise Tukey's a posteriori test. Wald's type II test results are provided (see also Table 1).

Fig. 5. Percentage of among-crop variance in yield decline explained by pollinator dependence and growth form in the absence of the other focal factor as estimated from models GLMM_1a and GLMM_1b, respectively, and the independent and shared percentage of the among-crop variance explained by these two focal factors as estimated from model GLMM_2. All these components of variance exclude any variation that could also be accounted by differences among crops in cultivated area (Table 1).

Fig. 6. Probability of yield decline in relation to pollinator dependence by geographical region. The figure depicts the mean estimates (+/- 1 SE) of the proportion of crops in each pollinator-dependence category with negative average growth rates in each region during the period 1961-2020 according to model GLMM_3. Means with the same letter in each panel do not provide evidence of statistical differences at the level of α = 0.05 according to a pairwise Tukey's a posteriori test.

Fig. 7. Probability of yield decline in relation to crop growth form by geographical region. The figure depicts the mean estimates (+/- 1 SE) of the proportion of crops in each growth-form category with negative average growth rates in each region during the period 1961-2020 according to model GLMM_3. Means with the same letter in each panel do not provide evidence of statistical differences at the level of α = 0.05 according to a pairwise Tukey's a posteriori test.

Table 1. Results of logistic mixed-model analysis assessing the effects of pollinator dependence, growth form, and geographical region on the probability of yield decline (*i.e.*, the probability that a given crop in a given country shows an average annual growth rate in yield <0 over the period 1961-2020). GLMM_0 only evaluates the extent of (random) variation in the probability of yield decline among crops and countries. All the other four models include the effects of geographical region, and (cumulative) cultivated area per crop and country as fixed effects. Model GLMM_1a and model GLMM_1b test whether variation among crops in the probability of yield decline can be accounted for by pollinator dependence or by growth form, respectively. Model GLMM_2 tests for potential confounding effects as both factors, pollinator dependence and growth form, are associated to some extent (Fig. 3). Last, model GLMM_3 assesses whether any effect of pollinator dependence depends on growth form, or whether any effect of crop pollinator dependence or growth form differs among regions. Fixed effects (A) are evaluated statistically using Wald's type II tests (X^2 estimates that have a $P<0.05$ are boldfaced). The table also includes (B) estimates of the crop and country random effects with their respective 95% confidence intervals, (C) estimates of the percentage of variation in yield decline explained by the fixed factors included in each model (*i.e.*, marginal R^2) and by each model as a whole (*i.e.*, conditional R^2), and (D) comparisons with homologous phylogenetic GLMM models and estimates of phylogenetic signal in models' residuals.

(A) Fixed effects

		GLMM_0		GLMM_1a		GLMM_1b		GLMM_2		GLMM_3	
	Df	X^2	P	X^2	P	X^2	P	X^2	P	X^2	P
Pollinator dependence (P)	2	–	–	**6.71**	0.035	–	–	2.54	0.28	2.08	0.35
Growth form (G)	2	–	–	–	–	**15.95**	0.00034	**11.64**	0.0029	**11.93**	0.0026
Region (R)	4	–	–	**26.37**	$<10^{-4}$	**26.16**	$<10^{-4}$	**26.39**	$<10^{-4}$	**24.75**	$<10^{-4}$
P x G	4	–	–	–	–	–	–	–	–	4.06	0.40
P x R	8	–	–	–	–	–	–	–	–	**19.53**	0.012
G x R	8	–	–	–	–	–	–	–	–	**28.53**	0.00038
Area_crop	1	–	–	**37.51**	$<10^{-8}$	**36.54**	$<10^{-8}$	**34.26**	$<10^{-8}$	**35.66**	$<10^{-8}$
Area_country	1	–	–	**26.19**	$<10^{-7}$	**25.55**	$<10^{-6}$	**25.76**	$<10^{-6}$	**26.40**	$<10^{-6}$
Estimate		Slope	SE	Slope	SE	Slope	SE	Slope	SE	Slope	SE
Area_crop		–	–	-0.481	0.079	-0.461	0.076	-0.445	0.076	-0.467	0.078
Area_country		–	–	-0.359	0.070	-0.353	0.070	-0.354	0.070	-0.362	0.071

(B) Random effects

	GLMM_0		GLMM_1a		GLMM_1b		GLMM_2		GLMM_3	
	Variance	95%CI	Variance	95%CI	Variance	95%CI	Variance	95%CI	Variance	95%CI
Crop	0.466	(0.30-0.71)	0.248	(0.15-0.41)	0.217	(0.13-0.36)	0.204	(0.12-0.35)	0.197	(0.12-0.33)
Country	0.695	(0.49-0.99)	0.455	(0.31-0.66)	0.448	(0.31-0.66)	0.449	(0.31-0.66)	0.453	(0.31-0.67)

(C) Model explanatory power

	GLMM_0	GLMM_1a	GLMM_1b	GLMM_2	GLMM_3
Marginal R^2	0.000	0.101	0.110	0.111	0.138
Conditional R^2	0.261	0.259	0.260	0.258	0.280

(D) Phylogenetic effects

	GLMM_0		GLMM_1a		GLMM_1b		GLMM_2		GLMM_3	
	AIC	ΔAIC	AIC	ΔAIC	AIC	ΔAIC	AIC	ΔAIC	AIC	ΔAIC
GLMM	4229.11		4148.85		4140.31		4141.82		4130.16	
Phylo-GLMM	4232.34	3.23	4152.43	3.58	4147.19	6.88	4150.26	8.44	4138.74	8.58

Phylogenetic signal in the model residuals	Estimate	P	Estimate	P	Estimate	P	Estimate	P	Estimate	P
K	$4.51*10^{-7}$	0.35	$6.79*10^{-7}$	0.13	$6.39*10^{-7}$	0.15	$6.28*10^{-7}$	0.15	$6.28*10^{-7}$	0.15

Appendix

Fig. S1. A randomly-chosen sample of yield trends (1961-2020) for different crops in different countries. Point colors represent different categories of pollinator dependence (*i.e.*, none=yellow, modest=orange, high=red). The difference in the natural logarithm of yield between any two given years estimates the relative yield growth during that period, whereas the slope of the regression line of (natural-log) yield as a function of year estimates the annual growth rate in yield. The sign of the regression was used to characterize whether the yield of a given crop in a given country showed evidence of long-term decline or not. For instance, maize in the USA showed no evidence of decline, whereas cocoa in Costa Rica did.

Fig. S2. Variation in the probability of yield decline among crops. As a crop estimate of the probability of yield decline, the figure depicts the expected proportion of countries (+/- 1SE) in which each crop showed average growth rates in yield <0 over the period 1961-2020. Estimates are based on model GLMM_0 (Table 1). Crops are classified based on their pollinator dependence, growth form, and type of organ consumed.

Fig. S3. Variation in the probability of yield decline among countries. As a country estimate of the probability of yield decline, the figure depicts the estimated proportion of crops (+/- 1SE) showing average growth rates in yield <0 for each country over the period 1961-2020. Estimates are based on model GLMM_0 (Table 1). Countries are classified according to geographical region.

Fig. S4. Probability of yield decline by geographical region as estimated by the main effects models. The figure depicts the mean estimates (+/- 1 SE) of the proportion of crops showing negative average growth rates for countries in each region of the world (*i.e.*, AF, Africa; AM, the Americas; AS, Asia; EU, Europe; OC, Oceania) during the period 1961-2020 according to the main-effects GLMMs that included this factor (*i.e.*, GLMM_1a, GLMM_1b, GLMM_2). Means with the same letter do not provide evidence of statistical differences at the level of $\alpha = 0.05$ according to a pairwise Tukey's a posteriori test. Wald's type II test results are provided (see also Table 1).

Fig. S5. Probability of yield decline and cumulative cultivated area per crop and country. The figure depicts the mean estimates (+/- 1 SE) of the proportion of countries in which a given crop showed negative average growth rates (upper two panels) or the proportion of crops in a given country showing negative average growth rates (lower panel) as a function of the cumulative cultivated area (1961-2020) per crop or country, respectively. Estimates were extracted from model GLMM_2, even though the effect of area was very similar in all models that included the effects of cultivated area per crop and country (Table 1). In the upper panels, crops were categorized by pollinator dependence (left) and growth form (right), and in the lower panel countries were categorized by geographical region.

Table S1. Two-way frequency tables showing the number of crops in each combined category of pollinator dependence x growth form according to (A) the five pollinator-dependence levels defined by Klein et al. (2007), and (B) the three pollinator-dependence levels considered in this contribution resulting from the lumping of the levels highlighted with the same color as in (A).

A.

Growth form→ Pollinator dependence	Herb	Shrub	Tree
None (0% reduction)	51	6	10
Little (>0 to <10%)	10	3	6
Modest (≥10 to <40%)	8	4	5
High (≥40 to <90%)	5	3	16
Essential (≥90%)	5	0	4

B.

Growth form→ Pollinator dependence	Herb	Shrub	Tree
None	51	6	10
Modest	18	7	11
High	10	3	20

Reference

Klein, A.-M., B. E. Vaissière, J. H. Cane, I. Steffan-Dewenter, S. A. Cunningham, C. Kremen, and T. Tscharntke. 2007. Importance of pollinators in changing landscapes for world crops. Proceedings of the Royal Society B: Biological Sciences 274:303–313. https://doi.org/10.1098/rspb.2006.3721

Table S2. List of countries and number of crops per country with uninterrupted yearly yield records (1961-2020).

	Country	Number of crops
1	Afghanistan	23
2	Albania	15
3	Algeria	35
4	Angola	25
5	Antigua and Barbuda	12
6	Argentina	46
7	Australia	60
8	Austria	26
9	Bahamas	7
10	Bahrain	6
11	Bangladesh	47
12	Barbados	12
13	Belize	8
14	Benin	28
15	Bhutan	11
16	Bolivia (Plurinational State of)	46
17	Botswana	7
18	Brazil	52
19	Brunei Darussalam	10
20	Bulgaria	36
21	Burkina Faso	18
22	Burundi	19
23	Cabo Verde	11
24	Cambodia	18
25	Cameroon	30
26	Canada	41
27	Central African Republic	21
28	Chad	16
29	Chile	47
30	China	68
31	China, Taiwan Province of	49
32	Colombia	40
33	Comoros	8
34	Congo	23
35	Cook Islands	10
36	Costa Rica	23
37	Côte d'Ivoire	32
38	Cuba	28
39	Cyprus	38

Table S2
(cont.)

40	Democratic People's Republic of Korea	29
41	Democratic Republic of the Congo	46
42	Denmark	16
43	Dominica	10
44	Dominican Republic	34
45	Ecuador	56
46	Egypt	56
47	El Salvador	27
48	Equatorial Guinea	7
49	Eswatini	16
50	Faroe Islands	1
51	Fiji	19
52	Finland	20
53	France	53
54	French Polynesia	21
55	Gabon	13
56	Gambia	7
57	Germany	24
58	Ghana	29
59	Greece	47
60	Grenada	13
61	Guatemala	30
62	Guinea	22
63	Guinea-Bissau	11
64	Guyana	14
65	Haiti	29
66	Honduras	29
67	Hungary	32
68	Iceland	6
69	India	67
70	Indonesia	44
71	Iran (Islamic Republic of)	47
72	Iraq	34
73	Ireland	12
74	Israel	49
75	Italy	49
76	Jamaica	39
77	Japan	53
78	Jordan	32
79	Kenya	39
80	Kiribati	3

Table S2
(cont.)

81	Kuwait	3
82	Lao People's Democratic Republic	23
83	Lebanon	47
84	Lesotho	8
85	Liberia	17
86	Libya	28
87	Madagascar	51
88	Malawi	24
89	Malaysia	32
90	Maldives	8
91	Mali	24
92	Malta	2
93	Mauritania	13
94	Mauritius	24
95	Mexico	70
96	Mongolia	4
97	Morocco	47
98	Mozambique	32
99	Myanmar	32
100	Namibia	4
101	Nauru	1
102	Nepal	16
103	Netherlands	21
104	New Caledonia	9
105	New Zealand	38
106	Nicaragua	20
107	Niger	17
108	Nigeria	40
109	Niue	6
110	Norway	16
111	Oman	6
112	Pakistan	42
113	Panama	23
114	Papua New Guinea	22
115	Paraguay	29
116	Peru	67
117	Philippines	42
118	Poland	22
119	Portugal	38
120	Puerto Rico	19

Table S2

(cont.)

121	Republic of Korea	39
122	Romania	27
123	Rwanda	17
124	Saint Kitts and Nevis	3
125	Saint Lucia	7
126	Saint Vincent and the Grenadines	11
127	Samoa	15
128	Sao Tome and Principe	10
129	Saudi Arabia	15
130	Senegal	19
131	Seychelles	7
132	Sierra Leone	18
133	Singapore	1
134	Solomon Islands	11
135	Somalia	17
136	South Africa	59
137	Spain	54
138	Sri Lanka	40
139	Suriname	18
140	Sweden	17
141	Switzerland	25
142	Syrian Arab Republic	45
143	Thailand	49
144	Timor-Leste	11
145	Togo	21
146	Tokelau	2
147	Tonga	12
148	Trinidad and Tobago	26
149	Tunisia	37
150	Turkey	76
151	Tuvalu	1
152	Uganda	29
153	United Arab Emirates	4
154	United Kingdom of Great Britain and Northern Ireland	28
155	United Republic of Tanzania	42
156	United States of America	72
157	Uruguay	29
158	Vanuatu	6
159	Venezuela (Bolivarian Republic of)	41
160	Viet Nam	32
161	Yemen	14

Table S2

(cont.)

162	Zambia	14
163	Zimbabwe	25

Table S3. Summary of statistical results associated with the effects of the focal factors pollinator dependence and growth form on the probability of yield decline (*i.e.*, the probability that a given crop in a given country shows an average annual growth rate in yield <0 over a period of time), as evaluated separately by models GLMM_1a, GLMM_1b, respectively, and jointly by GLMM_2. The comparisons involve (1) the estimation of yield decline and their analyses considering the entire period 1961-2020, as it was carried in this study and reported in Table 1, *vs.* considering the period 1962-2019, thus curtailing the modal and expected highly influential years 1961 and 2020 (Fig. 1); and (2) considering pollinator dependence as a categorical variable with three levels (*i.e.*, none, moderate, and high), as it was carried out in this study after lumping, *vs.* considering pollinator dependence as a numerical variable with five possible values (*i.e.*, 0, 5, 25, 65, 95% corresponding to the categories none, little, modest, high and essential, respectively), as analyzed in Deguines et al. (2014). As expected, the estimate of the slope of the probability of yield decline with increasing pollinator dependence, when this variable was treated as numerical, was positive (logit estimate $\pm$1SE = 0.005478 $\pm$ 0.002203)

Time series: Pollinator dependence:	1961-2020 Categorical			1962-2019 Categorical			1961-2020 Numerical		
Model/ Focal factor	Df	X^2	P	Df	X^2	P	Df	X^2	P
GLMM_1a									
Pollinator dependence	2	**6.71**	0.035	2	5.55	0.062	1	**6.14**	0.013
GLMM_1b									
Growth form	2	**15.95**	0.00034	2	**16.08**	0.00032	2	**15.95**	0.00034
GLMM_2									
Pollinator dependence	2	2.08	0.35	2	1.57	0.46	1	1.54	0.21
Growth form	2	**11.93**	0.026	2	**11.85**	0.0027	2	**11.04**	0.004

Reference

Deguines N, Jono C, Baude M, et al (2014) Large-scale trade-off between agricultural intensification and crop pollination services. Frontiers in Ecology and the Environment 12:212–217. https://doi.org/10.1890/130054