Main Manuscript and Supporting information for

CyAbrB2 is a nucleoid-associated protein in *Synechocystis* controlling hydrogenase expression during

fermentation

Ryo Kariyazono[1] and Takashi Osanai[1]*

[1]School of Agriculture, Meiji University

*Takashi Osanai

Postal address: 1-1-1 Higashimita, Tama-ku, Kawasaki, Kanagawa 214-8571, Japan

Telephone/Fax: +81-44-934-7103

Email: tosanai@meiji.ac.jp

Author Contributions: RK and TO designed the study; RK performed the study; RK analyzed the data;

RK and TO wrote the paper

Competing Interest Statement: The authors declare no conflict of interest.

Classification:

Major: Biological sciences

Minor: Plant Biology

Keywords: cyanobacteria, transcription, microoxic response, hydrogenase, nucleoid-associated protein,

sigma factor

This PDF file includes:

Main Text

Figures 1 to 8

Table 1

Figures S1 to S10

Tables S1 to S4

Abstract

The *hox* operon in *Synechocystis* sp. PCC 6803, encoding bidirectional hydrogenase responsible for H_2

production, is transcriptionally upregulated under microoxic conditions. Although several regulators for

hox transcription have been identified, their dynamics and higher-order DNA structure of *hox* region in

microoxic conditions remain elusive. We focused on key regulators for the *hox* operon: cyAbrB2, a

conserved regulator in cyanobacteria, and SigE, an alternative sigma factor. Chromatin

immunoprecipitation-sequencing revealed that cyAbrB2 binds to the *hox* promoter region under aerobic

conditions, with its binding being flattened in microoxic conditions. Concurrently, SigE exhibited increased

localization to the *hox* promoter under microoxic conditions. Genome-wide analysis revealed that cyAbrB2

binds broadly to AT-rich genome regions and represses gene expression. Moreover, we demonstrated the

physical interactions of the *hox* promoter region with its distal genomic loci. Both the transition to

microoxic conditions and the absence of cyAbrB2 influenced the chromosomal interaction. From these

results, we propose that cyAbrB2 is a cyanobacterial nucleoid-associated protein (NAP), modulating

chromosomal conformation, which blocks RNA polymerase from the *hox* promoter in aerobic conditions.

We further infer that cyAbrB2, with altered localization pattern upon microoxic conditions, modifies

chromosomal conformation in microoxic conditions, which allows SigE-containing RNA polymerase to

access the *hox* promoter. The coordinated actions of this NAP and the alternative sigma factor are crucial

for the proper *hox* expression in microoxic conditions. Our results highlight the impact of cyanobacterial

chromosome conformation and NAPs on transcription, which have been insufficiently investigated.

Introduction

Cyanobacteria perform fermentation, using glycolytic products as electron acceptors. (1). Cyanobacteria

have multiple fermentation pathways according to the environment. For example, the freshwater living

cyanobacterium *Synechocystis* sp. PCC 6803 (hereafter referred to as *Synechocystis*) produces acetate,

lactate, dicarboxylic acids, and hydrogen (1, 2).

Hydrogen is generated in quantities comparable to lactate and dicarboxylic acids as the result of electron

acceptance in the dark microoxic condition (3, 4). Bidirectional hydrogenase is a key enzyme for H_2

production from protons (5) and is commonly found in cyanobacteria (6). Cyanobacterial hydrogenase

comprises five subunits (HoxEFUHY) containing nickel and Fe-S clusters (7). This enzyme can utilize

NADH, reduced ferredoxin, and flavodoxin as substrates (8). Hydrogenase mainly receives reduced

ferredoxin from pyruvate-ferredoxin oxidoreductase (PFOR) in the microoxic condition (8, 9).

Although hydrogenase and PFOR are O_2 sensitive, they can work under aerobic conditions (10–12).

Therefore, uncontrolled expression of *hox* operon and *nifJ* (coding gene of PFOR) may hamper metabolism

under photosynthetic conditions. Furthermore, genetic manipulations on *Synechocystis* have demonstrated

that modulating the expression of certain enzymes including hydrogenase can alter fermentative metabolic

flow (3, 4, 13). This provides evidence that transcription regulates the fermentative pathway. Thus,

transcriptional regulation in response to the environment is essential for optimal energy cost performance.

Promoter recognition by RNA polymerases is an essential step in transcriptional regulation. Sigma factors,

subunits of RNA polymerase, recognize core promoter sequences. Transcription factors can also bind to

promoter regions to suppress or promote RNA polymerase transcription. As well as recruitment or blocking

of RNA polymerase, some transcriptional factors, known as Nucleoid-associated proteins (NAPs),

modulate chromosomal conformation to regulate transcription (14). NAPs are common in bacteria, but

cyanobacterial NAPs remain unidentified, and higher-order DNA structure in cyanobacteria is yet to be

shown. A recent study suggested that the manipulation of chromosomal supercoiling impacts transcriptional

properties in cyanobacteria (15). There is room for consideration of NAPs modulating chromosomal

conformation and regulating expression in cyanobacteria.

In *Synechosysits*, the coding genes of HoxEFUHY form a single operon (sll1220–1226), while PFOR is

encoded in the *nifJ* (sll0741) gene. Both *hox* and *nifJ* operons are highly expressed under microoxic

conditions (16). Genetic analysis has revealed that multiple global transcriptional regulators control *hox*

and *nifJ* expression. Sigma factor SigE (Sll1689) promotes the expression of *hox* and *nifJ* operons (17, 18),

while transcription factor cyAbrB2 (Sll0822) represses them (19, 20). Positive regulators for the *hox* operon

include LexA (Sll1626) and cyAbrB1 (Sll0359) (21–23).

SigE, an alternative sigma factor, controls the expression of genes involved in glycogen catabolism and

glycolysis, as well as PFOR/*nifJ* and hydrogenase (17). SigE shows a high amino acid sequence similarity

with the primary sigma factor SigA, which is responsible for transcribing housekeeping and photosynthetic

genes (24). A ChIP-seq study revealed that, while most SigE-binding sites are the same as SigA, SigE

exclusively occupies the promoters of glycogen catabolism and glycolysis (25).

CyAbrB2 and its homolog cyAbrB1 are transcription factors highly conserved in cyanobacteria. For

example, cyAbrB homologs in *Anabaena* sp. PCC7120 is involved in heterocyst formation (26). CyAbrB2

in *Synechocystis* regulates the expression of several genes involved in carbon metabolism, nitrogen

metabolism, and cell division (20, 27, 28). CyAbrB2 binds to the *hox* promoter *in vitro* and represses its

expression *in vivo* (19). CyAbrB1, an essential gene, physically interacts with the cyAbrB2 protein (29)

and binds the *hox* promotor region *in vitro* to promote its expression (21).

To explore the dynamics of those transcription factors governing the expression of *hox* operon, we

conducted a time-course analysis of the transcriptome and ChIP-seq of SigE and cyAbrB2. Our ChIP-seq

and transcriptome analysis showed the NAPs-like nature of cyAbrB2, which prompted us to conduct a

chromosomal conformation capture assay. 3C analysis explored the physical interaction between the *hox*

promoter region and its downstream and upstream genomic region in the aerobic condition, and some loci

changed interaction frequency upon entry to the microoxic condition. Furthermore, some interactions in

the Δ*cyabrB2* mutant were different from those of the wildtype. From those experiments, we propose that

cyAbrB2 modulates chromosomal conformation like NAPs, allowing access to the SigE-containing RNA

polymerase complex on the *hox* promoter, by which the *hox* operon achieves distinct expression dynamics.

Chromosomal conformation of bacteria is a growing area of interest, and the findings of this study have

brought insight into the transcriptional regulation of cyanobacteria.

Results

Transcriptomes on entry to dark microoxic conditions

To understand transcriptional regulation under microoxic conditions, we conducted a time-course

transcriptome capturing light aerobic and dark microoxic conditions at 1-, 2-, and 4-hours time points

(Figure 1A). Gene set enrichment analysis based on KEGG pathway revealed that many biological

pathways, including photosynthesis and respiration (oxidative phosphorylation), were downregulated by

the transition to dark microoxic conditions from light aerobic conditions (Figure 1B). Upregulated

pathways included butanoate metabolism and two-component systems. The enrichment in the butanoate

metabolism pathway indicates the upregulation of genes involved in carbohydrate metabolism. We further

classified genes according to their expression dynamics. Within 1 hour of switching from aerobic to

microoxic conditions, the expression levels of 508 genes increased more than 2-fold. Furthermore, genes

with increased expression levels were classified into four groups based on the time course (Figures 1C and

S1). Of the 508 genes, 28 were termed "transiently upregulated genes" due to their decreased expression

upon the comparison of 1- and 4-hours incubation under microoxic conditions (log2 fold change < −0.5),

and 119 were termed "continuously upregulated genes.", which continuously increased between 1- and 4-

hours incubation under microoxic conditions (log2-fold change > 0.5). Other than 508 genes 2-fold

upregulated within 1 hour, 28 genes showed more than 4-fold increment within 4 hours but not upregulated

within 1 hour. We combined those "Late upregulated genes" with 508 genes and referred to as "All

upregulated genes" in the subsequent analysis (Figure S1). Mapping the classified genes to central carbon

metabolism revealed that *nifJ* encoding PFOR and *hox* operon encoding a bidirectional hydrogenase

complex were transiently upregulated (Figure 1D and Table 1). RTqPCR verified the transient expression

of *hoxF*, *hoxH,* and *nifJ* (Figure S2).

SigE and cyAbrB2 control the expression of transiently upregulated genes

The functional correlation between hydrogenase and PFOR, encoded by the *hox* operon and *nifJ,* suggests

that transient upregulation has physiological significance. We focused on transiently upregulated genes and

attempted to reveal the regulatory mechanism underlying transient upregulation. While SigE promotes the

expression of *hox* and *nifJ*, cyAbrB2 represses them (17, 19, 20). We confirmed that the deletion of *sigE*

and *cyabrb2* (Δ*sigE* and Δ*cyabrb2*, respectively) affected the expression of *hoxF*, *hoxH*, and *nifJ* by RT-

qPCR (Figure S1). Thus, we conducted a time-course transcriptome analysis of Δ*sigE* and Δ*cyabrb2* under

aerobic conditions and after 1- and 2-hours cultivation in dark microoxic conditions (Figures 2A and S3).

The transcriptome data confirmed that SigE and cyAbrB2 regulated *hox* operon expression (Figure 2B). At

each time point, we searched for differentially expressed genes (DEGs) between mutants and wildtype with

a more than 2-fold expression change and a false discovery rate (FDR) less than 0.05. We found that deleting

sigE or *cyabrb2* preferentially affected the expression of transiently upregulated genes, not limited to *hox*

and *nifJ* operons (Figures 2C and 2D). Interestingly, *cyabrb2* deletion resulted in the upregulated expression

of transient genes under aerobic conditions, in contrast to 1-hour cultivation under microoxic conditions

(Figure 2C).

Genome-wide analysis of cyAbrB2, cyAbrB1, and SigE via ChIP-seq

To decipher the regulatory mechanism of transiently upregulated genes, we must first comprehend the

fundamental features and functions of these transcriptional regulators. Therefore, a genome-wide survey of

cyAbrB2 and SigE occupation (Figure S4) combined with transcriptome data was done. Specifically, we

generated a *Synechocystis* strain in which cyAbrB2 was epitope-tagged and performed a ChIP-seq assay

under aerobic and microoxic conditions (Figure S4A). SigE-tagged strains previously constructed and

analyzed elsewhere were also employed (25). The primary sigma factor SigA was also analyzed to

determine SigE-specific binding. In addition to cyAbrB2, we tagged and analyzed cyAbrB1, which is the

interactor of cyAbrB2 and positively regulates the *hox* operon.

CyAbrB2 binds to long tracts of the genomic region and suppresses genes in the binding region

The ChIP-seq data showed that cyAbrB2 bound to long tracts of the genomic region with lower GC

content than the whole genome *Synechocystis* (Figures 3A and 3B). Vice versa, regions exhibiting lower

GC contents showed a greater binding signal of cyAbrB2 (Figure 3C). This correlation was not a systematic

bias of next-generation sequencing because the binding signals of SigE, SigA, and control showed no

negative correlation to GC contents (Figures S5A and S5B). The binding regions of cyAbrB2 called by

peak-caller covered 15.7% of the entire genome length. 805 of 3614 genes overlapped with cyAbrB2

binding regions, and almost half (399 of 805 genes) were entirely covered by cyAbrB2 binding regions.

The cyAbrB2 binding regions included 80 of 125 insertion sequence elements (Figure 3D). Comparison

with the transcriptome of ΔcyabrB2 revealed that cyAbrB2 tended to suppress the genes overlapping with

its binding regions under aerobic conditions (Figures 3A and 3E). A survey of the genomic localization of

cyAbrB1 under aerobic conditions revealed that cyAbrB1 and cyAbrB2 shared similar binding patterns

(Figures 3A and S5C). Due to the essentiality of cyAbrB1, we did not perform transcriptome analysis on a

cyAbrB1-disrupted strain. Instead, we referred to the recent study performing transcriptome of cyAbrB1

knockdown strain, whose cyAbrB1 protein amount drops by half (30). Among 24 genes induced by

cyAbrB1 knockdown, 12 genes are differentially downregulated genes in cyabrb2Δ in our study (Figure

S5D).

CyAbrB2 binds to transiently upregulated genes

The binding regions of cyAbrB2 overlapped 17 of 28 transiently upregulated genes, showing significant

enrichment from all upregulated genes (Figure 4A). The transiently upregulated genes belong to 17

transcriptional units (TUs), according to the previous study(31), and cyAbrB2 tends to bind TUs with

transiently upregulated genes (Figure 4B). While cyAbrB2 covered the entire length of insertion sequences

and unknown proteins, its binding positions on other transient genes were diverse (Figure 4C). Specifically,

the *hox* and *nifJ* operons had two distinct binding regions located at the transcription start sites (TSSs) and

middle of operons, the *pntAB* operon had two binding regions in the middle and downstream of the operon,

and the *nrtABCD* operon had one binding region downstream of the operon (Figure 4C).

Localization of cyAbrB2 became blurry under the microoxic condition

When cells entered microoxic conditions, the relative ChIP-seq signals in the cyAbrB2 binding regions

slightly declined (Figures 5A and 5B). Notably, the total quantities of precipitated DNA by tagged cyAbrB2

did not decrease (Figure S6A), and qPCR confirmed that the cyAbrB2 binding signal increased in all

positions tested (Figure 5C). ChIP-seq data and ChIP-qPCR data indicate that the boundary between

cyAbrB2 binding region and cyAbrB2-free region became obscured when the cells entered microoxic

conditions due to increased binding of cyAbrB2 to both cyAbrB2 binding and cyAbrB2-free region. The

protein amount of cyAbrB2 was not altered on entry to the microoxic condition (Figure S6B). The cyAbrB2

binding signal around the transiently upregulated genes became less specific upon entry into microoxic

conditions, consistent with the general tendency (Figure 5B). The amount of DNA immunoprecipitated by

cyAbrB1 was also increased in the microoxic condition, and the protein amount was not increased (Figures

S6C and S6D).

Sigma factors SigE and SigA are excluded from cyAbrB2 binding regions regardless of GC contents

We searched for SigE and SigA binding sites under aerobic and microoxic conditions (Figures S7A and

S7B, left and right, respectively). The SigE and SigA peaks identified in this study predominantly covered

the previously identified peaks (Figure S7C), reproducing the previous study's conclusion (25); i.e., SigE

and the primary sigma factor SigA share localization on the promoters of housekeeping genes, but SigE

exclusively binds to the promoters of its dependent genes. SigE and SigA binding peaks were significantly

excluded from the cyAbrB2 binding regions (Figures 6A and 6B). SigE preferred the cyAbrB2-free region

in the aerobic condition more than SigA did (Odds ratios of SigE and SigA being in the cyAbrB2-free

region were 4.88 and 2.74, respectively). CyAbrB2 prefers AT-rich regions, but no correlation was found

between the GC content and binding signals of SigE and SigA (Figures S5A and S5B). Thus, SigA and

SigE are excluded from cyAbrB2 binding regions regardless of GC contents.

Dynamics of sigma factors upon exposure to the microoxic condition

When cells entered microoxic conditions, the binding signals of SigA and SigE were changed, although

most of their peaks observed under aerobic conditions were present under microoxic conditions (Figure

S7A). The preference of SigE for the cyAbrB2-free region was alleviated in the microoxic condition (Figure

6A). Next, we focused on sigma factor dynamics in transiently upregulated genes. SigE, but not SigA, binds

at the TSS of *pntAB* under aerobic and microoxic conditions (Figure 6C top). SigE binding summits were

not identified at the TSSs of the *hox* and *nifJ* operons under aerobic conditions. However, the SigE-specific

binding summit appeared at the TSS of *nifJ* when cells entered microoxic conditions (Figure 6C middle).

A bimodal peak of SigE was observed at the TSS of the *hox* operon in a microoxic-specific manner (Figure

6C bottom panel). The downstream side of the bimodal SigE peak coincides with the SigA peak and the

TSS of TU1715. Another side of the bimodal peak lacked SigA binding and was located at the TSS of the

hox operon (marked with an arrow in Figure 6C), although the peak caller failed to recognize it as a peak.

SigE binding without SigA on the promoters of *hox*, *nifj*, and *pntAB* is consistent with their SigE-dependent

expression (Figure 2B).

Chromatin conformation around *hox* operon and *nifJ* operon

We have shown that cyAbrB2 broadly binds to AT-rich genomic regions, including insertion element

sequences, and represses expression (Figure 3). This is functionally similar to the NAPs (14), which

makes us hypothesize that cyAbrB2 modulates chromosomal conformation. Therefore, we conducted the

chromatin conformation capture (3C) assay against wildtype and *cyabrb2*Δ strains at aerobic and

microoxic conditions. qPCR was performed with unidirectional primer sets, where the genomic fragment

containing *hox* operon and *nif* operon (hereinafter *hox* fragment and *nifJ* fragment, respectively) were

used as bait (Figure 7).

First, focusing on the aerobic condition of wildtype (Figure 7B, solid line), the *hox* fragment interacted

with its proximal downstream loci (loci (f) to (g)) and proximal upstream locus (locus (j)). The *hox*

fragment also interacts with the distal downstream locus (locus (c)). Meanwhile, the *nifJ* fragment shows

high interaction frequency with proximal upstream and downstream loci (Figure 7G, loci (i') and (j')),

and a distal downstream locus (locus (g')) showed higher interaction frequency with *nifJ* fragment than

proximal locus (h') did. The upstream regions of *nifJ* (loci (l') to (n') and (p')) showed comparable

frequency with locus (g').

The chromatin conformation is changed in *cyabrb2*Δ in some loci

Then we compared the chromatin conformation of wildtype and *cyabrb2*Δ. Although overall shapes of

graphs did not differ, some differences were observed in wildtype and *cyabrb2*Δ (Figures 7B and 7G);

interaction of locus (c) with *hox* region were slightly lower in *cyabrb2*Δ and interaction of loci (f') and

(g') with *nifJ* region were different in wildtype and *cyabrb2*Δ. Note that the interaction scores exhibit

considerable variability and we could not detect statistical significance at those loci.

Changes of chromatin conformation upon microoxic condition

When the cells entered the microoxic condition, proximal loci interacted more frequently (Figure 7D;

loci (f)-(h) and Figure 7I loci (j') and (k')). This tendency was more apparent in *cyabrb2*Δ (Figures 7E

and 7J). Furthermore, the interaction of *nifJ* upstream loci (l') – (n') increased in the microoxic condition

in *cyabrb2*Δ but not wildtype (Figures 7I and 7J). The locus (c) and locus (j) interacted less frequently

with *hox* fragment upon entry to the microoxic condition in the wildtype. While the interaction scores

exhibit considerable variability, the individual data over time demonstrate declining trends of the wildtype

at locus (c) and (j) (Figure S8). In Δ*cyabrb2*, by contrast, the interaction frequency of loci (c) and (j) was

unchanged in the aerobic and microoxic conditions (Figure 7E). The interaction frequency of locus (c) in

Δ*cyabrb2* was as low as that in the microoxic condition of wildtype, while that of locus (j) in Δ*cyabrb2*

was as high as that in the aerobic condition of wildtype (Figures 7B and 7C). In summary, 3C analysis

demonstrated cyAbrB2-dependent and independent dynamics of chromosomal conformation around the

hox and *nifJ* operon in response to the microoxic condition (Figure 8).

Discussion

Physiological significance of transient upregulation of *hox* and *nifJ* operons

As the transcriptional change can alter the metabolic flow, the transcriptional upregulation of fermentative

genes in response to the microoxic condition is expected to be adaptive for energy acquisition and the

maintenance of redox balance. Our time-course transcriptome showed upregulation of several genes

involved in catabolism upon exposure to the microoxic condition. The transient upregulation of *hox* and

nifJ operons is distinctive among them (Figure 1D).

One reason for transient upregulation is probably the resource constraints of inorganic cofactors.

Hydrogenase and PFOR (the product of *nifJ* gene) have iron-sulfur clusters, and hydrogenase requires

nickel for its activity (32, 33). Overexpression of the *hox* operon should be futile under physiological

conditions without an adequate nickel supply (34).

Another significance for transient upregulation may be the reusability of fermentative products. Hydrogen,

lactate, and dicarboxylic acids can be reused as the source of reducing power when cells return to aerobic

conditions (12, 35, 36). The substrate proton is abundant, but hydrogen is diffusive and difficult to store.

Therefore, hydrogenase may favor fermentation initiation, and the reductive branch of TCA-producing

dicarboxylic acids may become active subsequently. In fact, *citH*/*mdh* (sll0891) encoding a key enzyme of

the reductive branch of TCA was classified as continuously upregulated genes in this study (Figures 1C and

1D).

Mechanisms for transient expression mediated by SigE and cyAbrB2

265   SigE and cyAbrB2 can independently contribute to the transient transcriptional upregulation. This is

266   evident as the single mutants, Δ*sigE* or Δ*cyabrb2*, maintained transient expression of *hoxF* and *nifJ* (Figure

267   S2). We first discuss cyAbrB2 as the potential NAPs, and then the mechanism of transient upregulation

268   mediated by cyAbrB2 and SigE will be discussed.

cyAbrB2 is a novel nucleoid-associated protein of cyanobacteria

270   We have shown that cyAbrB2 broadly binds to AT-rich genomic regions, including IS elements (Figure

271   3). This is functionally similar to the histone-like nucleoid protein H-NS family; including H-NS in

272   *Enterobacteriaceae* (37, 38)*, and* Lsr2 in *Mycobacteria* (39). Like H-NS and Lsr2, cyAbrB2 may defend

273   against exogenous DNA elements, which often have different GC content. Interestingly, Lsr2 controls genes

274   responding to hypoxia, showing a functional analogy with cyAbrB2 (40).

The biochemistry of cyAbrB2 will shed light on the regulation of chromatin conformation in the

future

H-NS proteins often cause bound DNA to bend, stiffen, and/or bridge (14). DNA-bound cyAbrB2 is

expected to oligomerize, based on the long tract of cyAbrB2 binding region in our ChIP-seq data. However,

no biochemical data mentioned the DNA deforming function or oligomerization of cyAbrB2 in the previous

studies, and preference for AT-rich DNA is not fully demonstrated in vitro (19, 27, 41). Moreover, our 3C

data did not support bridging at least in *hox* region and *nifJ* region, as the high interaction locus and

cyAbrB2 binding region did not seem to correlate (Figure 7). Therefore, direct observation of the DNA-

cyAbrB2 complex by atomic force microscopy is the solution in the future.

Not only DNA structural change but also the effect of the post-translational modification can be

investigated by biochemistry. The previous studies report that cyAbrB2 is subject to phosphorylation and

glutathionylation (42, 43), and pH and redox state alters cyAbrB1's affinity to DNA (41). Those

modifications might respond to environmental changes and be involved in transient expression. Overall,

the biochemistry integrating assay conditions (PTM, buffer condition, and cooperation with cyAbrB1) and

output (DNA binding, oligomerization, and DNA structure) will deepen the understanding of cyAbrB2 as

cyanobacterial NAPs.

Cooperative and antagonistic function of cyAbrB1 and cyAbrB2

CyAbrB1, the homolog of cyAbrB2, may cooperatively work, as cyAbrB1 directly interacts with cyAbrB2

(29), their distribution is similar, and they partially share their target genes for suppression (Figures 3A S5C

and S5D). The possibility of cooperation would be examined by the electrophoretic mobility shift assay of

cyAbrB1 and cyAbrB2 as a complex. Despite their similar repressive function, cyAbrB1 and cyAbrB2

regulate *hox* expression in opposite directions, and their mechanism remains elusive. The stoichiometry of

cyAbrB1 and cyAbrB2 bound to DNA fluctuates in response to the environmental changes (28), but there

was no difference in the behavior of cyAbrB1 and cyAbrB2 around the *hox* region on entry to the microoxic

condition.

Localization pattern and function of cyAbrB2

Herein, we classified three types of binding patterns for cyAbrB2. The first is that cyAbrB2 binds a long

DNA tract covering the entire gene or operon, represented by the insertion sequence elements. CyAbrB2

suppresses expression in this pattern (Figure 3E). In the second pattern, cyAbrB2 binds on promoter regions,

such as *hox* operon and *nifJ*. The binding on those promoters fluctuates in response to environmental

changes, thus regulating expression. This pattern also applies to the promoter of *sbtA* (Na^+/HCO_3^-

symporter), where cyAbrB2 is bound in a CO_2 concentration-dependent manner (28). The last one is

cyAbrB2 binding in the middle or downstream of operons. The middle of *hox*, *pntAB*, and *nifJ* operons and

the downstream of *nrt* operon are the cases (Figure 4C). Our data show that genes in the same operon

separated by the cyAbrB2 binding region behave differently. In particular, *pntB* is classified as the

transiently upregulated gene, while *pntA* is not, despite being in the same operon. This might be explained

by the recent study which reported that cyAbrB2 affects the stability of mRNA transcribed from its binding

gene (41). The cyAbrB2-mediated stability of mRNA may also account for the decrease in transcript from

transient upregulated genes at 4 hours of cultivation. Hereafter, we will focus on the mechanism of the

second pattern, regulation by cyAbrB2 on the promoter.

Insight into the regulation of *hox* and *nifJ* operon by cyAbrB2

Genome-wide analysis indicates that the cyAbrB2-bound region blocks SigE and SigA (Figures 6A and

6B). This is presumably because sigma factors recognize the promoter as a large complex of RNA

polymerase. CyAbrB2 binds to the *hox and nifJ* promoter region and may inhibit access to RNA polymerase

complex under aerobic conditions. When cells entered microoxic conditions, the boundaries of the cyAbrB2

binding region and cyAbrB2-free region became obscure (Figure 5), and SigE binding peaks on those

promoters became prominent (Figure 6C). Notably, cyAbrB2 ChIP efficiency at the *hox* promoter is higher

in the microoxic condition than in the aerobic condition (Figure 5). Hence, while the exclusion by cyAbrB2

occupancy on promoter inhibits containing RNA polymerase in the aerobic condition, it is also plausible

that chromosomal conformation change governed by cyAbrB2 provides SigE-containing RNA polymerase

with access to the promoter region (Figure 8). Our 3C result demonstrated that cyAbrB2 influences the

chromosomal conformation of *hox* and *nifJ* region to some extent (Figure 7).

A recent study demonstrated that manipulating the expression of topoisomerase, which influences

chromosomal conformational change through supercoiling, affects transcriptional properties in

cyanobacteria (15). Moreover, Song et al. pointed out that DNA looping may inhibit transcription in

cyanobacteria (41) because artificial DNA looping by the LacI repressor of *E. coli* can repress

cyanobacterial transcription (44). Thus, we infer conformation change detected by the present 3C

experiment regulates expression of *hox* operon.

Generality for chromosomal conformation in cyanobacteria

Our 3C analysis revealed that local chromosomal conformation changes upon entry to the microoxic

conditions (Figure 8). As cyAbrB2 occupies about 15% of the entire genome and globally regulates gene

expression, cyAbrB2 likely governs the whole chromosomal conformation. Furthermore, the

conformational changes by deletion of cyAbrB2 were limited, suggesting there are potential NAPs in

cyanobacteria yet to be characterized. It is speculated that conformational change of the entire chromosome

occurs to deal with many environmental stresses.

The sigE-mediated mechanism for the transient expression

One possible SigE-mediated mechanism for transient expression is the post-transcriptional activation and

degradation of SigE in the dark; i.e., SigE is sequestered by anti-sigma factor under light conditions and

released under dark (45), enabling acute transcription of *hox* operon and *nifJ*. Transcripts of *sigE* were

continuously downregulated in our time-course transcriptome, while *sigB* (sll0306) and *sigC* (sll0184) were

classified as continuous upregulated genes (Supplemental Table S1). It is possible that upregulated SigB

and SigC outcompete SigE in prolonged incubation under microoxic conditions. Finally, SigE is degraded

under dark within 24 hours (46).

Another reason for the microoxic specific expression may exist in the sequence of the *hox* promoter. We

previously determined the consensus sequence of -10 element for SigE regulon in the aerobic condition as

"TANNNT," where N is rich in cytosine (25). The -10 sequence of the *hox* promoter "TAACA<u>A</u>" (23)

deviates from the consensus, and no hexamer precisely fitting the consensus is found in the *nifJ* promoter.

This deviation can inhibit SigE from binding during aerobic conditions, aside from cyAbrB2-mediated

inhibition. Under the microoxic condition, transcription factors LexA (23) and Rre34 (16) may aid SigE

binding to the promoter of *hox* and *nifJ*, respectively.

Moreover, SigE seems susceptible to the blocking from cyAbrB2 during the aerobic condition compared

with SigA. This is supported by the odds ratio of SigE being in the cyAbrB2-free region was higher than

that of SigA in the aerobic condition (Figures 6A and 6B). The higher exclusion pressure of cyAbrB2 on

SigE may contribute to sharpening the transcriptional response of hox and nifJ on entry to microoxic

conditions. Overall, multiple environmental signals are integrated into the *hox* and *nifJ* promoter through

the cyAbrB2 and SigE dynamics.

Materials and Methods

Bacterial strains and plasmids

The glucose-tolerant strain of *Synechocystis* sp. PCC 6803 (47) was used as a wildtype strain in this study.

The *sigE* (sll1689)-disrupted strain (G50), SigE FLAG-tagged strain, and SigA FLAG-tagged strain were

constructed in a previous study (17, 25). Disruption and epitope-tagging of *cyabrb1*(sll0359) and

cyabrb2(sll0822) were performed by homologous double recombination between the genome and PCR

fragment (47). The resulting transformants were selected using three passages on BG-11 plates containing

5 µg/mL kanamycin. Genomic PCR was used to confirm the insertion of epitope tag fragments and gene

disruption (Figure S9). Supplemental Tables S2, S3, and S4 contain the cyanobacterial strains,

oligonucleotides, and plasmids used in this study.

Aerobic and microoxic culture conditions

For aerobic conditions, cells were harvested after 24 hours cultivation in HEPES-buffered BG-11$_0$ medium

(48), which was buffered with 20 mM HEPES-KOH (pH 7.8) containing 5 mM NH$_4$Cl under continuous

exposure to white light (40 µmol m^{-2}s^{-1}) and bubbled with air containing 1% CO$_2$ (final OD$_{730}$ = 1.4–1.8).

For the dark microoxic culture, the aerobic culture cell was concentrated to an OD$_{730}$ of 20 with the

centrifuge and resuspended in the culture medium. The concentrated cultures were poured into vials,

bubbled with N$_2$ gas, and sealed. The sealed vials were shaded and shaken at 30˚C for the described times.

Antibodies and immunoblotting

Sample preparation for immunoblotting was performed as previously described (25), and FLAG-tagged

proteins were detected by alkaline-phosphatase-conjugated anti-FLAG IgG (A9469, Sigma Aldrich, St.

Louis, MO) and 1-Step NBT/BCIP substrate solution (Thermo Fisher Scientific, Waltham, MA).

RNA isolation

Total RNA was isolated with ISOGEN (Nippon gene, Tokyo, Japan) following the manufacturer's

instructions and stored at −80˚C until use. The extracted RNA was treated with TURBO DNase (Thermo

Fisher Scientific) for 1 hour at 37˚C to remove any genomic DNA contamination. We confirmed that the

A260/A280 of the extracted RNA was >1.9 by NanoDrop Lite (Thermo Fisher Scientific). We prepared

triplicates for each timepoint for the RNA-seq library. RT-qPCR was performed as described elsewhere

(46).

ChIP assay

Two biological replicates were used for each ChIP-seq experiment, and one untagged control ChIP was

performed. ChIP and qPCR analyses were performed using the modified version of a previous method (25).

FLAG-tagged proteins were immunoprecipitated with FLAG-M2 antibody (F1804 Sigma-Aldrich)

conjugated to protein G dynabeads (Thermo Fisher Scientific).

Library preparation and next-generation sequencing

For the ChIP-seq library, input and immunoprecipitated DNA were prepared into multiplexed libraries

using NEBNext Ultra II DNA Library Prep Kit for Illumina (New England Biolabs, Ipswich, MA). For the

RNA-seq library, isolated RNA samples were deprived of ribosomal RNA with Illumina Ribo-Zero Plus

rRNA Depletion Kit (Illumina, San Diego, CA) and processed into a cDNA library for Illumina with the

NEBNext Ultra II Directional RNA Library Prep Kit for Illumina (New England Biolabs). Dual-index

primers were conjugated with NEBNext Multiplex Oligos for Illumina (Set1, New England Biolabs). We

pooled all libraries, and the multiplexed libraries were dispatched to Macrogen Japan Inc. and subjected to

paired-end sequencing with HiSeqX. Adapter trimming and quality filtering of raw sequence reads were

conducted with fastp (ver. 0.21.0) (49) under default conditions. The paired-end sequences were mapped

onto the *Synechocystis* genome (ASM972v1) using Bowtie2 (50) (ver. 2.4.5 paired-end). Supplementary

Table S5 contains the read counts that passed via fastp quality control and were mapped by Bowtie2.

RNA-seq analysis

Mapped reads were counted by HT-seq count (ver. 2.0.2) (51) for the GFF file of ASM972v1, with the

reverse-strandedness option. EdgeR package (ver. 3.40.1) (52) was used to perform the differential

expression analysis. Fold changes in expression and FDR were used for gene classification. Supplemental

Table S6-S8 contains fold change in gene expression calculated by edgeR.

Genome-wide analyses

Peaks were called using the MACS3 program (ver. 3.0.0b1) (53). For paired-end reads for SigE, SigA,

and untagged control ChIP, narrow peaks were called with <1e−20 of the q-value cut-off and "--call-

summits" options. The peak summits from two replicates and the untagged control were merged if summits

were located within 40 bp of each other. Peak summits identified in both replicates but not in the control,

were considered for further analysis. The midpoint of the peak summits for the two merged replicates was

further analyzed.

Broad peak calling methods were applied to paired-end reads for cyAbrB2, cyAbrB1, and untagged

control ChIP using the "–broad" option, with a q-value cut-off of < 0.05 and a q-value broad cut-off of <

0.05. The intersection of broad peaks from two replicates, excluding those called by the control, was used

in subsequent analyses.

The positions of the TSS, including internal start sites, were obtained as reported by (31). The read count,

merging, and intersection of the binding region were calculated using BEDTools (ver. 2.30.0) (54).

Supplemental Tables S9–S15 contain SigA and SigE peaks and the broad binding regions of cyAbrB2 and

cyAbrB1, respectively.

Binding signals in every 100 bp bin for scatter plots were calculated as (IP read counts within 100 bp

window) / (input read counts within 100 bp window) * (total input read counts/total IP read counts). GC

contents were calculated within 500 bp in 100 bp sliding windows by seqkit (ver. 2.3.0) (55).

Genome extraction, digestion, and ligation for 3C assay

A 3C assay was conducted based on the previous prokaryotic Hi-C experiment (56, 57), with certain steps

modified. To begin, *Synechocystis* were fixed with 2.5% formaldehyde for 15 minutes at room temperature.

Fixation was terminated by adding a final concentration of 0.5M of glycine, and cells were stored at -80˚C

until use. Fixed cells were disrupted using glass beads and shake master NEO (Bio Medical Science, Tokyo,

Japan), following the previous study's instructions for preparing cell lysate for ChIP. The lysates were

incubated with buffer containing 1mM Tris-HCl (pH7.5), 0.1mM EDTA, and 0.5% SDS for 10min at room

temperature, and 1% Triton X-100 quenched SDS. Genomes in the cell lysate were digested by 600U/mL

of HindIII (Takara Bio, Shiga, Japan) for 4 hours at 37˚C, and RNA in the lysate was simultaneously

removed by 50 µg/mL of RNaseA (Nippon genetics, Tokyo, Japan). The digestion was terminated by adding

1% SDS and 22 mM EDTA. The fill-in reaction and biotin labeling steps were omitted from the procedure.

The digested genomes were diluted by ligation buffer containing 1% triton-X100 to the final concentration

of approximately 1μg/mL and incubated for 10 min at room temperature. Ligation was performed with 2

U/mL of T4 DNA ligase (Nippon Gene) overnight at 16˚C. Crosslinking was reversed under 65˚C for 4

hours in the presence of 2.5 mg/mL of proteinase K (Kanto Chemical, Tokyo, Japan), and DNA was purified

with the phenol-chloroform method and ethanol precipitation method.

Preparation of calibration samples for 3C qPCR

Based on a previous study, calibration samples for possible ligated pairs were prepared in parallel with 3C

ligation (58). In brief, the purified genome of *Synechocystis* was digested by HindIII, and DNA was purified

with the phenol-chloroform and ethanol precipitation. Purified DNA was dissolved into the ligation buffer

at a concentration of about 600ng/μL and ligated with 2 U/mL of T4 DNA ligase at 16˚C overnight.

Quantification of crosslinking frequency for 3C assay

Before the real-time PCR assay, we confirmed that each primer set amplified single bands in a ligation-

dependent manner by GoTaq Hot Start Green Master Mix (Promega, Madison, WI) (Figure S10). Real-

time PCR was performed with StepOnePlus (Applied Biosystems, Foster City, CA) and Fast SYBR Green

Master Mix (Thermo Fisher Scientific) according to the manufacturer's instructions. Interaction frequency

was calculated by ΔΔCt method using dilution series of calibration samples described above. We confirmed

each primer set amplified DNA fragment with a unique Tm value. The amount of the bait fragment

containing *hox* operon were quantified and used as an internal control. Supplemental Table S4 contains the

list of primers used in the 3C quantification. Interaction frequency for each primer position was calculated

as the relative abundance of ligated fragments against the calibration samples and normalized among

samples by internal control.

Statistical analysis

Statistical analyses were performed with R version 4.2.2 (59). The "fisher.test" function was used for

Fisher's exact test, and p-values < 0.05 were denoted as asterisks in the figure. Multiple comparisons of

Fisher's exact test were conducted using "fisher. Multcomp" function in the RVAideMemoire package (60),

where p-values were adjusted by the "fdr" method and FDRs <0.05 are shown in the figures. Multiple

comparisons of the Wilcoxon-rank test were conducted by "pairwise.wilcox.test," and p-values were

adjusted by the "fdr" method. Adjusted p-values < 0.05 are shown in the figure. The correlation coefficient

was calculated with the "cor" function. Gene set enrichment analysis (GSEA) was performed by

culsterPlofiler package (61) in R with p-value cut-off of 0.05. The enriched pathways detected by GSEA

are listed in Supplemental Table S16.

Manuscript writing

We employed GPT-3.5 to refine the expression of the manuscripts.

Accession Numbers

Raw ChIP sequencing and RNA sequencing reads were deposited in the Sequence Read Archive (accession

ID: PRJNA956842).

Declaration of interest

The authors declare no competing interests.

Acknowledgments

This study was supported by the following grants to T.O.: Grant-in-Aid for Scientific Research (B) (grant

no. 20H02905), JST-ALCA of the Japan Science and Technology Agency (grant number JPMJAL1306),

the Asahi Glass Foundation, and JST-GteX (grant number JPMJGX23B0). We thank Dr. Kohki Yoshimoto

for providing laboratory instruments and Ms. Kaori Iwazumi for the support of bacterial culture and the

medium preparation.

Author contributions

R.K. and T.O. designed the study; R.K. conducted the experiment; R.K. analyzed the data; R.K. and T.O.

wrote the paper.

References

1. L. J. Stal, R. Moezelaar, Fermentation in cyanobacteria. *FEMS Microbiol. Rev.* **21**, 179–211 (1997).

2. T. Osanai, *et al.*, Genetic manipulation of a metabolic enzyme and a transcriptional regulator

increasing succinate excretion from unicellular cyanobacterium . *Front. Microbiol.* **6**, 1064

(2015).

3. H. Iijima, *et al.*, Metabolomics-based analysis revealing the alteration of primary carbon

metabolism by the genetic manipulation of a hydrogenase HoxH in *Synechocystis* sp. PCC 6803.

Algal Res. **18**, 305–313 (2016).

4.   M. Akiyama, T. Osanai, Regulation of organic acid and hydrogen production by NADH/NAD(+)

ratio in *Synechocystis* sp. PCC 6803. *Front. Microbiol.* **14**, 1332449 (2023).

5.   D. Carrieri, K. Wawrousek, C. Eckert, J. Yu, P.-C. Maness, The role of the bidirectional

hydrogenase in cyanobacteria. *Bioresour. Technol.* **102**, 8368–8377 (2011).

6.   V. Puggioni, S. Tempel, A. Latifi, Distribution of Hydrogenases in Cyanobacteria: A Phylum-Wide

Genomic Survey. *Front. Genet.* **7**, 223 (2016).

7.   C. Cassier-Chauvat, T. Veaudor, F. Chauvat, Advances in the function and regulation of

hydrogenase in the cyanobacterium *Synechocystis* PCC6803. *Int. J. Mol. Sci.* **15**, 19938–19951

(2014).

8.   K. Gutekunst, *et al.*, The bidirectional NiFe-hydrogenase in *Synechocystis* sp. PCC 6803 is reduced

by flavodoxin and ferredoxin and is essential under mixotrophic, nitrate-limiting conditions. *J.*

Biol. Chem. **289**, 1930–1937 (2014).

9.   J. H. Artz, *et al.*, The structure and reactivity of the HoxEFU complex from the cyanobacterium

Synechocystis sp. PCC 6803. *J. Biol. Chem.* **295**, 9445–9454 (2020).

10. Y. Wang, *et al.*, Pyruvate:ferredoxin oxidoreductase and low abundant ferredoxins support aerobic

photomixotrophic growth in cyanobacteria. *Elife* **11**, 2021.08.27.457892 (2022).

11.   H. Burgstaller, *et al.*, *Synechocystis* sp. PCC 6803 Requires the Bidirectional Hydrogenase to

515   Metabolize Glucose and Arginine Under Oxic Conditions. *Front. Microbiol.* **13**, 896190 (2022).

516   12.   J. Appel, S. Phunpruch, K. Steinmüller, R. Schulz, The bidirectional hydrogenase of *Synechocystis*

517   sp. PCC 6803 works as an electron valve during photosynthesis. *Arch. Microbiol.* **173**, 333–338

518   (2000).

519   13.   H. Iijima, *et al.*, Four-carbon dicarboxylic acid production through the reductive branch of the open

520   cyanobacterial tricarboxylic acid cycle in *Synechocystis* sp. PCC 6803. *Metab. Eng.* **65**, 88–98

521   (2021).

522   14.   J. Hołówka, J. Zakrzewska-Czerwińska, Nucleoid Associated Proteins: The Small Organizers That

523   Help to Cope With Stress. *Front. Microbiol.* **11**, 590 (2020).

524   15.   A. Behle, *et al.*, Manipulation of topoisomerase expression inhibits cell division but not growth

525   and reveals a distinctive promoter structure in Synechocystis. *Nucleic Acids Res.* **50**, 12790–12808

526   (2022).

527   16.   T. C. Summerfield, S. Nagarajan, L. A. Sherman, Gene expression under low-oxygen conditions

528   in the cyanobacterium *Synechocystis* sp. PCC 6803 demonstrates Hik31-dependent and -

529   independent responses. *Microbiology* **157**, 301–312 (2011).

530   17.   T. Osanai, *et al.*, Positive regulation of sugar catabolic pathways in the cyanobacterium

531   *Synechocystis* sp. PCC 6803 by the group 2 sigma factor sigE. *J. Biol. Chem.* **280**, 30653–30659

532   (2005).

533 18. T. Osanai, *et al.*, Genetic engineering of group 2 sigma factor SigE widely activates expressions of

534   sugar catabolic genes in *Synechocystis* species PCC 6803. *J. Biol. Chem.* **286**, 30962–30971 (2011).

535 19. J. Dutheil, *et al.*, The AbrB2 autorepressor, expressed from an atypical promoter, represses the

536   hydrogenase operon to regulate hydrogen production in *Synechocystis* strain PCC6803. *J. Bacteriol.*

537   **194**, 5423–5433 (2012).

538 20. C. Leplat, *et al.*, Genome-wide transcriptome analysis of hydrogen production in the

539   cyanobacterium Synechocystis: Towards the identification of new players. *Int. J. Hydrogen Energy*

540   **38**, 1866–1872 (2013).

541 21. P. Oliveira, P. Lindblad, An AbrB-Like protein regulates the expression of the bidirectional

542   hydrogenase in *Synechocystis* sp. strain PCC 6803. *J. Bacteriol.* **190**, 1011–1019 (2008).

543 22. K. Gutekunst, *et al.*, LexA regulates the bidirectional hydrogenase in the cyanobacterium

544   *Synechocystis* sp. PCC 6803 as a transcription activator. *Mol. Microbiol.* **58**, 810–823 (2005).

545 23. P. Oliveira, P. Lindblad, LexA, a transcription regulator binding in the promoter region of the

546   bidirectional hydrogenase in the cyanobacterium *Synechocystis* sp. PCC 6803. *FEMS Microbiol.*

547   *Lett.* **251**, 59–66 (2005).

548 24. S. Imamura, M. Asayama, Sigma factors for cyanobacterial transcription. *Gene Regul. Syst. Bio.* **3**,

549   65–87 (2009).

550   25.   R. Kariyazono, T. Osanai, Identification of the genome-wide distribution of cyanobacterial group-

551   2 sigma factor SigE, accountable for its regulon. *Plant J.* **110**, 548–561 (2022).

552   26.   A. Higo, E. Nishiyama, K. Nakamura, Y. Hihara, S. Ehira, cyAbrB Transcriptional Regulators as

553   Safety Devices To Inhibit Heterocyst Differentiation in Anabaena sp. Strain PCC 7120. *J.*

554   *Bacteriol.* **201** (2019).

555   27.   A. Ishii, Y. Hihara, An AbrB-like transcriptional regulator, Sll0822, is essential for the activation

556   of nitrogen-regulated genes in *Synechocystis* sp. PCC 6803. *Plant Physiol.* **148**, 660–670 (2008).

557   28.   J. Lieman-Hurwitz, *et al.*, A cyanobacterial AbrB-like protein affects the apparent photosynthetic

558   affinity for CO2 by modulating low-CO2-induced gene expression. *Environ. Microbiol.* **11**, 927–

559   936 (2009).

560   29.   Y. Yamauchi, Y. Kaniya, Y. Kaneko, Y. Hihara, Physiological roles of the cyAbrB transcriptional

561   regulator pair Sll0822 and Sll0359 in *Synechocystis* sp. strain PCC 6803. *J. Bacteriol.* **193**, 3702–

562   3709 (2011).

563   30.   A. Hishida, *et al.*, CRISPRi knockdown of the cyabrB1 gene induces the divergently transcribed

564   icfG and sll1783 operons related to carbon metabolism in the cyanobacterium *Synechocystis* sp.

565   PCC 6803. *J. Gen. Appl. Microbiol.* (2024) https:/doi.org/10.2323/jgam.2024.01.001.

566   31.   M. Kopf, *et al.*, Comparative genome analysis of the closely related *Synechocystis* strains PCC

567   6714 and PCC 6803. *DNA Res.* **21**, 255–266 (2014).

568   32.   K. Uyeda, J. C. Rabinowitz, Pyruvate-ferredoxin oxidoreductase. 3. Purification and properties of

569   the enzyme. *J. Biol. Chem.* **246**, 3111–3119 (1971).

570   33.   P. M. Vignais, B. Billoud, Occurrence, classification, and biological function of hydrogenases: an

571   overview. *Chem. Rev.* **107**, 4206–4272 (2007).

572   34.   M. Ortega-Ramos, *et al.*, Engineering *Synechocystis* PCC6803 for hydrogen production: influence

573   on the tolerance to oxidative and sugar stresses. *PLoS One* **9**, e89372 (2014).

574   35.   N. Katayama, K. Iwazumi, H. Suzuki, T. Osanai, S. Ito, Malic Enzyme, not Malate Dehydrogenase,

575   Mainly Oxidizes Malate That Originates from the Tricarboxylic Acid Cycle in Cyanobacteria.

576   *MBio* **13**, e0218722 (2022).

577   36.   S. A. Angermayr, *et al.*, Chirality Matters: Synthesis and Consumption of the d-Enantiomer of

578   Lactic Acid by *Synechocystis* sp. Strain PCC6803. *Appl. Environ. Microbiol.* **82**, 1295–1304

579   (2016).

580   37.   W. W. Navarre, M. McClelland, S. J. Libby, F. C. Fang, Silencing of xenogeneic DNA by H-NS-

581   facilitation of lateral gene transfer in bacteria by a defense system that recognizes foreign DNA.

582   *Genes Dev.* **21**, 1456–1471 (2007).

583   38.   T. Oshima, S. Ishikawa, K. Kurokawa, H. Aiba, N. Ogasawara, Escherichia coli Histone-Like

584   Protein H-NS Preferentially Binds to Horizontally Acquired DNA in Association with RNA

585   Polymerase. *DNA Res.* **13**, 141–153 (2006).

39.   B. R. G. Gordon, *et al.*, Lsr2 is a nucleoid-associated protein that targets AT-rich sequences and

virulence genes in Mycobacterium tuberculosis. *Proc. Natl. Acad. Sci.* **107**, 5154–5159 (2010).

40.   M. Kołodziej, *et al.*, Lsr2 and Its Novel Paralogue Mediate the Adjustment of Mycobacterium

smegmatis to Unfavorable Environmental Conditions. *mSphere* **6** (2021).

41.   K. Song, *et al.*, Expression of the Cyanobacterial F(o)F(1) ATP Synthase Regulator AtpΘ Depends

on Small DNA-Binding Proteins and Differential mRNA Stability. *Microbiol. Spectr.* **10**,

e0256221 (2022).

42.   P. Spät, V. Krauspe, W. R. Hess, B. Maček, N. Nalpas, Deep Proteogenomics of a Photosynthetic

Cyanobacterium. *J. Proteome Res.* **22**, 1969–1983 (2023).

43.   S. Sakr, *et al.*, The activity of the *Synechocystis* PCC6803 AbrB2 regulator of hydrogen production

can be post-translationally controlled through glutathionylation. *Int. J. Hydrogen Energy* **38**,

13547–13555 (2013).

44.   D. Camsund, T. Heidorn, P. Lindblad, Design and analysis of LacI-repressed promoters and DNA-

looping in a cyanobacterium. *J. Biol. Eng.* **8**, 4 (2014).

45.   T. Osanai, *et al.*, ChlH, the H subunit of the Mg-chelatase, is an anti-sigma factor for SigE in

Synechocystis sp. PCC 6803. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 6860–6865 (2009).

46.   H. Iijima, *et al.*, Changes in primary metabolism under light and dark conditions in response to

overproduction of a response regulator RpaA in the unicellular cyanobacterium *Synechocystis* sp.

PCC 6803 . *Front. Microbiol.* **6**, 888 (2015).

47.   J. G. K. Williams, "Construction of Specific Mutations in Photosystem II Photosynthetic Reaction

Center by Genetic Engineering Methods in *Synechocystis* 6803" in *Methods in Enzymology*,

(Academic Press, 1988), pp. 766–778.

48.   R. Rippka, J. Deruelles, J. B. Waterbury, Generic assignments, strain histories and properties of

pure cultures of cyanobacteria. *J. Gen. Microbiol.* **111**, 1–61 (1979).

49.   S. Chen, Y. Zhou, Y. Chen, J. Gu, fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics **34**, i884–i890 (2018).

50.   B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**, 357–359

(2012).

51.   S. Anders, P. T. Pyl, W. Huber, HTSeq--a Python framework to work with high-throughput

sequencing data. *Bioinformatics* **31**, 166–169 (2015).

52.   M. D. Robinson, D. J. McCarthy, G. K. Smyth, edgeR: a Bioconductor package for differential

expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140 (2010).

53.   Y. Zhang, *et al.*, Model-based analysis of ChIP-Seq (MACS). *Genome Biol.* **9**, R137 (2008).

54.   A. R. Quinlan, I. M. Hall, BEDTools: a flexible suite of utilities for comparing genomic features.

Bioinformatics **26**, 841–842 (2010).

55.   W. Shen, S. Le, Y. Li, F. Hu, SeqKit: A Cross-Platform and Ultrafast Toolkit for FASTA/Q File

Manipulation. *PLoS One* **11**, e0163962 (2016).

56.   N. Takemata, R. Y. Samson, S. D. Bell, Physical and Functional Compartmentalization of Archaeal

Chromosomes. *Cell* **179**, 165-179.e18 (2019).

57.   N. Takemata, S. D. Bell, Chromosome conformation capture assay combined with biotin

enrichment for hyperthermophilic archaea. *STAR Protoc.* **2**, 100576 (2021).

58.   M. Abou El Hassan, R. Bremner, A rapid simple approach to quantify chromosome conformation

capture. *Nucleic Acids Res.* **37**, e35 (2009).

59. ,   R: A Language and Environment for Statistical Computing (2022).

60.   M. Hervé, RVAideMemoire: Testing and Plotting Procedures for Biostatistics (2022).

61.   T. Wu, *et al.*, clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. *Innov.*

(Cambridge **2**, 100141 (2021).

Table 1 List of transiently upregulated genes

		Operon
Oxidoreductase		
sll0741	*nifJ*/"pyruvate-ferredoxin/flavodoxin oxidoreductase"	TU3296
sll0743	hypothetical protein	
sll0744	dihydroorotate dehydrogenase (fumarate)	
sll1221	*hoxF*/"bidirectional [NiFe] hydrogenase diaphorase subunit"	TU1714
sll1222	unknown protein	
sll1223	*hoxU*/"bidirectional [NiFe] hydrogenase diaphorase subunit"	
sll1224	*hoxY*/"NAD-reducing hydrogenase small subunit"	
sll1225	unknown protein	
sll1226	*hoxH*/"NAD-reducing hydrogenase large subunit"	
slr1434	*pntB*/"H+-translocating NAD(P) transhydrogenase subunit beta"	TU1089
Transporter		
sll1450	*nrtA*/"nitrate/nitrite transport system substrate-binding protein"	TU1023
sll1451	*nrtB*/"nitrate/nitrite transport system permease protein"	
sll1452	*nrtC*/"nitrate/nitrite transport system ATP-binding protein"	
sll1453	*nrtD*/"nitrate/nitrite transport system ATP-binding protein"	
Two-component system		
slr1214	twitching motility two-component system response regulator PilG	TU905
slr1215	unknown protein	TU907
Glycosyl transferase		
slr2116	*spsA*/"spore coat polysaccharide biosynthesis protein; SpsA"	TU1673

Protease		
sll1009	*frpC*/"iron-regulated protein"	TU491
Insertion sequence (Transposase)		
slr1523	transposase	TU1659
sll1985	transposase	TU1589
sll7001	transposase	NA
sll7003	toxin FitB	TU7001
ssl0172	transposase	TU3163
Other		
slr1260	hypothetical protein	TU1446
slr0668	unknown protein	TU3532
slr5127	unknown protein	TU5127
sll0710	unknown protein	TU97
sll1307	unknown protein	TU1224

635   The list of transiently upregulated genes was merged by transcriptional units and sorted by function. The

636   transcriptional unit information was obtained from a previous study (Kopf et al., 2014).

Figure 1

Figure 1

(A) Schematic diagram for the sampling of cells under aerobic and microoxic conditions. (B) Gene set

enrichment analysis on time-course transcriptome data. KEGG pathways enriched in upregulated or

downregulated genes after 1-, 2-, and 4-h incubation under microoxic conditions are shown. (C) (left)

Heatmap showing expression change in all upregulated genes over the time-course. Genes classified into

Transient (striped square), Plateau (open square), Continuous (filled square), and Late (dotty square) were

clustered into subgroups and sorted by the gene name. (right) Examples of genes are classified into each

expression pattern. (D) The classified genes were mapped to central carbon metabolism, centered on

pyruvate. PEP: phosphoenolpyruvate, PYR: pyruvate, AcCoA: acetyl CoA, Ac-P: acetyl phosphate,

OXA: oxaloacetate, PHB: polyhydroxy butyrate. TCA: tricarboxylic acid cycle.

Figure 2

A

B

Expression fold change (Log₂)

C

Fraction of genes under cyAbrB2 dependent expression

Gene expression upon cyAbrB2 deletion
☐ down-regulated
▨ none
■ up-regulated

D

Fraction of genes under SigE dependent expression

Gene expression upon SigE deletion
☐ down-regulated
▨ none
■ up-regulated

Figure 2

(A) MA plot showing fold change (y-axis) and average (x-axis) of gene expression between wildtype and

mutant strains at each timepoint. Red dots indicate defined DEGs ($|Log_2$ FC| >1 with FDR < 0.05). (B)

Log2 scaled expression fold change in genes in the *hox* and *nifJ* operons upon Δ*cyabrb2* and Δ*sigE* under

aerobic conditions (0 h), 1 hour after microoxic condition (1 hr), and 2 hours after microoxic condition (2

hr). DEGs are marked with sky blue (downregulated upon deletion) or red (upregulated upon deletion).

(C and D) Fraction of upregulated and downregulated genes upon the (C) Δ*cyabrb2* and (D) Δ*sigE* at the

timepoints of aerobic conditions (0 hr), 1 hour after anoxic condition (1 hr), and 2 hour after anoxic

condition (2 hr). Genes are classified according to Figure 1C. Asterisk (*) and dagger (†) denote

statistically significant enrichment and anti-enrichment compared with all upregulated genes tested by

multiple comparisons of Fisher's exact test (FDR < 0.05).

Figure 3

Figure 3

(A) Snapshot of ChIP-seq data for cyAbrB2 and cyAbrB1 under aerobic conditions. The heatmap in the

second column indicates expression fold change upon Δ*cyabrb2* under aerobic conditions. Positive values

(colored in red) indicate that the gene expression is higher in wildtype than in Δ*cyabrb2*, and negative

values (colored in blue) indicate the opposite. The positions for the insertion elements are marked with

red in the third column. The heatmap in the fourth column indicates GC contents. High GC contents are

colored in blue and low GC contents are colored in blue. (B) GC contents and region length of cyAbrB2

binding regions (black dots). The horizontal dotted line indicates the genomic average of GC content. (C)

Scatter plot of GC content and binding signal of cyAbrB2. The x-axis is the binding signal of cyAbrB2 in

each 100 bp region, and the y-axis indicates GC contents within 500 bp windows sliding every 100 base

pairs. CyAbrB2 binding regions are marked with red dots. (D) Histogram of genes showing the extent of

occupancy (not bound, partially overlapped, or entirely overlapped) by the cyAbrB2 binding region. The

gray bars indicate non-IS genes, and the count numbers of the non-IS genes are displayed on the gray

bars. The black bars indicate the IS genes, and the count numbers of the IS genes are displayed above the

black bars. (E) Boxplot showing fold change in gene expression by Δ*cyabrb2* under aerobic conditions.

Genes are classified according to the extent of occupancy by the cyAbrB2 binding region. Asterisk (*)

denotes statistical significance tested by multiple comparisons of the Wilcoxon-rank test.

Figure 4

A

Fraction of abrB2 binding genes

- ■ abrB2 binding
- □ non-abrB2 binding

All genes (n=3608), All up-regulated (n=536), Transient up-regulated (n=28)

*

B

cyAbrB2 occupancy of Transcriptional Units (TUs)

extent of cyAbrB2 overlap
- no overlap
- 0~33%
- 33~66%
- 66~100%
- 100%

All TUs
- 265TUs
- 108TUs
- 120TUs
- 170TUs
- 3428TUs (83.8%)

TUs including transient up-regulated genes
- 2TUs (11.7%)
- 5TUs (29.4%)
- 5TUs (29.4%)
- 2TUs (11.7%)
- 3TUs (17.6%)

C

□ **no overlap**

Genes &operons / cyAbrB2-3xFLAG / binding region

slr1260, sll1009, slr1214 slr1215

□ **0~33% overlap**

(narB) nrtD nrtC nrtB nrtA (sll1450-sll1454)
sll0744 sll0743 nifJ(sll0741)
(pntA) pntB(sll1434)

□ **33~66% overlap**

slr5127
hoxH hoxY hoxU hoxF (hoxE) (sll1220-sll1226)

10kb

□ **66~100% overlap**

slr0668, sll1307, ssl0172

slr1523, sll1710

■ **100% overlap**

sll1985, slr2116

Figure 4

(A) Fraction of genes overlapped or non-overlapped with cyAbrB2 binding regions at the timepoints of

aerobic conditions. Genes are classified according to Figure S1. Asterisk (*) denotes statistically

significant enrichment compared with all upregulated genes tested by multiple comparisons of Fisher's

exact test. (B) Pie charts of transcriptional units (TUs) classified by extent of overlapping with cyAbrB2

binding region. The left pie represents all TUs, and the right pie represents only TUs containing the

transient upregulated genes. (C) Distribution of cyAbrB2 in the aerobic condition around transiently

upregulated genes. Arrows with bold lines indicate transiently upregulated genes. Shaded arrows indicate

operons whose data were obtained from a previous study. The bars below the graph indicate the binding

regions of each protein. The black bar at the top of the figure indicates a length of 10 kbp.

Figure 5

Figure 5

(A) Scatter plot showing changes of the binding signal by 1hr cultivation in the microoxic condition. The

binding signal of each 100bp window is plotted. Red dots are cyAbrB2 binding regions in either aerobic

or microoxic conditions. The dotty lines indicate Log2 fold enrichment of 0.5, 0, and -0.5 between

aerobic and microoxic conditions. (B) Distribution of cyAbrB2 around *hox* operon and *nifJ* operon. ChIP-

seq data in aerobic (L + O_2) and dark microoxic (D − O_2) conditions are overlayed. The bars below the

graph indicate the binding regions of each protein. (C) Quantification for IP efficiency of cyAbrB2 (top)

and cyAbrB1 (middle) by qPCR in the aerobic and microoxic conditions. The position of primers and

ChIP-seq data of cyAbrB2 are shown at the bottom. Scores are normalized by the IP% at position #2 in

the aerobic condition.

Figure 6

A SigE summits vs cyAbrB2 binding regions

B SigA summits vs cyAbrB2 binding regions

Figure 6

(A, B) Anti-co-occurrence of cyAbrB2 binding regions and sigma factors. Mosaic plots of cyAbrB2

binding regions and SigE peaks (A) or SigA binding peaks (B) are shown. Odds and p-values were

calculated by Fisher's exact test. (C) Snapshots of ChIP-seq data for CyAabrB2, SigE, and SigA at the

nifJ region (top) and *hox* region (bottom). ChIP-seq data for cyAbrB2, SigE, and SigA under aerobic and

dark microoxic conditions are overlayed. ChIP-seq data of cyAbrB2 under aerobic and microoxic

conditions are colored blue and pink, respectively. ChIP-seq data for SigE and SigA are shown in solid

lines (aerobic conditions) and the area charts (microoxic conditions). The positions of TSSs were obtained

from a previous study (Kopf et al., 2014) and indicated by vertical dotted lines. Open triangles indicate

peak summits under aerobic conditions, and solid triangles indicate peak summits under microoxic

conditions.

Figure 7

Figure 7

726 (A) and (F)Schematic diagram of 3C analysis around *hox* operon (A) and *nifJ* operon (F). In the panels

727 (A) and (F), the black horizontal arrow shows the location of the bait primer, and white horizontal arrows

728 ((a) to (n) in *hox* operon (A) and (a') to (t') in *nifJ* operon (F)) indicate loci where the interaction

729 frequency with bait were assayed. Vertical black arrowheads indicate the position of HindIII sites. ChIP-

730 seq data of cyAbrB2 in the aerobic condition is displayed in the bottom, and cyAbrB2 binding regions are

731 marked with shade. (B - E) The line plot showing the interaction frequency of each locus with *hox*

732 fragment. Two of data sets are presented; (B) wildtype vs Δ*cyabrb2* in aerobic condition, (C) wildtype vs

733 Δ*cyabrb2* in 1hr of microoxic condition, (E) wildtype in aerobic vs 1hr of microoxic condition, and (E)

734 Δ*cyabrb2* in aerobic vs 1hr of microoxic condition are compaired. (G – J) The line plot showing the

735 interaction frequency of each locus with *nifJ* fragment. Two data sets are selected and presented; (G)

736 wildtype vs Δ*cyabrb2* in aerobic condition, (H) wildtype vs Δ*cyabrb2* in 1hr of microoxic condition, (I)

737 wildtype in aerobic vs 1hr of microoxic condition, and (J) Δ*cyabrb2* in aerobic vs 1hr of microoxic

738 conditions are compared. The line plots indicate the average interaction frequency over the random

739 ligation (n=3). Individual data are plotted as dots.

743 **Figure 8**

744 Schematic diagram of the dynamics of transcription factors governing fermentative gene expression.

Figure S1

Figure S1

Schematic diagram showing the classification of genes according to the time-course transcriptome.

Transient (striped square), Plateau (open square), Continuous (filled square), and Late (dotty square) are

denoted as all upregulated genes.

Figure S2

Figure S2

RT-qPCR validated the transiently upregulated genes classified by RNAseq. Transcripts extracted from

wildtype (solid line), Δ*sigE* mutant (dotty line), Δ*cyabrb2* mutant (dashed line), and Δ*sigE* Δ*cyabrb2*

double mutant (dot-dashed line) were assayed in the aerobic condition (0hr) and 1,2,4 hour incubation of

microoxic conditions. As the representative of the transiently upregulated genes, expression of *hoxF*,

hoxY, nifJ, and *sll0744* were quantified by RT-qPCR. The line represents the mean of n=3, and individual

data points are shown as dot plots. Data of each gene is normalized by the mean score of wildtypes in the

aerobic condition.

Figure S3

Figure S3

Primary component scatter plot showing the profiles of RNA-seq data.

Figure S4

A

B

Figure S4 (continuing)

Figure S4 The procedure for ChIP-seq of flag-tagged proteins and Overview of ChIP-seq data

Validation of procedure for ChIP-seq of flag-tagged cyAbrB2, SigE, and SigA.

(A) The immunoblot for inputs and immunoprecipitants (IP) of ChIP for cyAbrB2-FLAG and cyAbrB1-

FLAG. Input lysate of untagged control (GT) is also loaded. Inputs equivalent to the indicated portion of

IP were loaded. (B)Scatter plots showing the reproducibility of two replicates for ChIP-seq assay. ChIP-

seq data of SigE, SigA, and cyAbrB2 in aerobic and microoxic conditions and ChIP-seq data of cyAbrB1

in the aerobic condition are shown. Dots indicate normalized IP read count / normalized input read count

in each 100bp window. X-axis is the value of replicate1, and Y-axis is the value of replicate 2. (C) and (D)

Overview for ChIp-seq of flag tagged cyAbrB2, cyAbrB1, SigE, and SigA. Y-axis indicates [normalized

IP read count / normalized input read count at each 25bp window], and X-axis indicates chromosome

position. (C) Distribution of cyAbrB2, cyAbrB1, SigE, and SigA across the whole genome of

Synechocystis. Aerobic (L +O_2) and dark microoxic (D -O_2) data are displayed. (D) Magnified image for

chromosome position of 1,550kb-1,800kb. ChIP-seq data of cyAbrB2, cyAbrB1, SigE, and SigA in

aerobic and dark microoxic conditions are overlayed. The dots below the graph indicate the binding

region of each protein calculated by peak caller.

Figure S5

Figure S5 GC content vs ChIP enrichment score of SigA and SigE

(A) Scatter plot showing GC contents in each 100bp vs. binding signal of SigA, SigE, and control IP.

Data are displayed as in Figure 3C. (B) GC content in each 100bp of (left) SigE peaks and non-SigE

peaks, and (right) SigA peaks and non-SigA peaks. (C) Venn diagram showing overlap of the binding

region of cyAbrB1 and cyAbrB2 (left), and scatter plot showing ChIP binding signal of cyAbrB2 (y-axis)

and cyAbrB1(x-axis) in the aerobic condition. Data is plotted as in Figure 5A. (D) Fractions of

upregulated and downregulated genes upon the Δ*cyabrb2* mutant in the aerobic conditions. Fractions of

all genes (left n=3608) and genes induced by cyAbrB1 knockout (right n=24) are shown. Genes induced

by cyAbrB1 knockout are from the previous study (30).

Figure S6

A

Quantity of DNA (ng)
by cyAbrB2 Immunoprecipitation

B

Normalized signal of
cyAbrB2-3FLAG

C

Quantity of DNA (ng)
by cyAbrB1 Immunoprecipitation

D

Normalized signal of
cyAbrB1-3FLAG

Figure S6 Alteration of cyAbrB2 binding to genome under the microoxic condition

(A) and (C) Amount of precipitated DNA by cyAbrB2 ChIP and cyAbrB1 ChIP, respectively. Three

experiments were performed in the aerobic and microoxic conditions. (B) and (D) Western blot images of

cyAbrB2-3FLAG and cyAbrB1-3FLAG, respectively. Proteins were extracted in the aerobic condition

and 1hr and 4hr incubation under microoxic conditions. The total protein concentration of each sample

was adjusted to 4mg/mL, measured by the BCA method. Quantification of cyAbrB2 from western blot

image was performed by ImageJ (ver. 2.0.0-rc-65) and plotted in the right graph.

Figure S7

A

Number of SigE binding peaks

L +O$_2$ D -O$_2$

283 | 347 | 202

Number of SigA binding peaks

L +O$_2$ D -O$_2$

160 | 453 | 116

B

Binding signal of SigE

Log2 FC > 0.5 : 37.6%

(Dark -O$_2$ 1hr)

Log2 FC < -0.5 : 34.7%

(Light +O$_2$)

Binding signal of SigA

Log2 FC > 0.5 : 31.5%

(Dark -O$_2$ 1hr)

Log2 FC < -0.5 : 31.4%

(Light +O$_2$)

C

SigE binding peaks

This study (aerobic)   Previous study

195 | 435 | 43

SigA binding peaks

This study (aerobic)   Previous study

91 | 522 | 333

binding signals for SigE peaks

peaks called only in previous

peaks newly called

Peak enrichment scores for previous ChIP−seq

Peak enrichment scores for ChIP−seq of this study

binding signals for SigA peaks

peaks called only in previous

peaks newly called

Peak enrichment scores for previous ChIP−seq

Peak enrichment scores for ChIP−seq of this study

Figure S7

(A) Venn diagram showing the number of peaks of SigE (left) and SigA (right) in aerobic (L + O_2) and

dark microoxic (D − O_2) conditions. (B) Scatter plot showing changes in the binding signal of SigE and

SigA by 1 hour cultivation under microoxic conditions. The binding signal of each 100 bp window is

plotted. (C) Reproducibility of ChIP-seq data of SigA and SigE, compared with the previous study

(Kariyazono and Osanai 2022). (top) Venn diagrams show the overlapping of peaks called in this study

and the previous study. (bottom) Scatter plot comparing ChIP binding signals of SigA and SigE peaks

commonly called in present and previous studies. Plots boxed by dashed lines are peaks called only in the

present or previous study.

Figure S8
hox region (Figures 7A, 7B and 7C)

Figure S8 (continuing)
  nifJ region (Figures 7D, 7E and 7F)

Figure S8 Dynamics of individual 3C scores

Re-plotting of Figure 5B and 5C with the x-axis showing time (0,1,4 hr in microoxic conditions) and the

y-axis showing the interaction frequency. Plots from the individual samples are connected by solid

(Wildtype) or dotty (Δ*cyabrb2*) lines.

Figure S9

Figure S9

Confirmation of genomic deletion and the epitope tagging of *abrB2* (#1-#3), the epitope tagging of *abrB1* (#4 and #5), and deletion of *sigE* (#6 and #7). Dotty lines are homologous regions between plasmids and the genome. Arrows indicate the position of check primers.

Figure S10

A   **3C primers for *hox* operon region**

M:100bp ladder (DM2100 SMOBIO)
1 Calibration sample x500
2 Calibration sample x2500
3 3C sample x100
4 Unligated sample x100

GoTaq® Hot Start
Program

95˚C 2min
95˚C 15sec
60˚C 30sec x35
4˚C   Hold

Figure S10 (continuing)

B   **3C primers for *nifJ* region**

M:100bp ladder (DM2100 SMOBIO)
1 Calibration sample x1000
2 3C sample x100
3 Unligated sample x100

GoTaq® Hot Start
Program

95˚C 2min
95˚C 15sec
60˚C 30sec
4˚C   Hold

Figure S10 Validation of unidirectional primer sets for 3C assay

The validation of unidirectional primer sets for 3C assay is shown in Figure 7. The 3C sample in this

assay is the mixture of all 3C samples assayed in Figure 7.

Table S1 Fold changes of transcripts from *sigA, sigB, sigC, sigD, and sigE*

| Sigma factor | locus | 0hr vs 1hr | | 1hr vs 4hr | |
		Log2FC	FDR	Log2FC	FDR
SigA	slr0653	-0.873248	0.00766486	-0.0013514	0.99797563
SigB	sll0306	1.38098826	8.42E-06	0.77453605	0.04057775
SigC	sll0184	2.97101055	1.75E-16	1.30743549	0.00067892
SigD	sll2012	0.4701823	0.1498473	-0.4522181	0.32402556
SigE	sll1689	-1.9111759	1.96E-11	-1.1223298	0.00633142

Data is extracted from Supplemental Table S7.

Table S2 List of strains used in this study

Strain ID	Host strain	genotype	Reference
GT			Williams, 1988
G50	GT	Δ*sigE::*KmR	T. Osanai, et al,2005
KR93	GT	SigA-8His-KmR	(Kariyazono and Osanai 2022)
KR94	GT	SigA-3FLAG-KmR	(Kariyazono and Osanai 2022)
KR340	GT	Δ*cyabrb2::*KmR	in this study
KR338	GT	cyAbrB(sll0359)-3xFLAG-KmR	in this study
KR339	GT	cyAbrB2(sll0822)-3xFLAG-KmR	in this study
KR359	KR340	Δ*cyabrB2::*KmR Δ*sigE::*CmR	in this study

Table S3 plasmids used in this study

Plasmid ID		backbone	
VK82	*sigE*ΔCmR	pTA2 (Toyobo)	transformation for KR359
VK200	AbrB1-3F-KmR	pTA2 (Toyobo)	transformation for KR338

VK201	AbrB2-3F-KmR	pTA2 (Toyobo)	transformation for KR339
VK203	*cyabrB2*ΔKmR	pTA2 (Toyobo)	transformation for KR340

842 **Table S4 oligonucleotides used in this study**

Oligonucleotides ID		sequence
qPCR_hoxF_F	Figure S2	GTCATCTGCAATGCTGACGAA
qPCR_hoxF_R	Figure S2	CCAACACACTGCGGTCCAT
qPCR_hoxH_F	Figure S2	TCCCGTTACCCGGATTGA
qPCR_hoxH_R	Figure S2	GCCCTGGTCGTTGAGGAAA
qPCR_nifJ_sll0741_F	Figure S2	GATTCGGGGCAAAGTTTACG
qPCR_nifJ_sll0741_R	Figure S2	TACCTGGATGGTGAACTTAG
qPCR_sll0744_F	Figure S2	GCAGCTTAAATGGTTCTACC
qPCR_sll0744_R	Figure S2	CCAAATCTGTGGGTACGTAG
qPCR_nrpB_F	Figure S2	AAAGGGTAAGGGTGCAAAGG
qPCR_nrpB_R	Figure S2	AATTCCTCAAGCGGTTCCAC
new_abrB_0822_del_chkF	Figure S9	AATCCTCCGCCAAACTTTG
abrB_0822_tag4	Figure S9	GACTGATGGTGAAAACTCC
abrB_0359_chkF	Figure S9	AGAGGGAAAAATTCCTCGGC
abrB_0359_tag4	Figure S9	TCGCCAGGGAAAGTAATTGG
SigE_whole_del_chkF	Figure S9	GGAACGCTTGGAATGGCAAC
sigE_tag4	Figure S9	TGAAGGGGGTAGATTTAAGCC
ChIP_slr2118_F1	Figure 5C #1	CATTTCAAGCGAGATAGCAG
ChIP_slr2118_R1	Figure 5C #2	TGGAAAGTGCCTGTTCAAAG
ChIP_ThoxH_F	Figure 5C #2	CATCACCGAGGGCATATCTAG
ChIP_ThoxH_R	Figure 5C #2	ACATAAGGCTATGGAAACC
ChIP_pHox_2F	Figure 5C #3	ACAGAGGATGGACAAATGTC
ChIP_pHox_2R	Figure 5C #3	TACCGCTTCGTCATTCTGCT
ChIP_pHox_up_2R	Figure 5C #4	TCGATTACTTCCCTTTAGTG
ChIP_pHox_up_2F	Figure 5C #4	CTTCGGCTCCATGTATTC
ChIP_sll0814_F	Figure 5C #5	GTTACTACCGCTATCTATCG

ChIP_sll0814_R	Figure 5C #5	ATGACGTAAACCGGCTCTGAC
3Chox_Hcommon	Figure 7 (hox region)	CGGTCCCTTTGGACGTAATC
3Chox_H01_pos_+1	Figure 7 (hox region)	GGGTTGGGCGCAACATTAAG
3Chox_H02_pos_+3	Figure 7 (hox region)	CAAAGCGATTTTGGCTGGTG
3Chox_H03_pos_+4	Figure 7 (hox region)	GATGTGGTGGAAGATACGGAC
3Chox_H04_pos_+5	Figure 7 (hox region)	GGAGCCACAGGGGAATTTAAC
3Chox_H05_pos_+6	Figure 7 (hox region)	GGTCAGATTTTATTGGTTGCATGG
3Chox_H06_pos_+7	Figure 7 (hox region)	TTGTCAATGTCAACCGCAAC
3Chox_H07_pos_+8	Figure 7 (hox region)	TTATTGCCCAATCTGGCGAC
3Chox_H08_pos_+11	Figure 7 (hox region)	AGGCAAGACAGATAAACCTTCTG
3Chox_H09_pos_-2	Figure 7 (hox region)	GGGGCATAGGGAGCAAACC
3Chox_H10_pos_-4	Figure 7 (hox region)	GGGGTAGTACCGGAAGTGG
3Chox_H11_pos_-6	Figure 7 (hox region)	TTGGGCTGATTCTTGCAATTC
3Chox_H12_pos_-11	Figure 7 (hox region)	AGCTTTTGTCTAACACCGTATCG
3Chox_H13_pos_-13	Figure 7 (hox region)	AAGCCATCCTTCAAAACCCG
3Chox_H14_pos_-17	Figure 7 (hox region)	GCCCCATCGGTGAGAAAAAG
qPCR_hoxF_F	Figure 7 (hox internal control)	GTCATCTGCAATGCTGACGAA
qPCR_hoxF_R	Figure 7 (hox internal control)	CCAACACACTGCGGTCCAT
3CnifJ_Hcommon	Figure 7 (nifJ region)	ATTCTTTTCGCCAGTTTCTCCAC
3CnifJ_H01_pos-16	Figure 7 (nifJ region)	CATGCTGTTCTGTGCCCATC
3CnifJ_H02_pos-14	Figure 7 (nifJ region)	TCAGTGATATTTTGACCGCTGG
3CnifJ_H03_pos-13	Figure 7 (nifJ region)	AAAACCCCTGCCAACAAGAC
3CnifJ_H04_pos-12	Figure 7 (nifJ region)	CTCCCCACAATAAAGGGCTG
3CnifJ_H05_pos-10	Figure 7 (nifJ region)	TCATAGAAGCGGCGGATACC
3CnifJ_H06_pos-8	Figure 7 (nifJ region)	AGATAACGCCCCGGAACC
3CnifJ_H07_pos-7	Figure 7 (nifJ region)	GTATGGGTCGCCAAAGTACG
3CnifJ_H08_pos-6	Figure 7 (nifJ region)	AGGGATGGGGATCTTTGCTG
3CnifJ_H09_pos-4	Figure 7 (nifJ region)	TTATTGCTTTCACCCCGGC
3CnifJ_H10_pos-3	Figure 7 (nifJ region)	GCCCTAGGCCAAAGAATACC
3CnifJ_H11_pos-1	Figure 7 (nifJ region)	CCAGCCAGGGAGATGACC
3CnifJ_H12_pos+1	Figure 7 (nifJ region)	GCCATTGGGGGTTTATTGGC
3CnifJ_H13_pos+2	Figure 7 (nifJ region)	GTAAACAGATAGGTGGTGCGG

3CnifJ_H14_pos+6	Figure 7 (nifJ region)	TACAAAGCCAGGTAAGCGTC
3CnifJ_H16_pos+8	Figure 7 (nifJ region)	TCTGACCTATTGCCCTGACC
3CnifJ_H17_pos+9	Figure 7 (nifJ region)	CTGTTGTCATTGCTTGCTCC
3CnifJ_H18_pos+11	Figure 7 (nifJ region)	TGTAGCCAGTGAAGGGGAAG
3CnifJ_H19_pos+14	Figure 7 (nifJ region)	TCCCGGTATTAATGGCATTGG
3CnifJ_H20_pos+15	Figure 7 (nifJ region)	GTTGTGTGATTCCCCAAGCC
3CnifJ_H22_pos+17	Figure 7 (nifJ region)	CTTATTTTGTACCCGGCACCC