1 **Knockout of GMC-oxidoreductase genes reveals functional redundancy in mimivirus**

3   **Authors:** Jean-Marie Alempic[1]†, Hugo Bisio[1]†, Alejandro Villalta[1]†, Sébastien Santini[1],

4   Audrey Lartigue[1], Alain Schmitt[1], Claire Bugnot[1#], Anna Notaro[2], Lucid Belmudes[3], Annie

5   Adrait[3], Olivier Poirot[1], Denis Ptchelkine[4], Cristina De Castro[2], Yohann Couté[3], Chantal Ab-

6   ergel[1*]

Affiliations :

[1]Aix–Marseille University, Centre National de la Recherche Scientifique, Information Géno-

mique & Structurale, Unité Mixte de Recherche 7256 (Institut de Microbiologie de la Médi-

terranée, FR3479, IM2B, IOM), 13288 Marseille Cedex 9, France.

[2]Department of Agricultural Sciences, University of Naples Federico II, Via Università 100,

80055, Portici, Italy

[3]Univ. Grenoble Alpes, CEA, INSERM, UA13 BGE, CNRS, CEA, FR2048, 38000 Grenoble,

France

[4]Aix–Marseille University, Centre National de la Recherche Scientifique, Architecture et

Fonction des Macromolécules Biologiques, Unité Mixte de Recherche 7257 (IM2B), 13288

Marseille Cedex 9, France.

*Corresponding author. Email: Chantal.Abergel@igs.cnrs-mrs.fr

†These authors contributed equally to the work

Present address: Sorbonne Université, Centre National de la Recherche Scientifique, BIOM

UMR7232, Observatoire Océanologique, 1 avenue Pierre Fabre, 66650 Banyuls sur mer,

France.

One-Sentence Summary: Functional redundancy warrants mimivirus genomic fiber and fibrils

formation.

Subject Area: Biological sciences/Microbiology/Virology/Viral evolution; Biological

sciences/Microbiology/Virology/Virus structures

Keywords: Mimivirus, giant virus, dsDNA, viral genome organization, genomic fiber, cryo-EM, helical

reconstruction, MS-based proteomics.

Abstract

The Mimivirus 1.2Mb genome is organized into a nucleocapsid-like genomic fiber composed of two closely related paralogs of GMC-oxidoreductase (1). Surprisingly, these proteins also compose the layer of fibrils that decorate the virions and are likely to play a key role in the infection (2). Each of these genes was inactivated using an original homologous recombination-based knockout protocol, with the aim of studying their essentiality by reverse genetics. None of these genes was found to be essential and each mutant could form a genomic fiber but with different stabilities. The most stable one led to the same structures as the wild type (5- and 6-start helices) and these results allowed us to propose a model explaining the coexistence of the two different helices in the virions. Furthermore, all mutants, including the double mutant, showed equivalent fitness in laboratory conditions, and the resulting virions presented the same glycosylation as the wild type, whereas the protein composition in the fibrils of the double mutant was different. Within the family, there is variability in the glycan composition of the fibril layer between clades, and even intra-clade (3, 4), and this variability now extends to the proteins composing the fibrils. These results obtained on two distinct essential processes (genome packaging and virion infectivity) illustrate the exceptional functional redundancy of the *Mimiviridae*, which may be the major evolutionary force behind their giant genomes.

Significance statement

Exaptation and genetic redundancy are key factors in giant virus evolution. Exaptation is exemplified by the recruitment of metabolic enzymes, GMC oxidoreductases, for DNA compaction and virion attachment. Despite their involvement in such essential processes these proteins can be lost during experimental evolution experiments. Here, we developed genetic tools that will allow addressing the function of thousands of genes in *Mimiviridae*. They were used to directly characterize the essentially of GMC oxidoreductases for the genomic fiber and virion external layer of fibril assembly. As a result, this study unveils an extensive toolkit of viral genes that can alternatively be used to secure essential functions as well as the unprecedented plasticity of mimivirus for virion assembly.

Introduction

Mimivirus is the prototype of the *Mimiviridae* family part of the *Nucleocytoviricota* phylum encompassing large and giant DNA viruses infecting eukaryotes (5). Members of the *Mimiviridae* family infecting amoeba have dsDNA genomes up to 1.5 Mb encoding over

1000 proteins, including complete glycosylation machinery (3, 4, 6–9). Mimivirus virions penetrate the cell through phagocytosis, and the acidic vacuole mediates the opening of the stargate structure at one vertex of its icosahedral capsid (10, 11). The internal membrane unwraps and fuses with the vacuole membrane, allowing the transfer of the nucleoid compartment into the host cytoplasm, while empty capsids remain in the cell vacuole (6, 10, 12). The infectious cycle occurs in the cytoplasm where a large viral factory is developed (6, 12–14). At the late stage of the cycle, neo-synthesized virions bud at the periphery of the viral factory where they are filled by the genome. Finally, a glycosylated surface layer of fibrils, composed of two large capsular-like polysaccharides synthesized by the virally encoded machinery, is added to the capsids (4, 15, 16). As a result, the 750 nm diameter virions appear as Russian dolls made of the surface layer of reticulated glycosylated fibrils decorating the icosahedral capsids. Inside, the nucleoid compartment encases the 1.2 Mb dsDNA genome organized into a 30 nm large genomic fiber with a nucleocapsid-like structure, made of a protein shell internally lined by the folded DNA, and the central channel can accommodate large proteins such as the viral RNA polymerase (1). Three independent genomic fiber structures have been determined by cryo-electron microscopy (cryo-EM): two compact 5-start and 6-start DNA containing helices and a 5-start relaxed helix without DNA (1). Unexpectedly, the protein shell was found to be composed of two GMC-oxidoreductases sharing 69% sequence identity, and the resolution of the reconstructions (3.7 and 4.4 Å) prevented to determine whether each or both paralogs were composing each helix. Interestingly, the same GMC-oxidoreductases also compose the surface fibrils of the virions(4, 9, 17). However, despite their involvement in both the mimivirus genomic fiber structures and the virion surface layer, homologs of these proteins are puzzlingly absent in the laboratory evolved mutant (strain M4) (17), suggesting they may not be essential. The M4 strain also lacks the glycosylation machinery responsible for synthesizing the glycan composing the polysaccharides and branching them on the proteins composing the virion fibrils. In the present study, we developed tools to knock out (KO) mimiviral genes in order to investigate the impact of the depletion of the GMC-oxidoreductases for the formation of the genomic fiber and fibrils. We assessed the fitness of the different mutants compared to a wild-type mimivirus (strain reunion). Cryo-EM was used to determine the structure of the genomic fiber made by the less abundant GMC-oxidoreductase in the wt virus. Nuclear magnetic resonance (NMR) and Gas Chromatography Mass Spectrometry (GC-MS) were used to complement the study by comparing the glycan compositions and structures of each mutant with the wt. We finally used Mass spectrometry (MS)-based proteomics to analyze the protein

96 composition of the fibrils of members of the clade B and C which glycosylate their fibrils

97 with different glycans using clade-specific glycosylation machineries (3, 9). While confirming

98 the non-essentiality of the two GMC-oxidoreductases, our results document the unexpected

99 global resilience of Mimivirus replication to significant disruptions in both its genome

100 packaging and virion structure.

Results

None of the two GMC-oxidoreductases is essential

To achieve genetic manipulation of mimiviral genes, we adapted the genetic tools

based on homologous recombination in combination with selection cassettes previously

described for melbournevirus and pandoravirus (18, 19). In particular, we built plasmids that

introduce a selection cassette (either a nourseothricin N-acetyl transferase (NAT) or a

Neomycin resistance gene (NEO)) to disrupt the gene of interest, allowing to select

recombinant viruses when the gene is non-essential (Fig. 1A). The resistance cassettes were

expressed under the control of a megavirus promoter (Fig. 1A). Consistently, the addition of

nourseothricin or geneticin (the neomycin resistance gene also confers resistance to geneticin)

allows to enrich the recombinant viruses, which can be cloned to obtain homogeneous

populations (Fig. 1B). Using these tools, we generated KO of each of the two genes encoding

the GMC oxidoreductases (qu_946 and qu_143). Each mutant was easily produced and

genotyped to confirm the mutation (Fig. S1). Using the second selection cassette we were able

to KO both genes (Fig. S1) and, surprisingly, the resulting mutant was easy to produce

demonstrating that the two GMC-oxidoreductases were not essential. The absence of

additional mutations in every mutant was confirmed by genome sequencing.

Mutants' fitness

To assess whether a fitness cost was associated with the mutations, we performed

competition assays against wt mimivirus by measuring the abundance of each mutant over

several cycles in the presence and absence of selection. Compared to the wild type, each

mutant presents a common nourseothricin resistance gene, with, for the double KO (2KO), an

additional geneticin resistance gene. As a result, in the presence of nourseothricin, each

mutant ratio increased, with the disappearance of the wild-type virus after 5 passages (Fig.

1C). These data allowed competition assays to be validated as an effective tool to assess the

fitness of recombinant viruses. In the absence of selection, the relative abundance of the

mutants compared to the wild type remained around 0.5 over 5 passages, supporting the

absence of a fitness cost, even for the 2KO mutant (Fig. 1C).

Figure 1: Mimivirus reunion mutants' generation and phenotypic characterization. A]
Schematic representation of the vector and strategy utilized for knock out of qu_143
(KO_qu143) and qu_946 (KO_qu946) in mimivirus reunion. Selection cassette was
introduced by homologous recombination and recombinant viruses were generated as shown
in Fig. 1B. Primer annealing sites are also shown and the sequence of the primers is included
in Table S1. B] Cartoon depicting the strategy for the selection of recombinant viruses. Viral
infection was performed 1-hour post-transfection. Ntc: Nourseothricin. P= passage. C]
Growth competition assays revealed no significant defects in the lytic cycle of KO strains.
The competition was also performed in presence of Nourseothricin which allows the out
competition of the recombinant strains due to the expression of a Nourseothricin selection
cassette. Measurements were performed by qPCR of an endogenous locus (present in wt and
recombinant strains) and the Nourseothricin selection cassette (only present in recombinant
viruses).

Composition of the genomic fiber of single mutants

We managed to extract and purify the genomic fiber of every single mutant and

assessed their protein composition using MS-based proteomics. This confirmed the most

abundant protein was the remaining GMC-oxidoreductase and that proteins related to

transcription and oxidative stress were still present in the genomic fiber. Negative staining

microscopy highlighted surprising differences between the two structures in terms of stability

(Fig. 2). The two mimivirus reunion proteins qu_946 and qu_143 share 69% sequence

identity (81% similarity) and the genomic fiber made of qu_946 (KO_qu143), the most

abundant in the wild type, was mostly in the unwound state (Fig. 2B), while the one made of

qu_143 (KO_qu946), resulted in very long and stable helices (Fig. 2C). The use of the qu_143

protein in the wt genomic fiber could thus contribute to its stabilization (Fig. 2A). In the case

of the double mutant, the protocol for capsid opening to extract the genomic fiber of wt and
single mutants did not allow the clean opening of the capsids. An optimized protocol allowed
the extraction of a possible thinner genomic fiber, but in poor yield, preventing its purification
and compositional characterization (Fig. 2D).

158
Figure 2: Micrograph of negative stained genomic fiber of A] Wild-type mimivirus, B]
KO_qu143 mutant, C] KO_qu946 mutant, and D] 2KO double mutant. Scale bars 100 nm.

Cryo-EM single particle analysis of the qu_143 genomic fiber

We performed cryo-EM single-particle analysis of the most stable genomic fiber made
of qu_143. The 2D classification (using Relion (20, 21)) confirmed the heterogeneity of the
sample, and 2D classes were sorted by applying our already described clustering protocol (1).
The two main clusters corresponding in width to the compact (Cl1a) and (Cl2) structures of
the wt genomic fiber (Fig. 3) were respectively named Cl1 and Cl2.

Figure 3: **Clustering analysis of the 2D classes obtained with qu_143 genomic fiber:**
Automatic sorting of the 2D classes using the fiber width W1 and pairwise correlations of the
2D classes resulting in 2 main clusters (5-start Cl1 cyan; 6-start Cl2 in orange).

For each, we confirmed the helical symmetry parameters were the same as the wt

genomic fiber by image power spectra analyses and performed structure determination and

refinement (Fig. 4). A less populated thinner cluster (Cl0, dark blue, ~25 nm), absent from the

wt genomic fiber 2D-classes, was also identified but, due to the lower number of segments

and the resulting lower resolution power spectra compared to Cl1 and Cl2, we failed to

identify helical parameters. For Cl1, after 3D-refinement, we obtained a 4.3 Å resolution

helical structure (FSC threshold 0.5, masked) corresponding to the same 5-start left-handed

helix as the wt, made of a ~8 nm-thick proteinaceous external shell with 5 dsDNA strands

lining the interior of the shell, leaving a ~9 nm wide central channel (Fig. 4). The 4.2 Å

resolution Cl2 map obtained after 3D refinement (Fig. 4) corresponds to the same 6-start left-

handed helix as the wt, made of the ~8 nm-thick external shell, with 6 dsDNA strands lining

it, leaving a ~12 nm wide inner channel (Fig.4). The Fourier shell correlation (FSC) curves

for the final 3D reconstructions of the helical assemblies are presented in Fig. S2.

184
Figure 4: **Workflow of the 5- and 6-start helices reconstruction processes.** Segment
extraction was performed with a box size of 400 pixels (pix) binned (box size 100 pix, 4.3436
Å/pix). The distance between consecutive boxes was equal to the axial rise calculated by
indexation of the power spectrum. After clustering, 2D classes were selected for Cl1 and Cl2
and 3D classification was carried out using the selected segments, helical symmetry
parameters from the power spectrum indexation, and a 300 Å or 330 Å featureless cylinder as
3D reference for Cl1 and Cl2, respectively. 3D-refinement of the 2 boxed 3D classes was
achieved using one low pass filtered 3D class as reference on the unbinned segments. A first
3D-Refinement was performed with solvent flattening followed by CTF refinement and
polishing of the selected segments. A last 3D refinement was achieved with solvent flattening.
The EM maps colored by local resolution from 5 Å (blue) to 3 Å (red) with Euler angle
distribution of the particles used for the 3D reconstruction are presented.

We used the qu_143 dimeric structure refined in the Cl1a focus refined map of the wt

genomic fiber structure for refinement into the mutant Cl1 and Cl2 maps. Initial models were

further refined using the real-space refinement program in PHENIX 1.18.2 (22) (Table S3).

The density that can be attributed to the FAD cofactor is present in both maps (Fig. S3) and

the models of Cl1 and Cl2 dimers are superimposable with a core RMSD of 0.436 Å based on

Cα atoms, comparable with the ones obtained with the wt Cl1a and Cl2 dimers (Table S4).

The relaxed Cl3 cluster in the wt was never observed with the mutant. As a result, the qu_143

helices appear more stable than the wt, composed of both GMC-oxidoreductases, themselves

being more stable than the qu_946-only structures, mostly found in an unfolded state.

Model explaining the co-existence of 5- and 6-start helices

In the KO_qu946 genomic fiber, despite the presence of a single GMC-oxidoreductase

to build the shell, there is a mixture of 5 and 6 strands of DNA in the genomic fiber as for the

wt genomic fiber. We estimated the ratio of 5-start and 6-start from the clustering (Fig. 3) and

propose a model that reconciles the co-occurrence of the two structures. In this model, the

whole genome would be folded into 6 parallel strands, 5 longer than the last one. The helix

would then be formed initially as a 6-start helix until the sixth strand ends and, from that point

becomes, a 5-start helix (Fig. 5). According to this model, assuming the length of the genomic

fiber is limited by the size of mimivirus nucleoid compartment, we can estimate that the

maximum genome length would be ~1.4 Mb for a full 6-start helix with an 8 nm thick protein

shell. The last cluster (Cl0, Fig. 3) could correspond to a 4-start with an additional shorter

DNA strand. Proteins, such as the chromosome condensation factor identified in the proteome

of the genomic fiber (1), could seed the folding of the DNA into 6 strands before assembly in

the helical structure.

Figure 5: Model explaining the transition from a 6- to a 5-start helix. The transition from
a 6- to a 5-starts involves a decrease of the helix diameter by ~3.5 nm. The small cluster
around 250 nm could thus correspond to a 4-start helix.

The N-terminal cys-pro-rich domain is an addressing signal to the virion surface

The cys-pro-rich N-ter domain of the GMC-oxidoreductases is absent in the genomic

fiber but present in the fibrils (1). To assess whether this cys-pro-rich N-ter domain could be

an addressing signal to the fibrils, we replaced the second GMC-oxidoreductase (qu_143) in

the genome of the mutant KO_qu946 by inserting the sequence of the GFP and a gentamicin

selection cassette, in fusion with the sequence of the qu_143 N-terminal domain (Nqu143-

GFP). We then analyzed the resulting mutant by MS-based proteomics (Tables 1 and S2) and

observed the virions by fluorescence microscopy. Purified Nqu143-GFP virions, but not

virions produced in cells expressing GFP, showed a strong fluorescence indicating the

incorporation of the chimeric protein into the viral particles (Fig. 6). Moreover, Nqu143-GFP

was identified in purified virions (ranked 53[rd] in terms of relative abundance, Table S2) and

was found enriched in the fraction containing the fibrils (ranked 11[th], Tables 1 and S2), with

peptides covering the N-terminal domain (Fig. S5). Taken together, these data indicate the N-

terminal cys-pro-rich domain of the GMC oxidoreductase is necessary and sufficient to direct

the proteins to the fibrils of mimivirus particles.

Figure 6: Quantification of fluorescent virion in infected cells overexpressing GFP (Vc2)
and in cells infected by the mutant Nqu143-GFP. Scale bar 5 μm. (CTVF: corrected total
virion fluorescence).

Protein composition of the fibrils for members of the clades A, B and C

While the cys-pro-rich N-ter domain is conserved in all members of clade A, it is

absent in the orthologous proteins in members of clade B and C, and the two GMC-

oxidoreductases are pseudogenized in tupanviruses, suggesting another protein was

composing their fibrils (Fig. S6). The proteomic analysis of enriched fibrils of the double KO

mutant identified a group of proteins as the most abundant in the fibrils fraction (qu_734,

qu_757, qu_384 and qu_482, Tables 1 and S2). These proteins were ranked 14[th], 59[th], 13[th] and

15th in the entire capsids and some were also highly ranked in the Nqu143-GFP mutant.

Interestingly, the orthologous of qu_734 is also the most abundant in the M4 natural mutant

fibrils (696-L688, Tables 1 and S2). Since this mutant does not encode a glycosylation

machinery, it is not glycosylated and its capsid lacks the large reticulated crown of fibrils

decorating mimivirus capsid (4, 17). The qu_734 protein also possesses a cys-pro-rich N-ter

domain which is conserved in all clades (Fig. S6). It was not possible to determine if one of

these proteins was also the building block composing the 2KO genomic fiber, given the

difficulty to open its capsids. We concluded that the change in surface protein composition

provided different properties to the capsids compared to wt and single mutants and

optimization of capsid opening will be necessary to recover enough genomic fiber to be able

to purify it and characterize its composition.

The proteomic analysis of fibrils purified from B clade members, moumouvirus

australiensis and maliensis, show similar composition for the most abundant proteins, but with

slight differences in their relative abundances (Tables 1 and S2). The moumouvirus maliensis

protein mm_751, ranked 1st in its fibrils, is absent from clade A and in the top 3 in clade B

and C. There are cys and pro in the N-ter domain of mm_751, but not as rich as in the GMC-

oxidoreductases. For ma_195 (qu_738 in mimivirus reunion, Tables 1 and S2), ranked 1st in

the fibrils of moumouvirus australiensis, the N-ter domain is not cys-pro-rich. The same

proteins are also the best ranked in the fibrils of members of clade C. The first ranked in

megavirus chilensis is mg749 (qu_657 in mimivirus reunion) which presents a cys-pro-rich

N-ter domain, and the first ranked protein in megavirus vitis, mvi_646 (qu_600 in mimivirus

reunion), is lacking a cys-pro-rich N-ter domain. Yet, for each virus in each clade, there is at

least one protein with a cys-pro-rich N-terminal domain among the most abundant in the

fibrils.

Table 1: Rank of the most abundant proteins in the fibrils of members of clade A, B and C

Rank	Mimvirus reunion V	Mimvirus reunion F	2KO V	2KO F	2KO-GFP V	2KO-GFP F	M4 V	M4 F	Megavirus chilensis V	Megavirus chilensis F	Megavirus Vitis V	Megavirus vitis F	Moumouvirus Australiensis V	Moumouvirus ausraliensis F	Moumouvirus maliensis V	Moumouvirus maliensis fibrils
qu_143	6	1	NA	NA	NA	NA	NA	NA	182	73	177	46	91	45	225	130
qu_757	54	2	59	2	26	9	89	7	134	44	164	24	47	7	115	19
qu_734	14	3	14	1	3	5	20	1	10	7	12	14	10	14	10	16
qu_482	16	4	15	4	194	1	8	54	23	42	32	90	43	9	11	42
qu_880	29	5	33	11	64	18	NA	NA	21	6	26	18	82	18	54	24
qu_384	15	6	13	3	29	2	16	2	27	9	22	4	18	13	23	18
qu_773	7	14	9	10	75	6	8	3	5	8	3	27	3	5	3	8
qu_657	135	15	84	27	12	4	30	27	7	1	8	7	15	4	4	2
qu_600	17	16	18	7	5	9	23	4	35	10	34	1	13	3	15	4
qu_738	22	17	24	12	4	15	17	5	11	3	16	2	7	1	8	3
qu_752	41	35	16	55	86	3	2	21	NA	NA	NA	NA	NA	NA	NA	NA
qu_582	12	39	5	43	1	103	3	28	4	26	6	55	29	33	12	39
mm_751	NA	NA	NA	NA	NA	NA	NA	NA	14	2	25	3	6	2	5	1
Nqu143-GFP	NA	NA	NA	NA	53	11	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

V: Virion; F: Fibrils (grey); NA: Not applicable, the corresponding gene is absent in the corresponding genome.

Sequences of the relevant orthologues of mimivirus reunion in other viruses are provided in Fig. S6.

Glycan composition of the mimivirus reunion mutants' fibrils

We recently established that the fibrils were coated with glycans (3, 9) and

demonstrated that for mimivirus, the prototype of the A clade, they were made of two distinct

polysaccharides (4). Since the cluster of 12 genes responsible for the biosynthesis of the

polysaccharides is preserved in the A clade (3), we suspected that the fibrils of the mimivirus

reunion isolate had the same composition as mimivirus prototype. To assess whether the

knockout of the most abundant protein composing the fibrils could also affect the branching

of these polysaccharides and their composition, we analyzed the sugar content of the viral

particles of the three mutants (KO_qu946, KO_qu143, and 2KO) and wt mimivirus reunion,

as reported in (3, 4). The chemical characterization revealed for each of them the presence of

sugars (Figure 7A), with rhamnose (Rha), viosamine (Vio), 2-OMe-Vio, glucose (Glc), and

glucosamine (GlcN) (Fig. 7A), confirming that mimivirus reunion and all mutants had the

same glycan composition as mimivirus prototype (3). We then analyzed the fibrils of the

mutants and wt by ^{1}H NMR spectroscopy (23) and compared them with mimivirus prototype.

As a result, in the ^{1}H NMR spectra, the anomeric region (23) (5.5-4.4 ppm) perfectly

overlapped with that of the previously studied mimivirus strain (4) (Fig. 7B, Fig. S7),

confirming that the different sugars were assembled the same way as in the reference

mimivirus to produce the same two polysaccharides (Fig. 7C). Indeed, the signal at 4.80 ppm

is the result of the overlap of two anomeric protons, the one of a 3-α-Rha (labeled **C**) and of a

3-β-GlcNAc (**D**) modified with a pyruvic acid at the hydroxyl functions 4 and 6. These two

residues are the building blocks of the repeating unit of polysaccharide 1 (**poly_1**, **Fig. 7C**)

(4). The other anomeric signals, labeled with the capital letters **A, A'** (2,3-α- L-Rha), **B** (3-β-

D-GlcNAc), **E** (2OMe-β-D-VioNAc) and **E'**(β-D-Vio4NAc), are part of the polysaccharide 2

(**poly_2**) that presents a backbone with a disaccharide repeating unit of 2)-α-L-Rha-(1→3)-β-

D-GlcNAc-(1→, with the rhamnose residue further substituted with a viosamine methylated

at position 2 (**E**) or not methylated (**E'unit**), thus taking the labels **A** and **A'**, respectively

(Fig. 7C) (4).

304
Figure 7: Compositional and NMR analysis of the fibrils of mimivirus reunion wt and
mutants A] GC-MS chromatogram profiles of the sugars composing the fibrils of wt (a),
KO_qu946 (b), KO_qu143 (c), and 2KO (d). B] Comparison of the ^{1}H NMR spectra of
mimivirus reunion wt and related mutants with that of mimivirus reference strain. The
anomeric signals related to poly_1 (**C** and **D** units, in black), and those of poly_2 are in red.
C] Structures of mimivirus polysaccharides as reported in Notaro *et al*, 2021 (4).

It was hypothesized that the mimivirus GMC oxidoreductases were the major target
for glycosylation at the fibrils of mimivirus (4, 17). The results presented here indicated that
the mutations, including the deletion of the two GMC-oxidoreductase genes, did not affect the
glycosylation of the fibrils. These data suggest either the presence of multiple proteins being

targeted for glycosylation or that the glycosylation target is not a GMC oxidoreductase.
Alternatively, it is also possible that the glycosylation machinery was able to use the qu_734
protein as support for the two polysaccharides. No glycans were identified in the M4 natural
mutant fibrils.

Discussion

Genome packaging is essential for the propagation of viruses. For instance, the
packaging ATPase of poxviruses (24) or the histones of marseilleviruses (19, 25) are essential
for the productive infection by their virions. We have previously demonstrated that the
genomic fiber of mimivirus is composed of two GMC oxidoreductases (1), but in contrast to
previous examples, the absence of both GMC oxidoreductases in the M4 laboratory-generated
strain suggested a non-essential function for these genes (17). Implementation of the genetics
for exclusively cytoplasmic giant viruses, based on homologous recombination, allowed us to
assess the essentiality of GMC-oxidoreductases in mimivirus reunion for the assembly of the
genomic fiber. We have previously proposed an increased genome redundancy as a
contributing factor to the appearance of viral giantism (18). The data presented here validate
such premise and extend these predictions outside of the *Pandoraviridae*. Indeed,
unsurprisingly, given the homology between the two enzymes, the KO of the most abundant
one in the genomic fiber induced its replacement by the second one, with no apparent cost for
the virus, at least in laboratory conditions. Regardless, despite the exchangeable nature of
these proteins, the appearance and physical features of the genomic fiber were significantly
different between recombinant viruses and the wt, indicating a certain degree of functional
specialization. The KO_qu946 genomic fiber presents two helices, a 5- and a 6-start helix,
with the same helical parameters as the wt, which led us to propose a model to explain the co-
existence of both structures. But, while the wt genomic fiber was highly heterogeneous, with
fully relaxed helices having lost DNA and even helices that lost a strand, the mutant appeared
more stable with no occurrence of fully relaxed structures. Yet, despite our efforts to improve
the resolution, the helices appeared less defined compared to the wild type, which may be due
to internal disorders or to the cryo-sample preparation. Moreover, the deletion of qu_143, the
less abundant GMC-oxidoreductase in the wt, seemed to increase the genomic fiber
instability, suggesting a stabilizing role for this second GMC-oxidoreductase in the wt virus.
Finally, the deletion of both GMC-oxidoreductases revealed they were not essential, with no
fitness cost associated with their loss.

Importantly, GMC oxidoreductases are not only present in the genomic fibers but also

in the virion surface fibril layer and have been previously predicted as targets for

glycosylation (4, 17). The only difference between the genomic fiber and the surface fibril

protein compositions being that the cys-pro-rich N-ter domain of the GMC-oxidoreductases

are only present in the fibrils and absent in the genomic fiber. Interestingly, when fused to

GFP, the cys-pro-rich N-ter domain of qu_143 is sufficient to produce fluorescent virions,

suggesting it is an addressing signal to the fibrils. Unexpectedly, the virions of all three

mutants were still decorated by long, reticulated fibrils. The MS-based proteomic analysis of

the 2KO fibrils revealed they were composed by other proteins which were also enriched in

the M4 natural mutant fibrils, some, including qu_734, also presenting a cys-pro-rich N-ter

domain (Fig. S6). The glycan analysis of the GMC-oxidoreductases single and double

mutants confirmed that their fibrils were still glycosylated by the same two polysaccharides as

the wt virions. As previously reported, M4 was not glycosylated as it lacks a virally encoded

glycosylation machinery (3, 4, 17). Moreover, the resulting change in the composition of the

surface fibril layer in the 2KO virions seemed to impact the capsid sensitivity to the

proteolytic treatment we originally used to recover wt and single mutants genomic fibers and

we were not able to recover the 2KO genomic fiber to purify it in quantities compatible with

structural and compositional analysis. Yet, a thinner genomic fiber was observed (~20-25nm,

Fig. 2D).

The analysis of the purified fibrils of members of clades B and C confirmed this

versatility, with fibrils composed of different proteins inter- and even intra-clade. Clades B

and C viruses present fibrils with a protein composition closer to each other than that of clade

A viruses. In a previous study, we reported that the glycosylation machinery was clade-

specific and this is now the overall composition of the fibrils that appears to vary. The

glycosylated fibrils bear capsular-like glycans echoing the bacterial ones, recognized by the

amoeba which feeds on bacteria (2, 3). They thus appear as key for infection, by fooling the

amoeba which mistake the virions for bacteria due to their size and "taste". In a given

environment, the composition of the fibrils could correspond to a variable, favoring the

engulfment of a given virion by a given amoeba compared to others virions and even bacteria.

The protein content of the fibrils also appears to have been optimized in a given clade but

their complex composition suggests there is a large set of proteins that can make the fibrils.

Having a flexible toolbox for glycan synthesis and fibril composition would reflect the need

to continuously adjust the capsids composition to outcompete other parasites and secure
infection. We can thus hypothesize that in the population of virions resulting from a single
infection, this composition could also be variable, providing to the virus with a toolbox to
ensure the productive infection of at least one of the potential hosts present in that
environment. The question of the use of the same protein to make the fibrils and the genomic
fiber remains unanswered given the difficulties we encountered in optimizing the capsid
opening protocol to recover the 2KO genomic fiber in quantities compatible with its
purification and analysis. Yet, a thinner structure appears (~20-25 nm). Given that the central
part of the genomic fiber has to corresponds to a central channel of at least 9 nm with 4nm of
DNA following the helical symmetry and thus, this is the protein making the shell that defines
the final dimension of the genomic fiber. The GMC-oxidoreductases being 8 nm that leads to
a ~29 nm helix. The most abundant protein composing the fibril of M4 and the 2KO
corresponds to the 222 amino-acids qu_734 protein, which according to the high confidence
alphaFold (26) prediction could be almost twice smaller (~4.5 nm) and if this same protein
makes the shell of the mutants, we expect a genomic fiber of 22 nm. Overall, our results
reveal the resilience of mimivirus, with redundant solutions securing essential functions, such
as infectiousness and genome packaging. Functional redundancy, well documented in the
cellular world as a way to preserve essential function such as cell division (27) can now be
expanded to the viral world.

Material and methods

Cloning of DNA constructs

Gene knock-out vectors. The plasmid for gene knock-out was generated by sequential cloning
of the 3' UTR of mg_18 (megavirus chiliensis), the promoter of mg_741 (megavirus
chiliensis), and a Nourseothricin N-acetyl transferase (NAT) or a neomycin selection cassette
(NEO). Each cloning step was performed using the Phusion Taq polymerase (ThermoFisher)
and InFusion (Takara). Finally, 500bp homology arms were introduced at the 5' and 3' end of
the cassette to induce homologous recombination with the viral DNA. Before transfection,
plasmids were digested with EcoRI and NotI. All primers are shown in Table S1.

Establishment of viral lines.

Gene knock-out. A detailed method for gene manipulation is provided in (Manuscript
accepted, Nature Protocols). Gene knockout strategy was performed as previously described

for pandoravirus (18). Briefly, 1.5×10^5 *Acanthamoeba castellanii cells* were transfected with

6 µg of linearized plasmid using Polyfect (QIAGEN) in phosphate saline buffer (PBS). One

hour after transfection, PBS was replaced with PPYG, and cells were infected with 1.5×10^7

mimivirus reunion particles for 1 hour with sequential washes to remove extracellular virions.

24h after infection the new generation of viruses (P0) was collected and used to infect new

cells. An aliquot of P0 viruses was utilized for genotyping to confirm the integration of the

selection cassette. Primers used for genotyping are shown in Table S1. A new infection was

allowed to proceed for 1 hour, then washed to remove extracellular virions and nourseothricin

was added to the media. Viral growth was allowed to proceed for 24 hours. This procedure

was repeated one more time before removing the nourseothricin selection to allow viruses to

expand more rapidly. Once, the viral infection was visible, the selection procedure was

repeated one more time. Viruses produced after this new round of selection were used for

genotyping and cloning. Double knockout of the GMC oxidoreductases was obtaining by

using a clonal population of qu_143 knockout viruses as parental strain. The locus of qu_946

was replaced by a neomycin resistance cassette. The transfection and selection of recombinant

viruses' procedure was performed identical to the process to generate single knockout but

replacing nourseothricin by geneticin.

Cloning and genotyping. 150,000 *A. castellanii* cells were seeded on 6 well plates with 2mL

of PPYG. After adhesion, viruses were added to the well at a multiplicity of infection (MOI)

of 1. One-hour post-infection, the well was washed 5 times with 1mL of PPYG, and cells

were recovered by well scraping. Amoebas were then diluted until obtaining a suspension of 1

amoeba/µL. 1µL of such suspension was added in each well of a 96-well plate containing

1000 uninfected *A. castellanii* cells and 200µL of PPYG. Wells were later monitored for cell

death and 100µL collected for genotyping. Genotyping was performed using Terra PCR

Direct Polymerase Mix (Takara) following the manufacturer's specifications. Primers used for

genotyping are shown in Table S1.

Competition assay and quantitative PCR analysis.

Equal infectious particles of wild-type and recombinant mimivirus reunion were mixed and

used to infect *A. castellanii* at an approximate multiplicity of infection of 0.8. Viruses were

allowed to grow overnight in the presence or absence of nourseothricin. Subsequent viral

progenies were used to infect new *A. castellanii* cells in reiterative passages. A fraction of

each passage was collected for genomic DNA extraction.

Viral genomes were purified using Wizard genomic DNA purification kit (PROMEGA). To

determine the amplification kinetic, the fluorescence of the EvaGreen dye incorporated into

the PCR product was measured at the end of each cycle using SoFast EvaGreen Supermix 2×

kit (Bio-Rad, France). A standard curve using the gDNA of purified viruses was performed in

parallel with each experiment. For each point, a technical triplicate was performed.

Quantitative real-time PCR (qRT-PCR) analyses were performed on a CFX96 Real-Time

System (Bio-Rad).

Genome sequencing and assembly of mutants' genomes

Genomic DNA was extracted from 10^{10} virus particles using the PureLink TM Genomic DNA

mini kit (Invitrogen) according to the manufacturer's protocol. Clones of individual mutants

and wt were sequences on an illumine platform (Novogen). For mimivirus wt, we obtained

4,819,885 150nt paired-ended reads. For KO_qu946, KO_qu946 and 2KO, 4,662,744,

4,572,873 and 5,068,030 150nt paired-ended reads, respectively. Genomes were assembled

using spades v3.13, with the option "careful" and we obtained 5 contigs for the wt, 4 for

KO_qu946, and 7 for KO_qu143 and 2KO. The wt sequence was consistent with the official

version of mimivirus reunion (GI MW004169) and the mutants only differed in the mutated

regions (in qu_946 and/or qu_143 genes). For the clone KO_qu946 the gene is interrupted

from position 1151358 to 1152985 relative to wt genome (position 302 to 1741 in the

corresponding qu_946 gene) and for the clone KO_qu143 the gene is interrupted from

position 165934 to 167572 relative to wt genome (position 306 to 1741 in the corresponding

qu_143 gene). Th region is replaced in both cases by the nourseothricin cassette construction

(1637 nt). For the 2KO, the gene qu_143 is interrupted from the positions 306 to 1741

(165934 to 167572 on the wt genome) and replaced by the nourseothricin cassette

construction (1637 nt), and gene qu_946 is interrupted from the positions 302 to 1206

(1151551 to 1153410 on the genome) and replaced by the geneticin cassette construction

(1859 nt). To confirm the KO of each mutant, the reads were mapped on the wt genome

resulting in a regular coverage along the genome, excepted for the qu_143 and qu_946 central

positions thus confirming the mutations. In addition, we used the central part of the KO genes

(deleted in mutants) as blast queries against the reads of each mutant genome, which also

confirmed the absent of the central portion.

Extraction and purification of the qu_946 and qu_143 mutants' genomic fiber

The genomic fiber of the mimivirus reunion single mutants of qu_946 (KO_qu946) and
qu_143 (KO_qu143), were extracted as described in Villalta et al, 2022 for the wild-type
virus (1). The genomic fiber was extracted from 12 mL of purified single KO mutant virions
at 2 x 10^{10} particles/mL, split into 12x1 mL samples processed in parallel. Trypsin (Sigma
T8003) in 40 mM Tris-HCl pH 7.5 buffer was added at a final concentration of 50 µg/mL and
the virus-enzyme mix was incubated for 2h at 30°C in a heating dry block (Grant Bio PCH-1).
DTT was then added at a final concentration of 10 mM and incubated at 30°C for 16h.
Finally, 0.001% Triton X-100 was added to the mix and incubated for 4h at 30°C. Each tube
was centrifuged at 4,000 x g for 5 min to pellet the opened capsids. The supernatant was
recovered, and the fibers were concentrated by centrifugation at 15,000 x g for 4h at 4°C.
Most of the supernatant was discarded leaving 12x~200 µL of concentrated fibers that were
pooled and layered on top of ultracentrifuge tubes of 4 mL (polypropylene centrifuge tubes,
Beckman Coulter) containing a discontinuous cesium chloride gradient (1.4, 1.3, 1.2 g/cm^3 in
40 mM Tris-HCl pH 7.5 buffer). The gradients were centrifuged at 200,000 x g for 16h at 4
°C. Since no visible band was observed, successive 0.5 mL fractions were recovered from the
bottom of the tube. Each fraction was dialyzed using 20 kDa Slide-A-Lyzers (ThermoFisher)
against 40 mM Tris-HCl pH 7.5 to remove the CsCl. These fractions were further
concentrated by centrifugation at 15,000 x g, at 4°C for 4h, and most of the supernatant was
removed, leaving ~100 µL of sample at the bottom of each tube. At each step of the extraction
procedure, the sample was imaged by negative staining transmission electron microscopy
(TEM) to assess the integrity of the genomic fiber. Each fraction of the gradient was finally
controlled by negative staining TEM.

Negative stain TEM

300 mesh ultra-thin carbon-coated copper grids (Electron Microscopy Sciences, EMS) were
prepared for negative staining by adsorbing 4-7 µL of the sample for 3 min., followed by
blotting excess liquid and staining for 2 min in 2% uranyl acetate to image the genomic fiber.
For fibrils, staining is performed with a drop of 1% uranyl followed by blotting after 10-15 s,
and a drop of uranyl acetate coupled with methylcellulose (0.2% and 2%, respectively) is
added twice and left for 5-10 s before blotting.

The grids were imaged either on an FEI Tecnai G2 microscope operated at 200 keV and
equipped with an Olympus Veleta 2k camera (IBDM microscopy platform, Marseille,

France); an FEI Tecnai G2 microscope operated at 200 keV and equipped with a Gatan
OneView camera (IMM, microscopy platform, France).

Single-particle analysis by cryo-EM

Sample preparation

508 For the KO_qu946 mutant, 3 µL of the purified sample were applied to glow-discharged
509 Quantifoil R 2/1 Cu grids, blotted for 2 s using a Vitrobot Mk IV (Thermo Scientific) and
510 applying the following parameters: 4°C, 100% humidity, blotting force 0, and plunge frozen
511 in liquid ethane/propane cooled to liquid nitrogen temperature.

Data acquisition

Grids were imaged using a Titan Krios (Thermo Scientific) microscope operated at 300 keV
and equipped with a K3 direct electron detector and a GIF BioQuantum energy filter
(Gatan). 2,224 movie frames were collected using the EPU software (Thermo Scientific) at a
nominal magnification of 81,000x with a pixel size of 1.0859 Å and a defocus range of -0.6
to -2.8 µm. Micrographs were acquired using EPU (Thermo Scientific) with 2.3 s exposure
time, fractionated into 40 frames, and 18.25 e$^-$/pixel/s (total fluence of 35.597 e$^-$
/Å²).

2D classification and clustering of 2D classes

All movie frames were aligned using MotionCor2 (28) and used for contrast transfer function
(CTF) estimation with CTFFIND-4.1 (29). Helical segments of the purified genomic fiber
manually picked with Relion 3.1.0 , were extracted with 400 pixels box sizes (decimated to
100 pixels) using a rise of 7.93 Å and a tube diameter of 300 Å. Particles were subjected to
reference-free 2D classification in Relion 3.1.0 (20, 21). We then performed additional cluster
analysis of the initial 2D classes provided by Relion to aim for more homogeneous clusters
(1), eventually corresponding to different states (Fig. 4).

Identification of candidate helical parameters

529 Fourier transform analysis methods have been used to confirm the helical parameters were the
530 same as in wt (1, 30–32) for the Cl1 and Cl2.

Cryo-EM data processing and 3D reconstruction

After helical parameters determination, segments were extracted with a box size of 400 pixels

(decimated to 100 pixels) using the proper rises for the Cl1 (7.93 Å, cylinder 300 Å, 442,237

segments) and the Cl2 (20.47 Å, cylinder 330 Å, 172,431 segments). A dedicated 2D

classification protocol was performed independently on each extraction. For the Cl2, one

round of 50 expectation-maximization (E-M) iterations was sufficient to produce 133

homogeneous 2D classes submitted to cluster analysis of which 44 2D classes were selected

(21,186 segments). For the Cl1 extraction, three iterative 2D classification/selection rounds

were performed (25, 50 and 100 E-M iterations) producing 61 classes from which 30

(149,593 segments) were finally selected on the basis of the cluster analysis.

Values of the helical parameters, rise and twist (Cl1: 7.9475 Å, 138.921°; Cl2: 20.48 Å,

49.43°), were then used for Relion 3D classification (20, 21), with a +/-0.5 units freedom

search range, using a featureless cylinder as initial reference (diameter of 300 Å and C1

symmetry for Cl1 and 330 Å and C3 symmetry for Cl2). In real space, the helical symmetry

was searched using 50% of the central part of the box for both structures. The number of

asymmetrical units in each segment box was set to 1 for Cl1 and 6 for Cl2. The entire helical

reconstruction and protein shell dimensions were obtained using in house developed program.

The superimposable 3D classes (same helical parameters, same helix orientation) were then

selected, reducing the data set to 98,882 segments for the 5-start fiber (Cl1) and 20,899

segments for the 6-start fiber (Cl2). After re-extraction of the selected segments without

scaling, a first unmasked 3D refinement was performed with a rescaled 3D classification

output low pass filtered to 15 Å as reference, followed by a 3D refinement step using solvent

flattened, FSCs and CTF refinement using the standard procedure described in Relion. To

further improve resolution of the maps, a Bayesian polishing was applied using 10,000

segments for training and default values for polishing. A last round of 3D refinement with

solvent flattening was applied to the previous refined map using the polished particles. At

that point, the maps were resolved enough (Cl1: 4.3 Å, Cl2: 4.2; FSC threshold 0.5) to

identify secondary structure elements (with visible cylinders corresponding to the helices) and

were used to compute local resolution.

Structures refinement

The best resolution Cl2 map was used to fit the qu_143 dimeric structure (PDB 7YX3) using

UCSF ChimeraX 1.5 (33). Each monomer was rigid-body fitted independently into the map

for a best result. The entire protein was then inspected within coot 0.9.7 (34) to fix local

inconsistencies and was further refined against the map using the real-space refinement
program in PHENIX 1.20.1 (22). The protein was submitted to 5 cycles of rigid body
refinement (with each chain defined) followed by twice 10 cycles of refinement with the
default PHENIX options. The resulting structure was manually corrected within coot. The
resulting protein model was submitted to the same steps refinement in PHENIX. This final
model was then fitted into the Cl1 map, inspected with coot and refined using 5 cycles of rigid
body refinement and simulated annealing followed by twice 10 cycles of refinement with the
default PHENIX options. Validations were performed into PHENIX using the comprehensive
validation program (table S3). RMSD between different structures (monomers and dimers,
Table S4) were computed using the align procedure in Pymol suite (Schrödinger, L., &
DeLano, W. (2020). *PyMOL*).

Extraction and purification of the mutants' fibrils

Defibrillation was performed by incubating purified virions (1.5 x 10^{10}) in Eppendorf tubes in
500 µL 40 mM Tris-HCl pH 7.5 buffer, 500 mM DTT for 2h at 30°C. Tubes are then
centrifuged at 14,000 g for 10 min. The supernatants containing the fibrils were recovered and
concentrated on vivaspin® 3 KDa (Sartorius, VS04T91) at 3,000 g. The pellet is washed
twice with 40 mM Tris-HCl pH 7.5 buffer and centrifuged at 14,000 g for 10 min and finally
resuspended in the same buffer. Both pellet and fibrils were imaged by negative staining
TEM.

For the 3 mutants and wt mimivirus reunion, 4 x 10^{11} viral particles were centrifuged at
14,000g for 10 min, the supernatant was discarded and the pellet was re-suspended in 10 ml
of 50 mM DTT. Fibril extraction was performed at 100°C under stirring. After 2 h, the tube
was centrifuged at 14,000 g for 15 min, at 4°C, and the fibrils recovered with the supernatant.
The fibrils were then dried and purified on Biogel P10, followed by subsequent NMR analysis
of each mutant and wt.

Sugar composition of viral particles of Mimivirus reunion wt and mutants

Monosaccharide composition analysis as acetylated methyl glycoside (AMG) was performed
on the intact viral particles (1,25 x 10^{10}, ~250 µl) of mimivirus reunion wt and mutants,
following the procedure reported by De Castro *et al* 2010 (35). The obtained AMG were
analyzed via gas chromatography-mass spectrometry (GC-MS) on an Agilent instrument (GC
instrument Agilent 6850 coupled to MS Agilent 5973) equipped with a SPB-5 capillary

column (Supelco, 30 m × 0.25 i.d., flow rate, 0.8 mL min–1) and He as the carrier gas. The

identification of the monosaccharides derivatized as AMG, was obtained by studying the

fragmentation pattern corresponding to each peak of the chromatogram and by comparison

with suitable standards.

Purification and ^{1}H NMR analysis of the fibrils

The fibrils of mimivirus reunion wt and mutants, extracted as reported above, were purified to

remove the excess of DTT used for the extraction procedure.

Briefly, the glycoproteins (protein/s carrying the polysaccharides) were precipitated with cold

acetone at 80%, at 4 °C, for 16 hours, for 2 times. The supernatant containing DTT and salts

was discarded, while the precipitate was dissolved in water and freeze-dried. Then, the

precipitate was purified by size exclusion chromatography (Biogel P10, flow: 12 ml / h) to

completely remove the DTT. The eluted fractions were checked by ^{1}H NMR, revealing that

the glycan-containing material was eluted at one-third of the column volume (full spectra are

shown in Fig. 7B).

The ^{1}H NMR measurements were carried out on 600 MHz Bruker instrument, equipped with

a CryoProbe™ at 310 K. The intensity of the solvent signal was reduced by measuring a

mono-dimensional DOSY spectrum, setting δ and Δ to 2.4 ms and 100 ms, respectively, and

the variable gradient to 50% of its maximum power. Spectra were processed and analyzed

using Bruker TopSpin 4.0.9 program.

genomic fiber. Each fraction of the gradient was finally controlled by negative staining TEM.

Mass spectrometry-based proteomic analyses

Proteins were solubilized with Laemmli buffer (4 volumes of sample with 1 volume of

Laemmli 5X - 125 mM Tris-HCl pH 6.8, 10% SDS, 20% glycerol, 25% β-mercaptoethanol

and traces of bromophenol blue) and heated for 10 min at 95 °C. The extracted proteins were

stacked in the top of an SDS-PAGE gel (4-12% NuPAGE, Life Technologies), stained with

Coomassie blue R-250 (Bio-Rad) before in-gel digestion using modified trypsin (Promega,

sequencing grade) as previously described (36). The resulting peptides were analyzed by

online nanoliquid chromatography coupled to tandem MS (UltiMate 3000 RSLCnano and Q-

Exactive Plus or Q-Exactive HF, Thermo Scientific). Peptides were sampled on a 300 µm x 5

mm PepMap C18 precolumn and separated on a 75 µm x 250 mm C18 column (Reprosil-Pur

120 C18-AQ, 1.9 µm, Dr. Maisch, except for Nqu143-GFP mutant samples separated on

Aurora, 1.7 μm, IonOpticks) using a 140-min gradient (except for fibrils from Nqu143-GFP mutant for which a 60-min gradient was used). MS and MS/MS data were acquired using Xcalibur 4.0 (Thermo Scientific). Peptides and proteins were identified using Mascot (version 2.8.0, Matrix Science) through concomitant searches against homemade *A. castellanii* protein sequence database, homemade virus-specific protein sequence databases, and a homemade database containing the sequences of classical contaminant proteins found in proteomic analyses (human keratins, trypsin...). Trypsin/P was chosen as the enzyme and two missed cleavages were allowed. Precursor and fragment mass error tolerances were set at respectively at 10 and 20 ppm. Peptide modifications allowed during the search were: Carbamidomethyl (C, fixed), Acetyl (Protein N-term, variable) and Oxidation (M, variable). The Proline software version 2.2.0 (37) was used for the compilation, grouping, and filtering of the results: conservation of rank 1 peptides, peptide score ≥ 25, peptide length ≥ 6 amino acids, peptide-spectrum-match identification false discovery rate < 1% (38), and minimum of 1 specific peptide per identified protein group. Proline was then used to perform a MS1-based quantification of the identified protein groups. Intensity-based absolute quantification (iBAQ) (39) values were calculated for each protein group from the MS intensities of razor and specific peptides.

Acknowledgments

We thank Jean-Michel Claverie for his comments on the manuscript and discussions all along the project. The data were collected at the Cryo-EM Swedish National Facility funded by the Knut and Alice Wallenberg, Family Erling Persson and Kempe Foundations, SciLifeLab, Stockholm University, and Umeå University under the expert assistance of Matthieu Coinçon. The preliminary electron microscopy experiments were performed on the PiCSL-FBI core facility (Aïcha Aouane, IBDM, AMU-Marseille), a member of the France-BioImaging national research infrastructure, sample freezing conditions and cryo-EM preliminary acquisitions were performed at the AFMB microscopy platform and on the IMM imaging platform (Dr. Artemis Kosta). We thank the PACA Bioinfo platform for computing support.

Data and materials availability: Genome sequences of the mutants of mimivirus reunion have been deposited to NCBI (accession numbers: KO_qu_946: OQ700912; KO_qu143: OQ700913, 2KO: OQ700914). 3D reconstruction maps and the corresponding PDB have been deposited to EMDB (Deposition number Cl1: 8ORS, EMD-17131; Cl2: 8ORH, EMD-

17125). The mass spectrometry proteomics data have been deposited to the ProteomeXchange

Consortium via the PRIDE (40) partner repository with the dataset identifier PXD041298.

AlphaFold predictions were performed using HPC/AI resources from GENCI-IDRIS (Grant

2022-AD011013526).

Competing interests: Authors declare that they have no competing interests

References

1. A. Villalta, *et al.*, The giant mimivirus 1.2 Mb genome is elegantly organized into a 30-nm
diameter helical protein shield. *eLife* **11**, e77607 (2022).

2. R. A. L. Rodrigues, *et al.*, Mimivirus Fibrils Are Important for Viral Attachment to the Microbial
World by a Diverse Glycoside Interaction Repertoire. *J. Virol.* **89**, 11812–11819 (2015).

3. A. Notaro, *et al.*, Giant viruses of the *Megavirinae* subfamily possess biosynthetic pathways to
produce rare bacterial-like sugars in a clade-specific manner. *microLife* **3**, uqac002 (2022).

4. A. Notaro, *et al.*, Expanding the Occurrence of Polysaccharides to the Viral World: The Case of
Mimivirus. *Angew. Chem. Int. Ed.* **60**, 19897–19904 (2021).

5. E. V. Koonin, *et al.*, Global Organization and Proposed Megataxonomy of the Virus World.
Microbiol. Mol. Biol. Rev. **84**, e00061-19 (2020).

6. D. Raoult, *et al.*, The 1.2-megabase genome sequence of Mimivirus. *Science* **306**, 1344–1350
(2004).

7. C. Abergel, J. Rudinger-Thirion, R. Giegé, J.-M. Claverie, Virus-encoded aminoacyl-tRNA
synthetases: structural and functional characterization of mimivirus TyrRS and MetRS. *J. Virol.*
81, 12406–12417 (2007).

8. J. Abrahão, *et al.*, Tailed giant Tupanvirus possesses the most complete translational apparatus
of the known virosphere. *Nat. Commun.* **9**, 749 (2018).

9. I. Speciale, *et al.*, The Astounding World of Glycans from Giant Viruses. *Chem. Rev.* **122**, 15717–
15766 (2022).

10. N. Zauberman, *et al.*, Distinct DNA exit and packaging portals in the virus Acanthamoeba
polyphaga mimivirus. *PLoS Biol.* **6**, e114 (2008).

11. J. R. Schrad, J. S. Abrahão, J. R. Cortines, K. N. Parent, Structural and Proteomic Characterization
of the Initiation of Giant Virus Infection. *Cell* **181**, 1046-1061.e6 (2020).

12. J.-M. Claverie, C. Abergel, Mimivirus: the emerging paradox of quasi-autonomous viruses.
Trends Genet. **26**, 431–437 (2010).

13. M. Suzan-Monti, B. L. Scola, L. Barrassi, L. Espinosa, D. Raoult, Ultrastructural Characterization
of the Giant Volcano-like Virus Factory of Acanthamoeba polyphaga Mimivirus. *PLoS ONE* **2**,
e328 (2007).

14. Y. Mutsafi, N. Zauberman, I. Sabanay, A. Minsky, Vaccinia-like cytoplasmic replication of the
giant Mimivirus. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 5978–5982 (2010).

15. Y. G. Kuznetsov, T. Klose, M. Rossmann, A. McPherson, Morphogenesis of mimivirus and its
viral factories: an atomic force microscopy study of infected cells. *J. Virol.* **87**, 11200–11213
(2013).

16. I. L. M. de Aquino, *et al.*, Diversity of Surface Fibril Patterns in Mimivirus Isolates. *J. Virol.*,
e01824-22 (2023).

17. M. Boyer, *et al.*, Mimivirus shows dramatic genome reduction after intraamoebal culture. *Proc.*
Natl. Acad. Sci. U. S. A. **108**, 10296–10301 (2011).

18. H. Bisio, *et al.*, Evolution of giant pandoravirus revealed by CRISPR/Cas9. *Nat. Commun.* **14**, 428
(2023).

19. Y. Liu, *et al.*, Virus-encoded histone doublets are essential and form nucleosome-like structures.
Cell **184**, 4237-4250.e19 (2021).

20. S. He, S. H. W. Scheres, Helical reconstruction in RELION. *J. Struct. Biol.* **198**, 163–176 (2017).

21. S. H. W. Scheres, RELION: Implementation of a Bayesian approach to cryo-EM structure
determination. *J. Struct. Biol.* **180**, 519–530 (2012).

22. D. Liebschner, *et al.*, Macromolecular structure determination using X-rays, neutrons and
electrons: recent developments in *Phenix*. *Acta Crystallogr. Sect. Struct. Biol.* **75**, 861–877
(2019).

23. I. Speciale, *et al.*, Liquid-state NMR spectroscopy for complex carbohydrate structural analysis:
A hitchhiker's guide. *Carbohydr. Polym.* **277**, 118885 (2022).

24. M. C. Cassetti, M. Merchlinsky, E. J. Wolffe, A. S. Weisberg, B. Moss, DNA Packaging Mutant:
Repression of the Vaccinia Virus A32 Gene Results in Noninfectious, DNA-Deficient, Spherical,
Enveloped Particles. *J. Virol.* **72**, 5769–5780 (1998).

25. M. I. Valencia-Sánchez, *et al.*, The structure of a virus-encoded nucleosome. *Nat. Struct. Mol.*
Biol. **28**, 413–417 (2021).

26. J. Jumper, *et al.*, Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583–
589 (2021).

27. H. P. Erickson, M. Osawa, Cell division without FtsZ--a variety of redundant mechanisms. *Mol.*
Microbiol. **78**, 267–270 (2010).

28. S. Q. Zheng, *et al.*, MotionCor2: anisotropic correction of beam-induced motion for improved
cryo-electron microscopy. *Nat. Methods* **14**, 331–332 (2017).

29. A. Rohou, N. Grigorieff, CTFFIND4: Fast and accurate defocus estimation from electron
micrographs. *J. Struct. Biol.* **192**, 216–221 (2015).

30. N. Coudray, *et al.*, Deducing the symmetry of helical assemblies: Applications to membrane
proteins. *J. Struct. Biol.* **195**, 167–178 (2016).

31.   R. Diaz, W. J. Rice, D. L. Stokes, Fourier-Bessel reconstruction of helical assemblies. *Methods*
Enzymol. **482**, 131–165 (2010).

32.   C. Sachse, Single-particle based helical reconstruction—how to make the most of real and
Fourier space. *AIMS Biophys.* **2**, 219–244 (2015).

33.   E. F. Pettersen, *et al.*, UCSF Chimera-A visualization system for exploratory research and
analysis. *J. Comput. Chem.* **25**, 1605–1612 (2004).

34.   P. Emsley, B. Lohkamp, W. G. Scott, K. Cowtan, Features and development of *Coot*. *Acta*
Crystallogr. D Biol. Crystallogr. **66**, 486–501 (2010).

35.   C. De Castro, M. Parrilli, O. Holst, A. Molinaro, "Microbe-Associated Molecular Patterns in
Innate Immunity" in *Methods in Enzymology*, (Elsevier, 2010), pp. 89–115.

36.   M. G. Casabona, Y. Vandenbrouck, I. Attree, Y. Couté, Proteomic characterization of
Pseudomonas aeruginosa PAO1 inner membrane. *Proteomics* **13**, 2419–2423 (2013).

37.   D. Bouyssié, *et al.*, Proline: an efficient and user-friendly software suite for large-scale
proteomics. *Bioinforma. Oxf. Engl.* **36**, 3148–3155 (2020).

38.   Y. Couté, C. Bruley, T. Burger, Beyond Target-Decoy Competition: Stable Validation of Peptide
and Protein Identifications in Mass Spectrometry-Based Discovery Proteomics. *Anal. Chem.* **92**,
14898–14906 (2020).

39.   B. Schwanhäusser, *et al.*, Global quantification of mammalian gene expression control. *Nature*
473, 337–342 (2011).

40.   Y. Perez-Riverol, *et al.*, The PRIDE database resources in 2022: a hub for mass spectrometry-
based proteomics evidences. *Nucleic Acids Res.* **50**, D543–D552 (2022).

Funding:

This project has received funding from the European Research Council (ERC) under the

European Union's Horizon 2020 research and innovation program (grant agreement No

832601).

Proteomic experiments were partly supported by ProFI (ANR-10-INBS-08-01) and GRAL, a

program from the Chemistry Biology Health (CBH) Graduate School of University Grenoble

Alpes (ANR-17-EURE-0003).

France-BioImaging national research infrastructure (ANR-10-INBS-04).

CDC gratefully acknowledges STARPLUS 2020 (project no. 21-UNINA-EPIG-042) from the

University of Napoli for financial support.