Title:

Variability and Bias in Microbiome Metagenomic Sequencing: an Interlaboratory
Study Comparing Experimental Protocols

Authors:
Samuel P. Forry[1], Stephanie L. Servetas[1], Jason G. Kralj[1], Keng Soh[2], Michalis
Hadjithomas[3], Raul Cano[4], Martha Carlin[4], Maria G de Amorim[5], Benjamin Auch[6],
Matthew G Bakker[7], Thais F Bartelli[5.], Juan P. Bustamante[8,9], Ignacio Cassol[8],
Mauricio Chalita[10], Emmanuel Dias-Neto[5], Aaron Del Duca[11], Daryl M. Gohl[6,12],
Jekaterina Kazantseva[13], Muyideen T. Haruna[14], Peter Menzel[15], Bruno S Moda[5,16],
Lorieza Neuberger-Castillo[17], Diana N Nunes[5], Isha R. Patel[18], Rodrigo D. Peralta[9,19],
Adrien Saliou[20], Rolf Schwarzer[15], Samantha Sevilla[21,22], Isabella K T M Takenaka[5],
Jeremy R. Wang[23], Rob Knight[24], Dirk Gevers[25], Scott A. Jackson[1]

Affiliation
1. Complex Microbial Systems Group, National Institute of Standards and Technology
(NIST), Gaithersburg, MD
2. Novo Nordisk
3. LifeMine Therapeutics, Cambridge Discovery Park, 30 Acorn Park Drive, Cambridge, MA,
02140
4. The BioCollective, LLC, 5650 Washington Street, Suite C9, Denver, CO 80216
5. Laboratory of Medical Genomics, A.C.Camargo Cancer Center, Sao Paulo 01508-010, SP,
Brazil.
6. University of Minnesota Genomics Center, Minneapolis, MN 55455, USA

7. Department of Microbiology, University of Manitoba, Winnipeg, MB R3T 2N2 Canada
8. Facultad de Ingeniería, Universidad Austral, Pilar, Argentina.
9. Instituto de Investigación y Desarrollo en Bioingeniería y Bioinformática (IBB), CONICET-
UNER, Oro Verde, Argentina.
10. CJ Bioscience, South Korea
11. OMX Advisors, Inc., Ottawa, Canada
12. Department of Genetics, Cell Biology, and Development, University of Minnesota,
Minneapolis, MN 55455, USA
13. Center of Food and Fermentation Technologies (TFTAK), Akadeemia tee 15A, Tallinn
12618, Estonia
14. Bio-Environmental Sciences Program, Morgan State University, Baltimore, MD
15. Labor Berlin Charité Vivantes GmbH, Sylter Str. 2, 13353 Berlin, Germany
16. Laboratory of Computational Biology and Bioinformatics, A.C.Camargo Cancer Center,
Sao Paulo 01508-010, SP, Brazil.
17. Integrated Biobank of Luxembourg (IBBL), Luxembourg Institute of Health (LIH),
Dudelange, Luxembourg
18. U. S. Food and Drug Administration, Center for Food Safety and Applied Nutrition, Office
of Applied Research and Safety Assessment, Laurel, MD 20708, USA

19. Laboratorio de Investigación, Desarrollo y Transferencia de la Facultad de Ingeniería de
la Universidad Austral (LIDTUA), CIC-Austral, Pilar, Argentina.
20. OMICS Hub, BIOASTER, Microbiology Research Institute, Lyon, France
21. CCR Collaborative Bioinformatics Resource, Center for Cancer Research, National Cancer
Institute, National Institutes of Health, Bethesda, MD 20892, USA
22. Advanced Biomedical Computational Sciences, Frederick National Laboratory for Cancer
Research, Leidos Biomedical Research, Inc., Frederick, MD 21701, USA
23. Department of Genetics, University of North Carolina at Chapel Hill, Chapel Hill, NC
27599, USA
24. University of California at San Diego, Departments of Pediatrics, Bioengineering and
Computer Science & Engineering, and Center for Microbiome Innovation, 9500 Gilman
Drive, MC 0763, La Jolla, CA 92093-0763 USA
25. Seed Health

Abstract

Background
Several studies have documented the significant impact of methodological choices in
microbiome analyses. The myriad of methodological options available complicate the
replication of results and generally limit the comparability of findings between independent
studies that use differing techniques and measurement pipelines. Here we describe the Mosaic
Standards Challenge (MSC), an international interlaboratory study designed to assess the
impact of methodological variables on the results. The MSC did not prescribe methods but
rather asked participating labs to analyze 7 shared reference samples (5x human stool samples
and 2x mock communities) using their standard laboratory methods. To capture the array of
methodological variables, each participating lab completed a metadata reporting sheet that
included 100 different questions regarding the details of their protocol. The goal of this study
was to survey the methodological landscape for microbiome metagenomic sequencing (MGS)
analyses and the impact of methodological decisions on metagenomic sequencing results.

Results
A total of 44 labs participated in the MSC by submitting results (16S or WGS) along with
accompanying metadata; thirty 16S rRNA gene amplicon datasets and 14 WGS datasets were
collected. The inclusion of two types of reference materials (human stool and mock
communities) enabled analysis of both MGS measurement variability between different
protocols using the biologically-relevant stool samples, and MGS bias with respect to ground
truth values using the DNA mixtures. Owing to the compositional nature of MGS
measurements, analyses were conducted on the ratio of Firmicutes: Bacteroidetes allowing us
to directly apply common statistical methods. The resulting analysis demonstrated that
protocol choices have significant effects, including both bias of the MGS measurement
associated with a particular methodological choices, as well as effects on measurement
robustness as observed through the spread of results between labs making similar
methodological choices. In the analysis of the DNA mock communities, MGS measurement bias
was observed even when there was general consensus among the participating laboratories.

Conclusion

This study was the result of a collaborative effort that included academic, commercial, and
government labs. In addition to highlighting the impact of different methodological decisions
on MGS result comparability, this work also provides insights for consideration in future
microbiome measurement study design.
90
Introduction
Over the last decade, advances in DNA sequencing technology (Next-Generation Sequencing or
NGS) have led to its widespread adoption by the scientific community for myriad applications.
One such application, known as metagenomic sequencing (MGS), has led to a transformation in
how we measure and characterize complex microbial communities of microbiomes. MGS has
emerged as an important and powerful tool as we seek to comprehend the roles of microbes
inside complex and dynamic communities that are both capable of maintaining and harming
human and environmental health. MGS measurements are able to 'see' whole classes of
microorganisms present in a microbiome sample (e.g., all bacteria by 16S rRNA gene amplicon
sequencing (16S), or all dsDNA by whole-genome shotgun WGS); MGS can also assign a relative
abundance to each microorganism in complex samples. [1-4] Because of these advantages,
MGS is being increasingly adopted across diverse application spaces including infectious disease
diagnostics, [5-12] epidemiological investigations, [13-15] food safety, [16] and biothreat
surveillance. [9, 17-19]. The results of MGS measurements have been used to diagnose
infectious diseases that were missed by conventional methods.[20, 21] As such, regulatory
agencies are actively developing new guidance and policies regarding the use of MGS in the
clinic and in other regulated spaces.

While MGS measurements hold great promise in monitoring and understanding microbial
communities, the current impact is often hampered by a lack of reproducibility and
comparability, particularly between different research centers. [22-24] MGS measurement
results are the product of complex workflows incorporating multiple distinct steps and involving
a multitude of methodological choices (e.g., sample collection and storage, DNA extraction and
purification, NGS library preparation either for WGS or 16S, DNA sequencing platform, data
cleanup and processing, bioinformatic analysis, interpretation). Throughout this workflow,
measurement bias (deviation from ground truth) and measurement noise (experimental
variability) are potentially introduced with each step and will depend on the particular
methodological choices made.[25] It is widely recognized that the interlaboratory
reproducibility of MGS microbiome measurements is poor, and there have been numerous
efforts aimed at benchmarking the analytical performance of MGS measurements in terms of
sensitivity, specificity, precision, reproducibility, etc. [26-33] These challenges are well-
documented, and the community has long recognized the need for studies to prioritize and
investigate the sources of variability and bias in the experimental workflow [27] and the need
for standardized materials and methods to improve the comparability and scope of MGS
measurement results.

Designing the studies to identify sources of variability and bias as outlined above comes with its
own set of challenges including: sufficient numbers and diversity of reference samples to help
power the study; testing of a wide range of variables; a lack of consistent data analysis; cost &
coordination. While the task may seem daunting, several groups have taken up the call to begin

to address these challenges. In recognition of the complexity of the workflow, some groups
have broken the MGS workflow into more manageable sections with most of the focus being
directed at characterizing the effect of data processing and analysis either using *in silico*
datasets [32, 33] or metagenomic DNA control material [26, 28, 34-38]. Other groups have
sought to capture bias throughout the workflow by distributing sets of identical microbiome
samples [39, 40].

Herein, we describe the Mosaic Standards Challenge (MSC). The MSC brought together
academic, federal, and private industry partners in an international interlaboratory study
focused on capturing the diversity of protocols and methodological choices involved in NGS-
based microbiome measurements and understanding their impact on observed taxonomic
profiles. To achieve this, we produced a panel of homogeneous microbiome samples,
developed a custom cloud-based web portal for collecting sequencing data and metadata, [41,
42] and statistically evaluated the MGS results. The microbiome samples included human feces
from multiple donors and DNA mock communities. For every sample analyzed, nearly 100
metadata parameters describing the MGS protocol were collected, with participation from 44
MGS laboratories. The resulting analysis demonstrated that various protocol choices have
significant effects that range from skewing MGS measurement results (e.g., WGS or 16S
analysis) to increasing measurement robustness (e.g., homogenizer use during DNA extraction).
The ground truth DNA mock community samples revealed that MGS measurement bias can
persist, even when there is consensus (measurement agreement) among results from different
laboratories.

Results
Briefly, the study consisted of three components: reference material selection and production,
broad participation from the microbiome community including metadata reporting and MGS
data uploads, and common analysis pipelines applied to the raw sequencing data alongside the
methodological metadata from each participating laboratory. The timeline and overall
workflow of the MSC are shown in Figure 1.

156
157 *Figure 1 Study design timeline. Inset image shows material received by participants.*

Material Production

The reference samples selected and distributed in this study included 5 human stool samples and 2 DNA mixtures (mock DNA communities). The five stool samples were selected from a pool of potential donors based on the dissimilarity of their microbiome composition (Figure 2).

Figure 2 Principal coordinate analysis of donor samples in the BioCollective stool collection. Stool samples included in the Mosaic study (green points) were selected based on their PCoA diversity within the constellation of samples available from the BioCollective. All selected donors self-reported being healthy except BC001485, who reported Parkinson's Disease.

For each sample, multiple stool donations from an anonymous individual were homogenized in the presence of a stabilization buffer to produce 1-liter of homogenized, stabilized fecal material. Two allochthonous microorganisms, *Aliivibrio fischeri* and *Leifsonia xyli,* were also added to each batch of stool (~10^8 cells/mL) and homogenized. Approximately 700 aliquots (1 ml per aliquot) were prepared from each of the 5 batches, and aliquots were stored at -20 °C until ready to ship to participants. To verify that these materials were sufficiently homogenous, 10 aliquots were selected randomly from each of the 5 batches and subjected to both 16S and WGS analysis. These results (Figures 3 and Si-1) indicate that (i) each individual sample donor has a unique microbiome composition, and (ii) the stool samples are suitably homogenized (fit-for-purpose).

Figure 3 Metagenomic sequencing analysis of Mosaic stool samples to determine homogeneity of samples. The bar chart shows the relative abundance as measured by 16S rRNA MGS at the genus level for 10 replicate tubes from each stool sample (stool 1-5). Taxa colors denote the 17 most abundant genera overall, as well as an exogenously added internal standard; all other genera are grouped as 'other' and shown in grey. MGS analysis by WGS also exhibited good homogeneity (Figure SI-1).

181 Additionally, 2 DNA-based mixtures were prepared for the MSC where ground-truth taxa
182 abundances could be assigned. Both materials (Mix A and Mix B) were mixtures of genomic
183 DNA (mock communities) that were extracted from pure cultures of 13 bacterial species mixed
184 at roughly equal genomic ratios (Mix A) or at varying abundances across 3-orders of magnitude
185 (Mix B).

Recruitment and Community Participation

To kick-off the study, a targeted media campaign was launched to recruit participation in the
MSC; study enrollment was open from May 2018 until December 2019. [43, 44] Each lab that
volunteered to participate was shipped the 5 stool samples and two DNA samples free of
charge.   By design, the MSC did not prescribe any required methods or instrumentation to the
participants. Rather, participants were instructed to use their own in-house protocols and
encouraged to explore new methods. To capture these methodological details, a
comprehensive standardized metadata reporting sheet was developed and deployed where
participants could record the details of their protocols. This metadata reporting form included
over 100 questions and was intended to capture the most intricate details of each step in the
measurement process (The metadata capture questions are available in the supplemental file
1.) Both methodological data and raw data were then captured using a custom web-based
cloud analytics portal that enabled the collection, storage, analysis, and visualization of MGS
data generated by the MSC participants [41]. This not only facilitated analysis within a single
bioinformatics pipeline; it also enabled participants to view their results in the context of all
other MSC results immediately following upload. Thus, participants could quickly visualize how
their methods compared to others in the community.

A total of 44 labs participated in the MSC by submitting MGS results (16S or WGS) along with
accompanying metadata (Table 1). Most labs analyzed all samples, though some only analyzed
the stool samples. Of the 44 MGS submissions, 30 were 16S rRNA datasets and 14 were WGS
datasets (Table 1). On average, 16S rRNA MGS datasets had $\geq 10^5$ reads, while the WGS analyses
were typically a log higher with $>10^6$ reads (Figure SI-2). Significant variation in read number
was observed both between participating labs and individual samples (Figure SI-2).

Table 1 A total of 44 labs submitted 16S or WGS analyses of the Mosaic Stool and DNA samples with accompanying metadata.
Two sequencing datasets with incomplete metadata were dropped.

MGS Analysis	Samples analyzed	Number of Labs
16S	5x Stool	8
	2x DNA	0
	7x (Stool and DNA)	22
WGS	5x Stool	0
	2x DNA	0
	7x (Stool and DNA)	14

211
212
213
214

Metagenomic Sequencing (MGS) Interlaboratory Comparison

A Bray-Curtis principal coordinate analysis (PCoA) for both the 16S (n=150) and WGS (n=70) datasets (Figure 4a and 4b) demonstrate that the biological variability (i.e., stool sample ID) was the major factor influencing the overall ordination of the data, as expected.   The impact of methodological variability can be seen via the dispersal of datasets within each stool sample. ~~It's noteworthy that the outputs from the 16S datasets and the WGS datasets were so dissimilar that separate PCoA analysis plots were required.~~ From the PCoA plot of the 16S data (Figure 4a), we observed that one of the participating labs made an apparent transposition in the labeling of samples 3, 4, and 5.   Based on this apparent error, we excluded all the data (stool samples 1-5) from this lab for the remainder of the analyses described in this manuscript.

Figure 4: Principal coordinate plots of the Bray-Curtis dissimilarities for 16S and WGS analyses exhibits clustering by Stool sample. Each data point represents a distinct laboratory analysis of each sample. The separation in the clusters is attributed to methodological differences between labs.

Firmicutes:Bacteroidetes ratio

Because of the compositional nature of MGS results, individual taxa relative abundances are not directly comparable between different samples.[45, 46] Instead, ratios of taxa within each sample were expected to be more reliable because the effects of sample composition on each taxa relative abundance could cancel out. One ratio that has been of interest in the field is the ratio of phyla Firmicutes:Bacteroidetes; therefore, we chose this ratio to demonstrate the utility of using ratios of taxa to compare data between samples [39-42].   Thus, this ratio was utilized and included in our results purely for its bioinformatic utility and is not intended to serve as an indicator of gut health or dysbiosis. The Firmicutes:Bacteroidetes ratio was calculated for each Mosaic stool sample and compared among the individual laboratory results (Figure 5). As was expected since each laboratory used their individual MGS protocols (e.g., methodological choices for DNA extraction, library preparation, and sequencing), the Firmicutes:Bacteroidetes ratio varied substantially both between stool samples within each lab, as well as between labs.

Amplicon vs. Shotgun sequencing

One goal of the MSC was to determine how the selection of different methodological parameters during MGS would lead to observed differences in the taxonomic profiles and relative abundances. The highest-level methodological choice was between 16S MGS or WGS MGS. Indeed, the Firmicutes:Bacteroidetes ratio was affected by the type of sequencing performed, with 16S MGS analyses reporting significantly higher Firmicutes:Bacteroidetes ratios (Figure 6a). While the majority of the 16S MGS datasets indicated that Firmicutes were present at a higher relative abundance than Bacteroidetes, WGS data found the inverse with Bacteroidetes being present at a higher relative abundance than Firmicutes. The magnitude of this effect was quantified by averaging the results from all labs reporting each methodological parameter (e.g., 16S or WGS for sequencing strategy) divided by the average result overall and plotted as a fold change on a log scale (Figure 6b). The dependence of the Firmicutes:Bacteroidetes ratio on analysis strategy that was observed in this dataset could explain recent reports that question the reliability of the Firmicutes:Bacteroidetes ratio as a diagnostic indicator of gut health [47]. This dependence was consistent across all five stool samples (Figure SI-3), and an analysis of each sample's alpha diversity also yielded a similar stratification with respect to the methodological choice of 16S or WGS analysis (Figure SI-4).

264
Figure 6 The effect of analysis strategy (16S versus WGS) on the Firmicutes:Bacteroidetes ratio was readily observed for just
one stool sample by simple grouping (A), and the effect was quantified (B) by dividing the average results among labs reporting
the specified parameter level by the average results overall. In B, this parameter effect was plotted on a log (base 2) scale, such
that the horizontal line at 0 denotes the null hypothesis of no effect; error bars show the 99% confidence interval. Quantified
effects for the other stool samples were similar and are included in Figure SI-3. Similar stratification was observed when
measuring each sample's Inverse Simpson alpha diversity (Figure SI-4).

<u>Other metadata parameters</u>
When submitting results, participating labs were asked to complete a standardized metadata
reporting sheet that included 100 different questions regarding the details of their protocol.
Some questions were generally applicable like "what sequencing instrument did you use" while
others were more nuanced like "what was the PCR primer set used." As such, some fields were
required, and others were optional. Because of the large impact generated by the 16S
vs. WGS methodological variable (Figure 6) and the hierarchical nature of other methodological
choices (e.g., 'What was the target gene amplicon'), we chose to analyze each data set
separately. The effect on the Firmicutes:Bacteroidetes ratio on the 16S MGS results was
quantified for each subsequent methodological choice (Figure 7) in a similar manner to that
employed in Figure 6b. While there were many methodological variables that appeared to have
a significant impact on the results (Figure 7; similar analysis for other stool samples is included
in Figure SI-5), many of these were only reported by a single lab (n=1).   Of the 30 labs
submitting 16S MGS data, there were 14 methodological differences in their protocols. Of the
14 labs submitting WGS data, there were 9 methodological differences in their protocols (Figure
SI-6). Not all methodologic variables had a significant impact on the result. Methodological
variables that were observed to have a significant impact on the 16S MGS results for 2 or more
stool samples (parameter effect and 99% confidence interval) included the manufacturer of the
DNA extraction kit and the target gene for amplification (Figure 7 and SI-5). Methodological
variables that were observed to have a significant impact on the WGS results (parameter effect)
for 2 or more stool samples included the DNA extraction protocol, the manufacturer of the DNA
extraction kit, and the library kit for shotgun sequencing. In addition to their impact on the
parameter effect as described above, some methodological variables were observed to have a
significant impact on the robustness of the measurement (observed as a lack of variability when
other parameters are varied). For example, when asked whether a homogenizer or shaking

apparatus was used, those labs that reported "Yes" displayed much greater robustness than
those reporting "No" (Figure 7 and SI-5 and SI-6).

298
299 *Figure 7 Within labs performing 16S amplicon sequencing, the parameter effect on the Firmicutes:Bacteroidetes ratio was*
300 *calculated as described in Figure 6 for each relevant metadata parameter. Shown here from just one stool sample, results from*
301 *other stool samples were similar and are provided in Figure SI-5.*

'Spike-in' organisms

An additional attribute of the fecal materials used for this interlaboratory study was the
inclusion of two exogenous organisms to serve as whole-cell internal controls (i.e., spike-ins).
Since these organisms were added during the bulk homogenization step, their abundance
should be constant across all the stool sample aliquots.   As such, it was expected that the ratio
of *A. fisherii* to *L. xyli* would be constant for each particular methodology (e.g., within a lab).
Surprisingly, *L. xyli* was not identified in any of the submitted 16S datasets and was only
observed at a low abundance (approximately 0.001 %) by WGS analysis. When the
A. fisherii:*L. xyli* ratio (by WGS) was plotted for each participating laboratory (Figure SI-7),
significant variability between samples was observed. These data were unexpected and could
have resulted from poor database representation of *L. xyli* in the commercially available
bioinformatic pipeline used, inefficient DNA extraction, or low or inconsistent distribution
during material manufacture, among other possible explanations.

Genomic DNA mixtures

Another control included in the interlaboratory study were mixtures of purified microbial
genomic DNA. These were included alongside the stool samples in the Mosaic Kit to serve as
parallel processing controls and included two different mixtures, one equigenomic between
taxa (Mix A) and one with ten-fold dilutions of the various taxa (Mix B). These genomic DNA
mixtures were validated for genome copy number using ddPCR (droplet digital PCR) and serve
as 'ground truth' for the MGS measurements.   For comparison to the MGS measurements,

genome copy number (as measured by ddPCR) was scaled by the assembled genomes of the
individual strains (i.e., rRNA copy number or genome size) to yield ground truth values for
comparison to 16S or WGS results, respectively. As with the Firmicutes:Bacteroidetes ratio
described above, ratios of individual taxa were used to characterize the DNA mock communities
and remove the compositional dependence of the raw relative abundance assignments. Since
these analyses included 16S sequencing results, we focused on strains that were unique at the
genus level, yielding 6 distinct ratios within each sample. The independent determination of
actual DNA concentration (ddPCR) was compared to the results of MGS analyses (Figure 8).
While there was some agreement among participating laboratories (consensus), their results
generally differed from the actual abundances. Overall, this indicates that even when
consensus exists among MGS results, significant unidentified bias can remain. Further, this was
taxa-dependent, with some taxa (e.g., Kp by 16S or Pa by WGS) producing particularly
significant variability and deviation from ground truth.

335
Figure 8 For the DNA mixtures, independently measured 'ground truth' results (black 99% confidence intervals) for the ratios
between taxa relative abundances can be compared to each individual lab's amplicon (red points) or shotgun (blue points)
metagenomic sequencing results, as well as the range of results (grey boxplots) among participating labs. The taxa in Mix A
were roughly equally abundant, while the Mix B sample exhibited groups of taxa added at 10-fold dilutions. The horizontal axis
identifies the taxa (known to be present in the DNA mixtures) whose observed relative abundances were ratioed. The ground
truth values were scaled to account for known 16S copy numbers (for amplicon sequencing) or genome sizes (for shotgun
sequencing), so the 'actual' ratios vary slightly between the two analyses even though the DNA concentrations are identical.
Genus-level taxonomic bar charts by (16S and WGS analysis) show the average composition observed for each DNA mixture
(Figure SI-8).

Discussion

The MSC represented the third in a series of community challenges of increasing complexity
hosted by Janssen's Human Microbiome Institute (JHMI) as an effort to improve the overall
quality of microbiome MGS measurements. This study was designed and implemented through
a collaborative effort that included the Janssen Human Microbiome Institute (JHMI), The
BioCollective, LLC (TBC), DNAGenotek, DNANexus, and the National Institute of Standards and

Technology (NIST) which serves as the National Metrology Institute for the U.S. These
organizations in turn represent biopharmaceutical companies, biotechnology companies, data
analytics companies, and Federal Government laboratories, all of whom have a vested interest
in reliable and comparable microbiome measurements. The goal of the MSC was to capture the
diversity of protocols for MGS-based microbiome measurements in an effort to begin to
elucidate the impact of these methodological variables on the resulting taxonomic profiles and
guide the development of future reference materials.

The MGS workflow required for microbiome analyses is complex.   Therefore, designing an
interlaboratory study that includes a multitude of the methodological variables and assesses
their effect on the results is an ambitious project. Several teams have sought to address the
question of methodological bias and variability over the years. [26-28, 31-33, 39, 40] These
investigations have taken a variety of approaches from prescribing locked-down SOPs and
analyzing specific samples to more open-ended data collection. The interlaboratory study
presented herein specified seven samples for analysis (5 different stool samples and 2 DNA
mixtures) while intentionally leaving protocol choices up the participating labs, both to sample
a diverse set of methodological parameters as well as to survey common methodological
choices.

The design and implementation of this project can be broken into three major areas: (1)
reference material selection and production, (2) capturing metadata and MGS raw data, and (3)
comparing results between participating laboratories.

Reference Material Production
One of the first decisions was the identification of reference material(s) to include. There are
two primary types of materials that have been used for this type of study: (1) biologically
derived microbiome samples and (2) mock communities. Both types of materials were included
in the current investigation because they are useful in different ways for comparing between
diverse analytical workflows.

For biologically derived microbiome reference materials, a natural community (e.g. sludge, soil,
fecal material) is collected, homogenized, and aliquoted. Previous interlaboratory studies have
used these homogenized real-world materials [31, 39]; however, the number of units needed
and the associated costs of a large-scale study are often prohibitive. Further, while biologically-
derived materials represent the complexity and diversity of real-world samples, they currently
lack ground-truth value assignments (e.g. actual taxonomic abundances) due to a lack of
unbiased analytical methods (e.g., DNA extraction, PCR amplification) and the inherent
ambiguity associated with microbial taxonomy that hinders our ability to define clear
measurands (e.g., Escherichia coli vs. Shigella, or the recent reclassification of Lactobacillus into
23 novel genera). [48] The addition of allochthonous bacteria ("spike-ins") at consistent
abundances into biologically-derived materials can provide some ground truth values to
facilitate the assessment of MGS measurements.

Nevertheless, these biologically-derived materials remain useful for comparing methods and
assessing measurement precision within individual laboratories and across different
laboratories. In the current study, five stool samples were selected based on their dissimilarity

from one another among a constellation of potential stool donors (Figure 2), with the intention
of representing the variability of naturally-occurring samples. Preliminary in-house analysis of
individual aliquots demonstrated (Figure 3) that the material collection and preparation
resulted in samples with reliable between-aliquot homogeneity, even given the inherently
inhomogeneous starting point of multiple donations of human stool.

Mock community reference materials are laboratory-prepared mixtures of defined constituents
(typically DNA from individually cultured bacteria; sometimes mixtures of whole cells) at
specified amounts. Thus, these materials are useful as 'ground truth' for analysis workflows,
allowing quantitative assessment of analytical performance (e.g., accuracy, bias, precision, etc.).
However, these mock community materials are inherently non-biomimetic of actual
microbiome samples (e.g., feces, soil, etc), namely due to their low complexity and the absence
of a matrix-effect, which can limit their utility for assessing analysis workflows. [26] By including
both types of reference materials (5x stool samples and 2x mock communities) in the MSC, we
sought to include the benefits of both, using the biologically-derived materials to assess MGS
measurement variability between different protocols, and using the DNA mixtures to assess
MGS bias with respect to ground truth values.

Capturing Metadata
The universe of discrete MGS methodologies is quite large. Preliminary projections during
project planning estimated that several hundred samples would be needed to fully explore this
methodological space. Thus, the MSC set-out to host an international interlaboratory study on
an unprecedented scale. 700 units of reference material were prepared and made available
free-of-charge, where each unit consisted of 5 distinct, biologically-derived human fecal
microbiome samples and 2 DNA mixtures (mock communities). To our knowledge, there has
never been an MGS interlab study designed on such a massive scale. However, during the 19-
month marketing campaign, only ~100 units were distributed. Further, from these recipients,
only 44 sets of raw sequencing data and metadata were submitted (Table 1), limiting the
statistical power of the resulting analyses. Nevertheless, the unused units remain currently
available from The BioCollective, allowing interested researchers to analyze with their own
methods using the same samples that have been characterized and reported on here.

Alongside the raw sequencing data submitted, participating laboratories filled out a metadata
questionnaire (available in supplemental file 1) with ~100 discrete questions about the
methods employed, most of which allowed selection from drop-down options describing the
most common methodological choices. However, it must be noted that even these in-depth
options were not sufficient to encompass all experimental possibilities, and many metadata
selections represented 'Other' or 'Internal Method' options. And, of course, the number of
potential methodologies continues to expand as new techniques are developed or made
commercially available. It was also apparent within the submitted metadata that the observed
methodological choices were not randomly distributed. There was no effort made in this
investigation to encourage exploration of a diverse set of methodologies, and groups tended to
cluster around common methods. The resulting metadata reflect the most employed methods
during the timeframe of this study (Figure. 1). For instance: nearly half of participants analyzing
samples by 16S reported using the same DNA extraction kit (there were ~15 other pre-

identified options, as well as 'In-house' and 'other' possibilities); and only 2 labs (~4%) used
non-Illumina sequencing platforms.

Comparing Results Between Laboratories
To help assess the impact of methodological choices in the context of compositionally-sensitive
MGS measurements, we focused here on ratios between Phyla (e.g., the
Firmicutes:Bacteroidetes ratio: Figure 5) instead of the raw relative abundances [46]. By using
this strategy to remove the compositional dependence of MGS results, common statistical tools
(e.g., mean, standard deviation, confidence intervals) could be directly applied. However, it
must be noted that the Firmicutes:Bacteroidetes ratio only reveals the impact of particular
methodological choices on the tested phyla (Firmicutes and Bacteroidetes). Thus, a
methodological choice that only impacted Proteobacteria, as well as one that affected
Firmicutes and Bacteroidetes similarly, would not be noted herein. Nevertheless, significant
variability in the Firmicutes:Bacteroidetes ratio was observed (Figure 5) both between samples
(presumably due to real differences between the samples) and between participating
laboratories (presumably due to differences in measurement methodology).

When comparing between methodologies, the most basic experimental choice is between 16S
and WGS, and this choice had further implications for how subsequent steps were performed
(e.g., PCR conditions, library prep, sequencing depth, bioinformatic analysis). Thus, we first
compared the Firmicutes:Bacteroidetes ratio between analysis methods (Figure 5). In this case,
it turned out that the most basic choice of how to analyze samples had a statistically significant
effect (Figure 6). Analysis of each of the stool samples individually (Figure SI-3) or statistical
analysis using alpha diversity instead of the Firmicutes:Bacteroidetes ratio (Figure SI-4)
confirmed the significant impact of analysis strategy on observed results. Practically, this raises
real concerns about the comparability of data results between laboratories whose analyses
differ between 16S and WGS analysis. More generally, researchers should use utmost caution
when trying to compare between data sets collected using divergent experimental methods.

Within the data collected for the MSC, the significant effect observed for the choice of analysis
strategy had the specific implication of further limiting statistical power (e.g., of the 44
participating labs, 30 reported 16S results and 14 reported WGS results). Nevertheless, the
observed effects of other methodological choices could be similarly assessed for 16S (Figure 7)
or WGS (Figure SI-4) results. Interestingly, while the statistical power was limited in this study,
some methodologies still appeared to have either large effect sizes or large impacts on
variability/precision. While it is tempting to draw firm conclusions from the current
investigation, caution is warranted due to the limited sample sizes. Instead, it is hoped that this
investigation will help guide further investigations.

'Spike-in' organisms
During production of the stool samples, two exogeneous, 'spike-in', whole cell bacterial strains
were included, *A. fischeri* and *L. xyli*. Both strains are typically absent in human stool. With the
addition of 10^8 cells/mL, each organism was expected to comprise approximately 1 % of the
total stool relative abundance, providing sufficient signal for identification without significantly
affecting the overall sample profile. Unfortunately, while this expectation proved accurate for
A. fischeri, *L. xyli* was not identified in any the of 16S MGS results and was only observed at a

very low relative abundance by WGS (Figure SI-7). This absence or low-level detection could be
the result of a number of sources including lack of representation in the databases, bias in the
DNA extraction of *L. xyli,* or the amount of *L. xyli* added to the samples. However, multiple
coauthors were individually able to reliably detect *L. xyli* using alternate bioinformatic pipelines,
so it is likely that its limited detection in this dataset reflects a shortcoming in the reference
database used [data not shown and manuscript in preparation]. This explanation is also
supported by the observation that for WGS analyses, the variability of the ratio of spike-in
relative abundances between samples was somewhat improved among the labs with the
deepest sequencing results (Figure SI-2). It is worth noting that all raw fastq data submitted
through the Mosaic Standards Challenge has been archived and made publicly available for the
exploration of alternate bioinformatic methods.

The inability to reliably detect *L. xyli* within the framework of this project impacts our ability to
accurately and confidently use *A. fischeri* as well since observing a constant ratio between the
two spike-in organisms is fundamental to trusting their utility (Figure SI-7). Nevertheless, key
considerations were identified for future experimental design and implementation of internal,
spike-in controls. First, the strain should normally be absent in the sample, but still identifiable
by the analysis/database used. This can be tricky because databases often focus on the
organisms commonly encountered in each type of sample, and because the users of
bioinformatic pipelines may not have easy access to the underlying reference databases at the
time of analysis. Second, spike-in abundance should be sufficiently high that it can withstand
potential losses in the processing and still be identified, while not significantly compromising
the fraction of sample reads allocated the organisms native to each sample. This is in turn
complicated by the dependence of the observed relative abundance of any spike-in organism
on the MGS methods to be employed and their potential for bias with respect to each spike-in
organism. And third, the inclusion of additional spike-in organisms (e.g., 3-4 spike-ins total)
should be considered when MGS workflows have not been identified and tested a priori. This
provides redundancy to accommodate wide ranges of MGS methodologies and biases. In this
study, the inclusion of additional organisms could have avoided the problematic absence of *L.*
xyli in the reference database.

DNA Mock Communities
The DNA mixtures provided the ground-truth component in this study. Here, measurement bias
was observed as a disagreement between the actual ratios (black bars show the 99 %
confidence interval) and observed ratios (red and blue points) in Figure 8. This bias depends on
both the particular taxa analyzed, as well as the methods employed (16S vs. WGS is broken out
here). Interestingly, even where there was consensus between participating labs (i.e., a narrow
boxplot indicating strong consensus), substantial bias was still observed (low accuracy). The
consensus between participating labs is particularly apparent in the WGS analysis of the equi-
genomic DNA mixtures (upper right panel, Figure 8) suggesting some systematic bias affecting
each lab. Of note, these mixtures were comprised of genomic DNA from a prototype reference
material. Since the time of the MSC, NIST has completed a full characterization of DNA from 19
bacterial strains; NIST Reference Material, RM 8376, is now available for researchers to
construct their own DNA based mock communities [49].

Conclusion

From 2017 to 2020, the MSC provided a set of biologically-derived and mock community
microbiome samples, at no charge, to any interested MGS research group in an effort to
identify the extent of methodological variability between researchers and assess its impact on
measured taxonomic profiles. 44 research groups submitted both raw MGS data and detailed
metadata about their in-house sample-handling protocols; although this represents a large
number of participating laboratories by most interlaboratory efforts, it remained statistically
limiting for the large number of metadata parameters ($\approx$100) that were explored. Initial choices
about analysis strategy (i.e., amplicon vs. shotgun) significantly impacted the observed
Firmicutes:Bacteroidetes ratios across all samples. The null hypothesis of no significant effect
could not be ruled out for most methodological choices within this study, though some
appeared to have real effects on results (i.e., bias) or measurement precision (i.e., variability).
Thus, the results collated herein should help refine the scope of future assessments of
methodological choices. To this end, researchers at NIST have undertaken a pairwise approach
to systematically compare select steps within the metagenomic workflow (manuscript in
preparation). Additionally, through the inclusion of DNA mock communities with
independently-measured ground-truth abundances, we were able to assess the accuracy of
MGS measurements and observe significant and systematic measurement bias, even when
participating laboratories achieved similar results. Overall, the MSC effort has significantly
expanded our understanding of the impact of methodological choices on MGS measurement
results and precision.

Methods[1]
Selection of Stool Donors
A total of 5 donors were selected from a donor pool maintained by TBC. Figure 2 shows a Bray-
Curtis PCoA ordination plot the entire donor pool, including the 5 donors selected, based on
their gut microbiome composition.   The 5 donors were selected based on the dissimilarities of
their microbiome composition (Figure 2).

Sample Collection and Processing
All stool samples were collected in accordance with TBC's Institutional Review Board protocol
and have been de-identified. The donors were provided with collection kits, and samples were
returned to the TBC via overnight shipping for processing. Upon receipt, the samples were
aseptically transferred to a zip-top bag for dispensing. The samples were stored at -80 °C in 30 g
aliquots until further processing. Multiple bowel movements were collected and pooled from
each donor. Material from each donor was processed individually (to avoid cross
contamination) and inside a biological safety cabinet. Using a Ninja blender, 150 g of fecal

---

1. Certain commercial equipment, instruments, or materials are identified in this paper to foster understanding. Such identification does not imply recommendation or endorsement by the National Institute of Standards and Technology, nor does it imply that the materials or equipment identified are necessarily the best available for the purpose. The reference materials used in this study were not certified by NIST and are not official NIST Reference Materials.

material was combined with 150 g to 300 g of dry ice and homogenized into a fine powder. The
blender was loosely covered with a sterile lab tissue and placed in a -20 °C freezer overnight to
allow the remaining dry ice to sublime. For each sample, before the addition of OMNIgene
Stabilizing Solution (OGS), 50 g of neat powder was set aside and stored at -80 °C.
Approximately 90 g of stool powder was added to 750 mL of OGS. The solution was covered
and left to stir overnight at room temperature. The following morning, 1 mL aliquots were
prepared and stored at –80 °C.

Addition of Spike-In Bacteria and Aliquoting of Samples
Spike-in bacteria, *Aliivibrio fischeri* (formerly known as *Vibrio fischeri*, Gram negative) and
Leifsonia xyli (Gram positive), were grown to an approximate density of 10^8 CFU/mL and
10^9 CFU/mL, respectively. Cell concentration was confirmed via plate count and optical density.
The spike-in bacteria were concentrated by centrifugation, resuspended, and added to each
stool solution 1 hour prior to aliquoting to ensure thorough homogenization. Working in a
biological safety cabinet, the solution was aliquoted using wide-bore pipette tips into (800 to
850) aliquots. Final concentration of stool after addition of the spike-in was 100 mg/mL and
final concentration of each spike-in organism was 10^8 CFU/mL. Samples were stored at -80 °C
until distribution.

Sample QC
To assess the homogeneity of the stool samples, ten aliquots from each donor pool were
subjected to 16S rRNA amplicon sequencing and shotgun metagenomic sequencing. All sample
processing, DNA extraction, library preparation and sequencing steps were conducted at
CosmosID (Germantown, MD) using proprietary protocols. For the 16S sequence data, reads
were demultiplexed using split_libraries.py with default filtering parameters. 16S rRNA gene
sequences were then sorted based on sample ID using the QIIME script
extract_seqs_by_sample_id.py. Bacterial operational taxonomic units were selected using
pick_open_reference_otus.py workflow. 16S rRNA taxonomy was defined by ≥ 97 % similarity to
reference sequences using the *core_diversity_analyses.py script*. Alpha diversity, alpha
rarefaction curves, and taxonomy assignments were determined using the core_diversity.py
workflow. Data were rarefied to 100,000 sequences per sample to minimize the effect of
disparate sequence number on the results. Alpha diversity metrics were computed from the
average of 100 iterations from the alpha collated results. Microbiome features were quantified
from metagenome data using existing [Metaphlan2, HUMAnN2, etc.] and in-house pipelines to
identify strain-level taxonomic markers for all samples.

DNA Mixtures
Mixtures of purified genomic DNA from thirteen ATCC-derived strains were prepared in 1X TE
buffer at a final concentration of ≈100 ng/μL.   The two mixtures were made by combining the
genomic DNA from the following bacterial strains: *Staphylococcus aureus* ATCC BAA 44,
Staphylococcus aureus ATCC 12600, *Pseudomonas aeruginosa* ATCC BAA 47, *Enterococcus*
faecalis ATCC 19433, *Salmonella enterica* ATCC 700720, *Salmonella enterica* ATCC 12324,
Escherichia coli ATCC 43895, *Staphylococcus epidermidis* ATCC 12228*, Klebsiella pneumoniae*
ATCC 13883, *Shigella sonnei* ATCC 25931, *Streptococcus pyogenes* ATCC 12344,
Corynebacterium amycolatum ATCC 49386. The individual genomic DNA components were part

of a prototype reference material and were not fully characterized at the time of the MSC.
Subsequent analysis revealed some of the materials were cross contaminated with other
components from the prototype materials including *Achromobacter xylosoxidans.* Mix A was
designed to be equi-genomic with calculated relative abundances by mass of each strain
ranging from ≈6.8 % to ≈10 %. Mix B was designed as a log-dilution of the genomes varying
across 3 orders of magnitude (from ≈0.01 % to ≈30 % by mass). For each mixture (A and B), we
prepared a single pool and then distributed across 700 aliquots where each contained
approximately 20 μL (2 μg) of DNA per aliquot. An average (across all participating laboratories)
relative abundance plot for each sample by amplicon or shotgun sequencing is included in the
supporting information (Figure SI-8). We performed digital droplet PCR (ddPCR) to measure the
absolute abundance as ground truth for the following species in the mixture: *Enterococcus*
faecalis, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, and *Streptococcus pyogenes*. These
species were selected because they were taxonomically distinct within the mixtures at the
Genus level, facilitating MGS discrimination. Pairwise ratios of these abundances provided the
ground truth values depicted in Figure 8. The validated ddPCR assays were reported previously
[49].

Interlaboratory Study Execution:
Recruitment**:** Starting in the Spring of 2018, we launched a media campaign that targeted the
scientific community via social media and email blasts as an attempt to recruit a large and
international cohort of participants. After the MSC launched in May 2018, we continued the
outreach campaign via public speaking engagements at various international microbiology
conferences. We actively recruited volunteers up until January 2020 when the MSC officially
closed. MSC reference materials were shipped to any lab in the world, upon request, from May
2018 till January 2020.

Data Availability: The Mosaicbiome.com web portal was used during MSC to store, analyze,
visualize, and share all the raw data and metadata that was submitted by the MSC participants.
However, in the Spring of 2022, the site was discontinued due to the costs associated with data
storage and maintenance. Therefore, these data (fastq files and metadata summaries) have
been made available via https://data.nist.gov. [exact link TBD]

Sample Availability: At the time of publication, many aliquots of the stool and DNA materials
generated through the Mosaic Standards Challenge still remain available from TBC.
626
Taxonomic Profiling of Interlab Data
All raw sequence data (fastq files) generated by interlab participants were downloaded from
the MosaicBiome web portal and subsequently analyzed via the CosmosID
(www.CosmosID.com) taxonomic classification tool using the CosmosID reference genome
databases (WGS version: 1.0.2; 16S version: 1.1.0). The MGS results (taxonomic profiles) for all
the MSC data are publicly available and can be found by visiting https://app.cosmosid.com and
following the directory structure: *Datasets -> Example_Datasets -> Mosaic_Microbiome*.

Analysis of mosaic data results and methodological parameters
A total of 50 datasets were received. Two datasets were dropped due to incomplete metadata
and an additional 4 labs were The MGS results (taxonomic profiles) and associated metadata

from all submitted data sets were analyzed using R. The raw data and code used for analysis
and to generate the figures in this manuscript have been shared via https://data.nist.gov. [exact
link TBD]

Acknowledgements

641 Funding for the production of the fecal reference materials and reference material shipping was
642 generously provided by the Janssen Human Microbiome Institute (JMHI) .

References

1.	Quince, C., et al., *Shotgun metagenomics, from sampling to analysis.* Nat Biotechnol, 2017.
35(9): p. 833-844.
2.	Lu, J., et al., *Bracken: estimating species abundance in metagenomics data.* PeerJ Computer
Science, 2017. **3**.
3.	Knight, R., et al., *Best practices for analysing microbiomes.* Nat Rev Microbiol, 2018. **16**(7): p.
410-422.
4.	Jovel, J., et al., *Characterization of the Gut Microbiome Using 16S or Shotgun Metagenomics.*
Front Microbiol, 2016. **7**: p. 459.
5.	Simner, P.J., S. Miller, and K.C. Carroll, *Understanding the Promises and Hurdles of Metagenomic
Next-Generation Sequencing as a Diagnostic Tool for Infectious Diseases.* Clin Infect Dis, 2018.
66(5): p. 778-788.
6.	Qin, N., et al., *Alterations of the human gut microbiome in liver cirrhosis.* Nature, 2014.
513(7516): p. 59-64.
7.	Mokili, J.L., F. Rohwer, and B.E. Dutilh, *Metagenomics and future perspectives in virus discovery.*
Curr Opin Virol, 2012. **2**(1): p. 63-77.
8.	Gu, W., S. Miller, and C.Y. Chiu, *Clinical Metagenomic Next-Generation Sequencing for Pathogen
Detection.* Annu Rev Pathol, 2019. **14**: p. 319-338.
9.	Deurenberg, R.H., et al., *Application of next generation sequencing in clinical microbiology and
infection prevention.* J Biotechnol, 2017. **243**: p. 16-24.
10.	Chiu, C.Y. and S.A. Miller, *Clinical metagenomics.* Nat Rev Genet, 2019. **20**(6): p. 341-355.
11.	Blauwkamp, T.A., et al., *Analytical and clinical validation of a microbial cell-free DNA sequencing
test for infectious disease.* Nat Microbiol, 2019. **4**(4): p. 663-674.
12.	Besser, J., et al., *Next-generation sequencing technologies and their application to the study and
control of bacterial infections.* Clin Microbiol Infect, 2018. **24**(4): p. 335-341.
13.	Land, M., et al., *Insights from 20 years of bacterial genome sequencing.* Funct Integr Genomics,
2015. **15**(2): p. 141-61.
14.	Scholz, M., et al., *Strain-level microbial epidemiology and population genomics from shotgun
metagenomics.* Nat Methods, 2016. **13**(5): p. 435-8.
15.	Truong, D.T., et al., *Microbial strain-level population structure and genetic diversity from
metagenomes.* Genome Res, 2017. **27**(4): p. 626-638.
16.	Jagadeesan, B., et al., *The use of next generation sequencing for improving food safety:
Translation into practice.* Food Microbiol, 2019. **79**: p. 96-115.
17.	Be, N.A., et al., *Metagenomic analysis of the airborne environment in urban spaces.* Microb Ecol,
2015. **69**(2): p. 346-55.
18.	Hsu, T., et al., *Urban Transit System Microbial Communities Differ by Surface Type and
Interaction with Humans and the Environment.* mSystems, 2016. **1**(3).
19.	Russell, J.A., et al., *Unbiased Strain-Typing of Arbovirus Directly from Mosquitoes Using
Nanopore Sequencing: A Field-forward Biosurveillance Protocol.* Sci Rep, 2018. **8**(1): p. 5417.

20.   Greninger, A.L., et al., *Clinical metagenomic identification of Balamuthia mandrillaris*
encephalitis and assembly of the draft genome: the continuing case for reference genome
sequencing. Genome Med, 2015. **7**: p. 113.

21.   Somasekar, S., et al., *Viral Surveillance in Serum Samples From Patients With Acute Liver Failure*
By Metagenomic Next-Generation Sequencing. Clin Infect Dis, 2017. **65**(9): p. 1477-1485.

22.   Berg, G., et al., *Microbiome definition re-visited: old concepts and new challenges.* Microbiome,
2020. **8**(1): p. 103.

23.   Schloss, P.D., *Identifying and Overcoming Threats to Reproducibility, Replicability, Robustness,*
and Generalizability in Microbiome Research. Mbio, 2018. **9**(3).

24.   Brooks, J.P., *Challenges for case-control studies with microbiome data.* Annals of Epidemiology,
2016. **26**(5): p. 336-341.e1.

25.   Plant, A.L., et al., *Improved reproducibility by assuring confidence in measurements in biomedical*
research. Nat Methods, 2014. **11**(9): p. 895-8.

26.   Amos, G.C.A., et al., *Developing standards for the microbiome field.* Microbiome, 2020. **8**(1): p.
1-13.

27.   Poussin, C., et al., *Interrogating the microbiome: experimental and computational considerations*
in support of study reproducibility. Drug Discov Today, 2018. **23**(9): p. 1644-1657.

28.   O'Sullivan, D.M., et al., *An inter-laboratory study to investigate the impact of the bioinformatics*
component on microbiome analysis using mock communities. Sci Rep, 2021. **11**(1): p. 10590.

29.   Zaiko, A., et al., *Towards reproducible metabarcoding data: Lessons from an international cross-*
laboratory experiment. Molecular Ecology Resources, 2022. **22**(2): p. 519-538.

30.   Sinha, R., et al., *The microbiome quality control project: baseline study design and future*
directions. Genome Biology, 2015. **16**(1): p. 1-6.

31.   Sinha, R., et al., *Assessment of variation in microbial community amplicon sequencing by the*
Microbiome Quality Control (MBQC) project consortium. Nature Biotechnology, 2017. **35**(11): p.
1077-1086.

32.   McIntyre, A.B.R., et al., *Comprehensive benchmarking and ensemble approaches for*
metagenomic classifiers. Genome Biol, 2017. **18**(1): p. 182.

33.   Sczyrba, A., et al., *Critical Assessment of Metagenome Interpretation-a benchmark of*
metagenomics software. Nature Methods, 2017. **14**(11): p. 1063-+.

34.   Kennedy, K., et al., *Evaluating bias of illumina-based bacterial 16S rRNA gene profiles.* Appl
Environ Microbiol, 2014. **80**(18): p. 5717-22.

35.   Schirmer, M., et al., *Insight into biases and sequencing errors for amplicon sequencing with the*
Illumina MiSeq platform. Nucleic Acids Res, 2015. **43**(6): p. e37.

36.   Gohl, D.M., et al., *Systematic improvement of amplicon marker gene methods for increased*
accuracy in microbiome studies. Nat Biotechnol, 2016. **34**(9): p. 942-9.

37.   D'Amore, R., et al., *A comprehensive benchmarking study of protocols and sequencing platforms*
for 16S rRNA community profiling. BMC Genomics, 2016. **17**: p. 55.

38.   Jones, M.B., et al., *Library preparation methodology can influence genomic and functional*
predictions in human microbiome research. Proc Natl Acad Sci U S A, 2015. **112**(45): p. 14024-9.

39.   Zaiko, A., et al., *Towards reproducible metabarcoding data: Lessons from an international cross-*
laboratory experiment. Molecular Ecology Resources, 2022. **22**(2): p. 519-538.

40.   Sinha, R., et al., *The microbiome quality control project: baseline study design and future*
directions. Genome Biol, 2015. **16**: p. 276.

41.   Westreich, S., *Examining Variation from Wet-Lab Protocol Choices in Microbiome Data through*
the Mosaic Standards Challenge, in *Inside DNANEXUS*. 2019.

42.   *Mosaicbiome*. 2017; Available from:
https://web.archive.org/web/20220302022932/https://mosaicbiome.com/.

43.   DNANexus, *Mosaic Community Challenge: Standards*. 2018, YouTube. p.
731   https://youtu.be/oZyGrIbLHxk.
732   44.   DNANexus, *Mosaic Standards Tutorial 1 - Joining the Challenge*. 2018, YouTube. p.
733   https://youtu.be/3-KuAOywWK8.
734   45.   Gloor, G.B., et al., *Microbiome Datasets Are Compositional: And This Is Not Optional.* Frontiers in
735   Microbiology, 2017. **8**.
736   46.   McLaren, M., A. Willis, and B. Callahan, *Consistent and correctable bias in metagenomic*
737   *sequencing experiments.* Elife, 2019. **8**.
738   47.   Magne, F., et al., *The Firmicutes/Bacteroidetes Ratio: A Relevant Marker of Gut Dysbiosis in*
739   *Obese Patients?* Nutrients, 2020. **12**(5): p. 1474.
740   48.   Salvetti, E., et al., *Comparative Genomics of the Genus Lactobacillus Reveals Robust Phylogroups*
741   *That Provide the Basis for Reclassification.* Appl Environ Microbiol, 2018. **84**(17).
742   49.   Kralj, J., et al., *Reference Material 8376 Microbial Pathogen DNA Standards for Detection and*
743   *Identification* NIST Special Publication 260-225, 2022.
744
745