Full title:

**Exploring Ethylene-Related Genes in Cannabis sativa: Implications for Sexual Plasticity**

Short title:

**Ethylene and *Cannabis* sexual plasticity**

**Adrian S. Monthony[123], Maxime de Ronne[123] & Davoud Torkamaneh[1234]***

**[1]***Département de phytologie, Université Laval, Québec City, Québec, Canada*
**[2]***Institut de Biologie Intégrative et des Systèmes (IBIS), Université Laval, Québec, Canada*
**[3]***Centre de recherche et d'innovation sur les végétaux (CRIV), Université Laval, Québec, Canada*
**[4]***Institut intelligence et données (IID), Université Laval, Québec, Canada*

Monthony ORCID:  0000-0003-0536-691X

de Ronne ORCID: 0000-0003-4062-3512

*Torkamaneh ORCID:  0000-0002-9782-5695

*Corresponding Author

Email:  davoud.torkamaneh.1@ulaval.ca

**Author Contributions:**

Conceptualization:  Adrian S. Monthony

Data Curation:  Adrian S. Monthony

Formal Analysis:  Adrian S. Monthony, Maxime de Ronne

Funding Acquisition: Adrian S. Monthony, Davoud Torkamaneh

Investigation: Adrian S. Monthony, Davoud Torkamaneh

Methodology: Adrian S. Monthony, Maxime de Ronne, Davoud Torkamaneh

Project Administration:  Adrian S. Monthony, Davoud Torkamaneh

Resources: Davoud Torkamaneh

Software: Adrian S. Monthony, Maxime de Ronne

Supervision: Davoud Torkamaneh

Validation: N/A

Visualization: Adrian S. Monthony

Writing- Original Draft Preparation: Adrian S. Monthony, Davoud Torkamaneh

Writing- Review& Editing: Adrian S. Monthony, Maxime de Ronne, Davoud Torkamaneh

**Data Availability Statement:**

All scripts and raw data used in this study are available from The Open Science Framework (OSF) under the DOI:  DOI 10.17605/OSF.IO/7G9VR at OSF.io. or from the NCBI, under previously published projects.

# Abstract

Sexual plasticity describes an organism's ability to change its phenotypic sex in response to environmental, chemical or physiological cues without a change in sex chromosome karyotype. *Cannabis sativa* L. (cannabis) is a medically important dioecious crop, with a pair of heteromorphic XY sex chromosomes. Cannabis has demonstrated sexual plasticity in response to chemical treatments which target the biosynthesis and signalling of ethylene. To date, ethylene biosynthesis and signalling has not been specifically studied in *C. sativa,* and the role of ethylene-related genes (ERGs) in sexual plasticity in the species has not been investigated. In the present study, we performed gene orthology analysis based on the model species *Arabidopsis thaliana* L. to reconstruct the *C. sativa* Yang Cycle, ethylene biosynthesis and ethylene signalling pathways, whose canonical composition appear conserved in cannabis. Additionally, we employ two transcriptomic datasets containing male, female and chemically induced male flowers to investigate patterns of expression for ethylene-related genes which may be involved in sexual plasticity. Comparison of these transcriptomes highlighted CsERGs whose differential expression was associated with sexual determination and those which we propose are associated with the sexual plasticity. These sexual plasticity ERGs could be classified into two patterns of expression: karyotype concordant (KC) and unique (uERG). We show that CsERGs involved in sexual plasticity are distributed throughout the genome and are not exclusive to the sex chromosomes, suggesting a broad control of sexual plasticity in *C. sativa*.

**Keywords:** ethylene, sexual plasticity, cannabis, STS, biosynthesis, sex expression, sex determination

# Introduction

   *Cannabis sativa* L. (cannabis) belongs to the estimated 5-6% of dioecious angiosperm species (Renner, 2014). Despite their rarity, dioecious plants provide an excellent opportunity to study the mechanisms involved in sex expression and the evolution of sex-determining regions and sex chromosomes. Cannabis (2n=20) is predominantly dioecious with a XY sex-determination system; however, monoecious cultivars of *C. sativa* have been bred, especially for the production of seeds and oil used in industrial hemp (Ferfuia *et al.*, 2021). Additionally, dioecious cannabis cultivars have been known to display hermaphroditism (Monthony *et al.*, 2021c), reversion from a sexual to vegetative state (Monthony *et al.*, 2021a), and sometimes complete changes in sex expression in response to certain environmental factors and chemical reagents (Ram & Jaiswal, 1972; Mohan Ram & Sett, 1982; Lubell & Brand, 2018; Moon *et al.*, 2020a). Taken together, these observations suggest that cannabis sex is plastic and not strictly controlled by its XX or XY chromosomal complement.

   Sexual plasticity is the ability of an organism to change its phenotypic sex in response to environmental or physiological cues without altering its sex chromosomes (Liu *et al.*, 2017). Vertebrates exhibit sexual plasticity on a continuum from genotypic sex determination (GSD) to environmental sex determination (ESD), with some organisms influenced by both (GSD + ESD; Lambert *et al.*, 2019). Non-mammalian vertebrates, such as fish (Senthilkumaran, 2015; Liu *et al.*, 2017) and frogs (Lambert *et al.*, 2019), as well as insects like aphids and bees (Morgan, 1909; Charlesworth & Mank, 2010), exhibit sexual plasticity triggered by diverse factors. The mechanisms responsible for these changes vary, but can involve gene expression changes related to hormone synthesis and signaling, germ cell development and maintenance as well as

microRNAs (Perry & Grober, 2003; Hayes *et al.*, 2010; Ashby *et al.*, 2016; Cardoso-Júnior *et al.*,

2017). For instance, social factors such as population density, mating competition, and hierarchical

structures can affect sexual plasticity in fish (Warner & Swearer, 1991; Liu *et al.*, 2017).

Amphibians can exhibit sexual plasticity under normal and extreme conditions (Lambert *et al.*,

2015, 2019), while in some insects, the mode of sexual reproduction can affect offspring sex

(Morgan, 1909; Charlesworth & Mank, 2010). Although the term *sexual plasticity* has largely been

used to describe fluid sex determination in the animal kingdom (Schlesinger *et al.*, 2010;

Senthilkumaran, 2015; Liu *et al.*, 2017), the present study highlights how sexual plasticity can also

be used to describe the interaction between GSD and ESD in dioecious plants, such as *C. sativa*.

Similar to amphibians, informal research conducted by enthusiast breeders prior to

legalization has demonstrated that chemical exposure can influence sexual plasticity in plants such

as *C. sativa*. (Green, 2005; Thomas, 2012). This pre-legalization exploratory work identified

silver-containing compounds such as silver thiosulfate (STS), a potent inhibitor of ethylene

signalling, as promoting the production of male flowers on female plants (induced male flowers;

IMF) and aqueous ethephon ((2-Chloroethyl) phosphonic acid), which rapidly decomposes to

produce ethylene gas, as promoting the production of female flowers on male plants (induced

female flowers; IFF) as shown in Figure 1. Following the legalization of cannabis in some

jurisdictions, the inverse effects of STS and ethephon on cannabis sex have been re-demonstrated,

quantified and published by a handful of research groups (Lubell & Brand, 2018; Kurtz *et al.*,

2020; Moon *et al.*, 2020a; Flajšman *et al.*, 2021), pointing to ethylene as being an effector of sexual

plasticity. Although the phenotypic outcomes of these treatments in cannabis have been quantified,

to date there are no studies which have sought to understand the underlying molecular mechanisms

controlling ethylene-related sexual plasticity in cannabis.

**Figure 1** Representative photos (unpublished data) of four floral phenotypes in *Cannabis sativa* (cv. La Rosca), grown at Université Laval. a) 4-week-old female flowers on a genetically female plant; b) 4-week-old STS-induced male flowers (IMFs) on a genetically female plant; c) 3-week-old male flowers on a genetically male plant and d) ethephon-induced female flowers (IFFs) on a genetically male plant. Scale bars = 5 mm.

Ethylene was first reported as a plant growth regulator in 1901 and has since been

extensively studied (Coulter & Barnes, 1901; Gane, 1934; Kende, 1998; Chang, 2016). It is

involved in various biological processes in plants, including seed germination, root development,

shoot and root growth, flowering, fruit ripening, leaf and fruit abscission, and responses to abiotic

and biotic stresses and senescence (Tieman *et al.*, 2000; Resnick *et al.*, 2006; Argueso *et al.*, 2007;

Yang *et al.*, 2015; Dhakal *et al.*, 2019; Binder, 2020). The biosynthesis pathways of ethylene are

highly conserved across the plant kingdom, and considerable time and resources have been

dedicated to its study  (Yang *et al.*, 2015). Ethylene is synthesized via a two-step pathway from S-

adenosyl methionine (SAM) derived from the Yang Cycle. The first step is catalyzed by 1-

aminocyclopropane-1-carboxilic acid (ACC) synthase (ACS), which converts SAM to ACC and

5'-methylthioadenosine (MTA; Argueso *et al.* 2007; Pattyn *et al.* 2021). ACS is encoded by a

multigene family in most plants, with the genes being differentially regulated depending on various

environmental and developmental factors (Argueso *et al.*, 2007; Dubois *et al.*, 2018). Arabidopsis

has eight identified ACS genes (Argueso *et al.*, 2007). The final step of ethylene biosynthesis is

catalyzed by ACC oxidase (ACO), which converts ACC to ethylene, cyanide, and $CO_2$ as by-

products (Argueso *et al.*, 2007; Houben & Van de Poel, 2019). Arabidopsis has an estimated five

ACO protein family members (Houben & Van de Poel, 2019). In cucumber, ACO plays a key role

in sex determination, with four ACO genes identified as rate-limiting in the biosynthetic pathway

(Chen *et al.*, 2016a). The genes involved in ethylene signaling are also highly conserved across

plant species and can be represented by a largely linear model, with recent studies finding

secondary non-canonical pathways that play a minor role  (Dubois *et al.*, 2018; Binder, 2020).

Ethylene receptors found in the endoplasmic reticulum (ER) are negatively regulated by the Golgi-

associated reversion to ethylene sensitivity 1 (RTE1; Zhou *et al.*, 2007) and an ER-localized family

of proteins called auxin-regulated genes involved in organ size (ARGOS; Rai *et al.* 2015; Shi *et*

*al.* 2016). In the presence of ethylene, these receptors, ethylene response sensor 1 (ERS1), ERS2,

ethylene resistance 1 (ETR1), ETR2, and ethylene insensitive 4 (EIN4) bind to ethylene to initiate

signaling (Dubois *et al.*, 2018). Ethylene functions as an inverse agonist, promoting further

ethylene production by altering downstream activity of the biosynthetic pathway and reducing the

inhibition of downstream signaling by lowering the activity of constitutive triple response 1

(CTR1) in the presence of a copper ion cofactor (Mayerhofer *et al.*, 2012; Binder, 2020). Inhibition

of CTR1 activity by ethylene results in dephosphorylation of EIN2 and cleavage of its c-terminal

domain (C-END), allowing for repression of EBF1/2 translation by binding to the 3' untranslated

regions of their mRNA (Chang, 2016; Dubois *et al.*, 2018; Binder, 2020). In the absence of

ethylene, EBF1/2 target the primary ethylene-responsive transcription factors EIN3 and EIN3-

LIKE (EIL1) for protein degradation (Dubois *et al.*, 2018). Repression of EBF1/2 by ethylene

allows for expression of EIN3 and EIN3-like transcription factors, which induce expression of

ethylene response factors (ERFs), promoting further ethylene biosynthesis. Although the canonical

signaling pathways are largely understood, the study of ERFs remains an area that requires further

investigation (Olsen *et al.*, 2015; Dubois *et al.*, 2018).

Compared with cannabis, the role of ethylene in sexual plasticity is much more well-

understood in the Cucurbitaceae family, where ethylene biosynthesis and signalling have also

been shown to be manipulated by STS or ethephon applications (Li *et al.*, 2021). In most

cucurbits, ethylene (usually applied as ethephon) has a feminizing effect, increasing the ratio of

female flowers relative to males, while the use of STS has the inverse effect, causing a

masculinizing effect and results in increased production of male flowers. Numerous studies of

sexual plasticity in cucurbits have pointed to the role of ethylene biosynthesis genes such as

*ACS4* and *ACS11*, as well as ethylene signalling genes such as the ethylene receptor *ETR1* as

contributing to sexual plasticity (Boualem *et al.*, 2015; Zhang *et al.*, 2017; Li *et al.*, 2020). The

manipulation of cucumber sex has allowed growers to control and optimize the ratio of female to

male flowers to improve yield and to control production and labour constraints (Oda *et al.*,

2022). In cannabis, control of plant sex is important as medicinally valuable cannabinoids are

primarily produced in female flowers. However, without a pathway model for ethylene

biosynthesis and signalling in the species, study of cannabis sexual plasticity remains impossible.

The highly conserved nature of ethylene biosynthesis and signalling and well-characterized

pathways in model species *Arabidopsis thaliana* present a valuable starting point for the

reconstruction of the pathway in an under-researched plant such as *C. sativa* where ethylene

biosynthesis and signalling has yet to be explored. In the current study, we use homology

analysis and RNA-seq data to present the first *C. sativa* putative Yang cycle, ethylene

biosynthesis and ethylene signalling pathways, which will serve as an important foundation for

expanding our understanding of the role of ethylene in sexual plasticity of this under-researched

dioecious species.

## Materials & Methods

### *In silico* identification of ethylene-related gene orthologs in *C. sativa*

*Arabidopsis thaliana* was chosen as the model from which a list of genes related to ethylene

biosynthesis and signalling, as well as the production of ethylene precursors derived from the Yang

cycle were compiled following a review of the literature (Pommerrenig *et al.*, 2011; van de Poel

*et al.*, 2012; Olsen *et al.*, 2015; Chang, 2016; Pattyn *et al.*, 2021). The resulting list of *A. thaliana*

ethylene-related genes (AtERGs; **Error! Reference source not found.**) were used for the

identification of orthologous *C. sativa* ERGs (CsERGs). Nucleotide and protein sequences for

AtERGs were obtained from the National Center for Biotechnology Information database (NCBI;

Bethesda, MD). Ortholog identification was conducted on the *C. sativa* cs10 reference genome

(RefSeq assembly accession: GCF_900626175.2; Grassa *et al.*, 2021). A list of putative CsERG

orthologs was compiled using two different methods. The first method employed OrthoFinder

(Emms & Kelly, 2019) adapted from Carey *et al.* (2021) and the second method consisted of a

BLASTP search of each individual AtERG against the cs10 reference genome (Altschul *et al.*,

1990), followed by a manual selection of putative orthologs from the of BLASTP result adapted

from Barcaccia *et al.* (2020). This method is described in further detail in the **Error! Reference**

**source not found.**. The chromosome map (Figure 4a) was generated with chromoMap (v4.1.1;

Anand & Rodriguez Lopez, 2022) and bar plots of gene distribution (Figure 4b) was generated

with ggplot2 (v3.4.1; Wickham, 2016) using the colour palette Spectral from RColorBrewer

(Brewer *et al.*, 2002)

## RNA-seq data collection and processing

The RNA-seq reads from Adal *et al.* (2021) and Prentout *et al.* (2020), hereafter referred

to as dataset 1 and dataset 2, were downloaded from the Sequence Read Archive (SRA) on NCBI

under the BioProject accessions PRJNA669389 and PRJNA549804, respectively. RNA was

isolated from male flowers (MFs), female flowers (FFs) and STS-induced IMFs for dataset 1, and

from MFs and FFs for dataset 2. To perform a relevant comparison between the 2 datasets, only

data from similar tissues were considered, i.e., the young flower buds. The SRA sample accessions

used in this study are indicated in **Error! Reference source not found.**. FastQC (Andrews, 2010)

was used for validating the quality of the data. Then, reads were aligned to the *C. sativa* cs10

reference genome (version GCF_900626175.2; Grassa *et al.*, 2021) using STAR (v2.7.2b; Dobin

*et al.*, 2013) as reported in Cai *et al.* (2020). The STAR index was built with the –

sjdbOverhang=100 and the genomeSAindexbases=13 arguments. Splice junctions from the *C.*

*sativa* cs10 RefSeq assembly GTF version GCF_900626175.2 were provided via the --sjdbGTFfile

argument. STAR alignments were generated using the default parameters including the --

outSAMtype BAM SortedByCoordinate arguments. Quality control of the mapped reads was

performed using Qualimap (v2.2.1; Okonechnikov *et al.*, 2015). One IMF sample from dataset 1

was excluded based on abnormal distribution of GC-content among mapped reads which included

large shoulders at 30% and 60 % GC content, indicating contamination of the sample (see **Error!**

**Reference source not found.**). Finally, quantification of reads overlapping genetic features was

performed with HTSeq (v2.0.1; Putri *et al.*, 2022) based on the cs10 GFF file obtained from NCBI

([https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/900/626/175/GCF_900626175.2_cs10/GCF_9006](https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/900/626/175/GCF_900626175.2_cs10/GCF_9006)

[26175.2_cs10_genomic.gff.gz](https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/900/626/175/GCF_900626175.2_cs10/GCF_900626175.2_cs10_genomic.gff.gz)) in order to generate a gene quantification table required for

subsequent differential analysis with DEBrowser (v1.22.5; Kucukural *et al.*, 2019).

## Differential expression analysis of CsERGs

Differentially expressed CsERGs were identified using DESeq2 (Love *et al.*, 2014) with a

FDR-adjusted *p*-value < 0.05 on the DEBrowser  graphical user interface in RStudio (RStudio

Inc., MA, USA) using v4.1.1 of R. The analysis was adapted from DEBrowser's DESeq2 method

outlined on the DEBrowser Github ([https://github.com/UMMS-Biocore/debrowser](https://github.com/UMMS-Biocore/debrowser)). In short, gene

quantification tables generated in HTSeq and a manually generated metadata file for the samples

were uploaded to DEBrowser. Genes with count per million values less than 1 (CPM < 1) in 7 out

of 15 replicates in dataset 1 and 6 out of 12 replicates in dataset 2 were removed. Replicates were

then grouped and pairwise comparisons were performed based on the floral type. In DEBrowser

the following DESeq2 parameters were used following the method outline by Kucukural *et al.*

(2019) and the DEBrowser Github: *Fit Type- parametric; betaPrior- FALSE; Test Type- Wald;*

*Shrinkage- None.* Differential expression levels were then normalized using a Median Ratio

Normalization (MRN) and pairwise comparisons between treatments were considered

significantly different in subsequent comparisons if they had an adjusted *p*-value <0.05 according

to the Wald test and a |log2foldchange| > 1. Heatmaps comparing differential gene expression were

then generated using DEBrowser heatmap function. Tables from pairwise comparisons were

generated from DEBrowser and then used for the generation of gene expression bar plots using

ggplot2 (Wickham, 2016) in RStudio.

# Results

## ERGs are largely conserved in *C. sativa*

Based on 47 AtERGs previously identified (**Error! Reference source not found.**),

ortholog analyses identified 43 putative CsERGs (Table 1). At least one ortholog of each key

gene in the canonical ethylene biosynthesis and signalling pathways, as well as the Yang cycle,

was identified *in silico* except for auxin-regulated genes involved in organ size (AtARGOS;

**Error! Reference source not found.**). To our knowledge this represents this first time that the

genes coding for the various enzymes involved in the putative Yang cycle (Figure 2a), ethylene

biosynthesis (Figure 2b) and signalling pathway (Figure 3) have been put forward in *C. sativa.*

**Table 1** *Cannabis sativa* ethylene-related genes (CsERGs) identified through ortholog analysis.

| Cs10 ERG Locus[‡] | Name | Chromosome | *A. thaliana* homologs | Pathway | Identification method |
|---|---|---|---|---|---|
| LOC115704991 | CsACO1 | 1 | AT2G19590 | Biosynthesis | BLASTP, OrthoFinder |
| LOC115702109 | CsACO1-LIKE1 | Un[§] | AT2G19590 | Biosynthesis | BLASTP, OrthoFinder |
| LOC115707397 | CsACO2[*†] | 1 | AT1G62380, AT1G12010, AT1G05010 | Biosynthesis | BLASTP, OrthoFinder |
| LOC115699412 | CsACO3[*†] | 8 | AT1G62380, AT1G12010, AT1G05010 | Biosynthesis | BLASTP, OrthoFinder |
| LOC115699408 | CsACO5[*†] | X | AT1G77330 | Biosynthesis | BLASTP, OrthoFinder |
| LOC115704849 | CsACS1 | 1 | AT3G49700, AT4G08040, AT4G11280, AT1G01480 | Biosynthesis | BLASTP |
| LOC115717044 | CsACS2[*†] | 5 | AT4G08040, AT1G01480, AT4G11280, AT3G49700 | Biosynthesis | BLASTP, OrthoFinder |
| LOC115712396 | CsACS3 | 4 | AT5G65800, AT4G37770, AT4G08040, AT3G49700 | Biosynthesis | BLASTP, OrthoFinder |
| LOC115704963 | CsACS7-LIKE | 1 | AT4G26200, AT1G01480, AT4G11280, AT4G08040, AT3G49700 | Biosynthesis | BLASTP, OrthoFinder |
| LOC115696400 | CsACS9[†] | X | AT4G08040, AT5G65800, AT4G37770, AT3G49700 | Biosynthesis | BLASTP |
| LOC115716540 | CsACS9-LIKE | 5 | AT5G65800, AT4G37770, AT4G08040, AT3G49700, AT2G22810 | Biosynthesis | BLASTP |
| LOC115697744 | CsCTR1[*†] | X | AT5G03730 | Signalling | BLASTP |
| LOC115706236 | CsCTR1-LIKE[*†] | 1 | AT5G03730 | Signalling | BLASTP, OrthoFinder |
| LOC115720532 | CsEBF1[*†] | X | AT5G25350, AT2G25490 | Signalling | BLASTP, OrthoFinder |
| LOC115724585 | CsEBF2[*†] | 6 | AT5G25350, AT2G25490 | Signalling | BLASTP |
| LOC115711236 | CsEIL1[*†] | X | AT2G27050 | Signalling | OrthoFinder |
| LOC115719373 | CsEIL3[*†] | 2 | AT1G73730 | Signalling | OrthoFinder, BLASTP |
| LOC115709621 | CsEIN2[*†] | 3 | AT5G03280 | Signalling | BLASTP, OrthoFinder |
| LOC115706079 | CsEIN3[*†] | 1 | AT2G27050, AT3G20770 | Signalling | BLASTP, OrthoFinder |
| LOC115723050 | CsERF1 | 9 | AT3G23240 | Signalling | BLASTP, OrthoFinder |
| LOC115697291 | CsERS1[*†] | 7 | AT2G40940 | Signalling | BLASTP, OrthoFinder |
| LOC115705474 | CsETP1[*†] | 1 | AT3G18980 | Signalling | BLASTP |
| LOC115720323 | CsETP2 | 2 | AT3G18910 | Signalling | BLASTP |
| LOC115702329 | CsETR1[*†] | X | AT1G66340, AT2G40940 | Signalling | BLASTP, OrthoFinder |
| LOC115721785 | CsETR2[*†] | 9 | AT3G04580, AT1G04310, AT3G23150 | Signalling | BLASTP, OrthoFinder |
| LOC115698184 | CsRAN1[*†] | 7 | AT5G44790 | Signalling | BLASTP, OrthoFinder |
| LOC115707134 | CsRTE1[*†] | X | AT2G26070 | Signalling | BLASTP, OrthoFinder |

| Cs10 ERG Locus[‡] | Name | Chromosome | *A. thaliana* homologs | Pathway | Identification method |
|---|---|---|---|---|---|
| LOC115721731 | CsARD1[*†] | 9 | AT5G43850, AT4G14716, AT4G14710, AT2G26400 | Yang Cycle | BLASTP, OrthoFinder |
| LOC115722194 | CsARD2[*†] | 9 | AT4G14716, AT2G26400, AT5G43850, AT4G14710 | Yang Cycle | BLASTP, OrthoFinder |
| LOC115706479 | CsMTI1[*†] | 1 | AT2G05830 | Yang Cycle | BLASTP, OrthoFinder |
| LOC115703014 | CsMTK1[*†] | X | AT1G49820 | Yang Cycle | BLASTP, OrthoFinder |
| LOC115702983 | CsMTK1-LIKE[*†] | X | AT1G49820 | Yang Cycle | BLASTP, OrthoFinder |
| LOC115704327 | CsMTN[*†] | X | AT4G34840, AT4G38800 | Yang Cycle | BLASTP, OrthoFinder |
| LOC115706293 | CsDEP1[*†] | 1 | AT5G53850 | Yang Cycle | BLASTP, OrthoFinder |
| LOC115711364 | CsSAM1[*†] | X | AT1G02500, AT4G01850, AT3G17390, AT2G36880 | Yang Cycle | BLASTP, OrthoFinder |
| LOC115719259 | CsSAM2[*†] | 2 | AT1G02500, AT4G01850, AT3G17390, AT2G36880 | Yang Cycle | BLASTP |
| LOC115711343 | CsSAM3[*†] | 3 | AT2G36880, AT3G17390, AT4G01850, AT1G02500 | Yang Cycle | BLASTP, OrthoFinder |
| LOC115700576 | CsFYF1[*†] | 8 | AT5G62165 | Ethylene Adjacent | BLASTP |
| LOC115706939 | CsFYF2[*†] | 1 | AT5G62165 | Ethylene Adjacent | BLASTP |
| LOC115716986 | CsFYF3[*†] | 5 | AT5G62165 | Ethylene Adjacent | BLASTP |
| LOC115717309 | CsFYF4[*†] | 5 | AT5G62165 | Ethylene Adjacent | BLASTP |
| LOC115701489 | CsORE1[*†] | 8 | AT5G39610 | Ethylene Adjacent | BLASTP, OrthoFinder |
| LOC115709772 | CsORE2[*†] | 3 | AT5G39610 | Ethylene Adjacent | BLASTP |

202 ‡ Hyperlinks to the NCBI cs10 reference genome information page for the specific locus.

203 * and † ERGs for which transcripts were detected from dataset 1 and/or in dataset 2, respectively.

204 § A gene located on an unclassified scaffold.

**Figure 2** The *C. sativa* Yang cycle and ethylene biosynthesis pathway determined by homology analysis with *A. thaliana* genome. (a) Yang cycle; abbreviations: *S*-adenosylmethionine (SAM), 5-methylthioadenosine (MTA), 5-methylthioribose (MTR), 5-methyltioriobose-1-P (MTR-1-P), 5-methylthioribulose-1-P (MTRu-1-P), 1,2-dihydroxy-3-keto-5-methylthiopentene (DHKMP), 2-keto-4-methylthiobutyrate (KMTB), methionine (Met). (b) Ethylene biosynthesis pathway.

**Figure 3** The *C. sativa* ethylene signalling pathway inferred by homology analysis with *A. thaliana* genome. The pathway was elucidated using the canonical pathway from *A. thaliana*, genes from *A. thaliana* that were not identified in *C. sativa* are outlined by a dashed line. Figure design was based on existing ethylene signalling models (Binder *et al.*, 2010; Dubois *et al.*, 2018; Binder, 2020).

## ERGs are distributed genome-wide in *C. sativa*

The 43 CsERGs found in the homology analysis were distributed throughout the genome

(Figure 4 and **Error! Reference source not found.**). A majority of the CsERGs were located on

autosomes (Figure 4b). Yang cycle genes were the most numerous CsERGs present on the sex

chromosomes, with 40% of the identified *C. sativa* Yang cycle genes located on the X

chromosome. In contrast, none of the genes that were classified as ethylene adjacent (ERGs not

belonging to the canonical ethylene biosynthesis or signaling pathways) were located on the sex

chromosomes (Figure 4). Of the 43 CsERGs identified all mapped to chromosomes in the reference

genome, with the exception of  CsACO1-LIKE1, which mapped to an unassigned scaffold.

**Figure 4** Position (a) of ERGs in the cannabis cs10 reference genome and the proportional frequency (b) of those genes per chromosome. An interactive version of (a) can be found in the supplemental materials or on the OSF page for this project.

## Differential expression analysis reveals three patterns of expression

Among the 43 putative CsERGs, transcripts of 35 of these genes were expressed in both dataset 1 and dataset 2 (Table 1). Of these, transcripts of 15 CsERGs in dataset 1 and 11 CsERGs in dataset 2 showed differential expression between flower types (**Error! Reference source not found.**). In 7 ERGs common to both datasets, differential expression between MFs and FFs was of similar magnitude (**Error! Reference source not found.**).

**Figure 5** Heatmaps of differentially expressed CsERGs (identified by locus ID; **Table 1**) for dataset 1 (a, c and d) and dataset 2 (b). X-axis labels are the flower type, followed by the last two digits of the SRA accession numbers, which can be found in **Error! Reference source not found.**. Y-axis contains the CsERG names. Heatmap scale is in log of the fold change in expression for the pairwise comparison of the flower type listed on the x-axis: a) MF vs. FF

comparison in dataset 1; b) MF vs. FF comparison in dataset 2; c) IMF vs. FF in dataset 1; d)
IMF vs. MF in dataset 1.

Differential expression analyses of CsERGs in FF, MF and IMF tissues from dataset 1

reveal three distinct patterns of expression which can be used to create a framework for thinking

of ethylene-related sexual plasticity in cannabis. We have named these cases of differential

expression as follows: karyotype concordant (KC; Figure 7), floral-organ concordant (FOC; Figure

8) and unique ERG (uERG; Figure 9).

**Karyotype concordant (KC) expression**

This first case, karyotype concordant (KC) expression, describes expression patterns where

CsERG expression is the same between plants with the sex chromosome karyotype (i.e., FF and

IMF). KC genes make up the largest number of genes identified by differential analysis of dataset

1, making up nine genes in total (Figure 7). KC expression represents cases where the pairwise

comparison between FF and IMF gene are not significantly different, but where the IMF

significantly differs from its biologically male counterpart (MF). In this case of KC expression,

both FF and IMF have XX chromosomes and normalized gene counts for a given ERG do not

significantly differ between FF and IMF individuals.

**Figure 6** Differential analysis of *C. sativa* ERGs revealed nine genes with a karyotype concordant (KC) expression pattern. This describes expression patterns shared between organisms with the same sex chromosome karyotype, but not the same sex expression. In these cases, female flowers (FF; XX karyotype) and induced male flowers (IMF; XX karyotype) showed KC expression.

### Floral organ concordant expression (FOC)

In this second case, genes from organisms sharing the same floral organs show similar patterns of expression. From the 15 differentially expressed CsERGs identified in dataset 1, five were identified as belonging to this category (Figure 8). In these five cases, there was no differential expression between male-flowers: i.e., IMFs and MFs. However, in all cases expression significantly differed between MFs and FFs, showing a pattern of expression that is conditioned by the floral organs that are present.

**Figure 7** Differential analysis of *C. sativa* ERGs revealed five cases of genes showing floral-organ concordant (FOC) expression. FOC refer to genes whose expression pattern is shared between organisms with the same floral organ, but not necessarily the same sex-chromosome karyotype designated by ns in transcription levels for male flowers on XY plants and induced male flowers on XX plants.

### Unique ERG (uERG) expression

In this final case, expression of CsERGs in the sex-induced plant (IMF) is not different from plants with shared sex-organs (MF vs IMF) or those with a like sex karyotype (FF vs IMF). Two CsERGs were identified with this pattern of expression.

**Figure 8** Differential expression analysis of dataset 1 revealed 2 cases of unique ERG (uERG) expression, which are cases where gene expression in the sex-changed plant does not match either that of the plant with which is shares the same sex chromosome karyotype or the plant with which is shares the same floral organs.

# Discussion

To date, sexual plasticity has been more widely studied in the animal kingdom, whereas in

angiosperms most efforts to understand plant sexual plasticity has been focused on the

Cucurbitaceae family (Yamasaki *et al.*, 2001; Boualem *et al.*, 2015; Oda *et al.*, 2022). Studies of

dioecious sexual plasticity are limited, but *C. sativa* is one dioecious species where sexual

plasticity has been frequently observed, both in nature and in research settings, and it is a species

for which methods inducing sexual plasticity have been established (Green, 2005; Lubell & Brand,

2018; Kurtz *et al.*, 2020; Moon *et al.*, 2020a; Flajšman *et al.*, 2021; Zheng, 2022). To date, these

reported occurrences of cannabis sexual plasticity have linked the phenomenon to the plant growth

regulator ethylene. Existing studies show that inhibition of ethylene in female plants causes the

induction of male flowers (Lubell & Brand, 2018) and the treatment of male plants with ethylene

causes the induction of female flowers on male plants (Moon *et al.*, 2020a).

Both STS and ethephon affect endogenous ethylene concentrations in plants. STS is

hypothesized to inhibit ethylene signaling by interacting with the ethylene receptor on the

endoplasmic reticulum, reducing the plant's ability to perceive ethylene (Binder, 2020). The exact

mode of inhibition is debated, with two prevailing ideas: in one, silver binds to the receptor's

copper ion cofactor site to competitively inhibit signaling (Altman & Solomos, 1995; Rodríguez

*et al.*, 1999); in the other, silver ions non-competitively inhibit the receptors from binding ethylene,

possibly by binding the active site or reducing the number of binding sites available to ethylene

(Beyer, 1976; Blat, 2010; McDaniel & Binder, 2012). Ethephon, on the other hand, increases

endogenous ethylene concentrations by breaking down to produce ethylene when applied to plant

tissues (Zhang & Wen, 2010). It triggers an autocatalytic increase in ethylene biosynthesis

throughout the plant (Hume & Lovell, 1983a).

Studying the role of ethylene in sexual plasticity is an obvious choice for understanding

sexual plasticity in dioecious plants, due to its well-known and consistent biosynthesis and

signaling pathways, as well as its proven impact on sexual plasticity in the Cucurbitaceae family

(Hume & Lovell, 1983b; Manzano *et al.*, 2014; Cebrián *et al.*, 2022). However, given the

numerous roles of ethylene in plants, untangling the specific ways with which ethylene affects

sexual plasticity also faces some challenges. Here, we presented a framework of three putative

CsERG expression patterns, two of which we propose are linked to sexual plasticity in cannabis.

We also described the first report of the *C. sativa* Yang cycle and ethylene biosynthesis and

signalling pathways and discuss which key genes in these pathways belong to sex plastic patterns

of expression. We highlight how the distribution of these CsERGs whose expression patterns are

associated with sexual EGRs are not solely located on sex chromosomes, but rather distributed

throughout the genome. Lastly we end by discussing limitations and future opportunities revealed

by this study.

## Analyses help distinguish sexual plasticity genes from deterministic

## genes

*C. sativa* transcriptomes analyses confirmed the expression of 35 of the 43 CsERGs

identified in this study (datasets 1 and 2). Additionally, the analysis helped identify candidate sex

plastic CsERGs, following the production of IMF using the known ethylene inhibitor STS (Dataset

1). Analyses comparing CsERGs in MF, FF and IMF revealed three sex-linked patterns of

expression, which have been named as follows: karyotype concordant (KC; Figure 7), floral organ

concordant (FOC; Figure 8) and unique ERG (uERG; Figure 9). Two of these three patterns are of

particular interest (FOC and UERG), and below we outline why we believe that CsERGs with

these patterns are involved in sexual plasticity of *C. sativa*.

The largest number of CsERGs belong to the KC case, containing a total of 9 genes (Figure 7). In

this case, CsERG expression is similar between individuals with the same sex chromosome

karyotype. The findings from the current analysis suggest that KC CsERG expression is unchanged

in XX plants regardless of the sex organs the plant has. As such, we suggest that genes belonging

to the KC case may have sex-specific patterns of expression, but these are unaffected by STS

treatments which induces sexual plasticity in *C. sativa*. Taken together, we propose that KC genes

are non-sex-plastic.

In FOC expression (Figure 8), the CsERG expression is the same between individuals with shared

phenotypic sex but carrying different sex chromosome karyotype (i.e., MF and IMF). Genes with

FOC had no significant difference in expression levels between individuals producing male

flowers, regardless of their genotypic sex (XX or XY sex chromosome karyotype). However, FOC

gene expression in IMFs were significantly different in FF. A total of 5 CsERGs were identified

as FOC. Three genes, *CsACS2*, *CsFYF2* and *CsFYF4* showed higher gene expression in FF than

the phenotypic male (MF and IMF) plants, while two genes (*CsETP2* and *CsMTN*) had higher

expression in the MF and IMF plants than in the FF plants (Figure 8).

In the last case, UERG (Figure 9), we observed a differential expression of CsERGs between all

three treatments. Two genes, *CsFYF1* and *CsACO5* showed UERG, with the sex manipulated

plant's (IMF) gene expression not matching either of the plant with the same genotypic sex (MF)

or the plant with the phenotypic sex (MF). We hypothesize that FOC and UERG  are indicative

that the genes is a sex-plastic ERG.

**Multiple *CsACS* and *CsACO* genes; a potential regulatory framework**

**for sexual plasticity**

An expansive body of literature exists which has established a canonical, highly conserved

ethylene biosynthesis, and signalling pathway in a number of widely studied plant species (Binder,

2020; Pattyn *et al.*, 2021). Ethylene biosynthesis and signalling has yet to be studied in cannabis

or closely related dioecious species, making *A. thaliana* a logical choice for the first attempts at

assembling the canonical *C. sativa* ethylene biosynthesis and signalling pathways. Ethylene

biosynthesis is regulated by a set of conserved enzymes that are found in all plant species. The key

enzyme in this two-step process is ACC synthase (ACS), which converts SAM, derived from the

Yang cycle, into 1-aminocyclopropane-1-carboxylic acid (ACC). ACC is then converted into

ethylene by the enzyme ACC oxidase (ACO). These enzymes are conserved across all plant

species, and their regulation is tightly controlled by various factors such as hormonal and

environmental cues (Bleecker & Kende, 2000). Ortholog analysis conducted in this study showed

that both the Yang cycle, responsible for SAM production and ethylene biosynthesis are conserved

in *C. sativa*, with orthologs identified for each key step in these two pathways (**Figure 2**). In the

present study, we found that in cannabis, *ACS* and *ACO* belong to multi-gene families like in

Arabidopsis. In *C. sativa*, six *ACS* and five *ACO* gene homologs were identified from the eight

*ACS* and five *ACO* genes in *A. thaliana* respectively (**Figure 2**). The presence of *ACS* and *ACO*

are critical to biosynthesis, but studies have shown that the number of genes in each family can

vary between species. For example, in corn (*Zea mays* L.) 5 *ACS* and 15 *ACO* genes have been

identified, in rice (*Oryza sativa* L.) those numbers are 6 and 9 respectively and in tomato (*Solanum*

*lycopersicum* L. Karst. ex Farw) there are 9 *ACS* and 7 *ACO* genes (Houben & Van de Poel, 2019;

Wu *et al.*, 2022). The importance of having multiple functionally redundant ethylene biosynthesis

genes has been demonstrated in multiple plants, with studies showing *ACS* and *ACO* genes are

expressed at different levels and in different tissues as a function of the plant organ's needs and

developmental stage. In ripening tomatoes van de Poel *et al.* (2012) reported a higher level of

*ACS6, ACO2* and *ACO4* expression during baseline (system 1) ethylene production during

immature fruit development prior to ripening, but that auto-catalytic increases in ethylene during

ripening (system 2) were correlated with increases in *ACS2*, *ACS4*, *ACO3* and *ACO5* expression,

while abundance of *ACO2* and *ACO4* transcripts decreased at that same stage. The

induction/repression of specific *ACS* genes has also been shown to play a role in sexual plasticity

of some species in the *Cucurbitaceae* species (Manzano *et al.*, 2014; Boualem *et al.*, 2015; Chen

*et al.*, 2016a; Zhang *et al.*, 2017). One study found that expression of *ACS11* was only observed

in the vascular bundles of female flowers of monoecious and in hermaphroditic flowers of

andromonoecious plants (plants producing male and hermaphrodite flowers), pointing to the role

of *ACS11* in carpel development (Boualem *et al.*, 2015). Sexual plasticity has also been shown in

cucumber and melon by way of disruption of a specific *ACO* genes (*ACO2* in cucumber; *ACO3* in

melon) resulting in decreased ethylene production in the carpel region of flowers and causing a

shift from a monoecy to androecy resulting in unisexual flower production in cucumber and

melons (Chen *et al.*, 2016a).

The existence of multiple, functionally redundant *ACS* and *ACO* genes, combined with

spatial and temporal patterns of expression allows for the fine tuning of ethylene biosynthesis and

giving a plant a high level of control over its development, including effects on a plant's sexual

plasticity. In the present study, we demonstrate that *C. sativa* also contains numerous *ACS* and

*ACO* genes, spread across its genome (Table 1, Figure 4), which leads us to hypothesize that their

expression and regulation is also controlled spatially and temporally in cannabis. In particular,

*CsSAM3, CsACS2, CsACO3* and *CsACO5* showed distinct expression cases (Figures Figure

**7**,**Figure 8** and**Figure 9**) suggesting these CsERGs are linked to sexual plasticity. *CsACS2* shows

FOC expression, linking it to sexual plasticity and is responsible for the first committed step in

ethylene biosynthesis. It has been demonstrated to catalyze the rate-limiting reaction in most cases

of ethylene production (van de Poel *et al.*, 2012; Houben & Van de Poel, 2019; Pattyn *et al.*, 2021).

This gene is downregulated in IMFs in comparison to female plants, and shows similar patterns of

expression to male plants with whom they share the same floral organ despite having an XX

karyotype. Since *ACS* is thought to be rate-limiting, and given the association of ethylene with

'feminization' of floral phenotypes we hypothesize that the downregulation of *CsACS2* in IMF

relative to FF, points to reduction in ethylene biosynthesis *in planta* which may play a role in the

induction of male flowers in genetically female plants.

*CsACO5* has a unique expression pattern in all three forms of sex expression (UERG;

Figure 9), with a significantly larger expression in IMFs compared with MFs or FFs (Figure 9).

Here, dosage compensation could play a role with downstream inhibition of ethylene signalling by

STS leading to increased transcription of *ACO* as IMF plants which are genetically female, fail to

perceive ethylene levels normally required for female floral development. Lastly, *CsSAM3*

(produces SAM required for the biosynthesis of ethylene) and *CsACO3* appear to be linked to

genotypic sex and uninfluenced by the phenotypic sex changes (KC expression; Figure 7) induced

by STS, with activity in XY plants much higher for both than in plants with the XX chromosomal

complement. Further study should be performed to investigate if this KC  expression pattern is

unique to floral tissues or if this karyotype specific pattern of expression is also present in

vegetative tissues of flowering plants.

Ethylene is known to affect the sexual plasticity of many plants (Hume & Lovell, 1983b;

Hyde *et al.*, 2020), and the present study found patterns of differentially expressed CsERGs

between male and female plants, suggesting their implication in cannabis sexual plasticity.

However, CsERGs that did not show a difference in expression between male and female plants

may have limited or no impact on sexual plasticity at this stage of development, although their

regulation may fluctuate before or during floral induction. The study identified 43 CsERGs in the

cs10 reference genome but only detected transcripts for 35 genes in two transcriptomic datasets,

possibly due to some genes being unexpressed during the specific time points when the floral tissue

was collected in their respective studies. The spatial and temporal expression of genes during floral

development is well (Trivellini *et al.*, 2016), and future studies should examine gene expression

before and during flowering in response to ethylene and its agonists to better understand the

temporal control of sexual plasticity by ethylene. The identification of the *C. sativa* biosynthesis

pathway is an important first step toward developing targeted functional studies to understand the

role of these biosynthesis genes during sex determination in cannabis.

## Ethylene signalling is conserved in *C. sativa*

Like ethylene biosynthesis, ethylene signalling has been shown to be highly conserved

across the plant kingdom (Chang, 2016). Ethylene binds to a family of receptors known as ethylene

receptors (ETRs), which are found in the membrane of the endoplasmic reticulum of plant cells

(Binder, 2020). These receptors are responsible for transducing the ethylene signal to downstream

signalling components. The downstream signalling pathway involves a series of conserved

components, including *CTR1*, *EIN2*, and *EIN3*, which are responsible for activating the expression

of ethylene-responsive genes (Shakeel *et al.*, 2013). In the present study, we were able to identify

the canonical ethylene pathway genes in *C. sativa* responsible for each step of the signalling

pathway, making this the first pathway model for the species to date (Figure 3).

The ethylene signalling pathway is of particular interest in the current study as potentially

playing a role in cannabis sexual plasticity that is induced by the action of STS, an inhibitor of

ethylene signaling. Treatment of genotypically female plants results in male floral phenotypes (i.e.,

IMFs). Transcriptomes of MF, FF and IMF in this study revealed numerous CsERGs

upstream/downstream of or directly involved in the signalling pathway which have FOC

expression (*CsETP2, CsFYF2* and *CsFYF4*) and UERG (*CsFYF1*). In *Arabidopsis EIN2*

*TARGETING PROTEIN1 and 2* (*ETP1/2*) acts to suppress ethylene responses in the signalling

pathway by carrying out ubiquitinating with Skp1 Cullen F-box E3 ubiquitin ligase (*SCF E3*) of

*EIN2*, marking *EIN2* for proteosome degradation (Qiao *et al.*, 2009; Binder, 2020). Conversely,

ethylene acts as a suppressor of *ETP1/2*, thereby allowing ethylene signalling to proceed by

preventing the *ETP1/2* mediated degradation of *EIN2* (Qiao *et al.*, 2009; Yang *et al.*, 2015). In the

present study, both *CsETP1* and *CsETP2* transcripts were detected, however only *CsETP2* showed

sex-plastic differential expression. *CsETP2* expression was lowest in FFs compared to MFs and

IMFs (Figure 8). There was no significant difference between *CsETP2* expression in MFs and

IMFs, despite IMFs sharing the same sex chromosomes as FF plants. These patterns of gene

expression fit with the hypothesized role of ethylene as a feminizing agent. If increased

endogenous ethylene is responsible for femaleness, then we should expect lower levels of *CsETP2*

in female flower producing plants, as observed.

438   The differential expression analysis also revealed three *FOREVER YOUNG FLOWER*

(*FYF*) genes which belong to the FOC expression (*CsFYF2* and *CsFYF4*) and uERG (*CsFYF1*)

expression patterns. FYF genes are expressed in young flowers prior to pollination (Chen *et al.*,

2011) and acts as a suppressor of floral abscission by inhibiting ethylene triggered responses

downstream of the signalling pathway (Chen *et al.*, 2022). FYF does this by repressing *Ethylene*

*Response DNA-binding Factors (EDFs)* which are downstream *Ethylene-Responsive Binding*

*Factors* (*ERFs*) that trigger floral senescence and abscission in maturing flowers (Chen *et al.*,

2015). Similar patterns of expression were observed for *CsFYF1*, *CsFYF2* and *CsFYF4* in the

present study. In these three genes expression was highest by several orders of magnitude in FF

compared to MF and IMF tissues. Typically expression of these genes is highest in early flowering

stages and diminishes over time, and it has also shown that different FYF genes are expressed

differentially depending on tissues (Chen *et al.*, 2021). However, variations in FYF transcription

linked to sex expression have not yet been investigated. These findings suggest that in dioecious

plants FYF expression may vary depending on sex expression, however it remains to be seen if

this is a cause or effect of sex plasticity. Alternatively, the reason for high levels of FYF in FF

tissues relative to MF and IMF might also be explained by differences in the lifespan of female

and male cannabis flowers, the latter which have a much shorter life span and which would

encounter senescence sooner. Since FYF is a suppressor of senescence, the presence of lower

levels of *CsFYF1/2/4* in MF and IMF flowers may be due to earlier senescence or later

developmental stages of the flower compared with FF tissues. An alternative hypothesis for this

observed difference could be that higher anticipated levels of ethylene in female plants and flowers

may result in more downstream expression of ethylene response factors such as the *EDF*s

associated with floral senescence, and in order to prevent premature senescence caused by

ethylene, female flowers required higher endogenous concentrations of FYF.

## Distribution of CsERGs is not limited to sex chromosomes

*Cannabis sativa* contains a pair of heteromorphic sex chromosomes, XX for female

individuals and XY for males. However, treatment of male plants with ethylene and female plants

with the ethylene inhibitor STS has been reported to cause plants to produce flowers opposite than

that implied by the genotypic sex of the plant (Green, 2005; Lubell & Brand, 2018; Moon *et al.*,

2020a). This sexual plasticity suggests that the sex-expression of cannabis is not solely influenced

by the sex-chromosome complement. However there exist few studies in cannabis which explore

the link between sex chromosomes and sex-expression, and to the author's knowledge, no studies

which explicitly explore the link between cannabis sexual plasticity and ethylene. In a study

comparing the transcriptomes of male and female flowers, Prentout *et al.* (2020) reported that 35%

of sex-linked genes mapped to autosomes while 65% mapped to the sex chromosomes of *C. sativa*.

The distribution of their reported sex-linked genes in the chromosome is in stark contrast to the

distribution of CsERGs that we identified with only 26.2% of CsERGs mapping to the sex

chromosomes. The authors suggest that mapping of their sex-linked genes to autosomes was the

result of false positives caused by permissive mapping or incorrect placement of gene in the

genome assembly of *C. sativa* (Prentout *et al.*, 2020). Adal *et al.*(2021), however, proposes a

different hypothesis suggesting that sex determination may be the result of an X:autosome dosage

mechanism, where male flowers may also have a small number of unique Y-chromosome genes

affecting their sex. We propose that cannabis sexual plasticity is brought about by ethylene's

regulation through differential expression of CsERGs displaying FOC expression and uERG

present in autosomal and sex chromosomes of cannabis.

## Limitations and future directions

The present study represents an important first step in to understanding the ethylene-mediated

control of sex expression in cannabis. However, in the case of this exploratory study, there are

some limitations that we would like to acknowledge. Only two transcriptomic datasets from floral

tissues were available which could be used begin identifying changes in gene expression based on

floral organ type and plant sex that may be linked to sexual plasticity(Prentout *et al.*, 2020; Adal

*et al.*, 2021). In an effort to begin probing the effect of CsERGs during chemically induced sex

plasticity, it was important that we also include transcriptomes from IMFs, which were only

available in one of the two datasets used (Adal *et al.*, 2021). Additionally, no existing

transcriptomic datasets were available for IFFs, which limited our ability to test whether the

mechanisms for sex plasticity in genetically male plants are the inverse of those genetically female

plants. Future studies on sexual plasticity in cannabis should include IFFs in order to provide a

more complete picture of the mechanisms underlying sexual plasticity in the species. Despite these

limitations, we were able to use the differential expression findings between MF and FFs in both

studies to find a set of consensuses differentially expressed CsERGs (**Error! Reference source**

**not found.**), which we believe are strong candidate CsERGs with a role in sexual plasticity.

As noted, there were a number of conditions which differed between the datasets employed in this

study, which may have caused some differentially expressed CsERGs important to sexual

plasticity to not be detected. Transcriptomes from both datasets were generated under different

experimental conditions: plants were grown under different conditions, flowers sampled at

different stages of growth, and the sampling location on the plants were different. Importantly,

plants used to generate the two datasets were not clonal, and each dataset was obtained from a

different genotype: one drug-type (cv. MS-17-338; Adal *et al.*, 2021) and one hemp-type cannabis

(cv. Zenitsa; Prentout *et al.*, 2020). The limitations of single genotype studies in a highly variable

species such as *C. sativa* has been well documented in Monthony et al. (2021b), and thus we

propose that future studies that further probe the role of ethylene in sexual plasticity should include

a greater number of cannabis genotypes. However, it is important to note that most of the CsERGs

identified in the ortholog analysis were present in both datasets, suggesting that the models for

ethylene biosynthesis and signalling in cannabis which we have proposed in this study are robust

and similar to the canonical pathway in *Arabidopsis thaliana*. The fact that these findings are the

same across two datasets with very different experimental conditions (outlined above) can allow

us to confidently say that CsERG expression is affected by a plant's karyotype and that the

canonical ethylene biosynthesis and signalling pathways is conserved in this species.

There are currently a limited number of biotechnological tools (e.g., *in vitro* culture, CRISPR,

VIGS, agrobacterium, etc.) which are optimized in order to functionally validate CsERGs in

cannabis (Hesami *et al.*, 2021) and many of these tools have been hampered by the lack of a reliable

*in vitro* regeneration protocol for the species (Monthony *et al.*, 2021b). Although such functional

validation is beyond the scope of the current study, it represents an important next step in probing

the role of ethylene in cannabis sexual plasticity. Recently progress has been made in establishing

methods for gene silencing in cannabis, through the employment of the *Cotton leaf crumple virus*

(*CLCrV*) in a virus-induced gene silencing protocol reported by Schachtsiek *et al.* (2019) and

through new methods using agrobacterium (Sorokin *et al.*, 2020; Galán-Ávila *et al.*, 2021).

However due to their recent nature, these methods have yet to be replicated independently. Given

the genotypic variability of cannabis and the role it plays in the replicability of *in vitro* methods

(Monthony *et al.*, 2021b), it remains to be seem how effectively these methods are adapted to

multiple genotypes across different research settings.

# Conclusion

This study represents the first step in exploring sexual plasticity in the dioecious plant

*C. sativa* and the important role that Yang cycle, ethylene biosynthesis and signaling pathways

play in sex determination. Here, we present a framework consisting of three distinct gene

expression patterns (KC, FOC and uERG) which can be used to categorize the role of CsERGs in

sexual plasticity in dioecious plants. These findings are based on our analysis of transcriptomes of

male, female and induced male flowers obtained from two different *Cannabis sativa* genotypes

produced under two different experimental conditions. From the ortholog analysis undertaken in

this study and with support from transcriptomic data, we propose that the canonical Yang Cycle

and ethylene biosynthesis and signalling pathways are conserved in *C. sativa* and we identified

seven genes belonging to two the FOC and uERG classes which we hypothesize play a role in

ethylene-induced sexual plasticity in cannabis. This work lays the groundwork for more detailed

study of the role of CsERGs in sexual plasticity in multiple cultivars of cannabis. Given the

commercial importance of cannabis, as well as a number of other dioecious plants such as kiwi,

papaya and persimmon, understanding the underlying factors of sexual plasticity and its link with

ethylene are important to future efforts in breeding and crop improvement.

# Acknowledgments

547 The authors gratefully acknowledge the support of the Natural Sciences and Engineering

548 Research Council (NSERC) of Canada. ASM has also been supported by a NSERC Canada

549 Vanier Graduate Scholarship. The authors would also like to thank Eric Normandeau for his help

550 with script writing.

# Bibliography

**Adal AM, Doshi K, Holbrook L, Mahmoud SS**. **2021**. Comparative RNA-Seq analysis reveals
genes associated with masculinization in female Cannabis sativa. *Planta* **253**: 1–17.

**Altman SA, Solomos T**. **1995**. Differential respiratory and morphological responses of
carnations pulsed or continuously treated with silver thiosulfate. *Postharvest Biology and*
*Technology* **5**: 331–343.

**Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ**. **1990**. Basic local alignment search
tool. *Journal of Molecular Biology* **215**: 403–410.

**Anand L, Rodriguez Lopez CM**. **2022**. ChromoMap: an R package for interactive visualization
of multi-omics data and annotation of chromosomes. *BMC Bioinformatics* **23**: 1–9.

**Andrews S**. **2010**. FastQC: A Quality Control Tool for High Throughput Sequence Data.

**Argueso CT, Hansen M, Kieber JJ**. **2007**. Regulation of ethylene biosynthesis. *Journal of*
*Plant Growth Regulation* **26**: 92–105.

**Ashby R, Forêt S, Searle I, Maleszka R**. **2016**. MicroRNAs in Honey Bee Caste
Determination. *Scientific Reports* **6**: 1–15.

**Barcaccia G, Palumbo F, Scariolo F, Vannozzi A, Borin M, Bona S**. **2020**. Potentials and
challenges of genomics for breeding *Cannabis* cultivars. *Frontiers in Plant Science* **11**: 1–19.

**Beyer EM**. **1976**. A Potent Inhibitor of Ethylene Action in Plants. *Plant Physiology* **58**: 268–
271.

**Binder BM**. **2020**. Ethylene signaling in plants. *Journal of Biological Chemistry* **295**: 7710–
7725.

**Binder BM, Rodríguez FI, Bleecker AB**. **2010**. The copper transporter RAN1 is essential for
biogenesis of ethylene receptors in Arabidopsis. *Journal of Biological Chemistry* **285**: 37263–
37270.

**Blat Y**. **2010**. Non-competitive inhibition by active site binders. *Chemical Biology and Drug*
*Design* **75**: 535–540.

**Bleecker AB, Estelle MA, Somerville C, Kende H**. **1988**. Insensitivity to Ethylene Conferred
by a Dominant Mutation in Arabidopsis thaliana Published by : American Association for the
Advancement of Science Stable URL : https://www.jstor.org/stable/1702414 REFERENCES
Linked references are available on JSTOR for. *Science* **241**: 1086–1089.

**Bleecker AB, Kende H**. **2000**. Ethylene: A gaseous signal molecule in plant. *Annual Review of*
*Cell and Developmental Biology* **16**: 1–18.

**Boualem A, Troadec C, Camps C, Lemhemdi A, Morin H, Sari MA, Fraenkel-Zagouri R,**
**Kovalski I, Dogimont C, Perl-Treves R,** *et al.* **2015**. A cucurbit androecy gene reveals how
unisexual flowers develop and dioecy emerges. *Science* **350**: 688–691.

**Brenner WG, Romanov GA, Köllmer I, Bürkle L, Schmülling T**. **2005**. Immediate-early and
delayed cytokinin response genes of Arabidopsis thaliana identified by genome-wide expression
profiling reveal novel cytokinin-sensitive processes and suggest cytokinin action through
transcriptional cascades. *Plant Journal* **44**: 314–333.

**Brewer C, Harrower M, Sheesley B, Woodruff A, Heyman D**. **2002**. ColorBrewer 2.0.

**Bürstenbinder K, Rzewuski G, Wirtz M, Hell R, Sauter M**. **2007**. The role of methionine
recycling for ethylene synthesis in Arabidopsis. *Plant Journal* **49**: 238–249.

**Cai Z, Cao C, Ji L, Ye R, Wang D, Xia C, Wang S, Du Z, Hu N, Yu X,** *et al.* **2020**. RIC-seq
for global in situ profiling of RNA–RNA spatial interactions. *Nature* **582**: 432–437.

**Cardoso-Júnior CAM, Fujimura PT, Santos-Júnior CD, Borges NA, Ueira-Vieira C,**
**Hartfelder K, Goulart LR, Bonetti AM**. **2017**. Epigenetic modifications and their relation to
caste and sex determination and adult division of labor in the stingless bee Melipona scutellaris.
*Genetics and Molecular Biology* **40**: 61–68.

**Carey SB, Jenkins J, Lovell JT, Maumus F, Sreedasyam A, Payton AC, Shu S, Tiley GP,**
**Fernandez-Pozo N, Healey A,** *et al.* **2021**. Gene-rich UV sex chromosomes harbor conserved
regulators of sexual development. *Science Advances* **7**.

**Cebrián G, Iglesias-Moya J, Romero J, Martínez C, Garrido D, Jamilena M**. **2022**. The
Ethylene Biosynthesis Gene CpACO1A: A New Player in the Regulation of Sex Determination
and Female Flower Development in Cucurbita pepo. *Frontiers in Plant Science* **12**: 1–16.

**Chang C**. **2016**. Q&A: How do plants respond to ethylene and what is its importance? *BMC*

*Biology* **14**: 7.

**Chao Q, Rothenberg M, Solano R, Roman G, Terzaghi W, Ecker JR**. **1997**. Activation of the
ethylene gas response pathway in arabidopsis by the nuclear protein ETHYLENE-
INSENSITIVE3 and related proteins. *Cell* **89**: 1133–1144.

**Charlesworth D, Mank JE**. **2010**. The birds and the bees and the flowers and the trees: Lessons
from genetic mapping of sex determination in plants and animals. *Genetics* **186**: 9–31.

**Chen MK, Hsu WH, Lee PF, Thiruvengadam M, Chen HI, Yang CH**. **2011**. The MADS box
gene, FOREVER YOUNG FLOWER, acts as a repressor controlling floral organ senescence and
abscission in Arabidopsis. *Plant Journal* **68**: 168–185.

**Chen WH, Jiang ZY, Hsu HF, Yang CH**. **2021**. Silencing of FOREVER YOUNG FLOWER-
Like Genes from Phalaenopsis Orchids Promotes Flower Senescence and Abscission. *Plant and
Cell Physiology* **62**: 111–124.

**Chen WH, Li PF, Chen MK, Lee YI, Yang CH**. **2015**. FOREVER YOUNG FLOWER
negatively regulates ethylene response DNA-binding factors by activating an ethylene-
responsive factor to control arabidopsis floral organ senescence and abscission. *Plant Physiology*
**168**: 1666–1683.

**Chen WH, Lin PT, Hsu WH, Hsu HF, Li YC, Tsao CW, Hsu MC, Mao WT, Yang CH**.
**2022**. Regulatory network for FOREVER YOUNG FLOWER-like genes in regulating
Arabidopsis flower senescence and abscission. *Communications Biology* **5**.

**Chen H, Sun J, Li S, Cui Q, Zhang H, Xin F, Wang H, Lin T, Gao D, Wang S, *et al.* 2016a**.
An ACC Oxidase Gene Essential for Cucumber Carpel Development. *Molecular Plant* **9**: 1315–
1327.

**Chen Y, Zou T, McCormick S**. **2016b**. S-adenosylmethionine synthetase 3 is important for
pollen tube growth. *Plant Physiology* **172**: 244–253.

**Cheng WH, Chiang MH, Hwang SG, Lin PC**. **2009**. Antagonism between abscisic acid and
ethylene in Arabidopsis acts in parallel with the reciprocal regulation of their metabolism and
signaling pathways. *Plant Molecular Biology* **71**: 61–80.

**Coulter JM, Barnes CR (Eds.)**. **1901**. *Botanical Gazette*. Chicago: University of Chicago Press.

**Dhakal S, Karki M, Subedi P, GC A**. **2019**. Effect Of Ethephon Doses On Vegetative
Characters, Sex Expression and Yield Of Cucumber (Cucumis sativus cv. Bhaktapur Local) In
Resunga Municipality, Gulmi, Nepal. *International Journal of Applied Sciences and
Biotechnology* **7**: 370–377.

**Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M,
Gingeras TR**. **2013**. STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* **29**: 15–21.

**Dubois M, Van den Broeck L, Inzé D**. **2018**. The Pivotal Role of Ethylene in Plant Growth.
*Trends in Plant Science* **23**: 311–323.

**Emms DM, Kelly S**. **2019**. OrthoFinder: Phylogenetic orthology inference for comparative
genomics. *Genome Biology* **20**: 1–14.

**Ferfuia C, Zuliani F, Danuso F, Piani B, Cattivello C, Dorigo G, Baldini M**. **2021**. Performance and stability of different monoecious hemp cultivars in a multi-environments trial in north-eastern Italy. *Agronomy* **11**: 1–15.

**Flajšman M, Slapnik M, Murovec J**. **2021**. Production of Feminized Seeds of High CBD Cannabis sativa L. by Manipulation of Sex Expression and Its Application to Breeding. *Frontiers in Plant Science* **12**: 1–12.

**Galán-Ávila A, Gramazio P, Ron M, Prohens J, Herraiz FJ**. **2021**. A novel and rapid method for Agrobacterium-mediated production of stably transformed Cannabis sativa L. plants. *Industrial Crops and Products* **170**: 113691.

**Gane R**. **1934**. Production of Ethylene by Some Ripening Fruits. *Nature* **134**: 1008–1008.

**Grassa CJ, Weiblen GD, Wenger JP, Dabney C, Poplawski SG, Timothy Motley S, Michael TP, Schwartz CJ**. **2021**. A new Cannabis genome assembly associates elevated cannabidiol (CBD) with hemp introgressed into marijuana. *New Phytologist* **230**: 1665–1679.

**Green G**. **2005**. *The Cannabis Breeder's Bible- The definitive guide to marijuana genetics, cannabis botany and creating strains for the seed market*. San Francisco: Green Candy Press.

**Guo H, Ecker JR**. **2003**. Plant responses to ethylene gas are mediated by SCFEBF1/EBF2-dependent proteolysis of EIN3 transcription factor. *Cell* **115**: 667–677.

**Guzman P, Ecker JR**. **1990**. Exploiting the Triple Response of Arabidopsis to Identify Ethylene-Related Mutants. *The Plant Cell* **2**: 513.

**Hayes TB, Khoury V, Narayan A, Nazir M, Parka A, Brown T, Adame L, Chan E, Buchholz D, Stueve T, *et al.* 2010**. Atrazine induces complete feminization and chemical castration in male African clawed frogs (Xenopus laevis). *Proceedings of the National Academy of Sciences of the United States of America* **107**: 4612–4617.

**Hesami M, Baiton A, Alizadeh M, Pepe M, Torkamaneh D, Jones AMP**. **2021**. Advances and Perspectives in Tissue Culture and Genetic Engineering of Cannabis. *International Journal of Molecular Sciences* **22**: 5671.

**Hirayama T, Kieber JJ, Hirayama N, Kogan M, Guzman P, Nourizadeh S, Alonso JM, Dailey WP, Dancis A, Ecker JR**. **1999**. RESPONSIVE-TO-ANTAGONIST1, a Menkes/Wilson disease-related copper transporter, is required for ethylene signaling in Arabidopsis. *Cell* **97**: 383–393.

**Houben M, Van de Poel B**. **2019**. 1-aminocyclopropane-1-carboxylic acid oxidase (ACO): The enzyme that makes the plant hormone ethylene. *Frontiers in Plant Science* **10**: 1–15.

**Hume B, Lovell P**. **1983a**. Role of aminocyclopropane-l-carboxylic acid in ethylene release by distal tissues following localized application of ethephon in Cucurbita pepo. *Physiologia Plantarum* **58**: 101–106.

**Hume RJ, Lovell PH**. **1983b**. The control of sex expression in cucurbits by ethephon. *Annals of Botany* **52**: 689–695.

**Hyde PT, Guan · Xian, Abreu V, Setter TL**. **2020**. The anti-ethylene growth regulator silver

thiosulfate (STS) increases flower production and longevity in cassava (Manihot esculenta
Crantz). *Plant Growth Regulation* **90**: 441–453.

**Kende H**. **1998**. Plant Biology and the Nobel Prize. *Science* **282**: 627–627.

**Kieber JJ, Rothenberg M, Roman G, Feldmann KA, Ecker JR**. **1993**. CTR1, a negative
regulator of the ethylene response pathway in arabidopsis, encodes a member of the Raf family
of protein kinases. *Cell* **72**: 427–441.

**Kucukural A, Yukselen O, Ozata DM, Moore MJ, Garber M**. **2019**. DEBrowser: interactive
differential expression analysis and visualization tool for count data. *BMC Genomics* **20**.

**Kurtz LE, Brand MH, Lubell-Brand JD**. **2020**. Production of tetraploid and triploid hemp.
*HortScience* **55**: 1703–1707.

**Lambert MR, Giller GSJ, Barber LB, Fitzgerald KC, Skelly DK**. **2015**. Suburbanization,
estrogen contamination, and sex ratio in wild amphibian populations. *Proceedings of the*
*National Academy of Sciences of the United States of America* **112**: 11881–11886.

**Lambert MR, Tran T, Kilian A, Ezaz T, Skelly DK**. **2019**. Molecular evidence for sex
reversal in wild populations of green frogs (Rana clamitans). *PeerJ* **2019**: 1–21.

**Li Q, Guo W, Chen B, Pan F, Yang H, Zhou J, Wang G, Li X**. **2021**. Transcriptional and
Hormonal Responses in Ethephon-Induced Promotion of Femaleness in Pumpkin. *Frontiers in*
*Plant Science* **12**.

**Li Q, Zhang L, Pan F, Guo W, Chen B, Yang H, Wang G, Li X**. **2020**. Transcriptomic
analysis reveals ethylene signal transduction genes involved in pistil development of pumpkin.
*PeerJ*: 1–17.

**Liu H, Todd E V., Lokman PM, Lamm MS, Godwin JR, Gemmell NJ**. **2017**. Sexual
plasticity: A fishy tale. *Molecular Reproduction and Development* **84**: 171–194.

**Love MI, Huber W, Anders S**. **2014**. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biology* **15**: 1–21.

**Lubell JD, Brand MH**. **2018**. Foliar sprays of silver thiosulfate produce male flowers on female
hemp plants. *HortTechnology* **28**: 743–747.

**Manzano S, Martínez C, García JM, Megías Z, Jamilena M**. **2014**. Involvement of ethylene
in sex expression and female flower development in watermelon (Citrullus lanatus). *Plant*
*Physiology and Biochemistry* **85**: 96–104.

**Mayerhofer H, Panneerselvam S, Mueller-Dieckmann J**. **2012**. Protein kinase domain of
CTR1 from arabidopsis thaliana promotes ethylene receptor cross talk. *Journal of Molecular*
*Biology* **415**: 768–779.

**McDaniel BK, Binder BM**. **2012**. Ethylene receptor 1 (ETR1) is sufficient and has the
predominant role in mediating inhibition of ethylene responses by silver in Arabidopsis thaliana.
*Journal of Biological Chemistry* **287**: 26094–26103.

**Meng J, Wang L, Wang J, Zhao X, Cheng J, Yu W, Jin D, Li Q, Gong Z**. **2018**.
METHIONINE ADENOSYLTRANSFERASE4 mediates DNA and histone methylation. *Plant*

*Physiology* **177**: 652–670.

**Mohan Ram HY, Sett R**. **1982**. Induction of Fertile Male Flowers in Genetically Female Cannabis sativa Plants by Silver Nitrate and Silver Thiosulphate Anionic Complex. *Theoretical and Applied Genetics* **62**: 369–275.

**Monthony AS, Bagheri S, Zheng Y, Jones AMP**. **2021a**. Flower power: floral reversion as a viable alternative to nodal micropropagation in Cannabis sativa. *In Vitro Cellular & Developmental Biology - Plant* **21**: 1–13.

**Monthony AS, Kyne ST, Grainger CM, Jones AMP**. **2021b**. Recalcitrance of Cannabis sativa to de novo regeneration; a multi-genotype replication study (KR Davis, Ed.). *PLOS ONE* **16**: e0235525.

**Monthony AS, Page SR, Hesami M, Jones AMP**. **2021c**. The Past, Present and Future of *Cannabis sativa* Tissue Culture. *Plants* **10**: 185.

**Moon Y-H, Lee YJ, Koo SC, Hur M, Huh YC, Chang J-K, Park WT**. **2020a**. Effect of Timing of Ethephon Treatment on the Formation of Female Flowers and Seeds from Male Plant of Hemp (Cannabis sativa L.). *The Plant Resources Society of Korea* **33(6)**: 682–688.

**Moon J, Park YJ, Son SH, Roh J, Youn JH, Kim SY, Kim SK**. **2020b**. Brassinosteroids signaling via BZR1 down-regulates expression of ACC oxidase 4 to control growth of Arabidopsis thaliana seedlings. *Plant Signaling and Behavior* **15**.

**Morgan TH**. **1909**. Sex Determination and Parthenogenesis in Phylloxerans and Aphids. *Science* **29**: 234–237.

**Oda A, Nomura-Ando K, Ahn D-H, Higashide T**. **2022**. Temporary Reduction and Control of Female Flower Expression in Cucumber (*Cucumis sativus* L.) by Application of 1-Methylcyclopropene. *The Horticulture Journal* **91**: 42–48.

**Okonechnikov K, Conesa A, García-Alcalde F**. **2015**. Qualimap 2: advanced multi-sample quality control for high-throughput sequencing data.

**Olsen A, Lütken H, Hegelund JN, Müller R**. **2015**. Ethylene resistance in flowering ornamental plants – improvements and future perspectives. *Horticulture Research* **2**: 15038.

**Pattyn J, Vaughan-Hirsch J, Van de Poel B**. **2021**. The regulation of ethylene biosynthesis: a complex multilevel control circuitry. *New Phytologist* **229**: 770–782.

**Peleman J, Boerjan W, Engler G, Seurinck J, Botterman J, Alliotte T, Van Montagu M, Inzé D**. **1989a**. Strong cellular preference in the expression of a housekeeping gene of Arabidopsis thaliana encoding S-adenosylmethionine synthetase. *The Plant cell* **1**: 81–93.

**Peleman J, Saito K, Cottyn B, Engler G, Seurinck J, Van Montagu M, Inzé D**. **1989b**. Structure and expression analyses of the S-adenosylmethionine synthetase gene family in Arabidopsis thaliana. *Gene* **84**: 359–369.

**Perry AN, Grober MS**. **2003**. A model for social control of sex change: Interactions of behavior, neuropeptides, glucocorticoids, and sex steroids. *Hormones and Behavior* **43**: 31–38.

**van de Poel B, Bulens I, Markoula A, Hertog MLATM, Dreesen R, Wirtz M, Vandoninck**

S, Oppermann Y, Keulemans J, Hell R, *et al.* **2012**. Targeted systems biology profiling of tomato fruit reveals coordination of the Yang cycle and a distinct regulation of ethylene biosynthesis during postclimacteric ripening. *Plant Physiology* **160**: 1498–1514.

**Pommerrenig B, Feussner K, Zierer W, Rabinovych V, Klebl F, Feussner I, Sauera N**. **2011**. Phloem-Specific expression of Yang cycle genes and identification of novel Yang cycle enzymes in Plantago and Arabidopsis. *Plant Cell* **23**: 1904–1919.

**Prentout D, Razumova O, Rhoné B, Badouin H, Henri H, Feng C, Käfer J, Karlov G, Marais GAB**. **2020**. An efficient RNA-seq-based segregation analysis identifies the sex chromosomes of Cannabis sativa. *Genome Research* **30**: 164–172.

**Putri GH, Anders S, Pyl PT, Pimanda JE, Zanini F**. **2022**. Analysing high-throughput sequencing data in Python with HTSeq 2.0. *Bioinformatics* **38**: 2943–2945.

**Qiao H, Chang KN, Yazaki J, Ecker JR**. **2009**. Interplay between ethylene, ETP1/ETP2 F-box proteins, and degradation of EIN2 triggers ethylene responses in Arabidopsis. *Genes and Development* **23**: 512–521.

**Qiu K, Li Z, Yang Z, Chen J, Wu S, Zhu X, Gao S, Gao J, Ren G, Kuai B, *et al.* 2015**. EIN3 and ORE1 Accelerate Degreening during Ethylene-Mediated Leaf Senescence by Directly Activating Chlorophyll Catabolic Genes in Arabidopsis. *PLoS Genetics* **11**: 1–20.

**Rai MI, Wang X, Thibault DM, Kim HJ, Bombyk MM, Binder BM, Shakeel SN, Schaller GE**. **2015**. The ARGOS gene family functions in a negative feedback loop to desensitize plants to ethylene. *BMC Plant Biology* **15**: 1–14.

**Ram HYM, Jaiswal VS**. **1972**. Induction of male flowers on female plants of Cannabis sativa by gibberellins and its inhibition by abscisic acid. *Planta* **105**: 263–266.

**Raz V, Ecker JR**. **1999**. Regulation of differential growth in the apical hook of Arabidopsis. *Development* **126**: 3661–3668.

**Renner SS**. **2014**. The relative and absolute frequencies of angiosperm sexual systems: Dioecy, monoecy, gynodioecy, and an updated online database. *American Journal of Botany* **101**: 1588–1596.

**Resnick JS, Wen C-K, Shockey JA, Chang C**. **2006**. REVERSION-TO-ETHYLENE SENSITIVITY1, a conserved gene that regulates ethylene receptor function in Arabidopsis. *Proceedings of the National Academy of Sciences* **103**: 7917–7922.

**Rodríguez FI, Esch JJ, Hall AE, Binder BM, Schaller GE, Bleecker AB**. **1999**. A Copper Cofactor for the Ethylene Receptor ETR1 from Arabidopsis. *Science* **283**: 996–998.

**Roman G, Lubarsky B, Kieber JJ, Rothenberg M, Ecker JR**. **1995**. Genetic analysis of ethylene signal transduction in Arabidopsis thaliana: Five novel mutant loci integrated into a stress response pathway. *Genetics* **139**: 1393–1409.

**Sakai H, Hua J, Chen QG, Chang C, Medrano LJ, Bleecker AB, Meyerowitz EM**. **1998**. ETR2 is an ETR1-like gene involved in ethylene signaling in Arabidopsis. *Proceedings of the National Academy of Sciences of the United States of America* **95**: 5812–5817.

**Schachtsiek J, Hussain T, Azzouhri K, Kayser O, Stehle F**. **2019**. Virus-induced gene
silencing (VIGS) in Cannabis sativa L. *Plant Methods* **15**: 1–9.

**Schlesinger A, Kramarsky-Winter E, Rosenfeld H, Armoza-Zvoloni R, Loya Y**. **2010**.
Sexual plasticity and self-fertilization in the sea anemone Aiptasia diaphana. *PLoS ONE* **5**: 1–7.

**Senthilkumaran B**. **2015**. Pesticide- and sex steroid analogue-induced endocrine disruption
differentially targets hypothalamo–hypophyseal–gonadal system during gametogenesis in
teleosts – A review. *General and Comparative Endocrinology* **219**: 136–142.

**Shakeel SN, Wang X, Binder BM, Schaller GE**. **2013**. Mechanisms of signal transduction by
ethylene: Overlapping and non-overlapping signalling roles in a receptor family. *AoB PLANTS* **5**:
1–16.

**Shi J, Drummond BJ, Wang H, Archibald RL, Habben JE**. **2016**. Maize and Arabidopsis
ARGOS proteins interact with ethylene receptor signaling complex, supporting a regulatory role
for ARGOS in ethylene signal transduction. *Plant Physiology* **171**: 2783–2797.

**Shi J, Habben JE, Archibald RL, Drummond BJ, Chamberlin MA, Williams RW, Renee
Lafitte H, Weers BP**. **2015**. Overexpression of ARGOS genes modifies plant sensitivity to
ethylene, leading to improved drought tolerance in both arabidopsis and maize. *Plant Physiology*
**169**: 266–282.

**Solano R, Stepanova A, Chao Q, Ecker JR**. **1998**. Nuclear events in ethylene signaling: A
transcriptional cascade mediated by ETHYLENE-INSENSITIVE3 and ETHYLENE-
RESPONSE-FACTOR1. *Genes and Development* **12**: 3703–3714.

**Sorokin A, Yadav NS, Gaudet D, Kovalchuk I**. **2020**. Transient expression of the β-
glucuronidase gene in Cannabis sativa varieties. *Plant Signaling and Behavior* **15**: 1–8.

**Thomas M**. **2012**. *Cannabis cultivation: A Complete Grower's Guide*. San Francisco: Green
Candy Press.

**Tieman DM, Taylor MG, Ciardi JA, Klee HJ**. **2000**. The tomato ethylene receptors NR and
LeETR4 are negative regulators of ethylene response and exhibit functional compensation within
a multigene family. *Proceedings of the National Academy of Sciences of the United States of
America* **97**: 5663.

**Trivellini A, Cocetta G, Hunter DA, Vernieri P, Ferrante A**. **2016**. Spatial and temporal
transcriptome changes occurring during flower opening and senescence of the ephemeral
hibiscus flower, Hibiscus rosa-sinensis. *Journal of Experimental Botany* **67**: 5919–5931.

**Warner RR, Swearer SE**. **1991**. Social control of sex change in the bluehead wrasse,
Thalassoma bifasciatum (Pisces: Labridae). *Biological Bulletin* **181**: 199–204.

**Wickham H**. **2016**. *ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York*.

**Wu Z, Zhang X, Zhang N, Gao X, Feng X, Zeng Q, Chen X, Wu J, Qi Y**. **2022**. Identification
of ACC Synthetase Genes in Saccharum and their Expression Profiles During Plant Growth and
in Response to Low-nitrogen Stress. *Tropical Plant Biology* **15**: 197–210.

**Yamagami T, Tsuchisaka A, Yamada K, Haddon WF, Harden LA, Theologis A**. **2003**.

Biochemical diversity among the 1-amino-cyclopropane-1-carboxylate synthase isozymes
encoded by the Arabidopsis gene family. *The Journal of biological chemistry* **278**: 49102–49112.

**Yamasaki S, Fujii N, Matsuura S, Mizusawa H, Takahashi H**. **2001**. The M Locus and
Ethylene-Controlled Sex Determination in Andromonoecious Cucumber Plants. *Plant and Cell*
*Physiology* **42**: 608–619.

**Yang C, Lu X, Ma B, Chen SY, Zhang JS**. **2015**. Ethylene signaling in rice and arabidopsis:
Conserved and diverged aspects. *Molecular Plant* **8**: 495–505.

**Zhang J, Shi J, Ji G, Zhang H, Gong G, Guo S, Ren Y, Fan J, Tian S, Xu Y**. **2017**.
Modulation of Sex Expression in Four Forms of Watermelon by Gibberellin, Ethephone and
Silver Nitrate. *Horticultural Plant Journal* **3**: 91–100.

**Zhang W, Wen CK**. **2010**. Preparation of ethylene gas and comparison of ethylene responses
induced by ethylene, ACC, and ethephon. *Plant Physiology and Biochemistry* **48**: 45–53.

**Zheng Y**. **2022**. *Handbook of Cannabis Production in Controlled Environments* (Y Zheng, Ed.).
Boca Raton: CRC Press.

**Zhou X, Liu Q, Xie F, Wen CK**. **2007**. RTE1 is a Golgi-associated and ETR1-dependent
negative regulator of ethylene responses. *Plant Physiology* **145**: 75–86.