Full title:

**Exploring Ethylene-Related Genes in *Cannabis sativa*: Implications for Sexual Plasticity**

Short title:

**Ethylene and *Cannabis* sexual plasticity**

**Adrian S. Monthony, Maxime de Ronne & Davoud Torkamaneh**[*]

*Département de phytologie, Université Laval, Québec City, Québec, Canada*

*Institut de Biologie Intégrative et des Systèmes (IBIS), Université Laval, Québec, Canada*

*Centre de recherche et d'innovation sur les végétaux (CRIV), Université Laval, Québec, Canada*

*Institut intelligence et données (IID), Université Laval, Québec, Canada*

Monthony ORCID:  0000-0003-0536-691X

de Ronne ORCID: 0000-0003-4062-3512

*Torkamaneh ORCID:  0000-0002-9782-5695

*Corresponding Author

Email:  davoud.torkamaneh.1@ulaval.ca

# Summary

Sexual plasticity describes an organism's ability to change its phenotypic sex in response to environmental, chemical or physiological cues without a change in sex chromosome karyotype. *Cannabis sativa* L. (cannabis) is a medically important dioecious crop, with a pair of heteromorphic XY sex chromosomes. Cannabis has demonstrated sexual plasticity in response to chemical treatments which target the biosynthesis and signalling of ethylene. To date, ethylene biosynthesis and signalling has not been specifically studied in *C. sativa,* and the role of ethylene-related genes (ERGs) in sexual plasticity in the species has not been investigated. In the present study, we performed gene orthology analysis based on the model species *Arabidopsis thaliana* L. to reconstruct the *C. sativa* Yang Cycle, ethylene biosynthesis and ethylene signalling pathways, whose canonical composition appear conserved in cannabis. Additionally, we employ two transcriptomic datasets containing male, female and chemically induced male flowers to investigate patterns of expression for ethylene-related genes which may be involved in sexual plasticity. Comparison of these transcriptomes highlighted CsERGs whose differential expression was associated with sexual determination and those which we propose are associated with the sexual plasticity. These sexual plasticity ERGs could be classified into two patterns of expression: floral organ concordant (FOC) and unique (uERG). We propose a third pattern, involved in sex determination, which we have named karyotype concordant (KC) expression. We show that CsERGs involved in sexual plasticity are distributed throughout the genome and are not exclusive to the sex chromosomes, suggesting a broad control of sexual plasticity in *C. sativa*.

**Key words:** ethylene, sexual plasticity, cannabis, STS, biosynthesis, sex expression, sex determination, transcriptomics

# Introduction

Cannabis sativa L. (cannabis) belongs to the estimated 5-6% of dioecious angiosperm species (Renner, 2014). Despite their rarity, dioecious plants provide an excellent opportunity to study the mechanisms involved in sex expression and the evolution of sex-determining regions and sex chromosomes. Cannabis (2n=20) is predominantly dioecious with a XY sex-determination system; however, monoecious genotypes of *C. sativa* have been bred, especially for the production of seeds and oil used in industrial hemp (Ferfuia *et al.*, 2021). Additionally, dioecious cannabis genotypes have been known to display hermaphroditism (Monthony *et al.*, 2021c), reversion from a sexual to vegetative state (Monthony *et al.*, 2021a), and sometimes complete changes in sex expression in response to certain environmental factors and chemical reagents (Ram & Jaiswal, 1972; Mohan Ram & Sett, 1982; Lubell & Brand, 2018; Moon *et al.*, 2020a). Taken together, these observations suggest that cannabis sex is plastic and not strictly controlled by its XX or XY chromosomal complement.

Sexual plasticity is the ability of an sex chromosome containing organism to change its phenotypic sex in response to environmental or physiological cues without altering its sex chromosomes (Liu *et al.*, 2017). Vertebrates exhibit sexual plasticity on a continuum from genotypic sex determination (GSD) to environmental sex determination (ESD), with some organisms influenced by both (GSD + ESD; Lambert *et al.*, 2019). Non-mammalian vertebrates, such as fish (Senthilkumaran, 2015; Liu *et al.*, 2017) and frogs (Lambert *et al.*, 2019), as well as insects like aphids and bees (Morgan, 1909; Charlesworth & Mank, 2010), exhibit sexual plasticity triggered by diverse factors. The mechanisms responsible for these changes vary, but can involve gene expression changes related to hormone synthesis and signalling, germ cell development and maintenance as well as microRNAs (Perry & Grober, 2003; Hayes *et al.*, 2010; Ashby *et al.*, 2016; Cardoso-Júnior *et al.*, 2017). For instance, social factors such as population density, mating competition, and hierarchical structures can affect sexual plasticity in fish (Warner & Swearer, 1991; Liu *et al.*, 2017). Amphibians can exhibit sexual plasticity under normal and extreme conditions (Lambert *et al.*, 2015, 2019), while in some insects, the mode of sexual reproduction can affect offspring sex (Morgan, 1909; Charlesworth & Mank, 2010). Although the term sexual plasticity has largely been used to describe fluid sex expression in the animal kingdom (Schlesinger *et al.*, 2010; Senthilkumaran, 2015; Liu *et al.*, 2017), the present study highlights how

sexual plasticity can also be used to describe the interaction between GSD and ESD in dioecious

plants containing sex chromosomes, such as *C. sativa*.

The present study defines sexual plasticity as a deviation from the typical sexual expression

dictated by their sex chromosomes of dioecious plants. This deviation occurs when certain genes,

influenced by external factors like chemicals, stress, or climate changes, induce a switch from the

plant's chromosomal sex to a contrasting phenotypic sex, akin to what has been reported in

vertebrates (Liu *et al.*, 2017; Lambert *et al.*, 2019). These genes involved in sexual plasticity

override the expected floral sex determined by the sex chromosomes. In dioecious plants with sex

chromosomes, sex determination genes ensure the proper development of floral organs in

accordance with the plant's sex chromosome karyotype. However, in monoecious plants that

produce both male and female flowers, only genes involved in sex determination are present, as

their expression directly influences the sex or sex ratio of the flowers produced. This distinction

underscores the role of sexual plasticity in dioecious plants with sex chromosomes, such as

cannabis, and highlights how it complements the traditional understanding of sex determination

mechanisms of monoecious plants.

Informal research conducted by enthusiast cannabis breeders prior to legalization has

demonstrated that chemical exposure can cause sexual plasticity in plants, not unlike in

amphibians. (Green, 2005; Thomas, 2012). This pre-legalization exploratory work identified

silver-containing compounds such as silver thiosulfate (STS), a potent inhibitor of ethylene

signalling, as promoting the production of male flowers on female plants (induced male flowers;

IMF) and aqueous ethephon ((2-Chloroethyl) phosphonic acid), which rapidly decomposes to

produce ethylene gas, as promoting the production of female flowers on male plants (induced

female flowers; IFF) as shown in Figure 1. Following the legalization of cannabis in some

jurisdictions, the inverse effects of STS and ethephon on cannabis sex have been re-demonstrated,

quantified and published by a handful of research groups (Lubell & Brand, 2018; Kurtz *et al.*,

2020; Moon *et al.*, 2020a; Flajšman *et al.*, 2021), pointing to ethylene as being an effector of sexual

plasticity. Although the phenotypic outcomes of these treatments in cannabis have been quantified,

to date there are no studies which have sought to understand the underlying molecular mechanisms

controlling ethylene-related sexual plasticity in cannabis.

**Figure 1** Representative photos of four floral phenotypes in *Cannabis sativa* (cv. La Rosca), grown at Université Laval. a) 4-week-old female flowers on a genetically female plant; b) 4-week-old STS-induced male flowers (IMFs) on a genetically female plant; c) ethephon-induced female flowers (IFFs) on a genetically male plant and d) 3-week-old male flowers on a genetically male plant. Scale bars = 5 mm.

Ethylene was discovered in 1901 and has been extensively studied since then (Coulter &
Barnes, 1901; Gane, 1934; Kende, 1998; Chang, 2016). It plays a crucial role as a plant growth
regulator in numerous plant processes such as seed germination, root and shoot growth, flowering,
fruit ripening, leaf and fruit abscission, as well as responses to environmental stresses and
senescence (Tieman *et al.*, 2000; Resnick *et al.*, 2006; Argueso *et al.*, 2007; Yang *et al.*, 2015;
Dhakal *et al.*, 2019; Binder, 2020). Ethylene is synthesized from S-adenosyl methionine (SAM)
through a two-step pathway. The first step involves the enzyme 1-aminocyclopropane-1-carboxilic
acid (ACC) synthase (ACS), which converts SAM to ACC and 5'-methylthioadenosine (MTA;
Argueso *et al.* 2007; Pattyn *et al.* 2021). *ACS* genes are regulated differently based on
environmental and developmental factors (Argueso et al., 2007; Dubois et al., 2018). Arabidopsis
has eight identified *ACS* genes (Argueso *et al.*, 2007). The final step is catalyzed by ACC oxidase
(ACO), which converts ACC to ethylene, cyanide, and $CO_2$ (Argueso *et al.*, 2007; Houben & Van
de Poel, 2019). Arabidopsis has around five *ACO* genes (Houben & Van de Poel, 2019), while
cucumber has four *ACO* genes involved in sex determination (Chen *et al.*, 2016a).

The genes involved in ethylene signalling are highly conserved among plants. Ethylene
receptors, including *ethylene response sensor 1* (*ERS1*), *ERS2*, *ethylene resistance 1* (*ETR1*),
*ETR2*, and *ethylene insensitive 4* (*EIN4*), found in the endoplasmic reticulum, bind to ethylene to
initiate signalling (Dubois *et al.*, 2018). Ethylene acts as an inverse agonist, promoting its own
production by altering the activity of the biosynthetic pathway and reducing inhibition of
downstream signalling (Mayerhofer *et al.*, 2012; Binder, 2020). Ethylene inhibits *constitutive*
*triple response 1* (*CTR1*) activity, leading to dephosphorylation of EIN2 and cleavage of its c-
terminal domain (C-END). This allows for the repression of *EBF1/2* translation by binding to the
3' untranslated regions of their mRNA (Chang, 2016; Dubois *et al.*, 2018; Binder, 2020). In the
absence of ethylene, *EBF1/2* target the primary ethylene-responsive transcription factors *EIN3* and
*EIN3-LIKE* (*EIL1*) for degradation (Dubois *et al.*, 2018). Repression of *EBF1/2* by ethylene
enables the expression of *EIN3* and *EIN3-like* transcription factors, which induce the expression
of ethylene response factors (ERFs), further promoting ethylene biosynthesis. Although the
canonical signalling pathways are well understood, further investigation is needed regarding ERFs
(Olsen *et al.*, 2015; Dubois *et al.*, 2018).

Compared with dioecious cannabis, the role of ethylene in sex determination in
monoecious members of the Cucurbitaceae family has been much more well-defined. In cucurbits
ethylene biosynthesis and signalling have also been shown to be manipulated by STS or ethephon
applications (Li *et al.*, 2021). Typically, ethylene (usually applied as ethephon) has a feminizing
effect, increasing the ratio of female flowers relative to males, while the use of STS has the inverse
effect, causing a masculinizing effect and results in increased production of male flowers.
Numerous studies of sex determination in cucurbits have pointed to the role of ethylene
biosynthesis genes such as *ACS4* and *ACS11*, as well as ethylene signalling genes such as the
ethylene receptor *ETR1* as contributing to sex determination (Boualem *et al.*, 2015; Zhang *et al.*,
2017; Li *et al.*, 2020). This has led to the labelling of some key loci in Cucurbitaceae as sex-
determining loci, with changes of expression of these particular loci determining the sex of the
plant, in contrast to *C. sativa* where the chromosome karyotype determines plant sex. The
manipulation of cucumber sex has allowed growers to control and optimize the ratio of female to
male flowers to improve yield and to control production so as to meet labour constraints (Oda *et
al.*, 2022). In cannabis, control of plant sex is important as medicinally valuable cannabinoids are
primarily produced in female flowers. However, without a pathway model for ethylene
biosynthesis and signalling in the species, study of cannabis sexual plasticity remains impossible.
The highly conserved nature of ethylene biosynthesis and signalling and well-characterized
pathways in model species *Arabidopsis thaliana* present a valuable starting point for the
reconstruction of the pathway in an under-researched plant such as *C. sativa* where ethylene
biosynthesis and signalling has yet to be explored. In the current study, we use homology analysis
and RNA-seq data to present the first *C. sativa* putative Yang cycle, ethylene biosynthesis and
ethylene signalling pathways, which will serve as an important foundation for expanding our
understanding of the role of ethylene in sexual plasticity of this under-researched dioecious
species.

# Materials & Methods

## *In silico* identification of ethylene-related gene orthologs in *C. sativa*

*Arabidopsis thaliana* was chosen as the model from which a list of genes related to ethylene biosynthesis and signalling, as well as the production of ethylene precursors derived from the Yang cycle were compiled following a review of the literature (Pommerrenig *et al.*, 2011; van de Poel *et al.*, 2012; Olsen *et al.*, 2015; Chang, 2016; Pattyn *et al.*, 2021). The resulting list of *A. thaliana* ethylene-related genes (AtERGs; Supplementary Table 2) were used for the identification of orthologous *C. sativa* ERGs (CsERGs). Nucleotide and protein sequences for AtERGs were obtained from the National Center for Biotechnology Information database (NCBI; Bethesda, MD). Ortholog identification was conducted on the *C. sativa* cs10 reference genome (RefSeq assembly accession: GCF_900626175.2; Grassa *et al.*, 2021). A list of putative CsERG orthologs was compiled using two different methods. The first method employed OrthoFinder (Emms & Kelly, 2019) adapted from Carey *et al.* (2021) and the second method consisted of a BLASTP search of each individual AtERG against the cs10 reference genome (Altschul *et al.*, 1990), followed by a manual selection of putative orthologs from the of BLASTP result adapted from Barcaccia *et al.* (2020). This method is described in further detail in the supporting information. The chromosome map (Figure 4a) was generated with chromoMap (v4.1.1; Anand & Rodriguez Lopez, 2022) and bar plots of gene distribution (Figure 4b) was generated with ggplot2 (v3.4.1; Wickham, 2016) using the colour palette Spectral from RColorBrewer (Brewer *et al.*, 2002)

## RNA-seq data collection and processing

The RNA-seq reads from Adal *et al.* (2021) and Prentout *et al.* (2020), hereafter referred to as dataset 1 and dataset 2, were downloaded from the Sequence Read Archive (SRA) on NCBI under the BioProject accessions PRJNA669389 and PRJNA549804, respectively. RNA was isolated from male flowers (MFs), female flowers (FFs) and STS-induced IMFs for dataset 1, and from MFs and FFs for dataset 2. To perform a relevant comparison between the 2 datasets, only data from similar tissues were considered, i.e., the young flower buds. The SRA sample accessions used in this study are indicated in Supplementary Table 1. FastQC (Andrews, 2010) was used for validating the quality of the data. Then, reads were aligned to the *C. sativa* cs10 reference genome

(version GCF_900626175.2; Grassa *et al.*, 2021) using STAR (v2.7.2b; Dobin *et al.*, 2013) as

reported in Cai *et al.* (2020). The STAR index was built with the –sjdbOverhang=100 and the

genomeSAindexbases=13 arguments. Splice junctions from the *C. sativa* cs10 RefSeq assembly

GTF version GCF_900626175.2 were provided via the --sjdbGTFfile argument. STAR alignments

were generated using the default parameters including the --outSAMtype BAM

SortedByCoordinate arguments. Quality control of the mapped reads was performed using

Qualimap (v2.2.1; Okonechnikov *et al.*, 2015). One IMF sample from dataset 1 was excluded

based on abnormal distribution of GC-content among mapped reads which included large

shoulders at 30% and 60 % GC content, indicating contamination of the sample (see

Supplementary Table 1). Finally, quantification of reads overlapping genetic features was

performed with HTSeq (v2.0.1; Putri *et al.*, 2022) based on the cs10 GFF file obtained from NCBI

([https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/900/626/175/GCF_900626175.2_cs10/GCF_9006](https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/900/626/175/GCF_900626175.2_cs10/GCF_9006)

[26175.2_cs10_genomic.gff.gz](26175.2_cs10_genomic.gff.gz)) in order to generate a gene quantification table required for

subsequent differential analysis with DEBrowser (v1.22.5; Kucukural *et al.*, 2019).

## Differential expression analysis of CsERGs

Differentially expressed CsERGs were identified using DESeq2 (Love *et al.*, 2014) with

an FDR-adjusted *p*-value $< 0.05$ on the DEBrowser  graphical user interface in RStudio (RStudio

Inc., MA, USA) using v4.1.1 of R. The analysis was adapted from DEBrowser's DESeq2 method

outlined on the DEBrowser Github ([https://github.com/UMMS-Biocore/debrowser](https://github.com/UMMS-Biocore/debrowser)). In short, gene

quantification tables generated in HTSeq and a manually generated metadata file for the samples

were uploaded to DEBrowser. Genes with count per million values less than 1 (CPM $< 1$) in 7 out

of 15 replicates in dataset 1 and 6 out of 12 replicates in dataset 2 were removed. Replicates were

then grouped and pairwise comparisons were performed based on the floral type. In DEBrowser

the following DESeq2 parameters were used following the method outline by Kucukural *et al.*

(2019) and the DEBrowser Github: *Fit Type- parametric; betaPrior- FALSE; Test Type- Wald;*

*Shrinkage- None*. Differential expression levels were then normalized using a Median Ratio

Normalization (MRN) and pairwise comparisons between treatments were considered

significantly different in subsequent comparisons if they had an adjusted *p*-value $<0.05$ according

to the Wald test and a $|log2foldchange| > 1$. Heatmaps comparing differential gene expression were

then generated using DEBrowser heatmap function. Tables from pairwise comparisons were

generated from DEBrowser and then used for the generation of gene expression bar plots using
ggplot2 (Wickham, 2016) in RStudio.

# Results

## ERGs are largely conserved in *C. sativa*

Based on 47 AtERGs previously identified (Supplementary Table 2), ortholog analyses
identified 43 putative CsERGs (Table 1). At least one ortholog of each key gene in the canonical
ethylene biosynthesis and signalling pathways, as well as the Yang cycle, was identified *in silico*
except for auxin-regulated genes involved in organ size (*AtARGOS*; Supplementary Table 2). To
the best of our knowledge, this is thethe first time that the genes coding for the various enzymes
involved in the putative Yang cycle (Figure 2a), ethylene biosynthesis (Figure 2b) and signalling
pathway (Figure 3) have been proposed in *C. sativa.*

**Table 1** *Cannabis sativa* ethylene-related genes (CsERGs) identified through ortholog analysis.

| Cs10 ERG Locus[*] | Name | Chromosome | *A. thaliana* orthologs | Pathway | Identification method |
|---|---|---|---|---|---|
| LOC115704991 | *CsACO1* | 1 | *AT2G19590* | Biosynthesis | BLASTP, OrthoFinder |
| LOC115702109 | *CsACO1-LIKE1* | Un[§] | *AT2G19590* | Biosynthesis | BLASTP, OrthoFinder |
| LOC115707397 | *CsACO2*[‡†] | 1 | *AT1G62380, AT1G12010, AT1G05010* | Biosynthesis | BLASTP, OrthoFinder |
| LOC115699412 | *CsACO3*[‡†] | 8 | *AT1G62380, AT1G12010, AT1G05010* | Biosynthesis | BLASTP, OrthoFinder |
| LOC115699408 | *CsACO5*[‡†] | X | *AT1G77330* | Biosynthesis | BLASTP, OrthoFinder |
| LOC115704849 | *CsACS1* | 1 | *AT3G49700, AT4G08040, AT4G11280, AT1G01480* | Biosynthesis | BLASTP |
| LOC115717044 | *CsACS2*[‡†] | 5 | *AT4G08040, AT1G01480, AT4G11280, AT3G49700* | Biosynthesis | BLASTP, OrthoFinder |
| LOC115712396 | *CsACS3* | 4 | *AT5G65800, AT4G37770, AT4G08040, AT3G49700* | Biosynthesis | BLASTP, OrthoFinder |

| Cs10 ERG Locus* | Name | Chromosome | *A. thaliana* orthologs | Pathway | Identification method |
|---|---|---|---|---|---|
| LOC115704963 | *CsACS7-LIKE* | 1 | AT4G26200, AT1G01480, AT4G11280, AT4G08040, AT3G49700 | Biosynthesis | BLASTP, OrthoFinder |
| LOC115696400 | *CsACS9†* | X | AT4G08040, AT5G65800, AT4G37770, AT3G49700 | Biosynthesis | BLASTP |
| LOC115716540 | *CsACS9-LIKE* | 5 | AT5G65800, AT4G37770, AT4G08040, AT3G49700, AT2G22810 | Biosynthesis | BLASTP |
| LOC115697744 | *CsCTR1*ǂ† | X | AT5G03730 | Signalling | BLASTP |
| LOC115706236 | *CsCTR1-LIKE*ǂ† | 1 | AT5G03730 | Signalling | BLASTP, OrthoFinder |
| LOC115720532 | *CsEBF1*ǂ† | X | AT5G25350, AT2G25490 | Signalling | BLASTP, OrthoFinder |
| LOC115724585 | *CsEBF2*ǂ† | 6 | AT5G25350, AT2G25490 | Signalling | BLASTP |
| LOC115711236 | *CsEIL1*ǂ† | X | AT2G27050 | Signalling | OrthoFinder |
| LOC115719373 | *CsEIL3*ǂ† | 2 | AT1G73730 | Signalling | OrthoFinder, BLASTP |
| LOC115709621 | *CsEIN2*ǂ† | 3 | AT5G03280 | Signalling | BLASTP, OrthoFinder |
| LOC115706079 | *CsEIN3*ǂ† | 1 | AT2G27050, AT3G20770 | Signalling | BLASTP, OrthoFinder |
| LOC115723050 | *CsERF1* | 9 | AT3G23240 | Signalling | BLASTP, OrthoFinder |
| LOC115697291 | *CsERS1*ǂ† | 7 | AT2G40940 | Signalling | BLASTP, OrthoFinder |
| LOC115705474 | *CsETP1*ǂ† | 1 | AT3G18980 | Signalling | BLASTP |
| LOC115720323 | *CsETP2* | 2 | AT3G18910 | Signalling | BLASTP |
| LOC115702329 | *CsETR1*ǂ† | X | AT1G66340, AT2G40940 | Signalling | BLASTP, OrthoFinder |
| LOC115721785 | *CsETR2*ǂ† | 9 | AT3G04580, AT1G04310, AT3G23150 | Signalling | BLASTP, OrthoFinder |
| LOC115698184 | *CsRAN1*ǂ† | 7 | AT5G44790 | Signalling | BLASTP, OrthoFinder |
| LOC115707134 | *CsRTE1*ǂ† | X | AT2G26070 | Signalling | BLASTP, OrthoFinder |

| Cs10 ERG Locus[*] | Name | Chromosome | *A. thaliana* orthologs | Pathway | Identification method |
|---|---|---|---|---|---|
| LOC115721731 | *CsARD1*[‡†] | 9 | *AT5G43850, AT4G14716, AT4G14710, AT2G26400* | Yang Cycle | BLASTP, OrthoFinder |
| LOC115722194 | *CsARD2*[‡†] | 9 | *AT4G14716, AT2G26400, AT5G43850, AT4G14710* | Yang Cycle | BLASTP, OrthoFinder |
| LOC115706479 | *CsMTI1*[‡†] | 1 | *AT2G05830* | Yang Cycle | BLASTP, OrthoFinder |
| LOC115703014 | *CsMTK1*[‡†] | X | *AT1G49820* | Yang Cycle | BLASTP, OrthoFinder |
| LOC115702983 | *CsMTK1-LIKE*[‡†] | X | *AT1G49820* | Yang Cycle | BLASTP, OrthoFinder |
| LOC115704327 | *CsMTN*[‡†] | X | *AT4G34840, AT4G38800* | Yang Cycle | BLASTP, OrthoFinder |
| LOC115706293 | *CsDEP1*[‡†] | 1 | *AT5G53850* | Yang Cycle | BLASTP, OrthoFinder |
| LOC115711364 | *CsMAT1*[‡†] | X | *AT1G02500, AT4G01850, AT3G17390, AT2G36880* | Yang Cycle | BLASTP, OrthoFinder |
| LOC115719259 | *CsMAT2*[‡†] | 2 | *AT1G02500, AT4G01850, AT3G17390, AT2G36880* | Yang Cycle | BLASTP |
| LOC115711343 | *CsMAT3*[‡†] | 3 | *AT2G36880, AT3G17390, AT4G01850, AT1G02500* | Yang Cycle | BLASTP, OrthoFinder |
| LOC115700576 | *CsFYF1*[‡†] | 8 | *AT5G62165* | Ethylene Adjacent | BLASTP |
| LOC115706939 | *CsFYF2*[‡†] | 1 | *AT5G62165* | Ethylene Adjacent | BLASTP |
| LOC115716986 | *CsFYF3*[‡†] | 5 | *AT5G62165* | Ethylene Adjacent | BLASTP |
| LOC115717309 | *CsFYF4*[‡†] | 5 | *AT5G62165* | Ethylene Adjacent | BLASTP |
| LOC115701489 | *CsORE1*[‡†] | 8 | *AT5G39610* | Ethylene Adjacent | BLASTP, OrthoFinder |
| LOC115709772 | *CsORE2*[‡†] | 3 | *AT5G39610* | Ethylene Adjacent | BLASTP |

227    * Hyperlinks to the NCBI cs10 reference genome information page for the specific locus.

228    ‡ and † ERGs for which transcripts were detected from dataset 1 and/or in dataset 2, respectively.

229 § A gene located on an unclassified scaffold.

**Figure 2** *C. sativa* model of the Yang cycle and ethylene biosynthesis determined by homology analysis from *A. thaliana.* Products of enzymatic reactions are italicized. Genes encoding enzymes are in boxes. (a) Yang cycle- product abbreviations are as follows: *S-*adenosylmethionine (SAM), 5-methylthioadenosine (MTA), 5-methylthioribose (MTR), 5-methyltioriobose-1-P (MTR-1-P), 5-methylthioribulose-1-P (MTRu-1-P), 1,2-dihydroxy-3-keto-5-methylthiopentene (DHKMP), 2-keto-4-methylthiobutyrate (KMTB), methionine (Met). (b) Ethylene biosynthesis- product abbreviations are as follows: 1-aminocyclopropane-1-carboxilic acid (ACC). Differentially expressed genes are annotated with Mars and Venus symbols, identifying their proposed roles as either involved in sexual determinism (sex determination; denoted with diverging arrows) or sexual plasticity (denoted with reciprocal arrows) based on RNA-seq analysis.

**Figure 3** A simplified model of ethylene signalling in *C. sativa*, constructed through computational analysis using the *A. thaliana* pathway as a template. Pathway is depicted in the presence (a) and absence- (b) of ethylene. Dashed lines represent genes not found in *C. sativa*. Pointed arrows indicate promotion, while blunt-end arrows represent repression. The figure design is based on existing ethylene signalling models (Binder *et al.*, 2010; Dubois *et al.*, 2018; Binder, 2020). Differentially expressed genes are marked with Mars and Venus symbols, indicating their proposed roles in sexual determinism (sex determination, diverging arrows) or sexual plasticity (reciprocal arrows) based on RNA-seq analysis.

**Figure 4** Position of CsERGs in the cs10 reference genome identified from A. thaliana orthologs. (a) Distribution of CsERGs across the genome. Different colors were used to described their functional annotation. (b) Proportion of CsERGs distributed on autosomes or ChrX.An interactive version of (a) can be found in the on the OSF page for this project.

## Differential expression analysis reveals three patterns of expression

Among the 43 putative CsERGs analyzed, a total of 35 were found to be expressed in both dataset 1 and dataset 2, as indicated in Table 1. Upon further investigation, we found that among these 35 CsERGs, 16 exhibited differential expression between flower types in dataset 1, while 11 showed differential expression in dataset 2, as shown in Supplemental Figure 2. Seven CsERGs that consistently displayed differential expression between MFs and FFs were identified across both datasets, with comparable levels of magnitude, as illustrated in Supplemental Figure 3. These 7 CsERGs are considered high confidence genes involved in cannabis sex determination or sexual plasticity. In conducting pairwise comparisons of CsERG expression between different flower types, we observed three distinct patterns of differential expression. These patterns are described below and represent a newly proposed framework for understanding the role of CsERGs in sex determination and sexual plasticity. These patterns have been classified as karyotype concordant

(KC- sex determination; Figure 5), floral organ concordant (FOC- sexual plasticity; Figure 6), and
unique ERG (uERG- sexual plasticity; Figure 7).

KC expression refers to CsERGs that displayed no significant differences in expression
between plants with the same sex chromosome karyotype, such as FFs and IMFs. Among the 16
CsERGs exhibiting differential expression, 9 were identified to have KC expression, primarily in
dataset 1, as depicted in Figure 5. Conversely, FOC expression describes genes with similar
expression between MFs and IMFs, which possess the same floral organs, but that had significantly
differen expression when compared with FF. Out of the 16 differentially expressed CsERGs, five
displayed FOC expression, as indicated in Figure 6. Notably, three genes (*CsACS2, CsFYF2*, and
*CsFYF4*) exhibited higher expression in FFs compared to phenotypic male flowers, whereas two
genes (*CsETP2* and *CsMTN*) demonstrated higher expression in MFs and IMFs compared to FFs,
as illustrated in Figure 6. Lastly, uERG expression referred to CsERGs that exhibited significant
differential expression among each floral type, including FFs, MFs, and IMFs. Two CsERGs,
namely *CsACO5* and *CsFYF1*, were identified to possess uERG expression, as depicted in Figure
7. By proposing this framework, we aim to deepen our understanding of the involvement of
CsERGs in sex determination and sexual plasticity in cannabis.

**Figure 5** Differential expression analysis of *C. sativa* ERGs revealed nine genes with a karyotype concordant (KC) expression pattern. Among these genes, female flowers (FF; XX karyotype) and induced male flowers (IMF; XX karyotype) showed not significant (ns) differential expression. Significant pairwise comparisons are denoted by asterisks: ns = not significant; * for $p \leq 0.05$; ** for $p \leq 0.01$; *** for $\leq 0.001$ as determined by the Wald test.

**Figure 6** Differential expression analysis of *C. sativa* ERGs revealed five cases of genes showing floral-organ concordant (FOC) expression. FOC refer to genes whose expression pattern is shared between organisms with the same floral organ, but not necessarily the same sex-chromosome karyotype designated by ns in transcription levels for male flowers on XY plants and induced male flowers on XX plants. S0 Significant pairwise comparisons are denoted by asterisks: ns = not significant; * for p ≤ 0.05; ** for p ≤ 0.01; *** for ≤ 0.001 as determined by the Wald test.

**Figure 7** Differential expression analysis of revealed unique ERG (uERG) expression, which are cases where gene expression in the sex-changed plant does not match either that of the plant with which is shares the same sex chromosome karyotype or the plant with which is shares the same floral organs. Significant pairwise comparisons are denoted by asterisks: ns = not significant; * for $p \leq 0.05$; ** for $p \leq 0.01$; *** for $\leq 0.001$ as determined by the Wald test.

## ERGs are distributed genome-wide in *C. sativa*

The 43 CsERGs found in the homology analysis were distributed throughout the genome
(Figure 4 and Supplementary Table 3) independently of autosomal or sex chromosomes. A
majority of the CsERGs were located on autosomes (Figure 4b). Yang cycle genes were the most
numerous CsERGs present on the sex chromosomes, with 40% of the identified *C. sativa* Yang
cycle genes located on the X chromosome. In contrast, none of the genes that were classified as
ethylene adjacent (ERGs from the literature which do not belong to the canonical ethylene
biosynthesis or signalling pathways) were located on the sex chromosomes (Figure 4). Of the 43
CsERGs identified all mapped to chromosomes in the reference genome, except *CsACO1-LIKE1*,
which mapped to an unassigned scaffold (Figure 4a). CsERGS with significant differential
expression linked to sex determination (KC) and sexual plasticity (FOC and uERG) were found
throughout the genome (Figure 4a) and included genes from the Yang Cycle and ethylene
biosynthesis pathway (Figure 2) as well as the ethylene signalling pathway (Figure 3). Of the 16
significantly differentially expressed CsERGs, a minority, 11% and 29% exhibiting KC and
FOC/uERG expressions were distributed on the sex chromosome, respectively.

# Discussion

To date, sexual plasticity has only been discussed in the context of the animal kingdom (Liu *et al.*, 2017). While studies have focused on sex determination in angiosperms without sex chromosomes, such as the Cucurbitaceae family (Yamasaki *et al.*, 2001; Boualem *et al.*, 2015; Oda *et al.*, 2022), sexual plasticity in plants remains relatively unexplored. However, as a dioecious plant with sex chromosomes, cannabis represents a uncommon example of a plant species that exhibits sexual plasticity. Previous research has demonstrated the induction of male flowers in female plants through the inhibition of ethylene using STS (Lubell & Brand, 2018), and the induction of female flowers in male plants through ethylene (ethephon) treatment (Moon *et al.*, 2020a). Considering ethylene's known involvement in sex determination across plant species, it serves as a promising starting point for investigating sexual plasticity in dioecious plants like C. sativa (Hume & Lovell, 1983; Manzano *et al.*, 2014; Cebrián *et al.*, 2022). However, unraveling the specific mechanisms by which ethylene influences sexual plasticity presents challenges due to its multifaceted roles in plants.

To understand the influence of CsERGs on sex determination and sexual plasticity, we propose a framework consisting of three putative CsERG expression patterns: karyotype concordant (KC; Figure 5), floral organ concordant (FOC; Figure 6), and unique ERG (uERG; Figure 7). KC expression is similar between individuals with the same sex chromosome karyotype and CsERGs exhibiting KC expression are unaffected by STS treatments that induce sexual plasticity in *C. sativa*. This type of expression suggests that genes belonging to the KC case may have sex-specific patterns of expression, playing a role in sex determination. Conversely, FOC expression (Figure 6) is the same between individuals with shared floral organ sex, but carrying different sex chromosome karyotypes. This pattern suggests that CsERGs showing FOC expression are involved in sexual plasticity, as their expression appears uncoupled from the sex chromosome karyotypes. uERGs (Figure 7) shows a pattern of different levels of expression of CsERGs in all three treatments. Two genes, *CsFYF1* and *CsACO5*, showed uERG expression, with the gene expression in the sex-manipulated plant (IMF) not matching either of the plants with the same genotypic sex (FF) or the plant with the phenotypic sex (MF). Below, we discuss how each of these patterns is found throughout the biochemical journey of ethylene, from its precursor producing in the Yang cycle (Figure 2a) all the way through ethylene biosynthesis (Figure 2b) and

312    signalling (Figure 3). We also highlight how the distribution of these differentially expressed
313    CsERGs are not exclusively found on sex chromosomes, but rather distributed throughout the
314    genome (Figure 4). Lastly, we end by discussing limitations and future opportunities revealed by
315    this study.

**High expression of key Yang Cycles genes; a potential indicator of masculinity in *C. sativa***

In previous studies, it has been demonstrated that changes in the expression of Yang cycle
genes regulate ethylene production in plants. Here, we explore the potential role of the Yang cycle
in cannabis sex determination and sexual plasticity. In the present study, we found that two
CsERGs from the Yang cycle showed KC expression (Figure 2a and Figure 5), methionine
adenosyltransferase 3 (*CsMAT3*) and acireductone dioxygenase 2 (*CsARD2*). Most higher
eukaryotes possess small *ARD* gene families (Sauter *et al.*, 2005). *A. thaliana* has four *ARD* genes
isoforms (Pommerrenig *et al.*, 2011) from which we identified two *CsARD* genes homologs. In
the dataset 1, *CsARD2* was more highly expressed in MFs than FFs and IMFs (Figure 5). In
contrast *CsARD1* transcripts were found in all treatments, in both datasets, but expression did not
vary significantly between MFs, FFs or IMFs. These findings are consistent with previous studies
in rice that showed that the different ARD isoforms are regulated differently *in planta*, with
*OsARD1* undergoing a transient increase in expression in response to submergence, while *OsARD2*
is constitutively expressed, and unaffected by changes in ethylene concentration (Sauter *et al.*,
2005). Similarly, Liang *et al.* (2019) showed that transgenic rice overexpressing *OsARD1*
produced higher levels of ethylene versus wild-type rice.

Methionine adenosyltransferase (MAT, or SAM synthetase; SAMS) is responsible for the
biosynthesis of SAM, an important methyl donor and precursor for ethylene, lignin and polyamine
synthesis (Arraes *et al.*, 2015). From the four *AtMAT* used to construct the putative Yang cycle
model in cannabis, we identified three putative *CsMAT* genes: *CsMAT1, CsMAT2, CsMAT3,* of
which *CsMAT3* was found to have a KC expression. *CsMAT3* was most expressed in MFs, with
significantly lower transcription levels in FFs and IMFs. The association *MAT* genes with male
floral characteristics have been reported in *Arabidopsis* by Chen *et al.* (2016b) who found that
expression of *AtMAT3,* a *CsMAT3* ortholog, is responsible for normal pollen germination and
pollen tube formation. Although critical male fertility and pollen germination in *Arabidopsis*, the
role of *MAT* genes in controlling sex determination has yet to be reported. The present findings

that *CsMAT3* gene expression is higher in MFs than in FFs, suggest that it may play a role in sex determination. One possible way that *CsMAT3* may affect sex determination by altering levels of DNA methylation. SAM, the product of *MAT* is a methyl donor for DNA and histone methylation, and it has been reported that *mat4* knockouts in *Arabidopsis* have lower genome-wide methylation levels as well as decreased histone methylation (Meng *et al.*, 2018). Similar findings have been echoed by Chen *et al.* (2016b) who reported that *MAT3* was required for proper histone and tRNA methylation. It is possible that higher expression of *CsMAT3* in male plants could lead to increased methylation of key genes involved in plant masculinity. To date, there has been a scarcity of research on the methylation patterns of cannabis (Aina *et al.*, 2004; Mayer *et al.*, 2015), with no studies investigating its role in sex determination and sexual plasticity. Investigating the methylation status of floral tissues in cannabis may yield valuable insights into the mechanisms underlying sex determination and sex plasticity in this species.

Here, we report the identification of one CsERG from the Yang cycle, *CsMTN,* with a FOC expression. MTN recycles MTA after the first step of the ethylene biosynthesis cycle where SAM is converted to ACC by ACS. The ortholog analysis for both *AtMTN1* and *AtMTN2* revealed a single ortholog in cannabis (Table 1), which is similar to both rice (*Oryza sativa* L.) and tomato (*Solanum lycopersicum* L. Karst. ex Farw) in which MTN is only encoded by one gene (Rzewuski *et al.*, 2007; van de Poel *et al.*, 2012). In Arabidopsis, *MTN* shows preferential expression in the phloem (Pommerrenig *et al.*, 2011), suggesting that tissue sampling may also play a role in the expression patterns of *MTN*. In dataset 1 *CsMTN* transcripts were most abundant in MFs (Supplemental Figure 3), and in dataset 2 (Figure 6) *CsMTN* transcripts were significantly more expressed in MFs and IMFs than in FFs, suggesting that high abundance of *CsMTN* may be associated with masculinity and sexual plasticity in cannabis.

**Multiple biosynthesis genes linked to sexual plasticity**

Ethylene biosynthesis is regulated by a set of conserved enzymes that are found in all plant species. The key enzyme in this two-step process is ACS, which converts SAM, derived from the Yang cycle, into ACC. ACC is then converted into ethylene by the enzyme ACO. These enzymes are conserved across all plant species, and their regulation is tightly controlled by various factors such as hormonal and environmental cues (Bleecker & Kende, 2000). The ortholog analysis found that *ACS* and *ACO* belong to multi-gene families in cannabis, similar to Arabidopsis. In *C. sativa*,

6 *ACS* and 5 *ACO* gene homologs were identified from the 8 *ACS* and 5 *ACO* genes in *A. thaliana*

respectively (Figure 2 and Table 1). The presence of *ACS* and *ACO* are critical to ethylene

biosynthesis, but the number of genes in each family can vary between species. For example, in

*Pyrus ussuriensis* L. (pear) 13 *ACS* and *11 ACO* genes have been identified (Yuan *et al.*, 2020), in

*Zea mays* L. 5 *ACS* and 15 *ACO* genes have been identified, in rice 6 *ACS* and 9 *ACO* genes, and

in tomato there are 9 *ACS* and 7 *ACO* genes (Houben & Van de Poel, 2019; Wu *et al.*, 2022). The

importance of having multiple functionally redundant ethylene biosynthesis genes has been

demonstrated in multiple plants, with studies showing *ACS* and *ACO* genes are expressed at

different levels and in different tissues as a function of the plant organ's needs and developmental

stage (van de Poel *et al.*, 2012; Van de Poel *et al.*, 2014). In ripening tomatoes van de Poel *et al.*

(2012) reported a higher level of *ACS6, ACO2* and *ACO4* expression during baseline (system 1)

ethylene production during immature fruit development prior to ripening, but that auto-catalytic

increases in ethylene during ripening (system 2) were correlated with increases in *ACS2*, *ACS4*,

*ACO3* and *ACO5* expression, while abundance of *ACO2* and *ACO4* transcripts decreased at that

same stage. The induction/repression of specific *ACS* genes has also been shown to play a role in

sex determination of some *Cucurbitaceae* species (Manzano *et al.*, 2014; Boualem *et al.*, 2015;

Chen *et al.*, 2016a; Zhang *et al.*, 2017). One study found that expression of *ACS11* was only

observed in the vascular bundles of female flowers of monoecious and in hermaphroditic flowers

of andromonoecious plants (plants producing male and hermaphrodite flowers), pointing to the

role of *ACS11* in carpel development (Boualem *et al.*, 2015). Changes in sex determination have

also been shown in cucumber and melon by way of disruption of a specific *ACO* genes (*ACO2* in

cucumber; *ACO3* in melon) resulting in decreased ethylene production in the carpel region of

flowers and causing a shift from a monoecy to androecy resulting in unisexual flower production

in cucumber and melons (Chen *et al.*, 2016a). The existence of multiple, functionally redundant

*ACS* and *ACO* genes, combined with spatial and temporal patterns of expression allows for the

fine tuning of ethylene biosynthesis and giving a plant a high level of control over its development,

including effects on a plant's sex determination. The present study demonstrates evidence of

numerous *ACS* and *ACO* genes distributed throughout the genome of *C. sativa* (Table 1 and Figure

4). Notably, the expression patters of *CsACS2* (Figure 6)*, CsACO3* (Figure 5) and *CsACO5* (Figure

7) represent the three distinct cases of gene expression proposed in this study. These findings

collectively suggest a diverse range of effects exerted by the ethylene biosynthesis pathway on
cannabis sex determination and sexual plasticity.

*CsACS2*, which is part of the *ACS* gene family responsible for the first step of ethylene
biosynthesis, shows a FOC expression linking it to sexual plasticity. *ACS* has been demonstrated
to catalyze the rate-limiting reaction in most cases of ethylene production (van de Poel *et al.*, 2012;
Houben & Van de Poel, 2019; Pattyn *et al.*, 2021). This gene is downregulated in MFs and IMFs
in comparison to FFs. Comparisons in *CsACS2* expression show similar low expression levels
between MFs and IMFs despite their different genotypic sex. Since *ACS* is thought to be rate-
limiting, and given the association of ethylene with 'feminization' of floral phenotypes we
hypothesize that the downregulation of *CsACS2* in IMF relative to FF, points to reduction in
ethylene biosynthesis *in planta* which may play a role in the induction of male flowers in
genetically female plants. *CsACO3* shows KC expression, with activity in MFs much higher for
both than in IMF and FFs. Further study which incorporates multiple tissue sources could prove
helpful in understanding *CsACO3*'s KC expression and how this plays a putative role in sex
determination, as it remains to be seen if this expression pattern is unique to floral tissues or if this
KC expression is also present in vegetative tissues of flowering plants prior to and during early
floral development. Lastly, *CsACO5* has a unique expression pattern in all treatments (uERG;
Figure 7), with a significantly larger expression in IMFs compared with MFs or FFs. Here, dosage
compensation could play a role with downstream inhibition of ethylene signalling by STS leading
to increased transcription of *ACO* as IMF plants which are genetically female, fail to perceive
ethylene levels normally required for female floral development.

**Ethylene signalling is conserved in *C. sativa***

Ethylene signalling has also been highly conserved across the plant kingdom (Chang,
2016). Ethylene binds to a family of receptors known as ETRs, which are found in the membrane
of the endoplasmic reticulum of plant cells (Binder, 2020). These receptors are responsible for
transducing the ethylene signal to downstream signalling components. The downstream signalling
pathway involves a series of conserved components, including *CTR1*, *EIN2*, and *EIN3*, which are
responsible for activating the expression of ethylene-responsive genes (Shakeel *et al.*, 2013). In
the present study the putative ethylene signalling pathway in *C. sativa* was identified based on
orthology with Arabidopsis genes (Figure 3). Transcriptomes obtained from MFs, FFs and IMFs

revealed numerous CsERGs adjacent to, or directly involved in the signalling pathway which have

FOC expression (*CsETP2, CsFYF2* and *CsFYF4*) and uERG (*CsFYF1*). In *Arabidopsis EIN2*

*TARGETING PROTEIN1* and *EIN2 TARGETING PROTEIN2* (*ETP1/2*) act to suppress ethylene

responses in the signalling pathway by carrying out ubiquitinating of *EIN2*, marking *EIN2*

degradation (Qiao *et al.*, 2009; Binder, 2020). When ethylene is present, *ETP1/2* is supressed,

allowing ethylene signalling to proceed (Qiao *et al.*, 2009; Yang *et al.*, 2015). In the present study,

both *CsETP1* and *CsETP2* transcripts were detected, however only *CsETP2* had an FOC

expression associated with sexual plasticity (Figure 3). *CsETP2* expression was lowest in FFs

compared to MFs and IMFs (Figure 6). There was no significant difference between *CsETP2*

expression in MFs and IMFs and both had higher expression than FF tissues. These patterns of

gene expression fit with the hypothesized role of ethylene as a feminizing agent. If increased

endogenous ethylene is responsible for femaleness, then we should expect lower levels of *CsETP2*

in female flower producing plants, as observed.

The differential expression analysis also revealed three *FOREVER YOUNG FLOWER*

(*FYF*) genes which belong to the FOC expression (*CsFYF2* and *CsFYF4*) and uERG (*CsFYF1*)

expression. *FYF* genes are expressed in young flowers prior to pollination (Chen *et al.*, 2011) and

acts as a suppressor of floral abscission. They act by repressing *Ethylene Response DNA-binding*

*Factors (EDFs)* found downstream *ERFs* that trigger floral senescence and abscission in maturing

flowers (Chen *et al.*, 2015, 2022). In both datasets, expression of *CsFYF1* and *CsFYF2* was highest

in  FFs compared with MFs  (Supplemental Figure 3) and highest in FFs when compared with

IMFs present in dataset 1 (Figure 6 and Figure 7). While *FYF* expression generally decreases over

time in other plants such as orchids or Arabidopsis (Chen *et al.*, 2011, 2021), their expression

patterns have not been explored in relation to sex expression in those species. These present

expression patterns suggest that *FYF* may vary depending with sexual plasticity in dioecious

cannabis, but further investigation is needed to determine causality. It is also possible that the

discrepancy in *CsFYF*s levels between FF and other tissues is influenced by the differing lifespans

of male and female cannabis flowers. Female cannabis flowers, which have a longer lifespan, may

maintain higher levels of *CsFYF* to suppress senescence caused by ethylene. Exploring

endogenous ethylene levels and studying the expression of *CsFYF* genes throughout flower

development could provide more insights into these observed expression patterns.

Among          the          signalling          genes          analysed,
*CsERS1, CsEIL1* and *CsEIL3* exhibited KC expression (Figure 5). Both IMF and FF displayed

similar, lower levels of *CsERS1* expression compared to MFs, indicating a potential association

between high *CsERS1* expression and male plants in *C. sativa*. However, it is worth noting that

these observations contrast with a previous study by Yamasaki *et al.* (2000), which reported higher

transcript levels of ethylene receptors *ETR1*, *ETR2*, and *ERS* in gynoecious cucumbers compared

to monoecious cucumbers, indicating a possible connection between receptor expression and a

preference for female flower production. Ethylene receptors have been the subject of extensive

study in Arabidopsis, which has five isoforms of ethylene receptors. These are divided into two

subfamilies, with *ERS1* and *ETR1* belongs to the first subfamily and *ERS2*, *ETR2* and *EIN4*

belonging to the second subfamily (Binder, 2020). Presently no orthologs of *AtEIN4* or *AtERS2*

were identified  in cannabis (Figure 3). *ETR1* from subfamily 1 has largely been the focus of

existing studies on ethylene perception and plant sex determination, with one study finding that

*ETR1* alone could mediate silver induced inhibitions of ethylene response (McDaniel & Binder,

2012) while another study in *Cucurbita pepo* found that *ETR1* was responsible control for sex

determination (García *et al.*, 2020). Liu *et al.* (2012) also demonstrated that *etr1-1*$^{(C65Y)}$

Arabidopsis mutants had reduced ethylene responses independently of other receptors,

highlighting the significant role of *ETR1* in ethylene signalling. However, in both datasets analysed

*CsETR1* was not differentially expressed in any of the flower types. In contrast, *ERS1* which

exhibited KC expression in cannabis, has been described to have a relatively modest role as an

ethylene receptor in other plants species. Studies suggest that *ERS1* may act in coordination with

other ethylene receptors to affect signalling (Liu & Wen, 2012) and that its response to silver ions

is much smaller than *ETR1* (McDaniel & Binder, 2012).

Ethylene insensitive-3 (*EIN3*)/Ethylene insensitive-3-like (*EIL*) family is a small family of

transcription factors involved in ethylene signalling (Binder, 2020). We found that *CsEIL1* and

*CsEIL3* were differentially expressed between flower types. *EILs* have been shown to play a role

in physiological processes which rely on ethylene, including responses to environmental stresses

(Liu *et al.*, 2019), sex determination (Li *et al.*, 2021) and floral development (Zhu *et al.*, 2022).

*EIN3/EIL1* transcription factors have been implicated with increasing the ratio of female flowers

in monoecious pumpkin in response to ethephon application on seedlings (Li *et al.*, 2021) and

EIN3/EIL1 expression triggered by ethylene causes developmental defects and pollen sterility in

anthers of *Arabidopsis*. We found higher levels of *EIL1* in XX flowers than in those of XY plants,
and this raises the question as to whether high levels of *EIL1* detected in IMFs may affect pollen
viability of induced males. In a study which overexpressed mulberry *EIL3* in Arabidopsis, the
upregulation of *EIL3* also coincided with increases in ethylene biosynthesis; however, *EIL1* and
*EIL2* did not follow *EIL3* expression under drought and salt stress, suggesting that expression
within this family of transcription factors can vary (Liu *et al.*, 2019). These findings are similar
to ours, where *CsEIL1* showed clear KC expression, with highest levels of expression in FFs and
IMFs, whereas *CsEIL3* showed slightly more elevated expression in MFs compared to FFs (Figure
5).

Our results highlight the differential expression of *CsERS1* and *CsEILs* between male and
female flowers, supporting their involvement in sex determination mechanisms. Future studies
should explore the functional significance of *ERS1* and *EILs* in sex determination pathways and
investigate the cooperative interactions between *ERS1* and other ethylene receptors and
downstream signalling to transcription factors in these processes. Moreover, further investigations
are warranted to elucidate the specific mechanisms underlying the differential responses to silver
ions among receptor subfamilies and their implications for ethylene perception.

## 509　Genes linked to sexual plasticity found throughout the genome

*C. sativa* contains a pair of heteromorphic sex chromosomes. However, treatment of male
plants with ethylene and female plants with the ethylene inhibitor STS has been reported to cause
plants to produce flowers opposite than that of the sex chromosome karyotype (Green, 2005;
Lubell & Brand, 2018; Moon *et al.*, 2020a). The observed sexual plasticity in the species suggests
that cannabis sex is not solely determined by the sex chromosomes, but rather by a combination
of genetic and environmental factors (GSD+ESD), as has been suggested in the study of
mammalian sexual plasticity (Lambert *et al.*, 2019). There are few studies in cannabis which
explore the link between sex chromosomes and sex-expression, and to the authors' knowledge, no
studies which explicitly explore molecular mechanisms linking cannabis sexual plasticity and
ethylene. In a study comparing the transcriptomes of male and female flowers, Prentout *et al.*
(2020) reported that 35% of sex-linked genes mapped to autosomes while 65% mapped to the sex
chromosomes of *C. sativa*. The distribution of their reported sex-linked genes in the chromosome
is in stark contrast to the distribution of CsERGs that we identified with only 26.2% of CsERGs

mapping to the sex chromosomes. When considering only CsERGs which are differentially expressed, 20% (3/15) are located on the sex chromosomes, while the remainder are located on the autosomes (Figure 4). The ubiquity of ERGs in the autosomes is to be expected, as they play important roles in general plant growth with ethylene regulating a number of plant growth responses. Prentout *et al.* (2020) suggested that their observation of autosomal mapping for sex-linked genes might have been the result of false positives or incorrect placement of genes in the genome assembly of *C. sativa*. However, their assumption appears to overlook the possibility that many genes involved in plant sex might also be crucial for the production of essential growth regulators or metabolites, which are required in both genetically male and female *C. sativa* and whose loci are predominantly found on autosomes. In contrast, Adal *et al.* (2021) proposes an X:autosome dosage mechanism as the basis for sex determination, where male flowers might possess a limited number of unique Y-chromosome genes influencing their sex. We propose that the regulation of cannabis sexual plasticity is mediated by the differential expression of CsERGs, encompassing both FOC and uERG expression, on both autosomes and sex chromosomes of cannabis. The presence of CsERGs on autosomes aligns with ethylene's involvement in general plant metabolism, while their presence on sex chromosomes suggests some specific roles in sexual development. By considering the comprehensive distribution of CsERGs across the genome, we highlight the importance of both autosomal and sex chromosome-associated genes in the regulation of cannabis sexual plasticity.

## Limitations and future directions

Based on previous observations suggesting a link between ethylene and sexual plasticity/determinism, the present study constitutes a preliminary analysis to identify putative candidate genes involved in the sexual expression of cannabis. However, in the case of this exploratory study, there are some limitations that we would like to acknowledge. Only two transcriptomic datasets from floral tissues were available that contained multiple flower sexes (Prentout *et al.*, 2020; Adal *et al.*, 2021). In an effort to begin probing the effect of CsERGs during chemically induced sex plasticity, it was important that some of the transcriptomes include IMFs, which were only available in one of the two datasets used (Adal *et al.*, 2021). Additionally, no existing transcriptomes were available that contained IFFs, which limited our ability to test whether there is an inverse relationship in the mechanisms for sexual plasticity between MFs and

IFFs. Future studies on sexual plasticity in cannabis should include IFFs in order to provide a more

complete picture of the mechanisms underlying sexual plasticity in the species. Despite these

limitations, the differential expression findings between MF and FF in both datasets was used to

to find a set of differentially expressed CsERGs (Supplemental Figure 1 and Supplemental Figure

3), which are strong candidate CsERGs with a role in sex determination.

One limitation of the study is the use of datasets collected at a single timepoint, specifically

from flowers at a later stage of development. This approach may result in the potential omission

of differentially expressed CsERG candidates that are not active during that specific timepoint,

particularly those involved in early floral sex determination or sex differentiation. Furthermore, it

is important to acknowledge that some of the identified candidates may exhibit differential

expression at the selected timepoint for reasons unrelated to the intended analysis, which aims to

establish their role in sexuality. The datasets employed in this study were generated under different

experimental conditions, including variations in plant growth conditions, sampling stages of flower

development, and sampling locations on the plants. It is crucial to note that the datasets were

derived from distinct genotypes of *C. sativa*: one drug-type (cv. MS-17-338; Adal *et al.*, 2021) and

one hemp-type cannabis (cv. Zenitsa; Prentout *et al.*, 2020). As cannabis is known for its high

genetic variability (Lapierre *et al.*, 2023), studies based on a single genotype lack the robustness

necessary for generalization (Page *et al.*, 2021; Monthony *et al.*, 2021b). Therefore, future studies

on *C. sativa* should aim to include a broader range of cannabis genotypes, particularly those

exploring the role of ethylene in plasticity and sex determination. Notably, despite these

limitations, the transcriptomes from the two different cannabis populations facilitated the

validation of 34 out of the 43 predicted CsERGs listed in Table 1. This validation adds robustness

to the gene models proposed in this study for the Yang Cycle, ethylene biosynthesis, and signaling

pathways (Figure 2 and Figure 3). It is important to emphasize that the lack of expression of certain

predicted CsERGs in both datasets does not necessarily imply that they are non-functional

CsERGs. As previously discussed, the inclusion of transcriptomes covering a wider timeline,

including early floral induction stages, will contribute to the comprehensive validation of these

predicted models.

There are currently a limited number of biotechnological tools (e.g., *in vitro* culture, CRISPR,

VIGS, agrobacterium, etc.) which are optimized in order to functionally validate CsERGs in

cannabis (Hesami *et al.*, 2021) and many of these tools have been hampered by the lack of a reliable
*in vitro* regeneration protocol for the species (Monthony *et al.*, 2021b). Although such functional
validation is beyond the scope of the current study, it represents an important next step in probing
the role of ethylene in cannabis sexual plasticity. Recently progress has been made in establishing
methods for gene silencing in cannabis, through the employment of the *Cotton leaf crumple virus*
(*CLCrV*) in a virus-induced gene silencing protocol reported by Schachtsiek *et al.* (2019) and
through new methods using agrobacterium (Sorokin *et al.*, 2020; Galán-Ávila *et al.*, 2021).
However due to their recent nature, these methods have yet to be replicated independently. Given
the genotypic variability of cannabis and the role it plays in the replicability of *in vitro* methods
(Monthony *et al.*, 2021b), it remains to be seem how effectively these methods are adapted to
multiple genotypes across different research settings.

## 594   **Conclusion**

This study represents the first step in exploring sexual plasticity in the dioecious plant
*C. sativa* and the important role that Yang cycle, ethylene biosynthesis and signalling pathways
play in sex determination. Here, we present a framework consisting of three distinct gene
expression patterns (KC, FOC and uERG) which can be used to categorize the role of CsERGs in
sexual plasticity in dioecious plants. These findings are based on our analysis of transcriptomes of
male, female and induced male flowers obtained from two different *Cannabis sativa* genotypes
produced under two different experimental conditions. From the ortholog analysis undertaken in
this study and with support from transcriptomic data, we propose that the canonical Yang Cycle
and ethylene biosynthesis and signalling pathways are conserved in *C. sativa* and we identified
seven genes belonging to two the FOC and uERG classes which we hypothesize play a role in
ethylene-induced sexual plasticity in cannabis. This work lays the groundwork for more detailed
study of the role of CsERGs in sexual plasticity in multiple genotypes of cannabis. Given the
commercial importance of cannabis, as well as a number of other dioecious plants such as kiwi,
papaya and persimmon, understanding the underlying factors of sexual plasticity and its link with
ethylene are important to future efforts in breeding and crop improvement.

# Acknowledgments

611 The authors gratefully acknowledge the support of the Natural Sciences and Engineering Research
612 Council (NSERC) of Canada. ASM has also been supported by a NSERC Canada Vanier Graduate
613 Scholarship. The authors would also like to thank Eric Normandeau for his support with script
614 writing.

# Competing Interests

The authors have no competing interests to declare.

# Author Contributions:

Conceptualization: Adrian S. Monthony

Data Curation: Adrian S. Monthony

Formal Analysis: Adrian S. Monthony, Maxime de Ronne

Funding Acquisition: Adrian S. Monthony, Davoud Torkamaneh

Investigation: Adrian S. Monthony, Davoud Torkamaneh

Methodology: Adrian S. Monthony, Maxime de Ronne, Davoud Torkamaneh

Project Administration: Adrian S. Monthony, Davoud Torkamaneh

Resources: Davoud Torkamaneh

Software: Adrian S. Monthony, Maxime de Ronne

Supervision: Davoud Torkamaneh

Visualization: Adrian S. Monthony

Writing- Original Draft Preparation: Adrian S. Monthony, Davoud Torkamaneh

Writing- Review & Editing: Adrian S. Monthony, Maxime de Ronne, Davoud Torkamaneh

# Data Availability Statement:

All scripts and raw data used in this study are available from The Open Science Framework (OSF) under the DOI:  DOI 10.17605/OSF.IO/7G9VR at OSF.io. or from the NCBI, under previously published projects (dataset 1 and dataset 2).

# References

**Adal AM, Doshi K, Holbrook L, Mahmoud SS**. **2021**. Comparative RNA-Seq analysis reveals genes associated with masculinization in female Cannabis sativa. *Planta* **253**: 1–17.

**Aina R, Sgorbati S, Santagostino A, Labra M, Ghiani A, Citterio S**. **2004**. Specific hypomethylation of DNA is induced by heavy metals in white clover and industrial hemp. *Physiologia Plantarum* **121**: 472–480.

**Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ**. **1990**. Basic local alignment search tool. *Journal of Molecular Biology* **215**: 403–410.

**Anand L, Rodriguez Lopez CM**. **2022**. ChromoMap: an R package for interactive visualization of multi-omics data and annotation of chromosomes. *BMC Bioinformatics* **23**: 1–9.

**Andrews S**. **2010**. FastQC: A Quality Control Tool for High Throughput Sequence Data.

**Argueso CT, Hansen M, Kieber JJ**. **2007**. Regulation of ethylene biosynthesis. *Journal of Plant Growth Regulation* **26**: 92–105.

**Arraes FBM, Beneventi MA, Lisei de Sa ME, Paixao JFR, Albuquerque EVS, Marin SRR, Purgatto E, Nepomuceno AL, Grossi-de-Sa MF**. **2015**. Implications of ethylene biosynthesis and signaling in soybean drought stress tolerance. *BMC Plant Biology* **15**: 213.

**Ashby R, Forêt S, Searle I, Maleszka R**. **2016**. MicroRNAs in Honey Bee Caste Determination. *Scientific Reports* **6**: 1–15.

**Barcaccia G, Palumbo F, Scariolo F, Vannozzi A, Borin M, Bona S**. **2020**. Potentials and challenges of genomics for breeding *Cannabis* cultivars. *Frontiers in Plant Science* **11**: 1–19.

**Binder BM**. **2020**. Ethylene signaling in plants. *Journal of Biological Chemistry* **295**: 7710–7725.

**Binder BM, Rodríguez FI, Bleecker AB**. **2010**. The copper transporter RAN1 is essential for biogenesis of ethylene receptors in Arabidopsis. *Journal of Biological Chemistry* **285**: 37263–37270.

**Bleecker AB, Estelle MA, Somerville C, Kende H**. **1988**. Insensitivity to Ethylene Conferred

by a Dominant Mutation in Arabidopsis thaliana. *Science* **241**: 1086–1089.

**Bleecker AB, Kende H**. **2000**. Ethylene: A gaseous signal molecule in plant. *Annual Review of*
*Cell and Developmental Biology* **16**: 1–18.

**Boualem A, Troadec C, Camps C, Lemhemdi A, Morin H, Sari MA, Fraenkel-Zagouri R,**
**Kovalski I, Dogimont C, Perl-Treves R,** *et al.* **2015**. A cucurbit androecy gene reveals how
unisexual flowers develop and dioecy emerges. *Science* **350**: 688–691.

**Brenner WG, Romanov GA, Köllmer I, Bürkle L, Schmülling T**. **2005**. Immediate-early and
delayed cytokinin response genes of Arabidopsis thaliana identified by genome-wide expression
profiling reveal novel cytokinin-sensitive processes and suggest cytokinin action through
transcriptional cascades. *Plant Journal* **44**: 314–333.

**Brewer C, Harrower M, Sheesley B, Woodruff A, Heyman D**. **2002**. ColorBrewer 2.0.

**Bürstenbinder K, Rzewuski G, Wirtz M, Hell R, Sauter M**. **2007**. The role of methionine
recycling for ethylene synthesis in Arabidopsis. *Plant Journal* **49**: 238–249.

**Cai Z, Cao C, Ji L, Ye R, Wang D, Xia C, Wang S, Du Z, Hu N, Yu X,** *et al.* **2020**. RIC-seq
for global in situ profiling of RNA–RNA spatial interactions. *Nature* **582**: 432–437.

**Cardoso-Júnior CAM, Fujimura PT, Santos-Júnior CD, Borges NA, Ueira-Vieira C,**
**Hartfelder K, Goulart LR, Bonetti AM**. **2017**. Epigenetic modifications and their relation to
caste and sex determination and adult division of labor in the stingless bee Melipona scutellaris.
*Genetics and Molecular Biology* **40**: 61–68.

**Carey SB, Jenkins J, Lovell JT, Maumus F, Sreedasyam A, Payton AC, Shu S, Tiley GP,**
**Fernandez-Pozo N, Healey A,** *et al.* **2021**. Gene-rich UV sex chromosomes harbor conserved
regulators of sexual development. *Science Advances* **7**.

**Cebrián G, Iglesias-Moya J, Romero J, Martínez C, Garrido D, Jamilena M**. **2022**. The
Ethylene Biosynthesis Gene CpACO1A: A New Player in the Regulation of Sex Determination
and Female Flower Development in Cucurbita pepo. *Frontiers in Plant Science* **12**: 1–16.

**Chang C**. **2016**. Q&A: How do plants respond to ethylene and what is its importance? *BMC*
*Biology* **14**: 7.

**Chao Q, Rothenberg M, Solano R, Roman G, Terzaghi W, Ecker JR**. **1997**. Activation of the

ethylene gas response pathway in arabidopsis by the nuclear protein ETHYLENE-

INSENSITIVE3 and related proteins. *Cell* **89**: 1133–1144.

**Charlesworth D, Mank JE**. **2010**. The birds and the bees and the flowers and the trees: Lessons

from genetic mapping of sex determination in plants and animals. *Genetics* **186**: 9–31.

**Chen MK, Hsu WH, Lee PF, Thiruvengadam M, Chen HI, Yang CH**. **2011**. The MADS box

gene, FOREVER YOUNG FLOWER, acts as a repressor controlling floral organ senescence and

abscission in Arabidopsis. *Plant Journal* **68**: 168–185.

**Chen WH, Jiang ZY, Hsu HF, Yang CH**. **2021**. Silencing of FOREVER YOUNG FLOWER-

Like Genes from Phalaenopsis Orchids Promotes Flower Senescence and Abscission. *Plant and*

*Cell Physiology* **62**: 111–124.

**Chen WH, Li PF, Chen MK, Lee YI, Yang CH**. **2015**. FOREVER YOUNG FLOWER

negatively regulates ethylene response DNA-binding factors by activating an ethylene-

responsive factor to control arabidopsis floral organ senescence and abscission. *Plant Physiology*

**168**: 1666–1683.

**Chen WH, Lin PT, Hsu WH, Hsu HF, Li YC, Tsao CW, Hsu MC, Mao WT, Yang CH**.

**2022**. Regulatory network for FOREVER YOUNG FLOWER-like genes in regulating

Arabidopsis flower senescence and abscission. *Communications Biology* **5**.

**Chen H, Sun J, Li S, Cui Q, Zhang H, Xin F, Wang H, Lin T, Gao D, Wang S,** *et al.* **2016a**.

An ACC Oxidase Gene Essential for Cucumber Carpel Development. *Molecular Plant* **9**: 1315–

1327.

**Chen Y, Zou T, McCormick S**. **2016b**. S-adenosylmethionine synthetase 3 is important for

pollen tube growth. *Plant Physiology* **172**: 244–253.

**Cheng WH, Chiang MH, Hwang SG, Lin PC**. **2009**. Antagonism between abscisic acid and

ethylene in Arabidopsis acts in parallel with the reciprocal regulation of their metabolism and

signaling pathways. *Plant Molecular Biology* **71**: 61–80.

**Coulter JM, Barnes CR (Eds.)**. **1901**. *Botanical Gazette*. Chicago: University of Chicago Press.

**Dhakal S, Karki M, Subedi P, GC A**. **2019**. Effect Of Ethephon Doses On Vegetative
Characters, Sex Expression and Yield Of Cucumber (Cucumis sativus cv. Bhaktapur Local) In
Resunga Municipality, Gulmi, Nepal. *International Journal of Applied Sciences and*
*Biotechnology* **7**: 370–377.

**Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M,**
**Gingeras TR**. **2013**. STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* **29**: 15–21.

**Dubois M, Van den Broeck L, Inzé D**. **2018**. The Pivotal Role of Ethylene in Plant Growth.
*Trends in Plant Science* **23**: 311–323.

**Emms DM, Kelly S**. **2019**. OrthoFinder: Phylogenetic orthology inference for comparative
genomics. *Genome Biology* **20**: 1–14.

**Ferfuia C, Zuliani F, Danuso F, Piani B, Cattivello C, Dorigo G, Baldini M**. **2021**.
Performance and stability of different monoecious hemp cultivars in a multi-environments trial
in north-eastern Italy. *Agronomy* **11**: 1–15.

**Flajšman M, Slapnik M, Murovec J**. **2021**. Production of Feminized Seeds of High CBD
Cannabis sativa L. by Manipulation of Sex Expression and Its Application to Breeding. *Frontiers*
*in Plant Science* **12**: 1–12.

**Galán-Ávila A, Gramazio P, Ron M, Prohens J, Herraiz FJ**. **2021**. A novel and rapid method
for Agrobacterium-mediated production of stably transformed Cannabis sativa L. plants.
*Industrial Crops and Products* **170**: 113691.

**Gane R**. **1934**. Production of Ethylene by Some Ripening Fruits. *Nature* **134**: 1008–1008.

**García A, Aguado E, Martínez C, Loska D, Beltrán S, Valenzuela JL, Garrido D, Jamilena**
**M**. **2020**. The ethylene receptors CpETR1A and CpETR2B cooperate in the control of sex
determination in Cucurbita pepo. *Journal of Experimental Botany* **71**: 154–167.

**Grassa CJ, Weiblen GD, Wenger JP, Dabney C, Poplawski SG, Timothy Motley S, Michael**
**TP, Schwartz CJ**. **2021**. A new Cannabis genome assembly associates elevated cannabidiol
(CBD) with hemp introgressed into marijuana. *New Phytologist* **230**: 1665–1679.

**Green G**. **2005**. *The Cannabis Breeder's Bible- The definitive guide to marijuana genetics,*

*cannabis botany and creating strains for the seed market*. San Francisco: Green Candy Press.

**Guo H, Ecker JR**. **2003**. Plant responses to ethylene gas are mediated by SCFEBF1/EBF2-

dependent proteolysis of EIN3 transcription factor. *Cell* **115**: 667–677.

**Guzman P, Ecker JR**. **1990**. Exploiting the Triple Response of Arabidopsis to Identify

Ethylene-Related Mutants. *The Plant Cell* **2**: 513.

**Hayes TB, Khoury V, Narayan A, Nazir M, Parka A, Brown T, Adame L, Chan E,**

**Buchholz D, Stueve T, *et al.* 2010**. Atrazine induces complete feminization and chemical

castration in male African clawed frogs (Xenopus laevis). *Proceedings of the National Academy*

*of Sciences of the United States of America* **107**: 4612–4617.

**Hesami M, Baiton A, Alizadeh M, Pepe M, Torkamaneh D, Jones AMP**. **2021**. Advances

and Perspectives in Tissue Culture and Genetic Engineering of Cannabis. *International Journal*

*of Molecular Sciences* **22**: 5671.

**Hirayama T, Kieber JJ, Hirayama N, Kogan M, Guzman P, Nourizadeh S, Alonso JM,**

**Dailey WP, Dancis A, Ecker JR**. **1999**. RESPONSIVE-TO-ANTAGONIST1, a Menkes/Wilson

disease-related copper transporter, is required for ethylene signaling in Arabidopsis. *Cell* **97**:

383–393.

**Houben M, Van de Poel B**. **2019**. 1-aminocyclopropane-1-carboxylic acid oxidase (ACO): The

enzyme that makes the plant hormone ethylene. *Frontiers in Plant Science* **10**: 1–15.

**Hume RJ, Lovell PH**. **1983**. The control of sex expression in cucurbits by ethephon. *Annals of*

*Botany* **52**: 689–695.

**Kende H**. **1998**. Plant Biology and the Nobel Prize. *Science* **282**: 627–627.

**Kieber JJ, Rothenberg M, Roman G, Feldmann KA, Ecker JR**. **1993**. CTR1, a negative

regulator of the ethylene response pathway in arabidopsis, encodes a member of the Raf family

of protein kinases. *Cell* **72**: 427–441.

**Kucukural A, Yukselen O, Ozata DM, Moore MJ, Garber M**. **2019**. DEBrowser: interactive

differential expression analysis and visualization tool for count data. *BMC Genomics* **20**.

**Kurtz LE, Brand MH, Lubell-Brand JD**. **2020**. Production of tetraploid and triploid hemp.

*HortScience* **55**: 1703–1707.

**Lambert MR, Giller GSJ, Barber LB, Fitzgerald KC, Skelly DK**. **2015**. Suburbanization,

estrogen contamination, and sex ratio in wild amphibian populations. *Proceedings of the*

*National Academy of Sciences of the United States of America* **112**: 11881–11886.

**Lambert MR, Tran T, Kilian A, Ezaz T, Skelly DK**. **2019**. Molecular evidence for sex

reversal in wild populations of green frogs (Rana clamitans). *PeerJ* **2019**: 1–21.

**Lapierre É, Monthony AS, Torkamaneh D**. **2023**. Genomics-based taxonomy to clarify

cannabis classification. *Genome*.

**Li Q, Guo W, Chen B, Pan F, Yang H, Zhou J, Wang G, Li X**. **2021**. Transcriptional and

Hormonal Responses in Ethephon-Induced Promotion of Femaleness in Pumpkin. *Frontiers in*

*Plant Science* **12**.

**Li Q, Zhang L, Pan F, Guo W, Chen B, Yang H, Wang G, Li X**. **2020**. Transcriptomic

analysis reveals ethylene signal transduction genes involved in pistil development of pumpkin.

*PeerJ*: 1–17.

**Liang S, Xiong W, Yin C, Xie X, Jin YJ, Zhang S, Yang B, Ye G, Chen S, Luan WJ**. **2019**.

Overexpression of OsARD1 Improves Submergence, Drought, and Salt Tolerances of Seedling

Through the Enhancement of Ethylene Synthesis in Rice. *Frontiers in Plant Science* **10**: 1–19.

**Liu C, Li J, Zhu P, Yu J, Hou J, Wang C, Long D, Yu M, Zhao A**. **2019**. Mulberry EIL3

confers salt and drought tolerances and modulates ethylene biosynthetic gene expression. *PeerJ*

**2019**: 1–17.

**Liu H, Todd E V., Lokman PM, Lamm MS, Godwin JR, Gemmell NJ**. **2017**. Sexual

plasticity: A fishy tale. *Molecular Reproduction and Development* **84**: 171–194.

**Liu Q, Wen CK**. **2012**. Arabidopsis ETR1 and ERS1 differentially repress the ethylene response

in combination with other ethylene receptor genes. *Plant Physiology* **158**: 1193–1207.

**Love MI, Huber W, Anders S**. **2014**. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biology* **15**: 1–21.

**Lubell JD, Brand MH**. **2018**. Foliar sprays of silver thiosulfate produce male flowers on female

hemp plants. *HortTechnology* **28**: 743–747.

**Manzano S, Martínez C, García JM, Megías Z, Jamilena M**. **2014**. Involvement of ethylene
in sex expression and female flower development in watermelon (Citrullus lanatus). *Plant*
*Physiology and Biochemistry* **85**: 96–104.

**Mayer BF, Ali-Benali MA, Demone J, Bertrand A, Charron JB**. **2015**. Cold acclimation
induces distinctive changes in the chromatin state and transcript levels of COR genes in
Cannabis sativa varieties with contrasting cold acclimation capacities. *Physiologia Plantarum*
**155**: 281–295.

**Mayerhofer H, Panneerselvam S, Mueller-Dieckmann J**. **2012**. Protein kinase domain of
CTR1 from arabidopsis thaliana promotes ethylene receptor cross talk. *Journal of Molecular*
*Biology* **415**: 768–779.

**McDaniel BK, Binder BM**. **2012**. Ethylene receptor 1 (ETR1) is sufficient and has the
predominant role in mediating inhibition of ethylene responses by silver in Arabidopsis thaliana.
*Journal of Biological Chemistry* **287**: 26094–26103.

**Meng J, Wang L, Wang J, Zhao X, Cheng J, Yu W, Jin D, Li Q, Gong Z**. **2018**.
METHIONINE ADENOSYLTRANSFERASE4 mediates DNA and histone methylation. *Plant*
*Physiology* **177**: 652–670.

**Mohan Ram HY, Sett R**. **1982**. Induction of Fertile Male Flowers in Genetically Female
Cannabis sativa Plants by Silver Nitrate and Silver Thiosulphate Anionic Complex. *Theoretical*
*and Applied Genetics* **62**: 369–275.

**Monthony AS, Bagheri S, Zheng Y, Jones AMP**. **2021a**. Flower power: floral reversion as a
viable alternative to nodal micropropagation in Cannabis sativa. *In Vitro Cellular &*
*Developmental Biology - Plant* **21**: 1–13.

**Monthony AS, Kyne ST, Grainger CM, Jones AMP**. **2021b**. Recalcitrance of Cannabis sativa
to de novo regeneration; a multi-genotype replication study (KR Davis, Ed.). *PLOS ONE* **16**:
e0235525.

**Monthony AS, Page SR, Hesami M, Jones AMP**. **2021c**. The Past, Present and Future of
*Cannabis sativa* Tissue Culture. *Plants* **10**: 185.

**Moon Y-H, Lee YJ, Koo SC, Hur M, Huh YC, Chang J-K, Park WT**. **2020a**. Effect of

Timing of Ethephon Treatment on the Formation of Female Flowers and Seeds from Male Plant

of Hemp (Cannabis sativa L.). *The Plant Resources Society of Korea* **33(6)**: 682–688.

**Moon J, Park YJ, Son SH, Roh J, Youn JH, Kim SY, Kim SK**. **2020b**. Brassinosteroids

signaling via BZR1 down-regulates expression of ACC oxidase 4 to control growth of

Arabidopsis thaliana seedlings. *Plant Signaling and Behavior* **15**.

**Morgan TH**. **1909**. Sex Determination and Parthenogenesis in Phylloxerans and Aphids. *Science*

**29**: 234–237.

**Oda A, Nomura-Ando K, Ahn D-H, Higashide T**. **2022**. Temporary Reduction and Control of

Female Flower Expression in Cucumber (*Cucumis sativus* L.) by Application of 1-

Methylcyclopropene. *The Horticulture Journal* **91**: 42–48.

**Okonechnikov K, Conesa A, García-Alcalde F**. **2015**. Qualimap 2: advanced multi-sample

quality control for high-throughput sequencing data.

**Olsen A, Lütken H, Hegelund JN, Müller R**. **2015**. Ethylene resistance in flowering

ornamental plants – improvements and future perspectives. *Horticulture Research* **2**: 15038.

**Page SRG, Monthony AS, Jones AMP**. **2021**. DKW basal salts improve micropropagation and

callogenesis compared with MS basal salts in multiple commercial cultivars of Cannabis sativa.

*Botany* **99**: 269–279.

**Pattyn J, Vaughan-Hirsch J, Van de Poel B**. **2021**. The regulation of ethylene biosynthesis: a

complex multilevel control circuitry. *New Phytologist* **229**: 770–782.

**Peleman J, Boerjan W, Engler G, Seurinck J, Botterman J, Alliotte T, Van Montagu M,**

**Inzé D**. **1989a**. Strong cellular preference in the expression of a housekeeping gene of

Arabidopsis thaliana encoding S-adenosylmethionine synthetase. *The Plant cell* **1**: 81–93.

**Peleman J, Saito K, Cottyn B, Engler G, Seurinck J, Van Montagu M, Inzé D**. **1989b**.

Structure and expression analyses of the S-adenosylmethionine synthetase gene family in

Arabidopsis thaliana. *Gene* **84**: 359–369.

**Perry AN, Grober MS**. **2003**. A model for social control of sex change: Interactions of

behavior, neuropeptides, glucocorticoids, and sex steroids. *Hormones and Behavior* **43**: 31–38.

**van de Poel B, Bulens I, Markoula A, Hertog MLATM, Dreesen R, Wirtz M, Vandoninck**
**S, Oppermann Y, Keulemans J, Hell R, *et al.* 2012**. Targeted systems biology profiling of
tomato fruit reveals coordination of the Yang cycle and a distinct regulation of ethylene
biosynthesis during postclimacteric ripening. *Plant Physiology* **160**: 1498–1514.

**Van de Poel B, Vandenzavel N, Smet C, Nicolay T, Bulens I, Mellidou I, Vandoninck S,**
**Hertog MLATM, Derua R, Spaepen S, *et al.* 2014**. Tissue specific analysis reveals a
differential organization and regulation of both ethylene biosynthesis and E8 during climacteric
ripening of tomato. *BMC Plant Biology* **14**.

**Pommerrenig B, Feussner K, Zierer W, Rabinovych V, Klebl F, Feussner I, Sauera N**.
**2011**. Phloem-Specific expression of Yang cycle genes and identification of novel Yang cycle
enzymes in Plantago and Arabidopsis. *Plant Cell* **23**: 1904–1919.

**Prentout D, Razumova O, Rhoné B, Badouin H, Henri H, Feng C, Käfer J, Karlov G,**
**Marais GAB**. **2020**. An efficient RNA-seq-based segregation analysis identifies the sex
chromosomes of Cannabis sativa. *Genome Research* **30**: 164–172.

**Putri GH, Anders S, Pyl PT, Pimanda JE, Zanini F**. **2022**. Analysing high-throughput
sequencing data in Python with HTSeq 2.0. *Bioinformatics* **38**: 2943–2945.

**Qiao H, Chang KN, Yazaki J, Ecker JR**. **2009**. Interplay between ethylene, ETP1/ETP2 F-box
proteins, and degradation of EIN2 triggers ethylene responses in Arabidopsis. *Genes and*
*Development* **23**: 512–521.

**Qiu K, Li Z, Yang Z, Chen J, Wu S, Zhu X, Gao S, Gao J, Ren G, Kuai B, *et al.* 2015**. EIN3
and ORE1 Accelerate Degreening during Ethylene-Mediated Leaf Senescence by Directly
Activating Chlorophyll Catabolic Genes in Arabidopsis. *PLoS Genetics* **11**: 1–20.

**Ram HYM, Jaiswal VS**. **1972**. Induction of male flowers on female plants of Cannabis sativa
by gibberellins and its inhibition by abscisic acid. *Planta* **105**: 263–266.

**Raz V, Ecker JR**. **1999**. Regulation of differential growth in the apical hook of Arabidopsis.
*Development* **126**: 3661–3668.

**Renner SS**. **2014**. The relative and absolute frequencies of angiosperm sexual systems: Dioecy,
monoecy, gynodioecy, and an updated online database. *American Journal of Botany* **101**: 1588–
1596.

**Resnick JS, Wen C-K, Shockey JA, Chang C**. **2006**. REVERSION-TO-ETHYLENE
SENSITIVITY1, a conserved gene that regulates ethylene receptor function in Arabidopsis.
*Proceedings of the National Academy of Sciences* **103**: 7917–7922.

**Roman G, Lubarsky B, Kieber JJ, Rothenberg M, Ecker JR**. **1995**. Genetic analysis of
ethylene signal transduction in Arabidopsis thaliana: Five novel mutant loci integrated into a
stress response pathway. *Genetics* **139**: 1393–1409.

**Rzewuski G, Cornell KA, Rooney L, Bürstenbinder K, Wirtz M, Hell R, Sauter M**. **2007**.
OsMTN encodes a 5′-methylthioadenosine nucleosidase that is up-regulated during
submergence-induced ethylene synthesis in rice (Oryza sativa L.). *Journal of Experimental*
*Botany* **58**: 1505–1514.

**Sakai H, Hua J, Chen QG, Chang C, Medrano LJ, Bleecker AB, Meyerowitz EM**. **1998**.
ETR2 is an ETR1-like gene involved in ethylene signaling in Arabidopsis. *Proceedings of the*
*National Academy of Sciences of the United States of America* **95**: 5812–5817.

**Sauter M, Lorbiecke R, Ouyang B, Pochapsky TC, Rzewuski G**. **2005**. The immediate-early
ethylene response gene OsARD1 encodes an acireductone dioxygenase involved in recycling of
the ethylene precursor S-adenosylmethionine. *Plant Journal* **44**: 718–729.

**Schachtsiek J, Hussain T, Azzouhri K, Kayser O, Stehle F**. **2019**. Virus-induced gene
silencing (VIGS) in Cannabis sativa L. *Plant Methods* **15**: 1–9.

**Schlesinger A, Kramarsky-Winter E, Rosenfeld H, Armoza-Zvoloni R, Loya Y**. **2010**.
Sexual plasticity and self-fertilization in the sea anemone Aiptasia diaphana. *PLoS ONE* **5**: 1–7.

**Senthilkumaran B**. **2015**. Pesticide- and sex steroid analogue-induced endocrine disruption
differentially targets hypothalamo–hypophyseal–gonadal system during gametogenesis in
teleosts – A review. *General and Comparative Endocrinology* **219**: 136–142.

**Shakeel SN, Wang X, Binder BM, Schaller GE**. **2013**. Mechanisms of signal transduction by
ethylene: Overlapping and non-overlapping signalling roles in a receptor family. *AoB PLANTS* **5**:

1–16.

**Shi J, Habben JE, Archibald RL, Drummond BJ, Chamberlin MA, Williams RW, Renee**
**Lafitte H, Weers BP**. **2015**. Overexpression of ARGOS genes modifies plant sensitivity to
ethylene, leading to improved drought tolerance in both arabidopsis and maize. *Plant Physiology*
**169**: 266–282.

**Solano R, Stepanova A, Chao Q, Ecker JR**. **1998**. Nuclear events in ethylene signaling: A
transcriptional cascade mediated by ETHYLENE-INSENSITIVE3 and ETHYLENE-
RESPONSE-FACTOR1. *Genes and Development* **12**: 3703–3714.

**Sorokin A, Yadav NS, Gaudet D, Kovalchuk I**. **2020**. Transient expression of the β-
glucuronidase gene in Cannabis sativa varieties. *Plant Signaling and Behavior* **15**: 1–8.

**Thomas M**. **2012**. *Cannabis cultivation: A Complete Grower's Guide*. San Francisco: Green
Candy Press.

**Tieman DM, Taylor MG, Ciardi JA, Klee HJ**. **2000**. The tomato ethylene receptors NR and
LeETR4 are negative regulators of ethylene response and exhibit functional compensation within
a multigene family. *Proceedings of the National Academy of Sciences of the United States of*
*America* **97**: 5663.

**Warner RR, Swearer SE**. **1991**. Social control of sex change in the bluehead wrasse,
Thalassoma bifasciatum (Pisces: Labridae). *Biological Bulletin* **181**: 199–204.

**Wickham H**. **2016**. *ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.*

**Wu Z, Zhang X, Zhang N, Gao X, Feng X, Zeng Q, Chen X, Wu J, Qi Y**. **2022**. Identification
of ACC Synthetase Genes in Saccharum and their Expression Profiles During Plant Growth and
in Response to Low-nitrogen Stress. *Tropical Plant Biology* **15**: 197–210.

**Yamagami T, Tsuchisaka A, Yamada K, Haddon WF, Harden LA, Theologis A**. **2003**.
Biochemical diversity among the 1-amino-cyclopropane-1-carboxylate synthase isozymes
encoded by the Arabidopsis gene family. *The Journal of biological chemistry* **278**: 49102–49112.

**Yamasaki S, Fujii N, Matsuura S, Mizusawa H, Takahashi H**. **2001**. The M Locus and
Ethylene-Controlled Sex Determination in Andromonoecious Cucumber Plants. *Plant and Cell*

*Physiology* **42**: 608–619.

**Yamasaki S, Fujii N, Takahashi H**. **2000**. The ethylene-regulated expression of CS-ETR2 and
CS-ERS genes in cucumber plants and their possible involvement with sex expression in flowers.
*Plant and Cell Physiology* **41**: 608–616.

**Yang C, Lu X, Ma B, Chen SY, Zhang JS**. **2015**. Ethylene signaling in rice and arabidopsis:
Conserved and diverged aspects. *Molecular Plant* **8**: 495–505.

**Yuan H, Yue P, Bu H, Han D, Wang A**. **2020**. Genome-wide analysis of ACO and ACS genes
in pear (Pyrus ussuriensis). *In Vitro Cellular and Developmental Biology - Plant* **56**: 193–199.

**Zhang J, Shi J, Ji G, Zhang H, Gong G, Guo S, Ren Y, Fan J, Tian S, Xu Y**. **2017**.
Modulation of Sex Expression in Four Forms of Watermelon by Gibberellin, Ethephone and
Silver Nitrate. *Horticultural Plant Journal* **3**: 91–100.

**Zhu BS, Zhu YX, Zhang YF, Zhong X, Pan KY, Jiang Y, Wen CK, Yang ZN, Yao X**. **2022**.
Ethylene Activates the EIN2-EIN3/EIL1 Signaling Pathway in Tapetum and Disturbs Anther
Development in Arabidopsis. *Cells* **11**.