Lack of detection of SARS-CoV-2 in British wildlife 2020-21 and first description of a stoat (*Mustela erminea) Minacovirus*

Ternenge Apaa[1,2], Amy J. Withers[1,2], Laura McKenzie[3], Ceri Staley[1], Adam Blanchard[1], Malcolm Bennett[1], Samantha Bremner-Harrison[4], Elizabeth A. Chadwick[5], Frank Hailer[5], Stephen W.R. Harrison[4], Xavier Lambin[3], Mathew Loose[6], Fiona Mathews[7], Rachael Tarlinton[1*]

1. School of Veterinary Medicine and Science, University of Nottingham, Sutton Bonington UK

2. Animal and Plant Health Agency, Addlestone Surrey, UK

3. School of Biological Sciences, University of Aberdeen, Aberdeen UK

3. School of Animal, Rural and Environmental Sciences, Nottingham Trent University, Southwell UK

4. Organisms and Environment, School of Biosciences, Cardiff University, Cardiff UK

5. School of Life Sciences, University of Nottingham, Nottingham, UK

6. School of Life sciences, University of Sussex, Sussex UK

*Corresponding author: Rachael Tarlinton Rachael.tarlinton@nottingham.ac.uk

1.1   Keywords

Coronavirus, Mustelid, Cricetid rodent, *Minacovirus*, Stoat, SARS-CoV-2

1.2   Repositories:

2. Abstract

Repeat spill over of SARS-CoV-2 into new hosts has highlighted the critical role of cross species transmission of coronaviruses and establishment of new reservoirs of virus in pandemic and epizootic spread of coronaviruses. Species particularly susceptible to SARS-CoV-2 spill-over include Mustelidae (mink, ferrets and related animals) and cricetid rodents (hamsters and related animals). These predispositions led us to screen British wildlife with sarbecovirus specific qPCR and pan coronavirus PCR assays for SARS-CoV-2 using samples collected during the human pandemic to establish if widespread spill-over was occurring. Fourteen wildlife species (n=402) were tested, including : 2 Red Foxes (*Vulpes vulpes*), 101 Badgers (*Meles meles*), 2 wild American Mink (*Neovison vison*), 41 Pine Marten (*Martes martes*), 2 Weasels (*Mustela nivalis*), 7 Stoats (*Mustela erminea*), 108 Water Voles (*Arvicola amphibius*), 39 Bank voles (*Myodes glareolous*), 10 Field Voles (*Microtus agrestis*), 15 Wood Mice (*Apodemus sylvaticus*), 1 Common Shrew (*Sorex araneaus*), 2 Pygmy Shrews (*Sorex minutus*), 2 Hedgehogs *(Erinaceus europaeus)* and 75 Eurasian Otters (*Lutra lutra*). No cases of SARS-CoV-2 were detected in any animals, however a novel minacovirus related to mink and ferret alphacoronaviruses was detected in stoats recently introduced to the Orkney Islands. This group of viruses is of interest due to pathogenicity in ferrets. The impact of this virus on the health of stoat populations remains to be established.

3. Introduction

Coronaviruses are a large and diverse group of enveloped RNA viruses spread found in diverse vertebrate hosts. Well-studied mammalian hosts, such as humans and domestic dogs, have multiple coronaviruses of several different subfamilies, although most of those of concern in mammals are members of the *Alpha-* and *Betacoronavirus* genera (Woo, Lau et al. 2012). The SARS-CoV-2 pandemic is thought to have arisen via a spill over from horseshoe bats (the natural hosts of the *Betacoronavirus* subgenus sarbecoviruses), with human infection likely via a 'liaison host' such as the racoon dog (*Nyctereutes procyonoides*) or Malayan pangolin (*Manis javanica*) (Holmes, Goldstein et al. 2021, Crits-Christoph, Gangavarapu et al. 2023, Liu, Lin et al. 2023).

The overwhelming scale of the global human pandemic led to repeated spill overs and onward transmission into other mammalian species such as domestic cats (*Felis cattus*) (Hosie, Hofmann-Lehmann et al. 2021, Goletic, Goletic et al. 2022, Jairak, Chamsai et al. 2022), farmed American mink (*Neogale vison*) (Oreshkova, Molenaar et al. 2020, Domańska-Blicharz, Orłowska et al. 2021, Eckstrand, Baldwin et al. 2021, Hammer, Quaade et al. 2021, Wasniewski, Boué et al. 2023) and

Syrian hamsters (*Mesocricetus auratus*) (Kok, Wong et al. 2022, Yen, Sit et al. 2022), and the
establishment of a new reservoir in North American white tailed deer (*Odocoileus virginianus*) (Hale,
Dennis et al. 2022, Kuchipudi, Surendran-Nair et al. 2022). A wide range of other species are either
able to be infected experimentally or have been subjects of sporadic case reports of SARS-CoV-2
infection or seroconversion including: cricetid rodents (a variety of hamster species: *Phodopus*
campbelli, Cricetulus griseus, Phodopus sungorus, Phodopus roborovskii, Mesocricetus auratus, deer
mice: *Peromyscus maniculatus nebrascensis, P. m. rufinus , P. m. sonoriensis, P. m. bairdii, P.*
polionotus subgriseus, P. californicus, bushy tailed wood rats *Neotoma cinerea* and bank voles
Myodes glareolus), felids (lion *Panthera leo*, puma *Puma concolor*, tiger *Panthera tigris*, leopard
Panthera pardus, fishing cat *Prionailurus viverrinus,* Eurasian lynx *Lynx lynx*, Canada lynx *Lynx*
canadensis , snow leopard *Panthera uncia*), other small carnivores and mustelids (dogs *Canis lupus*
familiaris, racoon dogs *Nyctereutes procyonoides*, ferrets *Mustela furo*, red foxes *Vulpes vulpes*,
racoons *Procyon lotor*, skunks *Mephitis mephitis*, marten *Martes martes*, badger *Meles meles*,
spotted hyena *Crocuta crocuta*, otters *Lutra lutra*, coati *Nasua nasua*, Asian small clawed otter *Aonyx*
cinereus, binturong *Arctictis binturong*), primates (African green monkeys *Chlorocebus aethiops*,
rhesus macaques *Macacca mulatta*, baboons *Papio hamadryas*, marmosets *Callithrix jacchus*, gorilla
Gorilla gorilla, squirrel monkeys *Saimiri spp.*), bats (Egyptian fruit bats *Rousettus aegyptiacus*),
Chinese tree shrews *Tupaia belangeri chinensis*, cattle *Bos taurus*, mule deer *Odocoileus hemionus*,
rabbits *Oryctolagus cuniculus*, big hairy armadillo *Chaetophractus villosus ,* Virginia opossum
Didelphis virginiana, Eastern grey squirrel *Sciurus carolinensis*, hippopotamus *Hippopotamus*
amphibius, giant anteater *Myrmecophaga tridactyla*, West Indian Manatee *Trichechus manatus*
manatus, (Nielsen, Alvarez et al. 2023)

SARS-CoV-2 infection in Muridae such as house mice (*Mus musculus*) and brown rats (*Rattus*
norvegicus) may depend on the virus strain: initial studies with the original (Wuhan) strains of the
virus failed to infect them (Dinnon, Leist et al. 2020, Shuai, Chan et al. 2021) and field studies failed
to demonstrate evidence of infection in wild populations (27 *M.musculus and 97 R.Norvegicus)*.
Later variants did however cause infection in laboratory studies (Gu, Chen et al. 2020, Shuai, Chan et
al. 2021, Halfmann, Iida et al. 2022, Zhang, Cui et al. 2022) and there has been at least one field
report of infection of rats (Wang, Lenoch et al. 2023).

This study focused on species of wild animals present in Great Britain that were assessed to be of
higher risk for SARS-CoV-2 spillover in 2021, when the study was begun. These included, horseshoe
bats (subject of a separate report ((Apaa, Withers et al. 2023)) mustelids, small carnivores and
cricetid rodents.

Thus far, no infection with SARS-CoV-2 has been reported in Horseshoe bats (*Rhinopholus* spp) in
Britain or mainland Europe (Orłowska, Smreczak et al. 2022, Sander, Moreira-Soto et al. 2022, Apaa,
Withers et al. 2023) although other related coronaviruses have been detected in these species. Nor,
in contrast with N. America, has the virus been detected in European deer (Holding, Otter et al.
2022, Moreira-Soto, Walzer et al. 2022, Wernike, Fischer et al. 2022). Wild animal surveillance
studies in mainland Europe have indicated sporadic detection in wild mustelids, including by qPCR in
wild American mink (*N.vison*), particularly near farmed mink outbreaks, and one otter (*L. lutra*)
(Aguiló-Gisbert, Padilla-Blanco et al. 2021, Padilla-Blanco, Aguiló-Gisbert et al. 2022, Sikkema,

Begeman et al. 2022). Serological evidence of exposure has been described in 3/14 pine martens
(*M. martes*) and 2/10 badgers (*M. meles*) (Davoust, Guérin et al. 2022), but other studies found no
evidence of infection in 48 polecats (*Mustela putorius*), 163 badgers or cricetid and murid rodents (
694 *M. glareolus*, 2 *Microtus arvalis*, 27 *M. musculus*, 97 *R. norvegicus* and 8 *Apodemus* species)
(Wernike, Drewes et al. 2022, Carmona, Burgos et al. 2023, Zamperin, Festa et al. 2023).

110 4. Methods

111 4.1 Sample collection

A total of 402 animals from 14 species (Table 1) were collected through a network of wildlife
researchers and volunteers engaged in wildlife conservation, monitoring or pest control. The
majority of samples were collected during the human covid pandemic, with the exception of a small
number of historical otter samples. The species targeted were primarily mustelids (Otters *L. lutra*,
Badgers *M. meles*, Mink, Pine Martens *M. martes*, Weasels *Mustela nivalis*, Stoats *M. erminea*) or
cricetid rodents (Water Voles *Arvicola amphibius*, Field Voles *Microtus agrestis*, Bank Voles *M.
glareolus*) with a small number of other species collected opportunistically (Hedgehogs *Erinaceus
europaeus*, Common Shrews *Sorex araneus*, Pygmy Shrews *S. minutus*, Red Foxes *V. vulpes*, Wood
Mice *Apodemus sylvaticus*). Ethical approval was granted by the University of Nottingham School of
Veterinary Medicine and Science Committee for Animal Research and Ethics (CARE), and the
University of Sussex Animal Welfare and Ethical Review Board.

Lung samples were taken from postmortem cadavers submitted to the Cardiff Otter Project (otters)
or found dead badgers submitted for tuberculosis monitoring to the University of Nottingham
(badgers tested for covid were all confirmed culture-negative for *Mycobacterium tuberculosis*
complex infections). Mink and Stoat samples (oronasal and rectal swabs from cadavers) were
provided from programmes for invasive species control and were from animals that were live
trapped and euthanased (Mink) or lethally trapped (Stoats). Rodent samples were faecal samples
from animals live trapped in Longworth or Elliot traps for population monitoring (faecal samples
taken from traps or latrine/burrow sites), and from a small number of captive Water Voles from a
licenced breeding and release programme. A small number of lung samples were taken from
animals found dead. Weasel, Hedgehog, and Shrew samples were from a small number of animals
caught as bycatch in Longworth or Elliot traps. Pine Marten samples were all faecal samples (scat)
from environmental monitoring. Samples were collected from a variety of British locations
(Supplementary Information). Collectors were solicited by social media, notices in game and hunting
organisation newsletters, letters in the *Veterinary Record* and contact networks of wildlife
organisations working with study participants and were provided with sampling packs including
gloves, collection and shipping material and instructions. Samples were collected into RNAlater®
before either storage at -20°C or direct shipping to The University of Nottingham, depending on the
capacity of the collectors.

Table 1: Species and Sample Type Screened for SARS-CoV-2

Species	Oral Swab	Rectal swab	Lung sample	Faecal sample	Total number of animals	Sample collection dates
Mustelids						
Otter	-	-	75	-	75	Jan 2020-April 2022 (51 samples)
						April 2017-Dec 2019 (25 samples)
Badger	-	-	101	-	101	April 2021-April 2022
Mink	1	1	-	-	2	Aug 2021-June2022
Pine Marten	-	-	-	41	41	March 2020- Oct 2021
Weasel	-	-	2	-	2	Sept 2021
Stoat	7	7	-	-	7	Dec 2021
Cricetid Rodents						
Water Vole	-	-	-	108	108	August-Nov 2021
Bank Vole	-	-	7	32	39	Jan-Feb 2022
Field Vole	-	-	5	5	10	Nov 2021-Feb 2022
Other species						
Wood Mouse	-	-	-	10	10	July 2021-Feb 2022
Common Shrew	-	-	-	1	1	Jan 2022
Pygmy Shrew	-	-	-	2	2	Jan 2022
Hedgehog	-	-	-	2	2	Nov 2022

Red Fox	2	2	-	-	2	April 2022
Total					402	

4.2 RNA extraction, reverse transcriptase (RT) and RNA-dependent RNA polymerase (RDRP) gene coronaviruses generic conventional PCR and envelope gene sarbecovirus-specific real-time PCR

RNA extraction from lung tissue, faecal samples, rectal and oronasal swabs, and cell culture supernatant as positive control, was carried out using the Macherey-Nagel RNA tissue extraction kit as per manufacturer's instructions. The Wuhan SARS-CoV-2 strain positive control sample used throughout this study was kindly donated by Dr Christopher Coleman (Division of Infection, Immunity and Microbes, School of Life Sciences, University of Nottingham, UK). RT was performed in two steps, using M-MLV-RT and random hexamer primers (Promega) as per manufacturer's instructions. All cDNA products were stored at -20 $^{\circ}$C for conventional PCR.

A generic pan-coronavirus PCR assay published by (Woo, Lau et al. 2005) was used to amplify a 440 bp fragment of the coronavirus RDRP gene using Q5® Hot Start High-Fidelity DNA Polymerase (New England Biolabs cat no: M0493S). Primers were: F: GGTTGGGACTATCCTAAGTGTGA and R: CCATCATCAGATAGAATCATCATA. PCR products were purified using the Nucleospin® extract II kit (Macherey-Nagel) according to manufacturer's instructions and were Sanger sequenced (Eurofins UK).

Real-time PCR was carried out using the Promega GoTaq® Probe 1-Step RT-qPCR System (Promega) with Sarbecovirus-specific envelope gene primers from bat RNA samples as published by (Corman, Landt et al. 2020). Primers and probes were: F: ACAGGTACGTTAATAGTTAATAGCGT Probe: FAM-ACACTAGCCATCCTTACTGCGCTTCG-BBQ R: ATATTGCAGCAGTACGCACACA

RNA and cDNA quality control was assessed via partial amplification of 108 bp of the beta actin gene using a published conventional PCR protocol (Fischer, Freuling et al. 2014). Primers were F: CAGCACAATGAAGATCAAGATCATC and R: CGGACTCATCGTACTCCTGCTT

4.3 High throughput sequencing and genome analyses

RNA sequencing was performed on positive samples by Novogene UK, using the Illumina NovaSeq 6000 platform. Quality filtering and trimming to remove adapters, duplicates and low quality reads was achieved using fastp v0.23.1 (Chen, Zhou et al. 2018). Kraken2 v2.1.2 was used for taxonomic classification reads against the Kraken2 viral Refseq database (Wood, Lu et al. 2019) (retrieved on 9[th] June 2022). Reads were assembled using the coronaSPAdes option in SPAdes genome assembler v3.15.4 (Meleshko, Hajirasouliha et al. 2021) using default parameters. While CheckV v1.0.1, a fully automated command-line pipeline, was used for identification and quality assessment of contigs, contigs were also queried against the NCBI custom BLASTN (v2.12.0) viral database (Altschul, Gish et al. 1990) (retrieved on 3[rd] July 2022).

177 Assembled contigs were indexed, and those that were classified and assessed as complete
178 alphacoronavirus genomes were extracted for downstream analysis using SAMtools v1.16.1 faidx
179 option (Danecek, Bonfield et al. 2021). Assembled genomes were annotated in Geneious Prime®
180 (v.2022.2.2) using NCBI coronavirus reference sequences for minacoviruses and tegacoviruses.

4.4   Phylogenetic analysis

Complete coronavirus genomes, extracted RDRP, spike, and nucleocapsid nucleotide sequences
from alphacoronavirus genomes assembled in this study, and a total of 22 reference
alphacoronavirus genomes (all *Minacovirus* full genomes available and a selection of Refseq or well
characterised full length isolates of tegacoviruses with the reference sequence of porcine epidemic
diarrhoea virus, PEDV, as an outgroup, supplementary information) were downloaded from NCBI,
aligned using Mafft v7.490 (Katoh and Standley 2013). Maximum likelihood phylogenetic trees were
reconstructed based on complete coronavirus genomes, and four different genes using IQ-TREE
v2.0.7 (Minh, Schmidt et al. 2020), using 1000 ultrafast bootstrap approximations using UFBoot2
within IQ-TREE v2.0.7 to evaluate branch support (Hoang, Chernomor et al. 2018). The ModelFinder
function within IQ-Tree was used to   select the best fitting nucleotide substitution model for
phylogenetic reconstruction (Kalyaanamoorthy, Minh et al. 2017). Phylogenetic trees were visualized
and annotated in FigTree v1.4.4 (https://github.com/rambaut/figtree/).   Using the same approach
trees were also similarly constructed for individual genes (25 Spike and Nucleocapsid genes as the S
genes of this group of viruses are known to recombine) and for all available *Minacovirus* partial gene
fragments of RDRP (RNA dependent RNA polymerase) and Spike (where there are a lot more
sequences available than for other parts of the genome).

5. Results

No animal sample tested positive on the *Sarbecovirus*-specific E gene qPCR.

Four (out of seven) stoat rectal swab samples (57%) tested positive on the pancoronavirus PCR.
Sanger sequencing of PCR products indicated that these were alphacoronaviruses of the *Minacovirus*
group. No oral swab samples tested positive from the same animals. All stoat samples in this study
were from the same population, and were sourced from the Orkney Islands stoat eradication
programme (https://www.nature.scot/professional-advice/land-and-sea-management/managing-
wildlife/orkney-native-wildlife-project) and are from the same recently introduced and heavily
bottlenecked population . All bar one sample in this study were from adult animals with a mix of
male and females. Positive samples were from 1 adult male, 1 juvenile female and 2 adult females.

5.1   Illumina sequencing Taxonomic classification, genome assembly

Taxonomic classification using Kraken2 identified reads assigned to other viral operational
taxonomic units, however, only reads classified to the Coronaviridae viral family are reported in this
study. De novo assembly of datasets from the stoats yielded one full length coronavirus contig of
28.1kb (100% quality, 99.9% completeness) and two partial contigs (7.5kb, 27% quality, 26.6%
completeness and 27.9kb, 99% quality and 99.1% completeness) from a further two samples. The
remaining sample did not yield any coronavirus contigs. The sequences had closest homology to

alphacoronaviruses of the *Minacovirus* group. The complete full genome sequence of the most
complete contig has been deposited in Genbank (accession number OP933726).

5.2 Genome annotation and organization

Genome annotation demonstrated a typical *Alphacoronavirus* genome organisation consisting of 5'
and 3' UTRs, a large ORF1ab, encoding sixteen non-structural peptides (nsp1-16) making up about
2/3 of the viral genome, and genes encoding four structural proteins: the spike (S), membrane (M),
envelope (E) and nucleocapsid (N) (Figure 1). Accessory proteins included open reading frames with
homology to ORF 3c and 7b of feline coronavirus isolates. A number of smaller potential open
reading frames within these regions were also present, possibly corresponding to other accessory
proteins with homology to the ORF3 or 7 of other *Minacovirus* and *Tegacovirus* isolates

Figure 1: Genome organisation of the stoat sequences derived from this study . Open reading frames
are shown in light green, UTR's in grey and regions with homology to non-structural proteins derived
from feline coronavirus ORF1ab are marked in dark green.

5.3 Phylogenetic analysis

Results from the maximum likelihood phylogenetic trees drawn using all available complete
Minacovirus genomes and selected reference sequences for tegacoviruses, demonstrated that the
stoat sequence is most closely clustered with ferret isolates (Figure 2). This relationship is consistent
across analysis of individual genes (Spike, Nucleocapsid full genes) and in larger phylogenetic trees of
all partial fragments of *Minacovirus* RDRP and Spike available in the NCBI database(Supplementary
information).

30210524-file00.docx

Figure 2: Maximum likelihood phylogenetic tree of full genomes of minacoviruses and tegacoviruses
constructed with 1000 bootstrap approximation, rooted on the PEDV coronavirus reference
sequence. Twenty three full genomes were included. The sequence from this study is marked in
orange. Sequences are named with species of origin, name of virus (if applicable) and a three letter
code for geographic origin (GBR=Great Britain, USA= United States of American, CHI=China,
JAP=Japan, NET=Netherlands, DEN=Denmark) and Genbank ID. Bootstrap values shown on nodes

6. Discussion

This study found no evidence of widespread circulation of SARs-CoV-2 in our opportunistically
gathered British wild carnivores and cricetid rodents in 2021-22. The study was designed to detect
an incidence rate of 5% based on our previous studies of alphacoronaviruses in British rodents
(Tsoleridis, Onianwa et al. 2016) with a target of 73 samples per species calculated with the online
Epitools sample size calculator (Epitools 2020). For some species including Otters and Badgers, for
which there were ongoing post mortem studies of found dead (largely road kill) animals (Sandoval
Barron, Swift et al. 2018, Swift, Barron et al. 2021, O'Rourke, Hynes et al. 2022, Thomas, Hailer et al.
2022), this sample size target was readily achieved. This target was also achieved for Water Voles
facilitated by a network of wildlife monitoring and captive breeding for re-introduction (Kirkland and
Farré 2021). We were however reliant on volunteer submitters for other species and did not achieve

the target levels of fresh samples for, e.g., hedgehogs, foxes, martens and other species. We were
also restricted in the type of sample collected for most species as it was primarily non-invasive
(faeces or postmortem rectal and oronasal swabs) that could be collected by network members.
There remains a possibility that our study did not detect low level circulation of SARS-CoV-2 in the
species tested if infection is spatially restricted, that we targeted the wrong tissue sample type or
that our samples were too degraded to detect virus with RNA based methods. This is inherent to the
study design of a study such as this one seeking to provide the basis for a preliminary assessment
with have scope to implement a spatially stratified study.

The detection of another coronavirus in this study, however, indicates that the methods used and
sample preservation were adequate for viral detection, at least at high prevalence, even in small
populations of samples. Our study is also in line with other European wildlife studies indicating
absence of widespread SARS-CoV-2 circulation in wild small carnivores and rodents, including wild
American Mink (Davoust, Guérin et al. 2022, Keller, Peter et al. 2022, Sikkema, Begeman et al. 2022,
Villanueva-Saz, Giner et al. 2022, Wernike, Drewes et al. 2022, Carmona, Burgos et al. 2023,
Zamperin, Festa et al. 2023). Detection of SARS-CoV-2 in one Eurasian River Otter in lung tissue and
nasal swabs and detection of a Eurasian Badger specific coronavirus in lung tissue from similar post
mortem monitoring programmes indicates that lungs are an appropriate target tissue for
coronavirus monitoring in those species (Padilla-Blanco, Aguiló-Gisbert et al. 2022, Zamperin, Festa
et al. 2023). SARS-CoV-2 is also readily detected in faecal samples in laboratory studies of rodents
and carnivore species as well as field studies of farmed mink (Griffin, Chan et al. 2021, Adney,
Lovaglio et al. 2022, Wolters, de Rooij et al. 2022, Wurtzer, Lacote et al. 2022), indicating faecal
sample screening is also an appropriate sample for coronavirus monitoring. Faecal samples or rectal
swabs are also among the easiest and most acceptable samples for volunteer submitters to collect
safely promoting their use in disease surveillance.

The stoat alphacoronavirus reported in this study is a novel finding. The stoats all came from an
eradication programme on the Orkney Islands, an archipelago off the northeastern tip of mainland
Great Britain. They are not native to the islands having been first found in 2010, and cause
considerable negative impact on breeding bird populations (Project 2023). Despite the small number
of samples collected (7) more than half of them (4) were PCR positive on rectal swabs, suggesting
that the prevalence of this virus in this population is high. Wider studies of this population and
mainland populations would be warranted to gauge the prevalence of the virus in the larger (and
source) populations.

The novel virus we identified in Orkney stoats is a member of the *Minacovirus* subgroup of the
Genus *Alphacoronavirus,* and clusters closely to mink and ferret coronaviruses. The ferret and mink
viruses have a well described pathogenicity, causing diarrhoea and sometimes a systemic disease
syndrome, similar to feline infectious peritonitis in cats (Wise, Kiupel et al. 2010, Vlasova, Halpin et
al. 2011, Terada, Minami et al. 2014, Autieri, Miller et al. 2015, Lescano, Quevedo et al. 2015, Doria-
Torra, Vidaña et al. 2016, Lamers, Smits et al. 2016, Minami, Kuroda et al. 2016, Wills, Beaufrère et
al. 2018). Wider studies of the pathogenesis of this virus in stoats and the impact of this on the
species would be warranted. There are multiple reports of the ferret viruses in this group
recombining with each other (Lamers, Smits et al. 2016, Minami, Kuroda et al. 2016, Xu 2020) and

multiple reports of farmed mink infected with SARS-CoV-2 also being co-infected with minacoviruses
at a high prevalence rate (Ip, Griffin et al. 2021, Kwok, de Rooij et al. 2021, Wasniewski, Boué et al.
2023). While recombination between SARS-CoV-2 and minacoviruses has not been observed it is a
possibility that warrants monitoring, particularly in farmed mink outbreaks.

Of interest the tegacoviruses (the most closely related alphacoronavirus clade to the minacoviruses)
are known for readily recombining and jumping host species, with canine, feline and porcine
recombinants in this group reported in multiple separate events (Wang, Ma et al. 2006, Decaro, Mari
et al. 2010, Ntafis, Mari et al. 2011, Licitra, Duhamel et al. 2014, Chen, Liu et al. 2019, Pratelli,
Tempesta et al. 2022). Canine coronaviruses have also been reported multiple times in multiple
locations in people with respiratory disease (Lednicky, Tagliamonte et al. 2022, Vlasova, Diaz et al.
2022, Vlasova, Toh et al. 2022), making this group of viruses of concern for recombination and cross
species transmission potential and warranting monitoring.

Sequences in the *Tegacovirus* group have also been reported from racoon dogs (*Nyctereutes*
procyonoides) (Wang, Ma et al. 2006, Wang, Tian et al. 2022), suggesting that there could be cross-
species transmission between racoon dogs, domestic cats, dogs, pigs and people. This is of particular
concern as this racoon dogs are known to be able to be infected with and transmit SARS-CoV-2 and
are one of the main suspects for the origin of the SARS-CoV-2 outbreak in humans (Freuling,
Breithaupt et al. 2020, Crits-Christoph, Gangavarapu et al. 2023, Rao, Parthasarathy et al. 2023).

The *Minacovirus* sequences isolated to date are all associated with mustelids of the genus *Mustela*,
subfamily Mustelinae (ferrets, mink, stoats). However there are relatively few studies of
coronaviruses in mustelids. That is beginning to be rectified with the publication of SARs-CoV-2
monitoring studies, with a possible *Gammacoronavirus* identified in Chinese ferret badgers
(*Melogale moschata*) and, most recently, isolates of a possibly new Genus, *Epsioloncoronavirus, in*
Italian badgers (Dong, Liu et al. 2007, Zamperin, Festa et al. 2023). The potential host and geographic
ranges of these viruses remain unknown.

Overall this study adds to a growing picture of a lack of SARS-CoV-2 circulation in wild European
mammals. It has however reported a novel alpha coronavirus of the *Minacovirus* sub-genus in an
island population of stoats, adding much needed information on alphacoronavirus diversity in
mustelids. The wider disease impacts and epidemiology of this virus in this species is however
unknown and requires further study.

7. Author statements

7.1   Conflicts of interest

333 The authors declare that there are no conflicts of interest

30210524-file00.docx

7.2 Funding information

This work was funded by the Biotechnology and Biosciences Research Council (BBSRC) grant number BB/W009501/1. During this period, the Otter Project was supported by funding from the Environment Agency, and by the Waterloo Foundation. The badger post mortem study was funded by DEFRA as part of their ongoing Tuberculosis monitoring

7.3 Ethical approval

Ethical approval was granted by the University of Nottingham School of Veterinary Medicine and Science Committee for Animal Research and Ethics (CARE), and the University of Sussex Animal Welfare and Ethical Review Board.

7.4 Acknowledgements

We are very grateful to the many volunteers who collected samples for this study including Holly Broadhurst, Emma Bryce, Rachel Ewans, Martina Sekirnik, Ellen Bielinksi, Cristobal Castillo, Max Kampmann, John Bruce, Andrea Sartorius, the RSPB Orkney, RSPCA Oak and Furrows, the Vincent Wildlife Trust, Northumberland Wildlife Trust,   Thanks also go to the many individuals and organisations who contribute to the otter collection network, including the Environment Agency, Natural Resources Wales, Wildlife Trusts, Local County Ecologists, Trunk Road Agencies, and local mammal groups.

8. References

Adney, D. R., J. Lovaglio, J. E. Schulz, C. K. Yinda, V. A. Avanzato, E. Haddock, J. R. Port, M. G. Holbrook, P. W. Hanley, G. Saturday, D. Scott, C. Shaia, A. M. Nelson, J. R. Spengler, C. Tansey, C. M. Cossaboom, N. M. Wendling, C. Martens, J. Easley, S. W. Yap, S. N. Seifert and V. J. Munster (2022). "Severe acute respiratory disease in American mink experimentally infected with SARS-CoV-2." JCI Insight **7**(22).

Aguiló-Gisbert, J., M. Padilla-Blanco, V. Lizana, E. Maiques, M. Muñoz-Baquero, E. Chillida-Martínez, J. Cardells and C. Rubio-Guerri (2021). "First Description of SARS-CoV-2 Infection in Two Feral American Mink (Neovison vison) Caught in the Wild." Animals (Basel) **11**(5).

Altschul, S. F., W. Gish, W. Miller, E. W. Myers and D. J. Lipman (1990). "Basic local alignment search tool." Journal of molecular biology **215**(3): 403-410.

Apaa, T., A. J. Withers, S. Staley, A. Blanchard, M. Bennett, S. Bremner-Harrison, E. A. Chadwick, F. Hailer, S. W. R. Harrison, M. Loose, F. Mathews and R. Tarlinton (2023). "Sarbecoviruses of British Horseshoe Bats; Sequence Variation and Epidemiology." BioRxiv preprint.

Autieri, C. R., C. L. Miller, K. E. Scott, A. Kilgore, V. A. Papscoe, M. M. Garner, J. L. Haupt, V. Bakthavatchalu, S. Muthupalani and J. G. Fox (2015). "Systemic Coronaviral Disease in 5 Ferrets." Comp Med **65**(6): 508-516.

Carmona, G., T. Burgos, R. Barrientos, S. Martin-Garcia, C. Muñoz, M. Sánchez-Sánchez, J. Hernández-Hernández, C. Palacín, P. Quiles, A. Moraga-Fernández, V. Bandeira, E. Virgós, C. Gortázar and I. G. Fernandez de Mera (2023). "Lack of SARS-CoV-2 RNA evidence in the lungs from wild European polecats (Mustela putorius) from Spain." Eur J Wildl Res **69**(2): 33.

30210524-file00.docx

Chen, S., D. Liu, J. Tian, H. Kang, D. Guo, Q. Jiang, J. Liu, Z. Li, X. Hu and L. Qu (2019). "Molecular
characterization of HLJ-073, a recombinant canine coronavirus strain from China with an ORF3abc
deletion." Arch Virol **164**(8): 2159-2164.
Chen, S., Y. Zhou, Y. Chen and J. Gu (2018). "fastp: an ultra-fast all-in-one FASTQ preprocessor."
Bioinformatics **34**(17): i884-i890.
Corman, V. M., O. Landt, M. Kaiser, R. Molenkamp, A. Meijer, D. K. Chu, T. Bleicker, S. Brunink, J.
Schneider, M. L. Schmidt, D. G. Mulders, B. L. Haagmans, B. van der Veer, S. van den Brink, L.
Wijsman, G. Goderski, J. L. Romette, J. Ellis, M. Zambon, M. Peiris, H. Goossens, C. Reusken, M. P.
Koopmans and C. Drosten (2020). "Detection of 2019 novel coronavirus (2019-nCoV) by real-time
RT-PCR." Euro Surveill **25**(3).
Crits-Christoph, A., K. Gangavarapu, J. E. Pekar, N. Moshiri, R. Singh, J. I. Levy , S. A. Goldstein, M. A.
Suchard, S. Popescu, D. L. Robertson, P. Lemey, J. O. Wertheim, R. F. Garry, A. L. Rasmussen, K. G.
Andersen, E. C. Holmes, A. Rambaut, M. Worobey and F. Débarre (2023). " Genetic evidence of
susceptible wildlife in SARS-CoV-2 positive samples at the Huanan Wholesale Seafood Market,
Wuhan: Analysis and interpretation of data released by the Chinese Center for Disease Control. ."
Zenodo.
Danecek, P., J. K. Bonfield, J. Liddle, J. Marshall, V. Ohan, M. O. Pollard, A. Whitwham, T. Keane, S. A.
McCarthy, R. M. Davies and H. Li (2021). "Twelve years of SAMtools and BCFtools." Gigascience
10(2).
Davoust, B., P. Guérin, N. Orain, C. Fligny, F. Flirden, F. Fenollar, O. Mediannikov and S. Edouard
(2022). "Evidence of antibodies against SARS-CoV-2 in wild mustelids from Brittany (France)."
Transbound Emerg Dis **69**(5): e3400-e3407.
Decaro, N., V. Mari, G. Elia, D. D. Addie, M. Camero, M. S. Lucente, V. Martella and C. Buonavoglia
(2010). "Recombinant canine coronaviruses in dogs, Europe." Emerg Infect Dis **16**(1): 41-47.
Dinnon, K. H., S. R. Leist, A. Schäfer, C. E. Edwards, D. R. Martinez, S. A. Montgomery, A. West, B. L.
Yount, Y. J. Hou, L. E. Adams, K. L. Gully, A. J. Brown, E. Huang, M. D. Bryant, I. C. Choong, J. S. Glenn,
L. E. Gralinski, T. P. Sheahan and R. S. Baric (2020). "A mouse-adapted SARS-CoV-2 model for the
evaluation of COVID-19 medical countermeasures." bioRxiv.
Domańska-Blicharz, K., A. Orłowska, M. Smreczak, K. Niemczuk, E. Iwan, A. Bomba, A. Lisowska, J.
Opolska, P. Trębas, P. Potyrało, M. Kawiak-Sadurska and J. Rola (2021). "Mink SARS-CoV-2 Infection
in Poland - Short Communication." J Vet Res **65**(1): 1-5.
Dong, B. Q., W. Liu, X. H. Fan, D. Vijaykrishna, X. C. Tang, F. Gao, L. F. Li, G. J. Li, J. X. Zhang, L. Q.
Yang, L. L. Poon, S. Y. Zhang, J. S. Peiris, G. J. Smith, H. Chen and Y. Guan (2007). "Detection of a
novel and highly divergent coronavirus from asian leopard cats and Chinese ferret badgers in
Southern China." J Virol **81**(13): 6920-6926.
Doria-Torra, G., B. Vidaña, A. Ramis, S. P. Amarilla and J. Martínez (2016). "Coronavirus Infection in
Ferrets: Antigen Distribution and Inflammatory Response." Vet Pathol **53**(6): 1180-1186.
Eckstrand, C. D., T. J. Baldwin, K. A. Rood, M. J. Clayton, J. K. Lott, R. M. Wolking, D. S. Bradway and T.
Baszler (2021). "An outbreak of SARS-CoV-2 with high mortality in mink (Neovison vison) on multiple
Utah farms." PLoS Pathog **17**(11): e1009952.
Epitools. (2020). "Epitools sample size calculation ", from
https://epitools.ausvet.com.au/samplesize?page=SampleSize.
Fischer, M., C. M. Freuling, T. Müller, A. Wegelt, E. A. Kooi, T. B. Rasmussen, K. Voller, D. A. Marston,
A. R. Fooks, M. Beer and B. Hoffmann (2014). "Molecular double-check strategy for the identification
and characterization of European Lyssaviruses." J Virol Methods **203**: 23-32.
Freuling, C. M., A. Breithaupt, T. Müller, J. Sehl, A. Balkema-Buschmann, M. Rissmann, A. Klein, C.
Wylezich, D. Höper, K. Wernike, A. Aebischer, D. Hoffmann, V. Friedrichs, A. Dorhoi, M. H. Groschup,
M. Beer and T. C. Mettenleiter (2020). "Susceptibility of Raccoon Dogs for Experimental SARS-CoV-2
Infection." Emerg Infect Dis **26**(12).

30210524-file00.docx

Goletic, S., T. Goletic, A. Softic, A. Zahirovic, D. Rukavina, A. Kavazovic, J. Omeragic, S. Umihanic and M. Hukic (2022). "The Evidence of SARS-CoV-2 Human-to-Pets Transmission in Household Settings in Bosnia and Herzegovina." Front Genet **13**: 839205.

Griffin, B. D., M. Chan, N. Tailor, E. J. Mendoza, A. Leung, B. M. Warner, A. T. Duggan, E. Moffat, S. He, L. Garnett, K. N. Tran, L. Banadyga, A. Albietz, K. Tierney, J. Audet, A. Bello, R. Vendramelli, A. S. Boese, L. Fernando, L. R. Lindsay, C. M. Jardine, H. Wood, G. Poliquin, J. E. Strong, M. Drebot, D. Safronetz, C. Embury-Hyatt and D. Kobasa (2021). "SARS-CoV-2 infection and transmission in the North American deer mouse." Nat Commun **12**(1): 3612.

Gu, H., Q. Chen, G. Yang, L. He, H. Fan, Y. Q. Deng, Y. Wang, Y. Teng, Z. Zhao, Y. Cui, Y. Li, X. F. Li, J. Li, N. N. Zhang, X. Yang, S. Chen, Y. Guo, G. Zhao, X. Wang, D. Y. Luo, H. Wang, X. Yang, Y. Li, G. Han, Y. He, X. Zhou, S. Geng, X. Sheng, S. Jiang, S. Sun, C. F. Qin and Y. Zhou (2020). "Adaptation of SARS-CoV-2 in BALB/c mice for testing vaccine efficacy." Science **369**(6511): 1603-1607.

Hale, V. L., P. M. Dennis, D. S. McBride, J. M. Nolting, C. Madden, D. Huey, M. Ehrlich, J. Grieser, J. Winston, D. Lombardi, S. Gibson, L. Saif, M. L. Killian, K. Lantz, R. M. Tell, M. Torchetti, S. Robbe-Austerman, M. I. Nelson, S. A. Faith and A. S. Bowman (2022). "SARS-CoV-2 infection in free-ranging white-tailed deer." Nature **602**(7897): 481-486.

Halfmann, P. J., S. Iida, K. Iwatsuki-Horimoto, T. Maemura, M. Kiso, S. M. Scheaffer, T. L. Darling, A. Joshi, S. Loeber, G. Singh, S. L. Foster, B. Ying, J. B. Case, Z. Chong, B. Whitener, J. Moliva, K. Floyd, M. Ujie, N. Nakajima, M. Ito, R. Wright, R. Uraki, P. Warang, M. Gagne, R. Li, Y. Sakai-Tagawa, Y. Liu, D. Larson, J. E. Osorio, J. P. Hernandez-Ortiz, A. R. Henry, K. Ciuoderis, K. R. Florek, M. Patel, A. Odle, L. R. Wong, A. C. Bateman, Z. Wang, V. V. Edara, Z. Chong, J. Franks, T. Jeevan, T. Fabrizio, J. DeBeauchamp, L. Kercher, P. Seiler, A. S. Gonzalez-Reiche, E. M. Sordillo, L. A. Chang, H. van Bakel, V. Simon, D. C. Douek, N. J. Sullivan, L. B. Thackray, H. Ueki, S. Yamayoshi, M. Imai, S. Perlman, R. J. Webby, R. A. Seder, M. S. Suthar, A. García-Sastre, M. Schotsaert, T. Suzuki, A. C. M. Boon, M. S. Diamond and Y. Kawaoka (2022). "SARS-CoV-2 Omicron virus causes attenuated disease in mice and hamsters." Nature **603**(7902): 687-692.

Hammer, A. S., M. L. Quaade, T. B. Rasmussen, J. Fonager, M. Rasmussen, K. Mundbjerg, L. Lohse, B. Strandbygaard, C. S. Jørgensen, A. Alfaro-Núñez, M. W. Rosenstierne, A. Boklund, T. Halasa, A. Fomsgaard, G. J. Belsham and A. Bøtner (2021). "SARS-CoV-2 Transmission between Mink (Neovison vison) and Humans, Denmark." Emerg Infect Dis **27**(2): 547-551.

Hoang, D. T., O. Chernomor, A. Von Haeseler, B. Q. Minh and L. S. Vinh (2018). "UFBoot2: improving the ultrafast bootstrap approximation." Molecular biology and evolution **35**(2): 518-522.

Holding, M., A. D. Otter, S. Dowall, K. Takumi, B. Hicks, T. Coleman, G. Hemingway, M. Royds, S. Findlay-Wilson, M. Curran-French, R. Vipond, H. Sprong and R. Hewson (2022). "Screening of wild deer populations for exposure to SARS-CoV-2 in the United Kingdom, 2020-2021." Transbound Emerg Dis **69**(5): e3244-e3249.

Holmes, E. C., S. A. Goldstein, A. L. Rasmussen, D. L. Robertson, A. Crits-Christoph, J. O. Wertheim, S. J. Anthony, W. S. Barclay, M. F. Boni, P. C. Doherty, J. Farrar, J. L. Geoghegan, X. Jiang, J. L. Leibowitz, S. J. D. Neil, T. Skern, S. R. Weiss, M. Worobey, K. G. Andersen, R. F. Garry and A. Rambaut (2021). "The origins of SARS-CoV-2: A critical review." Cell **184**(19): 4848-4856.

Hosie, M. J., R. Hofmann-Lehmann, K. Hartmann, H. Egberink, U. Truyen, D. D. Addie, S. Belák, C. Boucraut-Baralon, T. Frymus, A. Lloret, H. Lutz, F. Marsilio, M. G. Pennisi, S. Tasker, E. Thiry and K. Möstl (2021). "Anthropogenic Infection of Cats during the 2020 COVID-19 Pandemic." Viruses **13**(2).

Ip, H. S., K. M. Griffin, J. D. Messer, M. E. Winzeler, S. A. Shriner, M. L. Killian, K. T. M, T. J. DeLiberto, B. R. Amman, C. M. Cossaboom, R. R. Harvey, N. M. Wendling, H. Rettler, D. Taylor, J. S. Towner, C. Barton Behravesh and D. S. Blehert (2021). "An Opportunistic Survey Reveals an Unexpected Coronavirus Diversity Hotspot in North America." Viruses **13**(10).

Jairak, W., E. Chamsai, K. Udom, K. Charoenkul, S. Chaiyawong, N. Techakriengkrai, R. Tangwangvivat, K. Suwannakarn and A. Amonsin (2022). "SARS-CoV-2 delta variant infection in domestic dogs and cats, Thailand." Sci Rep 12(1): 8403.

Kalyaanamoorthy, S., B. Q. Minh, T. K. F. Wong, A. von Haeseler and L. S. Jermiin (2017). "ModelFinder: fast model selection for accurate phylogenetic estimates." Nat Methods 14(6): 587-589.

Katoh, K. and D. M. Standley (2013). "MAFFT multiple sequence alignment software version 7: improvements in performance and usability." Mol Biol Evol 30(4): 772-780.

Keller, M., N. Peter, C. M. Holicki, A. V. Schantz, U. Ziegler, M. Eiden, D. D. Dörge, A. Vilcinskas, M. H. Groschup and S. Klimpel (2022). "SARS-CoV-2 and West Nile Virus Prevalence Studies in Raccoons and Raccoon Dogs from Germany." Viruses 14(11).

Kirkland, C. and M. Farré (2021). "Mitochondrial Genome Evolution, Genetic Diversity, and Population Structure in British Water Voles (Arvicola amphibius)." Genes (Basel) 12(2).

Kok, K. H., S. C. Wong, W. M. Chan, L. Wen, A. W. Chu, J. D. Ip, L. K. Lee, I. T. Wong, H. W. Lo, V. C. Cheng, A. Y. Ho, B. H. Lam, H. Tse, D. Lung, K. N. H. Ng, A. K. Au, G. K. Siu and K. Y. Yuen (2022). "Co-circulation of two SARS-CoV-2 variant strains within imported pet hamsters in Hong Kong." Emerg Microbes Infect 11(1): 689-698.

Kuchipudi, S. V., M. Surendran-Nair, R. M. Ruden, M. Yon, R. H. Nissly, K. J. Vandegrift, R. K. Nelli, L. Li, B. M. Jayarao, C. D. Maranas, N. Levine, K. Willgert, A. J. K. Conlan, R. J. Olsen, J. J. Davis, J. M. Musser, P. J. Hudson and V. Kapur (2022). "Multiple spillovers from humans and onward transmission of SARS-CoV-2 in white-tailed deer." Proc Natl Acad Sci U S A 119(6).

Kwok, K. T. T., M. M. T. de Rooij, F. F. Sinartio, L. A. M. Smit, M. P. G. Koopmans and M. V. T. Phan (2021). "Genome Sequence of a Minacovirus Strain from a Farmed Mink in The Netherlands." Microbiol Resour Announc 10(8).

Lamers, M. M., S. L. Smits, G. B. Hundie, L. B. Provacia, M. Koopmans, A. Osterhaus, B. L. Haagmans and V. S. Raj (2016). "Naturally occurring recombination in ferret coronaviruses revealed by complete genome characterization." J Gen Virol 97(9): 2180-2186.

Lednicky, J. A., M. S. Tagliamonte, S. K. White, G. M. Blohm, M. M. Alam, N. M. Iovine, M. Salemi, C. Mavian and J. G. Morris (2022). "Isolation of a Novel Recombinant Canine Coronavirus From a Visitor to Haiti: Further Evidence of Transmission of Coronaviruses of Zoonotic Origin to Humans." Clin Infect Dis 75(1): e1184-e1187.

Lescano, J., M. Quevedo, O. Gonzales-Viera, L. Luna, M. K. Keel and F. Gregori (2015). "First Case of Systemic Coronavirus Infection in a Domestic Ferret (Mustela putorius furo) in Peru." Transbound Emerg Dis 62(6): 581-585.

Licitra, B. N., G. E. Duhamel and G. R. Whittaker (2014). "Canine enteric coronaviruses: emerging viral pathogens with distinct recombinant spike proteins." Viruses 6(8): 3363-3376.

Liu, M. Q., H. F. Lin, J. Li, Y. Chen, Y. Luo, W. Zhang, B. Hu, F. J. Tian, Y. J. Hu, Y. J. Liu, R. D. Jiang, Q. C. Gong, A. Li, Z. S. Guo, B. Li, X. L. Yang, Y. G. Tong and Z. L. Shi (2023). "A SARS-CoV-2-Related Virus from Malayan Pangolin Causes Lung Infection without Severe Disease in Human ACE2-Transgenic Mice." J Virol 97(2): e0171922.

Meleshko, D., I. Hajirasouliha and A. Korobeynikov (2021). "coronaSPAdes: from biosynthetic gene clusters to RNA viral assemblies." Bioinformatics.

Minami, S., Y. Kuroda, Y. Terada, K. Yonemitsu, D. Van Nguyen, R. Kuwata, H. Shimoda, A. Takano and K. Maeda (2016). "Detection of novel ferret coronaviruses and evidence of recombination among ferret coronaviruses." Virus Genes 52(6): 858-862.

Minh, B. Q., H. A. Schmidt, O. Chernomor, D. Schrempf, M. D. Woodhams, A. von Haeseler and R. Lanfear (2020). "IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era." Mol Biol Evol 37(5): 1530-1534.

Moreira-Soto, A., C. Walzer, G. Czirják, M. H. Richter, S. F. Marino, A. Posautz, P. De Yebra Rodo, G. K. McEwen, J. F. Drexler and A. D. Greenwood (2022). "Serological Evidence That SARS-CoV-2 Has Not Emerged in Deer in Germany or Austria during the COVID-19 Pandemic." <u>Microorganisms</u> **10**(4).

Nielsen, S. S., J. Alvarez, D. J. Bicout, P. Calistri, E. Canali, J. A. Drewe, B. Garin-Bastuji, J. L. Gonzales Rojas, C. Gortázar, M. Herskin, V. Michel, M. Miranda Chueca, B. Padalino, P. Pasquali, H. C. Roberts, H. Spoolder, A. Velarde, A. Viltrop, C. Winckler, C. Adlhoch, I. Aznar, F. Baldinelli, A. Boklund, A. Broglia, N. Gerhards, L. Mur, P. Nannapaneni and K. Ståhl (2023). "SARS-CoV-2 in animals: susceptibility of animal species, risk for animal and public health, monitoring, prevention and control." <u>Efsa j</u> **21**(2): e07822.

Ntafis, V., V. Mari, N. Decaro, M. Papanastassopoulou, N. Papaioannou, R. Mpatziou, C. Buonavoglia and E. Xylouri (2011). "Isolation, tissue distribution and molecular characterization of two recombinant canine coronavirus strains." <u>Vet Microbiol</u> **151**(3-4): 238-244.

O'Rourke, E., J. Hynes, S. Losada, J. L. Barber, M. G. Pereira, E. F. Kean, F. Hailer and E. A. Chadwick (2022). "Anthropogenic Drivers of Variation in Concentrations of Perfluoroalkyl Substances in Otters (Lutra lutra) from England and Wales." <u>Environ Sci Technol</u> **56**(3): 1675-1687.

Oreshkova, N., R. Molenaar, S. Vreman, F. Harders, B. B. Oude Munnink, H. Hakze, N. Gerhards, P. Tolsma, R. Bouwstra, R. Sikkema, M. Tacken, M. M. T. de Rooij, E. Weesendorp, M. Engelsma, C. Bruschke, L. A. M. Smit, M. Koopmans, W. H. M. van der Poel and A. Stegeman (2020). "SARS-CoV2 infection in farmed mink, Netherlands, April 2020." <u>BioRxiv preprint</u>.

Orłowska, A., M. Smreczak, K. Thor, M. Niedbalska, D. Pawelec, P. Trębas and J. Rola (2022). "The Genetic Characterization of the First Detected Bat Coronaviruses in Poland Revealed SARS-Related Types and Alphacoronaviruses." <u>Viruses</u> **14**(9).

Padilla-Blanco, M., J. Aguiló-Gisbert, V. Rubio, V. Lizana, E. Chillida-Martínez, J. Cardells, E. Maiques and C. Rubio-Guerri (2022). "The Finding of the Severe Acute Respiratory Syndrome Coronavirus (SARS-CoV-2) in a Wild Eurasian River Otter (Lutra lutra) Highlights the Need for Viral Surveillance in Wild Mustelids." <u>Front Vet Sci</u> **9**: 826991.

Pratelli, A., M. Tempesta, G. Elia, V. Martella, N. Decaro and C. Buonavoglia (2022). "The knotty biology of canine coronavirus: A worrying model of coronaviruses' danger." <u>Res Vet Sci</u> **144**: 190-195.

Project, O. N. W. (2023). "Orkney Native Wilife Project." from https://www.orkneynativewildlife.org.uk/.

Rao, S. S., K. Parthasarathy, V. Sounderrajan, K. Neelagandan, P. Anbazhagan and V. Chandramouli (2023). "Susceptibility of SARS Coronavirus-2 infection in domestic and wild animals: a systematic review." <u>3 Biotech</u> **13**(1): 5.

Sander, A. L., A. Moreira-Soto, S. Yordanov, I. Toplak, A. Balboni, R. S. Ameneiros, V. Corman, C. Drosten and J. F. Drexler (2022). "Genomic determinants of Furin cleavage in diverse European SARS-related bat coronaviruses." <u>Commun Biol</u> **5**(1): 491.

Sandoval Barron, E., B. Swift, J. Chantrey, R. Christley, R. Gardner, C. Jewell, I. McGrath, A. Mitchell, C. O'Cathail, A. Prosser, S. Ridout, G. Sanchez-Cabezudo, N. Smith, D. Timofte, N. Williams and M. Bennett (2018). "A study of tuberculosis in road traffic-killed badgers on the edge of the British bovine TB epidemic area." <u>Sci Rep</u> **8**(1): 17206.

Shuai, H., J. F. Chan, T. T. Yuen, C. Yoon, J. C. Hu, L. Wen, B. Hu, D. Yang, Y. Wang, Y. Hou, X. Huang, Y. Chai, C. C. Chan, V. K. Poon, L. Lu, R. Q. Zhang, W. M. Chan, J. D. Ip, A. W. Chu, Y. F. Hu, J. P. Cai, K. H. Chan, J. Zhou, S. Sridhar, B. Z. Zhang, S. Yuan, A. J. Zhang, J. D. Huang, K. K. To, K. Y. Yuen and H. Chu (2021). "Emerging SARS-CoV-2 variants expand species tropism to murines." <u>EBioMedicine</u> **73**: 103643.

Sikkema, R. S., L. Begeman, R. Janssen, W. J. Wolters, C. Geurtsvankessel, E. de Bruin, R. W. Hakze-van der Honing, P. Eblé, W. H. M. van der Poel, J. M. A. van den Brand, R. Slaterus, M. La Haye, M. P. G. Koopmans, F. Velkers and T. Kuiken (2022). "Risks of SARS-CoV-2 transmission between free-ranging animals and captive mink in the Netherlands." <u>Transbound Emerg Dis</u> **69**(6): 3339-3349.

Swift, B. M. C., E. S. Barron, R. Christley, D. Corbetta, L. Grau-Roma, C. Jewell, C. O'Cathail, A. Mitchell, J. Phoenix, A. Prosser, C. Rees, M. Sorley, R. Verin and M. Bennett (2021). "Tuberculosis in badgers where the bovine tuberculosis epidemic is expanding in cattle in England." <u>Sci Rep</u> **11**(1): 20995.

Terada, Y., S. Minami, K. Noguchi, H. Y. Mahmoud, H. Shimoda, M. Mochizuki, Y. Une and K. Maeda (2014). "Genetic characterization of coronaviruses from domestic ferrets, Japan." <u>Emerg Infect Dis</u> **20**(2): 284-287.

Thomas, N. E., F. Hailer, M. W. Bruford and E. A. Chadwick (2022). "Country-wide genetic monitoring over 21 years reveals lag in genetic recovery despite spatial connectivity in an expanding carnivore (Eurasian otter, Lutra lutra) population." <u>Evol Appl</u> **15**(12): 2125-2141.

Tsoleridis, T., O. Onianwa, E. Horncastle, E. Dayman, M. Zhu, T. Danjittrong, M. Wachtl, J. M. Behnke, S. Chapman, V. Strong, P. Dobbs, J. K. Ball, R. E. Tarlinton and C. P. McClure (2016). "Discovery of Novel Alphacoronaviruses in European Rodents and Shrews." <u>Viruses</u> **8**(3): 84.

Villanueva-Saz, S., J. Giner, A. M. Palomar, M. A. Gómez, M. Põdra, M. D. C. Aranda, M. Jiménez, P. Lizarraga, R. Hernández, A. Portillo, J. A. Oteo, I. Ruíz-Arrondo, M. D. Pérez, A. P. Tobajas, M. Verde, D. Lacasta, D. Marteles, R. Hurtado-Guerrero, L. Santiago, H. Ruíz and A. Fernández (2022). "No Evidence of SARS-CoV-2 Infection in Wild Mink (Mustela lutreola and Neogale vison) from Northern Spain during the First Two Years of Pandemic." <u>Animals (Basel)</u> **12**(15).

Vlasova, A. N., A. Diaz, D. Damtie, L. Xiu, T. H. Toh, J. S. Lee, L. J. Saif and G. C. Gray (2022). "Novel Canine Coronavirus Isolated from a Hospitalized Patient With Pneumonia in East Malaysia." <u>Clin Infect Dis</u> **74**(3): 446-454.

Vlasova, A. N., R. Halpin, S. Wang, E. Ghedin, D. J. Spiro and L. J. Saif (2011). "Molecular characterization of a new species in the genus Alphacoronavirus associated with mink epizootic catarrhal gastroenteritis." <u>J Gen Virol</u> **92**(Pt 6): 1369-1379.

Vlasova, A. N., T. H. Toh, J. S. Lee, Y. Poovorawan, P. Davis, M. S. P. Azevedo, J. A. Lednicky, L. J. Saif and G. C. Gray (2022). "Animal alphacoronaviruses found in human patients with acute respiratory illness in different countries." <u>Emerg Microbes Infect</u> **11**(1): 699-702.

Wang, W., J. H. Tian, X. Chen, R. X. Hu, X. D. Lin, Y. Y. Pei, J. X. Lv, J. J. Zheng, F. H. Dai, Z. G. Song, Y. M. Chen and Y. Z. Zhang (2022). "Coronaviruses in wild animals sampled in and around Wuhan at the beginning of COVID-19 emergence." <u>Virus Evol</u> **8**(1): veac046.

Wang, Y., J. Lenoch, D. Kohler, T. J. DeLiberto, C. Y. Tang, T. Li, Y. J. Tao, M. Guan, S. Compton, C. Zeiss, J. Hang and X. F. Wan (2023). "SARS-CoV-2 Exposure in Norway Rats (Rattus norvegicus) from New York City." <u>mBio</u>: e0362122.

Wang, Y., G. Ma, C. Lu and H. Wen (2006). "Detection of canine coronaviruses genotype I and II in raised Canidae animals in China." <u>Berl Munch Tierarztl Wochenschr</u> **119**(1-2): 35-39.

Wasniewski, M., F. Boué, C. Richomme, E. Simon-Lorière, S. Van der Werf, F. Donati, V. Enouf, Y. Blanchard, V. Beven, E. Leperchois, B. Leterrier, S. Corbet, M. Le Gouil, E. Monchatre-Leroy and E. Picard-Meyer (2023). "Investigations on SARS-CoV-2 and other coronaviruses in mink farms in France at the end of the first year of COVID-19 pandemic." <u>bioRxiv</u>.

Wernike, K., S. Drewes, C. Mehl, C. Hesse, C. Imholt, J. Jacob, R. G. Ulrich and M. Beer (2022). "No Evidence for the Presence of SARS-CoV-2 in Bank Voles and Other Rodents in Germany, 2020-2022." <u>Pathogens</u> **11**(10).

Wernike, K., L. Fischer, M. Holsteg, A. Aebischer, A. Petrov, K. Marquart, U. Schotte, J. Schön, D. Hoffmann, S. Hechinger, A. Neubauer-Juric, J. Blicke, T. C. Mettenleiter and M. Beer (2022). "Serological screening in wild ruminants in Germany, 2021/2022: No evidence of SARS-CoV-2, bluetongue virus or pestivirus spread but high seroprevalences against Schmallenberg virus." <u>Transbound Emerg Dis</u> **69**(5): e3289-e3296.

Wills, S. E., H. H. Beaufrère, B. A. Brisson, R. S. Fraser and D. A. Smith (2018). "Pancreatitis and Systemic Coronavirus Infection in a Ferret (Mustela putorius furo)." <u>Comp Med</u> **68**(3): 208-211.

Wise, A. G., M. Kiupel, M. M. Garner, A. K. Clark and R. K. Maes (2010). "Comparative sequence
analysis of the distal one-third of the genomes of a systemic and an enteric ferret coronavirus." Virus
Res **149**(1): 42-50.
Wolters, W. J., M. M. T. de Rooij, R. J. Molenaar, J. de Rond, J. C. M. Vernooij, P. A. Meijer, B. B. Oude
Munnink, R. S. Sikkema, A. N. van der Spek, M. A. H. Spierenburg, R. W. Hakze-van der Honing, W. H.
M. van der Poel, M. P. G. Koopmans, J. A. Stegeman, L. A. M. Smit, M. Augustijn-Schretlen and F. C.
Velkers (2022). "Manifestation of SARS-CoV-2 Infections in Mink Related to Host-, Virus- and Farm-
Associated Factors, The Netherlands 2020." Viruses **14**(8).
Woo, P. C., S. K. Lau, C. M. Chu, K. H. Chan, H. W. Tsoi, Y. Huang, B. H. Wong, R. W. Poon, J. J. Cai, W.
K. Luk, L. L. Poon, S. S. Wong, Y. Guan, J. S. Peiris and K. Y. Yuen (2005). "Characterization and
complete genome sequence of a novel coronavirus, coronavirus HKU1, from patients with
pneumonia." J Virol **79**(2): 884-895.
Woo, P. C., S. K. Lau, C. S. Lam, C. C. Lau, A. K. Tsang, J. H. Lau, R. Bai, J. L. Teng, C. C. Tsang, M. Wang,
B. J. Zheng, K. H. Chan and K. Y. Yuen (2012). "Discovery of seven novel Mammalian and avian
coronaviruses in the genus deltacoronavirus supports bat coronaviruses as the gene source of
alphacoronavirus and betacoronavirus and avian coronaviruses as the gene source of
gammacoronavirus and deltacoronavirus." J Virol **86**(7): 3995-4008.
Wood, D. E., J. Lu and B. Langmead (2019). "Improved metagenomic analysis with Kraken 2."
Genome Biol **20**(1): 257.
Wurtzer, S., S. Lacote, S. Murri, P. Marianneau, E. Monchatre-Leroy, M. Boni, O. Ferraris, Y. Maday,
O. Kébé, N. Dia, C. Peyrefitte, H. Sokol, C. Obepine, L. Moulin and V. Maréchal (2022). "Reduction in
SARS-CoV-2 Virus Infectivity in Human and Hamster Feces." Viruses **14**(8).
Xu, Y. (2020). "Genetic diversity and potential recombination between ferret coronaviruses from
European and American lineages." J Infect **80**(3): 350-371.
Yen, H. L., T. H. C. Sit, C. J. Brackman, S. S. Y. Chuk, H. Gu, K. W. S. Tam, P. Y. T. Law, G. M. Leung, M.
Peiris and L. L. M. Poon (2022). "Transmission of SARS-CoV-2 delta variant (AY.127) from pet
hamsters to humans, leading to onward human-to-human transmission: a case study." Lancet
399(10329): 1070-1078.
Zamperin, G., F. Festa, E. Palumbo, E. Quaranta, I. Monne, C. Terregino, P. De Benedictis and S.
Leopardi (2023). "Discovery of a coronavirus in the Eurasian badger (Meles meles) belonging to a
putative new genus." Infect Genet Evol **109**: 105406.
Zhang, C., H. Cui, E. Li, Z. Guo, T. Wang, F. Yan, L. Liu, Y. Li, D. Chen, K. Meng, N. Li, C. Qin, J. Liu, Y.
Gao and C. Zhang (2022). "The SARS-CoV-2 B.1.351 Variant Can Transmit in Rats But Not in Mice."
Front Immunol **13**: 869809.

30210524-file00.docx