1   **Title:** Personalized *Clostridioides difficile* engraftment risk prediction and probiotic
2   therapy assessment in the human gut.
3

4   **Authors:** Alex Carr[1,2], Nitin S. Baliga[1,2,3,4], Christian Diener[1,*], and Sean M.
5   Gibbons[1,2,5,6,*]
6
7   **Affiliations:**
8
[1] Institute for Systems Biology, Seattle, WA, USA
[2] Molecular Engineering Program, University of Washington, Seattle, WA, USA
[3] Departments of Biology and Microbiology, University of Washington, Seattle, WA, USA
[4] Lawrence Berkeley National Lab, Berkeley, CA, USA
[5] Departments of Bioengineering and Genome Sciences, University of Washington,
Seattle, WA, USA
[6] eScience Institute, University of Washington, Seattle, WA, USA
[*] correspondence can be addressed to cdiener@isbscience.org and
sgibbons@isbscience.org

18
19 **Abstract:**
20

Clostridioides difficile colonizes up to 30-40% of community-dwelling adults without

causing disease [1,2]. *C. difficile* infections (CDIs) are the leading cause of antibiotic-

associated diarrhea in the U.S. [3,4] and typically develop in individuals following

disruption of the gut microbiota due to antibiotic or chemotherapy treatments [2]. Further

treatment of CDI with antibiotics is not always effective and can lead to antibiotic

resistance and recurrent infections (rCDI) [5,6]. The most effective treatment for rCDI is

the reestablishment of an intact microbiota via fecal microbiota transplants (FMTs) [7].

However, the success of FMTs has been difficult to generalize because the microbial

interactions that prevent engraftment and facilitate the successful clearance of *C.*

difficile are still only partially understood [8]. Here we show how microbial community-

scale metabolic models (MCMMs) accurately predicted known instances of *C. difficile*

colonization susceptibility or resistance. MCMMs provide detailed mechanistic insights

into the ecological interactions that govern *C. difficile* engraftment, like cross-feeding or

competition involving metabolites like succinate, trehalose, and ornithine, which differ

from person to person. Indeed, three distinct *C. difficile* metabolic niches emerge from

our MCMMs, two associated with positive growth rates and one that represents non-

growth, which are consistently observed across 14,862 individuals from four

independent cohorts. Finally, we show how MCMMs can predict personalized

engraftment and *C. difficile* growth suppression for a probiotic cocktail (VE303)

designed to replace FMTs for the treatment rCDI [9,10]. Overall, this powerful modeling

approach predicts personalized *C. difficile* engraftment risk and can be leveraged to

assess probiotic treatment efficacy. MCMMs could be extended to better understand

personalized engraftment of other opportunistic bacterial pathogens, beneficial probiotic

organisms, or more complex microbial consortia.

Main Text

47
48 **Introduction**
49
The human gut microbiome plays important roles in shaping host metabolism, in the

development of chronic diseases, and in preventing opportunistic pathogen colonization

and infection [11–13]. The metabolic versatility of gut bacteria allows for the stable

coexistence of hundreds of commensal species within the gastrointestinal tract [14]. Some

species extract energy and nutrition directly from indigestible dietary substrates, like

plant fibers or recalcitrant proteins, while others subsist largely on host-derived mucosal

glycans or on the vast array of metabolic byproducts produced by primary fiber, protein,

and mucus degraders [15,16]. Saturation of these metabolic niches by commensal

microbes can prevent colonization and engraftment by external microbes that may

share a similar niche, including pathobionts [17,18].

Perturbations to the gut microbiome (e.g., antibiotic use or diarrheal events)

provide a window of opportunity for pathobiont colonization [19], which could in turn lead

to the development of disease following subsequent perturbations [20,21]. One such

pathobiont, *Clostridioides difficile*, is the most common hospital acquired gastrointestinal

infection in the U.S. [3,4]. *C. difficile* colonizes as much as 30-40% of community-dwelling

adults without causing disease, lying in wait until the opportunity for infection arises [1,2].

During active *C. difficile* infection (CDI), antibiotic treatment can be effective in

suppressing *C. difficile* growth, but antibiotics also disrupt the ecology of the commensal

microbiota and potentiate reinfection if *C. difficile* is not completely cleared by the

treatment [20,21]. Thus, an intact gut microbiota that prevents *C. difficile* colonization and

engraftment is critical to the host's defense against CDIs [13]. This understanding has led

to the widespread use of fecal microbiota transplants (FMTs) as a means of combating

cases of recurrent CDI (rCDI), where antibiotic treatment proves insufficient [7]. While the

biology of *C. difficile* has been fairly well-characterized in the context of disease, the

pre-disease mechanisms of *C. difficile* colonization and engraftment are still poorly

understood, as are the factors that govern *C. difficile* decolonization and FMT efficacy

[19].

There are currently no mechanistically-grounded, generalizable approaches to

accurately predicting the engraftment of an exogenous bacterial taxon in the context of

a given microbiota. Previous work has leveraged machine learning (ML) to predict the

engraftment of FMT donor strains in FMT recipients [22]. While effective and relatively

accurate, this kind of quasi-black-box ML approach does not provide a means of

understanding the molecular mechanisms that facilitate or prevent engraftment. Here,

we present an alternative approach to engraftment prediction that leverages microbial

community-scale metabolic models (MCMMs), which provide detailed information on the

ecological interactions within individual microbiota that prevent or facilitate engraftment,

in addition to generating accurate engraftment predictions.

Genome-scale metabolic models and classical flux balance analysis (FBA) have

been invaluable tools for exploring how environmental conditions impact the metabolic

capacity of individual bacterial taxa grown *in vitro* [23]. However, extending these methods

to complex, multi-species communities has proved to be a challenge. Recently, we

developed an approach called cooperative tradeoff flux balance analysis (ctFBA), which

leverages microbiome compositional and dietary constraints to rapidly estimate steady-

state community-scale metabolic fluxes [24,25]. Here, we leverage publicly available 16S

amplicon data from studies with both known and unknown *C. difficile* colonization

dynamics, along with our community-scale metabolic modeling framework, called

MICOM [24], to build and test MCMMs to estimate *C. difficile* engraftment potential within

a given microbiome and dietary context. We present novel insights into how *C. difficile*

can occupy three discrete metabolic niches across individuals, what metabolic

interactions within gut communities promote or prevent colonization, and we show how

we can predict potential responders and non-responders to a defined probiotic cocktail

that has recently shown efficacy in the treatment of rCDI [10]. Overall, MCMMs provide a

novel path towards predicting *C. difficile* engraftment risk. Furthermore, these models

can be leveraged to design precision dietary or probiotic interventions aimed at

decolonizing individuals who are already carrying *C. difficile* and preventing engraftment

in those who are not current carriers. Finally, we suggest that MCMMs could enable

precision engineering of the gut microbiome through personalized engraftment

predictions for other pathobionts beyond *C. difficile*, probiotic bacterial strains, or for

entire microbial consortia (e.g., FMTs from different donors), in the context of a specific

diet.

110
111
112   **Results**
113

Development of an in silico invasion assay to simulate C. difficile colonization

To simulate the colonization of *C. difficile* we developed an *in silico* invasion assay that

leverages microbiome relative abundance data, manually curated genome-scale

metabolic models of gut bacteria from the AGORA database, and the MICOM modeling

framework [24,26]. Here, we focused on leveraging available 16S amplicon sequencing

data sets, which were by far more common than shotgun metagenomic data sets and

provided a wider array of samples for validating our approach [19,27–29]. Amplicon

sequencing data is often limited to genus-level resolution in the taxonomic

classifications of amplicon sequence variants (ASVs). Therefore, we constructed genus-

level MCMMs for our invasion assays (see Methods) [30]. Specifically, strain-level

metabolic models from AGORA were combined at the genus level, to account for

potential coexistence of multiple strains and species from a given genus within an

individual and to reduce potential bias from arbitrarily selecting individual strain models.

Using this approach ~80% of reads, on average, could be mapped to an NCBI genus-

level taxonomic annotation across samples, and ~75% of the total reads could be

mapped to a genus-level metabolic model within the AGORA database (Fig. S1A). To

simulate the invasion of *C. difficile* into these model communities, a pan-genus model of

Clostridioides, representing four common *C. difficile* strains (including hypervirulent and

non-epidemic strains), was introduced at a relative abundance of 10% (see below for

justification of this percentage), while other community relative abundances were

decreased proportionally to approximate a minor perturbation in community-wide

biomass (Figure 1A). Growth simulations were then performed using a medium

representing an average European diet (i.e., a standard developed-world diet

appropriate to the cohorts studied here), with fluxes of metabolites known to be

absorbed in the small intestine decreased by 90%, as previously described [24]. Growth

rates were estimated using ctFBA, as implemented in MICOM, which uses a

regularization step and allows for a suboptimal community growth rate in order to

achieve a more realistic growth rate distribution across the community [24,25]. Import and

export fluxes were estimated using parsimonious enzyme usage FBA (pFBA) [24].

Personalized MCMMs were constructed for each sample and the potential for *C.*

difficile engraftment was quantified as the model-inferred growth rate. ctFBA has a

single free parameter that needs to be chosen, the tradeoff between community-wide

growth rates and individual, taxon-specific growth rates. Assuming that most genera

detected at appreciable abundances in a gut microbiome are actively growing *in vivo*, a

trade-off value was selected by choosing the minimal deviation from optimal community

growth for which >90% of genera obtained non-zero growth rates on average (Fig.

S1B). We found that with a trade-off value of 0.8 (i.e., 80% of maximal community

biomass production) resulted in a median fraction of genera with non-zero growth of

>90%. Furthermore, at this tradeoff value, MCMM-inferred *C. difficile* growth rates

accurately reflected trends in estimated *C. difficile* abundance across a time series with

a known *C. difficile* colonization event (Fig. 1B) [19,31]. Specifically, we found that

estimated *C. difficile* growth rates were at or below the limit of solver accuracy ($<10^{-6}$,

which effectively indicates a growth rate of zero) in samples collected prior to

colonization and comparable to growth rates of other dominant genera in samples taken

after the initial colonization event (Fig. 1B). Furthermore, we saw patchy engraftment

predictions in a second individual that was known to be colonized by *C. difficile* at a low

level (i.e., near the limit of detection) throughout a time series (Fig. 1C). We also

assessed the importance of propagule pressure [32] (i.e., the relative abundance at which

the invasive taxon is introduced into the models) and found that below 10% relative

abundance, agreement between growth rate estimates and measured abundances

were poor (Fig. S1C). Thus, propagule pressure plays an important role in predicted

engraftment success [33]. Based on these results, we decided to use a fixed tradeoff

value of 0.8 and a *C. difficile* invasion fraction of 10% for all subsequent analyses.

in silico invasion assay accurately predicts C. difficile colonization potential in rCDI
patients pre- and post- FMT

We applied our *in silico* invasion model to a dataset of rCDI patients who received FMTs

and were subsequently followed over time [29]. These data provided an additional

validation of MCMM performance and a means to explore the metabolic features

associated with community-scale colonization susceptibility or resistance across a

larger population. Given that all individuals in the rCDI cohort had experienced multiple

recurrent infections, we expected samples representative of patient pre-FMT

microbiome to be susceptible to invasion. Additionally, Weingarden *et al.* showed that

all the patient microbiomes returned to a compositional state more emblematic of

healthy controls post-FMT[29]. Thus, we expected post-FMT samples would be less

susceptible to invasion but could show variation as a function of time. Patients with

rCDI, prior to FMT treatment, had significantly higher MCMM-predicted *C. difficile*

growth rates compared to healthy individuals or to the same individuals after their FMT

treatment (Fig. 2A; Welch's t-test $p<0.01$ for comparison of pre-FMT vs. post-FMT).

Furthermore, predicted *C. difficile* growth rates were negatively associated with

Shannon diversity, albeit weakly (Fig. 2B; ordinary least squares (OLS) $R^2=0.05$,

$p=0.01$), which is in line with prior empirical observations indicating that lower diversity

communities are more susceptible to *C. difficile* colonization and the development of

rCDI [33–35].

The community-scale import flux profile prior to *in silico* invasion was predictive

of *C. difficile* growth rate following invasion (Fig. 2C). High-dimensional community-

scale import flux profiles were projected into a two-dimensional space using the Uniform

Manifold Approximation and Projection (UMAP) technique (Fig. 2C) [36]. The UMAP

projection provides a visual means of identifying patterns in the high dimensional import

flux space. The closer points are to one another in this ordination the more similar their

import flux profiles are. Thus, clusters of points in the UMAP can represent distinct

metabolic environments across samples. The ordination plot indicated that *C. difficile*

appears to grow well in more than one metabolic environment, when colonizing different

individuals. Indeed, we saw that the predicted metabolic environments occupied by *C.*

difficile could vary within an individual over time (Fig. 2C). For most patients, there was

a transition from colonization-susceptibility pre-FMT to colonization-resistance post-FMT

(Fig. 2A,C). We next examined the different apparent niches that *C. difficile* was able to

exploit when colonizing individuals in this CDI-FMT cohort, to better understand this

phenotypic plasticity.

C. difficile is predicted to occupy three distinct metabolic niches within the human gut
microbiome

To characterize *C. difficile* colonization-associated niches and identify the potential for

multiple metabolic strategies associated with its growth, we examined *C. difficile* import

fluxes with high variance (log flux variance >= 4.5) across the CDI-FMT cohort.

Biclustering of the high variance import flux data and an examination of how the

apparent clusters associated with growth rates revealed that *C. difficile* makes use of

multiple metabolic strategies (Fig. 3A). Three major clusters were observed across

patient samples. We designate these three clusters as "high growth", "moderate growth"

and "no growth" (Fig. 3A). The high growth cluster included many of the pre-FMT

samples and was characterized by consistently high import fluxes for all the metabolites

identified as most strongly coupled to *C. difficile* growth across all models. The

moderate growth cluster showed a sparser metabolite consumption profile. For

example, ornithine and fructose were rapidly consumed in the high growth cluster, but

showed almost no consumption in the moderate growth cluster (Fig. 3A). As expected,

very few metabolites were consumed by *C. difficile* above the zero-threshold of 10^{-6} in

the no growth cluster (Fig. 3A).

The metabolic strategies employed by *C. difficile* within the MCMMs showed

convergence with several observations from the literature. For example, we found that

metabolites known to promote growth of *C. difficile in vivo* (e.g., succinate, ornithine,

and trehalose) were preferentially utilized when available and were associated with high

pathobiont growth rates [37–39]. In addition, the consumption of the amino acids valine,

glycine, glutamate, glutamine, and proline were associated with higher *C. difficile* growth

rates in the MCMMs, indicating that *C. difficile* employs Strickland fermentation as one

of its growth modes, which has been observed empirically [40].

Following up on these findings we examined how cooperative and competitive

interactions within MCMMs contributed to *C. difficile* colonization. To accomplish this,

we examined the import and export fluxes of metabolites associated with *C. difficile*

colonization (e.g., amino acids, ornithine, succinate, etc.). Genera that produced

metabolites consumed by *C. difficile* likely promote its growth, while those consuming *C.*

difficile growth-associated metabolites may be in direct competition. For ornithine and

succinate, we found that cooperative and competitive interactions are context-

dependent, varying across samples. The genus *Phocaeicola*, for instance, produces

ornithine in some samples, which is in turn consumed by *C. difficile*, while in other

contexts it consumes ornithine, competing with *C. difficile* (Fig. 3B). Meanwhile,

Roseburia, and *Faecalibacterium* compete with *C. difficile* for ornithine, but these

genera also produce succinate and cysteine in some contexts, which *C. difficile*

consumes (Fig. 3B). Thus, community context is an important factor in determining the

metabolic strategies used by *C. difficile* and can lead to competitive or cooperative

interactions, which may hinder or promote colonization.

Finally, we assessed whether or not compositional variation in the microbiome

alone could explain observed differences in MCMM-predicted *C. difficile* growth rates.

We found that compositional variation was a modest predictor of estimated *C. difficile*

growth rate (out-of-sample least absolute shrinkage and selection operator (LASSO)

regression R^2=0.37 for the CDI-FMT cohort, see Methods). Meanwhile, the import flux

derived growth clusters (e.g., "high growth", "medium growth", and "no growth" groups)

could explain the vast majority of the variance in predicted *C. difficile* growth rates

(analysis of variance (ANOVA) R^2=0.94 for the CDI-FMT cohort), suggesting that

composition alone is not be sufficient for consistently accurate engraftment predictions.

Associations with C. difficile growth provide insights into the role of community context

In order to assess the consistency of the *C. difficile* growth clusters, we leveraged four

independent data sets, including the time series and CDI-FMT studies presented above

(Figs. 1-3), along with two large cross-sectional cohorts (i.e., the American Gut and

Arivale cohorts), covering a total of 14,862 individuals [27,31,41,42]. We evaluated growth

and flux predictions generated across all four data sets and found that *C. difficile* fell into

the same three clusters as we identified in the CDI-FMT data set, representing no

growth, moderate growth, and high growth (Fig. 4A).

To further contextualize the metabolic niche of *C. difficile*, we integrated model

outputs for all four data sets. Specifically, we looked at import fluxes across all genera.

Most genera formed unique clusters in the UMAP projection, suggesting that each

genus had a single metabolic niche that was consistent across datasets (Fig. 4B).

Within this community context, we found that *C. difficile* still fell into three distinct

clusters (Fig. 4B). Three genera that showed some of the strongest competitive and

cooperative interactions with *C. difficile*, *Blautia*, *Faecalibacterium*, and *Eubacterium*,

clustered near to one another in import flux space, indicating that these taxa had a

similar metabolic niche (Fig. 4B). However, these same taxa clustered apart from *C.*

difficile in their overall import flux profiles, with the exception of a few scattered samples

(Fig. 4B).

We next explored gut community diversity and predicted *C. difficile* growth rates

across the four data sets. Specifically, we looked at Shannon diversity, which integrates

species richness and evenness and is commonly used to quantify gut microbiome

alpha-diversity. Lower Shannon diversity is commonly associated with disease states,

like diarrhea, while higher diversity has generally been associated with diverse plant-

based diets and overall better health [34]. However, constipated individuals generally

have higher gut microbiome alpha-diversity as well, suggesting that there may be an

optimal range of alpha-diversity across healthy individuals [43]. Our initial analysis using

the CDI-FMT cohort suggested a negative linear relationship between predicted *C.*

difficile growth rate and Shannon (Fig. 2B). However, the integrated data sets, which

spanned a wider range of diversity, showed a U-shaped relationship between Shannon

diversity and predicted *C. difficile* growth rate (Fig. 4C). Intermediate levels of Shannon

diversity were associated with the lowest predicted growth rates, on average, with

higher average growth at the upper and lower tails of the distribution (Fig. 4C). The

relationship between Shannon diversity and predicted growth rate suggests extremes in

either direction on the diversity scale are, on average, more permissive to *C. difficile*

engraftment.

Blood metabolites and clinical labs associated with MCMM-predicted C. difficile
colonization susceptibility

We next sought to identify potential blood-based markers that were significantly

associated with MCMM-predicted *C. difficile* growth rate. Previous work has shown that

circulating blood metabolites can be leveraged to predict gut microbiome alpha-diversity

[41]. We identified several blood metabolites and clinical chemistries significantly

associated with *C. difficile* growth rate in the Arivale cohort, after adjusting for common

covariates (i.e., sex, age, and BMI) and correcting for multiple tests (FDR q<0.05).

These included two secondary bile acids, an unannotated metabolite previously

associated with the abundance of the family *Eggerthellacea*, and several red blood cell-

associated clinical chemistries (Fig. S1D)[44]. Unfortunately, while significant, these

blood-based markers, along with sex, age, and BMI, collectively accounted for only ~5%

of the variance in MCMM-predicted *C. difficile* growth rates. Thus, it appears MCMM-

based estimates of *C. difficile* engraftment cannot be readily replaced with commonly

measured clinical chemistries or blood metabolites.

MCMMs predicts engraftment heterogeneity of probiotic cocktail designed to treat rCDI

As a final proof-of-concept for our modeling framework we simulated a probiotic

intervention using a previously validated probiotic cocktail designed to treat rCDI [9,10].

The probiotic, referred to as VE303, was composed of 8 commensal Clostridia strains

and shown to be effective at treating CDI in mice [9]. This probiotic was also shown to be

safe and well-tolerated and effective in reducing rCDI incidence in humans [10].

Furthermore, the authors demonstrated that, for effective engraftment, administration of

the probiotic needed to occur following antibiotic treatment [9]. With these facts in mind,

we designed a simulated intervention that mimicked the treatment found to be most

effective by Dsouza et al. We were able to identify metabolic models for 6 of 8 strains in

VE303 in the AGORA database [26]. We leveraged the CDI-FMT dataset to test this six-

member probiotic cocktail, paired with *in silico* invasion by *C. difficile*. The probiotic

cocktail was introduced to patient samples, alongside 10% *C. difficile*, at a total relative

abundance of 50%, which was evenly distributed among the six strains. We also

simulated vancomycin treatment by reducing the abundance of *C. difficile* and all

commensal genera known to be impacted by vancomycin treatment [45] by 90%. We

found that a combined probiotic and antibiotic intervention most effectively suppressed

the growth of *C. difficile* in both the moderate and high *C. difficile* growth rate clusters

(Fig. 5A).

To better understand the mechanism of action of the probiotic cocktail, we

assessed the growth characteristics and the niche proximity of the probiotic strains in

relation to *C. difficile.* We found that suppression of *C. difficile* growth occurred when the

average growth of the probiotic strains was high ($>10^{-4}$, Fig. 5B) and when the average

niche distance between the probiotic strains and *C. difficile* was low (<25, Fig. 5C). We

also found that, relative to other genera, several of the probiotic strains occupied niches

closer to *C. difficile*, although these niche distances could vary widely for each organism

depending on their community context (Fig. S2). Finally, we compared the import fluxes

of the probiotic strains and *C. difficile* for the metabolites identified as important for *C.*

difficile growth (Fig. 3). This analysis showed that, in addition to occupying niches

similar to *C. difficile*, several of the probiotic strains directly competed for metabolites

important for *C. difficile* growth, such as succinate, ornithine, and trehalose (Fig. 5D).

Cumulatively these results suggest that metabolic competition is the mechanism by

which the probiotic cocktail suppressed *C. difficile* growth, as was suggested in the

original study [9]. Finally, we found that certain probiotic strains were more or less likely to

engraft in an individual (Fig. 5D), and that this engraftment/growth was generally

associated with *C. difficile* suppression (Fig. 5B), which indicates that MCMMs can be

leveraged to identify responders and non-responders prior to these kinds of probiotic

interventions.

Discussion

344
In this study, we provide a framework for predicting *C. difficile* engraftment risk in the

human gut microbiome using MCMMs. While we focus on *C. difficile*, due to its clinical

importance, this approach could be extended to other opportunistic bacterial pathogens,

probiotic organisms, or even entire communities, in the case of FMTs. We were able to

show how our approach predicts expected longitudinal and cross-sectional variation in

C. difficile colonization potential and we provide insights into the metabolic strategies

leveraged by *C. difficile* in different ecological contexts. Our analysis not only

recapitulates known metabolic associations with *C. difficile* growth (e.g., consumption of

trehalose, ornithine, and succinate; Fig. 3), it suggests additional associations (e.g.,

importance of reduced sulfur compounds like cysteine, Stickland fermentation reactants,

and utilization of other sugars, like fructose; Fig. 3). Additionally, we show that

competition and cooperation with community members can prevent or promote

colonization of *C. difficile*, and that many of these associations are highly context-

dependent (Fig. 3).

Consistent with the idea that simple metrics of community structure and

composition alone are not effective predictors of colonization susceptibility, we found

that community compositional variation was a modest predictor of estimated *C. difficile*

growth rate and the relationship between alpha diversity and estimated *C. difficile*

growth rate was nonlinear (Fig. 4C). Not only did low diversity communities tend to be

more invasible, as might be expected due to putative non-saturation of the metabolic

niche space, but high diversity communities were also more prone to *C. difficile*

engraftment. In high diversity communities, successful invasion may be due the

construction of new niches or changes in the interaction landscape that are in line with

the diversity-begets-diversity hypothesis [46]. Thus, an intermediate range of alpha-

diversity appears to be optimal for mitigating *C. difficile* colonization potential (Figure

4C). Overall, these complex mappings between community composition and pathobiont

engraftment risk underscore the necessity of systems-scale tools, like MCMMs, that are

capable of synthesizing this complexity.

Several genera were identified that engage in cooperative and competitive

interactions with *C. difficile* across MCMMs. *Blautia*, *Faecalibacterium*, and *Eubacterium*

were all shown to benefit *C. difficile* through production of key metabolites that it

consumes, like succinate, but were also capable of competition for metabolic resources

(Fig. 3). Meanwhile *Rumminococcus*, *Bacteroides* and *Phocaeicola* were more often

competing for the same metabolites that *C. difficile* consumed (Fig. 3). Contextualizing

these results through analysis of individual taxon import fluxes across studies we found

that *Blautia*, *Faecalibacterium*, and *Eubacterium* share similar niches with one another.

In most cases, these niches did not overlap with *C. difficile*, but in a subset of individuals

all three occupied niches states in close proximity to *C. difficile* (Fig. 4B). Thus, while we

observed competition for some key metabolites, on a global scale the majority of the

metabolic niche space used by *C. difficile* tends not to overlap with its apparent

competitors (Fig. 4B). These results highlight how flexible commensal gut bacteria are

in adapting their import fluxes to the communities in which they reside, which in turn

suggests why so many taxa are able to coexist.

In addition to developing a simulation framework to predict engraftment we

sought to identify blood-based clinical chemistries and blood metabolites that were

associated with MCMM-inferred *C. difficile* growth rate. We found three blood

metabolites that were independently associated with predicted *C. difficile* growth rates.

These included two secondary bile acids and an unannotated metabolite. One of the

secondary bile acids, isoursodeoxycholate, has previously been positively associated

with the abundance of *Bacteroides* [41], and was negatively associated with predicted *C.*

difficile growth rate. This result is in line with the apparent competition between

Bacteroides and *C. difficile* in our MCMMs. We also identified several clinical labs

negatively associated with predicted growth rate (Fig. S1D). However, together with

age, sex, and BMI, these features only accounted for ~5% of the variance in predicted

growth rates. Thus, while these features may be signatures for colonization

susceptibility in the blood, their clinical relevance is limited at this time.

Finally, we demonstrated that a probiotic intervention (i.e., 6/8 strains from

VE303), which recently showed positive efficacy results in a double-blinded, placebo-

controlled clinical trial for the treatment of rCDI [10], suppresses the growth of *C. difficile in*

silico in most people (Fig. 5). We also showed that the mechanism of action for this

particular probiotic is likely competition for metabolites essential for the growth of *C.*

difficile, as many of the probiotic strains occupy niches close to *C. difficile* and directly

compete for metabolites, such as succinate and ornithine, in samples where growth

suppression was observed (Fig. 5). Furthermore, analysis of niche distances between

C. difficile and other genera across donors suggest selecting strains from *Blautia and*

Dorea (e.g., including *B. producta and D. longicatana,* from VE303), in addition to

Anaerostipes, *Roseburia*, and *Faecalibacterium*, could be leveraged to design

individual-specific probiotic cocktails capable of suppressing *C. difficile* and rescuing

VE303 non-responders (Figs. S2 and 5B). These results illustrate how MCMMs are

powerful tools for assessing the individual-specific efficacy of clinically-relevant

probiotics, in addition to understanding personalized pathobiont colonization

susceptibility.

Future work should test this MCMM framework in the design of precision

interventions to prevent engraftment of *C. difficile* and other pathobionts, to design

precision probiotic interventions, and to improve the production of beneficial microbial

metabolites, like short-chain-fatty-acids, or reduce the production of undesired

metabolites, such as hydrogen sulfide or trimethylamine N-oxide (TMAO) [47–49]. In

summary, this MCMM framework presents a new path forward in engineering the

ecological composition and metabolic outputs of microbiota to prevent or treat disease.

424
425   **Methods:**
426
427   *External Data Collection and Processing*
428
Data used in this study came from four sources. This included both cross sectional and

time series 16S amplicon sequence data from David *et al.*, Weingarden, A. *et al.*, the

American Gut (McDonald, D. *et al.*), and a former scientific wellness program run by

Arivale, Inc.[27,29,31,42]. Publicly available 16S amplicon sequence data and associated

metadata were downloaded from the sequence read archive (SRA). Additionally, de-

identified 16S amplicon sequence data, associated metadata, and paired blood-based

clinical chemistries and metabolomics were obtained for 2,687 research consenting

individuals that were formerly participants in the Arivale wellness program. Raw 16S

amplicon sequence data were processed using QIIME2 (v2020.11.1). In brief, the

QIIME2 workflow consisted of read demultiplexing using the command *qiime tools*

import, and an associated manifest table for each study describing read metadata

followed by read quality assessment using *qiime demux summarize*. Read quality

assessment was used to determine trimming parameters for subsequent denoising

using the QIIME2 implementation of DADA2 via the command, *qiime dada2 denoise-*

single or *qiime dada2 denoise-paired*, for single and paired reads respectively. The first

10 bases were trimmed from all reads and reads were truncated to a length where

median quality score was >20 (100-150 base pair for the data leveraged). Following

denoising, data were reformatted into a table format using the command *qiime*

metadata tabulate, and representative sequence taxonomy was inferred using a custom

NCBI classifier with the command *qiime feature-classifier classify-sklearn*. The NCBI

classifier was trained using 16S 515f-806r V4 regions extracted from all available

bacterial NCBI genomes. To train the classifier 515f-806r regions were extracted from

NCBI sequences using the command *qiime feature-classifier extract-reads*, followed by

the command *qiime feature-classifier fit-classifier-naive-bayes* using the extracted V4

sequences and a table of known taxonomies. For source code and tables of processed

data refer to the Github repository listed below in Data and source code availability.

455
Model Construction and growth simulations

To construct community level metabolic models sample specific taxonomic abundance

profiles inferred from 16S amplicon sequencing were summarized at the genus level

and mapped to genus level metabolic models from the AGORA database (v1.03) using

MICOM (v0.25.1). Genera with a relative abundance less than 0.1% were omitted from

community models. An *in silico* media previously designed to represent an average

western diet was applied which defined the bounds for metabolic imports by the model

communities [24,26]. Growth rates were then inferred using cooperative tradeoff flux

balance analysis (ctFBA). In brief, this is a two-step optimization scheme, where the first

step finds the largest possible biomass production rate for the full microbial community

and the second step infers taxon-specific growth rates and fluxes, while maintaining

community growth within a fraction of the theoretical maximum (i.e., the tradeoff

parameter), thus balancing individual growth rates and the community-wide growth

rate[24]. For all models in the manuscript, we used a tradeoff parameter of 0.8. This

parameter value was chosen by identifying the largest tradeoff which allowed most

(>90%) taxa to grow (growth rate > 10^{-6}). Import and export fluxes were estimated

using parsimonious enzyme usage FBA (pFBA) and a defined medium constructed to

represent an average European diet[24]. pFBA further constrained simulation results by

requiring genera to utilize the lowest overall flux through their networks to achieve

maximal growth [23]. For source code and tables of processed data refer to the Github

repository listed below in Data and source code availability.

Probiotic Intervention

To model probiotic intervention a combination of strains previously shown to be effective

at suppressing the growth of *C. difficile* in mice were used [9]. Metabolic models for six of

the eight stains in the VE303 cocktail described by Dsouza *et al.* were identified in the

AGORA database and intervention was simulated by introducing them along with *C.*

difficile to individual samples. A total probiotic fraction of 50% was used, which was

evenly distributed among the six strains. This fraction was determined to be the most

effective at suppressing the growth of *C. difficile* growth *in silico* for the samples tested

(data not shown). Vancomycin treatment was simulated by reducing the abundance of

C. difficile and all genera known to be impacted by vancomycin by 90% [45]. Growth

simulations were performed as described above. For source code and tables of

processed data refer to the GitHub repository listed below in Data and source code

availability.

Statistical analysis

Statistical analyses were performed using functions from the python scipy (v1.7.1),

seaborn (v0.11.2), sklearn-learn (v0.24.2), umap-learn (v0.5.1) and statsmodels

(v0.13.1) packages. Linear associations were performed using the *statsmodels* ordinary

least squares function *OLS*, and visualized using the *seaborn* function *regplot.* Least

absolute shrinkage and selection operator (LASSO) was performed using the *sklearn*

Lasso function and a training-test framework. Data were split into training and test sets

(70% of samples were randomly assigned to the training set) and model performance

was assessed across a range of regularization values spanning several orders of

magnitude. LASSO training and test set R^2 were used to select the model with the best

test set R^2 that did not overfit training data (training R^2 >> test R^2). Analysis of variance

(ANOVA) was performed using the *statsmodels OLS* and *anova_lm* functions. UMAP

dimensionality reduction was performed using the *umap* function from the *umap*

package and associated methods with default parameters (i.e., *n_components=2,*

n_neighbors=15, *metric='euclidean'*, etc.). Biclustering was performed using the

seaborn function *clustermap* and the Ward clustering algorithm. Hexagonal binning and

associated histograms were generated using the *seaborn* function *jointplot.* Locally

weighted scatterplot smoothing (LOWESS) curves were generated using the *lowess*

function from *statsmodels* with default parameters. Additional statistical tests included

the t-test and Wilcox rank sum test implemented in scipy as *ttest_ind* and *wilcoxon*

respectively. For source code and tables of processed data refer to the GitHub

repository listed below in Data and source code availability.

Data and source code availability

Processed data tables and source code to reproduce the findings presented in this

manuscript can be found at https://github.com/Gibbons-Lab/cdiff_invasion. Raw 16S

amplicon sequence data from David *et al.*, Weingarden, A. *et al.*, the American Gut

(McDonald, D. *et al.*) can be downloaded using the sequence read archive (SRA)

accession numbers PRJEB6518, PRJEB19996, and PRJEB11419 respectively.

Metadata were obtained from manuscript supplementary information. Qualified

researchers can access the full Arivale deidentified dataset, including all raw data,

supporting the findings in this study for research purposes through signing a Data Use

Agreement (DUA). Inquiries to access the data can be made at data-

access@isbscience.org and will be responded to within 7 business days.

526
527
528   **Acknowledgements**
529
We thank members of the Gibbons Lab for helpful feedback and suggestions on this

work. This work was supported by a Washington Research Foundation Distinguished

Investigator Award (to SMG). Research reported in this publication was also supported

by the National Institute of Diabetes and Digestive and Kidney Diseases of the National

Institutes of Health (NIH) under award no. R01DK133468 (to SMG.).

References:

1.   Crobach, M. J. T. *et al.* Understanding Clostridium difficile Colonization. *Clin.*

Microbiol. Rev. **31**, (2018).

2. Baron, S. W. *et al.* Screening of carriers in an urban academic medical center:

Understanding implications of disease. *Infect. Control Hosp. Epidemiol.* **41**, 149–

153 (2020).

3. Dubberke, E. R. & Olsen, M. A. Burden of Clostridium difficile on the healthcare

system. *Clin. Infect. Dis.* **55 Suppl 2**, S88–92 (2012).

4. Banaei, N., Anikst, V. & Schroeder, L. F. Burden of Clostridium difficile infection in

the United States. *The New England journal of medicine* vol. 372 2368–2369

(2015).

5. Maroo, S. & Lamont, J. T. Recurrent clostridium difficile. *Gastroenterology* **130**,

1311–1316 (2006).

6. Spigaglia, P. Recent advances in the understanding of antibiotic resistance in

Clostridium difficile infection. *Ther Adv Infect Dis* **3**, 23–42 (2016).

7. Khoruts, A., Staley, C. & Sadowsky, M. J. Faecal microbiota transplantation for

Clostridioides difficile: mechanisms and pharmacology. *Nat. Rev. Gastroenterol.*

Hepatol. **18**, 67–80 (2021).

8. Sung, J. J. Y. & Wong, S. H. What is unknown in using microbiota as a

therapeutic? *Journal of Gastroenterology and Hepatology* **37**, 39–44 (2022).

9. Dsouza, M. *et al.* Colonization of the live biotherapeutic product VE303 and

modulation of the microbiota and metabolites in healthy volunteers. *Cell Host*

Microbe **30**, 583–598.e8 (2022).

10. Louie, T. *et al.* VE303, a Defined Bacterial Consortium, for Prevention of Recurrent

Clostridioides difficile Infection: A Randomized Clinical Trial. *JAMA* (2023)

doi:10.1001/jama.2023.4314.

11. Fan, Y. & Pedersen, O. Gut microbiota in human metabolic health and disease. *Nat. Rev. Microbiol.* **19**, 55–71 (2021).

12. Zeevi, D. *et al.* Personalized Nutrition by Prediction of Glycemic Responses. *Cell* **163**, 1079–1094 (2015).

13. Schubert, A. M., Sinani, H. & Schloss, P. D. Antibiotic-Induced Alterations of the Murine Gut Microbiota and Subsequent Effects on Colonization Resistance against Clostridium difficile. *MBio* **6**, e00974 (2015).

14. Donaldson, G. P., Lee, S. M. & Mazmanian, S. K. Gut biogeography of the bacterial microbiota. *Nat. Rev. Microbiol.* **14**, 20–32 (2016).

15. Bell, A. & Juge, N. Mucosal glycan degradation of the host by the gut microbiota. *Glycobiology* **31**, 691–696 (2021).

16. Baxter, N. T. *et al.* Dynamics of Human Gut Microbiota and Short-Chain Fatty Acids in Response to Dietary Interventions with Three Fermentable Fibers. *MBio* **10**, (2019).

17. Gibbons, S. M. Defining Microbiome Health through a Host Lens. *mSystems* **4**, (2019).

18. Cheng, A. G. *et al.* Design, construction, and in vivo augmentation of a complex gut microbiome. *Cell* **185**, 3617–3636.e19 (2022).

19. VanInsberghe, D. *et al.* Diarrhoeal events can trigger long-term Clostridium difficile colonization with recurrent blooms. *Nat Microbiol* **5**, 642–650 (2020).

20. Shields, K., Araujo-Castillo, R. V., Theethira, T. G., Alonso, C. D. & Kelly, C. P. Recurrent Clostridium difficile infection: From colonization to cure. *Anaerobe* **34**,

584  59–73 (2015).

585 21. Song, J. H. & Kim, Y. S. Recurrent Clostridium difficile Infection: Risk Factors,

586  Treatment, and Prevention. *Gut Liver* **13**, 16–24 (2019).

587 22. Ianiro, G. *et al.* Variability of strain engraftment and predictability of microbiome

588  composition after fecal microbiota transplantation across different diseases. *Nat.*

589  *Med.* **28**, 1913–1923 (2022).

590 23. Lewis, N. E. *et al.* Omic data from evolved E. coli are consistent with computed

591  optimal growth from genome-scale models. *Mol. Syst. Biol.* **6**, 390 (2010).

592 24. Diener, C., Gibbons, S. M. & Resendis-Antonio, O. MICOM: Metagenome-Scale

593  Modeling To Infer Metabolic Interactions in the Gut Microbiota. *mSystems* **5**,

594  (2020).

595 25. Diener, C. & Gibbons, S. M. More is Different: Metabolic Modeling of Diverse

596  Microbial Communities. *mSystems* e0127022 (2023).

597 26. Magnúsdóttir, S. *et al.* Generation of genome-scale metabolic reconstructions for

598  773 members of the human gut microbiota. *Nat. Biotechnol.* **35**, 81–89 (2017).

599 27. McDonald, D. *et al.* American Gut: an Open Platform for Citizen Science

600  Microbiome Research. *mSystems* **3**, (2018).

601 28. Wilmanski, T. *et al.* Gut microbiome pattern reflects healthy ageing and predicts

602  survival in humans. *Nat Metab* **3**, 274–286 (2021).

603 29. Weingarden, A. *et al.* Dynamic changes in short- and long-term bacterial

604  composition following fecal microbiota transplantation for recurrent Clostridium

605  difficile infection. *Microbiome* vol. 3 Preprint at https://doi.org/10.1186/s40168-015-

606  0070-0 (2015).

30. Bolyen, E. *et al.* Reproducible, interactive, scalable and extensible microbiome data
science using QIIME 2. *Nat. Biotechnol.* **37**, 852–857 (2019).

31. David, L. A. *et al.* Host lifestyle affects human microbiota on daily timescales.
Genome Biol. **15**, R89 (2014).

32. Lockwood, J. L., Cassey, P. & Blackburn, T. The role of propagule pressure in
explaining species invasions. *Trends Ecol. Evol.* **20**, 223–228 (2005).

33. Hromada, S. *et al.* Negative interactions determine Clostridioides difficile growth in
synthetic human gut communities. *Mol. Syst. Biol.* **17**, e10355 (2021).

34. Chang, J. Y. *et al.* Decreased Diversity of the Fecal Microbiome in Recurrent
Clostridium difficile—Associated Diarrhea. *J. Infect. Dis.* **197**, 435–438 (2008).

35. Pakpour, S. *et al.* Identifying predictive features of Clostridium difficile infection
recurrence before, during, and after primary antibiotic treatment. *Microbiome* **5**, 148
(2017).

36. McInnes, L., Healy, J. & Melville, J. UMAP: Uniform Manifold Approximation and
Projection for Dimension Reduction. *arXiv [stat.ML]* (2018).

37. Ferreyra, J. A. *et al.* Gut microbiota-produced succinate promotes C. difficile
infection after antibiotic treatment or motility disturbance. *Cell Host Microbe* **16**,
770–777 (2014).

38. Pruss, K. M. *et al.* Oxidative ornithine metabolism supports non-inflammatory C.
difficile colonization. *Nat Metab* **4**, 19–28 (2022).

39. Buckley, A. M. *et al.* Trehalose-Induced Remodelling of the Human Microbiota
Affects Clostridioides difficile Infection Outcome in an In Vitro Colonic Model: A
Pilot Study. *Front. Cell. Infect. Microbiol.* **11**, 670935 (2021).

40. Bouillaut, L., Self, W. T. & Sonenshein, A. L. Proline-dependent regulation of
Clostridium difficile Stickland metabolism. *J. Bacteriol.* **195**, 844–854 (2013).

41. Wilmanski, T. *et al.* Blood metabolome predicts gut microbiome α-diversity in
humans. *Nat. Biotechnol.* **37**, 1217–1228 (2019).

42. Manor, O. *et al.* Health and disease markers correlate with gut microbiome
composition across thousands of people. *Nat. Commun.* **11**, 5206 (2020).

43. Tian, H., Chen, Q., Yang, B., Qin, H. & Li, N. Analysis of Gut Microbiome and
Metabolite Characteristics in Patients with Slow Transit Constipation. *Dig. Dis. Sci.*
66, 3026–3035 (2021).

44. Bar, N. *et al.* A reference map of potential determinants for the human serum
metabolome. *Nature* **588**, 135–140 (2020).

45. Nazzal, L. *et al.* Effect of Vancomycin on the Gut Microbiome and Plasma
Concentrations of Gut-Derived Uremic Solutes. *Kidney Int Rep* **6**, 2122–2133
(2021).

46. Madi, N., Vos, M., Murall, C. L., Legendre, P. & Shapiro, B. J. Does diversity beget
diversity in microbiomes? *Elife* **9**, (2020).

47. Liu, Y. & Dai, M. Trimethylamine N-Oxide Generated by the Gut Microbiota Is
Associated with Vascular Inflammation: New Insights into Atherosclerosis.
Mediators Inflamm. **2020**, 4634172 (2020).

48. Dalile, B., Van Oudenhove, L., Vervliet, B. & Verbeke, K. The role of short-chain
fatty acids in microbiota-gut-brain communication. *Nat. Rev. Gastroenterol.*
Hepatol. **16**, 461–478 (2019).

49. Murros, K. E. Hydrogen Sulfide Produced by Gut Bacteria May Induce Parkinson's

Disease. *Cells* **11**, (2022).

Figure Captions

Figure 1. *in silco* invasion assay accurately predicts *C. difficile* colonization in
patient time series. (**A**) Schematic illustrating the *in silico* invasion assay workflow
leveraged in this study. Personalized microbial community-scale metabolic models
(MCMMs) are supplemented with 10% of a pan-genus *Clostridioides* model to simulate
an invasion event and ctFBA was used to predict *C. difficile* engraftment and metabolic
fluxes. (**B**) Donor A time series taken from David *et al.* displaying daily fluctuations in
microbiome composition over a period of several months. Composition is displayed,
colored by phylum-level annotations (different shading indicates taxonomic families). At
day 150, Donor A experienced a diarrheal event and was subsequently colonized by *C.*
difficile. Estimates of *C. difficile* relative abundance from 16S sequencing and MICOM-
predicted *C. difficile* growth rates are displayed. (**C**) Time series from Donor B from the
same study, who was apparently colonized by *C. difficile* (at very low relative
abundances, near the limit of detection) throughout the sampling period.

Figure 2. *C. difficile* growth rate predictions capture importance of community
context for patient recovery from CDI. (**A**) Violin plots displaying predicted *C. difficile*
$\log_{10}$ growth rate distributions across patient disease status (gray shading indicates the
numerical accuracy of the simulation (values below 10^{-6} cannot be distinguished from
zero and are considered negligible). Bars indicate comparisons for which differences
were significant using the Welch test. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$. (**B**)
Relationship between predicted *C. difficile* $\log_{10}$ growth rate and Shannon diversity.
Ordinary least squares fit and 95% confidence interval are displayed, as well as

regression R^2 and *p*-value. (**C**) Two-dimensional representation of community import
fluxes prior to *in silico* invasion using UMAP colored by $\log_{10}$ growth rates of *C. difficile*
following *in silico* invasion. Patient trajectories are displayed, each with a red circle
representing the patient's starting point (prior to FMT), and a red star representing the
patient's end point (post recovery).
697
Figure 3. *C. difficile* occupies multiple metabolic niches across communities. (**A**)
Biclustered *C. difficile* $\log_{10}$ import fluxes, where each row is the import flux of a
particular metabolite and each column is a patient sample. Imports for which the log
variance across samples was >= 4.5 are displayed as the blue-to-yellow heatmap. (**B**)
Biclustered community import and export fluxes of specific metabolites associated with
C. difficile colonization, where each row is a genus and each column is a patient
sample. Genera for which the mean import or export flux across samples was $> 10^{-6}$ are
displayed. Fluxes across samples are displayed using blue-to-yellow heatmap coloring.
C. difficile $\log_{10}$ growth rate quantiles are displayed in white-to-red heatmap coloring for
each patient sample in the top row of each plot. Additionally, three coarse grain growth
clusters are noted. These growth clusters represent "high growth", "moderate growth",
and "no growth" phenotypes.
710
Figure 4. Growth niches in large healthy cohorts challenged with *C. difficile*. (**A**)
Two-dimensional representation of $\log_{10}$ *C. difficile* import fluxes using UMAP across
four independent data sets. Colors denote *C. difficile* growth rate ranging from low
(blue) to high (yellow). The position of the no growth cluster is indicated. (**B**) Two-
dimensional representation of $\log_{10}$ genus import fluxes using UMAP across all
datasets. Top panel displays $\log_{10}$ *C. difficile* growth rate within the context of all other
genera. Bottom panel colors *C. difficile* and three genera of interest: *Blautia*,
Faecalibacterium, and *Eubacterium*. The position of the no growth cluster is indicated.
(**C**) Two-dimensional hexagonal binning of $\log_{10}$ *C. difficile* growth rate and community
alpha diversity (Shannon index). Red trend line indicates a LOWESS fit to the $\log_{10}$ *C.*
difficile growth rate and community Shannon diversity data.
722
Figure 5. Simulated probiotic intervention effectively suppresses *C. difficile*
growth *in silico*. (**A**) Box plots displaying $\log_{10}$ *C. difficile* growth rate across growth
clusters and simulated interventions. Growth clusters are those identified by bi-
clustering of *C. difficile* import fluxes using the Weingarden data (Figure 3). Conditions
include +None (no intervention control), +Probiotic (introduction of 6 strain probiotic
previously identified as an effective treatment for CDI at a total relative abundance of
50% equally distributed across the strains), +Vancomycin (90% reduction of *C. difficile*
relative abundance as well as all genera known to be impacted by Vancomycin), and
+Vancomycin, +Probiotic (introduction of 6 strain probiotic in combination with simulated
vancomycin treatment). Bars indicate comparisons for which differences were

significant using the Wilcoxon signed-rank test. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$. (**B**) Relationship between *C. difficile* growth ratio and mean $\log_{10}$ probiotic growth rate. *C. difficile* growth ratio is the growth rate of samples in the +Vancomycin, +Probiotic intervention relative to +None. Values below 1 indicate growth suppression by the probiotic and values above 1 indicate growth stimulation. The dashed line marks the value at which no effect is observed (1). Orange trend line indicates a LOWESS fit to the *C. difficile* growth ratio and mean $\log_{10}$ probiotic growth rate. (**C**) Relationship between $\log_{10}$ *C. difficile* growth rate and mean probiotic niche distance. Niche distance was calculated using the euclidean distance of $\log_{10}$ import flux vectors of each probiotic strain relative to *C. difficile* on a per sample basis. Orange trend line indicates a LOWESS fit to the $\log_{10}$ *C. difficile* growth rate and mean probiotic niche distance. (**D**) Biclustered $\log_{10}$ import fluxes for *C. difficile* and probiotic strains for samples previously identified as "high growth", where each row is the import flux of a particular metabolite and each column is a patient sample. Imports displayed are those previously identified as important for *C. difficile*. Color bars indicate sample *C. difficile* growth ratio and strain specific $\log_{10}$ growth rate. Ordering of samples and metabolites is the same across heatmaps and based on biclustering of *C. difficile* data.

Supplemental Figure Captions

Figure S1. Development of *in silico C. difficile* invasion assay. (**A**) Histograms displaying the fraction of reads mapped at the genus level for the David *et al.* 16S amplicon data using an NCBI reference and genus level metabolic models using the AGORA database referred to as "Source Data" and "With models" respectively. (**B**) Median growth fraction across samples (e.g., fraction of taxa with estimated growth rate $> 10^{-6}$) as a function of model tradeoff value. Dashed line indicated the tradeoff value chosen for subsequent analyses. (**C**) Relationship between *C. difficile* invasion abundance and growth rate for one of the two David *et al.* time series. (**D**) Association coefficients for estimated *C. difficile* log growth rate, blood metabolite concentrations, and clinical labs for the Arivale cohort.

Figure S2. Probiotic strains and associated genera have niche distances close to *C. difficile* relative to unrelated genera. Niche distances of strains and genera, represented as the Euclidean distance between flux vectors, relative to *C. difficile* across CDI-FMT cohort samples for which the *C. difficile* growth rate $>10^{-6}$. Genera and strains are ordered by the median niche distance. Probiotic strains and associated genera are colored, consistent with the legend in Fig. 5D. *C. bolteae* and *C. innocuum* are both members of the genus *Clostridium*.

A) log *C. diff* growth for post-FMT, healthy, and Pre-FMT. ** markers shown.

B) log *C. diff* growth vs Shannon. $R^2=0.05$, $P=0.01$

C) UMAP1 vs UMAP2 plots for CD1, CD2, CD3, CD4. Log10 *C. diff* growth color scale from -1 to -6.