Title: Personalized *Clostridioides difficile* engraftment risk prediction and probiotic therapy assessment in the human gut.

Authors: Alex Carr[1,2], Nitin S. Baliga[1,2,3,4], Christian Diener[1,5,*], and Sean M. Gibbons[1,2,6,7,*]

Affiliations:

[1] Institute for Systems Biology, Seattle, WA, USA
[2] Molecular Engineering Program, University of Washington, Seattle, WA, USA
[3] Departments of Biology and Microbiology, University of Washington, Seattle, WA, USA
[4] Lawrence Berkeley National Lab, Berkeley, CA, USA
[5] Diagnostic and Research Institute of Hygiene, Microbiology and Environmental Medicine, Medical University of Graz, Graz, Austria
[6] Departments of Bioengineering and Genome Sciences, University of Washington, Seattle, WA, USA
[7] eScience Institute, University of Washington, Seattle, WA, USA
[*] correspondence can be addressed to cdiener@isbscience.org and sgibbons@isbscience.org

Abstract:

Clostridioides difficile colonizes up to 30-40% of community-dwelling adults without causing disease. *C. difficile* infections (CDIs) are the leading cause of antibiotic-associated diarrhea in the U.S. and typically develop in individuals following disruption of the gut microbiota due to antibiotic or chemotherapy treatments. Further treatment of CDI with antibiotics is not always effective and can lead to antibiotic resistance and recurrent infections (rCDI). The most effective treatment for rCDI is the reestablishment of an intact microbiota via fecal microbiota transplants (FMTs). However, the success of FMTs has been difficult to generalize because the microbial interactions that prevent engraftment and facilitate the successful clearance of *C. difficile* are still only partially understood. Here we show how microbial community-scale metabolic models (MCMMs) accurately predicted known instances of *C. difficile* colonization susceptibility or resistance *in vitro* and *in vivo*. MCMMs provide detailed mechanistic insights into the ecological interactions that govern *C. difficile* engraftment, like cross-feeding or competition involving metabolites like succinate, trehalose, and ornithine, which differ from person to person. Indeed, three distinct *C. difficile* metabolic niches emerge from our MCMMs, two associated with positive growth rates and one that represents non-growth, which are consistently observed across 15,204 individuals from five independent cohorts. Finally, we show how MCMMs can predict personalized engraftment and *C. difficile* growth suppression for a probiotic cocktail (VE303) designed to replace FMTs for the treatment rCDI. Overall, this powerful modeling approach predicts personalized *C. difficile* engraftment risk and can be leveraged to assess probiotic treatment efficacy. MCMMs could be extended to understand the mechanistic underpinnings of personalized engraftment of other opportunistic bacterial pathogens, beneficial probiotic organisms, or more complex microbial consortia.

Introduction

The human gut microbiome plays important roles in shaping host metabolism, in the development of chronic diseases, and in preventing opportunistic pathogen colonization and infection [1-3]. The metabolic versatility of gut bacteria allows for the stable coexistence of hundreds of commensal species within the gastrointestinal tract [4]. Some species extract energy and nutrition directly from indigestible dietary substrates, like plant fibers or recalcitrant proteins, while others subsist largely on host-derived mucosal glycans or on the vast array of metabolic byproducts produced by primary fiber, protein, and mucus degraders [5,6]. Saturation of these metabolic niches by commensal microbes can prevent colonization and engraftment by external microbes that may share a similar niche, including pathobionts [7-9].

Perturbations to the gut microbiome (e.g., antibiotic use or diarrhoeal events) provide a window of opportunity for pathobiont colonization [10], which could in turn lead to the development of disease following subsequent perturbations [11,12]. One such pathobiont, *Clostridioides difficile*, is the most common hospital acquired gastrointestinal infection in the U.S. [13,14]. *C. difficile* colonizes as much as 30-40% of community-dwelling adults without causing disease, lying in wait until the opportunity for infection arises [15,16]. During active *C. difficile* infection (CDI), antibiotic treatment can be effective in suppressing *C. difficile* growth, but antibiotics also disrupt the ecology of the commensal microbiota and potentiate reinfection if *C. difficile* is not completely cleared by the treatment [11,12]. Thus, an intact gut microbiota that prevents *C. difficile* colonization and engraftment is critical to the host's defense against CDIs [3]. This

understanding has led to the widespread use of fecal microbiota transplants (FMTs) as

a means of combating cases of recurrent CDI (rCDI), where antibiotic treatment proves

insufficient [17]. While the biology of *C. difficile* has been fairly well-characterized in the

context of disease, the pre-disease mechanisms of *C. difficile* colonization and

engraftment are still poorly understood, as are the factors that govern *C. difficile*

decolonization and FMT efficacy [10].

There are currently no mechanistically-grounded, generalizable approaches to

accurately predicting the engraftment of an exogenous bacterial taxon in the context of

a given microbiota. Previous work has leveraged machine learning (ML) to predict the

engraftment of FMT donor strains in FMT recipients [18]. While effective and relatively

accurate, this kind of quasi-black-box ML approach does not provide a means of

understanding the molecular mechanisms that facilitate or prevent engraftment. Here,

we present an alternative approach to engraftment prediction that leverages microbial

community-scale metabolic models (MCMMs), which provide detailed, mechanistic

information on the ecological interactions within individual microbiota that prevent or

facilitate engraftment, in addition to generating accurate engraftment predictions.

Genome-scale metabolic models and classical flux balance analysis (FBA) have

been invaluable tools for exploring how environmental conditions impact the metabolic

capacities of individual bacterial taxa grown *in vitro* [19]. However, extending these

methods to complex, multi-species communities has proved to be a challenge.

Recently, we developed an approach called cooperative tradeoff flux balance analysis

(ctFBA), which leverages microbiome compositional and dietary constraints to rapidly

estimate steady-state community-scale metabolic fluxes [20,21]. Here, we use publicly

available 16S amplicon and shotgun metagenomic data from *in vitro* and *in vivo* studies

with both known and unknown *C. difficile* colonization dynamics, along with our

community-scale metabolic modeling framework, called MICOM [20], to build and test

MCMMs to estimate *C. difficile* engraftment potential within a given microbiome and

dietary/environmental context. We present novel insights into how *C. difficile* can

occupy three discrete metabolic niches across commensal communities, what metabolic

interactions within gut bacterial communities promote or prevent colonization, and we

show how we can predict potential responders and non-responders to a defined

probiotic cocktail that has recently shown efficacy in the treatment of rCDI [22]. Overall,

MCMMs provide a novel path towards predicting *C. difficile* engraftment risk.

Furthermore, these models can be leveraged to design precision dietary or probiotic

interventions aimed at decolonizing individuals who are already carrying *C. difficile* and

preventing engraftment in those who are not. Finally, we suggest that MCMMs could

enable precision engineering of the gut microbiome through personalized engraftment

predictions for other pathobionts beyond *C. difficile*, probiotic bacterial strains, or for

entire microbial consortia (e.g., FMTs from different donors), in the context of a specific

diet.

108
109
Results

Development of an in silico invasion assay to simulate C. difficile colonization

To simulate the colonization of *C. difficile* we developed an *in silico* invasion assay that

leverages microbiome relative abundance data, manually curated genome-scale

metabolic models of gut bacteria from the AGORA database, constraints on the

diet/environment, and the MICOM modeling framework (Fig. 1A) [20,23]. Here, we used

several existing 16S amplicon and shotgun metagenomic sequencing data sets to

validate our approach [10,18,24–27]. Amplicon sequencing data is often limited to genus-

level resolution in the taxonomic classifications of amplicon sequence variants (ASVs).

Even with metagenomic sequencing data, species level resolution can be poor when

mapping to available species-level metabolic models. Generally, only the most

abundant or prevalent commensal bacterial species are well represented. Therefore, we

focused on genus-level MCMMs, to approximate the community metabolic context, for

our invasion assays (see Methods) [28]. However, we also evaluated the use of species-

level MCMMs using *in vitro* 16S amplicon and stool shotgun metagenomic data (Fig.

S1A). Briefly, strain-level metabolic models from AGORA were combined at the genus

or species level, to account for potential coexistence of multiple strains from a given

genus or species within an individual and to reduce potential bias from arbitrarily

selecting individual strain models. Using this approach ~75% of reads, on average,

could be mapped to a genus-level metabolic model within the AGORA database from

16S amplicon data, while ~90% of reads, on average, could be mapped using shotgun

metagenomic data. At the species level, only ~50% of reads could be mapped, on

average, using the same *in vivo* metagenomics data, while 100% of reads could be

mapped for *in vitro* communities profiles using 16S amplicon sequencing (i.e., because

type strains were used in these experiments), illustrating the inherent limitation of using

species level models for *in vivo* applications and further justifying our focus on genus-

level analyses (Fig. S1A). To simulate the invasion of *C. difficile* into these model

communities, a pan-species model of *Clostridioides*, representing all four common *C.*

difficile strains present in the AGORA database (including hypervirulent and non-

epidemic strains), was introduced at a relative abundance of 10% (see below for

justification for this percentage), while other community relative abundances were

decreased proportionally to approximate a minor perturbation in community-wide

biomass (Figure 1A). Growth simulations were then performed using a medium

representing an average European diet (i.e., a standard developed-world diet

appropriate to the cohorts studied here), with fluxes of metabolites known to be

absorbed in the small intestine decreased by 90%, as previously described for *in vivo*

samples [20]. To represent the anaerobic basal broth (ABB) used to culture the *in vitro*

communities, a curated medium was constructed using a previously described

representation of Luria-Bertani (LB) broth as a template and the known composition of

ABB (see Methods for further details) [27,29]. Growth rates were estimated using ctFBA,

as implemented in MICOM, which uses a regularization step and allows for a

suboptimal community growth rate in order to achieve a more realistic growth rate

distribution across the community [20,21]. Import and export fluxes were estimated using

parsimonious enzyme usage FBA (pFBA) [20].

Personalized MCMMs were constructed for each sample and the potential for *C.*

difficile engraftment was quantified as the model-inferred growth rate. ctFBA has a

single free parameter that needs to be chosen, the tradeoff between community-wide

growth rates and individual, taxon-specific growth rates. Assuming that most taxa

detected at appreciable abundances in a gut microbiome are actively growing *in vivo*

and *in vitro*, a trade-off value was selected by choosing the minimal deviation from

optimal community growth for which >70% of genera obtained non-zero growth rates on

average (Fig. S1B). We found that with a trade-off value of 0.8 (i.e., 80% of maximal

community biomass production) resulted in a mean fraction of genera with non-zero

growth of >70% using genus-level models constructed from 16S amplicon data. This

same non-zero growth fraction was achieved with a trade-off value of 0.9 for the genus-

level models constructed from metagenomics data. The 16S and metagenomics specific

tradeoff values were also used for the species level models derived from these data

types, respectively. At the species-level these values were associated with much higher

growth fractions (~90% and 95% for 16S amplicon and shotgun metagenomics,

respectively; Fig. S1B).

We first validated our approach using an *in vitro* 16S amplicon data set where *C.*

difficile was co-cultured with communities of human commensal gut bacteria at differing

levels of species richness [27]. We found that MCMM-inferred *C. difficile* biomass

(calculated using the MCMM-inferred *C. difficile* growth rates, assuming exponential

growth over a fixed interval, see Methods) accurately reflected a negative trend in

measured *C. difficile* abundance along a gradient of increasing species richness and

was significantly correlated with empirically measured *C. difficile* biomass after 48 hours

of growth (Fig. 1B; ordinary least squares (OLS) R^2=0.45, p <10^{-16}). Similarly, we found

that the normalized *C. difficile* growth rate (normalized to the overall MCMM-inferred

community growth rate, see Methods) decreased as a function of community richness

and was a even stronger predictor of the empirically-observed *C. difficile* biomass after

48 hours of growth (Fig. 1C; ordinary least squares (OLS) R^2=0.5, p <10^{-16}). Biclustering

of the inferred species growth rates suggested that growth of *C. difficile* was largely

suppressed by *B. thetaiotaomicron, P. vulgatus*, and *E. lenta in vitro*, as indicated by the

strong negative associations between the growth rates of these species and the growth

rate of *C. difficile* (Pearson *r* <-0.45, p <0.01 for species). Positive associations between

the growth rates of *C. difficile, C. scindens*, and *P. hiranonis* (Pearson *r* >0.8, p <10^{-6}

for all species) were inline with the observation that these species tend to share

overlapping metabolic niches. The previously-observed suppressive effects of *C.*

scindens and *P. hiranonis* on *C. difficile* were not apparent within our modeling

framework [27], likely due to the fact that pH and its effects on growth are not easily

captured by FBA models. However, while *in vitro* acidification is a strong environmental

force in anaerobic batch culture, we do not expect variation in pH to be as prominent in

the buffered environment of the human colon.

To further validate our approach *in vivo* we leveraged a time series with a known

C. difficile colonization event (Fig. 1E) [10,30]. We found that estimated *C. difficile* growth

rates were at or below the limit of solver accuracy (<10^{-6}, which effectively indicates an

absence of growth) in samples collected prior to colonization and comparable to growth

rates of other dominant genera in samples taken after the initial colonization event

(Welch's t-test *t*=-3.29, *p*=0.003, for comparison of $\log_{10}$ growth rates before and after

the known colonization event; Fig. 1E). Furthermore, we saw patchy engraftment

predictions in a second individual that was known to be colonized by *C. difficile* at a low

level (i.e., near the limit of detection) throughout a time series (Fig. 1F). We also

assessed the importance of propagule pressure [31] (i.e., the relative abundance at which

the invasive taxon is introduced into the models) and found that below 10% relative

abundance, agreement between growth rate estimates and measured abundances was

poor (Fig. S1C). Thus, propagule pressure plays an important role in predicted

engraftment success [27].

Based on these initial results (Fig. S1), we decided to use a fixed tradeoff value

of 0.8 and a *C. difficile* invasion fraction of 10% for all subsequent analyses of 16S

amplicon data (i.e., the highest value at which >70% of the community showed a

positive growth rate). Using the same heuristic, a tradeoff value of 0.9 was chosen for

the metagenomics analyses, along with a 10% *C. difficile* invasion fraction (Fig. S1B).

in silico invasion assay accurately predicts C. difficile colonization potential in rCDI
patients pre- and post- FMT

We applied our *in silico* invasion model to two separate datasets (16S amplicon and

metagenomic shotgun sequencing, respectively) of rCDI patients who received FMTs

and were subsequently followed over time [18,26]. These data provided an additional test

of MCMM performance and a means to explore the metabolic features associated with

community-scale colonization susceptibility or resistance across a larger population.

Given that all individuals in the rCDI cohorts had experienced multiple recurrent

infections, we expected pre-FMT microbiomes from these patients to be more

susceptible to invasion. Additionally, both of the original studies found that nearly all the

patient microbiomes transitioned to a compositional state that was much closer to the

healthy controls post-FMT than to their pre-FMT compositional states [18,26]. Thus, we

expected post-FMT samples would be less susceptible to invasion. For the sake of

comparison of results across studies, genus-level MCMMs were used for both analyses.

Across both data sets, rCDI patients had significantly higher MCMM-predicted *C.*

difficile growth rates prior to FMT treatment than they did post FMT treatment (Fig. 2A;

Welch's t-test t=-3.19, p=0.001 for comparison of pre-FMT vs. post-FMT). Predicted *C.*

difficile growth rates were negatively associated with Shannon diversity, albeit weakly

(Fig. 2B; OLS R^2=0.05, p=0.01), which is in line with prior empirical observations

indicating that lower diversity communities are more susceptible to *C. difficile*

colonization and the development of rCDI [9,27,32,33].

The community-scale import flux profile prior to *in silico* invasion was predictive

of *C. difficile* growth rate following invasion in Weingarden *et al.* (Fig. 2C). High-

dimensional community-scale import flux profiles were projected into a two-dimensional

space using the Uniform Manifold Approximation and Projection (UMAP) technique (Fig.

2C) [34]. The UMAP projection provides a visual means of identifying patterns in the high

dimensional import flux space. The closer points are to one another in this ordination

the more similar their import flux profiles are. Thus, clusters of points in the UMAP can

represent distinct metabolic environments across samples. The ordination plot indicated

that *C. difficile* appears to grow well in more than one metabolic environment, when

colonizing different individuals. Indeed, we saw that the predicted metabolic

environments occupied by *C. difficile* could vary within an individual over time (Fig. 2C).

For most patients in the Weingarden *et al.* cohort, there was a transition from

colonization-susceptibility pre-FMT to colonization-resistance post-FMT (Fig. 2A,C). We

next examined the different metabolic niches that *C. difficile* was able to exploit when

colonizing individuals across both rCDI-FMT cohorts, to better understand this

phenotypic plasticity.

C. difficile is predicted to occupy three distinct metabolic niches within the human gut
microbiome

To characterize *C. difficile* colonization-associated niches and identify the potential for

multiple metabolic strategies associated with its growth, we examined *C. difficile* import

fluxes with high variance ($\log_{10}$ flux variance >=4.5) across the rCDI-FMT cohorts. Using

this criteria the high variance metabolites identified across the 16S and metagenomic

data sets were highly consistent (Fig. 3). We identified 31 and 44 high-variance

metabolite fluxes from the 16S and metagenomics analyses, respectively. Of these, 27

were shared, 4 were unique to the Weingarden *et al.* (16S) analysis, and 17 were

unique to Iraniro *et al.* (metagenomics) analysis. Biclustering of the high variance import

flux data and an examination of how the apparent clusters associated with growth rates

revealed that *C. difficile* makes use of multiple metabolic strategies (Fig. 3). Three major

clusters were observed across patient samples and cohorts. We designate these three

clusters as "high growth", "moderate growth" and "no growth" (Fig. 3). The high growth

cluster in both cohorts included many of the pre-FMT samples and was characterized

by consistently high import fluxes for all the metabolites identified as most strongly

coupled to *C. difficile* growth across all models. The moderate growth cluster showed a

sparser metabolite consumption profile. For example, ornithine and fructose were

rapidly consumed in the high growth cluster, but showed almost no consumption in the

moderate growth cluster (Fig. 3). As expected, very few metabolites were consumed by

C. difficile above the zero-threshold of 10^{-6} in the no growth cluster (Fig. 3).

The metabolic strategies employed by *C. difficile* within the MCMMs showed

convergence with several observations from the literature. For example, we found that

metabolites known to promote growth of *C. difficile in vivo* (e.g., succinate, ornithine,

and trehalose) were preferentially utilized when available and were associated with

higher pathobiont growth rates [35–37]. In addition, the consumption of amino acids valine,

leucine, glycine, glutamate, glutamine, and proline were associated with higher *C.*

difficile growth rates in the MCMMs, implying that *C. difficile* employs Strickland

fermentation in one of its growth modes, which has been observed empirically [38].

Following up on these findings we examined how cooperative and competitive

interactions within MCMMs contributed to *C. difficile* colonization. To accomplish this,

we examined the import and export fluxes of metabolites associated with *C. difficile*

colonization (e.g., amino acids, ornithine, succinate, etc.). Genera that produced

metabolites consumed by *C. difficile* likely promote its growth, while those consuming *C.*

difficile growth-associated metabolites may be in direct competition. For ornithine and

succinate, we found that cooperative and competitive interactions are context-

dependent, varying across samples. The genus *Phocaeicola*, for instance, produces

ornithine in some samples, which is in turn consumed by *C. difficile*, while in other

contexts it consumes ornithine, competing with *C. difficile* (Fig. 4A). Meanwhile,

Roseburia, and *Faecalibacterium* compete with *C. difficile* for ornithine, but these

genera also produce succinate and leucine in some contexts, which *C. difficile*

consumes (Fig. 4A-B). Thus, the overall community context, rather than the presence or

absence of any single taxon, appears to be the most important factor in determining the

metabolic strategies used by *C. difficile* and can lead to emergent competitive or

cooperative interactions, which can either hinder or promote colonization.

Finally, we assessed whether or not compositional variation in the microbiome

alone could explain observed differences in MCMM-predicted *C. difficile* growth rates.

We found that compositional variation was a modest predictor of estimated *C. difficile*

growth rate (out-of-sample least absolute shrinkage and selection operator (LASSO)

regression R^2=0.37 for the Weingarden *et al.* rCDI-FMT cohort, see Methods).

Meanwhile, the import flux derived growth clusters (e.g., "high growth", "medium

growth", and "no growth" groups) could explain the vast majority of the variance in

predicted *C. difficile* growth rates (analysis of variance (ANOVA) R^2=0.94 for the CDI-

FMT cohort). However, this comparison between model-estimated growth rates and

growth clusters is a bit circular (i.e., both are derived from the model), so as mentioned

above, we also observed that MCMM-estimated *C. difficile* growth rates were able to

explain 50% of the variance in empirical *C. difficile* biomass measurements *in vitro* (Fig.

1C). In summary, these results suggest that compositional information alone is not

sufficient for consistently accurate engraftment predictions.

Associations with C. difficile growth provide insights into the role of community context

In order to assess the consistency of the *C. difficile* growth clusters, we leveraged four

independent *in vivo* 16S amplicon data sets, including the time series and the

Weingarden *et al.* rCDI-FMT studies presented above (Figs. 1-4), along with two large

cross-sectional cohorts (i.e., the American Gut and Arivale cohorts), covering a total of

14,862 individuals [24,30,39,40]. We evaluated growth and flux predictions generated across

all four data sets and found that *C. difficile* fell into the same three clusters that were

identified in the rCDI-FMT data sets, representing no growth, moderate growth, and

high growth (Fig. 5A).

To further contextualize the metabolic niche of *C. difficile*, we integrated model

outputs for all four 16S data sets. Specifically, we looked at import fluxes across all

genera. Most genera formed unique clusters in the UMAP projection, suggesting that

each genus had a single metabolic niche that was consistent across datasets (Fig. 5B).

Within this community context, we found that *C. difficile* still fell into three very distinct

clusters (Fig. 5B). Several genera that showed some of the strongest competitive and

cooperative interactions with *C. difficile* across data sets, *Blautia*, *Faecalibacterium*,

Ruminococcus and *Dorea*, clustered near one another in import flux space, with some

overlap, indicating that these commensal taxa shared a similar metabolic niche (Fig.

5B). Meanwhile, *Bacteroides* and *Phocaeicola*, two apparent *C. difficile* competitors,

formed clusters largely separate from those formed by the other genera. Several of

these taxa also occupied niche space near the *C. difficile* moderate growth cluster,

supporting the potential for competitive interactions between *C. difficile* and these

genera. The *C. difficile* high growth cluster was more separated from other commensal

genera, suggesting that this growth mode is more specialized in the context of the

surrounding community (Fig. 5B).

We next explored gut community diversity and predicted *C. difficile* growth rates

across the four data sets. Specifically, we looked at Shannon diversity, which integrates

species richness and evenness and is commonly used to quantify gut microbiome

alpha-diversity. Lower Shannon diversity is commonly associated with disease states,

like diarrhea, while higher diversity has generally been associated with diverse plant-

based diets and overall better health [32]. However, constipated individuals generally

have higher gut microbiome alpha-diversity as well, suggesting that there may be an

optimal range of alpha-diversity across healthy individuals [39,41]. Our initial analysis using

the rCDI-FMT cohort suggested a monotonically negative relationship between

predicted *C. difficile* growth rate and Shannon (Fig. 2B). However, the integrated data

sets, which spanned a wider range of diversity, showed a U-shaped relationship

between Shannon diversity and predicted *C. difficile* growth rate (Fig. 5C). Intermediate

levels of Shannon diversity were associated with the lowest predicted growth rates, on

average, with higher average growth at the upper and lower tails of the distribution (Fig.

5C). The relationship between Shannon diversity and predicted growth rate suggests

extremes in either direction on the diversity scale are, on average, more permissive to

C. difficile engraftment.

Blood metabolites and clinical labs associated with MCMM-predicted C. difficile
colonization susceptibility

We next sought to identify potential blood-based markers that were significantly

associated with MCMM-predicted *C. difficile* growth rates. Previous work has shown

that circulating blood metabolites can be leveraged to predict gut microbiome alpha-

diversity [39]. We identified several blood metabolites and clinical chemistries significantly

associated with MCMM-predicted *C. difficile* growth rates in the Arivale cohort, after

adjusting for common covariates (i.e., sex, age, and BMI) and correcting for multiple

tests (FDR q <0.05). These included two secondary bile acids, an unannotated

metabolite previously associated with the abundance of the family *Eggerthellacea*, and

several red blood cell-associated clinical chemistries (Fig. S1D)[42]. Unfortunately, while

significant, these blood-based markers, along with sex, age, and BMI, collectively

accounted for only ~5% of the variance in MCMM-predicted *C. difficile* growth rates.

Thus, it appears MCMM-based estimates of *C. difficile* engraftment, constrained by

fecal microbiome data, cannot be readily replaced with commonly measured clinical

chemistries or blood metabolites.

MCMMs predicts engraftment heterogeneity of probiotic cocktail designed to treat rCDI

As a final proof-of-concept for our modeling framework we simulated a probiotic

intervention using a previously validated probiotic cocktail designed to treat rCDI [22,43].

The probiotic, referred to as VE303, was composed of 8 commensal Clostridia strains

and shown to be effective at treating CDI in mice [43]. This probiotic was also shown to be

safe, well-tolerated, and effective in reducing rCDI incidence in humans [22]. An earlier

study in both mice and humans found that engraftment of VE303 strains was optimal

following antibiotic pretreatment [43]. With these facts in hand, we designed a simulated

intervention that mimicked the treatment found to be most effective by Dsouza *et al* [43].

We were only able to identify metabolic models for 6 of the 8 strains in VE303 in the

AGORA database [23]. We leveraged the Weingarden *et al.* rCDI-FMT dataset to test this

six-member probiotic cocktail, paired with *in silico* invasion by *C. difficile*. The probiotic

cocktail was introduced to patient samples, alongside 10% *C. difficile*, at a total relative

abundance of 50%, which was evenly distributed among the six strains. We also

simulated vancomycin treatment by reducing the abundance of *C. difficile* and all

commensal genera known to be impacted by vancomycin treatment [44] by 90%. We

found that a combined probiotic and antibiotic intervention most effectively suppressed

the growth of *C. difficile* in both the moderate and high *C. difficile* growth rate clusters

(Fig. 6A).

To better understand the mechanism of action of the probiotic cocktail, we

assessed the growth characteristics and the niche proximity of the probiotic strains in

relation to *C. difficile.* We found that suppression of *C. difficile* growth occurred more

readily when the average growth of the probiotic strains was high ($>10^{-4}$, Fig. 6B) and

when the average niche distance between the probiotic strains and *C. difficile* was low

(<25 import flux Euclidean distance, Fig. 6C). We also found that, relative to other

genera, several of the probiotics strains occupied niches closer to *C. difficile*, although

these niche distances could vary widely for each organism depending on their

community context (Fig. S2). Finally, we compared the import fluxes of the probiotic

strains and *C. difficile* for the metabolites identified as important for *C. difficile* growth

(Fig. 3). This analysis showed that, in addition to occupying niches similar to *C. difficile*,

several of the probiotic strains directly competed for metabolites important for *C. difficile*

growth, such as succinate, ornithine, and trehalose (Fig. 6D). Cumulatively these results

suggest that metabolic competition is the mechanism by which the probiotic cocktail

suppressed *C. difficile* growth, which is consistent with the emerging consensus in the

field [9,43,45]. Finally, we found that certain probiotic strains were more or less likely to

engraft in an individual (Fig. 6D), and that this engraftment/growth was generally

associated with *C. difficile* suppression (Fig. 6B), which indicates that MCMMs can be

leveraged to identify responders and non-responders prior to these kinds of probiotic

interventions.

Discussion

In this study, we provide a framework for predicting *C. difficile* engraftment risk in the

human gut microbiome using MCMMs. While we focus on *C. difficile*, due to its clinical

importance, this approach could be extended to other opportunistic bacterial pathogens,

probiotic organisms, or even entire communities, in the case of FMTs. We were able to

show how our approach predicts expected longitudinal and cross-sectional variation in

C. difficile colonization potential *in vitro* and *in vivo* (Figs. 1-2), using both shotgun

metagenomic and 16S amplicon sequencing data sets, and we provide insights into the

metabolic strategies leveraged by *C. difficile* in different ecological contexts (Figs. 3-4).

Our analysis not only recapitulates known metabolic associations with *C. difficile* growth

(e.g., consumption of trehalose, ornithine, and succinate; Fig. 3), it suggests additional

associations (e.g., importance of reduced sulfur compounds like cysteine, Stickland

fermentation reactants, and utilization of dietary sugars, like fructose; Fig. 3).

Additionally, we show that competition and cooperation with community members can

prevent or promote colonization of *C. difficile*, and that many of these associations are

highly context-dependent (Fig. 4).

Consistent with the idea that simple metrics of community structure and

composition alone are not effective predictors of colonization susceptibility, we found

that community compositional variation was a modest predictor of estimated *C. difficile*

growth rate and the relationship between alpha diversity and estimated *C. difficile*

growth rate was nonlinear (Fig. 5C). Not only did low diversity communities tend to be

more invasible, as might be expected due to putative non-saturation of the metabolic

niche space [9], but high diversity communities were also more prone to *C. difficile*

engraftment (Fig. 5C). In high diversity communities, successful colonization may be

due to changes in the interaction landscape, like a higher rate of niche-construction in

higher-diversity systems, which would be in line with the diversity-begets-diversity

hypothesis [46]. An intermediate range of alpha-diversity seems to be optimal for

mitigating *C. difficile* colonization potential (Figure 5C). Overall, these complex

mappings between community composition and pathobiont engraftment risk underscore

the necessity of systems-scale tools, like MCMMs, that are capable of synthesizing this

complexity.

Several genera were found to engage in cooperative and competitive interactions

with *C. difficile* across MCMMs. *Blautia*, *Faecalibacterium*, and *Dorea* were all shown to

benefit *C. difficile* through production of key metabolites that it consumes, like succinate

and leucine, but were also capable of competition for other metabolic resources (Fig. 4).

Meanwhile *Rumminococcus*, *Bacteroides* and *Phocaeicola* growth rates were often

negatively associated with *C. difficile* engraftment, competing for some of the same

metabolites that *C. difficile* consumed at high rates (Fig. 1D, 4). Contextualizing these

results through analysis of individual taxon import fluxes across studies we found that

Blautia, *Faecalibacterium*, *Phocaeicola, Rumminococcus,* and *Dorea* share similar

niches with one another, with *Phocaeicola being the most divergent from the others*. In

most cases, these niches did not overlap with *C. difficile*, but in a subset of individuals

all five occupied niches states overlapping or similar to *C. difficile* (Fig. 5B). Thus, while

we observed competition for some key metabolites, on a global scale, the majority of

the metabolic niche space used by *C. difficile* tends not to overlap with its apparent

competitors (Fig. 5B). These results highlight how flexible commensal gut bacteria are

in adapting their import fluxes to the communities in which they reside, which in turn

suggests why so many taxa are able to coexist. These results also suggest that

colonization resistance is an emergent feature of multi-species metabolic interactions,

and not strongly driven by any particular pairwise interaction.

In addition to developing a simulation framework to predict engraftment we

sought to identify blood-based clinical chemistries and blood metabolites that were

associated with MCMM-inferred *C. difficile* growth rate. We found three blood

metabolites that were independently associated with predicted *C. difficile* growth rates.

These included two secondary bile acids and an unannotated metabolite. One of the

secondary bile acids, isoursodeoxycholate, has previously been positively associated

with the abundance of *Bacteroides* [39], and was negatively associated with predicted *C.*

difficile growth rates. This result is in line with the apparent competition between

Bacteroides and *C. difficile* in our MCMMs. We also identified several clinical labs

negatively associated with predicted growth rate (Fig. S1D). However, together with

age, sex, and BMI, these features only accounted for ~5% of the variance in predicted

growth rates. Thus, while these features may be signatures for colonization

susceptibility in the blood, their clinical relevance is limited at this time.

Finally, we demonstrated that a probiotic intervention (i.e., 6/8 strains from

VE303), which recently showed positive efficacy results in a double-blinded, placebo-

controlled clinical trial for the treatment of rCDI [22], suppresses the growth of *C. difficile in*

silico in most people in the Weingarden *et al.* rCDI-FMT study (Fig. 6). We also showed

that the mechanism of action for this particular probiotic is likely competition for

metabolites essential for the growth of *C. difficile*, as many of the probiotic strains

occupy niches close to *C. difficile* and directly compete for these metabolites, such as

succinate and ornithine, in samples where growth suppression was observed (Fig. 6).

Furthermore, analysis of niche distances between *C. difficile* and other genera across

donors suggest selecting strains from *Blautia and Dorea* (e.g., including *B. producta and*

D. longicatana, from VE303), in addition to *Anaerostipes*, *Roseburia*, and

Faecalibacterium, could be leveraged to design better individual-specific probiotic

cocktails capable of suppressing *C. difficile* and rescuing VE303 non-responders (Figs.

S2 and 6B). These results illustrate how MCMMs may be powerful tools for assessing

the individual-specific efficacy of clinically-relevant probiotics, in addition to

understanding personalized pathobiont colonization susceptibility.

Future work should test this MCMM framework in the design of precision

interventions to prevent engraftment of *C. difficile* and other pathobionts, to design

precision probiotic interventions, and to improve the production of beneficial microbial

metabolites, like short-chain-fatty-acids, or reduce the production of undesired

metabolites, such as hydrogen sulfide or trimethylamine [47–49]. In summary, MCMMs

present a promising new path forward in engineering the ecological composition and

metabolic outputs of microbiota to prevent or treat disease.

497
498   **Methods**
499

External Data Collection and Processing

Data used in this study came from six sources. This included both cross-sectional and

time series 16S amplicon and shotgun metagenomics sequencing data from Hromada

et al., David *et al.*, Weingarden, A. *et al.*, Iraniro *et al.,* the American Gut (McDonald, D.

et al.), and a former scientific wellness program run by Arivale, Inc.[18,24,26,27,30,40]. Publicly

available 16S amplicon sequence data and associated metadata were downloaded from

the sequence read archive (SRA). Additionally, de-identified 16S amplicon sequence

data, associated metadata, and paired blood-based clinical chemistries and

metabolomics were obtained for 2,687 research consenting individuals that were

formerly participants in the Arivale wellness program. Raw 16S amplicon sequence data

were processed using QIIME2 (v2020.11.1). In brief, the QIIME2 workflow consisted of

read demultiplexing using the command *qiime tools import*, and an associated manifest

table for each study describing read metadata followed by read quality assessment

using *qiime demux summarize*. Read quality assessment was used to determine

trimming parameters for subsequent denoising using the QIIME2 implementation of

DADA2 via the command, *qiime dada2 denoise-single* or *qiime dada2 denoise-paired*,

for single and paired reads respectively. The first 10 bases were trimmed from all reads

and reads were truncated to a length where median quality score was >20 (100-150

base pair for the data leveraged). Following denoising, data were reformatted into a

table format using the command *qiime metadata tabulate*, and representative sequence

taxonomy was inferred using a custom NCBI classifier with the command *qiime feature-*

classifier classify-sklearn. The NCBI classifier was trained using 16S 515f-806r V4

regions extracted from all available bacterial NCBI genomes. To train the classifier 515f-

806r regions were extracted from NCBI sequences using the command *qiime feature-*

classifier extract-reads, followed by the command *qiime feature-classifier fit-classifier-*

naive-bayes using the extracted V4 sequences and a table of known taxonomies. Pre-

processed data from Iraniro *et al.* were provided by the authors upon request and

included sample taxonomic relative abundance inferred using Metaphlan4 and

associated sample metadata. See Iraniro *et al.* for further details on their data

processing pipelines [18]. Sample taxonomic relative abundance and associated

metadata from Hromada *et al.* were obtained from their supplemental data. See

Hromada *et al.* for further details [27]. For source code and tables of processed data refer

to the Github repository listed below in Data and Source Code Availability.

534
Model Construction and growth simulations

To construct community level metabolic models sample specific taxonomic abundance

profiles inferred from 16S amplicon or shotgun metagenomic sequencing were

summarized at the genus or species level and mapped to genus or species level

metabolic models from the AGORA database (v1.03) using MICOM (v0.25.1). Taxa with

a relative abundance less than 0.1% were omitted from community models. An *in silico*

media previously designed to represent an average western diet was applied which

defined the bounds for metabolic imports by the model communities for all *in vivo*

analyses [20,23]. For the *in vitro* communities a custom medium was developed to reflect

the ABB medium used experimentally. The custom medium was developed using a

previously defined template designed for the representation of Luria-Bertani (LB)

medium, this contained metabolite concentration and fluxes for both yeast extract and

tryptone [29]. These details were used to convert the quantities of yeast extract and

peptone present in the experimental medium anaerobic basal broth (ABB) to fluxes. The

additional defined components were included with fluxes equal to the millimolar

concentration per hour. Finally, additional components were added to the constructed

medium so that all 13 species used in the *in vitro* communities could achieve a growth

rate of 10^{-3} or greater using a routine implemented in the MICOM function

complete_db_medium. Across all conditions, growth rates were inferred using

cooperative tradeoff flux balance analysis (ctFBA). In brief, this is a two-step

optimization scheme, where the first step finds the largest possible biomass production

rate for the full microbial community and the second step infers taxon-specific growth

rates and fluxes, while maintaining community growth within a fraction of the theoretical

maximum (i.e., the tradeoff parameter), thus balancing individual growth rates and the

community-wide growth rate[20]. For all models in the manuscript that leveraged 16S

amplicon data we used a tradeoff parameter of 0.8. A tradeoff parameter of 0.9 was

used for all the results derived from shotgun metagenomics data. These parameter

values were chosen by identifying the largest tradeoff value (corresponding to the

smallest deviation from maximal community biomass) that allowed >70% taxa to grow

(growth rate $>10^{-6}$). Import and export fluxes were estimated using parsimonious

enzyme usage FBA (pFBA) and a defined medium constructed to represent an average

European diet or ABB [20]. pFBA further constrained simulation results by requiring

genera to utilize the lowest overall flux through their networks to achieve maximal

growth [19]. For source code and tables of processed data refer to the Github repository

listed below in Data and Source Code Availability.

Biomass estimation and growth rate normalization for in vitro communities

To allow better comparison between model predictions and experimental results two

growth rate transformations were employed. These methods were implemented to

account for the fact that batch growth (in which cultures grow on a limited quantity of

nutrients) is not compatible with the steady state assumption employed by the modeling

framework (which assumes continuous supply of nutrients). Estimation of biomass was

leveraged to compare growth rates with experimentally measured end point optical

densities. This consisted of using a simple exponential growth model of the form $OD_f =$

$OD_i e^{kt}$, where OD_f is the final predicted biomass, OD_i is the initial biomass, k is the

MCMM inferred growth rate, and t is time in hours. The second transformation

employed was a growth rate normalization procedure, which accounted for differences

in community growth rates across samples. Growth rates were divided by the MCMM

inferred community growth rate, which was computed from the sum of relative growth

rates of each taxon in the system (e.g., the product of growth rate and relative

abundance), $k_{norm} = \frac{k}{\sum_i^n k_i r_i}$.

Probiotic Intervention

To model probiotic intervention a combination of strains previously shown to be effective

at suppressing the growth of *C. difficile* in mice were used [43]. Metabolic models for six of

the eight stains in the VE303 cocktail described by Dsouza *et al.* were identified in the

AGORA database and intervention was simulated by introducing them along with *C.*

difficile to individual samples. A total probiotic fraction of 50% was used, which was

evenly distributed among the six strains. This fraction was determined to be the most

effective at suppressing the growth of *C. difficile* growth *in silico* for the samples tested

(data not shown). Vancomycin treatment was simulated by reducing the abundance of

C. difficile and all genera known to be impacted by vancomycin by 90% [44]. Growth

simulations were performed as described above. For source code and tables of

processed data refer to the Github repository listed below in Data and Source Code

Availability.

Statistical analysis

Statistical analyses were performed using functions from the python scipy (v1.7.1),

seaborn (v0.11.2), sklearn-learn (v0.24.2), umap-learn (v0.5.1), and statsmodels

(v0.13.1) packages. Linear associations were performed using the *statsmodels* ordinary

least squares function *OLS*, and visualized using the *seaborn* function *regplot*. Least

absolute shrinkage and selection operator (LASSO) was performed using the *sklearn*

Lasso function and a training-test framework. Data were split into training and test sets

(70% of samples were randomly assigned to the training set) and model performance

was assessed across a range of regularization values. LASSO training and test set R^2

values were used to select the model with the best test set R^2 that did not overfit

training data (training R^2 >> test R^2). Analysis of variance (ANOVA) was performed

using the *statsmodels OLS* and *anova_lm* functions. UMAP dimensionality reduction

was performed using the *umap* function from the *umap* package and associated

methods with default parameters (i.e., *n_components=2, n_neighbors=15,*

metric='euclidean', etc.). Biclustering was performed using the *seaborn* function

clustermap and the Ward clustering algorithm. Hexagonal binning and associated

histograms were generated using the *seaborn* function *jointplot.* Locally weighted

scatterplot smoothing (LOWESS) curves were generated using the *lowess* function from

statsmodels with default parameters. Additional statistical tests included the t-test and

Wilcox rank sum test implemented in scipy as *ttest_ind* and *wilcoxon* respectively. For

source code and tables of processed data refer to the Github repository listed below in

Data and Source Code Availability.

Data and Source Code Availability

Acknowledgements

We thank members of the Gibbons Lab for helpful feedback and suggestions on this

work. We thank Nicola Segata, Michal Punčochář, and Nicolai Karcher for providing

access to their post-processing data files. This work was supported by a Washington

Research Foundation Distinguished Investigator Award (to SMG). Research reported in

this publication was also supported by the National Institute of Diabetes and Digestive

and Kidney Diseases of the National Institutes of Health (NIH) under award no.

R01DK133468 (to SMG.).

References:

1. Fan, Y. & Pedersen, O. Gut microbiota in human metabolic health and disease. *Nat. Rev. Microbiol.* **19**, 55–71 (2021).

2. Zeevi, D. *et al.* Personalized Nutrition by Prediction of Glycemic Responses. *Cell* **163**, 1079–1094 (2015).

3. Schubert, A. M., Sinani, H. & Schloss, P. D. Antibiotic-Induced Alterations of the Murine Gut Microbiota and Subsequent Effects on Colonization Resistance against Clostridium difficile. *MBio* **6**, e00974 (2015).

4. Donaldson, G. P., Lee, S. M. & Mazmanian, S. K. Gut biogeography of the bacterial microbiota. *Nat. Rev. Microbiol.* **14**, 20–32 (2016).

5. Bell, A. & Juge, N. Mucosal glycan degradation of the host by the gut microbiota. *Glycobiology* **31**, 691–696 (2021).

6. Baxter, N. T. *et al.* Dynamics of Human Gut Microbiota and Short-Chain Fatty Acids in Response to Dietary Interventions with Three Fermentable Fibers. *MBio* **10**, (2019).

7. Gibbons, S. M. Defining Microbiome Health through a Host Lens. *mSystems* **4**, (2019).

8. Cheng, A. G. *et al.* Design, construction, and in vivo augmentation of a complex gut

microbiome. *Cell* **185**, 3617–3636.e19 (2022).

9.   Spragge, F. *et al.* Microbiome diversity protects against pathogens by nutrient

blocking. *Science* **382**, eadj3502 (2023).

10.   VanInsberghe, D. *et al.* Diarrhoeal events can trigger long-term Clostridium difficile

colonization with recurrent blooms. *Nat Microbiol* **5**, 642–650 (2020).

11.   Shields, K., Araujo-Castillo, R. V., Theethira, T. G., Alonso, C. D. & Kelly, C. P.

Recurrent Clostridium difficile infection: From colonization to cure. *Anaerobe* **34**,

59–73 (2015).

12.   Song, J. H. & Kim, Y. S. Recurrent Clostridium difficile Infection: Risk Factors,

Treatment, and Prevention. *Gut Liver* **13**, 16–24 (2019).

13.   Dubberke, E. R. & Olsen, M. A. Burden of Clostridium difficile on the healthcare

system. *Clin. Infect. Dis.* **55 Suppl 2**, S88–92 (2012).

14.   Banaei, N., Anikst, V. & Schroeder, L. F. Burden of Clostridium difficile infection in

the United States. *The New England journal of medicine* vol. 372 2368–2369

(2015).

15.   Crobach, M. J. T. *et al.* Understanding Clostridium difficile Colonization. *Clin.*

Microbiol. Rev. **31**, (2018).

16.   Baron, S. W. *et al.* Screening of carriers in an urban academic medical center:

Understanding implications of disease. *Infect. Control Hosp. Epidemiol.* **41**, 149–

153 (2020).

17.   Khoruts, A., Staley, C. & Sadowsky, M. J. Faecal microbiota transplantation for

Clostridioides difficile: mechanisms and pharmacology. *Nat. Rev. Gastroenterol.*

Hepatol. **18**, 67–80 (2021).

18. Ianiro, G. *et al.* Variability of strain engraftment and predictability of microbiome

composition after fecal microbiota transplantation across different diseases. *Nat.*

Med. **28**, 1913–1923 (2022).

19. Lewis, N. E. *et al.* Omic data from evolved E. coli are consistent with computed

optimal growth from genome-scale models. *Mol. Syst. Biol.* **6**, 390 (2010).

20. Diener, C., Gibbons, S. M. & Resendis-Antonio, O. MICOM: Metagenome-Scale

Modeling To Infer Metabolic Interactions in the Gut Microbiota. *mSystems* **5**,

(2020).

21. Diener, C. & Gibbons, S. M. More is Different: Metabolic Modeling of Diverse

Microbial Communities. *mSystems* e0127022 (2023).

22. Louie, T. *et al.* VE303, a Defined Bacterial Consortium, for Prevention of Recurrent

Clostridioides difficile Infection: A Randomized Clinical Trial. *JAMA* (2023)

doi:10.1001/jama.2023.4314.

23. Magnúsdóttir, S. *et al.* Generation of genome-scale metabolic reconstructions for

773 members of the human gut microbiota. *Nat. Biotechnol.* **35**, 81–89 (2017).

24. McDonald, D. *et al.* American Gut: an Open Platform for Citizen Science

Microbiome Research. *mSystems* **3**, (2018).

25. Wilmanski, T. *et al.* Gut microbiome pattern reflects healthy ageing and predicts

survival in humans. *Nat Metab* **3**, 274–286 (2021).

26. Weingarden, A. *et al.* Dynamic changes in short- and long-term bacterial

composition following fecal microbiota transplantation for recurrent Clostridium

difficile infection. *Microbiome* vol. 3 Preprint at https://doi.org/10.1186/s40168-015-

0070-0 (2015).

27. Hromada, S. *et al.* Negative interactions determine Clostridioides difficile growth in

synthetic human gut communities. *Mol. Syst. Biol.* **17**, e10355 (2021).

28. Bolyen, E. *et al.* Reproducible, interactive, scalable and extensible microbiome data

science using QIIME 2. *Nat. Biotechnol.* **37**, 852–857 (2019).

29. Marinos, G., Kaleta, C. & Waschina, S. Defining the nutritional input for genome-

scale metabolic models: A roadmap. *PLoS One* **15**, e0236890 (2020).

30. David, L. A. *et al.* Host lifestyle affects human microbiota on daily timescales.

Genome Biol. **15**, R89 (2014).

31. Lockwood, J. L., Cassey, P. & Blackburn, T. The role of propagule pressure in

explaining species invasions. *Trends Ecol. Evol.* **20**, 223–228 (2005).

32. Chang, J. Y. *et al.* Decreased Diversity of the Fecal Microbiome in Recurrent

Clostridium difficile—Associated Diarrhea. *J. Infect. Dis.* **197**, 435–438 (2008).

33. Pakpour, S. *et al.* Identifying predictive features of Clostridium difficile infection

recurrence before, during, and after primary antibiotic treatment. *Microbiome* **5**, 148

(2017).

34. McInnes, L., Healy, J. & Melville, J. UMAP: Uniform Manifold Approximation and

Projection for Dimension Reduction. *arXiv [stat.ML]* (2018).

35. Ferreyra, J. A. *et al.* Gut microbiota-produced succinate promotes C. difficile

infection after antibiotic treatment or motility disturbance. *Cell Host Microbe* **16**,

770–777 (2014).

36. Pruss, K. M. *et al.* Oxidative ornithine metabolism supports non-inflammatory C.

difficile colonization. *Nat Metab* **4**, 19–28 (2022).

37. Buckley, A. M. *et al.* Trehalose-Induced Remodelling of the Human Microbiota

Affects Clostridioides difficile Infection Outcome in an In Vitro Colonic Model: A

Pilot Study. *Front. Cell. Infect. Microbiol.* **11**, 670935 (2021).

38. Bouillaut, L., Self, W. T. & Sonenshein, A. L. Proline-dependent regulation of

Clostridium difficile Stickland metabolism. *J. Bacteriol.* **195**, 844–854 (2013).

39. Wilmanski, T. *et al.* Blood metabolome predicts gut microbiome α-diversity in

humans. *Nat. Biotechnol.* **37**, 1217–1228 (2019).

40. Manor, O. *et al.* Health and disease markers correlate with gut microbiome

composition across thousands of people. *Nat. Commun.* **11**, 5206 (2020).

41. Tian, H., Chen, Q., Yang, B., Qin, H. & Li, N. Analysis of Gut Microbiome and

Metabolite Characteristics in Patients with Slow Transit Constipation. *Dig. Dis. Sci.*

66, 3026–3035 (2021).

42. Bar, N. *et al.* A reference map of potential determinants for the human serum

metabolome. *Nature* **588**, 135–140 (2020).

43. Dsouza, M. *et al.* Colonization of the live biotherapeutic product VE303 and

modulation of the microbiota and metabolites in healthy volunteers. *Cell Host*

Microbe **30**, 583–598.e8 (2022).

44. Nazzal, L. *et al.* Effect of Vancomycin on the Gut Microbiome and Plasma

Concentrations of Gut-Derived Uremic Solutes. *Kidney Int Rep* **6**, 2122–2133

(2021).

45. Horrocks, V., King, O. G., Yip, A. Y. G., Marques, I. M. & McDonald, J. A. K. Role of

the gut microbiota in nutrient competition and protection against intestinal pathogen

colonization. *Microbiology* **169**, (2023).

46. Madi, N., Vos, M., Murall, C. L., Legendre, P. & Shapiro, B. J. Does diversity beget

diversity in microbiomes? *Elife* **9**, (2020).

47. Liu, Y. & Dai, M. Trimethylamine N-Oxide Generated by the Gut Microbiota Is

Associated with Vascular Inflammation: New Insights into Atherosclerosis.

Mediators Inflamm. **2020**, 4634172 (2020).

48. Dalile, B., Van Oudenhove, L., Vervliet, B. & Verbeke, K. The role of short-chain

fatty acids in microbiota-gut-brain communication. *Nat. Rev. Gastroenterol.*

Hepatol. **16**, 461–478 (2019).

49. Murros, K. E. Hydrogen Sulfide Produced by Gut Bacteria May Induce Parkinson's

Disease. *Cells* **11**, (2022).

767
768   **Figures:**
769

770
Figure 1. *in silco* invasion assay accurately predicts *C. difficile* engraftment   *in*

***vitro* and *in vivo*.** (**A**) Schematic illustrating the *in silico* invasion assay workflow

leveraged in this study. Personalized microbial community-scale metabolic models

(MCMMs) are supplemented with 10% of a pan-genus *Clostridioides* model to simulate

an invasion event and ctFBA was used to predict *C. difficile* engraftment and metabolic

fluxes. (**B**) Measured *C. difficile* biomass at 48 hours from *Hromada et al.* and predicted

biomass, computed from MCMM-predicted *C. difficile* growth rate and an exponential

growth model, across a gradient of *in vitro* community richness. Mean trend lines and

standard deviation are displayed for both measured (blue) and predicted (yellow)

biomass. Relationship between $\log_{10}$ predicted biomass and $\log_{10}$ biomass. Ordinary

least squares fit and 95% confidence interval are displayed, as well as regression R^2

and *p*-value (**C**) Normalized *C. difficile* growth rate along a community richness gradient

and the relationship between normalized *C. difficile* growth rate and measured $\log_{10}$

biomass. Mean trend line and standard deviation are displayed for the relationship

between normalized *C. difficile* growth rate and community richness. Ordinary least

squares fit and 95% confidence interval are displayed, as well as regression R^2 and *p*-

value, for the relationship between normalized *C. difficile* growth rate and measured

$\log_{10}$ biomass . (**D**) Biclustered normalized growth rates for species across conditions.

Species normalized growth rate is displayed using blue-to-yellow heatmap coloring for

each sample. Sample community richness is displayed using white-to-red heatmap

coloring on the top row of the plot. Pearson correlation coefficients between *C. difficile*

normalized growth rate and the normalized growth rate of each other species are

displayed using blue-to-red heatmap coloring in the leftmost column of the plot.

Colorbars at the bottom of the blot indicate the scales for the various measures. (**E**)

Donor A time series taken from David *et al.* displaying daily fluctuations in microbiome

composition over a period of several months. Composition is displayed, colored by

phylum-level annotations (different shading indicates taxonomic families). At day 150,

Donor A experienced a diarrheal event and was subsequently colonized by *C. difficile*.

Estimates of *C. difficile* relative abundance from 16S sequencing and MICOM-predicted

C. difficile growth rates are displayed. (**F**) Time series from Donor B from the same

study, who was apparently colonized by *C. difficile* (at very low relative abundances,

near the limit of detection) throughout the sampling period.

804
805
Figure 2. *C. difficile* growth rate predictions capture importance of community

context for patient recovery from CDI. (**A**) Violin plots displaying predicted *C. difficile*

$\log_{10}$ growth rate distributions across patient disease status categories. Data from

Weingarden *et al.* and Ianiro *et al.* were leveraged to validate the approach using both

16S amplicon and metagenomic sequencing. Gray shading indicates the numerical

accuracy of the solver (values below 10^{-6} can not be distinguished from zero and are

considered negligible). Bars indicate comparisons for which differences were significant

using Welch's t-test. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$. (**B**) Relationship between

predicted *C. difficile* $\log_{10}$ growth rate and Shannon diversity across datasets. Ordinary

least squares fit and 95% confidence interval are displayed, as well as regression R^2

and *p*-value. (**C**) Two-dimensional representation of community import fluxes prior to *in*

silico invasion using UMAP colored by $\log_{10}$ growth rates of *C. difficile* following *in silico*

invasion. Patient trajectories are displayed, each with a red circle representing the

patient's starting point (prior to FMT), and a red star representing the patient's end point

(post recovery).

Figure 3. *C. difficile* **occupies multiple metabolic niches across communities.** (**A**)

Biclustered *C. difficile* $\log_{10}$ import fluxes from Weingarden *et al.*, where each row is the

import flux of a particular metabolite and each column is a patient sample. (**B**)

Biclustered *C. difficile* $\log_{10}$ import fluxes from Ianiro *et al.*, where each row is the import

flux of a particular metabolite and each column is a patient sample. In each heatmap,

imports for which the log variance across samples was >=4.5 are displayed. Fluxes

across samples are displayed using blue-to-yellow heatmap coloring. *C. difficile* log$_{10}$

growth rate quantiles are displayed in white-to-red heatmap coloring for each patient

sample in the top row of each plot. Additionally, three coarse grain growth clusters are

noted. These growth clusters represent "high growth", "moderate growth", and "no

growth" phenotypes.

Figure 4. Import and export fluxes of key metabolites across communities highlight the context-dependency of key *C. difficile* competitors and cooperators. (**A**) Biclustered community import and export fluxes of specific metabolites associated with *C. difficile* colonization from Weingarden *et al.*, where each row is a genus and each column is a patient sample. (**B**) Biclustered community import and export fluxes of specific metabolites associated with *C. difficile* colonization from Ianiro *et al.*, where

each row is a genus and each column is a patient sample. For each heatmap the top 10

genera with the highest mean import or export flux across samples are displayed.

Fluxes across samples are displayed using blue-to-yellow heatmap coloring. *C. difficile*

$\log_{10}$ growth rate quantiles are displayed in white-to-red heatmap coloring for each

patient sample in the top row of each plot. Additionally, three coarse grain growth

clusters are noted. These growth clusters represent "high growth", "moderate growth",

and "no growth" phenotypes.

Figure 5. Growth niches across large four independent cohorts challenged *in*

***silico* with *C. difficile*.** (**A**) Two dimensional representation of $\log_{10}$ *C. difficile* import

fluxes using UMAP across four independent data sets. Colors denote *C. difficile* growth

rate ranging from low (blue) to high (yellow). The position of the no growth cluster is

indicated. (**B**) Two dimensional representation of $\log_{10}$ genus import fluxes using UMAP

across all datasets. Top panel displays $\log_{10}$ *C. difficile* growth rate within the context of

all other genera. Bottom panel colors *C. difficile* along with six genera of interest: *Dorea*,

Blautia, *Faecalibacterium*, *Ruminococcus*, *Bacteroides*, and *Phocaeicola*. The position

of the no growth cluster is indicated. (**C**) Two-dimensional hexagonal binning of $\log_{10}$ *C.*

difficile growth rate and community alpha diversity (Shannon index). Red trend line

indicates a LOWESS fit to the $\log_{10}$ *C. difficile* growth rate and community Shannon

diversity data.

Figure 6. Simulated probiotic intervention effectively suppresses *C. difficile*

growth *in silico*. (**A**) Box plots displaying $\log_{10}$ *C. difficile* growth rate across growth

clusters and simulated interventions. Growth clusters are those identified by bi-

clustering of *C. difficile* import fluxes using the Weingarden data (Figure 3). Conditions

include +None (no intervention control), +Probiotic (introduction of 6 strain probiotic

previously identified as an effective treatment for CDI at a total relative abundance of

50% equally distributed across the strains), +Vancomycin (90% reduction of *C. difficile*

relative abundance as well as all genera known to be impacted by Vancomycin), and

+Vancomycin, +Probiotic (introduction of 6 strain probiotic in combination with simulated

vancomycin treatment). Bars indicate comparisons for which differences were

significant using the Wilcoxon signed-rank test. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

(**B**) Relationship between *C. difficile* growth ratio and mean $\log_{10}$ probiotic growth rate.

C. difficile growth ratio is the growth rate of samples in the +Vancomycin, +Probiotic

intervention relative to +None. Values below 1 indicate growth suppression by the

probiotic and values above 1 indicate growth stimulation. The dashed line marks the

value at which no effect is observed (1). Orange trend line indicates a LOWESS fit to

the *C. difficile* growth ratio and mean $\log_{10}$ probiotic growth rate. (**C**) Relationship

between $\log_{10}$ *C. difficile* growth rate and mean probiotic niche distance. Niche distance

was calculated using the euclidean distance of $\log_{10}$ import flux vectors of each

probiotic strain relative to *C. difficile* on a per sample basis. Orange trend line indicates

a LOWESS fit to the $\log_{10}$ *C. difficile* growth rate and mean probiotic niche distance. (**D**)

Biclustered $\log_{10}$ import fluxes for *C. difficile* and probiotic strains for samples previously

identified as "high growth", where each row is the import flux of a particular metabolite

and each column is a patient sample. Imports displayed are those previously identified

as important for *C. difficile*. Color bars indicate sample *C. difficile* growth ratio and strain

specific $\log_{10}$ growth rate. Ordering of samples and metabolites is the same across

heatmaps and based on biclustering of *C. difficile* data.

Figure S1. Development of *in silico C. difficile* invasion assay. (**A**) Histograms

displaying the fraction of reads mapped at the genus and species levels for the David *et*

al. and Hromade *et al.* 16S amplicon data and Ianiro *et al.* metagenomics data using an

NCBI reference and metabolic models from the AGORA database. (**B**) Mean growth

fraction across samples, datasets, and taxonomic mappings (e.g., fraction of taxa with

estimated growth rate $>10^{-6}$) as a function of model tradeoff value. Dashed lines

indicated the tradeoff values chosen for subsequent analyses. The blue dashed line

indicates the value used for 16S data and green dashed line denotes the value used for

metagenomics data. Tradeoff values were chosen such that mean growth fractions of

~0.7 were achieved using genus level mapping. (**C**) Relationship between *C. difficile*

invasion abundance and growth rate for one of the two David *et al.* time series. (**D**)

Association coefficients for estimated *C. difficile* log growth rate, blood metabolite

concentrations, and clinical labs for the Arivale cohort.

Figure S2. Probiotic strains and associated genera have niche distances close to

***C. difficile* relative to unrelated genera.** Niche distances of strains and genera,

represented as the euclidean distance between flux vectors, relative to *C. difficile*

across CDI-FMT cohort samples for which the *C. difficile* growth rate >10^{-6}. Genera and

strains are ordered by the median niche distance. Probiotic strains and associated

genera are colored, consistent with the legend in Fig. 5D. *C. bolteae* and *C. innocuum*

are both members of the genus *Clostridium*.