Ca$_V$2.1 α$_1$ subunit motifs that control presynaptic Ca$_V$2.1 subtype abundance are distinct from Ca$_V$2.1 preference

Jianing Li[1,2], Priyadharishini Veeraraghavan[1], Samuel M. Young, Jr[1,3^]

[1]Department of Anatomy and Cell Biology, University of Iowa, Iowa City, IA, USA.
[2]Cell Developmental Biology Graduate Program, University of Iowa, Iowa City, IA 52242, USA
[3]Department of Otolaryngology, Iowa Neuroscience Institute, University of Iowa, Iowa City, IA, USA.

^Correspondence
Samuel M. Young, Jr. PhD
Department of Anatomy and Cell Biology
University of Iowa
PBDB 5322
169 Newton Road
Iowa City, IA 52242
Email: samuel-m-young@uiowa.edu

ORCIDs:

Jianing Li	0000-0001-5384-773X
Priyadharishini Veeraraghavan	0000-0003-2826-1686
Samuel M. Young, Jr.	0000-0002-7589-7612

26 **Key points summary:**

- Presynaptic Ca_V2 subtype abundance regulates neuronal circuit properties, however the mechanisms regulating presynaptic Ca_V2 subtype abundance and preference remains enigmatic.
- The Ca_V α_1 subunit determines subtype and contains multiple motifs implicated in regulating presynaptic subtype abundance and preference.
- The $Ca_V2.1$ α_1 subunit domain II-III loop and cytoplasmic C-terminus are positive regulators of presynaptic $Ca_V2.1$ abundance but do not regulate preference.
- The $Ca_V2.3$ α_1 subunit cytoplasmic C-terminus negatively regulates presynaptic Ca_V2 subtype abundance but not preference while the $Ca_V2.2$ α_1 subunit cytoplasmic C-terminus is not a key regulator of presynaptic Ca_V2 subtype abundance or preference.
- The Ca_V2 α_1 subunit motifs determining the presynaptic Ca_V2 preference are distinct from abundance.

Abstract

Presynaptic voltage-gated Ca^{2+} channels (Ca_V) subtype abundance at mammalian synapses regulates synaptic transmission in health and disease. In the mammalian central nervous system, most presynaptic terminals are $Ca_V2.1$ dominant with a developmental reduction in $Ca_V2.2$ and $Ca_V2.3$ levels, and Ca_V2 subtype levels are altered in various diseases. However, the molecular mechanisms controlling presynaptic Ca_V2 subtype levels are largely unsolved. Since the Ca_V2 α_1 subunit cytoplasmic regions contain varying levels of sequence conservation, these regions are proposed to control presynaptic Ca_V2 subtype preference and abundance. To investigate the potential role of these regions, we expressed chimeric $Ca_V2.1$ α_1 subunits containing swapped motifs with the $Ca_V2.2$ and $Ca_V2.3$ α_1 subunit on a $Ca_V2.1/Ca_V2.2$ null background at the calyx of Held presynaptic terminal. We found that expression of $Ca_V2.1$ α_1 subunit chimeras containing the $Ca_V2.3$ loop II-III region or cytoplasmic C-terminus (CT) resulted in a large reduction of presynaptic Ca^{2+} currents compared to the $Ca_V2.1$ α_1 subunit. However, the Ca^{2+} current sensitivity to the $Ca_V2.1$ blocker Agatoxin-IVA, was the same between the chimeras and the $Ca_V2.1$ α_1 subunit. Additionally, we found no reduction in presynaptic Ca^{2+} currents with $Ca_V2.1/2.2$ cytoplasmic CT chimeras. We conclude that the motifs in the $Ca_V2.1$ loop II-III and CT do not individually regulate $Ca_V2.1$ preference, but these motifs control $Ca_V2.1$ levels and the $Ca_V2.3$ CT contains motifs that negatively regulate presynaptic $Ca_V2.3$ levels. We propose that the motifs controlling presynaptic $Ca_V2.1$ preference are distinct from those regulating $Ca_V2.1$ levels and may act synergistically to impact pathways regulating $Ca_V2.1$ preference and abundance.

Introduction

At mammalian central nervous system (CNS) synapses, $Ca_V2.1$, $Ca_V2.2$, and $Ca_V2.3$, are key determinants of Ca^{2+} entry that controls synaptic transmission and neuronal circuit function (Dolphin & Lee, 2020; Young & Veeraraghavan, 2021). Among the three Ca_V2 subtypes, presynaptic $Ca_V2.1$ levels are dominant compared to $Ca_V2.2$ and $Ca_V2.3$ (Iwasaki & Takahashi, 1998; Iwasaki *et al.*, 2000; Nanou & Catterall, 2018; Dolphin & Lee, 2020). During neuronal circuit maturation, action-potential (AP) evoked synaptic transmission becomes increasingly $Ca_V2.1$-dependent due to a selective presynaptic reduction of $Ca_V2.2$ and $Ca_V2.3$ (Iwasaki & Takahashi, 1998; Eggermann *et al.*, 2011; Young & Veeraraghavan, 2021). In addition, presynaptic Ca_V2 subtype levels vary between synapses and are altered in various diseases (Dolphin & Lee, 2020; Young & Veeraraghavan, 2021). Since Ca_V2 subtypes differentially control Ca^{2+} influx, elucidating the molecular mechanisms controlling presynaptic Ca_V2 subtype abundance are critical to understanding brain function in health and disease.

Although presynaptic Ca_V2 subtype levels are highly variable, they are distinct between the neuronal soma and presynaptic terminal (Doughty *et al.*, 1998). In addition, loss of $Ca_V2.1$ leads to increases in presynaptic $Ca_V2.2$ levels, however total presynaptic Ca^{2+} currents are reduced (Jun *et al.*, 1999; Inchauspe *et al.*, 2004; Lubbert *et al.*, 2017). Therefore, it is proposed that mechanisms that control presynaptic Ca_V2 levels and preferences are distinct from those controlling somatic Ca_V2 levels (Weiss & Zamponi, 2017; Dolphin & Lee, 2020). However, whether the mechanisms that determine presynaptic Ca_V2 subtype preference and abundance are interrelated or separable are unknown.

The pore- forming Ca_V2 α_1 subunit determines presynaptic Ca_V2 preference and contains multiple cytoplasmic protein interaction motifs (Dolphin & Lee, 2020). Due to varying levels of sequence conservation between the three Ca_V2 α_1 subunit isoforms cytoplasmic regions, three regions are implicated in regulating presynaptic Ca_V2 channel levels. They are: 1) the I-II loop; the primary binding site to the $Ca_V\beta$ subunit (AID) 2) the II-III loop; the synprint domain; and 3)

the C-terminal region, which contains multiple active zone (AZ) protein binding sites and is

frequently mutated in Ca_V2 channelopathies (Dolphin & Lee, 2020; Young & Veeraraghavan,

2021). Since the loop II-III and cytoplasmic CT regions are not highly conserved between the

different Ca_V2 α_1 subunit isoforms and are subject to alternative splicing (Simms, 2014), they are

speculated to contain motifs that regulate the presynaptic Ca_V2 subtypes levels and preference.

However, the identity of the Ca_V2 α_1 subunit motifs that regulate presynaptic Ca_V2 subtype levels

and preference is unclear. Studies from cultured superior cervical ganglion neurons demonstrated

that the $Ca_V2.1$ loop II-III domain regulates presynaptic targeting (Mochida *et al.*, 2003). However,

it is not defined if the $Ca_V2.1$ loop II-III region is a key determinant of preference or if the II-III

region from other Ca_V2 subtypes regulates abundance and preference. Although motifs in the

$Ca_V2.1$ CT (aa. 2016-2368) that interact with AZ proteins are dispensable for controlling

presynaptic $Ca_V2.1$ abundance (Lubbert *et al.*, 2017), it is not known if other regions in the $Ca_V2.1$

CT control Ca_V2 subtype abundance and preference. Furthermore, whether the $Ca_V2.2$ and

$Ca_V2.3$ CT regulates abundance and preference is unresolved.

To elucidate the roles of the Ca_V2 α_1 subunit loop II-III and CT in regulating presynaptic

abundance and preference in a native neuronal circuit, we generated Ca_V2 α_1 subunit chimeras:

1) Ca$_V$2.1/2.3 chimeras, the $Ca_V2.1$ α_1 subunit II-III loop and/or CT region, were swapped with

these regions from the $Ca_V2.3$ α_1 subunit. 2) Ca$_V$2.1/2.2 chimeras, $Ca_V2.1$ α_1 subunit CT region

was swapped with the $Ca_V2.2$ α_1 subunit CT region. We then used helper-dependent Adenoviral

(HdAd) technology and delivery techniques to express $Ca_V2.1$ α_1 subunit chimeras on a

$Ca_V2.1$/$Ca_V2.2$ null background at the postnatal days (P) 9–11 calyx of Held using a

Cacna1a/Cacna1b flox/flox ($Ca_V2.1$ $Ca_V2.2$ CKO) mouse model. The calyx of Held is a large

glutamatergic presynaptic terminal in the auditory brainstem that at this developmental stage

contains multiple Ca_V2 subtypes which is similar to the majority of presynaptic terminals (Iwasaki

et al., 2000; Young & Veeraraghavan, 2021).

Using direct presynaptic Ca^{2+} current recordings at the calyx of Held, we found that

ablation of $Ca_V2.1$ and $Ca_V2.2$ resulted in a significant reduction in total presynaptic Ca^{2+} currents

and were insensitive ω-Agatoxin-IVA (Aga), a $Ca_V2.1$ specific blocker, and ω-Conotoxin GVIA

(Cono), a $Ca_V2.2$ specific blocker, and Nifedipine a Ca_V1 specific blocker, but were partially

sensitive to SNX-482, a $Ca_V2.3$ specific blocker. Expression of $Ca_V2.1/2.3$ chimeras with single

or combined region swaps resulted in a significant reduction in peak Ca^{2+} current amplitudes.

However, the remaining Ca^{2+} currents with the single swap $Ca_V2.1/2.3$ chimeras exhibited a

similar proportion of Aga sensitive currents as the full transcript (FT) $Ca_V2.1$ α_1 subunit, while the

combined swap had a significant increase in Aga insensitive Ca^{2+} currents. In addition, expression

of $Ca_V2.1/2.3$ chimeras which replaced the dispensable region of the $Ca_V2.1$ CT (aa. 2016-2368)

with corresponding $Ca_V2.3$ CT region resulted in a reduction in Ca^{2+} currents. Finally, we found

that expression of $Ca_V2.1/2.2$ CT chimeras revealed no reduction in presynaptic Ca^{2+} currents.

Based on our data, we conclude that individual motifs in the $Ca_V2.1$ α_1 subunit loop II-III

or CT are positive regulators of presynaptic $Ca_V2.1$ levels but do not individually control $Ca_V2.1$

preference. Furthermore, we conclude the $Ca_V2.3$ CT contains motifs that negatively regulate

presynaptic $Ca_V2.3$ levels. Therefore, we propose that the motifs that determine presynaptic

$Ca_V2.1$ abundance are distinct from those regulating $Ca_V2.1$ preference and these motifs may

synergistically impact pathways that regulate $Ca_V2.1$ preference and levels.

Methods and materials

Ethical approval

All procedures were performed according to the animal welfare law of the University of Iowa Office of Animal Resources (OAR) and Institutional Animals Care and Use Committee (IACUC, D16-00009 (A3021-01)). Mice are maintained at The University of Iowa Animal Care Facility which is approved under NIH assurance A3021-01 and certified by the State of Iowa for use of living animals. Animals were housed with 12 h light/dark cycle and *ab libitum* food/water supply. Animals of both sexes at postnatal day 9-11 were used for all experiments. All available measures were taken to minimize animal pain and suffering.

EXPERIMENTAL DESIGN

Generation of *Cacna1a/Cacna1b* flox/flox animal

Cacna1a fl/fl mice

Cacna1a$^{fl/fl}$ mice were generated as follows. The Cacna1a$^{tm1a(EUCOMM)Hmgu}$ targeting vector (EUMMCR) was electroporated into the JM8A3 ES-cells derived from C57BL/6N mice and clones were selected and PCR verified. The neo gene was removed by crossing to FLPo mice carrying the FLP recombinase gene under the control of the CAG promoter (C57BL/6N-Tg (CAG-Flpo)1Afst/Mmucd, stock 036512-UCD. (Mouse Biology Program KOMP). Genotyping of *Cacna1a* flox mice by PCR was carried out by the following primers,

754 5' GGAATCCCAAGTGCGTGAGTCTCC 3'

887 5' TCAGCCCCTTCTCTTTAATCTGACCC 3'

Cacna1a$^{fl/fl}$ mice were backcrossed to C57BL6/J background.

Cacna1b$^{fl/fl}$ mice

Cacna1b$^{fl/fl}$ mice (C57BL/6N-Cacna1b$^{tm1c(KOMP)Wtsi}$/H) were purchased from MRC Harwell Institute. Genotyping of *Cacna1b$^{fl/fl}$* was carried out using the following primers,

Cacna1b-5arm-WTF 5' TTGGTGTCCTGTTCTCCACA 3'

Cacna1b-Crit-WTR 5' CAGGGAATCCTGGGAAGTAA 3'

5mut-R1   5' GAACTTCGGAATAGGAACTTCG 3'

Cacna1b$^{fl/fl}$ mice were backcrossed to the C57BL6/J background.

Cacna1a$^{fl/fl}$, Cacna1b$^{fl/fl}$ mice

Cacna1a$^{fl/fl}$, Cacna1b$^{fl/fl}$ mice were generated by crossbreeding *Cacna1a$^{fl/fl}$* with *Cacna1b$^{fl/fl}$* to

homozygosity. Genotyping was performed by PCR.

Stereotactic P1 surgery

Stereotactic surgery was performed at P1 as described previously (Radulovic *et al.*, 2020). Briefly,

Cacna1a$^{fl/fl}$, *Cacna1b$^{fl/fl}$* mice were anesthetized by hypothermia, using ice bath for 5min.

Subsequently, ~2 μl HdAd in storage buffer (in Mm: 10 HEPES, 250 sucrose, 1 MgCl$_2$ at pH: 7.4

and 6.6% mannitol) was injected at 1 μl/min into the cochlear nucleus (CN) using pulled glass

pipettes with a 20 μm opening (Sutter Instrument, P-97, U.S.A.). Two viral vectors, expressing

Cre and Ca$_V$2 α$_1$ hybrid constructs were co-injected. The total viral particle delivered did not

exceed 2*10^9 viral particles. To dissipate pressure after injection, the needle was slowly removed.

Animals were then placed under a warming lamp. Upon full recovery, visual inspection and tail

pinch responsiveness, animals were returned to their respective cage with the dam. If animals did

not recover from the surgery, they were immediately euthanized. In the case of runting or if

animals were found in distress at later time points the veterinary staff was consulted and their

recommendation for further action was followed.

DNA constructs and recombinant viral vector production

Ca$_V$2.1 chimeras

Codon optimized (GENEart) Ca$_V$2.1 α$_1$ subunit cDNA Cacna1a (*Mus musculus*, Accession No.

NP_031604.3), Ca$_V$2.2 α$_1$ subunit cDNA Cacna1b (*Mus musculus,* Accession No.

NP_001035993.1) and Ca$_V$2.3 α$_1$ subunit cDNA Cacna1e (*Mus musculus*, Accession No.

NP_033912.2) were used to generate Ca$_V$2 α$_1$ hybrid constructs. All Ca$_V$2 hybrid constructs were

sequenced verified. Amino acid alignment of above constructs were were performed under Clustal

Omega function using the MegAlign Pro 17 program implemented in the Lasergene software

package (v.17, DNASTAR).The following Ca_V2 hybrid constructs were created.

<u>$Ca_V2.1/2.3$ II-III loop</u>: $Ca_V2.1$ α_1 subunit aa. 719-1210 replaced with $Ca_V2.3$ α_1 subunit aa. 707-

1151

<u>$Ca_V2.1/2.3$ CT</u>: $Ca_V2.1$ α_1 subunit aa. 1933-2368 replaced with $Ca_V2.3$ α_1 subunit aa. 1897-2273

<u>$Ca_V2.1/2.3$ II-III loop CT</u>: $Ca_V2.1$ α_1 subunit aa. 719-1210 replaced with $Ca_V2.3$ α_1 subunit aa. 707-

1151 and $Ca_V2.1$ α_1 subunit aa. 1933-2368 replaced with $Ca_V2.3$ α_1 subunit aa. 1897-2273

<u>$Ca_V2.1/2.2$ CT</u>: $Ca_V2.1$ α_1 subunit aa. 1933-2368 replaced by $Ca_V2.2$ α_1 subunit aa. 1870-2327

<u>$Ca_V2.1/2.3$ CT1</u>: $Ca_V2.1$ α_1 subunit aa. 2016-2368 replaced with $Ca_V2.3$ α_1 subunit 1952-2273

<u>$Ca_V2.1/2.3$ CT2</u>: $Ca_V2.1$ α_1 subunit aa. 1967-2368 replaced by $Ca_V2.3$ α_1 subunit 1929-2273

These constructs were individually cloned into the EcoRI and NotI sites of the 470 bp human

synapsin (hSyn) neurospecific expression cassette (Montesinos *et al.*, 2016; Dong *et al.*, 2018).

HdAd production

Helper-dependent Adenoviral vector (HdAd) (Cre) which co-expresses Cre recombinase (Cre and

EGFP or mCherry under separate neurospecific expression cassettes driven by the 470 bp

human synapsin promoter (hSyn) was produced as previously described (Montesinos *et al.*, 2016;

Dong *et al.*, 2018). To produce HdAd Ca_V2 chimeric viruses we followed standard protocols

(Montesinos *et al.*, 2016; Lubbert *et al.*, 2017). Briefly, Ca_V2 α_1 chimera cDNA constructs driven

by hSyn promoter expression casseete or a high-level overexpression pUNISHER (Pun) cassette

were then cloned into he AscI site, pdelta23E4 which contains a separate neuro-specific

expression of mCherry or myristolated EGFP (mEGFP) markers (Montesinos *et al.*, 2011).

Production of HdAd was carried out as previously described (Palmer & Ng, 2005; Montesinos *et*

al., 2016). The HdAd plasmid was linearized with PmeI and then transfected (Profection

Mammalian Transfection System, Promega, Madison, WI, USA) into 116 producer cells. Helper

virus (HV) was added the following day. Forty-eight hours post infection, cells were subjected to

three freeze/thaw cycles and lysate was collected. The HdAd vectors were amplified for total of

five serial coinfections from 3x6 cm tissue culture dishes followed by one 15 cm dish and finally

30x15 cm dishes of 116 cells (confluence ~90%). HdAd was purified by CsCl ultracentrifugation.

HdAd was stored at -80°C in storage buffer (10 mM Hepes, 1 mM MgCl$_2$, 250 mM Sucrose, pH

7.4).

HdAd titering and quality control

HdAds were titered by qPCR as described previously (Lubbert *et al.*, 2019). The purified reference

virus (RV); Ad5-CMV-EGFP (UNC, Gene Therapy Center, Chapel Hill, North Carolina) was titered

by endpoint dilution and by transducing units, as described previously (Bewig & Schmidt, 2000).

Human Adenovirus 5 Reference Material (ARM), (ATCC, VR-1516). To titer the HdAd by qPCR,

adenoviral gDNA from HdAd, RV and ARM was purified from 6 cm plates. Plates were seeded

with 3.75 x 10^6 HEK293 cells in DMEM with 10% FBS and coinfected with HdAd plus RV on the

following day at a multiplicity of 100 vp/cell. The cells were incubated for 1 hr at 37°C with 5%

CO$_2$ and washed to remove unabsorbed virions. Cells were harvested 4 hr post-infection, then

washed with 1 mL PBS and centrifuged for 5 min at 4°C. Pellets were resuspended in 100 mM

Tris-HCl, pH 8, lysed by freeze/thaw cycles and treated with 20 U/mL Benzonase for 10 min.

Subsequently, lysate was incubated in Proteinase K-SDS solution for 3 hr at 50°C. After protein

precipitation with 1.65 M NaCl on ice, DNA was precipitated from supernatants with isopropanol

and 3 M sodium acetate, pH 5.2. Samples were mixed well and incubated for 1 hr at -80°C then

spun at 4°C for 30 min. Pellets were washed with cold ethanol, dried and resuspended in 10 mM

Tris-Cl, pH 8.0. A plasmid standard containing three specific DNA sequences; Ad5 E1a

(replication-competent Adenovirus (RCA) detection), Ad5 E2B Iva2 (viral DNA polymerase and

terminal protein precursor, HV-exclusive), and Ad genomic stuffer sequence (Cosmid 346 and

HPRT1 gene, in all HdAd-exclusive) was used. Three specific primer/probe sets were used for

these 3 regions in the plasmid for the detection in qPCR:

1.) **Ad5 E1a (RCA):** probe: NED-5'-AGCACCCCGGGCACGGTTG-3'-MGBNFQ; fwd: 5'-

GGGTGAGGAGTTTGTGTTAGATTATG-3'; rev: 5'-TCCTCCGGTGATAATGACAAGA-3'

2.) **E2B Iva2 (HV):** probe: FAM-5'-TGTCTTTCAGTAGCAAGCT-3'-TAMRA, HV fwd primer:

5'-TGGGCGTGGTGCCTAAAA-3', rev: 5'-GCCTGCCCCTGGCAAT-3'

3.) **Genomic stuffer DNA (HdAd):** probe: VIC-5'- AGCCTCTCTCATCTCACAGT -3'-

MGBNFQ,   fwd: 5'-CCCCGCTACCCCAATCC-3',   rev: 5'-

TTAGCTTTTTTGGGTGATTTTTCC-3'.

Sample DNA was diluted $1:10^3$, $1:10^4$ or 10^5 in 10 mM Tris-Cl, pH 8.0. For HdAd and HV probes,

a 6-point standard curve was constructed with the plasmid standard using 10^7- 10^2

molecules/well. For RCA detection, two 7-point curves were constructed with 10^7 - 10

molecules/well (one plasmid standard and one using ARM gDNA spiked with 10^7 copies of HdAd

gDNA). Real-Time PCR parameters were 50°C for 2 min, 95°C for 10 min; 40 cycles of: 95°C for

15 sec, 60°C for 1 min. Outliers and results out of range of standard curve were discarded. Results

within range of the standard curve were corrected for their dilution and determined as HdAd vector

genomes/mL. All samples were negative for RCA.

To determine the IU copies/mL from purified coinfection samples, averaged results were

divided by 3 μL and multiplied by 25 μL giving IU copies/coinfection. Next, this number was divided

by the volume in μL of HdAd added to the coinfection giving IU copies/μL then multiplied by 1000

to obtain the i,nfectious HdAd genomes/mL.

Calculation to determine the titer of purified HdAd virus:

$$HdAd\ infectivity = RV\ infectivity * \frac{HdAd\ vector\ genomes/RV\ genomes}{infectious\ HdAd\ genomes/infectious\ RV\ genomes}$$

where Ad5-CMV-EGFP is RV, RV infectivity is VP/IU. IU is the average titer of end point dilution

and transducing unit assays, HdAd vector genomes and RV genomes are the number of genomes

from purified virus as determined by qPCR, and infectious HdAd genomes and infectious RV

genomes are the number of genomes in DNA purified from coinfections. The titers of HdAds carry

different cDNA constructs are listed as follow, viral particle/ml: Cre EGFP, $1.75*10^{12}$, Cre

mCherry, $2.41*10^{12}$, syn Ca$_V$2.1 FT, $2.40*10^{12}$, Pun Ca$_V$2.1 FT, $2.19*10^{12}$, Ca$_V$2.1/2.3 II-III loop

CT, $2.00*10^{12}$, $Ca_V2.1/2.3$ II-III loop, $2.07*10^{12}$, Syn $Ca_V2.1/2.3$ CT, $1.90*10^{12}$, Pun $Ca_V2.1/2.3$ CT,

$3.11*10^{12}$, $Ca_V2.1$ delta 2016, $1.40*10^{12}$, $Ca_V2.1/2.3$ CT 2016-2368, $4.70*10^{12}$, $Ca_V2.1/2.3$ CT

1967-2368, $3.29*10^{12}$, syn $Ca_V2.1/2.2$ CT, $3.17*10^{12}$, Pun $Ca_V2.1/2.2$ CT, $9.52*10^{12}$. Particle to

infectious unit ratio was 10:1~ 50:1. Either syn Cre syn EGFP or syn Cre syn mCherry was used

to make sure that the viruses expressing Cre carry different fluorescent protein than the viruses

expressing Ca_V2 constructs.

Western Blot

265   To determine expression of our $Ca_V2.1$ chimeric constructs, we performed Western Blot

analysis on HEK293 cells transduced with our HdAd vectors expressing the Ca_V2 α_1 chimeric

constructs. To do so, HEK293 cells were infected with our HdAd vectors, and 48 hrs later cells

were collected and lysed in RIPA buffer containing: 10 mM Tris-Cl pH8.0, 1mM EDTA, 1% Triton-

X, 0.1% sodium deoxycholate, 0.1% SDS, 140 mM NaCl, protease inhibitor (11836153001,

Roche). 20 µg of protein lysate were loaded and analyzed by SDS-PAGE followed by

immunoblotting: $Ca_V2.1$ α_1 CT antibody (Synaptic Systems, #152 203, rabbit polyclonal) diluted

1:1000 (final concentration 2 µg/ml) and $Ca_V2.1$ α_1 II-III loop antibody (Alomone Labs, ACC-001,

rabbit polyclonal) diluted 1:500 (final concentration 1.7 µg/ml) were incubated overnight at 4°C to

blot the expression of $Ca_V2.1/2.3$ II-III loop hybrid construct, $Ca_V2.3$ α_1 CT antibody (Synaptic

Systems, #152 411, mouse monoclonal) diluted 1:500 (final concentration 2 µg/ml) was incubated

overnight at 4°C to blot the expression of $Ca_V2.1/2.3$ II-III loop CT and $Ca_V2.1/2.3$ CT chimeras,

$Ca_V2.2$ CT antibody (Synaptic Systems, #152 311, mouse monoclonal) diluted 1:1000 (final

concentration 2 µg/ml) was incubated overnight at 4°C to detect the expression of $Ca_V2.1/2.2$ CT

chimera. β-actin antibody (sigma, A2228, mouse monoclonal) diluted 1:5000 (final concentration

0.4 µg /ml) was incubated overnight at 4°C used to blot the expression of β-actin. Goat anti mouse

IgG (H+L) poly-HRP (Thermo Fisher, #32230) diluted 1:2000 (final concentration 0.25 µg /ml) and

goat anti rabbit IgG (H+L) poly-HRP (Thermo Fisher, #32260) secondary antibody diluted 1:2000

(final concentration 0.25 ng/ml) was incubated at room temperature for 1 hour. Protein bands

were detected by chemiluminescence using ECL Western blot detection reagents (Bio-Rad, #170-

5060).

Preparation of acute slices

Acute brain slices were prepared as previously described (Chen *et al.*, 2013). Briefly, after

decapitation of P9-11 mice of either sex, the brains were immersed in ice-cold low Ca^{2+} artificial

cerebrospinal fluid (aCSF) containing (in mM): 125 NaCl, 2.5 KCl, 3 $MgCl_2$, 0.1 $CaCl_2$, 10 glucose,

25 $NaHCO_3$, 1.25 Na_2HPO_4, 0.4 L-ascorbic acid, 3 myo-inositol, and 2 Na-pyruvate, pH 7.3-7.4

(310 mosmol/l). aCSF solution was continuously bubbled with 95% O_2 and 5% CO_2. Acute brain

slices (200 μm thickness) containing MNTB were obtained using a vibrating tissue slicer

(Campden 7000 smz-2, Campden Instruments LTD, Loughborough, England). Slices were

immediately transferred to standard aCSF (37°C, continuously bubbled with 95% O_2-5% CO_2)

containing the same chemicals as the slicing buffer except that it contained 1mM $MgCl_2$ and 1 mM

$CaCl_2$. After 1 hour incubation for recovery, slices were transferred to a recording chamber with

the same aCSF at room temperature (RT: 25°C).

Acute slice electrophysiology

In all experiments, slices were continuously perfused with aCSF and visualized by upright

microscope (BX51W, Olympus) through 4X air or 60X water-immersion objective (LUMPlanFL N,

Olympus, Tokyo, Japan) and a CCD camera (QI-Click, QImaging, Surrey, BC, Canada). Patch

clamp recordings were performed by using an EPC 10/2 patch-clamp amplifier (HEKA,

Lambrecht, Germany), operated by PatchMaster version 2x90.5 (Harvard Instruments, Holliston,

MA, USA). Data were low-pass filtered at 6 kHz and sampled with a rate of 50 kHz. Calyces

transduced with HdAd $Ca_V2.1$ α_1 chimeric constructs were identified visually with two co-

expressed mEGFP and mCherry markers. To visualize mEGFP and mCherry with light of 470 nm

or 560 nm, respectively, a Polychrome V xenon bulb monochromator or a lumen Metal arc lamp

was used (TILL Photonics, Grafelfing, Germany).

Presynaptic Ca^{2+} current recordings

To isolate presynaptic Ca^{2+} currents, aCSF was supplemented with 1 or 1.5 µM tetrodotoxin (TTX,

Alomone labs), 100 µM 4-aminopyridin (4-AP, Tocris) and 20 mM tetraethylammonium chloride

(TEA, Sigma Aldrich) to block Na^+ and K^+ channels. Calyces were whole-cell voltage clamped at

-80 mV. Current-voltage (IV) relationships were recorded in the presence of 1 mM $CaCl_2$ and 1

mM $MgCl_2$ with 1.5 µM TTX while pharmacological isolation of Ca_V2 subtype components was

performed in 2 mM $CaCl_2$ and 1 mM $MgCl_2$ with 1 µM TTX. To characterize the in the presynaptic

terminal, we used 200 nM ω-agatoxin IVA (Alomone labs) for selective inhibition of $Ca_V2.1$

channels, 2 µM ω-conotoxin GIVA (Alomone labs) for selective inhibition of $Ca_V2.2$ channels, 200

nM SNX-482 (Alomone labs) for selective inhibition of $Ca_V2.3$ channels, 10 µM Nifedipine

(Alomone labs) for selective inhibition of Ca_V1 chnannels and supplemented 0.1 mg/mL

cytochrome C to prevent toxin absorption. 50µM $CdCl_2$ was used for non-selective inhibition of

all VGCCs. Presynaptic patch pipette with open tip diameters 4-6 MΩ resistance were pulled from

1.5 mm thin-walled borosilicate glass (Harvard apparatus) and were filled with following internal

solution (in mM): 145 Cs-gluconate, 20 TEA-Cl, 10 HEPES, 2 Na_2-phosphocreatine, 4 MgATP,

0.3 NaGTP, and 0.075 EGTA, pH 7.2, 325-340 mOsm. Pipettes were coated with dental wax to

minimize stray capacitance and improve voltage clamp quality. Presynaptic series resistance was

less than 15 MΩ and was compensated online to 3 MΩ or less for IV recordings or compensated

to 6 MΩ for Ca_V2 subtype characterization experiments with a time lag of 10 µs by the HEKA

amplifier. Leak and capacitive currents were subtracted online with a P/5 routine. Cells with series

resistance >15 MΩ and leak currents > 100 pA were excluded from the analysis.

Analysis of electrophysiological data

All data was analyzed by NeuroMatic (Rothman & Silver, 2018) in Igor Pro (version 8.0,

Wavemetrics, Portland, OR, USA). Voltage dependence of channel activation was described by

both peak and tail currents as functions of voltage. Peak currents were fitted according to a

Hodgkin-Huxley formalism with four independent gates assuming a Goldman-Hodgkin-Katz

(GHK) open-channel conductance T: with E_{rev} as reversal potential, V_m as half-maximal activation

voltage per gate, and k_m as the voltage dependence of activation. Tail currents were measured

as peaks minus baseline and fitted with a Boltzmann function:

$$I(V) \ = \ \text{T} \frac{1 - e^{-\frac{V - E_{rev}}{25mV}}}{1 - e^{-\frac{V}{25mV}}} \left(1 - e^{\frac{V - Vm}{Km}}\right)^{-4}$$

where $V_{1/2}$ represents the half-maximal voltage and k the corresponding slope factor.

Statistical analysis

All statistical tests were conducted in Prism 6 (GraphPad Software). Sample sizes for all

experiments were chosen based on assuming a population with a normal distribution sample size

of seven is sufficient to invoke the Central Limit Theorem. All data were tested for normal

distribution by performing a Shapiro-Wilk test for normality and variances of all data were

estimated and compared using Bartlett's test. Electrophysiological data were compared with one-

way analysis of variance (ANOVA) with a post hoc Dunnett's test, $Ca_V2.1$ FT dataset was used

as control group. In Figures and Table, data are reported as mean ± SD. Statistical significance

was accepted at *$p<0.05$; **$p<0.01$; ***$p<0.001$; ****$p<0.0001$.

Results

Ca$_V$2.1 and Ca$_V$2.2 channels can be ablated at the calyx of Held in Ca$_V$2.1/Ca$_V$2.2 CKO mouse model.

Since ablation of Ca$_V$2.1 channels results in partial compensation of Ca$_V$2 current levels by Ca$_V$2.2 channels at the calyx of Held (Inchauspe *et al.*, 2004; Lubbert *et al.*, 2017), we created a Ca$_V$2.1/ Ca$_V$2.2 CKO mouse model (Fig. 1A) to allow for spatial and temporal ablation of Ca$_V$2.1 and Ca$_V$2.2 channels. To determine how presynaptic loss of Ca$_V$2.1 and Ca$_V$2.2 channels impacted presynaptic Ca$_V$2 currents at the calyx of Held, we injected HdAd Cre at P1 into the cochlear nucleus (CN) of Ca$_V$2.1/ Ca$_V$2.2 CKO mice to create a Ca$_V$2.1$^{-/-}$/ Ca$_V$2.2$^{-/-}$ calyx of Held. Subsequently we performed whole-cell patch clamp recordings at the P9-11 calyx of Held and characterized the total Ca^{2+} current and current sensitivity to Ca$_V$2 subtype-specific blockers: ω-Agatoxin IVA (Aga), Ca$_V$2.1-specific, ω-Conotoxin GIVA (Cono), Ca$_V$2.2-specific, and Cadmium (Cd^{2+}) a non-specific blocker of Ca$_V$ currents (Fig. 1D-F, Table 1). Since sequential application of blockers takes ~10-15 minutes for complete characterization of Ca$_V$2 currents, we measured presynaptic Ca^{2+} current rundown at wild-type calyces in the absence of blockers after ~10-15 minutes (Fig. 1B, C). We found a ~20-70 pA (2-5%) reduction of presynaptic Ca^{2+} currents (Fig. 1J, Table 1, n=3). Subsequently, analysis of Ca^{2+} currents at the Ca$_V$2.1$^{-/-}$/ Ca$_V$2.2$^{-/-}$ calyx revealed a significant reduction (wt vs. CKO (pA): -1078 ± 260.9 (n=10) vs. -425.5 ± 219.5 (n=11), p<0.0001) (Fig. 1E, Table 2). We found an almost complete loss of Aga sensitive currents, 809 ± 255.6 (n=4) vs 99.22 ± 53.64 (n=5), p=0.0005; and Cono sensitive currents, 92 ± 142.2 (n=4) vs 50.8 ± 43.1 (n=5), p=0.5532). (Fig1. D and F, Table 1). Therefore, these results demonstrate successful ablation of Ca$_V$2.1 and Ca$_V$2.2 channels.

In the absence of Ca$_V$2.1 and Ca$_V$2.2, a significant Aga and Cono insensitive presynaptic Ca^{2+} current was present. Since Ca$_V$2.3 subtypes can compensate for the loss of Ca$_V$2.1 channels (Kaja *et al.*, 2006) and Ca$_V$1 channels can be found in some presynaptic terminals (Giugovaz-

Tropper *et al.*, 2011; Young & Veeraraghavan, 2021), it was important to determine if the

combined loss of $Ca_V2.1$ and $Ca_V2.2$ resulted in compensation by $Ca_V2.3$ or Ca_V1 channels.

Therefore, we characterized the sensitivity of presynaptic Ca^{2+} current to $Ca_V2.3$ and Ca_V1

specific blockers in the $Ca_V2.1^{-/-}$/ $Ca_V2.2^{-/-}$ calyx of Held. First, we used SNX-482, a $Ca_V2.3$

selective blocker and found a range of ~10% to 35% block of presynaptic Ca^{2+} currents (SNX-

482 sensitive fraction 24.33 ± 9.074 %) (n=3) (Fig. 1G, H, Table 1). Since there were variable

levels of Ca^{2+} current reduction with SNX-482, this indicated the remaining Ca^{2+} currents in the

absence of $Ca_V2.1$ and $Ca_V2.2$ could be mediated by a combination of $Ca_V2.3$ variants that are

SNX-482 sensitive and SNX-482 insensitive and/or presynaptic Ca_V1 channels. To determine if

Ca_V1 channels were present, we used nifedipine, a Ca_V1 selective blocker (Fig. 1I, J) and found

presynaptic Ca^{2+} currents were nifedipine insensitive as there was no difference compared to the

reduced fraction of Ca^{2+} current due to rundown (Fig. 1I, J, Table 1). Therefore, we conclude that

in the absence of $Ca_V2.1$ and $Ca_V2.2$ channels the compensatory presynaptic Ca^{2+} currents are

mediated by mix of SNX-482 sensitive and insensitive $Ca_V2.3$ channels.

The $Ca_V2.1$ α_1 subunit loop II-III domain, and CT are positive regulators of presynaptic

$Ca_V2.1$ abundance

The amino acid conservation of $Ca_V2.1$ α_1 subunit and $Ca_V2.3$ α_1 subunit loop II-III region

is 33.2% and CT region from aa. 1933 onwards is 39.8%, while the loop I-II is 87.3% conserved

(Figure 2A, MegAlign Pro 17, Clustal Omega). In fact, the $Ca_V2.3$ α_1 subunit loop II-III lacks a

syntaxin I binding site in the synprint region, while the $Ca_V2.3$ α_1 subunit CT lacks many of

proposed AZ protein binding motifs found in the $Ca_V2.1$ CT (Dolphin & Lee, 2020). Since the loop

II-III and CT regions are the most divergent, we created $Ca_V2.1$/$Ca_V2.3$ α_1 subunit chimeras that

swapped the loop II-III and CT domains to determine if these domains regulated presynaptic

$Ca_V2.1$ abundance and preference (Fig. 2B). We generated three HdAd $Ca_V2.1$/2.3 α_1 subunit

chimeric constructs (Figure 2C). There were Ca$_V$2.1/2.3 II-III loop, which contains the $Ca_V2.3$ α_1

subunit loop II-III domain (aa. 707 to aa. 1151), Ca$_V$2.1/2.3 CT, which contains the cytoplasmic

$Ca_V2.3$ CT (aa. 1897 to aa. 2273), and Ca$_V$2.1/2.3 II-III+CT which contains both the $Ca_V2.3$ α_1

subunit loop II-III and CT (Fig. 3C). To validate that the chimeras were expressed as full-length

proteins, we performed Western Blot analysis of HEK293 cells infected with our HdAd $Ca_V2.1/2.3$

chimeric constructs. Based on our results, we found that all constructs were successfully

expressed (Figure 2D-F).

Rescue experiments with an HdAd vector that expressed the $Ca_V2.1$ α_1 subunit FT (Mus

musculus NP_031604.3) (HdAd $Ca_V2.1$ FT) in the $Cav2.1^{-/-}$ calyx of Held in a *Cacna1a* CKO

mouse model has been validated to restore $Ca_V2.1$ current amplitudes, subtype levels and

synaptic vesicle release rates to similar levels as wild-type calyces (Lubbert *et al.*, 2017). To

validate our experimental approach, we used expressed the $Ca_V2.1$ α_1 subunit FT using HdAd

$Ca_V2.1$ FT on the $Ca_V2.1,Ca_V2.2$ CKO mouse line . Our rescue experiments show that HdAd

expression of the $Ca_V2.1$ FT in the $Ca_V2.1^{-/-}/$ $Ca_V2.2^{-/-}$ calyx of Held led to similar maximum Ca^{2+}

current amplitudes and voltage-dependent activation as the wild-type calyx of Held. These results

validated our experimental approach and confirming previous findings (Table 2).

Subsequently, we expressed the $Ca_V2.1/2.3$ chimeras and performed direct presynaptic

Ca^{2+} recordings. All chimera constructs were compared to $Ca_V2.1$ α_1 subunit FT and $Ca_V2.1^{-/-}/$

$Ca_V2.2^{-/-}$ calyces to determine rescue efficiency (Fig. 3). Since we did not find any differences in

the calyx of Held size with expression of the Ca_V2 chimeras (Table 8), we reported data as

maximal Ca^{2+} currents amplitude at all voltages. Current to voltage (IV) analysis revealed

significant reduction in peak Ca^{2+} current amplitudes with all three chimeras compared to $Ca_V2.1$

FT (Fig. 3, Table 3) However, only expression of the $Ca_V2.1/2.3$ loop II-III led to maximum Ca^{2+}

currents that were larger than that of the $Ca_V2.1^{-/-}/$ $Ca_V2.2^{-/-}$ (Maximum Ca^{2+} current amplitudes

(pA) $Ca_V2.1^{-/-}/$ $Cav2.2^{-/-}$: -425.5 ± 219.5 (n=11) vs. $Ca_V2.1/2.3$ II-III loop: -958.1 ± 331.3 (n=11),

p<0.0001) (Fig. 3E, Table 3). Based on these results, we demonstrate that $Ca_V2.1$ loop II-III and

$Ca_V2.1$ CT are positive regulators of presynaptic $Ca_V2.1$ abundance but differentially regulate

presynaptic $Ca_V2.1$ abundance.

The Ca$_V$2.1 α$_1$ subunit loop II-III and Ca$_V$2.1 α$_1$ subunit CT do not independently control Ca$_V$2.1 preference

Ablation of Ca$_V$2 subtypes leads to increased presynaptic levels of other Ca$_V$2 subtypes (Inchauspe *et al.*, 2004; Lubbert *et al.*, 2017) (Fig 1D, F), and overexpression of Ca$_V$2.1 results in increased presynaptic Ca$_V$2.1 levels and competes away Ca$_V$2.2, but not vice versa (Lubbert *et al.*, 2019). Therefore, it is postulated that presynaptic Ca$_V$2 abundance is correlated to Ca$_V$2 subtype preference. Although we found a reduction of presynaptic Ca^{2+} current with the Ca$_V$2.1/2.3 chimeras, whether these regions regulated Ca$_V$2.1 preference or just abundance is unknown. Therefore, it was important to elucidate if motifs that regulate presynaptic Ca$_V$2 subtype abundance are also utilized to determine Ca$_V$2 subtype preference or if they were distinct. The remaining Ca^{2+} currents from calyces expressing the Ca$_V$2.1/2.3 chimeras could be mediated by: 1) a mixture of Ca$_V$2.1/2.3 chimeras and Ca$_V$2.3 channels, 2) solely Ca$_V$2.1/2.3 chimeras, or 3) solely Ca$_V$2.3 channels. Since the presynaptic Ca^{2+} currents in the Ca$_V$2.1$^{-/-}$/Ca$_V$2.2$^{-/-}$ calyx of Held were completely Aga insensitive, and the Ca$_V$2.1/2.3 chimeras still maintain Aga binding sites and are the Aga sensitive, we measured Aga sensitivity of Ca2+ currents with the Ca$_V$2.1/2.3 chimeras to distinguish between the different possibilities. In scenario 1, the Ca$_V$2.1/2.3 chimeras are localized to the presynaptic membrane but do not completely away or only partially compete away Ca$_V$2.3 channels and would result in a mix of Aga sensitive and insensitive Ca^{2+} currents. In scenario 2, Ca$_V$2.1/2.3 chimeras are localized to the presynaptic membrane and compete away Ca$_V$2.3 channels and would result in similar Aga sensitivity as the Ca$_V$2.1 FT. In scenario 3, Ca$_V$2.1/2.3 chimeras are not localized to the presynaptic membrane with only Ca$_V$2.3 currents remaining and would result in similar Aga insensitivity as the Ca$_V$2.1$^{-/-}$/ Ca$_V$2.2$^{-/-}$ calyx of Held. Analysis of Aga sensitivity of Ca$_V$2.1/2.3 II-III or Ca$_V$2.1/2.3 CT showed no difference compared to the Ca$_V$2.1 FT (Fig. 4, Table 4). However, we found partial Aga sensitivity with Ca$_V$2.1/2.3 II-III+ CT (Aga sensitive fraction (%): Ca$_V$2.1 FT 89.5 ± 4.95 (n=3) vs. Ca$_V$2.1/2.3 II-III+ CT 67.67 ± 7.371 (n=3), p=0.0111) (Fig. 4, Table 4). Based on our data, we demonstrate that both Ca$_V$2.1/2.3

chimeras with the loop II-III or CT swap are able to compete away $Ca_V2.3$ channels (Fig. 4, Table

4). Therefore, we conclude that $Ca_V2.1$ loop II-III and the CT do not independently control $Ca_V2.1$

preference and motifs that regulate presynaptic $Ca_V2.1$ preference are distinct from those that

regulate presynaptic $Ca_V2.1$ abundance.

The $Ca_V2.3$ α_1 subunit CT negatively regulates presynaptic $Ca_V2.1$ abundance while the

$Ca_V2.1$ CT aa. 1933-1966 positively regulates presynaptic $Ca_V2.1$ abundance

The $Ca_V2.3$ CT contains a Calmodulin binding domain (CBD) but does not contain many

motifs postulated to mediate presynaptic $Ca_V2.1$ subtype abundance (Kamp *et al.*, 2012; Nanou

& Catterall, 2018). In addition, proposed $Ca_V2.1$ CT motifs (aa 2016 onwards) are not essential

determinants of presynaptic $Ca_V2.1$ abundance (Lubbert *et al.*, 2017). Thus, we hypothesized that

the 1) $Ca_V2.3$ CT acts as a negative regulator of presynaptic Ca_V2 abundance, and/or 2) the

$Ca_V2.1$ CBD regulates presynaptic Ca_V2 abundance. Therefore, to elucidate the mechanism of

action of the $Ca_V2.3$ CT, we created two additional $Ca_V2.1/2.3$ CT chimeras. They were the

<u>$Ca_V2.1/2.3$ CT 1967-2368</u>, which replaces the $Ca_V2.1$ CT with the $Ca_V2.3$ CT from $Ca_V2.1$ aa.

1967 onwards and contains the $Ca_V2.3$ CBD, and the <u>$Ca_V2.1/2.3$ CT 2016-2368</u> which replaces

the $Ca_V2.1$ CT with the $Ca_V2.3$ CT from $Ca_V2.1$ aa 2016 onwards (Fig. 5A). Subsequently, we

expressed these chimeras and $Ca_V2.1$ $\Delta2016$, which deletes all motifs, AZ protein binding sites,

and the secondary $Ca_V\beta$ interaction site(Lubbert *et al.*, 2017) and measured their maximum Ca^{2+}

amplitudes to determine rescue efficiency (Fig. 5).   As previously published. we found no

difference in maximum Ca^{2+} currents between $Ca_V2.1$ $\Delta2016$ and $Ca_V2.1$ FT (Fig. 5B-C, Table 5),

thereby confirming that motifs in the $Ca_V2.1$ CT aa. 2016 onwards are not essential for controlling

presynaptic $Ca_V2.1$ abundance. In contrast, expression of $Ca_V2.1/2.3$ CT 2016-2368 and

$Ca_V2.1/2.3$ CT 1967-2368 resulted in a reduction in the maximum Ca^{2+} current amplitudes ($Ca_V2.1$

$\Delta2016$: -1241 ± 207.4 (n=6) vs. $Ca_V2.1/2.3$ CT 2016-2368: -934.1 ± 210.6 (pA) (n=12), p=0.0209;

$Ca_V2.1$ $\Delta2016$: -1241 ± 207.4 (n=6) vs. $Ca_V2.1/2.3$ CT 1967-2368: -839.5 ± 251.9 (pA) (n=8),

p=0.0047) (Fig. 5D-F, Table 5), but the maximum Ca^{2+} current amplitudes with either $Ca_V2.1/2.3$

CT 1967-2368 or Ca$_V$2.1/2.3 CT 2016-2368 were larger than the Ca$_V$2.1/2.3 CT 1933-2368 ((pA)

Ca$_V$2.1/2.3 CT aa. 1933-2368: -557.7 ± 198.9 (n=9) vs. Ca$_V$2.1/2.3 CT 2016-2368: -934.1 ± 210.6

(n=12), p=0.0013; Ca$_V$2.1/2.3 CT aa. 1933-2368: -557.7 ± 198.9 (n=9) vs. Ca$_V$2.1/2.3 CT 1967-

2368: -839.5 ± 251.9 (n=8), p=0.0321) (Fig. 5D-F, Table 5). Therefore, we conclude that the

Ca$_V$2.3 CT aa. 1952-2273 contains a motif or motifs that negatively regulate presynaptic Ca$_V$2.3

abundance and the Ca$_V$2.1 CBD is not a positive regulator of presynaptic Ca$_V$2.1 abundance.

Furthermore, we identified the region between Ca$_V$2.1 CT aa. 1933-1966 positively regulates

presynaptic Ca$_V$2.1 abundance.

The Ca$_V$2.2 CT does not control presynaptic Ca$_V$2.1 abundance

Since we identified that the Ca$_V$2.3 CT was a negative regulator of presynaptic Ca$_V$2.3

abundance, we set out to determine if this was unique to the Ca$_V$2.3 CT. Thus, to address if

Ca$_V$2.2 CT negatively regulated presynaptic Ca$_V$2 abundance, we created a Ca$_V$2.1/2.2 CT

chimera and cloned this into our HdAd vectors. Western blot analysis demonstrated no issues

with expression of the Ca$_V$2.1/2.2 CT chimera (Fig. 6A). Subsequently we expressed the

Ca$_V$2.1/2.2 CT chimera and performed IV analysis on presynaptic Ca^{2+} currents. Results of our

IV analysis revealed that there was no difference in the maximum Ca^{2+} current between the

Ca$_V$2.1/2.2 CT chimera and Ca$_V$2.1 FT (Fig. 6B-E). However, compared to Ca$_V$2.1 FT, there

appeared to be unequal variances of maximum Ca^{2+} current amplitudes with Ca$_V$2.1/2.2 CT that

was marginally significant (F-test comparing variances p=0.0607) (Fig. 6B-E, Table 6). Therefore,

our data suggests that the Ca$_V$2.2 CT is not a negative regulator of presynaptic abundance

High level overexpression of Ca$_V$2.1/2.2 CT rescues presynaptic Ca^{2+} currents while

Ca$_V$2.1/2.3 CT does not.

High level overexpression of Ca$_V$2.1 α$_1$ subunit results in increased presynaptic Ca$_V$2.1

levels and compete away Ca$_V$2.2 and Ca$_V$2.3 channels, however high-level overexpression of

Ca$_V$2.2 α$_1$ subunit does not result in a reduction of Ca$_V$2.1 levels at the calyx of Held (Lubbert *et*

al., 2019). This suggests that upstream regulatory steps, such as protein trafficking and

membrane insertion steps, are integral to $Ca_V2.1$ level preference and abundance (Ferron *et al.*,

2021; Young & Veeraraghavan, 2021). It is known that high levels of overexpression can saturate

regulatory steps that control trafficking (Lubbert *et al.*, 2019). Therefore, to determine if these

regulatory step(s) could be saturated, we used the high-level overexpression Pun cassette

(Young & Neher, 2009; Montesinos *et al.*, 2011; Lubbert *et al.*, 2019) to drive expression of

$Ca_V2.1/2.2$ CT and $Ca_V2.1/2.3$ CT chimeras and determine if we could increase the presynaptic

Ca_V2 chimera current levels and compared their maximum Ca^{2+} current amplitudes to Pun $Ca_V2.1$

FT. We found no difference in presynaptic Ca^{2+} currents with Pun $Ca_V2.1/2.2$ CT compared to

Pun $Ca_V2.1$ FT (Fig. 7, Table 7). However, maximum Ca^{2+} current amplitudes with Pun $Ca_V2.1/2.3$

CT were lower than Pun $Ca_V2.1$ FT (Pun $Ca_V2.1$ FT: -1438 ± 263.9 (n=7) vs. Pun $Ca_V2.1/2.3$ CT:

-772.8 ± 291.8 (n=10), (pA), p=0.0006) (Fig. 7, Table 7). Moreover, the maximum Ca^{2+} current

amplitudes with Pun $Ca_V2.1/2.3$ CT showed no significant difference than syn $Ca_V2.1/2.3$ CT (Fig.

7, Table 7). Therefore, we demonstrate that the $Ca_V2.2$ CT is not a negative regulator of

presynaptic $Ca_V2.2$ abundance and the $Ca_V2.1/2.2$ CT chimera can be forced to increase

expression level, however, the negative regulator motif in $Ca_V2.3$ CT restrains the upstream

regulatory steps which results in low abundance of $Ca_V2.3$ channels in the presynaptic terminal.

Discussion

In this study, we provide insight into how presynaptic Ca_V2 subtype abundance is determined by the Ca_V2 α_1 subunit. Based on our data, we demonstrate that the cytoplasmic loop II-III and CT regions of the $Ca_V2.1$ α_1 subunit are positive regulators of presynaptic Ca_V2 abundance, while $Ca_V2.3$ CT negatively regulates presynaptic Ca_V2 abundance. In addition, we find that $Ca_V2.2$ CT is not a key determinant of presynaptic Ca_V2 abundance. Furthermore, we demonstrate that $Ca_V2.1$ α_1 subunit cytoplasmic loop II-III and CT regions do not independently control presynaptic $Ca_V2.1$ preference. Therefore, we propose that the motifs controlling presynaptic $Ca_V2.1$ preference are distinct from those regulating $Ca_V2.1$ levels and they may act synergistically in pathways that regulate presynaptic $Ca_V2.1$ preference and levels.

The Ca_V2 α_1 subunit II-III loop regulation of presynaptic $Ca_V2.1$ abundance.

By swapping the loop II-III domain between $Ca_V2.1$ and $Ca_V2.3$ we demonstrated that the $Ca_V2.1$ loop II-III region positively regulates presynaptic $Ca_V2.1$ abundance but does not regulate preference. All Ca_V2 subunit loop II-III domains contain a synprint region which is the site of interaction between the SNARE proteins syntaxin and SNAP-25 (Zhong *et al.*, 1999; Mochida *et al.*, 2003; Yokoyama *et al.*, 2005). Although $Ca_V2.3$ currents are modulated by syntaxin (Wiser *et al.*, 2002; Kamp *et al.*, 2005), the proposed syntaxin binding domain found in $Ca_V2.1$ and $Ca_V2.2$ is not found in the $Ca_V2.3$ α_1 subunit synprint region (Kamp *et al.*, 2005; Yokoyama *et al.*, 2005; Rajapaksha *et al.*, 2008). Our findings that $Ca_V2.1/2.3$ loop II-III chimera resulted in presynaptic Ca^{2+} current reduction indicates that potential differences in syntaxin binding between the different subtypes of Ca_V2 α_1 subunits regulates presynaptic Ca_V2 abundance. Studies from cultured hippocampal neurons, $Ca_V2.1$ dominant presynaptic terminals, found that partial deletions in the $Ca_V2.2$ synprint region that impact SNARE binding resulted in less efficient $Ca_V2.2$ targeting to presynaptic terminal (Szabo *et al.*, 2006). Similar partial deletions in the $Ca_V2.1$ synprint region reduced $Ca_V2.1$ localization to release sites in cultured superior cervical ganglion (SCG) neurons which utlize $Ca_V2.2$ dominant presynaptic terminals (Mochida *et al.*, 2003). In addition, splice

variants of $Ca_V2.1$ lacking the synprint region reduce incorporation into neuroendocrine cells

(Rajapaksha *et al.*, 2008). Since partial deletions of $Ca_V2.2$ synprint do not impair axonal

targeting(Szabo *et al.*, 2006), it is possible that the $Ca_V2.3$ synprint region does not regulate

axonal trafficking but rather regulates incorporation into the presynaptic terminal. Although the

$Ca_V2.3$ synprint region chimera resulted in reduced presynaptic Ca_V2 chimera levels, there was

no change in the proportion of Agatoxin sensitive currents. Therefore, this indicates the

mechanisms that control $Ca_V2.1$ preference are not determined by the loop II-III region binding

interactions.

The Ca_V2 $α_1$ subunit CT differentially regulates presynaptic Ca_V2 subtype abundance.

We found that the $Ca_V2.2$ CT and $Ca_V2.3$ CT had differential effects on presynaptic Ca_V2

abundance but did not impact Ca_V2 preference. Our data demonstrated that $Ca_V2.1/2.3$ CT

chimera results in a severe reduction in presynaptic Ca^{2+} currents and suggested that the $Ca_V2.3$

CT is a negative regulator of presynaptic $Ca_V2.3$ abundance. Prior studies (Lubbert *et al.*, 2017)

and our findings here revealed that the $Ca_V2.1$ CT Δ2016, which deletes multiple AZ protein

binding sites and other motifs restores presynaptic Ca^{2+} to $Ca_V2.1$ FT levels. These data

demonstrate these interactions and motifs are not essential for controlling presynaptic $Ca_V2.1$

abundance. However, replacing the $Ca_V2.1$ CT from aa. 2016 onwards with corresponding $Ca_V2.3$

CT aa. 1952-2273 resulted in a reduction of presynaptic Ca^{2+} currents compared to $Ca_V2.1$ CT

Δ2016. Furthermore, overexpression of $Ca_V2.1/2.3$ CT chimera by pUNISHER cassette was

similar to presynaptic Ca^{2+} currents levels with the low-level synapsin expression cassette. Based

on our data that showed the $Ca_V2.1$ CT aa. 2016-2368 is dispensable for $Ca_V2.1$ abundance, we

propose that $Ca_V2.3$ CT aa. 1952-2273 contains a motif that prevents presynaptic Ca_V2 targeting

or incorporation into the plasma membrane. In addition, our results indicate that CT dependent

mechanism of regulation of presynaptic Ca_V2 abundance is dominant to the loop II-III region

mechanisms of action as $Ca_V2.1/2.3$ CT chimera resulted in significantly lower presynaptic Ca^{2+}

currents than those with the $Ca_V2.1/2.3$ loop II-III.

Calmodulin (CaM) is considered a core component of Ca_V2 channel complexes and

interacts with the Cav2.1 channels through a bipartite interaction with the calmodulin binding

domain (CBD) and IQ-like motif (IM) domain to regulate calcium dependent facilitation (CDF) and

inactivation (CDI) (Lee *et al.*, 2003; Chaudhuri *et al.*, 2004; Mori *et al.*, 2008). However, how CaM

interactions regulate presynaptic $Ca_V2.1$ levels is unclear. The $Ca_V2.1/2.3$ CT aa. 1967-2368

chimera, which replaces the $Ca_V2.1$ CBD with the $Ca_V2.3$ CBD, showed no difference in rescuing

efficiency compared to $Ca_V2.1/2.3$ CT aa. 2016-2368 chimeras. However, we found that

presynaptic Ca^{2+} currents with $Ca_V2.1/2.3$ CT aa. 1967-2368 chimera were larger than that with

$Ca_V2.1/2.3$ CT aa. 1933-2368 chimera. Therefore, our results demonstrated that the $Ca_V2.1$ CBD

does not control presynaptic $Ca_V2.1$ abundance but the linker region of $Ca_V2.1$ CT aa. 1933-1966

between the IQ domain and the CBD plays a role in regulation of $Ca_V2.1$ levels. Since the

CaM/CBD complex interacts with the IM domain (Nanou *et al.*, 2016; Serra *et al.*, 2018), it is

possible that the $Ca_V2.3$ CBD and linker region with $Ca_V2.1$ IQ domain result in a different

structural conformation of the CT that impacts presynaptic Ca_V2 abundance. Interestingly, the

combination of the mutation (IM-AA) in the IM domain and CBD deletion in SCG neurons results

in an increased peak Ca^{2+} currents levels, while the IM-AA or CBD deletion alone did not alter

Ca^{2+} current levels (Mochida *et al.*, 2008). Therefore, the IQ domain and CBD may synergistically

regulate $Ca_V2.1$ level. However, the molecular mechanisms how these regions impact $Ca_V2.1$

abundance is unknown and future experiments are needed.

In contrast to the $Ca_V2.1/2.3$ CT chimera, we found no difference in the average

presynaptic Ca^{2+} current levels with the $Ca_V2.1/2.2$ CT chimera compared to $Ca_V2.1$ FT. However,

we observed a large variability of Ca^{2+} current amplitudes with $Ca_V2.1/2.2$ CT that is marginally

significant compared to $Ca_V2.1$ FT rescue. Therefore, our data implies that the $Ca_V2.2$ CT is not

a key determinant of presynaptic Ca_V2 abundance. Unlike the $Ca_V2.3$ CT, high level

overexpression of $Ca_V2.1/2.2$ CT resulted in similar presynaptic Ca^{2+} current levels compared to

high level overexpression of $Ca_V2.1$ FT level and no change in variability. Therefore, we propose

that the $Ca_V2.2$ CT does not contain motifs that prevents presynaptic $Ca_V2.2$ targeting or

incorporation into the plasma membrane.

Ca_V2 subtype abundance and preference are regulated by distinct pathways.

Although there were differences in the presynaptic Ca^{2+} current levels with the $Ca_V2.1/2.3$

loop II-III and $Ca_V2.1/2.3$ CT chimeras, they had similar Aga sensitivity as presynaptic Ca^{2+}

currents with the $Ca_V2.1$ FT. However, there was a significant Aga insensitive component of Ca^{2+}

currents with $Ca_V2.1/2.3$ loop II-III+ CT chimera despite having similar Ca^{2+} current levels as the

$Ca_V2.1/2.3$ CT and $Ca_V2.1^{-/-}/ Ca_V2.2^{-/-}$ background. Therefore, our data indicates that: 1)

Presynaptic Ca_V2 subtype abundance mechanisms are independent of those that control

preference. 2) The loop II-III region and the CT region do not individually encode preference. 3)

The $Ca_V2.1$ II-III loop and CT region may act synergistically to control Ca_V2 subtype preference.

4) Other motifs in the $Ca_V2.1$ α_1 subunit outside of the CT and synprint region are involved in

regulating presynaptic Ca_V2 subtype preference. Potential candidates that may control preference

are the Ca_V2 α_1 subunit at AID in domain I-II which binds to the $Ca_V\beta$ subunit with high affinity(De

Waard *et al.*, 1994). $Ca_V\beta$ subunits are critical for regulating Ca_V2 trafficking and $Ca_V\beta4$ and

$Ca_V\beta2a$ are located within the presynaptic terminal. $Ca_V2.1$ channels have higher specificity for

binding to $Ca_V\beta4$ than $Ca_V2.2$ and $Ca_V2.3$ channels (Xie *et al.*, 2007; Muller *et al.*, 2010). Another

possibility is that preference is not encoded by motifs in the cytoplasmic regions but is controlled

by the Cacna2d protein family (Dolphin, 2012; Hoppa *et al.*, 2012; Stephani *et al.*, 2019). Knockout

of $\alpha_2\delta3$ subunit in spiral ganglion neurons results in reduced $Ca_V2.1$ and $Ca_V2.3$ components with

no change in $Ca_V2.2$ subtype levels (Stephani *et al.*, 2019). In contrast overexpression of $\alpha_2\delta1$

subunit decreases $Ca_V2.2$ level in cultured hippocampal neurons (Pilch *et al.*, 2022), but did not

change other Ca_V2 subtype levels. Furthermore, the globular bushy cells which give rise to the

calyx of Held, only express *Cacna2d2* and *Cacna2d3*. In addition, other synapses that undergo

developmental reduction of $Ca_V2.2$ and $Ca_V2.3$ express *Cacna2d1*. Finally, the $\alpha_2\delta$ binding

preference does not correlate to presynaptic Ca$_V$2 subtype levels (Voigt *et al.*, 2016) while

CACNA2D expression patterns do not correlate to presynaptic Cav2 subytpe levels (Cole *et al.*,

2005). Therefore, given these conflicting findings more experiments will be needed to decipher if

the Cacna2d proteins regulate preference.

While it is possible that the loop II-III domain and the CT domain act in concert to regulate

presynaptic preference, it is also possible that they are not involved in preference mechanisms

but rather combination of the loop II-III and CT domain swap construct dramatically impair the

Ca$_V$2 chimera trafficking to the presynaptic terminal. In this scenario, the Ca$_V$2.1/2.3 II-III loop+

CT chimera does not effectively compete away native Ca$_V$2.3 channels, which are normally

restricted from the upstream pathways that regulate presynaptic trafficking. Therefore, native

Ca$_V$2.3 channels are also trafficked to the presynaptic terminals that express the Ca$_V$2.1/2.3 II-III

loop+ CT chimera. In this hypothetical scenario, the preference mechanism(s) has high affinity for

the Ca$_V$2.1/2.3 chimera channel and low affinity for the Ca$_V$2.3 channels, however by law of mass

action, Ca$_V$2.3 channels now participate in molecular pathways that determine preference and

results in the presence of presynaptic Ca$_V$2.3 channels. Based on our model, this demonstrates

how alternative splicing of the Ca$_V$2 α_1 subunits and potentially alternative splicing of molecules

in pathways that control Ca$_V$2 trafficking (Weiss & Zamponi, 2017), and insertion into the

membrane may lead to developmentally changes in presynaptic Ca$_V$2 subtype levels. However,

the molecular machinery and the motifs that determine preference remain to be identified.

By demonstrating that motifs in the Ca$_V$2 α_1 subunit that control presynaptic Ca$_V$2 subtype

abundance are distinct from those determining presynaptic Ca$_V$2 subtype preference, our findings

provide new insights into the regulation of presynaptic Ca$_V$2 subtype levels.

Figure legends

Figure 1. Presynaptic Ca$_V$2.1 and Ca$_V$2.2 channels can be ablated and the remaining Ca^{2+} currents are compensated by Ca$_V$2.3 channels at the calyx of Held.

(A) Illustration of CKO mouse model. Exon 4 of *Cacna1a*, and exons 5 and 6 of *Cacna1b* are flanked by LoxP sites. (B-C) Example traces of Ca^{2+} currents and bar graph show rundown fraction of electrophysiological recordings at duration of 10~15min at wildtype calyx of Held (n=3). (D) Pharmacological isolation of presynaptic Ca$_V$2 isoforms in wildtype and Ca$_V$2.1$^{-/-}$/ Ca$_V$2.2$^{-/-}$ calyces. Ca^{2+} current traces in absence of any blockers (black), after blocking Ca$_V$2.1 fraction with 200 nM Aga (orange), after blocking Ca$_V$2.2 fraction with 2 mM Cono (blue) and after blocking all Ca$_V$ channels with 50 mM Cd^{2+} (gray). (E) Bar graph shows total Ca^{2+} currents in wildtype and Ca$_V$2.1$^{-/-}$/ Ca$_V$2.2$^{-/-}$ calyces. (F) Relative Ca$_V$2 subtype fractions in wildtype, Ca$_V$2.1$^{-/-}$/ Ca$_V$2.2$^{-/-}$ calyces (n=3-5 for each condition). (G) Ca^{2+} current traces in absence of any blockers (black), after blocking Ca$_V$2.3 fraction with 200 nM SNX-482 (green). (H) Bar graph shows SNX-482 sensitive fraction at Ca$_V$2.1$^{-/-}$/ Ca$_V$2.2$^{-/-}$ calyces (n=3). (I) Ca^{2+} current traces in absence of any blockers (black), after blocking Ca$_V$1 fraction with 10 μM Nifedipine (red). (J) Bar graph shows the reduction fraction and Ca^{2+} current amplitudes in Ca$_V$2.1$^{-/-}$/ Ca$_V$2.2$^{-/-}$ calyces after Nifedipine treatment compared to wildtype rundown fraction.

Figure 2. Ca$_V$2.1/Ca$_V$2.3 Chimeras can be efficiently expressed

Western Blot validated protein expression of Ca$_V$2 hybrid constructs. HEK293 cells are infected by HdAd Ca$_V$2.1/2.3 chimeric constructs. (A) Illustration of Ca$_V$2 α$_1$ structure. Four transmembrane domains are highly conserved while the cytoplasmic loops are highly variant between Ca$_V$2.1 and Ca$_V$2.3, the I-II loop, II-III loop and CT share 87.3%, 33.2% and 39.8% consensus, respectively. (B) Illustration of swapped region between Ca$_V$2.1 and Ca$_V$2.3 channels, the amino acid numbers are labelled. (C) Illustration of three Ca$_V$2.1/2.3 chimeric constructs. Ca$_V$2.1 II-III loop and CT are both replaced by Ca$_V$2.3 II-III loop and CT in Ca$_V$2.1/2.3 II-III loop + CT chimeric construct. Ca$_V$2.1

II-III loop is replaced by $Ca_V2.3$ II-III loop in $Ca_V2.1/2.3$ II-III loop chimeric construct. $Ca_V2.1$ CT is

replaced by $Ca_V2.3$ CT in $Ca_V2.1/2.3$ CT chimeric construct. (D) Antibody targeting $Ca_V2.1$ CT

shows the protein expression of $Ca_V2.1$ FT and $Ca_V2.1/2.3$ II-III loop. (E) Antibody targeting

$Ca_V2.1$ II-III loop fails to detect protein expression of $Ca_V2.1/2.3$ II-III loop and $Ca_V2.1/2.3$ II-III

loop CT, indicating that the II-III loop of $Ca_V2.1$ is swapped by $Ca_V2.3$ II-III loop. (F) Antibody

targeting $Ca_V2.3$ CT shows the protein expression of $Ca_V2.1/2.3$ II-III loop CT and $Ca_V2.1/2.3$ CT.

Mock, negative control, protein lysis from uninfected HEK293 cells.

Figure 3. The $Ca_V2.1/2.3$ α_1 CT chimera cannot rescue Ca^{2+} currents to $Ca_V2.1$ FT levels,

however the $Ca_V2.1/2.3$ loop II-III chimera partially rescues Ca^{2+} currents.

(A) Sample traces of Ca^{2+} current triggered by 50 ms voltage steps from -60 mV to -30 mV in 10

mV steps, -25 mV to +20 mV in 5 mV steps, and +30 mV to +60 mV in 10 mV steps. Black: calyces

expressing $Ca_V2.1$ FT. Yellow: $Ca_V2.1^{-/-}/Ca_V2.2^{-/-}$ calyces. Red: calyces expressing $Ca_V2.1/2.3$ II-

III loop CT chimeric construct. Blue: calyces expressing $Ca_V2.1/2.3$ II-III loop chimeric construct.

Gray: calyces expressing $Ca_V2.1/2.3$ CT chimeric construct. (B-C) IV relationship of Ca^{2+} currents

(B) and normalized Ca^{2+} currents by maximal Ca^{2+} currents (I/I_{max}; C) plotted against voltage

steps. (D-E) Bar graph shows absolute peak Ca^{2+} current amplitudes, one-way ANOVA, Dunnett's

test, $Ca_V2.1$ FT is used as positive control (D) and $Ca_V2.1^{-/-}/Ca_V2.2^{-/-}$ is used as negative control

(E).

Figure 4. The $Ca_V2.1/2.3$ α_1 subunit chimeras compete away $Ca_V2.3$ channel in the calyx of

Held.

(A) Agatoxin isolation of presynaptic $Ca_V2.1/2.3$ chimeras in calyces expressing $Ca_V2.1$ FT (black,

n=3), $Ca_V2.1/2.3$ II-III loop CT (red, n=3), $Ca_V2.1/2.3$ II-III loop (blue, n=4), and $Ca_V2.1/2.3$ CT

(gray, n=3). Basal and reduced Ca^{2+} current traces after applying 200 nM ω-Aga IVA are labelled.

(B) Reduction of Agatoxin sensitive Ca^{2+} current over time. Ca^{2+} current amplitudes are plotted

against time, 200 nM ω-Aga IVA was applied at 5[th] pulse (50s). (C) Bar graph shows the ω-Aga

IVA sensitive reduction fraction in calyces expressing $Ca_V2.1$ FT, $Ca_V2.1/2.3$ II-III loop CT,

$Ca_V2.1/2.3$ II-III loop, and $Ca_V2.1/2.3$ CT.

Figure 5.   The $Ca_V2.3$ CT negatively regulates presynaptic Ca_V2 abundance, while the

$Ca_V2.1$ CT region aa. 1933-1966 promotes $Ca_V2.1$ abundance

(A) Amino acid alignment of $Ca_V2.1$ and $Ca_V2.3$ CT. Sequences on top and bottom represent

$Ca_V2.1$ CT and $Ca_V2.3$ CT, respectively. Amino acid numbers of swapped sites in three $Ca_V2.1/2.3$

chimeras are indicated in black. $Ca_V2.1$ CBD is highlighted in green. (B) Example Ca^{2+} current

traces to 50 ms depolarization from -80mV holding potential to the maximal activation voltage

step (-15mV to +15mV, 5 mV steps) for $Ca_V2.1$ FT (black, n=11) and $Ca_V2.1$ Δ2016 (red, n=6).

(C) Bar graph shows absolute peak Ca^{2+} currents amplitude between $Ca_V2.1$ CT and $Ca_V2.1$

Δ2016, unpaired t test. (D) Example Ca^{2+} current traces to 50ms depolarization from -80mV

holding potential to the maximal activation voltage step (-15mV to +15mV, 5 mV steps) for $Ca_V2.1$

Δ2016, $Ca_V2.1/2.3$ CT aa. 2016-2368 (n=12); $Ca_V2.1/2.3$ CT 1967-2368 (n=8) and $Ca_V2.1/2.3$ CT

1933-2368 (n=9). (E-F) Bar graph shows peak Ca^{2+} currents amplitude, one-way ANOVA,

Dunnett's test, $Ca_V2.1$ Δ2016 is used as positive control (E) and $Ca_V2.1/2.3$ CT aa. 1933-2368 is

used as negative control (F).

Figure 6. $Ca_V2.1/2.2$ CT chimera can be efficiently expressed and rescues Ca^{2+} currents to

$Ca_V2.1$ FT level with high variability.

(A) Western Blot validated protein expression of $Ca_V2.1/2.2$ CT. HEK293 cells are infected by

HdAd $Ca_V2.1/2.2$ CT and $Ca_V2.2$ FT. Antibody targeting $Ca_V2.2$ CT shows the protein expression

of $Ca_V2.2$ FT and $Ca_V2.1/2.2$ CT. Mock, negative control, protein lysis from uninfected HEK293

cells. (B) Exemplary Ca^{2+} currents triggered by 50 ms voltage steps from -60 mV to -30 mV in 10

mV steps, -25 mV to +20 mV in 5 mV steps, and +30 mV to +60 mV in 10 mV steps. Black: calyces

expressing $Ca_V2.1$ FT (n=11). Yellow: $Ca_V2.1^{-/-}/$ $Ca_V2.2^{-/-}$ calyces (n=11). Green: calyces

expressing $Ca_V2.1/2.2$ CT (n=11), two phenotypes are demonstrated. (C-D) IV relationship of

Ca^{2+} currents (C) and normalized Ca^{2+} currents by maximal Ca^{2+} currents (I/I$_{max}$; D) plotted against

voltage steps. (E) Bar graph shows absolute peak Ca^{2+} current amplitudes. One-way ANOVA,

Dunnett's test, Ca$_V$2.1 FT is used as positive control, Ca$_V$2.1$^{-/-}$/ Ca$_V$2.2$^{-/-}$ is used as negative

control.

Figure 7. High level overexpression of the Ca$_V$2.1/2.3 α$_1$ CT cannot rescue Ca^{2+} currents to

Pun Ca$_V$2.1 levels while Pun Ca$_V$2.1/2.2 α$_1$ CT can.

(A) High-level overexpression cassette pUNISHER (Pun) is used to drive Ca$_V$2.1 FT, Ca$_V$2.1/2.2

CT and Ca$_V$2.1/2.3 CT chimeric constructs. Example Ca^{2+} current traces to 50ms depolarization

from -80mV holding potential to the maximal activation voltage step (-15mV to +15mV, 5 mV

steps) for calyces expressing Pun Ca$_V$2.1 FT (green, n=7), Pun Ca$_V$2.1/2.2 CT (purple, n=9) and

Pun Ca$_V$2.1/2.3 CT (orange, n=10) (B) Bar graph shows averaged peak Ca^{2+} currents for Pun

Ca$_V$2.1 FT, Pun Ca$_V$2.1/2.2 CT and Pun Ca$_V$2.1/2.3 CT constructs. One-way ANOVA, Dunnett's

test, Pun Ca$_V$2.1 FT is used as positive control.

Table1. *Cacna1a, Cacna1b* flox/flox animal characterization.

Parameter	Mean	SD	n	P	statistics
Wildtype animal					
Agatoxin sensitive fraction (%)	79.09	14.98	4		
Agatoxin sensitive currents (pA)	809.3	255.6	4	control	
Conotoxin sensitive fraction (%)	7.828	10.5	4		
Conotoxin sensitive currents (pA)	92.00	142.2	4		
$Ca_V2.1^{-/-}$/ $Ca_V2.2^{-/-}$					
Agatoxin sensitive fraction (%)	13.64	6.24	5	p<0.0001 (****)	
Agatoxin sensitive currents (pA)	99.22	53.64	5	p=0.0005 (***)	
Conotoxin sensitive fraction (%)	11.92	4.722	5	p=0.4571 (n.s.)	Unpaired t test
Conotoxin sensitive currents (pA)	50.80	43.10	5	p=0.5532 (n.s.)	
SNX-482 sensitive fraction (%)	24.33	9.074	3		
SNX-482 sensitive currents (pA)	86.65	31.26	3		
Rundown fraction					
Wildtype animal (pA)	43.70	20.85	3	control	Unpaired t test
$Ca_V2.1^{-/-}$/ $Ca_V2.2^{-/-}$ (pA)	51.40	23.78	3		
Wildtype animal (%)	3.167	0.6525	3	control	Unpaired t test
$Ca_V2.1^{-/-}$/ $Ca_V2.2^{-/-}$ (%)	8.393	4.169	3	P=0.0985	

754
755
756

Table2. Electrophysiological parameters of IV relations of Ca^{2+} currents.

Parameter	Mean	SD	n	P	statistics
Max. Ca^{2+} current amplitude I_{max} (pA)					
Wildtype (positive control)	-1078	260.9	10	control	One-Way ANOVA
$Ca_V2.1$ FT	-1222	256.2	11	p=0.3140 (n.s.)	Dunnett's Test
$Ca_V2.1^{-/-}$/ $Cav2.2^{-/-}$	-425.5	219.5	11	p<0.0001 (****)	
Wildtype	-1078	260.9	10	p<0.0001 (****)	One-Way ANOVA
$Ca_V2.1$ FT	-1222	256.2	11	p<0.0001 (****)	Dunnett's Test
$Ca_V2.1^{-/-}$/ $Cav2.2^{-/-}$ (negative control)	-425.5	219.5	11	control	
IV fit: Half-maximal activation voltage V_m (mV)					

Wildtype (control)	-21.61	5.205	10	positive control	One-Way ANOVA
$Ca_V2.1$ FT	-19.75	3.212	11	p=0.5848 (n.s.)	Dunnett's test
$Ca_V2.1^{-/-}$/ $Cav2.2^{-/-}$	-12.38	5.791	11	p=0.0003 (***)	

IV fit: Voltage-dependence of activation K_m (mV)

Wildtype (control)	2.948	1.673	10	positive control	One-Way ANOVA
$Ca_V2.1$ FT	2.854	0.7658	11	p=0.9918 (n.s.)	Dunnett's test
$Ca_V2.1^{-/-}$/ $Cav2.2^{-/-}$	8.123	3.005	11	p<0.0001 (****)	

Boltzmann fit: Half-maximal activation voltage $V_{0.5}$ (mV)

Wildtype (control)	-19.6	5.029	10	positive control	One-Way ANOVA
$Ca_V2.1$ FT	-17.29	4.761	11	p=0.4829 (n.s.)	Dunnett's test
$Ca_V2.1^{-/-}$/ $Cav2.2^{-/-}$	-14.82	5.428	11	p=0.0716 (n.s.)	

Boltzmann fit: Voltage-dependence K (mV)

Wildtype (control)	3.654	1.981	10	positive control	One-Way ANOVA
$Ca_V2.1$ FT	5.052	5.07	11	p=0.5672 (n.s.)	Dunnett's test
$Ca_V2.1^{-/-}$/ $Cav2.2^{-/-}$	5.756	2.624	11	p=0.3036 (n.s.)	

758
759 **Table3. Electrophysiological parameters of IV relations of Ca^{2+} currents.**
760

Parameter	Mean	SD	n	P	statistics
Max. Ca^{2+} current amplitude I_{max} (pA)					
$Ca_V2.1$ FT (positive control)	-1222	256.2	11	control	
$Ca_V2.1^{-/-}$/ $Cav2.2^{-/-}$	-425.5	219.5	11	p<0.0001 (****)	One-Way ANOVA
Cav2.1/2.3 II-III loop CT	-470.5	112.6	8	p<0.0001 (****)	Dunnett's Test
Cav2.1/2.3 II-III loop	-958.1	331.3	11	p=0.0479 (*)	
Cav2.1/2.3 CT	-557.7	198.9	9	p<0.0001 (****)	
$Ca_V2.1$ FT	-1222	256.2	11	p<0.0001 (****)	
$Ca_V2.1^{-/-}$/ $Cav2.2^{-/-}$ (negative control)	-425.5	219.5	11	control	One-Way ANOVA
Cav2.1/2.3 II-III loop CT	-470.5	112.6	8	P=0.9849 (n.s.)	Dunnett's Test
Cav2.1/2.3 II-III loop	-958.1	331.3	11	p<0.0001 (****)	
Cav2.1/2.3 CT	-557.7	198.9	9	P=0.5723 (n.s.)	
IV fit: Half-maximal activation voltage V_m (mV)					

Ca$_V$2.1 FT (control)	-19.75	3.212	11	control	
Ca$_V$2.1$^{-/-}$/ Cav2.2$^{-/-}$	-12.38	5.791	11	p=0.0392 (*)	One-Way ANOVA
Cav2.1/2.3 II-III loop CT	-17.62	8.814	8	p=0.8972 (n.s.)	Dunnett's Test
Cav2.1/2.3 II-III loop	-15.52	7.123	11	p=0.3787 (n.s.)	
Cav2.1/2.3 CT	-12.85	7.367	9	p=0.0780 (n.s.)	
IV fit: Voltage-dependence of activation K$_m$ (mV)					
Ca$_V$2.1 FT (control)	2.854	1.673	11	control	One-Way ANOVA
Ca$_V$2.1$^{-/-}$/ Cav2.2$^{-/-}$	8.123	0.7658	11	p=0.0002 (***)	Dunnett's test
Cav2.1/2.3 II-III loop CT	10.48	4.514	8	p<0.0001 (****)	
Cav2.1/2.3 II-III loop	3.622	1.148	11	p=0.9159 (n.s.)	
Cav2.1/2.3 CT	7.886	3.293	9	p=0.0007 (***)	
Boltzmann fit: Half-maximal activation voltage V$_{0.5}$ (mV)					
Ca$_V$2.1 FT (control)	-17.29	4.761	11	control	One-Way ANOVA
Ca$_V$2.1$^{-/-}$/ Cav2.2$^{-/-}$	-14.82	5.428	11	p=0.7135 (n.s.)	Dunnett's test
Cav2.1/2.3 II-III loop CT	-11.25	6.953	8	p=0.0924 (n.s.)	One-Way ANOVA
Cav2.1/2.3 II-III loop	-14.25	5.943	11	p=0.5513 (n.s.)	Dunnett's test
Cav2.1/2.3 CT	-11.34	5.750	9	p=0.0841 (n.s.)	
Boltzmann fit: Voltage-dependence K (mV)					
Ca$_V$2.1 FT (control)	5.052	5.07	11	positive control	
Ca$_V$2.1$^{-/-}$/ Cav2.2$^{-/-}$	5.756	2.624	11	p=0.9880 (n.s.)	One-Way ANOVA
Cav2.1/2.3 II-III loop CT	6.670	4.116	8	p=0.8508 (n.s.)	Dunnett's test
Cav2.1/2.3 II-III loop	6.628	4.506	11	p=0.8231 (n.s.)	
Cav2.1/2.3 CT	7.556	5.261	9	p=0.5365 (n.s.)	

761
762
763 **Table4. Agatoxin sensitive fraction.**
764

Parameter	Mean	SD	n	P	statistics
Agatoxin sensitive fraction (%)					
Ca$_V$2.1 FT	89.5	4.95	3	control	
Cav2.1/2.3 II-III loop CT	67.67	7.371	3	p=0.0111 (*)	One-Way ANOVA
Cav2.1/2.3 II-III loop	93	1.414	4	p=0.8282 (n.s.)	Dunnett's Test
Cav2.1/2.3 CT	86.33	9.074	3	p=0.8787 (n.s.)	

Table5. Electrophysiological parameters of IV relations of Ca^{2+} currents.

Parameter	Mean	SD	n	P	statistics
Max. Ca^{2+} current amplitude I_{max} (pA)					
$Ca_V2.1$ FT	-1222	256.2	11	control	
$Ca_V2.1$ Δ2016	-1241	207.4	6	p=0.8827 (n.s.)	Unpaired t test
$Ca_V2.1$ Δ2016	-1241	207.4	6	Positive control	
Cav2.1/2.3 CT2016-2368	-934.1	210.6	12	p=0.0209 (*)	One-Way ANOVA
Cav2.1/2.3 CT1967-2368	-839.5	251.9	8	p=0.0047 (**)	Dunnett's Test
Cav2.1/2.3 CT1933-2368	-557.7	198.9	9	P<0.0001 (****)	
$Ca_V2.1$ delta 2016	-1241	207.4	6	P<0.0001 (****)	
Cav2.1/2.3 CT 2016-2368	-934.1	210.6	12	p=0.0013 (**)	One-Way ANOVA
Cav2.1/2.3 CT 1967-2368	-839.5	251.9	8	P=0.0321 (*)	Dunnett's Test
Cav2.1/2.3 CT 1933-2368	-557.7	198.9	9	Negative control	

Table6. Electrophysiological parameters of IV relations of Ca^{2+} currents.

Parameter	Mean	SD	n	P	statistics
Max. Ca^{2+} current amplitude I_{max} (pA)					
$Ca_V2.1$ FT	-1222	256.2	11	control	One-Way ANOVA
$Ca_V2.1/2.2$ CT	-1049	479.4	11	p=0.3885 (n.s.)	Dunnett's Test
$Ca_V2.1^{-/-}/$ Cav2.2$^{-/-}$	-425.5	219.5	11	P<0.0001 (****)	

Table7. Electrophysiological parameters of IV relations of Ca^{2+} currents.

Parameter	Mean	SD	n	P	statistics
Max. Ca^{2+} current amplitude I_{max} (pA)					
Pun $Ca_V2.1$ FT	-1438	263.9	7	control	One-Way ANOVA
Pun $Ca_V2.1/2.2$ CT	-1534	370.4	9	p=0.7617 (n.s.)	Dunnett's Test
Pun $Ca_V2.1/2.3$ CT	-772.8	291.8	10	P=0.0006 (***)	
Pun $Ca_V2.1$ FT	-1438	263.9	7	P<0.0001 (****)	One-Way ANOVA
Pun $Ca_V2.1/2.3$ CT	-772.8	291.8	10	p=0.1816 (n.s.)	Dunnett's Test
Syn $Ca_V2.1/2.3$ CT	-557.7	198.9	9	Negative control	

Table8. Capacitances of calyx of Held expressing Ca_V2 hybrid channels.

Parameter	Mean	SD	n	P	statistics
Membrane capacitance C_{slow} (pF)					
Wildtype (control)	20.59	4.220	10	p=0.9246 (n.s.)	

Ca$_V$2.1 FT	23.80	5.150	11	control	
Ca$_V$2.1$^{-/-}$/ Cav2.2$^{-/-}$	22.85	8.831	11	p=0.9996 (n.s.)	
Cav2.1/2.3 II-III loop CT	21.72	6.096	8	p=0.9990 (n.s.)	
Cav2.1/2.3 II-III loop	25.76	6.045	11	p=0.9960 (n.s.)	One-Way ANOVA
Cav2.1/2.3 CT	19.06	4.894	9	p=0.6135 (n.s.)	Dunnett's Test
Ca$_V$2.1 Δ2016	27.26	3.420	6	p=0.9522 (n.s.)	
Cav2.1/2.3 CT 2016-2368	28.22	7.622	12	p=0.6103 (n.s.)	
Cav2.1/2.3 CT 1967-2368	27.47	8.513	9	p=0.8646 (n.s.)	
Cav2.1/2.2 CT	21.94	9.095	11	p=0.9991 (n.s.)	
Pun Ca$_V$2.1 FT	21.44	5.828	7	p=0.9955 (n.s.)	
Pun Ca$_V$2.1/2.2 CT	23.94	4.943	9	p>0.9999 (n.s.)	
Pun Ca$_V$2.1/2.3 CT	26.04	7.309	10	p=0.9916 (n.s.)	

777
778

780
781 **References**

782 Bewig B & Schmidt WE. (2000). Accelerated titering of adenoviruses. *Biotechniques* **28,** 870-
783   873.

784
785 Chaudhuri D, Chang SY, DeMaria CD, Alvania RS, Soong TW & Yue DT. (2004). Alternative
786   splicing as a molecular switch for Ca^{2+}/calmodulin-dependent facilitation of P/Q-type
787   Ca^{2+} channels. *J Neurosci* **24,** 6334-6342.

788
789 Chen ZX, Cooper B, Kalla S, Varoqueaux F & Young SM. (2013). The Munc13 Proteins
790   Differentially Regulate Readily Releasable Pool Dynamics and Calcium-Dependent
791   Recovery at a Central Synapse. *Journal of Neuroscience* **33,** 8336-8351.

792
793 Cole RL, Lechner SM, Williams ME, Prodanovich P, Bleicher L, Varney MA & Gu G. (2005).
794   Differential distribution of voltage-gated calcium channel alpha-2 delta (alpha2delta)
795   subunit mRNA-containing cells in the rat central nervous system and the dorsal root
796   ganglia. *J Comp Neurol* **491,** 246-269.

797
798 De Waard M, Pragnell M & Campbell KP. (1994). Ca2+ channel regulation by a conserved beta
799   subunit domain. *Neuron* **13,** 495-503.

800
801 Dolphin AC. (2012). Calcium channel auxiliary alpha2delta and beta subunits: trafficking and
802   one step beyond. *Nat Rev Neurosci* **13,** 542-555.

803
804 Dolphin AC & Lee A. (2020). Presynaptic calcium channels: specialized control of synaptic
805   neurotransmitter release. *Nat Rev Neurosci* **21,** 213-229.

806
807 Dong W, Radulovic T, Goral RO, Thomas C, Montesinos MS, Guerrero-Given D, Hagiwara A,
808   Putzke T, Hida Y, Abe M, Sakimura K, Kamasawa N, Ohtsuka T & Young SM. (2018).
809   CAST/ELKS Proteins Control Voltage-Gated Ca2+ Channel Density and Synaptic
810   Release Probability at a Mammalian Central Synapse. *Cell Rep* **24,** 284-+.

811
812 Doughty JM, Barnes-Davies M, Rusznak Z, Harasztosi C & Forsythe ID. (1998). Contrasting
813   Ca2+ channel subtypes at cell bodies and synaptic terminals of rat anterioventral
814   cochlear bushy neurones. *J Physiol* **512 (Pt 2),** 365-376.

815
816 Eggermann E, Bucurenciu I, Goswami SP & Jonas P. (2011). Nanodomain coupling between
817   Ca(2)(+) channels and sensors of exocytosis at fast mammalian synapses. *Nat Rev*
818   *Neurosci* **13,** 7-21.

819
Ferron L, Koshti S & Zamponi GW. (2021). The life cycle of voltage-gated Ca(2+) channels in
neurons: an update on the trafficking of neuronal calcium channels. *Neuronal Signal* **5,**
NS20200095.

Giugovaz-Tropper B, Gonzalez-Inchauspe C, Di Guilmi MN, Urbano FJ, Forsythe ID & Uchitel
OD. (2011). P/Q-type calcium channel ablation in a mice glycinergic synapse mediated
by multiple types of Ca(2)+ channels alters transmitter release and short term plasticity.
Neuroscience **192,** 219-230.

Hoppa MB, Lana B, Margas W, Dolphin AC & Ryan TA. (2012). alpha2delta expression sets
presynaptic calcium channel abundance and release probability. *Nature* **486,** 122-125.

Inchauspe CG, Martini FJ, Forsythe ID & Uchitel OD. (2004). Functional compensation of P/Q
by N-type channels blocks short-term plasticity at the calyx of Held presynaptic terminal.
J Neurosci **24,** 10379-10383.

Iwasaki S, Momiyama A, Uchitel OD & Takahashi T. (2000). Developmental changes in calcium
channel types mediating central synaptic transmission. *J Neurosci* **20,** 59-65.

Iwasaki S & Takahashi T. (1998). Developmental changes in calcium channel types mediating
synaptic transmission in rat auditory brainstem. *J Physiol* **509 (Pt 2),** 419-423.

Jun K, Piedras-Renteria ES, Smith SM, Wheeler DB, Lee SB, Lee TG, Chin H, Adams ME,
Scheller RH, Tsien RW & Shin HS. (1999). Ablation of P/Q-type Ca(2+) channel
currents, altered synaptic transmission, and progressive ataxia in mice lacking the
alpha(1A)-subunit. *Proc Natl Acad Sci U S A* **96,** 15245-15250.

Kaja S, Van de Ven RC, Ferrari MD, Frants RR, Van den Maagdenberg AM & Plomp JJ. (2006).
Compensatory contribution of Cav2.3 channels to acetylcholine release at the
neuromuscular junction of tottering mice. *J Neurophysiol* **95,** 2698-2704.

Kamp MA, Krieger A, Henry M, Hescheler J, Weiergraber M & Schneider T. (2005). Presynaptic
'Ca$_V$2.3-containing' E-type Ca^{2+} channels share dual roles during neurotransmitter
release. *Eur J Neurosci* **21,** 1617-1625.

Kamp MA, Shakeri B, Tevoufouet EE, Krieger A, Henry M, Behnke K, Herzig S, Hescheler J,
Radhakrishnan K, Parent L & Schneider T. (2012). The C-terminus of human Ca$_V$2.3
voltage-gated calcium channel interacts with alternatively spliced calmodulin-2
expressed in two human cell lines. *Biochim Biophys Acta* **1824,** 1045-1057.

Lee A, Zhou H, Scheuer T & Catterall WA. (2003). Molecular determinants of
Ca(2+)/calmodulin-dependent regulation of Ca(v)2.1 channels. *Proc Natl Acad Sci U S A*
100, 16059-16064.

Lubbert M, Goral RO, Keine C, Thomas C, Guerrero-Given D, Putzke T, Satterfield R,
Kamasawa N & Young SM, Jr. (2019). Ca(V)2.1 alpha(1) Subunit Expression Regulates

Presynaptic Ca(V)2.1 Abundance and Synaptic Strength at a Central Synapse. *Neuron* **101,** 260-273 e266.

Lubbert M, Goral RO, Satterfield R, Putzke T, van den Maagdenberg AM, Kamasawa N & Young SM, Jr. (2017). A novel region in the Ca(V)2.1 alpha(1) subunit C-terminus regulates fast synaptic vesicle fusion and vesicle docking at the mammalian presynaptic active zone. *Elife* **6**.

Mochida S, Few AP, Scheuer T & Catterall WA. (2008). Regulation of presynaptic Ca(V)2.1 channels by Ca2+ sensor proteins mediates short-term synaptic plasticity. *Neuron* **57,** 210-216.

Mochida S, Westenbroek RE, Yokoyama CT, Zhong H, Myers SJ, Scheuer T, Itoh K & Catterall WA. (2003). Requirement for the synaptic protein interaction site for reconstitution of synaptic transmission by P/Q-type calcium channels. *Proc Natl Acad Sci U S A* **100,** 2819-2824.

Montesinos MS, Chen ZX & Young SM. (2011). pUNISHER: a high-level expression cassette for use with recombinant viral vectors for rapid and long term in vivo neuronal expression in the CNS. *J Neurophysiol* **106,** 3230-3244.

Montesinos MS, Satterfield R & Young SM, Jr. (2016). Helper-Dependent Adenoviral Vectors and Their Use for Neuroscience Applications. *Methods Mol Biol* **1474,** 73-90.

Mori MX, Vander Kooi CW, Leahy DJ & Yue DT. (2008). Crystal structure of the CaV2 IQ domain in complex with Ca2+/calmodulin: high-resolution mechanistic implications for channel regulation by Ca2+. *Structure* **16,** 607-620.

Muller CS, Haupt A, Bildl W, Schindler J, Knaus HG, Meissner M, Rammner B, Striessnig J, Flockerzi V, Fakler B & Schulte U. (2010). Quantitative proteomics of the Cav2 channel nano-environments in the mammalian brain. *Proc Natl Acad Sci U S A* **107,** 14950-14957.

Nanou E & Catterall WA. (2018). Calcium Channels, Synaptic Plasticity, and Neuropsychiatric Disease. *Neuron* **98,** 466-481.

Nanou E, Yan J, Whitehead NP, Kim MJ, Froehner SC, Scheuer T & Catterall WA. (2016). Altered short-term synaptic plasticity and reduced muscle strength in mice with impaired regulation of presynaptic CaV2.1 Ca2+ channels. *Proc Natl Acad Sci U S A* **113,** 1068-1073.

Palmer DJ & Ng P. (2005). Helper-dependent adenoviral vectors for gene therapy. *Hum Gene Ther* **16,** 1-16.

910   Pilch KS, Ramgoolam KH & Dolphin AC. (2022). Involvement of Ca(V) 2.2 channels and
911   alpha(2) delta-1 in homeostatic synaptic plasticity in cultured hippocampal neurons. *J*
912   *Physiol* **600,** 5333-5351.

913
914   Radulovic T, Dong W, Goral RO, Thomas CI, Veeraraghavan P, Montesinos MS, Guerrero-
915   Given D, Goff K, Lubbert M, Kamasawa N, Ohtsuka T & Young SM. (2020). Presynaptic
916   development is controlled by the core active zone proteins CAST/ELKS. *J Physiol-*
917   *London* **598,** 2431-2452.

918
919   Rajapaksha WR, Wang D, Davies JN, Chen L, Zamponi GW & Fisher TE. (2008). Novel splice
920   variants of rat CaV2.1 that lack much of the synaptic protein interaction site are
921   expressed in neuroendocrine cells. *J Biol Chem* **283,** 15997-16003.

922
923   Rothman JS & Silver RA. (2018). NeuroMatic: An Integrated Open-Source Software Toolkit for
924   Acquisition, Analysis and Simulation of Electrophysiological Data. *Front Neuroinform* **12,**
925   14.

926
927   Serra SA, Gene GG, Elorza-Vidal X & Fernandez-Fernandez JM. (2018). Cross talk between
928   beta subunits, intracellular Ca(2+) signaling, and SNAREs in the modulation of Ca(V) 2.1
929   channel steady-state inactivation. *Physiol Rep* **6**.

930
931   Simms BAZ, G. W. (2014). Neuronal voltage-gated calcium channels: structure, function, and
932   dysfunction. *Neuron* **82,** 24-45.

933
934   Stephani F, Scheuer V, Eckrich T, Blum K, Wang W, Obermair GJ & Engel J. (2019). Deletion
935   of the Ca(2+) Channel Subunit alpha(2)delta3 Differentially Affects Ca(v)2.1 and
936   Ca(v)2.2 Currents in Cultured Spiral Ganglion Neurons Before and After the Onset of
937   Hearing. *Front Cell Neurosci* **13,** 278.

938
939   Szabo Z, Obermair GJ, Cooper CB, Zamponi GW & Flucher BE. (2006). Role of the synprint
940   site in presynaptic targeting of the calcium channel CaV2.2 in hippocampal neurons. *Eur*
941   *J Neurosci* **24,** 709-718.

942
943   Voigt A, Freund R, Heck J, Missler M, Obermair GJ, Thomas U & Heine M. (2016). Dynamic
944   association of calcium channel subunits at the cellular membrane. *Neurophotonics* **3,**
945   041809.

946
947   Weiss N & Zamponi GW. (2017). Trafficking of neuronal calcium channels. *Neuronal Signal* **1,**
948   NS20160003.

949
Wiser O, Cohen R & Atlas D. (2002). Ionic dependence of Ca2+ channel modulation by syntaxin
1A. *Proc Natl Acad Sci U S A* **99,** 3968-3973.

Xie M, Li X, Han J, Vogt DL, Wittemann S, Mark MD & Herlitze S. (2007). Facilitation versus
depression in cultured hippocampal neurons determined by targeting of Ca2+ channel
Cavbeta4 versus Cavbeta2 subunits to synaptic terminals. *J Cell Biol* **178,** 489-502.

Yokoyama CT, Myers SJ, Fu J, Mockus SM, Scheuer T & Catterall WA. (2005). Mechanism of
SNARE protein binding and regulation of Cav2 channels by phosphorylation of the
synaptic protein interaction site. *Mol Cell Neurosci* **28,** 1-17.

Young SM, Jr. & Neher E. (2009). Synaptotagmin has an essential function in synaptic vesicle
positioning for synchronous release in addition to its role as a calcium sensor. *Neuron*
63, 482-496.

Young SM, Jr. & Veeraraghavan P. (2021). Presynaptic voltage-gated calcium channels in the
auditory brainstem. *Mol Cell Neurosci* **112,** 103609.

Zhong H, Yokoyama CT, Scheuer T & Catterall WA. (1999). Reciprocal regulation of P/Q-type
Ca2+ channels by SNAP-25, syntaxin and synaptotagmin. *Nat Neurosci* **2,** 939-941.

Additional information section

Conflict of Interest – the authors report no conflict of interest.

Author contributions: J.L. and P.V., carried out experiments and analyzed data. S.M.Y, Jr developed the mouse model S.M.Y, Jr planned the project and analyzed data. J.L. and S.M.Y, Jr. wrote the manuscript. All authors jointly revised the paper. All authors approved the final version of the manuscript, agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved, all persons designated as authors qualify for authorship, and all those who qualify for authorship are listed.

Funding sources: This work was supported by the National Institutes of Deafness and Communication Disorders (R01 DC014093), the National Institute of Neurological Disorders and Stroke (R01 NS110742), and the National Center for Advancing Translational Sciences. R03TR004161-01 and the University of Iowa Distinguished Scholar Award. (S.M.Y., Jr) and the American Heart Association (Predoctoral fellowship 23PRE1019152) (J. L)

Acknowledgements: We thank members of the Young lab for their comments and critiques of the manuscript and project. We thank Hao Li for help with for HdAd viral vector production. We thank Dr. Susan Stamnes and University of Iowa viral vector core, for help with HdAd vector production and tittering. We thank Dr. Phillip Ng and Dr. Brendan Lee for gifts of HdAd packing plasmids and HdAd stuffer DNA, respectively.

1000

1001

1002

1003

1004

1005

.

Fig. 1

1010

Fig. 2

1011

1012

1013

Fig. 3

Fig. 4

1015

1016

Fig. 5

A Alignment of Ca$_V$2.1 and Ca$_V$2.3 CT

B Ca$_V$2.1 full transcript   Ca$_V$2.1 delta2016

0.5nA
10ms

C ns

D Ca$_V$2.1Δ2016   Ca$_V$2.1/2.3 CT aa. 2016-2368   Ca$_V$2.1/2.3 CT aa. 1967-2368   Ca$_V$2.1/2.3 CT aa. 1933-2368

E Comparison to Ca$_V$2.1Δ2016

F Comparison to Ca$_V$2.1/2.3 CT aa. 1933-2368

Fig. 6

1018

1019

Fig. 7

A

B

1020