Title:

EBF1 limits the numbers of cochlear hair and supporting cells and forms the scala tympani

and spiral limbus during inner ear development

Abbreviated title: EBF1 controls the development of the cochlea

Author names:

Hiroki Kagoshima[1], Hiroe Ohnishi[1], Ryosuke Yamamoto[2], Akiyoshi Yasumoto[1], Yosuke Tona[1],

Takayuki Nakagawa[1], Koichi Omori[1], and Norio Yamamoto[1,3]

[1]Department of Otolaryngology, Head and Neck Surgery, Graduate School of Medicine, Kyoto

University, Kyoto, Japan

[2]Biological Sciences, Sunnybrook Research Institute, Sunnybrook Health Sciences Centre,

Toronto, ON, Canada.

[3]Department of Otolaryngology, Kobe City Medical Center General Hospital, Kobe, Japan.

Corresponding Author's name: Norio Yamamoto

Affiliation: Department of Otolaryngology, Kobe City Medical Center General Hospital

E-mail:2-1-1 Minatojimaminami-cho, Chuo-ku, Kobe, 650-0047, Japan

E-mail address: yamamoto@ent.kuhp.kyoto-u.ac.jp

Conflicts of Interest: The authors declare that they have no conflicts of interest.

Abstract

Early B-cell factor 1 (EBF1) is a basic helix-loop-helix transcription factor essential for the differentiation of various tissues. Our single-cell RNA sequencing data suggest that *Ebf1* is expressed in the sensory epithelium of the inner ear. Here, we found that the *Ebf1* gene and its protein are expressed in the prosensory domain of the inner ear, medial region of the cochlear floor, inner ear mesenchyme, and cochleo-vestibular ganglion. *Ebf1* deletion results in incomplete formation of the spiral limbus and scala tympani, increased number of cells in the organ of Corti and Kölliker's organ, and aberrant course of the spiral ganglion axons. The inner ear-specific deletion of *Ebf1* revealed that scala tympani formation depended on *Ebf1* expressed in the otic mesenchyme. *Ebf1* deletion in the cochlear epithelia caused the proliferation of SOX2-positive cochlear cells at E13.5, indicating that EBF1 suppresses the proliferation of the prosensory domain and cells of Kölliker's organ to facilitate the development of appropriate numbers of hair and supporting cells. Furthermore, mice with deletion of cochlear epithelium-specific *Ebf1* showed poor postnatal hearing function. Our results suggest that *Ebf1* is essential for normal auditory function in mammals.

Introduction

The inner ear is a unique and complex organ that consists of bony and membranous labyrinths. The membranous labyrinth contains multiple sensory organs, including the cochlea, which is responsible for hearing and comprises three compartments (scalae): the scala vestibuli, scala tympani, and scala media. The scala vestibuli and scala tympani develop from the mesenchyme surrounding the inner ear (Sher, 1971). The scala media is situated between the scala vestibuli and scala tympani, which are separated from the scala media by the tectorial and basilar

membranes, respectively. Moreover, the scala media contains sensory epithelia that transduce

sound into electrical signals via specialized sensory cells known as hair cells. Cochlear hair cells

are located within the organ of Corti in the middle part of the scala media epithelium, and consist

of one row of inner hair cells and three rows of outer hair cells. These hair cells are surrounded

by at least seven types of non-sensory supporting cells: inner boundary cells, inner phalangeal

cells, Deiters' cells, inner and outer pillar cells, Hensen's cells, and Claudius cells. The precise

number and placement of mechanosensory hair cells and non-sensory supporting cells enable the

accurate reception of mechanical stimulation of sound or body motion and its conversion into

neural signals.

Inner ear development in mice begins with the formation of an ectodermal thickening called the

otic placode, which is located adjacent to the hindbrain (Wu and Kelley, 2012). The otic placode

invaginates to form a spherical structure called an otocyst at approximately embryonic day (E)

9.5. The ventral side of the otocyst forms the future sensory epithelium, where the sex-

determining region Y-box transcription factor 2 (*Sox2*) is expressed (Kiernan et al., 2005). At

E10.5, the cochlear and endolymphatic ducts and semicircular canals begin to form on the

ventral and dorsal sides of the otocyst, respectively. After ventral cochlear duct formation

becomes evident at E11, the ventral side of the cochlear duct (cochlear duct floor) begins to

develop into a future sensory domain by expressing *Sox2* and *Jagged1* at E11.5 (Wu and Kelley,

2012). The *Sox2*-positive region becomes limited to the middle part of the ventral cochlear duct

and is recognized as a prosensory domain at E13.5 and E14.5 (Kiernan et al., 2005; Ohyama et

al., 2010). This region is located between two molecularly distinct nonsensory areas. The region

medial to the prosensory domain toward the axis of the cochlea (modiolus) is called the greater

epithelial ridge (GER) and transiently contains Kölliker's organ, which is composed of columnar

supporting cells during the developmental stage and becomes the inner sulcus with cuboidal cells

and the spiral limbus with interdental cells in the mature cochlea (Dayaratne et al., 2014).

Additionally, the GER is a source of sensory epithelia and has the potential to produce sensory

cells after the establishment of hair cells (Kubota et al., 2021).

The complex cellular structure and developmental processes of the inner ear depend on the

highly regulated expression patterns of signaling molecules and transcription factors. However,

the mechanisms underlying inner ear development are not fully understood. To comprehensively

elucidate these mechanisms, we analyzed the single-cell RNA-seq data of the inner ear epithelial

cells. In this study, we found that the early B-cell factor 1 gene (*Ebf1*) was upregulated in

clusters of sensory epithelial progenitors in the cochlea, macula, and crista, and confirmed that it

was expressed on the medial side of the cochlear duct floor, the prosensory area of the macula

and crista, and the spiral ganglion (Yamamoto et al., 2021).

EBF1 belongs to the EBF family of transcription factors and encodes four paralogous genes in

mammals, including mice and humans (Liberg et al., 2002). EBF family proteins comprise a

functional DNA-binding domain, transcription factor immunoglobulin, helix-loop-helix (HLH)

domain, and transactivation domain (Hagman et al., 1995), which are the characteristics of basic

HLH (bHLH) transcription factors. Similar to other bHLH transcription factors, EBF1 is

involved in various developmental processes, including the determination of cell fate and

differentiation of B lymphocytes and olfactory epithelia (Liberg et al., 2002).

Considering the various roles of EBF1 as a bHLH transcription factor, the importance of bHLH

transcription factors—such as ATOH1—in inner ear development, and the results of our

previous study confirming *Ebf1* expression in the inner ear sensory epithelium, we analyzed the

function of EBF1 in inner ear development. In the present study, we confirmed the

spatiotemporal expression of *Ebf1* during inner ear development and examined the effects of

Ebf1 deletion on inner ear development and hearing.

Results

Ebf1 is expressed in developing mouse cochlea

In silico analysis of embryonic inner ear epithelia suggested that *Ebf1* is predominantly

expressed in the inner ear sensory epithelium during early development (Yamamoto et al., 2021).

To quantify *Ebf1* expression at each stage of inner ear development, we performed quantitative

reverse transcription polymerase chain reaction (qRT-PCR) using whole embryonic inner ears

from E9.5 to P0 (Fig. 1A). The expression of *Ebf1* mRNA transcripts began to increase at E10.5

and reached a maximum at E13.5 (Fig. 1A). The relative expression level at E13.5 was

approximately 10-fold higher than that at E9.5. The expression level then decreased but

remained 7.5 times higher than that at E9.5 even at P0.

To describe the spatiotemporal expression patterns of *Ebf1* during inner ear development, we

performed *in situ* hybridization (ISH) and immunohistochemistry (IHC) analyses on sections of

the developing inner ear of wild-type mice from E9.5 to P0 (Fig. 1 B–D, Supplemental Fig. S1).

We stained *Sox2*, which is expressed in the sensory progenitor region of the inner ear from the

early developmental stages (Kiernan et al., 2005), as well as *Ebf1* on adjacent sections to specify

the location of *Ebf1* expression, and compared the expression of the two genes and their products.

First, we examined *Ebf1* expression at E13.5 (Fig. 1B), which is when the sensory epithelium of

the inner ear forms and *Ebf1* expression level is maximized during inner ear development (Fig.

1A). *Ebf1* was expressed on the medial side of the cochlear floor, including the prosensory

domain (white arrows in Fig. 1B), spiral ganglion (white asterisks in Fig. 1B), otic mesenchyme

(black asterisks in Fig. 1B), and parts of the prosensory regions of the vestibule and crista (black

arrows in Fig. 1B). Compared with *Sox2*, *Ebf1* was expressed more medially within the cochlear

floor, which developed into Kölliker's organ and the organ of Corti, and its expression in the

vestibule was more restricted (Fig. 1B).

Subsequently, we examined the spatiotemporal expression of *Ebf1* throughout inner ear

development, including the onset of expression in the inner ear epithelium, using inner ear

sections from E9.5 to P0 (Fig. 1C, Supplemental Fig. S1). At E9.5, *Ebf1* was not expressed in the

otocyst but was expressed in the progenitor cells of the cochleo-vestibular ganglion (CVG) (data

not shown), which delaminate from the ventral side of the otocyst into the otic mesenchyme (Wu

and Kelley, 2012). At E10.5, *Ebf1* expression was observed on the ventromedial side of the

otocyst, which develops into the cochlear duct, and in the ventrolateral epithelium of the otocyst,

which develops into the crista (black arrowheads in Fig. 1C). Additionally, *Ebf1* expression was

detected in the otic mesenchyme (black asterisk in Fig. 1C) and CVG (white asterisk in Fig. 1C),

which persisted until later stages (Supplemental Fig. S1). *Ebf1* was expressed in the border

region, where the cochlear duct begins to elongate, at E11.5, in the medial side of the cochlear

floor at E12.5, and in the future crista region in the vestibule at E11.5 and 12.5 (Supplemental

Fig. S1). In the cochlea at E16.5 and E18.5, *Ebf1* was expressed throughout the organ of Corti

and Kölliker's organ, whereas *Sox2* was expressed in the organ of Corti and the lateral half of

Kölliker's organ (Fig. 1C and Supplemental Fig. S1), consistent with a previous report (Urness et

al., 2015). *Ebf1* was expressed in the spiral ligaments, tympanic border cells (white arrowheads

in Fig. 1C) (Taniguchi et al., 2012), vestibules, and crista (Fig. 1C). *Ebf1* expression was

maintained until P0 (Supplemental Fig. S1).

IHC analysis showed that EBF1 was expressed throughout Kölliker's organ and the prosensory

domain, whereas SOX2 was expressed in a part of Kölliker's organ and the prosensory domain

(upper panels of Fig. 1D), which is consistent with the ISH results. The disappearance of the

EBF1 signal from the cochlear epithelia and mesenchyme in conventional *Ebf1* knockout (*Ebf1$^{-/-}$*

) mice confirmed the specificity of the anti-EBF1 antibody used in this study (Fig. 1D).

Ebf1 deletion altered the structure of the cochlear duct

Our ISH and IHC analyses, which showed the expression of *Ebf1* and its protein in both the

developing inner ear epithelia and mesenchyme, suggest that *Ebf1* is involved in the

development of both the inner ear sensory epithelium and otic mesenchyme. We used two

mutant mouse strains to examine the roles of *Ebf1* in developing inner ears: an *Ebf1* conventional

knockout (*Ebf1$^{-/-}$*) mouse (Lin and Grosschedl, 1995) and a Foxg1-Cre-mediated inner ear

epithelia-specific conditional knockout mouse (*Foxg1Cre;Ebf1$^{fl/fl}$*) (Hébert and McConnell,

2000; Gyory et al., 2012) in which *Ebf1* expression remains localized in the inner ear

mesenchyme.

Comparison of the gross morphology of the membranous labyrinth of the inner ear at E18.5

revealed no difference between *Ebf1$^{+/+}$* and *Ebf1$^{+/-}$* mice (Fig. 2A). However, after removing the

lateral wall and roof of the cochlea to expose the cochlear floor, we found that *Ebf1$^{-/-}$* mice had

fewer turns in the cochlear duct than *Ebf1$^{+/+}$* mice (white arrowhead in Fig. 2B).

Hematoxylin-eosin (HE) staining of the cochlea at E18.5 revealed incomplete formation of the

scala tympani, particularly in the middle and apical turns of the cochlea of *Ebf1$^{-/-}$* mice compared

with those of control mice (arrowheads in Fig. 2C). Moreover, *Ebf1$^{-/-}$* mice lacked a spiral limbus

(sl; Fig. 2C); a lower number of cochlear turns was also observed in the cochlear sections of

these mice (arrows in the upper right panels in Fig. 2C), supporting the gross morphological

observations.

Ebf1 was expressed in the otic mesenchyme from early to later developmental stages. To

elucidate whether the hypoplastic scala tympani, fewer cochlear turns, and lack of spiral limbus

were caused by the *Ebf1*-deficient mesenchyme, we examined the cochlear morphology of

Foxg1Cre;Ebf1$^{fl/fl}$ and *Foxg1Cre;Ebf1$^{fl/+}$* mice via HE staining. In contrast to the hypoplastic

scala tympani of *Ebf1$^{-/-}$* mice, the scala tympani of *Foxg1Cre;Ebf1$^{fl/fl}$* mice was completely

formed throughout the cochlear turns (left panels of Fig. 2C). The number of cochlear turns in

Foxg1Cre;Ebf1$^{fl/fl}$ mice was similar to that in the control mice (*Foxg1Cre;Ebf1$^{fl/+}$* mice).

However, *Foxg1Cre;Ebf1$^{fl/fl}$* mice lacked a spiral limbus, as observed in *Ebf1$^{-/-}$* mice (Fig. 2C).

These results suggest that *Ebf1* controls scala tympani formation and the turning of the cochlear

duct through the mesenchyme, and the lack of the spiral limbus in *Ebf1*-deficient mice is

secondarily caused by EBF1 malfunction within the epithelia.

Ebf1 deletion caused an increase in the number of cochlear hair, supporting, and Kölliker's

organ cells

Observation of the cochlear epithelia in HE-stained samples revealed that both *Ebf1$^{-/-}$* and

Foxg1Cre;Ebf1$^{fl/fl}$ mice had deformed Kölliker's organs and organs of Corti (Ko and oC in Fig.

2C). To examine these phenotypes more comprehensively, we performed IHC analysis on inner

ear sections and cochlear whole-mount samples from E18.5 (Fig. 3 and Supplemental Figs. S2

and S3).

IHC analysis of cochlear sections showed that *Ebf1* deletion increased the number of MYO7A-

positive hair cells as well as SOX2-positive supporting and Kölliker's organ cells from the basal

to the apical region at E18.5 (Fig. 3A). An increase in the number of SOX2-positive cells within

the medial region of the cochlear floor was also observed at E14.5 (arrowhead in Fig. 1D). In

contrast, *Ebf1* deletion had no morphological effects on the vestibular sensory epithelium

(Supplemental Fig. S2). Additionally, the apical region of *Ebf1*$^{-/-}$ mouse cochleae contained

multiple layers of SOX2-positive cells (asterisk in Fig. 3A). IHC analysis of whole-mount

cochlear samples showed that *Ebf1*$^{-/-}$ mice had an increased number of MYO7A-positive hair

cells (Fig. 3B) and ectopic MYO7A-positive cells within the GER (arrows in Fig. 3B). Although

the normal cochlea has one and three rows of inner and outer hair cells, respectively, the mutant

cochlea had eight to nine rows of hair cells. We found ectopic hair cells in 7 of the 12 examined

cochleae. These ectopic MYO7A-positive cells contained stereocilia-like structures, as indicated

by phalloidin staining (arrows in Fig. 3C). Increased numbers of supporting cells were confirmed

in whole-mount cochlear samples by IHC staining of SOX2 (Fig. 3D), a supporting and

Kölliker's organ cell marker, and PROX1 (Fig. 3E), a pillar and Deiters' cell marker

(Bermingham-McDonogh et al., 2006).

To quantify the number of hair and supporting cells, we counted the cells in three regions within

the cochlea (basal, middle, and apical regions; Fig. 4A) and measured the number of MYO7A- or

PROX1-positive cells per 200 μm at E18.5. For SOX2-positive cells, we counted the cell number

per 100 μm only in the basal regions at E18.5. The results showed that the cochlear hair cell

number was significantly increased in *Ebf1*$^{-/-}$ mice compared to *Ebf1*$^{+/+}$ mice in all three regions

(Fig. 4B). *Ebf1*$^{+/-}$ mice exhibited a significantly higher number of cochlear hair cells than *Ebf1*$^{+/+}$

mice in the middle and apical regions (Fig. 4B, $p < 0.0001$). The hair cell numbers per 200 μm of

Ebf1$^{+/+}$, *Ebf1*$^{+/-}$, and *Ebf1*$^{-/-}$ mice were 142.3 ± 9.0, 150.0 ± 2.8, and 367.5 ± 30.8 in the basal

regions; 152.8 ± 8.1, 190.3 ± 7.0, and 363.0 ± 22.4 in the middle regions; and 154.5 ± 5.1, 183.5

± 6.8, and 440.3 ± 23.8 in the apical regions, respectively.

SOX2-positive cells, constituting a part of Kölliker's organ cells and supporting cells within

organs of Corti, also increased in number by 1.7 times in $Ebf1^{-/-}$ mice compared to $Ebf1^{+/+}$ mice

(Fig. 4C). PROX1-positive cell numbers significantly increased in $Ebf1^{-/-}$ mice compared to

$Ebf1^{+/+}$ mice only in the basal and middle regions (Fig. 4D). In the apical region, the number of

PROX1-positive cells was similar to that in $Ebf1^{-/-}$ and $Ebf1^{+/+}$ mice. The numbers of PROX1-

positive cells in $Ebf1^{+/+}$ and $Ebf1^{-/-}$ mice were 203.3 ± 2.2 and 439.0 ± 15.4, 228.0 ± 16.8 and

389.8 ± 10.4, and 196.0 ± 27.3 and 204.5 ± 19.6 in the basal, middle, and apical regions,

respectively.

We performed IHC analysis using more specific markers to reveal which population of hair or

supporting cells increased in number in $Ebf1^{-/-}$ mice (Supplemental Fig. S3). We immunostained

whole-mounted cochlea at E18.5 with anti-VGLUT3 and anti-p75 antibodies, which indicate

inner hair (Li et al., 2018) and pillar cells (von Bartheld et al., 1991), respectively. VGLUT3-

positive inner hair cells, which are arranged in a single row in wild-type mice, were found to be

increased in number in $Ebf1^{-/-}$ mice (Supplemental Fig. S3A), indicating an increase in both inner

and outer hair cells. The number of p75-positive pillar cells, which separate inner and outer hair

cells, did not increase in $Ebf1^{-/-}$ mice (Supplemental Fig. S3B). Considering that PROX1-positive

cells indicate pillar and Deiters' cells, the number of Deiters' cells increased in $Ebf1^{-/-}$ mice.

However, the arrangement of pillar cells was disrupted in $Ebf1^{-/-}$ mice (arrows in Supplemental

Fig. S3B), which was reflected in the appearance of VGLUT3 cells in the outer hair cell region

of $Ebf1^{-/-}$ mice (arrowheads in Supplemental Fig. S3A).

The total cochlear length, measured based on the length of the MYO7A-positive region

(Supplemental Fig. S4A), was slightly, but significantly, shorter in $Ebf1^{-/-}$ mice than in $Ebf1^{+/+}$

and $Ebf1^{+/-}$ mice (Supplemental Fig. S4B, $p < 0.001$).

Ebf1 deletion caused the aberrant spiral ganglion and nerve fibers

As *Ebf1* is expressed in the spiral ganglion and the number of hair cells, a target of the spiral

ganglion cell axon, increased in $Ebf1^{-/-}$ mice, we examined the spiral ganglion morphology and

innervation of cochlear hair cells with IHC using anti-Tubulin β 3 (TUJ1) antibodies at E18.5

(Supplemental Fig. S5).

Compared with $Ebf1^{+/+}$ mice, which exhibited axons extending from the spiral ganglion cells to

the cochlear hair cells, $Ebf1^{-/-}$ mice had spiral ganglion cells (sg in Supplemental Fig. S5A) under

the organs of Corti (arrows in Supplemental Fig. S5A), as well as in their normal position. The

axons, which usually run parallel to the rows of outer hair cells, formed a reticulation within the

$Ebf1^{-/-}$ mouse cochlear hair cell regions (Supplemental Fig. S5B). Moreover, the innervation

reached Kölliker's organ (arrowheads and brackets in Supplemental Fig. S5A) as well as the

organ of Corti.

Ebf1 deletion changed the distribution of JAG1-positive Kölliker's organ cells, the

differentiation timing of a prosensory domain, and the proliferation of SOX2-positive cells

As $Ebf1^{-/-}$ mice had increased numbers of cochlear hair cells, which were differentiated from the

prosensory domain, we investigated its specification, differentiation, proliferation, and cell death

in the $Ebf1^{-/-}$ mouse cochlear duct floor.

First, to determine whether formation of the prosensory domain was affected by *Ebf1* deletion,

we examined the formation of regions medial and lateral to the prosensory domain. Because

these regions express FGF10 and BMP4 to induce non-sensory or sensory epithelia in the

cochlear floor (Ohyama et al., 2010; Urness et al., 2015), respectively, we performed ISH for

Fgf10 and *Bmp4* in the basal region of the cochlear duct of *Ebf1*$^{+/+}$ *and Ebf1*$^{-/-}$ mice at E13.5 (Fig.

5A). The formation of both regions was similar in *Ebf1*$^{+/+}$ and *Ebf1*$^{-/-}$ mice, suggesting that *Ebf1*

is not involved in the development of cell populations expressing *Fgf10* or *Bmp4*. To verify the

medial cell population more precisely, we immunostained E13.5 and E14.5 cochleae with an

anti-JAG1 antibody (Fig. 5A), as JAG1 is exclusively expressed in Kölliker's organ, a part of the

medial region, at this stage (Ohyama et al., 2010). JAG1-positive cells in *Ebf1*$^{-/-}$ mouse cochlea

expanded to the more medial region compared with those in *Ebf1*$^{+/+}$ mouse cochlea at E13.5 and

E14.5 (arrowheads in Fig. 5A).

Subsequently, we examined the expression of *Atoh1* within the prosensory domain via ISH at

E14.5 and E15.5 (Fig. 5B), as *Atoh1* is necessary for hair cells to differentiate from the

prosensory cell population (Bermingham et al., 1999) and its expression indicates the initiation

of hair cell development from the prosensory domain. Compared to *Ebf1*$^{+/+}$ mice that expressed

Atoh1 within the prosensory domain from E14.5, *Atoh1* mRNA was not detected in the basal to

middle region of the E14.5 *Ebf1*$^{-/-}$ mouse cochlea, although the vestibular organs expressed

Atoh1 within the prosensory epithelia. However, E15.5 *Ebf1*$^{-/-}$ mice exhibited an *Atoh1* signal

within the prosensory domain of the cochlea. This result suggests that while the cell fate

specification of sensory epithelia in the cochlea is not affected, its timing is delayed by *Ebf1*

deletion. Considering that the differentiation of cochlear sensory epithelia promotes the transition

from the basal to apical turns of the cochlea (Sher, 1971), the expression of hair and supporting

cell markers at a later stage, E18.5, also indicated delayed differentiation of *Ebf1*^{-/-} mouse

cochleae (Supplemental Fig. S3). These markers were found to be detected in more basal

cochlear regions in *Ebf1*^{-/-} mice than in *Ebf1*^{+/+} mice.

The expansion of the JAG1-positive cell area and the increased numbers of hair and supporting

cells suggest that the enhancement of proliferation or suppression of cell death occurs within the

prosensory domain and Kölliker's organ of *Ebf1*^{-/-} mouse cochlea. To identify the mechanisms

that correlate with the functions of EBF1 within the cochlea, we tested the proliferation and

apoptotic status of *Ebf1*^{-/-} mouse cochlea. To evaluate the proliferation status of the prosensory

domain and Kölliker's organ, we immunostained cochlear sections with SOX2, a marker of the

prosensory domain and a part of Kölliker's organ, and 5-ethynyl-2′-deoxyuridine (EdU) at E12.5,

13.5, 14.5, and 16.5 after administering EdU to pregnant mice (Fig. 6A). Quantification of

SOX2-positive cells showed that their number decreased in *Ebf1*^{+/+} mice from E12.5 onward

(Fig. 6B), and their location was limited to the prosensory domain (brackets in Fig. 6A). In

contrast, SOX2-positive cells in *Ebf1*^{-/-} mouse cochlea were found both in the prosensory domain

and the medial region, even at E13.5, which was consistent with the results of JAG1

immunostaining (Fig. 5A). The number of SOX2-positive cells in *Ebf1*^{-/-} mice was similar to that

in *Ebf1*^{+/+} mice at E12.5, but increased at E13.5 and returned to the E12.5 level at E14.5.

Therefore, the number of SOX2-positive cells in *Ebf1*^{-/-} mice was significantly higher than those

in *Ebf1*^{+/+} mice at E13.5 and E14.5 (Fig. 6B). The number of EdU-positive proliferating cells

within SOX2-positive cells was significantly higher in *Ebf1*^{-/-} mice than in *Ebf1*^{+/+} mice at E13.5

and E14.5 (Fig. 6C). As the number of SOX2-positive cells increased in *Ebf1*^{-/-} mice after E13.5

(Fig. 6B), normalization of SOX2-positive cell numbers was necessary to correctly evaluate the

proliferation status of SOX2-positive cells. We calculated the proportion of EdU-positive cells

among SOX2-positive cells and found that the proliferation was enhanced in the SOX2-positive cells of $Ebf1^{-/-}$ mouse cochlea only at E13.5 (49.8 ± 4.3%) compared with that in $Ebf1^{+/+}$ mouse cochlea (37.8 ± 4.2%) (Fig. 6D). These results suggested that EBF1 suppressed the proliferation of SOX2-positive cells within a limited time window. Morphologically, a difference in proliferation was observed in the prosensory domain, as indicated by EdU immunostaining (brackets at E13.5; Fig. 6A). EdU staining was observed in the Kölliker's organs of $Ebf1^{-/-}$ mice, even at E16.5, but not observed in $Ebf1^{+/+}$ mice (E16.5 of Fig. 6A).

The loss of proliferation within the prosensory domain around E13.5 (bracket in the $Ebf1^{+/+}$ sample at E13.5, Fig. 6A) has been well documented in previous studies (Chen and Segil, 1999; Chen et al., 2002). The post-mitotic domain is called the zone of non-proliferating cells (ZNPC), and is characterized by the expression of the cyclin-dependent kinase inhibitor p27Kip1. To confirm the EdU immunostaining results, we performed p27Kip1 immunostaining at E13.5 and E14.5 (Fig. 6E). Although p27Kip1 was detected in the prosensory domain of $Ebf1^{+/+}$ mice at E13.5, it was not expressed in $Ebf1^{-/-}$ mouse cochlea at this stage (arrows in Fig. 6E), which was consistent with the results of EdU detection. At E14.5, p27Kip1 was detected in a larger area of the middle part of $Ebf1^{-/-}$ mouse cochlear floors than in those of $Ebf1^{+/+}$ mice.

In contrast, apoptosis within the inner ear or cochlear duct did not increase in $Ebf1^{-/-}$ mice at E11.5 and E13.5, compared with that in $Ebf1^{+/+}$ mice (Supplemental Fig. S6).

Ebf1 deletion impairs auditory function

The aberrant cochlear sensory epithelia observed in *Ebf1*-deleted mice suggest that hearing ability is impaired in these mice. To evaluate the effect of *Ebf1* deletion on auditory function, we measured the auditory brainstem response (ABR) (Fig. 7A) and distortion product of otoacoustic

emissions (DPOAE) (Fig. 7B) in P21 *Foxg1Cre;Ebf1*^{fl/fl} mice. We did not use *Ebf1*^{-/-} mice for

this analysis to avoid embryonic lethality and to eliminate the effects of the hypoplastic scala

tympani observed in *Ebf1*^{-/-} mice on auditory function. The phenotype of *Foxg1Cre;Ebf1*^{fl/fl} mice

was evaluated using whole-mount cochlear samples collected at P23 (Fig. 7C). We observed a

marked increase in the number of cochlear hair cells in *Foxg1Cre;Ebf1*^{fl/fl} mice, comparable to

the morphology of E18.5 *Ebf1*^{-/-} mice (Fig. 3B and Fig. 7C).

ABR measurement showed significant elevations of thresholds of the response to sound in

Foxg1Cre;Ebf1^{fl/fl} mice at all frequencies examined (10 kHz, 93.3 ± 2.5 dB; 20 kHz, 86.7 ± 3.8

dB; 40 kHz, 105.0 ± 0.0 dB) (Fig. 7A) compared with control mice, indicating severe hearing

loss in *Ebf1*-deleted mice. Subsequently, we performed DPOAE tests to assess the function of

the increased number of outer hair cells caused by *Ebf1* deletion because DPOAE detects

nonlinear responses of outer hair cells to sound. The DPOAE responses in *Foxg1Cre;Ebf1*^{fl/fl}

mice were significantly lower than those in control mice (Fig. 7B). The decreased DPOAE

response in *Foxg1Cre;Ebf1*^{fl/fl} mice also suggests that the increased number of hair cells caused

by *Ebf1* deletion did not function as outer hair cells.

Discussion

The results of this study indicate a novel and interesting role of *Ebf1* in the development of the

cochlear epithelia and otic mesenchyme. *Ebf1* is involved in the formation of appropriate

numbers of hair and supporting cells within the cochlear sensory epithelia. Moreover, *Ebf1* is

important for the development of the scala tympani, which is formed through *Ebf1* expression

within the mesenchyme, and spiral ganglion cells. The *Ebf1* deficiency in the cochlear epithelia

resulted in impaired auditory function, suggesting that *Ebf1* is necessary for normal auditory

function.

Since EBF1 was originally identified from the regulators of early B cell differentiation (Hagman

et al., 1991) and olfactory-specific genes (Wang and Reed, 1993), the expression patterns and

functions of EBF1 have been studied in B lymphocytes (Lin and Grosschedl, 1995), olfactory

epithelia (Wang et al., 1997), striatum neurons (Garel et al., 1999), the retina (Jin et al., 2010; Jin

and Xiang, 2011), the kidney (Fretz et al., 2014; Nelson et al., 2019), bone marrow stromal cells

(Derecka et al., 2020), osteoblasts (Hesslein et al., 2009), and adipose tissue (Jimenez et al.,

2007). The expression of *Ebf1* and its protein product has been reported in all three germinal

layers, and the function of EBF1 varies (Liberg et al., 2002). EBF1 is involved in cell fate

specification (Jin et al., 2010; Nechanitzky et al., 2013), cell differentiation (Lin and Grosschedl,

1995; Wang et al., 1997; Garel et al., 1999; Jimenez et al., 2007; Jin and Xiang, 2011; Nieminen-

Pihala et al., 2021), cell maturation (Fretz et al., 2014; Nelson et al., 2019), cell migration

(Derecka et al., 2020), and path finding by neurons (Garel et al., 2000; Jin and Xiang, 2011),

mainly during the developmental stage in mammals.

The present study revealed that *Ebf1* is expressed not only in the inner ear epithelium and spiral

ganglion but also in the otic mesenchyme. As the inner ear epithelium and spiral ganglion cells

are derived from the ectoderm (Wu and Kelley, 2012), *Ebf1* is expressed in both ectodermal and

mesenchymal tissues within the inner ear. Its expression in the inner ear began at approximately

E10.5, as confirmed by qRT-PCR and ISH (Fig. 1A and C). Within the inner ear epithelia, *Ebf1*

showed different expression patterns in the cochlear and vestibular organs from E13.5 to E18.5.

In the vestibule and crista, *Ebf1* was expressed in parts of *Sox2*-positive prosensory regions (Fig.

1B, black arrows and C). However, its expression in the cochlea was not limited to a part of the

Sox2-positive prosensory domain (white arrows in Fig. 1B) but expanded toward a more medial

region in the cochlear floor, where Kölliker's organ exists (Fig. 1B–D). Thus, the *Ebf1*

expression area comprised most of the GER. This difference suggests distinct functions of *Ebf1*

in the development of cochlear and vestibular organs.

To elucidate the function of *Ebf1* in inner ear development, we examined the inner ear

morphology of *Ebf1* conventional and inner ear-specific conditional knockout mice. As

suggested by the expression study, the cochlear and vestibular organs exhibited different

morphological phenotypes. No differences were observed in the vestibular morphology of

Ebf1$^{+/+}$ and *Ebf1$^{-/-}$* mice (Supplemental Fig. S2), despite *Ebf1* expression. As observed in the

olfactory epithelia and retina (Wang and Reed, 1993; Wang et al., 1997; Jin et al., 2010), the

deletion of *Ebf1* may be compensated for by other *Ebf* family genes, and studies in other mutant

mice, including double- or triple-knockout mice and mice with the dominant negative form of

the EBF protein, are needed to elucidate the function of EBF1 in the vestibular organ.

In contrast, the cochlea of *Ebf1*-deleted mice showed various phenotypes, indicating that other

Ebf subtypes do not have redundant functions with *Ebf1* in the cochlea as in B cells, osteoblasts,

and the striatum (Lin and Grosschedl, 1995; Garel et al., 1999; Nieminen-Pihala et al., 2021). HE

staining revealed loss of the scala tympani and spiral limbus (Fig. 2C). Because both structures

are derived from mesenchymal tissues (Sher, 1971; Phippard et al., 1999), we hypothesized that

these phenotypes reflect the roles of EBF1 in cochlear mesenchymal tissue. To confirm this, we

compared the formation of the scala tympani and spiral limbus in two *Ebf1*-deleted mouse strains,

conventional (*Ebf1$^{-/-}$* mice) and epithelia-specific conditional (*Foxg1Cre;Ebf1$^{fl/fl}$* mice) *Ebf1*

knockout mice (Fig. 2C). Although the scala tympani developed normally in *Foxg1Cre;Ebf1$^{fl/fl}$*

mice, the spiral limbus was hypoplastic in both *Foxg1Cre;Ebf1$^{fl/fl}$* and *Ebf1$^{-/-}$* mice. These results

clearly indicate that EBF1 in the cochlear mesenchyme is involved in the formation of the scala

tympani. In contrast, spiral limbus formation depends on both mesenchymal and epithelial or

epithelial expression of *Ebf1*. Because *Ebf1* was expressed in both the epithelia and mesenchyme

(Fig. 1B and Supplemental Fig. S1), we were unable to determine which was the case in this

study; however, mesenchyme-specific deletion of *Ebf1* will confirm the tissue responsible for

inducing the spiral limbus. Gross morphological observation (Fig 2B) and MYO7A staining of

the whole-mount cochlea (Supplemental Fig. S4) showed that the length of the cochlear duct was

shorter in *Ebf1$^{-/-}$* mice. As the number of cochlear turns in the cochlear section was lower in

Ebf1$^{-/-}$ mice than in *Foxg1Cre;Ebf1$^{fl/fl}$* mice, this phenotype reflects the function of mesenchymal

EBF1 in regulating the length of the cochlear duct.

In addition to the mesenchyme, more prominent roles of *Ebf1* have been found in the cochlear

epithelia. In both *Ebf1$^{-/-}$* and *Foxg1Cre;Ebf1$^{fl/fl}$* mice, the numbers of both hair and supporting

cells increased throughout cochlear turns at E18.5 (Fig. 3 and 4). We observed an increase in the

numbers of both inner and outer hair cells (Supplemental Fig. S3A). However, more detailed

IHC analysis of supporting cells showed that only the Deiters' cells increased in number among

PROX1-positive and SOX2-positive cells located medial to the organs of Corti, and the number

of pillar cells did not increase (Supplemental Fig. S3B). This phenotype suggests that EBF1 is

involved in the regulation of hair and supporting cell number during cochlear development. To

determine the mechanisms of hair and supporting cell number increase under *Ebf1* knockout

conditions, we evaluated the specifications and differentiation of the cochlear prosensory domain

and its proliferation and cell death status. We found that the formation of cochlear non-sensory

regions medial and lateral to the prosensory domain were normal in *Ebf1$^{-/-}$* mice (Fig. 5A).

Although the lateral region marker BMP4 has been reported to affect the development of the

prosensory domain (Ohyama et al., 2010), the normal formation of this region indicates that the

phenotypes of *Ebf1*⁻/⁻ mouse cochlear sensory epithelia were not caused by aberrant BMP4

expression, but by factors within the prosensory domain. In contrast to markers outside the

prosensory domain, the molecules expressed in the prosensory domain and Kölliker's organs,

JAG1 and SOX2, showed abnormal expression patterns (Fig. 1D, Fig. 3A, and Fig. 5A). These

two molecules were expressed in a more medial region of the *Ebf1*⁻/⁻ mouse cochlear floor at

E14.5 and E18.5. The fact that EBF1 was expressed in a more medial region than SOX2 in wild-

type mice indicates that it plays a role in suppressing the localization of JAG1- and SOX2-

positive cells in the most medial region. The study of proliferation status within the SOX2-

positive cells showed that *Ebf1* deletion enhanced the proliferation of SOX2-positive cells

specifically at E13.5 (Fig. 6 D), which was supported by the loss of p27Kip1 expression in the

possible prosensory domain of the *Ebf1*⁻/⁻ mice at E13.5 (Fig. 6E). The highest *Ebf1* expression

level at E13.5 (Fig. 1A) may be related to these phenotypes in *Ebf1*⁻/⁻ mice. This aberrant

proliferation within SOX2-positive cells was suggested to increase the numbers of hair and

supporting cells at later stages (Fig. 3A, B, D, and E). Evaluation of hearing ability at the

postnatal stage showed that an increase in hair and supporting cell numbers resulted in an

increased hearing threshold (Fig. 7). These results indicate that EBF1 suppresses the proliferation

of SOX2-positive cells and thus contributes to the development of appropriate numbers of hair

and supporting cells, resulting in the development of normal auditory function. Because the

SOX2-positive cells within the E13–14 cochlea become Kölliker's organ as well as lateral and

medial prosensory domains at a later stage (Kolla et al., 2020) and the GER has the potential to

form cochlear sensory epithelia (Kubota et al., 2021), EBF1 expressed in SOX2-positive cells

regulates the number of hair and supporting cells. Several lines of evidence support the function

of EBF1 to suppress cell proliferation. Human EBF1 has been reported to suppress the

proliferation of malignant tumors (Shen et al., 2020), and the deletion of *Rb1*, a known tumor

suppressor and cell cycle regulator (Lipinski and Jacks, 1999; Classon and Harlow, 2002), results

in the same morphology in the cochlea as that caused by *Ebf1* deletion (Sage et al., 2005).

The expression of mature cochlear cell markers, including MYO7A, VGULT3, and p75, in *Ebf1*$^{-}$

$^{/-}$ mice indicated that each cell type developed with normal cell fate specification. Although some

GER cells in *Ebf1*$^{-/-}$ mice expressed the hair cell marker MYO7A (Fig. 3B and C) and cell fate

specification appeared to deteriorate, the penetrance of this phenotype was low. Cell fate may be

regulated by EBF1 in the cochlea to a small extent; however, cell specification does not play a

prominent role in cochlear EBF1, which is different from other tissues, including B lymphocytes

(Nechanitzky et al., 2013). In contrast, several results from our study indicate that the speed of

differentiation appears to deteriorate in the *Ebf1*$^{-/-}$ mouse cochlea. The expression of *Atoh1*

within the cochlear prosensory domain, which was observed at E13.5 in wild-type mice, was

detected as late as E14.5 in *Ebf1*$^{-/-}$ mice (Fig. 5B). The regions where VGLUT3- and p75-

positive cells were detected in *Ebf1*$^{-/-}$ mouse cochlea were more basal than those in *Ebf1*$^{+/+}$

mouse cochlea at E18.5 (Supplemental Fig. S3). This delay in differentiation may be caused by

the aberrant proliferation of the prosensory domain in *Ebf1*$^{-/-}$ mice, as the deterioration of

proliferation affects the differentiation of cells (Ruijtenberg and van den Heuvel, 2016).

In addition to the cochlear epithelia and mesenchyme, deletion of *Ebf1* resulted in an altered

neuroaxonal composition of spiral ganglion neuronal cells in the organ of Corti, suggesting that

EBF1 may also affect the pathfinding of spiral ganglion cells within the cochlea, as observed in

the aberrant pathfinding of facial branchiomotor neurons and retina (Garel et al., 2000; Jin and

Xiang, 2011). The present study was limited to IHC analysis of TUJ1, and further studies are

needed to elucidate the role of *Ebf1* in spiral ganglion differentiation and development. Spiral

ganglion axons synapse with hair cells from within the otic mesenchyme, and axons extend

through the mesenchyme to innervate the organ of Corti (Coate and Kelley, 2013). Thus, deletion

of *Ebf1* in the mesenchyme may also contribute to the phenotype observed in the spiral ganglion

cells.

The expression of *Ebf1* in the medial region of the cochlear floor (Fig. 1B, C, and D) and the

spiral limbus loss (Fig. 2C) and ectopic MYO7A-positive cells within the GER (Fig. 3B and C)

in *Ebf1*-deleted mice suggest that EBF1 has functions within the GER or Kölliker's organ.

Recently, PRDM16 was reported to be a marker of Kölliker's organ and an important molecule

in its development (Ebeid et al., 2022). *Prdm16* null mutant mice showed phenotypes similar to

those of *Ebf1*-deleted mice, including shortened cochlear length, spiral limbus loss, and ectopic

hair cells within the GER. Moreover, the *Prdm16* null mutant mouse cochlear duct expressed

approximately half the amount of *Ebf1* compared to the wild-type mouse cochlear duct (Ebeid et

al., 2022), suggesting that EBF1 is involved in the development of Kölliker's organ. However, if

PRDM16 is the direct upstream regulator of EBF1, the *Ebf1* expression level should be much

less in *Prdm16* null mutant mice, and these mice should exhibit increased hair and supporting

cells and loss of scala tympani, as observed in $Ebf1^{-/-}$ mice. Thus, PRDM16 does not control the

expression of EBF1.

The phenotypes of the increased numbers of hair and supporting cells suggest the involvement

of previously reported molecules crucial for the development of cochlear sensory epithelia,

including Notch signal-related molecules (Yamamoto et al., 2011) and SOX2 (Kiernan et al.,

2005), in the regulation of *Ebf1* expression. However, *Foxg1Cre*-mediated *Notch 1*- or *Sox2*-

deleted mice do not show lower *Ebf1* expression levels in striatal neurons (Mason et al., 2005;

Ferri et al., 2013), suggesting that Notch signaling and SOX2 do not control EBF1 expression.

Identification of the molecules upstream and downstream of EBF1 will be the next step in

revealing the precise function of EBF1 in the cochlea and the grand scheme of inner ear

development.

In conclusion, our study confirms that *Ebf1* and its protein are expressed in the epithelia of the

inner ear prosensory domain as well as in Kölliker's organ, the mesenchyme, and CVG cells

within the cochlea, and play important roles in the formation of each structure. Epithelial EBF1

regulates the number of cochlear hair and supporting cells by suppressing the proliferation of the

prosensory domain and Kölliker's organ cells, mainly at E13.5, whereas mesenchymal EBF1

contributes to scala tympani formation. Therefore, epithelial EBF1 is crucial for normal hearing

in mammals.

Material and Methods

Animals

Slc: ICR mice were purchased from Japan SLC (Hamamatsu, Japan). *Ebf1$^{-/-}$* mice (Lin and

Grosschedl, 1995) and *Ebf1$^{fl/fl}$* (Gyory et al., 2012) were used in this study. *Ebf1$^{fl/+}$* mice were

crossed with *Foxg1$^{Cre/+}$ mice* (*Foxg1Cre*) (Hébert and McConnell, 2000) and *Ebf1$^{-/-}$* or

Foxg1Cre;Ebf1$^{fl/fl}$ mice were used as experimental animals. Additionally, *Ebf1$^{+/+}$*, *Ebf1$^{+/-}$*, or

Foxg1Cre;Ebf1$^{fl/+}$ mice were used as controls.

Ebf1$^{+/-}$, *Foxg1Cre*, and *Ebf1$^{fl/fl}$* mice were maintained on a C57BL/6 background. All

experimental protocols were approved by the Animal Research Committee of Kyoto University

(Med Kyo 20132, Kyoto, Japan). All animal experiments were performed according to the

National Institutes of Health Guidelines for the Care and Use of Laboratory Animals. All the

animals used in this study were maintained at the Institute of Laboratory Animals, Graduate

School of Medicine, Kyoto University. The mice were mated in the evening, and vaginal plugs

were checked early in the morning. The day a vaginal plug was detected was defined as E0.5.

qRT-PCR

Inner ears were dissected from E9.5, E10.5, E11.5, E12.5, E13.5, E14.5, E16.5, E18.5, and P0

ICR mice. After the surrounding tissue was removed from the inner ears, at least four samples

were immersed in TRIzol Reagent (Thermo Fischer Scientific, Waltham, MA, USA) and

preserved at −80 ℃ until RNA extraction. Total RNA was extracted using the RNeasy™ Mini

Kit (QIAGEN, Venlo, Netherlands) and reverse transcribed using the ReverTra Ace™ qPCR RT

Master Mix with gDNA Remover (TOYOBO, Osaka, Japan). The cDNA was mixed with

PowerUp SYBR Green Master Mix (Applied Biosystems) and various sets of gene-specific

forward and reverse primers and subsequently subjected to real-time PCR quantification using a

StepOnePlus Real-Time PCR System (Applied Biosystems, Waltham, MA, USA). The following

primer sequences were used: *Ebf1* forward, AACTCCAAGCACGGGCGGAG; *Ebf1* reverse,

CGGGCTGATGGCTTTGATACAGG; *Rplp0* forward, CACTGGTCTAGGACCCGAGAAG;

Rplp0 reverse, GGTGCCTCTGGAGATTTTCG. Relative mRNA expression levels were

calculated using the standard curve method, and the mouse housekeeping gene *Rplp0* was used

as an invariant control.

ISH

Whole embryos (E9.5–E11.5) and whole heads (E12.5–P0) were fixed in 4% paraformaldehyde

(PFA; Nacalai Tesque, Kyoto, Japan) in 0.1 M phosphate-buffered saline (PBS) at 4 ℃

overnight. Samples were cryoprotected in 30% sucrose/PBS (Nacalai Tesque), embedded in

Tissue-Tek® O.C.T.™ compound (Sakura Finetek Japan, Tokyo, Japan), and sectioned at 10-μm

thickness using a cryostat (CryoStar™ NX70; Thermo Fisher Scientific). The sections were

subsequently mounted on silane-coated glass slides (Matsunami Glass, Osaka, Japan).

cDNA fragments were generated by PCR using E13.5 inner ear cDNA of Slc:ICR mice and

subsequently cloned into the pCR®-Blunt II-TOPO® vector (Invitrogen, Waltham, MA, USA) to

prepare RNA probe templates. We synthesized digoxigenin (DIG)-labeled sense and antisense

RNA probes using a DIG RNA Labeling Kit (Roche, Basel, Switzerland) after digestion with the

appropriate restriction enzymes *Bam*HI-HF, *Hind*III, *Not*I-HF, *Sac*I, or *Xho*I (New England

Biolabs, Ipswich, MA, USA). The following probes were used for ISH: *Ebf1* (NM_001,290,709,

nucleotides 1436–2269), *Sox2* (IMAGE clone: 6413283), *Bmp4* (NM_007554.3, nucleotides

1013-1876), *Atoh1* (NM_007500.5, nucleotides 13-2111), and *Fgf10* (NM_008002.5, nucleotides

571-1027). Each corresponding sense probe was used as a negative control.

Sections were fixed with 4% PFA and 0.2% glutaraldehyde (Nacalai Tesque) in PBS at room

temperature (RT) for 10 min, bleached with 6% hydrogen peroxidase (FUJIFILM Wako Pure

Chemical Corporation, Osaka, Japan) in 0.1% Tween-20 in PBS (PBST) at RT for 10 min,

treated with proteinase K (20 μg/μL, Roche) for 5 min, and re-fixed with 4% PFA and 0.2%

glutaraldehyde in PBS at RT for 10 min.

The prehybridization was performed in hybridization solution containing 50% formamide

(Nacalai Tesque), 5× saline sodium citrate buffer (SSC, Nacalai Tesque; adjusted to pH 4.5 with

citrate), 1% sodium dodecyl sulfate (Sigma-Aldrich, St. Louis, MO, USA), 50 μg/mL yeast RNA

(Invitrogen), and 50 μg/mL heparin (Sigma-Aldrich) at 70 °C for 1 h. For hybridization, we

incubated the sections in a hybridization solution with a 0.2 μg/mL DIG-labelled RNA probe at

70 °C for 16 h in sealed plastic bags.

Sections were rinsed first in 50% formamide with 6× SSC and 1% sodium dodecyl sulfate at

70 °C, then in 50% formamide with 2.4× SSC at 65 °C, and finally in 1× Tris-buffered saline

(Nacalai Tesque) with 0.1% Tween 20 (TBST) at RT. Sections were blocked with 5% sheep

serum (Sigma-Aldrich) and incubated with a 1:4000 dilution of Anti-Digoxigenin-AP Fab

fragments (Roche) at 4 °C overnight.

After rinsing with TBST and NTMT containing 100 mM NaCl (Nacalai Tesque), 100 mM Tris-

HCl pH 9.5, 10 mM $MgCl_2$ (FUJIFILM Wako Pure Chemical Corporation), 0.1% Tween-20, and

480 μg/mL levamisole (Sigma-Aldrich), the sections were incubated with nitro-blue tetrazolium

chloride (Roche) and 5-bromo-4-chloro-3-indolyl phosphate solution (Roche). Images were

captured using a BX-50 microscope (Olympus Corp., Tokyo, Japan).

IHC analysis

IHC sections were prepared in a manner similar to that used for ISH.

After washing with PBS, all samples were incubated with Blocking One Histo (Nacalai Tesque)

for 10 min at RT and 10% normal donkey serum (Sigma-Aldrich) in PBS/0.5% Triton X- 100

with 5% Blocking One Histo for 30 min at RT.

The samples were stained with primary antibodies at 4 °C overnight or RT for 1 h. After washing

with PBST, the samples were incubated with Alexa Fluor secondary antibodies. F-actin (actin

filaments) was stained with phalloidin 647 (1:500; Thermo Fisher Scientific) at RT for 1 h.

Nuclei were stained with 4′,6-diamidino-2-phenylindole (DAPI; Thermo Fisher Scientific).

The following primary antibodies were used in this study: rabbit anti-EBF1 antibody (1:1000,

AB10523; Millipore, Darmstadt, Germany,), mouse anti-MYO7A antibody (1:1000, 138-1;

Developmental Studies Hybridoma Bank, Iowa City, IA, USA), rabbit anti-MYO7A antibody

(1:1000, 25-6790; Proteus BioSciences, Waltham, MA, USA), goat anti-SOX2 antibody (1:250,

AF2018; R&D Systems, Minneapolis, MN, USA), rabbit anti-SOX2 antibody (1:100, 11064-1-

AP; Proteintech, Manchester, UK), rabbit anti-VGLUT3 antibody (1:500, 135 203; Synaptic

systems, Goettingen, Germany), goat anti-JAG1 antibody (1:500, sc-6011; Santa Cruz, Dallas,

TX, USA), mouse anti-p27Kip1 antibody (1:200, 610242; BD Transduction Laboratories), rabbit

anti-Tubulin β 3 (TUJ1) antibody (1:1000, PRB-435P; Biolegend, San Diego, CA, USA), rabbit

anti-PROX1 antibody (1:500, AB5475; Millipore), rabbit anti-p75 antibody (1:500, AB1554;

Millipore), and rabbit anti-cleaved caspase 3 (CC3) antibody (1:400, Asp175; Cell Signaling

Technology, Danvers, MA, USA).

Antigen retrieval was performed for p27Kip1 staining by heating sections in HistoVT one

(Nacalai Tesque) at 90 °C for 10 min prior to the addition of the primary antibodies.

The secondary antibodies used were Alexa Flour 488 donkey anti-rabbit IgG, Alexa Flour 488

donkey anti-goat IgG, Alexa Flour 488 donkey anti-mouse IgG, Alexa Flour 568 donkey anti-

rabbit IgG, Alexa Flour 568 donkey anti-goat IgG, Alexa Flour 633 donkey anti-rabbit IgG,

Alexa Flour 633 donkey anti-goat IgG, and Alexa Flour 633 donkey anti-mouse IgG (1:500;

Thermo Fisher Scientific).

The sections were mounted using Fluoromount-G® Anti-Fade (Southern Biotechnology

Associates Inc., Birmingham, AL, USA). Images of the specimens were captured using an

Olympus BX50 microscope (Olympus), an Olympus DP70 digital camera (Olympus), and a

Zeiss LSM900 with Airyscan2 (Carl Zeiss AG, Oberkochen, Germany).

HE staining

Frozen sections (10-μm thick) were immersed in Mayer's Hematoxylin Solution (FUJIFILM

Wako Pure Chemical Corporation) at RT for 3 min and in Eosin Alcohol solution (FUJIFILM

Wako Pure Chemical Corporation) at RT for 1 min. Dehydration was performed using graded

ethanol solutions (70%, 80%, 90%, and 100%) and clearing was performed three times using

xylene.

Cochlea whole-mount preparation

The inner ears were dissected from mice heads at E18.5 and fixed in 4% PFA in PBS at RT for 1

h. After fixation and before primary antibody staining, the outer membrane, including the

tectorial membrane, was removed to expose the organs of Corti. After staining with a secondary

antibody, the organ of Corti was dissected and mounted on a glass slide for imaging.

Proliferation and apoptosis assays

Cell proliferation in the cochlea was assessed by detecting the incorporated EdU (Thermo Fisher

Scientific) on frozen sections. EdU was detected using the Click-iT Plus EdU Cell Proliferation

Kit for Imaging Alexa 555 Dye (Thermo Fisher Scientific), according to the manufacturer's

instructions. Pregnant mice were injected with EdU (Thermo Fisher Scientific) at E12.5, E13.5,

and E14.5 (three injections at 50 μg/g at 2-h intervals) and at E16.5 (a single injection at 100

μg/g). E12.5, E13.5, and E14.5 embryos were collected 8 h after the first injection. E16.5

embryos were collected 4 h after the injection. The basal or basal-to-middle regions of the

cochlear duct were observed at E12.5 or at E13.5, E14.5, and E16.5, respectively.

Apoptotic cells were detected by identifying the expression of CC3 in frozen sections via IHC

staining.

ABR and DPOAE

ABR measurements were performed under general anesthesia as described previously (Kada et al., 2009) at P21 (n = 3 for each genotype). The thresholds for 10, 20, and 40 kHz were determined based on the responses at different intensities with 5 dB sound pressure level intervals. DPOAE recordings were performed as described previously (Hamaguchi et al., 2012) at P21 (n = 4 for each genotype). Two primary tones (f1, f2, f1<f2) were used as input signals, with f2 set at eight frequency points (4, 6, 8, 12, 16, 24, 32, and 40 kHz), maintaining a frequency ratio of f2/f1 = 1.2. The intensity levels of the stimulatory sounds were 65 and 55 dB sound pressure level for f1 and f2, respectively. DPOAE was detected as a peak at 2f1–f2 in the spectrum.

Quantification

Cell quantification and measurements were performed at E18.5 using the Cell Counter plugin of ImageJ (Schneider et al., 2012). The total length of the cochlea was measured based on the region with MYO7A-positive hair cells from the basal to apical turns. Cochlear hair cells were identified by phalloidin and MYO7A labeling. Two types of cells, PROX1- and SOX2-positive cells, were counted to quantify the supporting cells of the cochlea. The cochlear duct was divided into three regions: basal, middle, and apical, and we counted the number of cells per 200 μm in each region for MYO7A- and PROX1-positive cells or 100 μm in the basal region for SOX2-positive cells.

Statistical analysis

For all statistical analyses, at least three samples from each experimental group were analyzed.

Student's *t*-test was used to determine the differences between two experimental groups. One-

way or two-way analysis of variance was performed to assess the differences between more than

two experimental groups and *p*-values less than 0.05 were considered statistically significant.

Statistical analyses were performed using R version 4.2.2 (2022-10-31). All details of statistical

analyses are provided in the figures and legends.

Competing Interest Statement

The authors declare that they have no known competing financial interests or personal

relationships that could have appeared to influence the work reported in this paper

Acknowledgements

We thank Dr. Rudolf Grosschedl for providing $Ebf1^{-/-}$ and $Ebf1^{fl/fl}$ mice and Dr. Ryoichiro

Kageyama for providing *Atoh1* ISH probe. This study was supported by JSPS KAKENHI (grant

number: JP22H03234) awarded to NY.

Author Contributions

N.Y. and H.O. designed the study. H.K., H.O., R.Y., and Y.T. performed the experiments with

advice from N.Y., T.K., and K.O.. A.Y. performed ABR and DPOAE. H.K. and N.Y. wrote the

manuscript with contribution from all authors.

References

Bermingham-McDonogh O, Oesterle EC, Stone JS, Hume CR, Huynh HM, Hayashi T (2006)
Expression of Prox1 during mouse cochlear development. The Journal of comparative
neurology 496:172-186.
Bermingham NA, Hassan BA, Price SD, Vollrath MA, Ben-Arie N, Eatock RA, Bellen HJ,
Lysakowski A, Zoghbi HY (1999) Math1: an essential gene for the generation of inner
ear hair cells. Science 284:1837-1841.
Chen P, Segil N (1999) p27(Kip1) links cell proliferation to morphogenesis in the developing
organ of Corti. Development 126:1581-1590.
Chen P, Johnson JE, Zoghbi HY, Segil N (2002) The role of Math1 in inner ear development:
Uncoupling the establishment of the sensory primordium from hair cell fate
determination. Development 129:2495-2505.
Classon M, Harlow E (2002) The retinoblastoma tumour suppressor in development and cancer.
Nature Reviews Cancer 2:910-917.
Coate TM, Kelley MW (2013) Making connections in the inner ear: Recent insights into the
development of spiral ganglion neurons and their connectivity with sensory hair cells.
Seminars in Cell and Developmental Biology 24:460-469.
Dayaratne MW, Vlajkovic SM, Lipski J, Thorne PR (2014) Kolliker's organ and the
development of spontaneous activity in the auditory system: implications for hearing
dysfunction. BioMed research international 2014:367939.
Derecka M, Herman JS, Cauchy P, Ramamoorthy S, Lupar E, Grun D, Grosschedl R (2020)
EBF1-deficient bone marrow stroma elicits persistent changes in HSC potential. Nature
immunology 21:261-273.
Ebeid M, Barnas K, Zhang H, Yaghmour A, Noreikaite G, Bjork BC (2022) PRDM16
expression and function in mammalian cochlear development. Developmental Dynamics.
Ferri A, Favaro R, Beccari L, Bertolini J, Mercurio S, Nieto-Lopez F, Verzeroli C, La Regina F,
De Pietri Tonelli D, Ottolenghi S, Bovolenta P, Nicolis SK (2013) Sox2 is required for
embryonic development of the ventral telencephalon through the activation of the ventral
determinants Nkx2.1 and Shh. Development 140:1250-1261.
Fretz JA, Nelson T, Velazquez H, Xi Y, Moeckel GW, Horowitz MC (2014) Early B-cell factor
1 is an essential transcription factor for postnatal glomerular maturation. Kidney
International 85:1091-1102.
Garel S, Garcia-Dominguez M, Charnay P (2000) Control of the migratory pathway of facial
branchiomotor neurones. Development 127:5297-5307.
Garel S, Marin F, Grosschedl R, Charnay P (1999) Ebf1 controls early cell differentiation in the
embryonic striatum. Development 126:5285-5294.
Gyory I, Boller S, Nechanitzky R, Mandel E, Pott S, Liu E, Grosschedl R (2012) Transcription
factor Ebf1 regulates differentiation stage-specific signaling, proliferation, and survival
of B cells. Genes and Development 26:668-682.
Hagman J, Travis A, Grosschedl R (1991) A novel lineage-specific nuclear factor regulates mb-1
gene transcription at the early stages of B cell differentiation. The EMBO journal
10:3409-3417.
Hagman J, Gutch MJ, Lin H, Grosschedl R (1995) EBF contains a novel zinc coordination motif
and multiple dimerization and transcriptional activation domains. EMBO Journal
14:2907-2916.

Hamaguchi K, Yamamoto N, Nakagawa T, Furuyashiki T, Narumiya S, Ito J (2012) Role of
PGE-type receptor 4 in auditory function and noise-induced hearing loss in mice.
Neuropharmacology 62:1841-1847.
Hébert JM, McConnell SK (2000) Targeting of cre to the Foxg1 (BF-1) locus mediates loxP
recombination in the telencephalon and other developing head structures. Developmental
Biology 222:296-306.
Hesslein DG, Fretz JA, Xi Y, Nelson T, Zhou S, Lorenzo JA, Schatz DG, Horowitz MC (2009)
Ebf1-dependent control of the osteoblast and adipocyte lineages. Bone 44:537-546.
Jimenez MA, Akerblad P, Sigvardsson M, Rosen ED (2007) Critical role for Ebf1 and Ebf2 in
the adipogenic transcriptional cascade. Molecular and Cellular Biology 27:743-757.
Jin K, Xiang M (2011) Ebf1 deficiency causes increase of Müller cells in the retina and
abnormal topographic projection at the optic chiasm. Biochemical and Biophysical
Research Communications 414:539-544.
Jin K, Jiang H, Mo Z, Xiang M (2010) Early B-cell factors are required for specifying multiple
retinal cell types and subtypes from postmitotic precursors. The Journal of neuroscience :
the official journal of the Society for Neuroscience 30:11902-11916.
Kada S, Nakagawa T, Ito J (2009) A Mouse Model for Degeneration of the Spiral Ligament.
Journal of the Association for Research in Otolaryngology 10:161-172.
Kiernan AE, Pelling AL, Leung KKH, Tang ASP, Bell DM, Tease C, Lovell-Badge R, Steel KP,
Cheah KSE (2005) Sox2 is required for sensory organ development in the mammalian
inner ear. Nature 434:1031-1035.
Kolla L, Kelly MC, Mann ZF, Anaya-Rocha A, Ellis K, Lemons A, Palermo AT, So KS, Mays
JC, Orvis J, Burns JC, Hertzano R, Driver EC, Kelley MW (2020) Characterization of the
development of the mouse cochlear epithelium at the single cell level. Nature
communications 11:2389.
Kubota M, Scheibinger M, Jan TA, Heller S (2021) Greater epithelial ridge cells are the principal
organoid-forming progenitors of the mouse cochlea. Cell Reports 34:108646.
Li C, Shu Y, Wang G, Zhang H, Lu Y, Li X, Li G, Song L, Liu Z (2018) Characterizing a novel
vGlut3-P2A-iCreER knockin mouse strain in cochlea. Hear Res 364:12-24.
Liberg D, Sigvardsson M, Akerblad P (2002) The EBF/Olf/Collier family of transcription
factors: regulators of differentiation in cells originating from all three embryonal germ
layers. Molecular and Cellular Biology 22:8389-8397.
Lin H, Grosschedl R (1995) Failure of B-cell differentiation in mice lacking the transcription
factor EBF. Nature 376:263-267.
Lipinski MM, Jacks T (1999) The retinoblastoma gene family in differentiation and development.
Oncogene 18:7873-7882.
Mason HA, Rakowiecki SM, Raftopoulou M, Nery S, Huang Y, Gridley T, Fishell G (2005)
Notch signaling coordinates the patterning of striatal compartments. Development
132:4247-4258.
Nechanitzky R, Akbas D, Scherer S, Gyory I, Hoyler T, Ramamoorthy S, Diefenbach A,
Grosschedl R (2013) Transcription factor EBF1 is essential for the maintenance of B cell
identity and prevention of alternative fates in committed cells. Nature immunology
14:867-875.
Nelson T, Velazquez H, Troiano N, Fretz JA (2019) Early B Cell Factor 1 (EBF1) Regulates
Glomerular Development by Controlling Mesangial Maturation and Consequently COX-
2 Expression. Journal of the American Society of Nephrology : JASN 30:1559-1572.

Nieminen-Pihala V, Tarkkonen K, Laine J, Rummukainen P, Saastamoinen L, Nagano K, Baron
R, Kiviranta R (2021) Early B-cell Factor1 (Ebf1) promotes early osteoblast
differentiation but suppresses osteoblast function. Bone 146:115884.
Ohyama T, Basch ML, Mishina Y, Lyons KM, Segil N, Groves AK (2010) BMP Signaling Is
Necessary for Patterning the Sensory and Nonsensory Regions of the Developing
Mammalian Cochlea. Journal of Neuroscience 30:15044-15051.
Phippard D, Lu L, Lee D, Saunders JC, Crenshaw EB, 3rd (1999) Targeted mutagenesis of the
POU-domain gene Brn4/Pou3f4 causes developmental defects in the inner ear. The
Journal of neuroscience : the official journal of the Society for Neuroscience 19:5980-
5989.
Ruijtenberg S, van den Heuvel S (2016) Coordinating cell proliferation and differentiation:
Antagonism between cell cycle regulators and cell type-specific gene expression. Cell
Cycle 15:196-212.
Sage C, Huang M, Karimi K, Gutierrez G, Vollrath MA, Zhang DS, Garcia-Anoveros J, Hinds
PW, Corwin JT, Corey DP, Chen ZY (2005) Proliferation of functional hair cells in vivo
in the absence of the retinoblastoma protein. Science 307:1114-1118.
Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image
analysis. Nat Methods 9:671-675.
Shen Z, Chen Y, Li L, Liu L, Peng M, Chen X, Wu X, Sferra TJ, Wu M, Lin X, Cheng Y, Chu J,
Shen A, Peng J (2020) Transcription Factor EBF1 Over-Expression Suppresses Tumor
Growth in vivo and in vitro via Modulation of the PNO1/p53 Pathway in Colorectal
Cancer. Front Oncol 10:1035.
Sher AE (1971) The embryonic and postnatal development of the inner ear of the mouse. Acta
oto-laryngologica Supplementum 285:1-77.
Taniguchi M, Yamamoto N, Nakagawa T, Ogino E, Ito J (2012) Identification of tympanic
border cells as slow-cycling cells in the cochlea. PloS one 7:e48544.
Urness LD, Wang X, Shibata S, Ohyama T, Mansour SL (2015) Fgf10 is required for
specification of non-sensory regions of the cochlear epithelium. Developmental Biology
400:59-71.
von Bartheld CS, Patterson SL, Heuer JG, Wheeler EF, Bothwell M, Rubel EW (1991)
Expression of nerve growth factor (NGF) receptors in the developing inner ear of chick
and rat. Development 113:455-470.
Wang MM, Reed RR (1993) Molecular cloning of the olfactory neuronal transcription factor
Olf-1 by genetic selection in yeast. Nature 364:121-126.
Wang SS, Tsai RY, Reed RR (1997) The characterization of the Olf-1/EBF-like HLH
transcription factor family: implications in olfactory gene regulation and neuronal
development. Journal of Neuroscience 17:4149-4158.
Wu DK, Kelley MW (2012) Molecular mechanisms of inner ear development. Cold Spring
Harbor Perspectives in Biology 4:a008409.
Yamamoto N, Chang W, Kelley MW (2011) Rbpj regulates development of prosensory cells in
the mammalian inner ear. Developmental Biology 353:367-379.
Yamamoto R, Ohnishi H, Omori K, Yamamoto N (2021) In silico analysis of inner ear
development using public whole embryonic body single-cell RNA-sequencing data.
Developmental Biology 469:160-171.
795

 Figures and Figure Legends

Fig. 1. Quantitative and spatiotemporal expression of *Ebf1* during inner ear development.

A. Results of quantitative reverse transcription polymerase chain reaction (qRT-PCR) analysis

for *Ebf1* in the inner ear of wild-type mice from embryonic day (E) 9.5 to P0. The value of each

date is normalized to the value of E9.5. Box plot representing the medians and interquartile

ranges of the relative mRNA expression of *Ebf1*.

B and **C**. Result of *in situ* hybridization for *Ebf1* and *Sox2* in cross sections of the inner ear of

wild-type mice at E13.5 (B) and E10.5 and E18.5 (C). Areas enclosed by dashed lines indicate

the inner ear epithelium. Low-magnification images of the cochlear basal turn and vestibule are

presented in the uppermost panels of **B**. High-magnification images of the apical turn of the

cochlear duct and vestibules are presented in the middle and lower panels of **B**. Images of E10.5

otocysts and E18.5 cochleae and vestibules are presented in the uppermost, middle, and

lowermost images of **C**, respectively.

D. Immunohistochemical images of E14.5 $Ebf1^{+/+}$ (upper panels) and $Ebf1^{-/-}$ (lower panels)

mouse cochlear ducts.

One-way analysis of variance (ANOVA) with Tukey–Kramer *post-hoc* tests was performed. *p

<0.05, **p <0.01. D, dorsal; L, lateral. cd, cochlear duct; sg, spiral ganglion; Ut, utricle; Sa,

saccule; Lc, lateral crista; SL, spiral ligament; Ko, Kölliker's organ; oC, organ of Corti; Pd,

prosensory domain. Scale bars: 100 μm.

Fig. 2 Development of the cochlear duct is deteriorated in *Ebf1⁻ᐟ⁻* mice.

A. Gross morphology of membranous labyrinth of the inner ear of *Ebf1⁺ᐟ⁺* and *Ebf1⁻ᐟ⁻* mice at embryonic day (E) 18.5. **B.** Gross morphology of the cochlear floor after the lateral wall and roof of the cochlea were removed at E18.5. **C.** Hematoxylin and eosin-stained cross sections of the cochlea at E18.5 from *Ebf1⁺ᐟ⁺*, *Ebf1⁻ᐟ⁻*, *Foxg1Cre;Ebf1ᶠˡᐟ⁺*, and *Foxg1Cre;Ebf1ᶠˡᐟᶠˡ* mice. The upper panels are low-magnification images, and the middle and lower panels are high-magnification images. Areas enclosed by dashed lines indicate the spiral limbus. cd, cochlear duct; Ut, utricle; Sa, saccule; Lc, lateral crista; Pc, posterior crista; sv, scala vestibuli; sm, scala media; st, scala tympani; sl, spiral limbus: Ko, Kölliker's organ; oC, organ of Corti. Scale bars: 0.5 mm in **A** and **B** and 200 μm in **C**.

Fig. 3. *Ebf1* deletion increases the number of MYO7A-, SOX2-, and PROX1-positive cells in the cochlea.

A. Cross sections of the basal and apical turns of the cochlea of *Ebf1*⁺/⁺ and *Ebf1*⁻/⁻ mice at embryonic day (E) 18.5, labeled with MYO7A (green) and SOX2 (magenta). Magnified images of the organ of Corti (oC) and Kölliker's organ (Ko) are presented in the four right panels. **B**–**E.** Whole mount cochlear images from *Ebf1*⁺/⁺ and *Ebf1*⁻/⁻ mice at E18.5, labeled with MYO7A

835 (green, **B** and **C**), phalloidin (white, **C**), SOX2 (magenta, **D**), and PROX1 (green, **E**). Scale bars:

836 50 μm in **A**, **D**, and **E**; 100 μm in **B**; 10 μm in **C**.

Fig. 4. Increases in the number of hair and supporting cells.

A. Schematic diagram of the cochlea duct showing the positions of the basal, middle, and apical

regions of the cochlea. **B**. Total hair cell numbers per 200 μm in the basal, middle, and apical

regions of the cochlear ducts of *Ebf1*$^{+/+}$, *Ebf1*$^{+/-}$, and *Ebf1*$^{-/-}$ mice at embryonic day (E) 18.5. **C**.

Numbers of SOX2-positive cells per 100 μm in the basal region of the cochlea. **D**. Numbers of

PROX1-positive cells per 200 μm in the basal, middle, and apical regions of the cochlea. Two-

way analysis of variance (ANOVA) test with Bonferroni *post-hoc* test (**B** and **D**) and Student's *t*-

846   test (**C**) were performed. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, and ****$p < 0.0001$; ns, not

847   significant. Error bars represent mean $\pm$ standard deviation (n = 4).

 Fig. 5. *Ebf1* deletion causes JAG1 expression to spread inward, delaying *Atoh1* expression

during cochlear development.

A. Results of *in situ* hybridization of *Fgf10* and *Bmp4* and immunostaining of JAG1 (green) on

cross sections of the basal-to-middle region of the cochlea of *Ebf1*[+/+] and *Ebf1*[-/-] mice at

embryonic day (E) 13.5 and E14.5. The prosensory domain is shown in brackets. **B**. Result of *in*

situ hybridization of *Atoh1* on cross sections of the basal-to-middle region of the cochlea of

Ebf1[+/+] and *Ebf1*[-/-] mice at E14.5 and E15.5. Areas enclosed by dashed lines indicate cochlear

ducts and vestibules. Scale bars: 100 μm.

Fig. 6. Effect of *Ebf1* deletion on cell proliferation during inner ear development.

A. Cross sections of the cochlear basal regions at embryonic day (E) 12.5, E13.5, E14.5, and E16.5 from *Ebf1⁺/⁺* and *Ebf1⁻/⁻* mice. Sections were immunostained with 5-ethynyl-2'-deoxyuridine (EdU, green) and SOX2 (magenta). E16.5 sections were counter-stained with 4',6-diamidino-2-phenylindole (DAPI, gray). Areas enclosed by dashed lines indicate the cochlear ducts and brackets indicate prosensory domains. **B–D**. Quantitative assessment of the SOX2-

positive region in the cochlea epithelia. The numbers of SOX2-positive cells (**B**) as well as EdU-

and SOX2-double positive cells (**C**) were counted, and the percentage of EdU-positive cells

among SOX2-positive cells (**D**) was calculated. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, and ****p

< 0.0001. Error bars represent mean ± standard deviation. n = 4. **E**. Cross sections of the cochlear

basal regions at E13.5 and E14.5 from *Ebf1*$^{+/+}$ and *Ebf1*$^{-/-}$ mice immunostained with p27Kip1

(green) and SOX2 (magenta). Scale bars: 100 μm.

Fig. 7. *Ebf1* deletion impairs auditory function.

A. Auditory brainstem response (ABR) thresholds of P21 *Foxg1Cre;Ebf1^{fl/fl}* (red) and

Foxg1Cre;Ebf1^{fl/+} (blue, control) mice. **B**. Distortion product optoacoustic emissions (DPOAE)

were measured at P21 from *Foxg1Cre;Ebf1^{fl/fl}* (red) and control (blue) mice. *$p < 0.05$, **$p <$

0.01, and ****$p < 0.0001$ from one-way analysis of variance (ANOVA) with Tukey's HSD *post-*

hoc test. Error bars represent mean ± standard error of mean. **C**. The whole mount images of the

cochlear basal regions from *Foxg1Cre;Ebf1^{fl/+}* and *Foxg1Cre;Ebf1^{fl/fl}* mice at P23 labeled with

phalloidin (green). Scale bar: 20 μm.

Supplemental Figures and Legends for Supplemental Figures

Supplemental Fig. S1. Spatiotemporal expression of *Ebf1* during inner ear development.

Result of *in situ* hybridization for *Ebf1* and *Sox2* on cross sections of the inner ear of wild-type

mice at embryonic day (E) 11.5, E12.5, E16.5, and P0. Areas enclosed by dashed lines indicate

the inner ear epithelium. From E11.5 to P0, *Ebf1* is expressed in the sensory epithelium of the

cochlea (white arrows), the vestibular and semicircular canals (arrows), the spiral ganglion

(white asterisks), and the surrounding mesenchymal tissues (asterisks and SL). sg, spiral

ganglion; Ut, utricle; Sa, saccule; Lc, lateral crista; SL, spiral ligament; Ko, Kölliker's organ; oC,

organ of Corti. Scale bars: 100 μm.

Supplemental Fig. S2. *Ebf1* deletion does not affect hair and supporting cells in the

vestibular macula and crista.

Cross sections of the embryonic day (E) 18.5 vestibular region immunostained for MYO7A

(green) and SOX2 (magenta). Low-magnification (left four panels) and high-magnification (right

panels) images of the saccule, lateral crista, and utricle are presented. Ut, utricle; Sa, saccule; Lc,

lateral crista. Scale bars: 100 μm.

Supplemental Fig. S3. *Ebf1* **deletion causes increased number of inner hair cells and delayed differentiation of hair and supporting cells.**

High-magnification images of the basal, middle, and apical regions of whole-mount cochlear samples from E18.5 *Ebf1*⁺/⁺ and *Ebf1*⁻/⁻ mice. **A**. Immunostaining with phalloidin (green) and an anti-VGLUT3 antibody (magenta). In the cochlea of *Ebf1*⁻/⁻ mice, the row number of VGLUT3-positive cells was 2 to 3, whereas this number was only 1 in wild-type mice. VGLUT3-positive cells were observed only in the basal region of the cochlea but not in the middle and apical regions of *Ebf1*⁻/⁻ mice. Several VGLUT3-positive cells were observed in the outer hair cell area (arrowheads). **B**. Immunostaining with anti-MYO7A (gray) and p75 (green) antibodies. In the

cochlea of *Ebf1*$^{-/-}$ mice, the row number of p75-positive cells was similar to that in *Ebf1*$^{+/+}$ mice,

although their arrangement was deteriorated in *Ebf1*$^{-/-}$ mice (arrow). p75-positive cells are

observed only in the basal and middle region in the *Ebf1*$^{-/-}$ mice. Scale bars: 20 μm.

Supplemental Fig. S4. Quantification of cochlear length.

A. Whole mount images of the cochlea of embryonic day (E) 18.5 *Ebf1$^{+/+}$*, *Ebf1$^{+/-}$*, and *Ebf1$^{-/-}$*

mice labeled with MYO7A (green). **B**. Quantification of cochlear duct length of E18.5 *Ebf1$^{+/+}$*,

Ebf1$^{+/-}$, and *Ebf1$^{-/-}$* mice. The cochlear length of *Ebf1$^{-/-}$* mice was significantly shorter than those

of *Ebf1$^{+/+}$* and *Ebf1$^{+/-}$* mice. Two-way analysis of variance (ANOVA) test with Bonferroni *post-*

hoc tests was performed. ***$p < 0.001$, and ****$p < 0.0001$; ns, not significant. Error bars

represent mean ± standard deviation. n = 6

Supplemental Fig. S5. *Ebf1* **deletion causes aberrant spiral ganglion development and axon**

outgrowth to cochlear hair cells.

A. Cross sections of the basal, middle, and apical regions of the cochlea and spiral ganglion in

embryonic day (E) 18.5 *Ebf1*$^{+/+}$ and *Ebf1*$^{-/-}$ mice stained with Tubulin β 3 (TUJ1, green) and

4',6-diamidino-2-phenylindole (DAPI, gray). The morphology of the spiral ganglion in *Ebf1*$^{-/-}$

mice differed from that in *Ebf1*$^{+/+}$ mice. TUJ1-positive cell bodies were observed below the

organ of Corti (arrow) and at their normal site. Additionally, innervation from the spiral ganglion

was observed in the hair cell part and in Kölliker's organ in the middle and apical of the cochlea

(arrowheads and brackets). **B**. High-magnification view of the basal region of the whole mount

cochlear image in E18.5 *Ebf1$^{+/+}$* and *Ebf1$^{-/-}$* mice labeled with TUJ1 (green) and MYO7A (gray).

In *Ebf1$^{+/+}$* mice, the neurons ran parallel to the outer hair cells, whereas in *Ebf1$^{-/-}$* mice, the

neurons formed a reticulation within the cochlear hair cell regions. Scale bars: 100 μm (**A**) and

20 μm (**B**).

Supplemental Fig. S6. *Ebf1* deletion did not affect apoptosis during inner ear development.

Cross sections of the embryonic day (E) 11.5 inner ear and E13.5 cochlear basal region from *Ebf1⁺/⁺* and *Ebf1⁻/⁻* mice. Sections were labeled with cleaved caspase 3 (CC3, green) and 4',6-diamidino-2-phenylindole (DAPI, blue). The number of CC3-positive cells was similar between *Ebf1⁻/⁻* and *Ebf1⁺/⁺* mice at E11.5 and E13.5. Scale bars: 100 μm.