An integrative trait-based framework to infer resource budgets and life-histories of long-

lived plants

Huw Cooksley[1,2], Matthias Schleuning[3], Alexander Neu[2,3,4], Karen J. Esler[2], Frank M. Schurr[1,5]

1: Institute of Landscape and Plant Ecology, University of Hohenheim, Stuttgart, Germany

2: Department of Conservation Ecology and Entomology, Stellenbosch University, Stellenbosch,

South Africa

3: Senckenberg Biodiversity and Climate Research Centre (SBiK.F), Frankfurt am Main,

Germany

4: Institute for Ecology, Evolution and Diversity, Goethe University Frankfurt, Frankfurt am

Main, Germany

5: KomBioTa – Center for Biodiversity and Integrative Taxonomy, University of Hohenheim &

State Museum of Natural History, Stuttgart, Germany

Contact & corresponding author: Huw Cooksley – huw.cooksley@uni-hohenheim.de

Abstract

A fundamental assumption of functional ecology is that functional traits determine life-histories. Yet correlations between traits and life-history components are often weak, especially for long-lived plants. This is because trade-offs, constraints, dynamic resource budgets and the scaling from single organs to entire plants cause complex relationships between traits and life-history. To elucidate these relationships, we present an integrated Trait-Resource-Life-History (TRL) framework that infers how functional traits affect organ-level costs and benefits of different life-history components, how these costs and benefits shape the dynamics of whole-plant resource acquisition and allocation, and how these dynamics translate into life history. We illustrate this framework by developing a TRL model for a functionally diverse group of woody plants (22 species of the genus *Protea* from the South African Greater Cape Floristic Region). Using hierarchical Bayesian latent state-space modelling, we statistically parameterise this model from data on year-to-year variation in growth, reproduction and maternal care (serotiny) for 600 individuals. The parameterised model reveals that higher resource acquisition translates into both larger absolute resource pools and greater proportional resource allocation to reproduction. Accordingly, specific leaf area, a key trait increasing resource acquisition, is associated with larger resource pools, an earlier age of maturity as well as increased vegetative and reproductive performance at young to intermediate ages. In contrast, seed nitrogen content has opposing effects on the benefits of different organs and thus only shows weak correlations with life-history components. Importantly, the TRL model identifies trait- and resource-mediated trade-offs at the level of organs, whole-plant resource budgets and life-histories. It can thus quantify key components of life-history theory that are so far largely inaccessible for long-lived plants. This

permits novel insights into ecological and evolutionary mechanisms shaping life-histories.

Application of the proposed framework to a broad range of plant systems should be facilitated by

the increasing availability of trait and demographic data, whole-plant phenotyping and high-

resolution remote sensing. The integration of the TRL framework with models of biotic

interactions further holds promise for a resource-based understanding of community dynamics

across trophic levels and a closer integration of functional ecology, evolutionary ecology,

community ecology and ecosystem science.

Introduction

Understanding the drivers of organismal performance is foundational to ecology and evolutionary biology (Stearns, 1992). How individuals take up resources and allocate them to different dimensions of performance (such as growth or reproduction) shapes their life-histories, fitness and demographic rates, which in turn influence the dynamics of populations and communities (Shipley *et al*., 2016; Ohse *et al*., 2023). The vital rates of an individual, and its fitness in a given environment, hence arise from its resource budget.

Trait-based approaches have proliferated as a means to predict plant demography and fitness (Lavorel & Garnier, 2002; McGill, *et al*. 2006; Violle *et al*., 2014; Evans *et al*., 2016; Shipley *et al*., 2016; Salguero-Gómez *et al*., 2018; Laughlin *et al*., 2020). These approaches assume that functional traits influence the growth, survival, or reproductive rate of an individual (Violle *et al*., 2007; Reich, 2014; Yang *et al*., 2018). Despite the promise of trait-based approaches, numerous studies find perplexingly weak correlations between traits and whole-plant performance (Adler *et al*., 2013; Pistón *et al*., 2019; Swenson *et al*., 2020). This is especially the case in long-lived plants, which show substantial year-to-year variation in growth and reproduction and may experience trade-offs between performance early versus late in life (Petit & Hampe, 2006; Yang *et al*., 2018). Three challenges to a trait-based understanding of life-histories are particularly pronounced in long-lived plants, such as trees and shrubs. These challenges are to understand how trait effects on resource costs and benefits scale up from organs to entire plants, how whole-plant costs and benefits shape resource allocation to different life-

history dimensions, and how resource allocation changes through the lifetime of an individual.

We expand on these three challenges in the following paragraphs.

Many functional traits are proxies for per-organ or per-unit resource investment and for the

expected return on that investment. For instance, specific leaf area (SLA) reflects the costs of

constructing and maintaining a given unit of leaf area, the expected energetic returns of this

investment, and the susceptibility of leaves to biotic and abiotic stress (such as herbivory and

desiccation) which reduce return on resource investment (Reich *et al*., 1997; Wright *et al*., 2004).

However, since SLA is a measure per unit leaf mass, it has to be combined with whole-plant leaf

biomass to determine plant-level construction costs and benefits in terms of carbon assimilation.

Hence, it is necessary to scale up per-unit or per-organ traits to the whole plant level (Yang *et al*.,

2018).

Constraints and trade-offs shape resource allocation in different ways (Stearns, 1992; Niklas,

1992). First, resource allocation is constrained by positive allometries if allocation to one organ

must be matched by allocation to another. For instance, trees and shrubs cannot produce flowers

and leaves without allocation to wood that provides mechanical support, water and nutrients

(Niklas, 1992; 1994). Secondly, resource-mediated trade-offs constrain phenotypes if resource

allocation to one organ comes at the expense of allocation to other organs, causing, for instance,

a trade-off between reproductive and vegetative performance (Levins, 1968). Thirdly, trade-offs

between the quantity and quality of organs can limit resource allocation (Moles & Westoby,

2006; Shipley *et al*., 2006). Such quantity-quality trade-offs arise if trait values that increase the

costs of an organ (e.g. large seed size) also increase the organ's benefits (e.g. higher germination

and recruitment success). Identifying how traits and resources mediate these constraints and

trade-offs is thus necessary to understand why organisms occupy a limited subset of the possible

life-history space.

Changes in resource acquisition and allocation over an organism's lifetime may furthermore lead

to variation in trait-performance relationships with size and age (Iida *et al*., 2014; Gibert *et al*.,

2016; Gray *et al*., 2019). For instance, the benefit of high SLA for vegetative growth may

decrease with size in long-lived plants, as the shorter lifespan of high SLA leaves implies more

investment into new support structures as plants grow (Gibert *et al*., 2016). As another example,

plants with high seed mass or nutrient content should initially grow faster than ones with lower

seed mass or nutrient content (Westoby, 1998; Cooksley *et al*., 2023). However, once they

become reproductively mature, they need to invest more resources to produce a given number of

seeds hence they may grow more slowly as adults and reproduce later (Stearns, 1992). Such size-

and/or age-dependent changes may thus contribute to weakening correlations between traits and

whole-plant performance.

Quantifying how life-histories arise from trait effects on whole-plant costs and benefits of

organs, from constraints and trade-offs governing resource budgets, and from age-dependent

shifts in resource allocation should better elucidate the true relationships between traits and

organismal performance and thereby improve the mechanistic understanding of life-history

variation (Kleyer & Minden, 2015; Swenson *et al*., 2020). While this research agenda is highly

promising, it faces considerable methodological challenges. First, it requires the quantification of

variables that are very difficult to observe (such as whole-plant costs and benefits and resource

budgets). Secondly, relationships between traits, whole-plant resource budgets and whole-plant

performance are likely to be non-linear. A promising approach to overcome these challenges are

latent state-space models (Clark, 2005). Latent state-space models can infer unobservable states

and incorporate non-linear functional relationships resulting from physiological and ecological

principles (such as allometric relationships or the principle of mass conservation; Enquist *et al*.,

1999).

Here, we develop a Trait-Resource-Life-History (TRL) framework for latent state-space models

to infer how functional traits shape whole-plant resource budgets and life-histories. In the

following, we first outline the general concept of this framework. We then apply the TRL

framework to the genus *Protea* from the South African Greater Cape Floristic Region (GCFR), a

global hotspot of woody plant diversity (Myers *et al*., 2000; Born *et al*., 2007). Members of this

genus exhibit a remarkable amount of variation in functional traits and life-history strategies

(Schurr *et al*., 2012) and are well suited to elucidate long-term dynamics of resource budgets and

life-history components. To this end, we develop a hierarchical Bayesian latent state-space

version of the TRL framework that we parameterise from data on the life-history schedules of

600 *Protea* individuals. We use this empirically-parameterised TRL model to address three key

questions of functional ecology and life-history theory: (1) do functional trait values that increase

the benefits of resource investment into a life-history component also increase the resource costs

of this investment? (2) What are the relationships between organ-level costs and benefits, whole-

plant resource budgets and whole-plant performance, and do these relationships change with

plant age? (3) Can functional traits be used to predict whole-plant performance and key life-

history characteristics? Building upon the *Protea* example, we then discuss the value of the TRL

framework for an in-depth understanding of plant life-histories, the application of the framework

to other study systems and the potential for its integration into process-based models of

community dynamics across trophic levels.

A Trait-Resource-Life-history (TRL) framework

The general structure of the TRL framework (Fig. 1) reflects a fundamental assumption of life-

history theory: in a given time interval, an individual acquires resources and can then allocate

them to different life-history components such as vegetative growth, reproduction, parental care,

survival and future resource acquisition (Stearns, 1992). Traits play a central role in the TRL

framework as they determine the time-invariant resource costs and performance benefits of

investment into a life-history component (Fig. 1). Since traits measure these costs and benefits at

organ level, an implementation of the TRL framework has to track the amount of different organs

per plant in order to quantify plant-level resource budgets and convert these into whole-plant

life-history components (Fig. 1).

Latent state-space modelling enables one to link this general framework to multiple types of data

(Clark, 2005). In general, measurements of functional traits and proxies for life-history

components are readily available whereas direct measurements of resource budgets are more

difficult to obtain. In many cases, the resource will thus be modeled as a hidden aggregate

variable that mediates trade-offs between life-history components rather than explicitly

describing amounts of a single compound (Tonnabel *et al*., 2012). Modeling resource acquisition

as a function of the amount of resource-capturing organs (e.g. leaves or roots) and expressing

resource amounts in units of an easily measurable life-history component (e.g. wood biomass

production), helps to constrain the model to biologically plausible parameter ranges and to

improve its statistical identifiability.

Application of the TRL framework to *Protea* plants

We applied the TRL framework to 20 species and 2 interspecific hybrids of the genus *Protea*

(Proteaceae). For simplicity, all study taxa are referred to as 'species'. They are endemic to the

South African Greater Cape Floristic Region and frequently dominate fire-prone fynbos

shrublands in this global biodiversity hotspot (Rebelo, 2001). These study species share the same

fire-driven life cycle and general growth architecture. Their regular growth pattern and the fact

that they form canopy seed banks enables one to reconstruct the temporal dynamics of growth,

reproduction and maternal care for multiple years into the past (Walter *et al*., 2023). Despite this

commonality, *Protea* species exhibit remarkable variation in life-histories and functional traits

(Schurr *et al*., 2012), with adult plant height and seed mass varying by more than an order of

magnitude (Treurnicht *et al*., 2016). In fact, Linnaeus named the genus *Protea* after the deity

Proteus who – according to Greek mythology – could take any form. *Protea* species are thus well

suited to test the TRL framework.

All study species are serotinous nonsprouters: fire kills them as adults but triggers the release of

seeds from fire-protected infructescences (hereafter cones) borne in the adult canopy (Bond *et*

al., 1984; Lamont *et al*., 1991). Almost all seeds then germinate in the first year after fire once

soil moisture and temperature are conducive (Lamont & Groom, 1998). Consequently, our study

species form even-aged stands. Within the first few years, *Protea* plants develop a deep taproot

that accesses reliable water supply in deeper soil layers (Lamont & Groom, 1998; Cramer et al.,

2014). Thereafter, they are very likely to survive until the next fire, unless this fire arrives

exceptionally late (Bond, 1980; West *et al*., 2012; Treurnicht *et al*., 2016). Growth and survival

during the critical juvenile phase increase with seed nutrient reserves (Cooksley *et al*., 2023).

Protea individuals show a regular growth pattern (Fig. 1) that is driven by strong seasonality, in

particular in terms of rainfall seasonality in the Mediterranean-type climate found across most of

the region (Bradshaw & Cowling, 2014). The branch increments produced in a given growth

bout are separated by branching points or by 'nodes' (branch segments with a high density of

leaves that are well visible by dense leaf marks even after leaves are shed). Branches typically

produce one increment per year so that the number of increments along major growth axes can

be used to age plants (Bond, 1985; Carlson *et al*., 2011; Treurnicht *et al*., 2016) although

individuals growing in very benign conditions can consistently show either two or three branch

increments per year. This regular growth pattern makes it possible to age individual branch

increments and the subtending leaves, inflorescences and cones (Fig. 2; Walter *et al*., 2023).

Moreover, it enables the reconstruction of past vegetative growth (see *Reconstructing whole-*

plant vegetative growth).

Protea species are evergreen, retaining their sclerophyllous leaves for several years (Rebelo,

2001). While light typically does not limit plant growth in fynbos ecosystems (Cramer *et al*.,

2014), Proteaceae are capable of using photosynthates to acquire limiting soil nutrients (notably

P and N). They achieve this by producing root clusters (or proteoid roots) consisting of a high

density of rootlets that actively excrete large amounts of carboxylates and phosphatases to

mobilize inorganic and organic P (Shane & Lambers, 2005; Cramer *et al*., 2014; Lambers *et al*.,

2015). Proteaceae can thereby effectively convert light into the acquisition of limiting soil

nutrients.

Reproductive maturity is reached after several years but the study species differ considerably in

age of maturity (Midgley & Rebelo, 2008). *Protea* plants are hermaphroditic, producing robust

inflorescences that contain multiple florets (again with substantial interspecific variation in

inflorescence size and floret number; Rebelo, 2001). Each floret contains a single ovule that can

develop into a single-seeded fruit. Inflorescences produce copious amounts of nectar (Nottebrock

et al., 2017) and are pollinated by a large diversity of insects and birds (Neu *et al*., 2023a).

After flowering, inflorescences turn into woody cones that can remain closed for several years

(Bond, 1985). Our study species differ in degree of serotiny, which varies from weak to moderate

(in comparison to strongly serotinous Australian Proteaceae, Lamont & Groom, 1998).

Proteaceae cones need to be supplied with water and carbon to remain closed. In fact, closed

cones continuously lose water and CO_2 so that cone maintenance implies costs (Cramer &

Midgley, 2009). Consequently, serotiny constitutes a case of maternal care in plants (Harris &

Pannell, 2010; Tonnabel *et al*., 2012). Cones open when the water supply to them stops because

the supporting branch died, because seed predators damaged xylem vessels or because the

mother plant was killed in a fire. Open cones remain on the plant for many years after, and scars

of abscised cones remain visible. Seedlings germinating from seeds released between fires face

intense competition from established vegetation and have low recruitment success (Lamont *et*

al., 1991). Moreover, our study species do not form persistent soil seed banks. Seeds that are

released from the canopy seed bank after fire and germinate in competition-poor post-fire

environments have relatively high establishment success. Consequently, the size of a plant's

canopy seed bank (all fertile seeds contained in closed cones) at the time of fire closely matches

its lifetime reproductive output (Treurnicht *et al*., 2016).

The canopy seed banks formed by *Protea* species hold a substantial proportion of the above-

ground nutrient pool in fynbos ecosystems (in particular for P, N and K; Low & Lamont, 1986).

This nutrient concentration results from the high nutrient acquisition ability of *Protea* plants (see

above) and makes their canopy seed banks attractive to insect seed predators that complete their

larval development inside *Protea* cones (Nottebrock *et al*., 2017). Seed predator communities

comprise a limited number of species (Coleoptera, Lepidoptera and Diptera), most of which

infest many different *Protea* species (Neu *et al*., 2023a). The majority of the seed predators are

borers that bite through the woody involucral bracts to enter and/or leave cones (Coetzee &

Giliomee, 1987).

Reconstructing whole-plant vegetative growth

The reconstruction of past whole-plant wood and leaf biomass of *Protea* individuals rests on four

findings (Fig. 3): (1) our study species conform to da Vinci's rule on the conservation of stem

cross-sectional area (Leonardo's notes 394, 395, Richter, 1970), (2) there are strong species-

specific allometric relationships between trunk basal area and total stem length, (3) the mass of

leaves produced on a branch segment is proportional to branch cross-sectional area (as predicted

by the pipe model theory, Shinozaki *et al.*, 1964), and (4) leaf retention can be described by

species-specific Weibull functions. In the following, we describe the evidence for these findings

and show how they can be used to reconstruct time series of whole-plant wood and leaf biomass

of *Protea* individuals.

Da Vinci's rule states that the total cross-sectional area of branches is conserved across branching

events (Richter, 1970), a prediction that is also derived from the pipe model theory (Shinozaki *et*

al., 1964). To test whether this holds in *Protea*, we measured the diameter of the mother branch

below a branching point and the diameters of all daughter branches above the branching point

using callipers. We then calculated branch cross-sectional area as the area of a circle. This was

done for 876 branching points across 425 individuals of all study species. We covered a wide

range of branch sizes with mother branch cross-sectional areas ranging from 0.05 cm^2 to 71.6

cm^2. The obtained data show a very tight relationship between mother branch cross-sectional

area (A_M) and the summed cross-sectional areas of the daughter branches (ΣA_D; Fig. 3a). In fact, a

standardised major axis regression (SMA, Warton *et al.*, 2007) relating log-transformed A_M and

ΣA_D estimated a relationship that is very close to identity, with a power-law coefficient of 0.004

(95% confidence interval: -0.017 – 0.026) and a power-law exponent of 1.007 (95% confidence

interval: 0.998 – 1.016). We thus find strong support for da Vinci's rule in *Protea*. Consequently,

the wood volume of a plant at time *t* can be approximated as the volume of a cylinder with basal

area equal to trunk basal area A_t and height equal to summed stem length SSL_t (i.e. the sum of

all stem length increments; Fig. 2).

To quantify allometric relationships between trunk basal area A_t and summed stem length in the

main growing direction SSL_t, we measured these quantities for 1,041 plants of all study species

(range of plant ages: 1-33 years). We then related log-transformed A_t to log-transformed SSL_t

with a SMA regression including random effects of species identity on the intercept (Brose *et al.*,

2019). This model of species-specific $A_t \sim SSL_t$ allometries had a high goodness of fit ($r^2 = 0.85$;

Fig. 3B; Appendix S1: Figure S1). Consequently, trunk basal area at any time in the past $t - \Delta t$

($A_{t-\Delta t}$) can be predicted from the observed summed stem length at this time point ($SSL_{t-\Delta t}$). With

the above-mentioned cylindrical approximation of wood volume, this enables the reconstruction

of whole-plant wood volume at time $t - \Delta t$ (Fig. 3). Multiplying this wood volume by population-

specific wood density, *WD*, yields whole-plant wood biomass as

$M_{t-\Delta t} = \alpha \cdot SSL_{t-\Delta t}^{\beta+1} \cdot WD$ (1),

where α and β are the species-specific allometric coefficients and exponents, respectively, of the

$A_t \sim SSL_t$ – relationship. Consequently, wood biomass production in time interval Δt is

Wood Mass$_{\Delta t} = \alpha \cdot SSL_t^{\beta+1} \cdot WD - \alpha \cdot SSL_{t-\Delta t}^{\beta+1} \cdot WD$ (2).

According to the pipe model theory (Shinozaki *et al.*, 1964), the leaf mass produced on a branch

is proportional to branch cross-sectional area. We tested this by estimating the allometric

relationship between the leaf mass produced on a one-year-old branch and the branch's cross-

sectional area. To this end, we collected all leaves from the branch and dried them at 70°C for 48

h before weighing them. In case leaves had been shed from the branch, we visually estimated the

proportion of leaf mass lost (accounting for the loss of both entire leaves and leaf parts). This

served to calculate the total dry mass of leaves produced on the branch. Additionally, we

calculated branch cross-sectional area from calliper measurements of branch diameter.

Measurements on multiple branches per plant (mean: 2.7 branches) were then summed for each

plant. We did this for 424 plants of all study species. A SMA regression of log-transformed total

leaf mass and cross-sectional area with random effects of species and site on the intercept

estimated a median allometric exponent of 1.0232 (CI: 0.9049 – 1.1459). Hence, there is a strong

proportionality between leaf mass production and branch cross-sectional area (Fig. 3c; Appendix

S1: Figure S2). In combination with da Vinci's rule, the total leaf mass that a plant produces in

year t, LP_t, is thus proportional to the plant's trunk basal area A_t and to SSL_t^β

$$LP_t = \alpha_{LP} \cdot A_t = \alpha_{LP} \cdot \alpha \cdot SSL_t^\beta \quad (3)$$

where α_{LP} is the species-specific coefficient relating leaf mass production to branch cross-

sectional area.

Since *Protea* plants are evergreen, their standing leaf biomass depends not only on leaf

production in a given year but also on leaf retention from previous years. To characterize leaf

retention, we visually estimated the proportion of leaf mass retained on one year old branches of

458 individuals of all study species. Along the same branches, we estimated the branch age at

which the proportion of retained leaf mass drops to 50% (cf. Treurnicht *et al*., 2020). These

estimates of median leaf retention time were combined with corresponding measures from the

FYNBASE database (Schurr *et al*., 2007; Treurnicht *et al*., 2020) in TRY (Kattge *et al*., 2020) to

calculate species-level medians of median leaf retention time. We fitted species-specific Weibull

functions to all leaf retention data for a species, since the Weibull distribution provides good

descriptions of leaf lifespan in evergreen Mediterranean shrubs (Dungan *et al*., 2008).

Specifically, we used non-linear least squares to fit the complement of the Weibull's cumulative

density function, which describes the proportion of leaf mass retained on branches of age k, LR_k

as $LR_k = \exp\dot\iota$, where a and s are the shape and scale parameter of the Weibull distribution,

respectively. From the resulting species-specific leaf retention functions (Fig. 3d; Appendix S1:

Figure S3), one can reconstruct a plant's *total* standing leaf mass, *TLM*, at time $t - \Delta t$ in the past

as:

327
$$TLM_{t-\Delta t} = \sum_{j=1}^{t-\Delta t} LP_j LR_{t-\Delta t-j} \qquad (4)$$

In summary, species-specific scaling relationships enable one to effectively reconstruct the

vegetative growth of *Protea* plants. In particular, measurements of stem length increments in past

growth bouts make it possible to calculate proxies of whole-plant aboveground wood biomass,

leaf biomass production and total leaf biomass at multiple time points in the past (Fig. 3).

<u>Measurement of life-history components</u>

We took time-stratified measures of vegetative growth, reproduction and maternal care for 600

individuals of the 22 *Protea* species on 21 study sites. By comparing the known time of last fire

at each study site to the number of growth increments per focal plant (Bond, 1985; Treurnicht *et*

al., 2016), we confirmed that the study individuals had established after the last fire. For 160

plants from five sites, the number of growth increments was either two- or three-times larger

than the site's post-fire age, indicating that the plants had undergone two or three growth bouts

per year, respectively. By combining known post-fire ages of each site and counts of growth

increments, we were thus able to age both individual increments and entire plants.

To reconstruct the vegetative growth of each individual, we measured the length of each growth

increment (the distance between nodes) along a representative stem in the plant's main growing

direction (starting from the youngest increment). For most individuals, these increments

represent annual growth. However, in seeing two or three growth bouts per year, we took these

measures at higher temporal resolution. In older stem sections, for which growth increments

could not be delineated unambiguously, we additionally recorded the total stem length from the

last resolvable increment to trunk base as a measure for total plant growth up to this time point.

Reproductive performance was quantified by cone production, retention of closed cones, seed set

in sampled closed cones, and cone defence from predation. Cone production and closed cone

retention was measured by counting the number of cones produced by each individual at each

time-step (at the same temporal resolution as for growth increments), and whether they were

closed, open or abscised at the time of observation. In certain older plants, it was not possible to

assess reproduction in the first years of life; for these initial years, cone number was recorded as

missing data. In very large plants, cones were counted in a representative subset of the canopy

and these counts were then extrapolated to the entire plant. From each cohort of closed cones on

an individual, we sampled one closed cone to measure seed set (seed number per cone). To this

end, the cone was cross-sectioned at the stratum where seeds are borne, and the number of fertile

seeds was counted (Treurnicht *et al*., 2016). In each cone, we also visually estimated cone

defence as the proportion (in steps of 5%) of the total seed area not consumed by seed predators.

In summary, our time-stratified measures of life-history components reached back 1-30 years,

thus encompassing much of the total lifespan of our focal plants, which ranged from 3-33 years

at the time of sampling.

<u>Measurement of functional traits</u>

To quantify the costs and benefits of resource allocation to different life-history components, we

measured four key functional traits representing investment into growth and reproduction, and

return on this investment: specific leaf area (SLA), leaf N concentration, cone mass and seed N

content (Appendix S1: Figure S4; Table S1). Additionally, we measured wood density of one

year old stem wood as a key determinant of whole-plant wood biomass (see eqn. 1). For SLA,

leaf N concentration and wood density, we followed the standard protocols of Pérez-

Harguindeguy *et al*. (2013; leaf N concentration was measured using elemental analysis (Vario

EL Cube, Elementar Analysensysteme GmbH, Langenselbold, Germany)). Cone dry mass was

measured from closed cones that were oven-dried at 70°C for 48 h and then weighed using

precision scales. Seed dry mass (g) was measured from fertile seeds oven-dried at 70°C for 48 h,

and weighed with precision scales. Seed N concentration (% mass) was determined by elemental

analysis of finely milled dried seeds (Vario EL Cube, Elementar Analysensysteme GmbH,

Langenselbold, Germany). Seed N content was then calculated as the product of seed mass and

seed N concentration. In addition, we used measurements of leaf and seed N concentration from

the FYNBASE database. Note that seed N and P contents are strongly correlated in these species

(r = 0.86; Cooksley *et al.*, 2023). Trait measurements were averaged at population level, except

for leaf N which were averaged at species level.

A Trait-Resource-Life-history (TRL) model for *Protea*

Based on the knowledge of the life cycle of our study species, we adapted the general TRL

framework (Fig. 1) in four ways (Fig. 4). First, we did not explicitly model resource investment

into survival (since all study species have very high rates of inter-fire survival). Secondly, we

resolved reproduction into cone and seed production, maternal care in the form of cone

maintenance and defence against seed predation. Thirdly, we modeled resource acquisition as a

function of leaf mass because *Protea* plants are capable of converting photosynthates into

acquisition of soil nutrients, water and pollination service (see above) and because leaf biomass

can be reconstructed for past time points (see above) and is allometrically related to root biomass

(Niklas, 2004). Fourthly, we expressed resource amounts in units of wood biomass which can

also be reconstructed for our study species (see above), with conversion parameters for the

relative costs of other tissues. In brief, this TRL model is an integration of organ-level costs and

benefits informed by traits, whole-plant resource pools at each time-step, and proportional

allocation of resources to wood, leaves, cones, seeds and serotiny, which we detail in the

following.

Organ-level costs and benefits are defined with respect to resources (in units of equivalent grams

of woody biomass). Costs of leaves, cones, seeds and serotiny are the resources required to

produce one gram of leaves, one cone or seed, or to maintain one closed cone for one year,

respectively. The benefits of leaves, cone defence, seeds and serotiny represent the respective

return on these resource investments. Leaf benefit is the amount of resources acquired per gram

of leaf and year. Seed benefit is the resource pool a seedling derives from the seed. Cone defence

benefit is the per-year and per-cone-area probability of successful defence against seed predators.

Serotiny benefit is the per-year probability of a cone remaining closed (and thus potentially

contributing to post-fire regeneration).

Costs and benefits are modelled as functions of key traits for the respective organ, with random

intercepts for species, site and individual (Table 1). While costs and benefits thus vary between

individuals, we assume that they do not change with the age of an individual. Leaf cost and

benefit are informed by SLA and leaf N concentration (Wright *et al.,* 2004). Cone cost is a

function of cone dry mass, as the dense wood and bracts that make up most of a cone and afford

protection for seeds are likely the substantial cost of this organ. Cone defence benefit is informed

by cone dry mass, representing the cost to the parent plant of defending seeds, and by seed N

content, representing the potential reward for granivorous insects. Seed cost and benefit are

functions of seed N content, a key limiting nutrient in impoverished fynbos soils and a good

predictor of juvenile performance (Cooksley *et al.*, 2023). We model serotiny cost and benefit as

functions of cone dry mass, assuming that costs of water replacement and carbon supply are

proportional to cone dry mass (water content and dry mass of cones are highly correlated:

Pearson's r=0.87). Since resources are measured in units of wood biomass, we do not explicitly

estimate the cost of wood production, but rather take it to be wood density.

Whole-plant performance components of individual i in time-step t are latent variables modelled

as a function of the resource pool, R, proportional resource allocations, Θ, to each component,

and the respective organ-level cost. The performance variables are grams of wood and leaves

produced, number of cones and seeds produced, as well as number of closed cones cared for.

We first model the reproductive maturity status, *Mature*, of a plant as a function of R

$\text{Mature}_{t,i} \sim \text{Bernoulli}\left(\text{p\_mat}_{t,i}\right)$ (5),

$\text{logit}\left(\text{p\_mat}_{t,i}\right) = \alpha\_m + \alpha\_m\_sp_{\text{species}[i]} + \alpha\_m\_si_{\text{site}[i]} + \left(\beta\_m + \beta\_m\_sp_{\text{species}[i]}\right) \cdot \ln\left(R_{t,i}\right)$ (6),

where α_m_sp and α_m_si are zero-mean normal random intercepts for species and site,

respectively, and β_m_sp are zero-mean normal random slopes for species.

Proportional resource allocation to wood, leaf, cone and seed production and cone maintenance

forms a five-element allocation vector $\boldsymbol{\Theta} = (\Theta_W, \Theta_L, \Theta_C, \Theta_S, \Theta_M)$ defined by

$\Theta_{t,i} \sim \text{Dirichlet}\left(\left(1, 1, \text{Mature}_{t,i}, \text{Mature}_{t,i}, \text{Mature}_{t,i}\right) + 10^{-100}\right)$ (7),

such that allocations are assigned a flat Dirichlet prior, except that allocations to cones, seeds and

serotiny are essentially zero if an individual is immature. We thus assume that all resources in a

given time-step are used, with no carry-over to the next time-step.

The five whole-plant latent performance variables are then defined by:

$\text{Performance}_{p,t,i} = \dfrac{\Theta_{p,t,i} \cdot R_{t,i}}{\text{cost}_{p,i}}$ (8),

where p denotes the respective performance variable.

An individual's resource pool in the first time step is a function of seed benefit

$\ln(R_{1,i}) = \ln(\text{Seed benefit}_i) + (\beta\_R1 + \beta\_R1\_sp_{species[i]} + \beta\_R1\_si_{site[i]}) \cdot \Delta t$ (9),

where β_R1 and the zero-mean normal random slopes for species and site of β_R1_sp and

β_R1_si define resource growth from seed resources to the resource pool in the first time-step, Δ

t (1, 1/2 or 1/3 year, depending on the number of growth bouts per year, see above).

The resource pool of an individual in subsequent time-steps is a random variable, the expected

value of which depends on leaf benefit and *TLM* in the current time-step. Thus:

$\ln(R_{t+1,i}) \sim \text{Normal}(\ln(\text{Leaf benefit}_i) + \ln(TLM_{t,i}), \sigma)$ (10),

for all $t > 1$, where σ is the standard deviation of the residual error.

The expected number of closed cones on a plant, *N closed cones*, is a function of *N cones cared*

for and *Serotiny benefit*, assuming all closed cones on a plant in a given year receive the same

allocation to serotiny

$\text{N closed cones}_{t,i} = \text{Serotiny benefit}_i \cdot \text{N cones cared for}_{t,i}$ (11).

Cone defence against granivorous insects is defined as the probability, p_def, that a given 5%

section of the total seed basal area is not eaten. This is modelled as a function of cone defence

benefit (*Defence benefit*), site age at the time of cone production (*SA*, reflecting post-fire changes

in predation; Walter *et al.*, 2023) and cone age, *CA*, the time a cone has been on a plant before

the year of observation:

$\text{logit}\left(\text{p\_def}_{t,i}\right)=\text{logit}\left(\text{Defence benefit}_i\right) + \beta\_sa \cdot \ln\left(\text{SA}_{t,i}\right) + \beta\_ca \cdot \ln\left(\text{CA}_{t,i}\right)$ (12).

We link latent performance variables to age-stratified observations of performance via stochastic

observation equations. The number of cones observed, *N cones obs*, was modelled as

$\text{N cones obs}_{t,i} \sim \text{Poisson}\left(\text{N cones}_{t,i} \cdot \text{Proportion counted}_i\right)$ (13),

where *Proportion counted* is an offset for the proportion of an individual for which cones were

counted. Observed cone defence, *Cone defence obs*, was modelled as

$\text{Cones defence obs}_{t,i} \sim \text{Binomial}\left(\text{p\_def}_{t,i}, N = 20\right)$ (14).

The number of seeds observed in a cone, *N seeds obs*, was modelled as a function of cone

defence as well as whole-plant seed and cone production

$\text{N seeds obs}_{t,i} \sim \text{Poisson}\left(\dfrac{\text{Cone defence}_{t,i}}{20} \cdot \dfrac{\text{N seeds}_{t,i}}{\text{N cones}_{t,i}}\right)$ (15).

The total number of closed cones observed on a plant, *N closed cones obs*, was modelled as

N closed cones obs$_{t,i}$ $\sim$ Poisson $\left(\text{N closed cones}_{t,i} \cdot \text{ Proportion counted}_i\right)$ (16),

and the observed cone-age-stratified number of closed cones, *N closed cones obs str*, was

N closed cones obs str$_{t,i}$ $\sim$ Binomial $\left(\text{Serotiny Benefit}_i^{Y_i} \text{ , N cones obs}_{t,i}\right)$ (17),

where Y_i is the age of cones of that cohort in years. By rearranging equation 2, we obtain

$\text{SSL}_{t,i} = \left(\dfrac{\text{Wood Mass}_{t,i}}{\alpha \cdot \text{WD}_i} + \text{SSL}_{t-1}^{\beta+1}\right)^{1/(\beta+1)}$　　　　(21),

so that the expected stem length increment, SL_t, is

$\text{SL}_{t,i} = \left(\dfrac{\text{Wood Mass}_{t,i}}{\alpha \cdot \text{WD}_i} + \text{SSL}_{t-1}^{\beta+1}\right)^{1/(\beta+1)} - \text{SSL}_{t-1}$　　　　(22).

Since the allometry of seedlings differs from that of older plants (Niklas, 2004), we allowed the

allometric relationship between SSL_t and A_t (Fig. 3) to differ between the first time-step and later

time-steps. To this end, we estimated an α of seedlings that was multiplied by a species-specific

random effect. Since in our study species, variance in stem length increments between branches

of a given cohort is approximately equal to the mean (see Appendix S1: Figure S5), we modelled

observed stem length increment as

SL obs$_{t,i}$ $\sim$ Normal $\left(\text{ SL}_{t,i} \text{ , sd} = \sqrt{\text{SL}_{t,i}}\right)$　　　(23).

We informed resource-based estimates of leaf production with the proportional relationship

between leaf mass and branch cross-sectional area by combining equations 3 and 9, such that

Leaf Mass$_{t,i} = \Theta_L \cdot \text{ R}_{t,i} \approx \alpha_{LP} \cdot \alpha \cdot \text{ SSL}_t^{\beta}$　(24).

By ln-transforming, incorporating uncertainty and rearranging to maintain a proper acyclical

graph, we obtain

$\ln\left(\Theta_L \cdot \text{ R}_{t,i}\right) - \ln\left(\alpha_{LP} \cdot \alpha \cdot \text{ SSL}_t^{\beta}\right) + \varepsilon = 0$　　　　　(25),

where ε is the residual error of this relationship, given an informative prior based on our

empirical estimate (see 'Reconstructing whole-plant vegetative growth'). Similar to above, we

allowed the proportionality between leaf mass and cross sectional area to vary for plants in the

first time-step, adding random intercepts for species and site to equation 25 for $t = 1$.

<u>Fitting the TRL model to data</u>

All data preparation, calculations and statistical analyses were performed in R 4.0.0 (R Core

Team, 2020). Our model was fitted to data via Markov chain Monte Carlo (MCMC) sampling,

using JAGS 4.3.0 (Plummer, 2003) interfaced via the R package saveJAGS 0.0.4.9002

(Meredith, 2021). Three MCMC chains were run, with 20,000 iterations of adaptation and

40,000 iterations of sampling, thinned to every 20th iteration, resulting in 6,000 samples from the

joint posterior distribution. We used weakly informative priors for all parameters, except for the

effect of seed N content on seed cost, and for the residual error of leaf mass per branch area,

which were informed by data for these processes. JAGS and R code used to fit the TRL model,

including prior distributions, are given in Appendix S2. Convergence to the stationary posterior

distribution was determined by the potential scale reduction factors for all process parameters

being < 1.1, and by visual inspection of trace plots.

<u>Analysis of the parameterised model</u>

Posterior predictions of observed performance measures were done as 'one-step-ahead'

predictions, conditioned upon the previous resource state. In short, we simulated resource

amounts from the whole-plant resource acquisition state-transition equation (equation 10) for the

previous time step and then drew all other parameters from the joint posterior distribution.

To assess whether functional trait values that increase the benefits of resource investment into a

life-history component also increase the resource costs of this investment, we compared posterior

distributions of effect sizes of traits on paired costs and benefits.

To assess relationships between organ-level costs and benefits, whole-plant resource budgets and

whole-plant performance, we calculated Pearson's correlation coefficients between posterior

medians of costs and benefits, resource pools, allocations and yearly whole-plant performance.

We conservatively chose to calculate these correlations beginning from the age of 6 years, as

prior to this numerous posterior medians were zero, distorting correlations, and ending at 18

years, as few species had older individuals in the dataset. Prior to calculating correlation

coefficients, costs and benefits were ln-transformed (except for cone and serotiny benefits which

were logit-transformed), cone, seed and closed cone number were $ln(x+1)$-transformed, leaf and

wood masses as well as traits were ln-transformed.

To address whether functional traits can predict whole-plant performance measures, we

calculated Pearson's correlation coefficients between them for ages from 6 to 18 years. We also

calculated posterior median total reproductive output (seed bank size) for each individual at each

time-step, integrating over cone and seed production, cone defence and cone maintenance, and

calculated correlation coefficients of this fitness proxy with traits across the above age range.

Finally, we calculated Pearson's correlations between traits and age of maturity (the earliest age

at which the posterior median of seed bank size was ≥ 1).

Results

<u>Predictions of life-history components and resource budgets</u>

The TRL model predicted most performance measures with high accuracy and little bias (Fig. 5).

For the four performance measures directly affected by resource allocation (stem length

increment, cone number, seed number and closed cone number), the squared correlation

coefficient (r^2) between observed and posterior-median one-step-ahead predicted values ranged

from 0.73 to 0.95 (Fig. 5a-d). Predictions of cone defence (which are not directly informed by

resource budgets) were still reasonably accurate ($r^2 = 0.54$) (Fig. 5e).

In addition to predicting observable performance components, the TRL model also estimates the

essentially unobservable dynamics of whole-plant resource budgets (Fig. 6a). Specifically, it

permits the individual- and age-specific quantification of resource acquisition that leads to

growth of the resource pool and of the allocation of this resource pool to different components

(Fig. 6a). Moreover, the TRL estimates the build-up of total reproductive output (which for

Protea individuals is measured as the size of the canopy seed bank; Fig. 6b) which integrates

over cone and seed production, cone defence and serotiny.

<u>Trait effects on organ-level costs and benefits of life-history components</u>

Functional traits affect organ-level costs and benefits of life-history components by-and-large in

the same direction, such that trait values conferring greater benefits also increase costs (Fig. 7).

Leaf construction cost increases with SLA, as does the rate of resource acquisition (Fig. 7a).

Heavier cones are more costly to produce and incur greater serotiny costs, but are more likely to

remain closed and to be defended from seed predators (Fig. 7b,d). The greater cost of N-rich

seeds is rewarded by the greater resource pool available to offspring (Fig. 7c) although this

trades-off with an increased risk of seed predation (Fig. 7b).

<u>Relationships between organ-level costs and benefits, whole-plant resource budgets and whole-</u>

<u>plant performance</u>

Resource pools increase strongly with the amount of resources plants acquire from the seed, even

well into the adult stage, and with the rate of resource acquisition from leaves (Fig. 8a). Resource

pools also increase with leaf costs (since costly leaves are also more beneficial) and moderately

with costs of cone production and serotiny (Fig. 8b). These patterns by-and-large hold with age,

although the relationship between leaf cost and resource pools does tend to get stronger with

time.

The larger the resource pool, the greater the share of resources plants allocate to reproduction

and maternal care, and the lesser the share they allocate to production of leaves (Fig. 8c). These

relationships only slightly weaken with age, at least for reproductive and maternal allocations. In

turn, resource investment (the product of resource pools and resource allocation) is closely

correlated with the respective life history component at all ages (Fig. 8d).

Relationships of traits to whole-plant performance and life history

Leaf and seed benefits play a central role for resource budgets and performance at the whole-

plant level (Fig. 8). Hence, one may expect that SLA and seed N content, which shape these

benefits (Fig. 7), should be correlated with whole-plant performance and other key life-history

characteristics. SLA is indeed a reasonable predictor of all performance measures at early ages,

but less so in later years, particularly for reproductive measures (Fig. 9a). Seed N content is,

however, only weakly correlated with performance (Fig. 9b): leaf and seed production and

serotiny decrease mildly with seed N, but wood and cone production are largely unrelated to seed

N.

Age of maturity as a key life-history characteristic shows clear relationships to a number of

functional traits (Fig. 10;Appendix S1: Figure S6). In particular, it is negatively related to SLA

across all study species and sites (Fig. 10a). Moreover, age of maturity is positively related with

seed N content, although this relationship is weaker than for SLA (Fig. 10b). Traits shaping the

costs and benefits of vegetative and reproductive organs thus play a substantial role for the pace

of *Protea* life-histories. Moreover, correlations between traits and age of maturity have

considerable conservation relevance since climate change is expected to increase fire frequency

in much of the study region (Wilson *et al*., 2015). This should increase the extirpation risk of

late-maturing populations (Enright *et al*., 2015) and lead to a functional shift of *Protea*

communities towards species with higher SLA and lower seed N content. This is one example of

how the TRL framework can contribute to a life-history based assessment of global change

impacts on biodiversity (Ohse *et al*., 2023).

Discussion

The TRL framework enables the formulation of integrated process-based models that predict the

temporal dynamics of reproductive and vegetative performance throughout the lifetime of long-

lived plants. A TRL model describes how these temporal dynamics are shaped by the costs and

benefits of plant organs, and by the dynamics of plant-level resource acquisition and allocation

(Fig. 1). Importantly, the model can be parameterised with data on plant performance and

functional traits (Fig. 4). This has three key benefits. First, we can predict performance measures

with notable accuracy, throughout the lifetime of individuals up to the time of observation (Fig.

5). Second, by using a latent-variable modelling framework, we can learn about unmeasured

variables that shape this process, namely, we can infer the resource costs and benefits of different

plant organs (Fig. 7), the dynamics of resource budgets (Figs. 6,8), and unobserved whole plant

performance measures such as the mass of leaves, the number of seeds produced annually, and

total reproductive output (Fig. 8d,9). Third, by being able to infer these unmeasured variables,

we can tease apart the processes linking organ-level traits to whole-plant-level performance and

life-history. In the following sections, we discuss how the TRL model helps to infer the dynamics

of whole-plant resource budgets and performance, how it contributes to a resource- and trait-

based understanding of plant life-histories, how it can be applied to other plant systems and how

it may promote a resource-based understanding of community dynamics.

<u>Inferring the dynamics of whole-plant resource budgets and performance</u>

Understanding how functional traits shape life histories requires an understanding of how organ-

level traits scale-up to the lifetime dynamics of whole-plant resource acquisition, resource

allocation and performance. By using hierarchical Bayesian latent variable modelling, the TRL

framework can infer essentially unobservable variables (e.g. resource budgets), close gaps in

performance observations for certain years (e.g. early growth and reproduction), and estimate

performance quantities that are challenging to measure (e.g. the total mass of leaves produced

per year). Further, by characterising all elements of a plant's resource economy in consistent

units (equivalents of woody biomass), with appropriate conversion parameters for other organs,

we can directly compare the processes shaping life-histories in a single currency. With our case

study, we have shown that the TRL framework can reconstruct the dynamics of resource budgets

and performance from germination up to 33 years of age for the oldest individuals in the *Protea*

dataset.

The construction costs of organs are often inferred from their mass and chemical composition at

the time of harvest, using fixed conversion parameters to express costs in units of glucose

(Poorter, 1994; Williams *et al*., 1997). The TRL framework takes an inverse approach, estimating

net costs and benefits over the lifetime of an organ, based on the effect of organs on whole plant

resource pools. This inherently accounts for the costs of organ maintenance, and the benefits of

nutrient recycling (e.g. from senescing leaves) that are not considered in once-off construction

cost calculations (Poorter, 1994). The inverse approach taken by TRL also accounts for the

different costs of nutrient acquisition experienced by different species in different environments,

a prohibitive task with fixed conversion parameters (Poorter, 1994). Further, within this

integrated framework one can estimate the benefits of leaf production for resource acquisition, of

seed resources for seedling growth, and of resource allocation to serotiny and to defence against

seed predation. While the first two of these could be estimated via extensive physiological

measurements, the presented integrative approach enables us to estimate these benefits from data

that are easier to obtain.

Resource- and trait-based understanding of plant life-histories

A resource-based understanding of life histories is facilitated by the TRL framework's

description of how organ-level costs and benefits shape whole-plant resource budgets and

performance (Fig. 1). Applying this framework to *Protea*, we found that higher resource

acquisition - be it initial acquisition from the seed or subsequent acquisition through leaves - is

linked to the build-up of larger resource pools at all ages (Fig. 8a). The size of the resource pool

is in turn linked to vegetative versus reproductive allocation: plants with more resources invest a

larger share of these resources into reproduction (Fig. 8c). Since performance is strongly

correlated with resource investment (the product of resource pool and resource allocation; Fig.

8d), a greater resource pool implies a disproportionate increase in reproductive performance.

Functional traits are relatively modest predictors of whole-plant performance (Fig. 9), even if

they play important roles for the organ-level costs and benefits that shape resource budgets and

life histories (Fig. 7). This is because multiple co-varying processes are involved in upscaling

from organ-level traits to whole plant resource budgets, which themselves vary through time and

are allocated to multiple dimensions of performance. For example, seed N content increases seed

benefits that are linked to greater whole-plant resource pools and performance, but it also

increases the costs of both seeds and cones (Figs. 7,8). Overall, this leads to negative and

relatively weak relationships between seed N content and all performance measures (Fig. 9b). On

the other hand, SLA as the key trait determining resource acquisition from leaves (Fig. 7), is

correlated with earlier maturity (Fig. 10a) and showed strongly positive relationships to all

performance measures at young to intermediate ages (Fig. 9a). However, the positive relationship

between SLA and total reproductive output clearly weakened for older ages (Fig. 9a). Hence,

traits do matter (Adler *et al*., 2013), but in ways that cannot necessarily be captured by simple

correlations with whole-plant performance (Kleyer & Minden, 2015; Messier *et al*., 2017;

Swenson *et al*., 2020).

The TRL framework should enable the detection of trade-offs at the level of organs, whole-plant

resource budgets and life histories. At organ level, we found clear trait-mediated trade-offs:

functional trait effects on the costs and benefits of an organ generally act in the same direction

(Fig. 7). Hence, if a plant is to gain the benefits of an organ, it has to invest resources to reap this

reward. This is a fundamental tenet of evolutionary life-history theory (Stearns, 1992) – resource

investment into an organ that is not sufficiently beneficial to fitness will be selected against.

When examining resource budgets, we also detected a clear trade-off between resource allocation

to growth versus reproduction (Fig. 8c). Such a growth-reproduction trade-off may give rise to a

life-history-level trade-off between early and late reproduction (Stearns, 1992; Salguero-Gómez

et al., 2016). This is expected if the earlier maturity of plants that start with many resources

reduces their subsequent resource acquisition and reproduction (Tonnabel *et al*., 2012). In our

case study, we found no evidence for this life-history trade-off: plants that matured earlier

generally also reproduced more later in life (Appendix S1: Figure S7). This may be because

plants performing well throughout their lives have continuously higher resource acquisition (van

Noordwijk & de Jong, 1986) that results from factors not considered in our analysis, e.g. because

they occupy more resource-rich microsites or experience less competition (Walter *et al*., 2023). A

comprehensive test for the trade-off between early and late reproduction should thus aim to

standardize resource availability and/or account for the competitive environment of plant

individuals. While we did not detect a trade-off between early and late reproduction in the *Protea*

system, there is evidence for a life-history trade-off between the quantity and quality of

offspring: plants with high seed N content mature later and produce fewer seeds (Fig. 9b, 10b),

but their seedlings have a higher survival probability (Cooksley *et al*., 2023).

<u>Application of the TRL framework to other plant systems</u>

Application of the TRL framework to other study systems will help to further develop the

framework and extend its general validity, advancing a general understanding of how functional

traits shape resource dynamics and performance of plants. This transfer to other study systems

requires data on functional traits as proxies for costs and benefits of resource investment into

organs, data on the temporal dynamics of life-history components and – potentially – data on the

temporal dynamics of resource budgets (Fig. 1). It seems worth noting that the parameterisation

of a TRL model using latent state-space modelling does not require a comprehensive coverage of

all of these data types at all times.

TRL considers functional traits as measures for the costs and benefits of investment into different

life-history components (Fig. 1). These life-history components are typically linked to a

particular organ type (such as cones, seeds, wood or leaves). Biomass or nutrient investment at

organ-level can be captured by a few well-established traits such as SLA, wood density or seed

(nutrient) mass. Hence, application of TRL models should typically require a limited number of

traits for which measurements are widely available (Kattge *et al*., 2020) or can be taken

relatively easily. Moreover, since a given trait often affects both the costs and the benefits of an

organ (Fig. 7; see also Westoby *et al*., 2002; Wright *et al*., 2004; Chave *et al*., 2009), application

of the TRL framework should typically require easily accessible data on a limited number of

traits.

Reconstruction of vegetative growth in *Protea* individuals relied on their regular growth pattern

(Fig. 3). Other species-rich genera in the Proteaceae show a similar growth pattern so that our

approach should be easily transferable to them (e.g. Australian *Banksia* and South African

Leucadendron; Lamont, 1985; Harris & Pannell, 2010). Beyond the specific case of Proteaceae,

vegetative growth of a broad range of tree species can be reconstructed by combining

dendrochronological data with allometric relationships between trunk diameter and wood or leaf

biomass (Babst *et al*., 2016). A similar approach should be feasible for herbaceous perennials

from seasonal environments that commonly form annual growth rings in the main root (Dietz &

Ullmann, 1997).

For long-lived plants, data on the temporal dynamics of reproduction are generally more difficult

to obtain than data on vegetative growth. In many serotinous plants, however, past reproduction

and maternal care can be quantified relatively easily. Serotiny is no exception in the plant

kingdom: it occurs in more than 40 genera that play important roles in both Mediterranean-type

ecosystems and coniferous forests (Lamont *et al.*, 1991; 2019). It thus seems both interesting and

relevant to apply the TRL framework to a broader range of serotinous species.

Non-serotinous trees arguably pose one of the biggest challenges for reconstructing lifetime

reproductive schedules. However, long-term records of individual reproductive output exist for

certain tree species (Bogdziewicz *et al.*, 2020). In other species, it is possible to reconstruct

components of past reproduction. For instance, annual cone production in pinyon pines (*Pinus*

edilis) can be reconstructed from cone abscission scars (Redmond *et al.*, 2016). Moreover,

remote sensing can be used to estimate current reproductive output. For instance, terrestrial lidar

scanning of a *Pinus pinea* stand quantified canopy characteristics that were informative of

between-individual variation in cone number and cone mass (Schneider *et al.*, 2020). Remote

sensing methods can thus yield snapshots of reproductive output for large numbers of

individuals. Such data may be used to infer reproductive schedules at population or species level

using individual-for-age substitution. Alternatively, the reproductive output of trees in a given

year can be estimated inversely by combining data on spatial distribution of adult trees and post-

dispersal seed densities (e.g. Clark *et al.*, 2019). Such estimates can be further improved by

marker-based genotyping of seeds and adults and by integrating dendrochronological data for

adults (Lamonica *et al.*, 2021).

In short-lived plants, it is possible to directly observe the temporal dynamics of reproduction and

vegetative growth. Notably, rapid developments in noninvasive plant phenotyping (Pieruschka &

Schurr, 2019) make it possible to efficiently observe the temporal dynamics of multiple life-

history components at high temporal resolution (e.g. Vasseur *et al*., 2019). Short-lived plants also

offer the possibility of experimentally controlling resource availability for individuals to

rigorously test for life-history trade-offs (see *Resource- and trait-based understanding of plant*

life-histories). This is particularly true for the model species *Arabidopsis thaliana*, for which it is

even possible to manipulate both resource acquisition and allocation (Schulze *et al*., 1994;

Bennett *et al*., 2012). This opens interesting avenues for experimentally testing the TRL

framework.

<u>Resource-based understanding of community dynamics</u>

A key aim of trait-based ecology is to use functional traits to explain the structure and dynamics

of higher levels of biological organisation (Lavorel & Garnier, 2002; McGill *et al.,* 2006;

Salguero-Gómez *et al*., 2018). This requires scaling-up through the levels of this hierarchy, in

which emergent properties at higher levels are shaped by the integration of, and covariances

between, multiple lower-level processes (Laughlin & Messier, 2015; Shipley *et al.,* 2016). By

combining the TRL framework, which describes the scaling-up from organs to individual-level

resource budgets, with descriptions of resource-based interactions among individuals (Mittelbach

& McGill, 2019), it should be possible to develop integrative, resource-driven models of

community dynamics.

In our model system, explicitly considering resource flows in analyses of demography and

community assembly has proven fruitful. Traits shaping competition for abiotic resources

determine interactions and performance at both the adult (Walter *et al*., 2023) and juvenile

(Cooksley *et al*., 2023) stage of *Protea* individuals. Furthermore, floral and seed resources at

plant-, local- and site-levels shape the interaction frequencies of *Protea* individuals with animal

mutualists and antagonists (Schmid *et al*., 2015; Nottebrock *et al*., 2017; Neu *et al*., 2023a,b) as

do the energetic requirements of the interacting animals (Neu *et al*., 2023a). In turn, the foraging

of animal mutualists and antagonists affects *Protea* fitness via pollination and seed predation,

respectively (Nottebrock *et al*., 2017; Walter *et al*., 2023).

An ambitious, but logical, next step would be to integrate these disparate analyses via a latent-

variable modelling framework, driven by the flow of resources from the abiotic environment

over competing plants to higher trophic levels. This could be done by integrating resource-driven

interactions into the TRL framework to scale up resource dynamics from organisms to entire

communities. Modelling such hierarchical dynamics via a universal currency of resources would

have clear advantages, particularly avoiding the intractability of quantifying pairwise interaction

coefficients among species in diverse communities (Broekman *et al*., 2019). Ultimately, such an

integrative resource-driven understanding of life-histories and community dynamics may feed

into a new generation of ecosystem models that represent biodiversity by general ecological

mechanisms rather than a predefined set of coarse functional strategies of organisms (Scheiter *et*

al. 2013; Harfoot *et al*., 2014; Higgins, 2017).

Conclusions

The proposed TRL framework enables data-driven inference on the processes linking traits,

resource budgets and life-histories of long-lived plants. Its application to a broad range of study

systems and its integration with models of community dynamics hold promise for a process-

based understanding of higher levels of biological organisation and a closer theory-founded

integration of functional ecology, evolutionary ecology, community ecology and ecosystem

science.

Acknowledgements

Fieldwork was conducted under Western Cape Nature Conservation Board permit 0028-

AAA008-00262 and the following landowners and reserves gave permission to work on their

properties: Ceres Mountain Fynbos Nature Reserve and Witzenberg Municipality, City of Cape

Town and Helderberg Nature Reserve, Fernkloof Nature Reserve and Overberg Municipality,

Flower Valley Conservation Trust, Grootbos Private Nature Reserve, Heuningklip Farm,

Hottentots Holland Nature Reserve, Limietberg Nature Reserve, Mont Rochelle Nature Reserve

and Riviersonderend Nature Reserve. Sarah de Gruchy, Madelein de Klerk, Nicolene Hellström,

Liezel Knight, Michael Leach, Marisa Noordergraaf, Hannah Oliphant, Barbara Seele, Adrian

D.A. Simmers, Megan Smith and Martina Treurnicht provided invaluable help with field and lab

work. We thank the Central Analytical Facility of the University of Stellenbosch for nutrient

determination, and Carsten Buchmann, Jörn Pagel and Hanna Walter for helpful discussions.

This research was funded by the German Research Foundation (DFG) project ProteaNet (grant

numbers SCHU 2259/3-3 and SCHL 1934/1-3).

Author contributions

Huw Cooksley, Matthias Schleuning and Frank Schurr designed the study and the conceptual

TRL framework. Huw Cooksley and Alex Neu collected data for the *Protea* case study, and Huw

Cooksley and Frank Schurr performed the analysis. Huw Cooksley and Frank Schurr led the

writing of the manuscript, and all authors contributed revisions and accepted the final version.

Conflict of interest statement

The authors declare no conflict of interest.

References

Adler, P. B., A. Fajardo, A. R. Kleinhesselink, and N. J. B. Kraft. 2013. Trait-based tests of

coexistence mechanisms. Ecology Letters 16:1294–1306.

https://doi.org/10.1111/ele.12157.

Babst, F., P. Bodesheim, N. Charney, A. D. Friend, M. P. Girardin, S. Klesse, D. J. P. Moore, K.

Seftigen, J. Björklund, O. Bouriaud, et al.. 2018. When tree rings go global: Challenges

and opportunities for retro- and prospective insight. Quaternary Science Reviews 197:1–

20. https://doi.org/10.1016/j.quascirev.2018.07.009.

Bennett, E., J. A. Roberts, and C. Wagstaff. 2012. Manipulating resource allocation in plants.

Journal of Experimental Botany 63:3391–3400. https://doi.org/10.1093/jxb/err442.

Bogdziewicz, M., D. Kelly, P. A. Thomas, J. G. A. Lageard, and A. Hacket-Pain. 2020. Climate

warming disrupts mast seeding and its fitness benefits in European beech. Nature Plants

6:88–94. https://doi.org/10.1038/s41477-020-0592-8.

Bond, W. 1980. Fire and Senescent Fynbos in the Swartberg, Southern Cape. South African

Forestry Journal 114:68–71. https://doi.org/10.1080/00382167.1980.9629837.

Bond, W. J. 1985. Canopy-stored seed reserves (serotiny) in Cape Proteaceae. South African

Journal of Botany 51:181–186. https://doi.org/10.1016/S0254-6299(16)31669-6.

Bond, W. J., J. Volk, and M. Viviers. 1984. Variation in Seedling Recruitment of Cape Proteaceae

after Fire. Journal of Ecology 72:209–221. https://doi.org/10.2307/2260014.

Born, J., H. P. Linder, and P. Desmet. 2007. The Greater Cape Floristic Region. Journal of

Biogeography 34:147–162. https://doi.org/10.1111/j.1365-2699.2006.01595.x.

Bradshaw, P. L., and R. M. Cowling. 2014. Landscapes, rock types, and climate of the Greater

Cape Floristic Region. Pages 26–46 *in* N. Allsopp, J. F. Colville, and G. A. Verboom,

editors. Fynbos: Ecology, Evolution, and Conservation of a Megadiverse Region. Oxford

University Press. https://doi.org/10.1093/acprof:oso/9780199679584.003.0002.

Broekman, M. J. E., H. C. Muller-Landau, M. D. Visser, E. Jongejans, S. J. Wright, and H. de

Kroon. 2019. Signs of stabilisation and stable coexistence. Ecology Letters 22:1957–

1975. https://doi.org/10.1111/ele.13349.

Brose, U., P. Archambault, A. D. Barnes, L.-F. Bersier, T. Boy, J. Canning-Clode, E. Conti, M.

Dias, C. Digel, A. Dissanayake, et al.. 2019. Predator traits determine food-web

architecture across ecosystems. Nature Ecology & Evolution 3:919–927.

https://doi.org/10.1038/s41559-019-0899-x.

869 Carlson, J. E., K. E. Holsinger, and R. Prunier. 2011. Plant Responses to Climate in the Cape

870   Floristic Region of South Africa: Evidence for Adaptive Differentiation in the Proteaceae.

871   Evolution 65:108–124. https://doi.org/10.1111/j.1558-5646.2010.01131.x

872 Chave, J., D. Coomes, S. Jansen, S. L. Lewis, N. G. Swenson, and A. E. Zanne. 2009. Towards a

873   worldwide wood economics spectrum. Ecology Letters 12:351–366.

874   https://doi.org/10.1111/j.1461-0248.2009.01285.x.

875 Clark, J. S. 2005. Why environmental scientists are becoming Bayesians. Ecology Letters 8:2–

876   14. https://doi.org/10.1111/j.1461-0248.2004.00702.x.

877 Clark, J. S., C. L. Nuñez, and B. Tomasek. 2019. Foodwebs based on unreliable foundations:

878   spatiotemporal masting merged with consumer movement, storage, and diet. Ecological

879   Monographs 89:e01381. https://doi.org/10.1002/ecm.1381.

880 Coetzee, J. H., and J. H. Giliomee. 1987. Seed predation and survival in the infructescences of

881   Protea repens (Proteaceae). South African Journal of Botany 53:61–64.

882   https://doi.org/10.1016/S0254-6299(16)31475-2.

883 Cooksley, H., L. Dreyling, K. J. Esler, S. Griebenow, G. Neumann, A. Valentine, M. Schleuning,

884   and F. M. Schurr. 2023, February 22. Functional traits shape plant-plant interactions and

885   recruitment in a hotspot of woody plant diversity. bioRxiv.

886   https://doi.org/10.1101/2023.02.22.529386.

887 Cramer, M. D., and J. J. Midgley. 2009. Maintenance costs of serotiny do not explain weak

888   serotiny. Austral Ecology 34:653–662. https://doi.org/10.1111/j.1442-9993.2009.01971.x.

889 Cramer, M., A. West, S. Power, R. Skelton, and W. D. Stock. 2014. Plant ecophysiological

890   diversity. Pages 248–272 *in* N. Allsopp, J. F. Colville, and G. A. Verboom, editors.

Fynbos: Ecology, Evolution and Conservation of a Megadiverse Region. Oxford

University Press. https://doi.org/10.1093/acprof:oso/9780199679584.003.0011

Dietz, H., and I. Ullmann. 1997. Age-determination of Dicotyledonous Herbaceous Perennials by

Means of Annual Rings: Exception or Rule? Annals of Botany 80:377–379.

https://doi.org/10.1006/anbo.1997.0423.

Dungan, R. J., M.-L. Navas, R. P. Duncan, and E. Garnier. 2008. Effects of leaf emergence on

leaf lifespan are independent of life form and successional status. Austral Ecology

33:932–939. https://doi.org/10.1111/j.1442-9993.2008.01875.x.

Enquist, B. J., G. B. West, E. L. Charnov, and J. H. Brown. 1999. Allometric scaling of

production and life-history variation in vascular plants. Nature 401:907–911.

https://doi.org/10.1038/44819.

Enright, N. J., J. B. Fontaine, D. M. Bowman, R. A. Bradstock, and R. J. Williams. 2015. Interval

squeeze: altered fire regimes and demographic responses interact to threaten woody

species persistence as climate changes. Frontiers in Ecology and the Environment

13:265–272. https://doi.org/10.1890/140231.

Evans, M. E. K., C. Merow, S. Record, S. M. McMahon, and B. J. Enquist. 2016. Towards

Process-based Range Modeling of Many Species. Trends in Ecology & Evolution

31:860–871.

Gibert, A., E. F. Gray, M. Westoby, I. J. Wright, and D. S. Falster. 2016. On the link between

functional traits and growth rate: meta-analysis shows effects change with plant size, as

predicted. Journal of Ecology 104:1488–1503. https://doi.org/10.1111/1365-2745.12594.

Gray, E. F., I. J. Wright, D. S. Falster, A. S. D. Eller, C. E. R. Lehmann, M. G. Bradford, and L.

A. Cernusak. 2019. Leaf:wood allometry and functional traits together explain substantial

growth rate variation in rainforest trees. AoB PLANTS 11:plz024.

https://doi.org/10.1093/aobpla/plz024.

Harfoot, M. B. J., T. Newbold, D. P. Tittensor, S. Emmott, J. Hutton, V. Lyutsarev, M. J. Smith, J.

P. W. Scharlemann, and D. W. Purves. 2014. Emergent Global Patterns of Ecosystem

Structure and Function from a Mechanistic General Ecosystem Model. PLOS Biology

12:e1001841. https://doi.org/10.1371/journal.pbio.1001841.

Harris, M. S., and J. R. Pannell. 2010. Canopy seed storage is associated with sexual dimorphism

in the woody dioecious genus Leucadendron. Journal of Ecology 98:509–515.

https://doi.org/10.1111/j.1365-2745.2009.01623.x.

Higgins, S. I. 2017. Ecosystem Assembly: A Mission for Terrestrial Earth System Science.

Ecosystems 20:69–77. https://doi.org/10.1007/s10021-016-0054-3.

Iida, Y., T. S. Kohyama, N. G. Swenson, S.-H. Su, C.-T. Chen, J.-M. Chiang, and I.-F. Sun. 2014.

Linking functional traits and demographic rates in a subtropical tree community: the

importance of size dependency. Journal of Ecology 102:641–650.

https://doi.org/10.1111/1365-2745.12221.

Kattge, J., G. Bönisch, S. Díaz, S. Lavorel, I. C. Prentice, P. Leadley, S. Tautenhahn, G. D. A.

Werner, T. Aakala, M. Abedi, et al.. 2020. TRY plant trait database – enhanced coverage

and open access. Global Change Biology 26:119–188. https://doi.org/10.1111/gcb.14904.

Kleyer, M., and V. Minden. 2015. Why functional ecology should consider all plant organs: An

allocation-based perspective. Basic and Applied Ecology 16:1–9.

https://doi.org/10.1016/j.baae.2014.11.002.

Lambers, H., P. M. Finnegan, R. Jost, W. C. Plaxton, M. W. Shane, and M. Stitt. 2015.

Phosphorus nutrition in Proteaceae and beyond. Nature Plants 1:1–9.

https://doi.org/10.1038/nplants.2015.109.

Lamonica, D., J. Pagel, and F. M. Schurr. 2021. Predicting the dynamics of establishing tree

populations: A framework for statistical inference and lessons for data collection.

Methods in Ecology and Evolution 12:1721–1733. https://doi.org/10.1111/2041-

210X.13656.

Lamont, B. 1985. Fire Responses of Sclerophyll Shrublands: A Population Ecology Approach,

with Particular Reference to the Genus Banksia. Pages 41–46 *in* Fire ecology and

management of Western Australian ecosystems. Curtin University, Perth.

http://scienceon.kisti.re.kr/srch/selectPORSrchArticle.do?cn=NPAP01749595.

Lamont, B. B., and P. K. Groom. 1998. Seed and Seedling Biology of the Woody-fruited

Proteaceae. Australian Journal of Botany 46:387–406. https://doi.org/10.1071/bt96135.

Lamont, B. B., T. He, and Z. Yan. 2019. Evolutionary history of fire-stimulated resprouting,

flowering, seed release and germination. Biological Reviews 94:903–928.

https://doi.org/10.1111/brv.12483.

Lamont, B. B., D. C. Le Maitre, R. M. Cowling, and N. J. Enright. 1991. Canopy Seed Storage in

Woody Plants. Botanical Review 57:277–317. https://www.jstor.org/stable/4354176.

Laughlin, D. C., J. R. Gremer, P. B. Adler, R. M. Mitchell, and M. M. Moore. 2020. The net

effect of functional traits on fitness. Trends in Ecology & Evolution 35:1037–1047.

https://doi.org/10.1016/j.tree.2020.07.010

Laughlin, D. C., and J. Messier. 2015. Fitness of multidimensional phenotypes in dynamic

adaptive landscapes. Trends in Ecology & Evolution 30:487–496.

https://doi.org/10.1016/j.tree.2015.06.003.

Lavorel, S., and E. Garnier. 2002. Predicting changes in community composition and ecosystem

functioning from plant traits: revisiting the Holy Grail. Functional Ecology 16:545–556.

https://doi.org/10.1046/j.1365-2435.2002.00664.x.

Levins, R. 1968. Evolution in Changing Environments: Some Theoretical Explorations. (MPB-

2). Princeton University Press. https://doi.org/10.2307/j.ctvx5wbbh.

Low, A. B., and B. B. Lamont. 1986. Nutrient allocation in winter rainfall Proteaceous

heathlands in relation to nutrient losses through wildflower picking and fire. Acta

Horticulturae:89–100. https://doi.org/10.17660/ActaHortic.1986.185.9.

McGill, B. J., B. J. Enquist, E. Weiher, and M. Westoby. 2006. Rebuilding community ecology

from functional traits. Trends in Ecology & Evolution 21:178–185.

https://doi.org/10.1016/j.tree.2006.02.002.

Meredith, M. 2021. saveJAGS: Run JAGS and Regularly Save Output to Files. R package

version 0.0.4.9002. https://github.com/mikemeredith/saveJAGS

Messier, J., M. J. Lechowicz, B. J. McGill, C. Violle, and B. J. Enquist. 2017. Interspecific

integration of trait dimensions at local scales: the plant phenotype as an integrated

network. Journal of Ecology 105:1775–1790. https://doi.org/10.1111/1365-2745.12755.

Midgley, J. J., and A. G. Rebelo. 2008. Life-history evolution as an explanation for the absence

of the tree life-form in Cape fynbos. South African Journal of Science 104:89–90.

http://www.scielo.org.za/scielo.php?script=sci_abstract&pid=S0038-

23532008000200005&lng=en&nrm=iso&tlng=en.

Mittelbach, G. G., and B. J. McGill. 2019. Community Ecology. Second Edition, Second Edition.

Oxford University Press, Oxford, New York.

https://global.oup.com/academic/product/community-ecology-9780198835851?

cc=de&lang=en&

Moles, A. T., and M. Westoby. 2006. Seed size and plant strategy across the whole life cycle.

Oikos 113:91–105. https://doi.org/10.1111/j.0030-1299.2006.14194.x.

Myers, N., R. A. Mittermeier, C. G. Mittermeier, G. A. B. da Fonseca, and J. Kent. 2000.

Biodiversity hotspots for conservation priorities. Nature 403:853–858.

https://doi.org/10.1038/35002501.

Neu, A., H. Cooksley, K. J. Esler, A. Pauw, F. Roets, F. M. Schurr, and M. Schleuning. 2023a.

Interactions between protea plants and their animal mutualists and antagonists are

structured more by energetic than morphological trait matching. Functional Ecology

37:176–189. https://doi.org/10.1111/1365-2435.14231.

Neu, A., H. Cooksley, K. J. Esler, A. Pauw, F. Roets, F. M. Schurr, and M. Schleuning. 2023b.

Bird and insect pollinators differ in specialization and potential pollination services along

disturbance and resource gradients. Austral Ecology 00.

https://doi.org/10.1111/aec.13324.

Niklas, K. J. 1992. Plant Biomechanics: An Engineering Approach to Plant Form and Function.

University of Chicago Press, Chicago, IL.

https://press.uchicago.edu/ucp/books/book/chicago/P/bo3631132.html.

Niklas, K. J. 1994. Plant Allometry: The Scaling of Form and Process. University of Chicago

Press, Chicago, IL.

https://press.uchicago.edu/ucp/books/book/chicago/P/bo3629790.html.

Niklas, K. J. 2004. Plant allometry: is there a grand unifying theory? Biological Reviews

79:871–889. https://doi.org/10.1017/S1464793104006499.

van Noordwijk, A. J., and G. de Jong. 1986. Acquisition and Allocation of Resources: Their

Influence on Variation in Life History Tactics. The American Naturalist 128:137–142.

https://www.jstor.org/stable/2461293.

Nottebrock, H., B. Schmid, K. Mayer, C. Devaux, K. J. Esler, K. Böhning-Gaese, M. Schleuning,

J. Pagel, and F. M. Schurr. 2017. Sugar landscapes and pollinator-mediated interactions in

plant communities. Ecography 40:1129–1138. https://doi.org/10.1111/ecog.02441.

Ohse, B., A. Compagnoni, C. E. Farrior, S. M. McMahon, R. Salguero-Gómez, N. Rüger, and T.

M. Knight. 2023. Demographic synthesis for global tree species conservation. Trends in

Ecology & Evolution 0. https://doi.org/10.1016/j.tree.2023.01.013

Pérez-Harguindeguy, N., S. Díaz, E. Garnier, S. Lavorel, H. Poorter, P. Jaureguiberry, M. S. Bret-

Harte, W. K. Cornwell, J. M. Craine, D. E. Gurvich, et al.. 2016. Corrigendum to: New

handbook for standardised measurement of plant functional traits worldwide. Australian

Journal of Botany 64:715–716. https://doi.org/10.1071/bt12225_co.

Petit, R. J., and A. Hampe. 2006. Some Evolutionary Consequences of Being a Tree. Annual

Review of Ecology, Evolution, and Systematics 37:187–214.

https://doi.org/10.1146/annurev.ecolsys.37.091305.110215

Pieruschka, R., and U. Schurr. 2019. Plant Phenotyping: Past, Present, and Future. Plant

Phenomics 2019. https://doi.org/10.34133/2019/7507131.

Pistón, N., F. de Bello, A. T. C. Dias, L. Götzenberger, B. H. P. Rosado, E. A. de Mattos, R.

Salguero-Gómez, and C. P. Carmona. 2019. Multidimensional ecological analyses

demonstrate how interactions between functional traits shape fitness and life history

strategies. Journal of Ecology 107:2317–2328. https://doi.org/10.1111/1365-2745.13190.

Plummer, M. 2003. JAGS: A Program for Analysis of Bayesian Graphical Models Using Gibbs

Sampling. Proceedings of the 3rd International Workshop on Distributed Statistical

Computing 124:1–10.

Poorter, H. 1994. Construction costs and payback time of biomass: a whole plant perspective.

Pages 111–127 *in* J. Roy and E. Garnier, editors. A whole plant perspective on carbon-

nitrogen interactions. The Hague, Netherlands: SPB Academic Publishing.

http://science.poorter.eu/1994_Poorter_C&Nrelations.pdf

R Core Team. 2020. R: A Language and Environment for Statistical Computing. Vienna, Austria:

R Foundation for Statistical Computing. https://www.R-project.org/

Rebelo, T. 2001. Sasol Proteas: A Field Guide to the Proteas of Southern Africa. Fernwood Press,

Vlaeberg, South Africa.

Redmond, M. D., P. J. Weisberg, N. S. Cobb, C. A. Gehring, A. V. Whipple, and T. G. Whitham.

2016. A robust method to determine historical annual cone production among slow-

growing conifers. Forest Ecology and Management 368:1–6.

https://doi.org/10.1016/j.foreco.2016.02.028.

Reich, P. B. 2014. The world-wide 'fast–slow' plant economics spectrum: a traits manifesto.

Journal of Ecology 102:275–301. https://doi.org/10.1111/1365-2745.12211.

Reich, P. B., M. B. Walters, and D. S. Ellsworth. 1997. From tropics to tundra: Global

convergence in plant functioning. Proceedings of the National Academy of Sciences

94:13730–13734. https://doi.org/10.1073/pnas.94.25.13730.

Richter, J. P. 1970. The Notebooks of Leonardo Da Vinci. Courier Corporation, Chelmsford,

Massachusetts.

Salguero-Gómez, R., O. R. Jones, E. Jongejans, S. P. Blomberg, D. J. Hodgson, C. Mbeau-Ache,

P. A. Zuidema, H. de Kroon, and Y. M. Buckley. 2016. Fast–slow continuum and

reproductive strategies structure plant life-history variation worldwide. Proceedings of

the National Academy of Sciences 113:230–235.

https://doi.org/10.1073/pnas.1506215112.

Salguero-Gómez, R., C. Violle, O. Gimenez, and D. Childs. 2018. Delivering the promises of

trait-based approaches to the needs of demographic approaches, and vice versa.

Functional Ecology 32:1424–1435. https://doi.org/10.1111/1365-2435.13148.

Scheiter, S., L. Langan, and S. I. Higgins. 2013. Next-generation dynamic global vegetation

models: learning from community ecology. New Phytologist 198:957–969.

https://doi.org/10.1111/nph.12210.

Schmid, B., H. Nottebrock, K. J. Esler, J. Pagel, A. Pauw, K. Böhning-Gaese, F. M. Schurr, and

M. Schleuning. 2015. Reward quality predicts effects of bird-pollinators on the

reproduction of African Protea shrubs. Perspectives in Plant Ecology, Evolution and

Systematics 17:209–217. https://doi.org/10.1016/j.ppees.2015.02.007

Schneider, R., R. Calama, and O. Martin-Ducup. 2020. Understanding Tree-to-Tree Variations in

Stone Pine (Pinus pinea L.) Cone Production Using Terrestrial Laser Scanner. Remote

Sensing 12:173. https://doi.org/10.3390/rs12010173.

Schulze, W., E.-D. Schulze, J. Stadler, H. Heilmeier, M. Stitt, and H. a. Mooney. 1994. Growth

and reproduction of Arabidopsis thaliana in relation to storage of starch and nitrate in the

wild-type and in starch-deficient and nitrate-uptake-deficient mutants. Plant, Cell &

Environment 17:795–809. https://doi.org/10.1111/j.1365-3040.1994.tb00174.x.

Schurr, F. M., K. J. Esler, J. A. Slingsby, and N. Allsopp. 2012. Fynbos Proteaceae as model

organisms for biodiversity research and conservation. South African Journal of Science

108:1–4. https://doi.org/10.10520/EJC127526.

Schurr, F. M., G. F. Midgley, A. G. Rebelo, G. Reeves, P. Poschlod, and S. I. Higgins. 2007.

Colonization and persistence ability explain the extent to which plant species fill their

potential range. Global Ecology and Biogeography 16:449–459.

https://doi.org/10.1111/j.1466-8238.2006.00293.x.

Shane, M. W., and H. Lambers. 2005. Cluster Roots: A Curiosity in Context. Plant and Soil

274:101–125. https://doi.org/10.1007/s11104-004-2725-7.

Shinozaki, K., K. Yoda, K. Hozumi, and T. Kira. 1964. A quantitative analysis of plant form; the

pipe model theory,1. Japanese Journal of Ecology 14:97–105.

https://doi.org/10.18960/seitai.14.3_97.

Shipley, B., F. De Bello, J. H. C. Cornelissen, E. Laliberté, D. C. Laughlin, and P. B. Reich.

2016. Reinforcing loose foundation stones in trait-based plant ecology. Oecologia

180:923–931. https://doi.org/10.1007/s00442-016-3549-x.

Shipley, B., M. J. Lechowicz, I. Wright, and P. B. Reich. 2006. Fundamental Trade-Offs

Generating the Worldwide Leaf Economics Spectrum. Ecology 87:535–541.

https://doi.org/10.1890/05-1051.

Stearns, S. C. 1992. The Evolution of Life Histories. Oxford University Press, Oxford, New

York. https://global.oup.com/academic/product/the-evolution-of-life-histories-

9780198577416?cc=de&lang=en&

Swenson, N. G., S. J. Worthy, D. Eubanks, Y. Iida, L. Monks, K. Petprakob, V. E. Rubio, K.

Staiger, and J. Zambrano. 2020. A Reframing of Trait–Demographic Rate Analyses for

Ecology and Evolutionary Biology. International Journal of Plant Sciences 181:33–43.

https://doi.org/10.1086/706189.

Tonnabel, J., T. J. M. Van Dooren, J. Midgley, P. Haccou, A. Mignot, O. Ronce, and I. Olivieri.

2012. Optimal resource allocation in a serotinous non-resprouting plant species under

different fire regimes. Journal of Ecology 100:1464–1474. https://doi.org/10.1111/j.1365-

2745.2012.02023.x.

Treurnicht, M., J. Pagel, K. J. Esler, A. Schutte-Vlok, H. Nottebrock, T. Kraaij, A. G. Rebelo, and

F. M. Schurr. 2016. Environmental drivers of demographic variation across the global

geographical range of 26 plant species. Journal of Ecology 104:331–342.

https://doi.org/10.1111/1365-2745.12508.

Treurnicht, M., J. Pagel, J. Tonnabel, K. J. Esler, J. A. Slingsby, and F. M. Schurr. 2020.

Functional traits explain the Hutchinsonian niches of plant species. Global Ecology and

Biogeography 29:534–545. https://doi.org/10.1111/geb.13048.

Vasseur, F., J. Bresson, G. Wang, R. Schwab, and D. Weigel. 2018. Image-based methods for

phenotyping growth dynamics and fitness components in Arabidopsis thaliana. Plant

Methods 14:63. https://doi.org/10.1186/s13007-018-0331-6.

Violle, C., M.-L. Navas, D. Vile, E. Kazakou, C. Fortunel, I. Hummel, and E. Garnier. 2007. Let

the concept of trait be functional! Oikos 116:882–892. https://doi.org/10.1111/j.0030-

1299.2007.15559.x.

Violle, C., P. B. Reich, S. W. Pacala, B. J. Enquist, and J. Kattge. 2014. The emergence and

promise of functional biogeography. Proceedings of the National Academy of Sciences

111:13690–13696. https://doi.org/10.1073/pnas.1415442111.

Walter, H. E., J. Pagel, H. Cooksley, A. Neu, M. Schleuning, and F. M. Schurr. 2023. Effects of

biotic interactions on plant fecundity depend on spatial and functional structure of

communities and time since disturbance. Journal of Ecology 111:110–124.

https://doi.org/10.1111/1365-2745.14018.

Warton, D. I., I. J. Wright, D. S. Falster, and M. Westoby. 2006. Bivariate line-fitting methods for

allometry. Biological Reviews 81:259–291. https://doi.org/10.1017/S1464793106007007.

West, A. G., T. E. Dawson, E. C. February, G. F. Midgley, W. J. Bond, and T. L. Aston. 2012.

Diverse functional responses to drought in a Mediterranean-type shrubland in South

Africa. New Phytologist 195:396–407. https://doi.org/10.1111/j.1469-8137.2012.04170.x.

Westoby, M. 1998. A leaf-height-seed (LHS) plant ecology strategy scheme. Plant and Soil

199:213–227. https://doi.org/10.1023/A:1004327224729.

Westoby, M., D. S. Falster, A. T. Moles, P. A. Vesk, and I. J. Wright. 2002. Plant Ecological

Strategies: Some Leading Dimensions of Variation Between Species. Annual Review of

Ecology and Systematics 33:125–159.

https://doi.org/10.1146/annurev.ecolsys.33.010802.150452.

Williams, K., F. Percival, J. Merino, and H. A. Mooney. 1987. Estimation of Tissue Construction

Cost from Heat of Combustion and Organic Nitrogen Content. Plant, Cell & Environment

10:725–734. https://doi.org/10.1111/1365-3040.ep11604754.

Wilson, A. M., A. M. Latimer, and J. A. Silander. 2015. Climatic controls on ecosystem

resilience: Postfire regeneration in the Cape Floristic Region of South Africa.

Proceedings of the National Academy of Sciences 112:9058–9063.

https://doi.org/10.1073/pnas.1416710112.

Wright, I. J., P. B. Reich, M. Westoby, D. D. Ackerly, Z. Baruch, F. Bongers, J. Cavender-Bares,

T. Chapin, J. H. C. Cornelissen, M. Diemer, et al.. 2004. The worldwide leaf economics

spectrum. Nature 428:821–827. https://doi.org/10.1038/nature02403.

Yang, J., M. Cao, and N. G. Swenson. 2018. Why Functional Traits Do Not Predict Tree

Demographic Rates. Trends in Ecology & Evolution 33:326–336.

https://doi.org/10.1016/j.tree.2018.03.003.

Table 1: Overview of formulae for costs and benefits of resource allocation to life-history

components.

Cost / Benefit	Link	Trait covariates
Leaf cost	ln – ln	SLA, Leaf N concentration
Cone cost	ln – ln	Cone mass
Seed cost	ln – ln	Seed N content
Serotiny cost	ln – ln	Cone mass
Leaf benefit	ln – ln	SLA, Leaf N concentration
Cone defence benefit	logit – ln	Cone mass, Seed N content
Seed benefit	ln – ln	Seed N content
Serotiny benefit	logit – ln	Cone mass

All formulae include normally-distributed, zero-mean random intercepts for species, site and

individual.

Figure 1: A trait-resource-life-history (TRL) framework for quantifying how functional traits

shape resource budgets and life-histories of plants. Functional traits (orange ovals) shape organ-

level costs and benefits of resource investment into different life-history components (green

circles). The framework requires quantifying how these functional traits scale up to whole-plant

resource flows and how the resulting resource budget (purple) causes time-dependent changes of

life-history components at the whole-plant level. While direct measurements of resource budgets

are difficult, a latent-state-space representation of the TRL framework can be fitted to data on

functional traits and life-history components (ovals and circles, respectively) and can model

resource budgets as hidden aggregate variables.

1156

1157   Figure 2: A stylised depiction of a *Protea* individual, showing proxies for life-history

1158   components measured in our study: stem length increments, and number and age of closed, open

1159   and abscised cones. The insert shows a stylised cross-section through a closed cone, showing

1160   fertile seeds and the proportion of the cone area consumed by insect seed predators.

1161

Figure 3: Reconstructing the vegetative growth of *Protea* individuals. (a) *Protea* plants follow da

Vinci's rule so that their wood volume can be approximated as the volume of a cylinder. (b)

Since species-specific allometries (line of fit shown for *P. repens*) relate basal area to stem

length, measurements of past stem length predict past wood volume and – in combination with

wood density - aboveground wood biomass in the past. (c) Species-specific proportionalities

(line of fit shown for *P. repens*) between leaf mass produced on a branch and branch cross-

sectional area permit the prediction of whole-plant leaf production. By combining this with (d)

species-specific leaf retention functions (line of fit shown for *P. repens*), one obtains proxies of

whole-plant leaf mass in the past. For (a) – (c), species are shown as symbols of differing shape

and colour.

1172

Figure 4: Process diagram of the TRL model for *Protea*. Data are represented by rectangles, and

unobserved nodes are represented by ovals. Costs and benefits of different organs are informed

by traits. In turn, organ costs, allocations and the resource pool determine the production of

cones, seeds, wood and leaves, and maternal care. These latent performance measures are then

linked to the observed performance data. Seed benefit determines the amount of resources

available to a seedling in the first time-step, and for subsequent time-steps leaf benefit and total

leaf mass determine resource acquisition and the size of the resource pool in the next time-step.

Serotiny benefit determines the proportion of cones surviving to the next time-step.

1181

Figure 5: Observed annual performance measures for 600 *Protea* individuals against predictions

of the TRL model for (a) annual stem length increment, (b) annual cone production, (c) seed

number per sampled cone at time of observation, (d) number of closed cones maintained (age-

stratified) at time of observation and (e) cone defence percent at time of observation (points are

jittered slightly for clarity). Predictions are posterior medians of one-step-ahead predictions

conditioned upon the resource state in the previous time-step. The square of the Pearson's

correlation between observed and predicted values is shown in the top-left corner of each plot,

and the dashed gold line represents the line of identity.

1190

Figure 6: TRL model estimates of the age-dependent dynamics of whole-plant resource pools,

resource allocation and total reproductive output (canopy seed bank size) for the 20 studied

individuals of three *Protea* species on study site 'jona_4'. (a) Posterior medians of resource pools

versus age. Insets show resource allocation versus age for one individual per species. (b)

Posterior medians of total reproductive output (size of the canopy seed bank) versus age for the

same individuals.

1197

Figure 7: The effects of functional traits on paired organ-level costs and benefits of life-history

components. Panels show posterior medians and 95% credible intervals of the effect of one

standard deviation increase in ln trait values on the link-scale costs and benefits of (a) leaves, (b)

cones, (c) seeds and (d) serotiny. Individual-level costs and benefits are estimated by the TRL

framework applied to 22 species of *Protea*, inferred in an inverse-modelling approach from

individual-level data on age-stratified performance measures and population-level trait values

(except for leaf N concentration measured at species-level). For definitions of costs and benefits

see 'A Trait-Resource-Life-history (TRL) model for *Protea*'.

1206

Figure 8: Age-specific relationships between organ-level costs and benefits, whole-plant resource

budgets and whole-plant performance. Panels show Pearson's correlation coefficients between

(a) organ-level benefits and whole-plant resource pools, (b) organ-level costs and whole-plant

resource pools, (c) whole-plant resource pools and allocations and (d) whole-plant resource

investments (resource pool x allocations) and performance measures. Correlations were

calculated from posterior median estimates for all 600 study individuals, using individual-level

estimates of costs and benefits, and individual- and age-specific estimates of resource pools,

allocations and performance measures.

1215

Figure 9: Age-dependent relationships between functional traits and whole-plant performance.

Panels show Pearson's correlation coefficients between whole-plant performance measures

(including total reproductive output, an individual' s seed bank) and (a) SLA and (b) seed N

content. Correlations were calculated from posterior median estimates for all 600 study

individuals, using individual-level estimates of whole-plant performance measures. Relationships

between performance and the remaining functional traits are shown in Appendix S1: Figure S8).

1222

Figure 10: The relationship of (a) SLA and (b) seed N content with age of maturity for 600

individuals of 22 *Protea* species. The depicted age of maturity is the posterior median age at

which an individual is predicted to have at least one seed in its canopy seed bank. Black lines

show the fitted linear relationships between these variables and grey shadings show the 95%

confidence intervals of these relationships. Pearson's correlation coefficients between variables

are shown in the top-left of each panel.

Figure 1

Figure 2

1232

1233   Figure 3

1234

1235   Figure 4

1237   Figure 5

1238

1239   Figure 6

1240

1241   Figure 7

1242

1247   Figure 10

1248