1

Brief Report:

Tricontinental detection of Streptococcus pyogenes M1$_{UK}$: A call for wider research and active surveillance.

Ana Vieira[1,2], Valerie WC Soo[1], Ho Kwong Li[1,2], Xiangyun Zhi[1,2], Lucy Reeves[1], Kristin K. Huse[1,2],

Kai Yi Mok[1], Oscar Cowen[1], Elita Jauneikaite[3,4], Juliana Coelho[3,5], Shiranee Sriskandan[1,2,3]

[1]Department of Infectious Disease, Faculty of Medicine, Imperial College London, London, UK.

[2]Centre for Bacterial Resistance Biology, Imperial College London, London, UK.

[3]NIHR Health Protection Research Unit in Healthcare-associated Infections and AMR, Imperial

College London, London, UK.

[4]School of Public Health, Faculty of Medicine, Imperial College London, London, UK.

[5]Centre for Infections, UK Health Security Agency, London, UK,

Correspondence: Shiranee Sriskandan s.sriskandan@imperial.ac.uk

Word count 1441

Summary

The *Streptococcus pyogenes* M1$_{UK}$ lineage, characterised by an intrinsic ability to express SpeA

toxin and defined by 27 single nucleotide polymorphisms in the core genome, dominates the

population of *emm*1 *S. pyogenes* isolates in England. The lineage has been identified elsewhere in

Europe, North America, and, most recently, Australia. SpeA however may not be the sole

deterministic factor underlying success of the M1$_{UK}$ lineage. Production of SpeA by strains belonging

to an intermediate *emm*1 sublineage, M1$_{23SNP}$, is indistinguishable from M1$_{UK}$ strains. Despite this,

in England at least, M1$_{UK}$ has outcompeted strains from the M1$_{23SNP}$ sublineage. We infer that the

fitness of M1$_{UK}$ resides in additional properties that confer an advantage to *S. pyogenes*, underlining

a need for further research. A single nucleotide polymorphism (SNP) in the ssrA leader sequence

upstream of *speA* is one of a limited number of SNPs that distinguish intermediate sublineages that

differ in SpeA production. Introduction of the ssrA SNP into representative isolates of the widely

disseminated M1$_{global}$ clone and the intermediate M1$_{13SNP}$ lineage (that cannot otherwise produce

readily-detectable SpeA in culture) resulted in SpeA expression, confirming the importance of the

ssrA SNP to SpeA phenotype. However, RNAseq analysis of clinical strains showed that presence

of the SNP was not invariably linked to read-through from the ssrA leader sequence or SpeA

expression. Literature review suggests that read through and *speA* mRNA transcript length may be

impacted by the two component regulator CovRS, pointing to a complex regulatory network

interaction between the bacterial chromosome and phage-encoded superantigens.

Introduction

A marked increase in serious *Streptococcus pyogenes* infections has been reported in several

countries following relaxation of public health interventions designed to limit the spread of COVID-

19 (1). In England, notifications of both scarlet fever and invasive *S. pyogenes* infections increased

throughout 2022; infections peaked at the end of the calendar year but remained high thereafter (2).

In the 30 weeks from mid September 2022-mid April 2023 there were 54,394 notifications of scarlet

fever and 2965 invasive *S. pyogenes* infections in England alone (2). The overall case fatality rate

of 14.2% (2) underlines the major public health impact of the observed increase, and the importance

of understanding the relative roles of strain pathogenicity, population immune susceptibility, co-infections, season, and access to healthcare.

*Emm*1 *S. pyogenes* strains are known to be inherently invasive and normally account for one quarter of invasive infections; currently however, *emm*1 isolates account for over half of invasive infections, rising to over 60% in children in England (2). The population of *emm*1 *S. pyogenes* isolates in England is dominated by a novel sublineage, $M1_{UK}$, that has an intrinsic ability to express the phage-encoded superantigenic toxin, streptococcal pyrogenic exotoxin (Spe) A (3, 4). $M1_{UK}$ strains can be differentiated from strains belonging to the highly successful *emm*1 *S. pyogenes* globally-disseminated clone that emerged in the 1980's ($M1_{global}$) by just 27 single nucleotide polymorphisms in the core genome (3). The $M1_{UK}$ lineage has also been identified elsewhere in Europe (5), North America (6, 7), and most recently in Australia, in isolates from Queensland and Victoria dating from 2013 (8), just 3 years after its first detection in England (3). While $M1_{UK}$ represented ~60% of Australian *emm*1 isolates by 2020, in England, this proportion reached 91% by 2020 (4,8). The detection of $M1_{UK}$ in Australia, including Queensland, underlines the ability of this lineage to spread in temperate and more tropical climates. Of added concern are reports of strains infected with a ΦHKU488.vir-like phage, that carries both *speC* and *ssa* superantigen genes in Australia (8). To date, this phage has not been identified in *emm*1 strains from England, although ~10% of *emm*1 strains in England do possess a phage encoding *speC* and *spd*.

Previous genomic analysis of *emm*1 *S. pyogenes* strains in England identified two intermediate sublineages in addition to $M1_{global}$ and $M1_{UK}$, with just 13 and 23 of the 27 SNPs that define $M1_{UK}$ (3). We recently reported that strains from the $M1_{13SNP}$ intermediate sublineage produce negligible amounts of SpeA, similar to $M1_{global}$ (9). Phylogenetic analysis of Australian sequences (9) shows the $M1_{13SNP}$ sublineage to be present in Australia too (Figure 1A). Strains from the $M1_{23SNP}$ sublineage, in contrast, produce SpeA at an increased level that is indistinguishable from $M1_{UK}$ isolates (9). Despite increased SpeA production, by 2020, we were unable to detect $M1_{23SNP}$ isolates when screening all *emm*1 isolates submitted to the reference laboratory in England (4). The $M1_{23SNP}$

intermediate sublineage appears to have been outcompeted by M1$_{UK}$, suggesting an added fitness

advantage in M1$_{UK}$ strains beyond production of SpeA alone. We postulate this further adaptation

may be conferred by the additional 4 SNPs that distinguish M1$_{UK}$ from M1$_{23SNP}$ isolates (3 non-

synonymous SNPs and an intergenic SNP 39bp upstream of *glpF2*, aquaporin, expression of which

is significantly reduced in M1$_{UK}$) (8, 9). In this report we seek to further understand the mechanisms

that differentiate SpeA-producers from non-producers among *emm*1 *S. pyogenes* strains that carry

the phage that encodes *speA*.

Results.

The M1$_{UK}$ SNP at position 983438 of the MGAS5005 reference genome, within the leader sequence

of *ssrA* and upstream of the *speA* start site (referred to hereafter as the 'ssrA SNP'), is present in

both M1$_{23SNP}$ and M1$_{UK}$ sublineages, but is absent in M1$_{global}$ and M1$_{13SNP}$ sublineages. Through

screening of natural mutants arising during outbreaks, the ssrA SNP was determined to be

specifically associated with increased SpeA production (9). We introduced the ssrA SNP into an

M1$_{global}$ strain, and into an intermediate M1$_{13SNP}$ strain, to determine if this would be sufficient to

enhance SpeA production. Consistent with the findings of Davies et al (8), SpeA production in culture

supernatants increased in both transformants (Figure 1B). Enhancement of SpeA was similar in both

the M1$_{global}$ and M1$_{13SNP}$ backgrounds, suggesting that the accumulated SNPs in the M1$_{13SNP}$ lineage,

which include almost half of the SNPs that characterize M1$_{UK}$, do not appreciably contribute to the

SpeA phenotype. We considered if our mutagenesis strategy itself might lead to increased SpeA

expression but this was ruled out by use of a control construct in each strain (Figure 1B).

To understand the basis for difference in *speA* expression between the lineages, we compared

RNAseq reads from four randomly selected M1$_{global}$ strains, which do not have the ssrA SNP, and

four randomly selected M1$_{UK}$ strains, which all have this SNP. None of the strains possessed

regulatory gene mutations other than those that characterize M1$_{UK}$. Similar to Davies et al (8), we

found that there was indeed evidence of read-through in the intergenic regions between the end of

ssrA and the start of *spea*. (Figure 1C). However, we did not see clear differentiation between the

four M1$_{global}$ strains and the four M1$_{UK}$ strains within the intergenic region, where read abundance in

two of the M1$_{global}$ strains was almost identical to M1$_{UK}$ strains (Figure 1C). Regardless, transcription

of *speA* was significantly reduced in all four M1$_{global}$ strains compared with M1$_{UK}$. Interestingly, *speA*

read abundance was somewhat higher in the two M1$_{global}$ strains that had more intergenic reads, but

not as high as the M1$_{UK}$ strains. These data provide support for intergenic read-through affecting

speA transcription. However, based on short read RNA sequencing data, we cannot definitively

ascribe read abundance in the intergenic region to the ssrA SNP in all cases, noting that all four of

the M1$_{global}$ strains studied lack any SNP in this region. Comparative WGS analysis between the four

M1$_{global}$ strains failed to detect genetic features associated with higher levels of read-through. Taken

together, transcriptional read-through was observed in this region, but the association with the ssrA

SNP and SpeA over-expression was not absolute. Despite not possessing the ssrA SNP, two of four

M1$_{global}$ strains tested showed similar read-through to M1$_{UK}$ strains, yet made less SpeA, pointing to

additional factors that influence SpeA release. The ssrA SNP however does indeed appear

important, at least in the strains examined.

Comment

A polycistronic transcript for speA was originally reported ~25 years ago by Cleary *et al*, who

identified both ~2kB and ~900bp *speA* mRNA transcripts in a single *emm*1 *S. pyogenes* strain (10).

In that case, the longer 2kB transcript was however associated with colonies that failed to produce

SpeA, while the shorter 900bp transcript, corresponding to the expected size for the *speA* transcript,

was associated with mucoid *emm*1 colonies that produced abundant SpeA. Based on the date of

publication, we infer the *emm*1 strain in that study (10) to be from an M1$_{global}$ background. The within-

strain variation in *speA* transcript length coupled with mucoid phenotype points to involvement of the

two-component regulator covRS (csrRS) which is now known to repress capsule and *speA*

transcription in *S. pyogenes* (11-13). Although M1$_{global}$ strains produce little detectable SpeA protein,

SpeA production can be markedly upregulated following mutation of *covS* (11,12) or *covR* (12, 13)

(Figure 1D), likely accounting for previous reports of upregulation of SpeA following *in vivo* passage

of *emm*1 *S. pyogenes* isolates (14).

Consistent with the above experimental observations, contemporary M1$_{global}$ invasive strains with

covRS mutations arising in patients produce SpeA at amounts equivalent to M1$_{UK}$ strains (9). A

historic *emm*1 isolate NCTC8198 (SF130; H250; H305) (15) produces 5-10 fold more SpeA than

M1$_{UK}$ strains (Figure 1E) yet does not have the M1$_{UK}$ ssrA SNP. It does however possess a stop

mutation in *covS* at residue 318. Northern blotting has previously identified only a single dominant

~750bp *speA* mRNA transcript for this strain (16), potentially consistent with abundant SpeA

production. We postulate that release from indirect or direct covRS repression may allow *speA*

transcription to start at a native promoter site proximal to *speA*, triggered by growth-phase related

factors. Alternatively, this may result in more rapid processing of any longer transcript. Whether

there is an interaction between the ssrA SNP and covRS, other signals, or RNAseY is unknown.

M1$_{UK}$ is associated with a recent upsurge in invasive infections in Europe (17,18). The two

intermediate sublineages, M1$_{13SNP}$ and M1$_{23SNP,}$ appear to be no longer present in England (4).

Despite the M1$_{23SNP}$ sublineage being able to produce SpeA at a level that is indistinguishable from

M1$_{UK}$ isolates (9), it has been fully outcompeted by M1$_{UK}$ in England (4). Although increased SpeA

production is predicted to perpetuate community transmission, the fitness advantage(s) of M1$_{UK}$ may

yet require additional features that promote *emm*1 survival in the human population, underlining a

need for further research and surveillance.

Author contributions.

AV, VWCS, HKL, and SS conceived the work described. AV, VS, HKL, XZ, EJ, LR, KKH, KYM, OC,

JC (UKHSA) undertook data collection, analysis, and data presentation. AV, VWCS, and SS wrote

the first draft of the report and all authors contributed to its revision.

The authors declare no competing interests

Acknowledgements. Funding was from UKRI (MR/P022669/1); Wellcome Trust Collaborative

award (215539/Z/19/Z); the NIHR Imperial Biomedical Research Centre (BRC). VWCS is supported

by an MRC Career Development Award (MR/X007421/1) and HKL was supported by an MRC CMBI

Clinical Training Fellowship. The authors acknowledge the support of the NIHR Health Protection

Research Unit in Healthcare-associated infections and AMR at Imperial and the Imperial BRC

Leonard and Dora Colebrook Laboratory. The authors thank all those who contribute strains to the

national and northwest London collections of *S. pyogenes* and the diagnostic and reference

laboratories that process these.

References

1. World Health Organization. Increase in invasive Group A streptococcal infections among children
in Europe, including fatalities. https://www.who.int/europe/news/item/12-12-2022-increase-in-invasive-group-a-streptococcal-infections-among-children-in-europe--including-fatalities#:~:text=The%20observed%20increases%20reported%20to,during%20the%20COVID%2D19%20pandemic. Accessed, April 30th 2023.

2. UK Health Security Agency. Group A Streptococcal infections: 13th update on seasonal activity in
England. April 23rd 2023 https://www.gov.uk/government/publications/group-a-streptococcal-infections-activity-during-the-2022-to-2023-season/group-a-streptococcal-infections-13th-update-on-seasonal-activity-in-england. Accessed April 30th 2023.

3. Lynskey NN, Jauneikaite E, Li HK, Zhi X, Turner CE, Mosavie M, Pearson M, Asai M, Lobkowicz
L, Chow JY, Parkhill J, Lamagni T, Chalker VJ, Sriskandan S. Emergence of dominant toxigenic
M1T1 *Streptococcus pyogenes* clone during increased scarlet fever activity in England: a population-based molecular epidemiological study. Lancet Infect Dis. 2019 Nov;19(11):1209-1218. doi: 10.1016/S1473-3099(19)30446-3.

4. Zhi X, Li HK, Li H, Loboda Z, Charles S, Vieira A, Huse K, Jauneikaite E, Reeves L, Mok KY,
Coelho J, Lamagni T, Sriskandan S. Emerging invasive group A Streptococcus M1$_{UK}$ lineage

detected by allele-specific PCR, England, 2020. Emerg Infect Dis. 2023 Apr 5;29(5). doi:

10.3201/eid2905.221887.

5. Rümke LW, de Gier B, Vestjens SMT, van der Ende A, van Sorge NM, Vlaminckx BJM,

Witteveen S, van Santen M, Schouls LM, Kuijper EJ. Dominance of M1$_{UK}$ clade among Dutch M1

Streptococcus pyogenes. Lancet Infect Dis. 2020 May;20(5):539-540. doi: 10.1016/S1473-

3099(20)30278-4.

6. Demczuk W, Martin I, Domingo FR, MacDonald D, Mulvey MR. Identification of *Streptococcus*

pyogenes M1$_{UK}$ clone in Canada. Lancet Infect Dis. 2019 Dec;19(12):1284-1285. doi:

10.1016/S1473-3099(19)30622-X.

7. Li Y, Nanduri SA, Van Beneden CA, Beall BW. M1$_{UK}$ lineage in invasive group A streptococcus

isolates from the USA. Lancet Infect Dis. 2020 May;20(5):538-539. doi: 10.1016/S1473-

3099(20)30279-6.

8.Davies MR, Keller N, Brouwer S, Jespersen MG, Cork AJ, Hayes AJ, Pitt ME, De Oliveira DMP,

Harbison-Price N, Bertolla OM, Mediati DG, Curren BF, Taiaroa G, Lacey JA, Smith HV, Fang NX,

Coin LJM, Stevens K, Tong SYC, Sanderson-Smith M, Tree JJ, Irwin AD, Grimwood K, Howden BP,

Jennison AV, Walker MJ. Detection of *Streptococcus pyogenes* M1$_{UK}$ in Australia and

characterization of the mutation driving enhanced expression of superantigen SpeA. Nat Commun.

2023;14(1):1051. doi: 10.1038/s41467-023-36717-4.

9. Li HK, Zhi X, Vieira A, Whitwell HJ, Schricker A, Jauneikaite E, Li H, Yosef A, Andrew I, Game L,

Turner CE, Lamagni T, Coelho J, Sriskandan S. Characterization of emergent toxigenic M1$_{UK}$

Streptococcus pyogenes and associated sublineages. Microb Genom. 2023 Apr;9(4). doi:

10.1099/mgen.0.000994. PMID: 37093716.

10. Cleary PP, McLandsborough L, Ikeda L, Cue D, Krawczak J, Lam H. High-frequency intracellular

infection and erythrogenic toxin A expression undergo phase variation in M1 group A streptococci.

Mol Microbiol. 1998 Apr;28(1):157-67. doi: 10.1046/j.1365-2958.1998.00786.x.

11. Sumby P, Whitney AR, Graviss EA, DeLeo FR, Musser JM. Genome-wide analysis of group A

streptococci reveals a mutation that modulates global phenotype and disease specificity. PLoS

Pathog. 2006 Jan;2(1):e5. doi: 10.1371/journal.ppat.0020005.

12. Treviño J, Perez N, Ramirez-Peña E, Liu Z, Shelburne SA 3rd, Musser JM, Sumby P. CovS

simultaneously activates and inhibits the CovR-mediated repression of distinct subsets of group A

Streptococcus virulence factor-encoding genes. Infect Immun. 2009 Aug;77(8):3141-9. doi:

10.1128/IAI.01560-08.

13. Horstmann N, Myers KS, Tran CN, Flores AR, Shelburne Iii SA. CovS inactivation reduces CovR

promoter binding at diverse virulence factor encoding genes in group A Streptococcus. PLoS Pathog.

2022 Feb 18;18(2):e1010341. doi: 10.1371/journal.ppat.1010341

14. Kazmi SU, Kansal R, Aziz RK, Hooshdaran M, Norrby-Teglund A, Low DE, Halim AB, Kotb M.

Reciprocal, temporal expression of SpeA and SpeB by invasive M1T1 group A streptococcal isolates

in vivo. Infect Immun. 2001 Aug;69(8):4988-95. doi: 10.1128/IAI.69.8.4988-4995.2001

15. Griffith F. Types of Haemolytic Streptococci in Relation to Scarlet Fever. The Journal of Hygiene, Vol.

25, No. 4 (Nov., 1926), pp. 385-397

16. Unnikrishnan M, Cohen J, Sriskandan S. Growth-phase-dependent expression of virulence

factors in an M1T1 clinical isolate of *Streptococcus pyogenes*. Infect Immun. 1999 Oct;67(10):5495-

9. doi: 10.1128/IAI.67.10.5495-5499.1999

17. Guy R, Henderson KL, Coelho J, Hughes H, Mason EL, Gerver SM, Demirjian A, Watson C,

Sharp A, Brown CS, Lamagni T. Increase in invasive group A streptococcal infection notifications,

England, 2022. Euro Surveill. 2023 Jan;28(1):2200942. doi: 10.2807/1560-

7917.ES.2023.28.1.2200942.

18. van der Putten BCL, Vlaminckx BJM, de Gier B, Freudenburg-de Graaf W, van Sorge NM. Group

A streptococcal meningitis with the $M1_{UK}$ variant in the Netherlands. JAMA. 2023 Apr 7:e235927. doi:

10.1001/jama.2023.5927.

Figure 1

245
246

Figure 1. M1ᴜₖ and sublineages; complex regulation of SpeA production

Phylogenetic tree constructed with 413 core SNPs (without recombination regions) obtained by mapping

269 *emm*1 *S. pyogenes* isolates selected from UK to demonstrate sublineages (ref. 7) with 319 isolates

from Australia (ref. 1) to the reference genome MGAS5005 **(A)**; coloured tips indicate the various

sublineages, with Australian strains indicated by lighter colours and a UK strain with 19SNPs in grey. **(B)**

Introduction of ssrA SNP into M1ɢₗₒᵦₐₗ strain H1499 and M1₁₃SNP strain H1750 results in detectable SpeA

in supernatants when tested using SpeA immunoblot. AphA3 (kanR) was introduced upstream of the ssrA

leader sequence as a selectable marker in all transformants. Only transformants with the ssrA SNP

introduced expressed detectable SpeA. For comparison, M1ᴜₖ strain H1490 also shown. **C.** RNA

sequencing read abundance across *ssrA*; an intergenic region which includes *ptx*; and *speA* comparing

four M1ɢₗₒᵦₐₗ strains (blue lines) and four M1ᴜₖ strains (red lines). All were non-invasive throat isolates.

M1ɢₗₒᵦₐₗ isolates were BHS0674 (H1499); BHS0162 (H1489); BHS0130 (H1504); and BHS0151 (H1488).

M1ᴜₖ isolates were BHS0170 (H1490); BHS0128 (H1503); BHS0258 (H1491); and BHS0581 (H1496).

D. SpeA immunoblot of supernatant from M1ɢₗₒᵦₐₗ strain H584 and an isogenic covR T65P mutant (H1565).

E. SpeA immunoblot of concentrated *emm*1 supernatants from M1ɢₗₒᵦₐₗ strains (H1488; H1489; H1492;

H1495; H1499) and M1ᴜₖ strains (H1490; H1491; H1493; H1539; H1496) (all non-invasive throat strains)

compared with non-concentrated supernatant from *emm*1 strain H305 (NCTC8198).