1

Multi-continental detection of *Streptococcus pyogenes* M1$_{UK}$:

Impact of ssrA SNP on SpeA expression in ancestral and

M1$_{UK}$ isolates.

6   Ana Vieira[1,2], Ho Kwong Li[1,2], Xiangyun Zhi[1,2], Lucy Reeves[1], Kristin K. Huse[1,2], Kai Yi Mok[1], Oscar
7   Cowen[1], Elita Jauneikaite[2,3], Juliana Coelho[4], Shiranee Sriskandan**[1,2], Valerie WC Soo**[1,2].
8

[1]Department of Infectious Disease, Faculty of Medicine, Imperial College London, London, UK.

[2]Centre for Bacterial Resistance Biology, Imperial College London, London, UK.

[3]School of Public Health, Faculty of Medicine, Imperial College London, London, UK.

[4]Centre for Infections, UK Health Security Agency, London, UK,

**Co-corresponding authors Valerie Soo v.soo@imperial.ac.uk; Shiranee Sriskandan

s.sriskandan@imperial.ac.uk;

Abstract

The *Streptococcus pyogenes* M1$_{UK}$ lineage, characterised by an intrinsic ability to express SpeA toxin and defined by 27 single nucleotide polymorphisms in the core genome, dominates the population of *emm*1 *S. pyogenes* isolates throughout Europe, Canada, South America, Japan and Oceania. While not the sole deterministic factor, enhanced SpeA expression is likely to have contributed to M1$_{UK}$ lineage expansion, but was not sufficient to support expansion of intermediate lineage M1$_{23SNP}$, that expresses SpeA at a similar level to M1$_{UK}$.

A single nucleotide polymorphism (SNP) in the *ssrA* leader sequence upstream of *speA* is one of a limited number of SNPs that distinguish intermediate sublineages that differ in SpeA production. It was recently shown that this SNP leads to increased *ssrA* terminator read-through, and consequent increased transcription of *speA*, which lies downstream of *ssrA*. In this work, introduction of the *ssrA* SNP into representative isolates of the widely disseminated M1$_{global}$ clone and the intermediate M1$_{13SNP}$ lineage, that cannot otherwise produce readily-detectable SpeA in culture, resulted in SpeA expression, confirming the importance of the *ssrA* SNP to SpeA phenotype. Consistent with this, correction of the *ssrA* SNP in M1$_{UK}$ abrogated SpeA expression. However, RNAseq analysis of 8 *emm*1 clinical pharyngitis strains showed that presence of the SNP was not invariably linked to read-through from the ssrA leader sequence or SpeA expression. Read-through was observed in isolates that did not possess the ssrA SNP. Critical review of existing data suggests that *speA* mRNA transcript length may be impacted by the two-component regulator CovRS, pointing to a complex regulatory network interaction between the bacterial chromosome and phage-encoded superantigens.

Introduction

A marked increase in serious *Streptococcus pyogenes* infections was reported in several countries following relaxation of public health interventions designed to limit the spread of COVID-19 [1]. In England, notifications of both scarlet fever and invasive *S. pyogenes* infections increased throughout 2022; infections peaked at the end of the calendar year but remained high thereafter [2]. In the 30 weeks from mid September 2022-mid April 2023 there were 54,394 notifications of scarlet fever and 2965 invasive *S. pyogenes* infections in England alone [2]. The high case fatality rate of 14.2% [2] underlines the major public health impact of the observed increase, and the importance of understanding the relative roles of strain pathogenicity, population immune susceptibility, co-infections, season, and access to healthcare.

*Emm*1 *S. pyogenes* strains are known to be inherently invasive and normally account for one quarter of invasive infections; however, in the 2022-2023 season, *emm*1 isolates accounted for over half of invasive infections, rising to over 60% in children in England [2]. The population of *emm*1 *S. pyogenes* isolates in England is dominated by a novel sublineage, $M1_{UK}$, that has an intrinsic ability to express the phage-encoded superantigenic toxin, streptococcal pyrogenic exotoxin (Spe) A [3,4]. $M1_{UK}$ strains can be differentiated from strains belonging to the highly successful *emm*1 *S. pyogenes* globally-disseminated clone that emerged in the 1980's ($M1_{global}$) by just 27 single nucleotide polymorphisms in the core genome [3]. The $M1_{UK}$ lineage has been identified elsewhere in Europe [5], North America [6,7], Japan [8], China, Russia [9] and Australia, where isolates from Queensland and Victoria date from 2013 [10], just 3 years after its first detection in England [3]. While $M1_{UK}$ represented ~60% of Australian *emm*1 isolates by 2020 [10], in England, this proportion reached 91% by 2020 [4]. The detection of $M1_{UK}$ Queensland, underlines the ability of this lineage to spread in temperate and more tropical climates, although notably *emm*1 strains appear to be uncommon in sub-Saharan Africa [11]. Of added concern are reports of strains infected with a ΦHKU488.vir-like phage, that carries both *speC* and *ssa* superantigen genes in Australia [10]. To date, this phage has not been identified in *emm*1 strains from England, although ~10% of *emm*1 strains in England do possess a phage encoding *speC* and *spd*.

Previous genomic analysis of *emm*1 *S. pyogenes* strains in England identified two intermediate

sublineages in addition to $M1_{global}$ and $M1_{UK,}$ with just 13 and 23 of the 27 SNPs that define $M1_{UK}$ [3],

namely $M1_{13SNP}$ and $M1_{27SNP}$ (Figure 1). We recently reported that strains from the $M1_{13SNP}$

intermediate sublineage produce negligible amounts of SpeA, similar to $M1_{global}$ [12]. Strains from

the $M1_{23SNP}$ sublineage, in contrast, produce SpeA at an increased level that is indistinguishable from

$M1_{UK}$ isolates [12]. Despite increased SpeA production, $M1_{23SNP}$ isolates could not be detected when

screening all *emm*1 isolates submitted to the reference laboratory in England in 2020 [4], suggesting

an added fitness advantage in $M1_{UK}$ strains beyond production of SpeA alone. We postulate this

further adaptation may be conferred by the additional 4 SNPs that distinguish $M1_{UK}$ from $M1_{23SNP}$

isolates (3 non-synonymous SNPs and an intergenic SNP 39bp upstream of *glpF2*, aquaporin,

expression of which is significantly reduced in $M1_{UK}$) [10,12]. In this report we seek to further

understand the mechanisms that differentiate SpeA-producers from non-producers among *emm*1 *S.*

pyogenes strains that carry the phage that encodes *speA*.

Results and Discussion

The $M1_{UK}$ SNP at position 983438 of the MGAS5005 reference genome, within the leader sequence

of *ssrA* and upstream of the *speA* start site (referred to hereafter as the 'ssrA SNP'), is present in

both $M1_{23SNP}$ and $M1_{UK}$ sublineages, but is absent in $M1_{global}$ and $M1_{13SNP}$ sublineages. Screening of

natural mutants arising during outbreaks identified the ssrA SNP to be associated with increased

SpeA production [12]. Davies et al [10] showed that introduction of the ssrA SNP resulted in

increased expression of SpeA by $M1_{global}$. We introduced the ssrA SNP into an $M1_{global}$ strain, and

into an intermediate $M1_{13SNP}$ strain (Figure 2A). Consistent with the findings of Davies et al [10], SpeA

production in culture supernatants increased in both transformants (Figure 2B). Enhancement of

SpeA was similar in both the $M1_{global}$ and $M1_{13SNP}$ backgrounds, suggesting that the accumulated

SNPs in the $M1_{13SNP}$ lineage, which include almost half of the SNPs that characterize $M1_{UK}$, do not

appreciably contribute to the SpeA phenotype. We considered if our mutagenesis strategy itself

might lead to increased SpeA expression but this was ruled out by use of a control construct in each

strain (Figure 2B). In contrast, the enhanced SpeA expression in an M1$_{UK}$ background could be

reduced considerably upon correction of the ssrA SNP (Figure 2B).

To understand the basis for difference in *speA* expression between the lineages, we compared

RNAseq reads from four randomly selected M1$_{global}$ strains, which do not have the ssrA SNP, and

four randomly selected M1$_{UK}$ strains, which all have this SNP. None of the strains possessed

regulatory gene mutations other than those that characterize M1$_{UK}$. Similar to Davies et al [10], we

found that there was indeed evidence of read-through in the intergenic regions between the end of

ssrA and the start of *speA*. (Figure 2C). However, we did not see clear differentiation between the

four M1$_{global}$ strains and the four M1$_{UK}$ strains within the intergenic region, where read abundance in

two of the M1$_{global}$ strains was almost identical to M1$_{UK}$ strains (Figure 2C). Regardless, transcription

of *speA* was significantly reduced in all four M1$_{global}$ strains compared with M1$_{UK}$. Interestingly, *speA*

read abundance was somewhat higher in the two M1$_{global}$ strains that had more intergenic reads, but

not as high as the M1$_{UK}$ strains. These data provide support for intergenic read-through affecting

speA transcription. However, based on short read RNA sequencing data, we cannot definitively

ascribe read abundance in the intergenic region to the ssrA SNP in all cases, noting that all four of

the M1$_{global}$ strains studied lack any SNP in this region. Comparative WGS analysis between the four

M1$_{global}$ strains failed to detect any other genetic features associated with higher levels of read-

through. Taken together, transcriptional read-through was observed in this region, but the association

with the ssrA SNP and SpeA over-expression was not absolute. Despite not possessing the ssrA

SNP, two of four M1$_{global}$ strains tested showed similar read-through to M1$_{UK}$ strains, yet made less

SpeA, pointing to additional regulatory factors that influence SpeA expression. The ssrA SNP

however was important, at least in the strains examined.

A polycistronic transcript for speA was originally reported ~25 years ago by Cleary *et al*, who

identified both ~2kB and ~900bp *speA* mRNA transcripts in a single *emm*1 *S. pyogenes* strain [13].

In that case, the longer 2kB transcript was associated with colonies that failed to produce SpeA,

while the shorter 900bp transcript, corresponding to the approximate expected size for the *speA*

transcript, was associated with mucoid *emm*1 colonies that produced abundant SpeA. Based on the date of publication, we infer the *emm*1 strain in that study [13] to be from an M1$_{global}$ background. The within-strain variation in *speA* transcript length coupled with mucoid phenotype points to involvement of the two-component regulator covRS (csrRS) which is now known to repress capsule and *speA* transcription in *S. pyogenes* [14–16]. Although M1$_{global}$ strains produce little detectable SpeA protein, SpeA production can be markedly upregulated following mutation of *covS* [14,15] or *covR* [15,16] (Figure 3A), likely accounting for previous reports of upregulation of SpeA following *in vivo* passage of *emm*1 *S. pyogenes* isolates [17].

Consistent with the above experimental observations, contemporary M1$_{global}$ invasive strains with *covRS* mutations arising in patients produce SpeA at amounts equivalent to M1$_{UK}$ strains [12]. A historic *emm*1 isolate NCTC8198 (SF130; H250; H305) [18,19] produces 5-10 fold more SpeA than M1$_{UK}$ strains (Figure 3B) yet does not have the M1$_{UK}$ ssrA SNP. It does however possess a stop mutation in *covS* at residue 318. Northern blotting has previously identified only a single dominant ~750bp *speA* mRNA transcript for this strain [20], potentially consistent with abundant SpeA production. We postulate that release from indirect or direct covRS repression may allow *speA* transcription to start at a native promoter site proximal to *speA*, triggered by growth-phase related factors. Alternatively, this may result in more rapid processing of any longer transcript. Whether there is an interaction between the ssrA SNP and covRS, other signals, or RNAse Y is unknown.

M1$_{UK}$ is associated with the recent upsurge in invasive infections in Europe [21,22]. The two intermediate sublineages, M1$_{13SNP}$ and M1$_{23SNP,}$ appear to be no longer present in England [4]. Despite the M1$_{23SNP}$ sublineage being able to produce SpeA at a level that is indistinguishable from M1$_{UK}$ isolates [12], it has been fully outcompeted by M1$_{UK}$ in England [4]. Although increased SpeA production is predicted to perpetuate community transmission, the fitness advantage(s) of M1$_{UK}$ may yet require additional features that promote *emm*1 survival in the human population, underlining a need for further research and surveillance.

Methods and Materials

Bacterial strains and growth conditions

All bacterial strains used in this study are described in Table 1. *S. pyogenes* isolates were routinely cultured in Todd Hewitt Broth (THB; Oxoid), on THY (THB supplemented with 1% (w/v) yeast extract) agar or Columbia blood agar at 37ºC in 5% CO_2. *Escherichia coli* One Shot™ TOP10 was grown in lysogeny broth (LB) or LB agar at 37ºC. For maintaining and selecting pUCMUT-containing *E. coli* and *S. pyogenes*, 50 µg/mL and 400 µg/mL of kanamycin were added to the growth medium, respectively.

Introduction of the ssrA G-to-C SNP into the M1$_{global}$ and M1$_{13SNP}$ strains

The ssrA G-to-C SNP was introduced separately into *S. pyogenes* M1$_{global}$ and *S. pyogenes* M1$_{13SNP}$ by allelic exchange (Figure 2A) using the suicide vector pUCMUT [19], which harbours the *aph3* gene (coding for kanamycin resistance cassette (Kan^R)). *aph3* is flanked by two combinations of restriction sites on either side: PstI + SalI and KpnI + EcoRI.

The 5' region upstream of ssrA was PCR-amplified from the genomic DNA of *S. pyogenes* M1$_{UK}$ H1490 using primers 993P1 (5'-AA<u>CTGCAG</u>CTATCATCCTTGCATTTGC-3') and 993P2 (5'-ACGC<u>GTCGAC</u>GGTTTGGTGTTAATTTGATAAAC-3'), and the resulting 554-bp PCR product was cloned into the PstI and SalI sites of pUCMUT, resulting in pUCMUT5s. Next, the 3' region containing the ssrA SNP was PCR-amplified using primers 994P3 (5'-GG<u>GGTACC</u>TTAATGCTGTAAGAACCAAGATATAA-3') and 994P4 (5'-AGC<u>GAATTC</u>ATAATGCCTGTCGAATCC-3'), and the 735-bp PCR product was cloned into the KpnI and EcoRI sites of pUCMUT5s to yield the suicide plasmid pUCMUTssrA_M1UK. All constructed plasmids were verified by sequencing at Full Circle Labs.

pUCMUTssrA_M1UK was electroporated into the recipient strains, *S. pyogenes* M1$_{global}$ H1499 and *S. pyogenes* M1$_{13SNP}$ H1750, and was allowed to cross into the chromosome by homologous

recombination. The presence of the ssrA G-to-C SNP in the selected strains resulting from a double allelic exchange was verified by PCR and Sanger sequencing (Genewiz).

Detection of SpeA via immunoblotting

SpeA production in various *S. pyogenes* isolates was assessed using immunoblotting with anti-rSpeA rabbit serum. An overnight culture of *S. pyogenes* (50 mL) was harvested by centrifugation (2,500 rpm, 10 min, 4ºC, swing bucket rotor). The supernatant was passed through an 0.2-µm filter (Sartorius) to obtain a cell-free supernatant. A 500-µL aliquot of this cell-free supernatant was concentrated 5x using a 10 kDa MWCO ultra centrifugal filter (Amicon). The concentrated supernatant was denatured in 1x Bolt™ LDS Sample Buffer (ThermoFisher) containing 300 mM DTT, prior to being fractionated on an 10% Bolt™ Bis-Tris Plus Gel (ThermoFisher). Fractionated proteins were electroblotted onto a nitrocellulose membrane. After blocking the membrane with 5% (w/v) skim milk buffer and 0.05% Tween-20, it was then probed with anti-rSpeA rabbit serum (1:1,000 dilution) [12], and followed by an HRP-conjugated anti-rabbit secondary antibody (1:80,000 dilution). Probed SpeA was detected using chemiluminescence reagents (Cytiva Amersham™ ECL™ Prime Western Blotting Detection Reagents), and the blot was visualised on a Li-COR Odyssey Fc imaging system.

Quantification of the RNA reads in M1$_{global}$ and M1$_{UK}$ strains

The average read coverage at genomic regions covering *ssrA*, *paratox*, and *speA* was determined according to Davies et al [10].

Author contributions

AV, VWCS, HKL, and SS conceived the work described. AV, VS, HKL, XZ, EJ, LR, KKH, KYM, OC, JC (UKHSA) undertook data collection, analysis, and data presentation. AV, VWCS, and SS wrote the first draft of the report and all authors contributed to its revision.

The authors declare no competing interests

Acknowledgements

Funding was from UKRI (MR/P022669/1); Wellcome Trust Collaborative award (215539/Z/19/Z); the NIHR Imperial Biomedical Research Centre (BRC). VWCS is supported by an MRC Career Development Award (MR/X007421/1) and HKL was supported by an MRC CMBI Clinical Training Fellowship. The authors acknowledge the support of the NIHR Health Protection Research Unit in Healthcare-associated infections and AMR at Imperial and the Imperial BRC Leonard and Dora Colebrook Laboratory. The authors thank all those who contribute strains to the national and northwest London collections of *S. pyogenes* and the diagnostic and reference laboratories that process these.

References

1. World Health Organization. Increase in invasive Group A streptococcal infections among children in Europe, including fatalities. In: World Health Organization [Internet]. [cited 23 Feb 2026]. Available: https://www.who.int/europe/news/item/12-12-2022-increase-in-invasive-group-a-streptococcal-infections-among-children-in-europe--including-fatalities

2. UK Health Security Agency. Group A streptococcal infections: 13th update on seasonal activity in England. [cited 23 Feb 2026]. Available: https://www.gov.uk/government/publications/group-a-streptococcal-infections-activity-during-the-2022-to-2023-season/group-a-streptococcal-infections-13th-update-on-seasonal-activity-in-england

3. Lynskey NN, Jauneikaite E, Li HK, Zhi X, Turner CE, Mosavie M, et al. Emergence of dominant toxigenic M1T1 *Streptococcus pyogenes* clone during increased scarlet fever activity in England: a population-based molecular epidemiological study. Lancet Infect Dis. 2019;19: 1209–1218. doi:10.1016/S1473-3099(19)30446-3

4. Zhi X, Li HK, Li H, Loboda Z, Charles S, Vieira A, et al. Emerging invasive Group A *Streptococcus* $M1_{UK}$ lineage detected by allele-specific PCR, England, 2020. Emerg Infect Dis. 2023;29: 1007–1010. doi:10.3201/eid2905.221887

5. Rümke LW, de Gier B, Vestjens SMT, van der Ende A, van Sorge NM, Vlaminckx BJM, et al. Dominance of $M1_{UK}$ clade among Dutch M1 *Streptococcus pyogenes*. Lancet Infect Dis. 2020;20: 539–540. doi:10.1016/S1473-3099(20)30278-4

6. Demczuk W, Martin I, Domingo FR, MacDonald D, Mulvey MR. Identification of Streptococcus pyogenes $M1_{UK}$ clone in Canada. Lancet Infect Dis. 2019;19: 1284–1285. doi:10.1016/S1473-3099(19)30622-X

7. Li Y, Nanduri SA, Van Beneden CA, Beall BW. $M1_{UK}$ lineage in invasive group A streptococcus isolates from the USA. Lancet Infect Dis. 2020;20: 538–539. doi:10.1016/S1473-3099(20)30279-6

8. Ujiie M. Emerging Trends in Streptococcal Toxic Shock Syndrome, Japan. Emerg Infect Dis. 2025;31: 847–849. doi:10.3201/eid3104.241076

9.   Khan H, Lun H, Xu G, Huang K. Global emergence and polycentric evolution of invasive

Streptococcus   pyogenes   M1UK   lineage.   J   Infect.   2025;91:   106653.

doi:10.1016/j.jinf.2025.106653

10.   Davies MR, Keller N, Brouwer S, Jespersen MG, Cork AJ, Hayes AJ, et al. Detection of

Streptococcus pyogenes M1UK in Australia and characterization of the mutation driving

enhanced   expression   of   superantigen   SpeA.   Nat   Commun.   2023;14:   1051.

doi:10.1038/s41467-023-36717-4

11.   Bah SY, Keeley AJ, Armitage EP, Khalid H, Chaudhuri RR, Senghore E, et al. Genomic

Characterization of Skin and Soft Tissue Streptococcus pyogenes Isolates from a Low-Income

and a High-Income Setting. mSphere. 2022; e0046922. doi:10.1128/msphere.00469-22

12.   Li HK, Zhi X, Vieira A, Whitwell HJ, Schricker A, Jauneikaite E, et al. Characterization of

emergent toxigenic $M1_{UK}$ *Streptococcus pyogenes* and associated sublineages. Microb Genom.

2023;9. doi:10.1099/mgen.0.000994

13.   Cleary PP, McLandsborough L, Ikeda L, Cue D, Krawczak J, Lam H. High-frequency

intracellular infection and erythrogenic toxin A expression undergo phase variation in M1 group

A streptococci: Co-variation of M protein, hyaluronic acid and erythrogenic toxin A in group A

streptococci. Mol Microbiol. 1998;28: 157–167. doi:10.1046/j.1365-2958.1998.00786.x

14.   Sumby P, Whitney AR, Graviss EA, DeLeo FR, Musser JM. Genome-wide analysis of group a

streptococci reveals a mutation that modulates global phenotype and disease specificity. PLoS

Pathog. 2006;2: e5. doi:10.1371/journal.ppat.0020005

15.   Treviño J, Perez N, Ramirez-Peña E, Liu Z, Shelburne SA 3rd, Musser JM, et al. CovS

simultaneously activates and inhibits the CovR-mediated repression of distinct subsets of group

A *Streptococcus* virulence factor-encoding genes. Infect Immun. 2009;77: 3141–3149.

doi:10.1128/IAI.01560-08

16.   Horstmann N, Myers KS, Tran CN, Flores AR, Shelburne SA Iii. CovS inactivation reduces CovR

promoter binding at diverse virulence factor encoding genes in group A *Streptococcus*. PLoS

Pathog. 2022;18: e1010341. doi:10.1371/journal.ppat.1010341

17. Kazmi SU, Kansal R, Aziz RK, Hooshdaran M, Norrby-Teglund A, Low DE, et al. Reciprocal,

temporal expression of SpeA and SpeB by invasive M1T1 group a streptococcal isolates in vivo.

Infect Immun. 2001;69: 4988–4995. doi:10.1128/IAI.69.8.4988-4995.2001

18. Griffith F. Types of haemolytic streptococci in relation to Scarlet fever. J Hyg (Lond). 1926;25:

385–397. doi:10.1017/s0022172400017502

19. Sriskandan S, Unnikrishnan M, Krausz T, Cohen J. Molecular analysis of the role of

streptococcal pyrogenic Exotoxin A (SPEA) in invasive soft-tissue infection resulting from

Streptococcus pyogenes. Mol Microbiol. 1999;33: 778–790. doi:10.1046/j.1365-

2958.1999.01525.x

20. Unnikrishnan M, Cohen J, Sriskandan S. Growth-phase-dependent expression of virulence

factors in an M1T1 clinical isolate of *Streptococcus pyogenes*. Infect Immun. 1999;67: 5495–

5499. doi:10.1128/IAI.67.10.5495-5499.1999

21. Guy R, Henderson KL, Coelho J, Hughes H, Mason EL, Gerver SM, et al. Increase in invasive

group A streptococcal infection notifications, England, 2022. Euro Surveill. 2023;28.

doi:10.2807/1560-7917.ES.2023.28.1.2200942

22. van der Putten BCL, Vlaminckx BJM, de Gier B, Freudenburg-de Graaf W, van Sorge NM.

Group A streptococcal meningitis with the M1$_{UK}$ variant in the Netherlands. JAMA. 2023;329:

1791–1792. doi:10.1001/jama.2023.5927

23. Turner CE, Dryden M, Holden MTG, Davies FJ, Lawrenson RA, Farzaneh L, et al. Molecular

analysis of an outbreak of lethal postpartum sepsis caused by *Streptococcus pyogenes*. J Clin

Microbiol. 2013;51: 2089–2095. doi:10.1128/JCM.00679-13

24. Lamb LE, Siggins MK, Scudamore C, Macdonald W, Turner CE, Lynskey NN, et al. Impact of

contusion injury on intramuscular emm1 group a streptococcus infection and lymphatic spread.

Virulence. 2018;9: 1074–1084. doi:10.1080/21505594.2018.1482180

27.102

Figure 1. M1$_{UK}$ and sublineages

Phylogenetic tree constructed with 413 core SNPs (without recombination regions) obtained by mapping

269 *emm*1 *S. pyogenes* isolates selected from UK to demonstrate sublineages [12] with 319 isolates from

Australia [10] to the reference genome MGAS5005. Coloured tips indicate the various sublineages, with

Australian strains indicated by lighter colours and a UK strain with 19SNPs in grey.

Figure 2. Complex regulation of SpeA production

A. The overall schematic for introducing the ssrA SNP into a *S. pyogenes* M1$_{global}$ and M1$_{13SNP}$ lineages.

B. Introduction of the ssrA SNP into M1$_{global}$ strain H1499 and M1$_{13SNP}$ strain H1750 results in detectable SpeA in supernatants when tested using SpeA immunoblot. AphA3 (KanR) was introduced upstream of the ssrA leader sequence as a selectable marker in all transformants. Only transformants with the ssrA SNP introduced expressed detectable SpeA. In contrast, M1UK strain H1490 expressed considerably lower SpeA upon correction of its ssrA SNP. **C.** RNA sequencing read abundance across *ssrA*; an intergenic region which includes *ptx (paratox)*; and *speA* comparing four M1$_{global}$ strains (blue lines) and four M1$_{UK}$ strains (red lines). All were non-invasive throat isolates. M1$_{global}$ isolates were BHS0674 (H1499); BHS0162 (H1489); BHS0130 (H1504); and BHS0151 (H1488). M1$_{UK}$ isolates were BHS0170 (H1490); BHS0128 (H1503); BHS0258 (H1491); and BHS0581 (H1496).

Figure 3. Regulation of SpeA production beyond ssrA SNP

A. SpeA immunoblot of supernatant from M1$_{global}$ strain H584 and an isogenic covR T65P mutant (H1565).

B. SpeA immunoblot of concentrated *emm*1 supernatants from M1$_{global}$ strains (H1488; H1489; H1492; H1495; H1499) and M1$_{UK}$ strains (H1490; H1491; H1493; H1539; H1496) (all non-invasive throat strains) compared with non-concentrated supernatant from *emm*1 strain H305 (NCTC8198).

320　**Table 1. Bacterial strains used in this study.**

Strain	Description	Reference/Source
S. pyogenes M1$_{global}$ H1499	Imperial ENT isolate collection	[3]
S. pyogenes M1$_{global}$ H1499 + Kan^R	Isogenic strain of H1499 with only the Kan^R introduced upstream of ssrA	This study
S. pyogenes M1$_{global}$ H1499 + Kan^R + ssrA SNP	Isogenic strain of H1499 with both the Kan^R and ssrA SNP (G-to-C) introduced	This study
S. pyogenes M1$_{13SNP}$ H1750	PHE iGAS collection	[12]
S. pyogenes M1$_{13SNP}$ H1750 + Kan^R	Isogenic strain of H1750 with only the Kan^R introduced upstream of ssrA	This study
S. pyogenes M1$_{13SNP}$ H1750 + Kan^R + ssrA SNP	Isogenic strain of H1750 with both the Kan^R and ssrA SNP (G-to-C) introduced	This study
S. pyogenes M1$_{UK}$ H1490	Imperial ENT isolate collection	[3]
S. pyogenes M1$_{global}$ H584	Invasive puerperal sepsis blood isolate	[23]
S. pyogenes M1$_{global}$ covR H1565	Isogenic CovR T65P mutant of H584	[24]
S. pyogenes M1$_{global}$ H1488	Imperial ENT isolate collection	[3]
S. pyogenes M1$_{global}$ H1489	Imperial ENT isolate collection	[3]

S. pyogenes M1$_{global}$ H1492	Imperial ENT isolate collection	[3]
S. pyogenes M1$_{global}$ H1495	Imperial ENT isolate collection	[3]
S. pyogenes M1$_{global}$ H1504	Imperial ENT isolate collection	[3]
S. pyogenes M1$_{global}$ H1488	Imperial ENT isolate collection	[3]
S. pyogenes M1$_{UK}$ H1491	Imperial ENT isolate collection	[3]
S. pyogenes M1$_{UK}$ H1493	Imperial ENT isolate collection	[3]
S. pyogenes M1$_{UK}$ H1539	Imperial ENT isolate collection	[3]
S. pyogenes M1$_{UK}$ H1496	Imperial ENT isolate collection	[3]
S. pyogenes M1$_{UK}$ H1503	Imperial ENT isolate collection	[3]
S. pyogenes M1$_{global}$ H305	NCTC8198	[19]
E. coli One Shot™ TOP10	Cloning host	Invitrogen