Breakthrough Report

Title: Multiplexed effector screening for recognition by endogenous resistance genes using

positive defense reporters in wheat protoplasts

Salome Wilson*, Bayantes Dagvadorj*, Rita Tam, Lydia Murphy, Nigel Heng, Emma Crean[$],

John P. Rathjen, and Benjamin Schwessinger

Research School of Biology, The Australian National University, Canberra, ACT, 2601,

Australia

*These authors contributed equally to this work

[$]Current address: Institute for Plant Sciences, University of Cologne, Cologne, Germany

Corresponding author: Benjamin Schwessinger

Email: benjamin.schwessinger@anu.edu.au

Short Title:

Defense signaling in wheat protoplasts

Abstract

Plant resistance (*R*) and pathogen avirulence (*Avr*) gene interactions play key roles in crop

resistance to pathogens. Efficient molecular screening tools for crop species are required to

define interactions that trigger host resistance and to understand the virulence mechanisms of

pathogenic fungi. *Avr* identification supports genome-based pathogen surveillance to enhance

pre-emptive breeding of disease resistant cereal crops. To this end, we have developed a novel

wheat protoplast assay that enables efficient screening of Avr/R interactions at scale. Our assay

allows access to the extensive gene pool of phenotypically described *R* genes because it does not

require the overexpression of cloned *R* genes and is applicable to all wheat cultivars tested so far.

It is suitable for multiplexed *Avr* screening, with interactions tested in pools of up to ten *Avr*

candidates. Our assay is based on newly identified Avr/R-induced marker genes which we used

to create promoter-luciferase reporter constructs. We combined these reporter constructs with a

dual-color ratiometric reporter system that normalizes read-outs accounting for experimental

variability and Avr/R-induced cell death. Our new assay increases the throughput of *Avr*

candidate screening, accelerating the study of cellular defense signaling and resistance gene

identification in wheat. We anticipate that the uptake of our assay by the community will

significantly accelerate *Avr* identification for many wheat pathogens, leading to improved

genomic pathogen surveillance and breeding of disease resistant crops.

Introduction

Plant pathogenic fungi cause severe damage to cereal crops. These pathogens secrete hundreds of

proteinaceous molecules called effectors during infection which manipulate plant processes to

facilitate disease (Dodds and Rathjen, 2010). Some effectors can be recognized by resistance (R)

proteins and are therefore also called avirulence (Avr) proteins. Avr recognition triggers a suite

of defense-related host responses to fight off pathogen infection including the hypersensitive

response (HR) that leads to cell-death at the site of infection and halts pathogen growth

(Laflamme et al., 2020). The deployment of *R* genes is a major way to reduce the negative

impact of pathogens on crop production including in wheat. Recent years have made significant

progress in identifying and cloning a wide variety of *R* genes in wheat (Wulff and Krattinger,

2022). Yet the identification of the correspoding *Avr* genes is lacking far behind. For example,

there are only four confirmed *Blumeria graminis* f. sp. *tritici Avr* genes (Praz et al., 2017;

Bourras et al., 2019; Hewitt et al., 2021; Müller et al., 2022), two for *Zymoseptoria tritici* (Zhong

et al., 2017; Meile et al., 2018), and five for *Puccinia graminis* f. sp. *tritici* (Chen et al., 2017;

Salcedo et al., 2017; Upadhyaya et al., 2021; Arndell et al., 2023). There are no confirmed *Avr*

genes for *Puccinia triticina* and *Puccunia striiformis* f. sp. *tritici* despite both belonging to the

top five wheat pathogens globally (Savary et al., 2019). This lack of confirmed *Avr* genes in

wheat pathogen and especially rust fungi is confounded by the absence of an efficient screening

tool for *Avr* recognition in wheat. Most current methods require to have *Avr* and *R* gene

candidates cloned, express them heterologously, and rely on HR as general read-out for Avr/R

interactions. For example, the model plant *Nicotiana benthamiana* is used to validate Avr/R

interactions as it produces visible HR upon transient expression of *Avr/R* genes through

agroinfiltration. Virus-mediated overexpression (VOX) can be used to screen *Avr* candidates in

wheat cultivars carrying the corresponding *R* gene as the Avr recognition will lead to visible HR

symptoms on the infected leaves (Lee et al., 2012; Bouton et al., 2018). However, the method is

only applicable to small proteins (Lee et al., 2012) and used viruses are not accessible globally

due to biosecurity regulations (Bouton et al., 2018). Recently, HR-based assays in wheat

protoplasts have emerged as a potential alternative to test Avr/R interactions (Saur et al., 2019).

In this transient assay, wheat protoplasts are co-transfected with *Avr/R* genes along with a

luciferase reporter gene. Avr/R interaction triggers cell death causing significant decreases in

luciferase enzyme activity. This assay has been used to validate *Avr* candidates from wheat rust

pathogens (Saur et al., 2019; Upadhyaya et al., 2021; Luo et al., 2021; Ortiz et al., 2022).

However, the HR-based wheat protoplast assay in its current form has several limitations that

make it impractical for high-throughput *Avr/R* screening. Firstly, it requires the transgenic

overexpression of both the *Avr* and *R* genes from plasmids (Saur et al., 2019). This prevents its

use when the *R* gene has not yet been cloned thus stopping access to large numbers of

phenotypically described but uncloned *R* genes (McIntosh et al., 1995). Moreover, it is well

known that luminescence is highly variable between biological replicates and experiments (Saur

et al., 2019; González-Grandío et al., 2021) which can make the results difficult to interpret for

high-throughput screening purposes. In addition, the assay in its current form relies on the

reduction of luciferase activity as read out which is unspecific and cannot be differentiated from

cell-death that is unrelated to HR. These limitations encouraged us to develop a novel assay in

wheat protoplast that relies on a positive defense specific read-out and accounts for experimental

variability via normalization. Our new assay allows the research community to access the whole

wheat gene pool to screen for Avr/R interactions and enables the investigation of downstream

defense signaling directly in wheat.

Results

Cell death triggered by Avr/R interactions in protoplasts is time sensitive

We first sought to characterize the time of HR onset in wheat protoplasts following *Avr/R*

recognition. We used the two published *Avr/R* gene pairs, *AvrSr35/Sr35* (Salcedo et al., 2017;

Saintenac et al., 2013) and *AvrSr50/Sr50* (Chen et al., 2017; Mago et al., 2015), from wheat stem

rust and *P. graminis* f. sp. *tritici* for initial method development. We aimed to identify the

optimal time point for RNAseq analyses that captures transcriptional changes prior to the onset

of HR triggered by Avr/R interactions. As previously described (Saur et al., 2019), we used

luminescence as a proxy for cell viability after overexpression of the *luciferase* reporter gene in

conjunction with each *Avr/R* combination and corresponding control. We expressed all *Avr* and *R*

genes under control of maize ubiquitin promoter (*UBI*) (Christensen et al., 1992) when expressed

from a plasmid. We used the wheat cultivar (cv.) Fielder for the analysis of *AvrSr35/Sr35* and

expressed both genes from a plasmid. For *AvrSr50/Sr50*, we made use of the introgression of

Sr50 into cv. Gabo (GaboSr50) and generated protoplast from cv. Gabo and cv. GaboSr50

(Jensen and Saunders, 2023). We only expressed *AvrSr50* from a plasmid. We also included the

controls, the reporter alone and the reporter with *Avr* or *R* alone (Zenodo Supplemental Data S1).

We measured luminescence at 3, 4, 5, 6, and 18 h post-transfection (hpt). We observed a gradual

increase in luminescence measurements in protoplast expressing luciferase alone or when

expressed together with an *Avr* or *R* gene only (Figure 1, A and B). The luciferase activity was

strongest at 18 hpt in all treatment groups except for the ones expressing *Avr/R* pairs. In cv.

Fielder expressing *AvrSr35/Sr35* and in cv. GaboSr50 expressing *AvrSr50*, luminescence

measurements only increased slightly early on during the time course when compared to mock

treatments of protoplasts transfected with no DNA (Figure 1 C). At about 4 to 5 hpt we observed

a divergence of luciferase activity from control treatment groups when compare with the Avr/R

interaction treatment groups (Figure 1A and B). The lack of continuous increase of luciferase

activity specifically in protoplast expressing *Avr/R* pairs suggests that their co-expressing leads

to cell death in wheat protoplasts soon after transfection.

Transfected protoplasts are suitable for replicable transcriptional analysis

We concluded that 4 hpt is a suitable time point to capture *Avr/R*-dependent transcriptional

changes before strong HR induction based on our time course analysis. We repeated the

experimental setup while omitting the reporter plasmid and collected protoplasts at 4 hpt for

RNA sequencing (Figure 2 A, Treatment groups table Zenodo Supplemental Data S2). We

included four biological replicates per treatment group and sequenced each sample on the

Illumina platform generating >20 million 150 bp paired-end reads per replicate. We used kallisto

(Bray et al., 2016) for transcript quantification and DESeq2 (Love et al., 2014) for downstream

differential expression analysis. As a first quality control step we quantified the expression of all

Avr and *R* genes to investigate their specific expression pattern at 4 hpt. This analysis clearly

showed that all genes are correctly expressed and at similar levels, including in treatment groups

that co-express *Avr/R* gene pairs (Figure 2 A). In addition, it revealed that the endogenous

expression of *Sr50* in cv. GaboSr50 is much lower level than the overexpressed of *Sr35* in cv.

Fielder from a plasmid under the *UBI* promoter. Next, we performed principal component

analysis (PCA) of all sample replicates combined. This revealed that wheat cultivar had the

strongest impact on the observed variation in transcript abundance and accounted for close to

95% of all variation on PC1 and PC2 combined (Figure 2 B). Consequently, we analyzed each

cultivar separately. The wheat cultivar specific PCA analysis showed a clear impact of Avr/R

interaction on transcript abundance variation for both *AvrSr35/Sr35* and *AvrSr50/Sr50* (Figure 2

C and D). This impact was replicable because all four biological replicates clustered closely

together and clearly separated treatment groups including Avr/R interactions sample types from

their controls. This is consistent with the cell death results seen in Figure 1 A and B. Overall,

these results suggest that these data could be used to identify reporter genes specifically

expressed downstream of Avr/R interactions in wheat protoplasts.

Avr/R recognition in wheat protoplasts leads to defense signaling

Next, we investigated if the observed gene expression changes in *Avr/R* expressing protoplast

overlapped with gene ontology (GO) terms commonly associated with defense signaling

pathways in wheat and other plant species. We identified all genes that are specifically

upregulated in response to Avr/R interactions in cv. Gabo/Gabo50 and cv. Fielder above a 2-fold

change threshold. We identified 309 and 922 genes in *AvrSr35/Sr35* and *AvrSr50/Sr50*

expressing protoplasts, respectively, when compared to all their control treatment groups.

Complete gene lists are available on Zenodo as Supplemental Data S3. We performed a

statistical enrichment analysis using these gene lists to test for overrepresentation of defense

signaling associated GO terms. Our results revealed significant enrichment for GO terms related

to cellular components of cell periphery (GO:0071944) and plasma membrane (GO:0005886) in

both cultivars (Figure 2 E). We identified defense response to other organisms (GO:0098542),

response to other organisms (GO:0051707), and biological process involved in interspecies

interaction between organisms (GO:0044419) when looking at cv. GaboSr50 alone (Zenodo

Supplemental Data S4). We found molecular functions GO terms being enriched for calcium ion

binding (GO:0005509), cation binding (GO:0043169), and calcium ion transmembrane

transporter activity (GO:0015085) in both cultivars (Figure 2 E). Our GO terms analysis overall

confirms that *Avr/R* expressing protoplasts undergo defense signaling before the onset of strong

cell death.

We were then interested in genes that are commonly upregulated by the Avr/R interaction in

both cv. GaboSr50 and cv. Fielder. We identified 66 genes commonly upregulated in both

cultivars, (Zenodo Supplemental Data S5) of which only 16 genes were associated with three

different GO terms, all related to the molecular function of ion binding (Zenodo Supplemental

Data S4). Our results showed that several of the identified genes have been previously implicated

in plant defense signaling including TaNHL10 (TRAESCS3D02G368800) (Century et al., 1995;

Aarts et al., 1998; Dagvadorj et al., 2022).

In summary, we observed the upregulation of genes involved in defense signaling in wheat

protoplasts expressing *Avr/R* pairs. The promoters of commonly upregulated genes, by two

Avr/pairs and in two wheat cultivars (Figure 2F), are promising candidates to generate *Avr/R*

inducible reporters.

Identification of positive defense reporters in wheat protoplasts

We further investigated the expression patterns of the 66 commonly upregulated genes to select a

subset whose promoters we could use as positive defense signaling reporters. We selected 13

promoters (D1-D7, D10-D12, and D14-D16) based on their genes' expression patterns in both

cultivars (Figure 2 F). To further evaluate these candidates, we cloned or synthesized their

promoters, consisting of 600-800 bp upstream of their respective genes' start codons. We

accounted for the impact of *Avr/R* induced HR upon luciferase activity (Figure 1) by adapting a

dual-color green/red luciferase reporter system (González-Grandío et al., 2021). This ratiometric

system uses green- and red-shifted luciferases (Sarrion-Perdigones et al., 2019), where green-

shifted luciferase is fused to a conditionally responsive promoter and red-shifted luciferase

Wilson et al., 2023

serves as a constitutively expressed reference for internal normalization, thus greatly reducing

intra experimental variation.

To test the activity of our thirteen candidate defense promoters, we generated promoter fusion

constructs driving the expression of green-emitting luciferase (E-Luc). These plasmids are

named *pDefense* (*pD*[*1-7, 10-12, 14-16*]), respectively. We co-transfected each of these plasmids

along with internal normalization plasmid *UBI-pRedf* and *UBI-AvrSr50* into protoplasts isolated

from cv. GaboSr50 and Gabo. Ten of these promoters did not result in specific and reproducible

promoter activation downstream of AvrSr50/Sr50 interactions when compared to their respective

controls. In contrast, the three promoters D2, D14, and D15 showed a strong increase in

normalized ratiometric luminescence specifically downstream of AvrSr50/Sr50 interactions. We

observed similar specific reporter activation for all three promoters downstream of *AvrSr35* in

wheat cv. C6969 (*Triticum aestivum*) carrying *Sr35* (Figure 3), which was introgressed from

Triticum monococcum (McIntosh et al., 1984). These results indicate that our new ratiometric

reporter system is able to detect specific defense activation downstream of two Avr/R

interactions when the *R* gene is expressed endogenously.

We next addressed whether the induction of D2, D14, and D15 promoters is specific to

AvrSr50/Sr50 and AvrSr35/Sr35 interactions. We extended our analysis to the recently published

AvrSr27/Sr27 (Upadhyaya et al., 2021) gene combination to test whether the *pD2, pD14* and

pD15 can serve as reporters of *Avr/R*-mediated response more broadly. We used protoplasts

isolated from cv. Coorong, which carries *Sr27* (Upadhyaya et al., 2021), and transfected them

with a plasmid that expresses *AvrSr27* under the control of the *UBI* promoter. We compared the

induction of all three defense reporters in cv. Coorong protoplasts transfected with *AvrSr50* and

cv. Fielder protoplasts, which does not carry *Sr27,* transfected with *UBI-AvrSr27*. We included

UBI-pRedf as internal normalization control. We observed that all *pD2, pD14,* and *pD15* have

high levels of normalized luminescence in cv. Coorong expressing *AvrSr27* compared to the

controls (Figure 3). These results support D2, D14, and D15 as commonly applicable defense

reporters in wheat independent of the wheat cultivar and the specific *Avr/R* combination used. In

addition, this suggests that our protoplast assay is applicable to testing wheat cultivars that carry

endogenous *R* genes and does not require overexpression of a cloned *R* gene.

The new positive defense reporters are broadly applicable for Avr/R interaction screening in many wheat cultivars

We aimed to illustrate the broad applicability of our new assay in a wide range of wheat cultivars as current cell-death based assays were mostly conducted in cv. Fielder (Saur et al., 2019, 5; Ortiz et al., 2022; Upadhyaya et al., 2021; Luo et al., 2021; Arndell et al., 2023) . We tested the defense signaling reporter D14 for its inducibility by three *Avr/R* combinations in protoplasts of nine different wheat cultivars. We tested cultivars that carry *R* genes (cv. GaboSr50, cv. Coorong and cv. C69.69) with their corresponding *Avr* construct *(AvrSr50, AvrSr27* and *AvrSr35, respectively).* The cultivars Clement, Compair, HeinesPeko, (Dracatos et al., 2016) and Kenya W1483 (McIntosh et al., 1995) were tested with overexpression of *AvrSr50/Sr50* combination. The cultivars Avocet (*Yr8)* and Fielder were tested with *AvrSr27/Sr27* and *AvrSr35/Sr35* combinations, respectively. We include a negative control in all cultivars by expressing the *R* gene together with an un-recognized *Avr* (Figure 4). In all cultivars we tested, a consistent robust induction of normalized luminescence of *pD14* was observed in protoplasts expressing cognate *Avr/R*-pair compared to the negative control. The relative normalized luminescence and its induction in the *Avr/R* treatment group varied between the different cultivars. These results indicate that the new wheat protoplast assay enables the detection of Avr/R interactions in all wheat cultivars tested.

The positive defense reporter assay enables multiplexed screening of *Avr* candidates in bulk

To explore multiplexed screening of *Avr* candidates using our protoplast assay, we transfected a pool of ten *Avr* candidates, including *AvrSr50*, *AvrSr27*, *AvrSr35*, and seven other wheat rust candidate effector genes used in our studies, into protoplasts isolated from cv. GaboSr50 and cv. Gabo. The same *Avr* pool excluding the *AvrSr50* was used as a negative control, and *AvrSr50* alone was used as a positive control. Each replicate included *pD14* reporter and *UBI-pRedf* co-transfection to detect *Avr/R*-triggered defense promoter induction and normalized luminescence measurements. In cv. GaboSr50 protoplasts, both *AvrSr50* alone and the pooled mix containing *AvrSr50* induced a high level of normalized luminescence of p*D14* when compared to the pooled mix without *AvrSr50* and all other treatment groups including controls in cv. Gabo protoplasts (Figure 5A). We observed a clear specific induction of the D14 promoter only in the presence of *AvrSr50* in cv. GaboSr50 when provided as pairs or as multiplexed mixture with no statistical

difference between the two treatment groups (Figure 5A). In contrast, there was a clear Avr-
unspecific reduction in red luminescence induced by both multiplexed mixtures in both cv. Gabo
and cv. GaboSr50 protoplast (Figure 5B). This indicates that the ratiometric dual-color
luciferase system combined with *pDefense* reporter is highly sensitive in detecting Avr/R
interactions and more specific than general luciferase based cell death based assays. Overall,
these results showed that our positive defense reporter protoplast assay can be used to screen a
pool of *Avr* candidates against a wheat cultivar endogenously expressing the recognizing *R* gene.

Discussion

Here, we describe an improved wheat protoplast assay to screen avirulence effector candidates
and study cellular defense signaling in wheat. We identify Avr/R-induced promoters and use
them to generate reporter constructs that produce a positive readout during Avr/R interaction. In
addition, we also reduce variability of luminescence measurement by adapting a ratiometric
dual-color luciferase reporter system (Sarrion-Perdigones et al., 2019). We show the broad
applicability of the positive defense reporters by testing three *Avr/R* pairs from the wheat stem
rust pathosystem on nine different wheat cultivars. The improved wheat protoplast assay is
highly sensitive because it allows for multiplexed screening of *Avr* candidates in wheat cultivars
expressing *R* genes endogenously.

This approach has several advantages over previous methods applied to identify Avr/R
interactions. Firstly, it does not require a cloned and overexpressed *R* gene to test the interaction.
We demonstrate that the positive defense reporter assay is suitable to test interactions directly in
a broad range of wheat cultivars encoding specific *R* genes in their genome. This will open
opportunities to screen *Avr* candidates directly on wheat-resistant cultivars, which was
previously considered not amenable using the cell-death based wheat protoplast assays (Saur et
al., 2019). Secondly, combining (generic) *Avr/R*-inducible promoters with the dual-reporter
system allows for generating normalized positive signals, which enables a much more sensitive
screening approach by greatly reducing variability between replicates. Thirdly, this approach
unlocks screening multiple *Avr* candidates in a single transfection, which saves time and
resources. Lastly, our new wheat protoplast reporter assay enables us to study a wide range of
plant defense signaling directly in wheat. This is highlighted by our RNAseq analysis of defense
signaling in wheat protoplasts. Our results show that genes responsible for calcium ion binding,

cation binding and calcium ion transmembrane transporter activity along with other defense-
related genes are specifically upregulated in response to both AvrSr35/Sr35 and AvrSr50/Sr50
interactions in two wheat cultivars. In line with our findings, there have been strong indications
that plant defense signaling triggered by Avr/R interactions involves plasma membrane
reorganization including formation of ion channels that regulate in plant cell death and disease
resistance. Recent studies of *Arabidopsis* ZAR1 (Bi et al., 2021) and wheat Sr35 (Förderer et al.,
2022) showed the activated ZAR1 and Sr35 proteins forming pentameric structures, so-called
resistosomes, that appear to function as non-selective calcium channels responsible for Ca^{2+}
uptake and leading to cell death. Similarly, in *Arabidopsis*, an activated NRG1.1 protein, known
as downstream signaling component of plant *R* proteins, compose a Ca^{2+} permeable non-selective
cation channel enabled by oligomerization, which control cell Ca^{2+} influx and induce plant cell
death (Jacob et al., 2021). Therefore, our findings support that the channel activity, especially
Ca^{2+} regulating channels, are conserved in broader Avr/R-induced downstream signaling.

Interestingly, one of our marker genes (the gene of promoter D14) encodes a predicted wheat
NDR1/HIN1-like 10 (NHL10) protein. The NHL proteins that are grouped in a large gene family
in *Arabidopsis thaliana* involved in plant defense, such as disease resistance (Aarts et al., 1998;
Century et al., 1995) and pathogen sensing (Boudsocq et al., 2010; Zheng et al., 2004; Zipfel et
al., 2004). The NHL proteins have one or two transmembrane domain(s) (Zheng et al., 2004) and
several of them were shown to localize to the plasma membrane (Selote et al., 2014; Coppinger
et al., 2004; Dagvadorj et al., 2022), which is thought to be crucial for their functioning in
disease resistance pathways. Moreover, *Arabidopsis NHL10* expression, which is highly induced
by flg22, is reported to be involved in Ca^{2+} signaling through the activation of calcium-
dependent protein kinases (Boudsocq et al., 2010). In a recent study, wheat blotch disease
effector ToxA was shown to interact with a wheat NHL protein, TaNHL10, to promote cell death
mediated by intracellular resistance like protein Tsn1 (Dagvadorj et al., 2022). Future studies
will benefit from our defense specific promoters, in particular D14, to investigate downstream
defense signaling components in wheat.

We acknowledge future improvement will further extend the applicability of our positive defense
reporter as described here. Similar to the previous wheat protoplast cell-death based assays (Saur
et al., 2019), the assay we developed uses a large amount of DNA material for transfection to

Wilson et al., 2023

obtain adequate luminescence signal induced by target Avr/R interactions. For example, in our

transfections, we typically use 10-15 µg of plasmid for each target, which requires a large

amount of pure plasmid DNA extraction. This could be time-consuming and expensive for

preparing a large list of *Avr* candidates for screening. Moreover, we observed a larger quantity of

DNA having a negative effect on the transfection efficiency when we transformed ten *Avr*

candidates in one transfection. Future experiments will address the sensitivity of our positive

defense reporters and the amount of plasmid required for robust responses and multiplexed

avirulence effector screening. It is likely that further optimization as reported recently (Arndell et

al., 2023) will allow us to reduce required plasmid amounts. Our study is complementary to a

recent report that describes highly multiplexed avirulence effector recognition screening using

cell death as a negative enrichment method. Multiplexed effector libraries (n>600) of *Pgt* were

co-transfected into cv. Fielder protoplasts overexpressing specific *Sr* genes from plasmids. The

author used the overexpression of YFP (Yellow Fluorescent Protein) as a cell viability marker to

separate live protoplast from dead protoplast via flow cytometry. This allowed for negative

enrichment analysis of expressed effectors when comparing live protoplast before and after flow

cytometry via RNAseq analysis. The author could identify new matching *Avr* genes for two out

of five *Sr* genes including *AvrSr13/Sr13* and *AvrSr22/Sr22* but not for *Sr21*, *Sr26*, and *Sr61*.

Such large scale avirulence effector screening approaches will be much more efficient when

combined with our positive reporter assays that will allow researchers to first identify avirulence

effector pools that are recognized by a specific wheat cultivar. In addition, further improvements

of the reported negative enrichment screen will likely enable screening in wheat cultivars

expressing *R* genes endogenously similar to our assay. Hence the combination of both

multiplexed avirulence effector screening approaches promises to usher in a new area of

avirulence gene identification of economically important pathogens of wheat and other crop

species.

Methods

Protoplast Isolation

Protoplast   isolation   and   plant   growth   protocol   are   described   here:

https://dx.doi.org/10.17504/protocols.io.q26g7r3zkvwz/v1. Wheat plants of cv. Fielder, Gabo

and GaboSr50 were grown for 7-9 days in a growth cabinet with 150 µmol/m^2/s light intensity

and at 21 ^{o}C with 16 hr day length. Protoplasts were isolated by making a shallow cut across the

leaf epidermis and peeling back the epidermal layer to expose the mesophyll layer beneath. The

cut segments were placed, peeled side down in 0.6 M Mannitol for 10 mins, then placed into a 7

cm diameter petri dish containing filtered enzyme solution (MES-pH5.7 (20 mM), Mannitol (0.6

M), KCI (10 mM), Cellulase (Yakult R-10) 1.5% w/v, Macerozyme (Yakult R-10) 0.75% w/v,

$CaCl_2$(10 mM). BSA 0.1% w/v). The dish was wrapped in aluminum foil to shield it from light

and placed on an orbital shaker, horizontally rotating at 60 RPM, for three hours. Protoplasts

were collected by gentle centrifugation at 100 x g for 3 mins in a 30 mL round-bottomed tube,

then the enzyme solution was removed and the pellet was washed in W5 solution (MES-pH5.7 (2

mM), KCI (5 mM), $CaCl_2$ (125 mM), NaCl (154 mM). 5 µL of concentrated protoplasts were

removed and used to quantify cells and check quality using a hemocytometer and light

microscope. Protoplasts were washed again with W5, this time left on ice for 45 minutes to

settle. W5 solution was removed, and protoplasts were resuspended to 300,000-350,000

protoplasts/mL using MMG (MES-pH5.7 (4 mM), Mannitol (0.4 mM), $MgCl_2$ (15 mM)).

Protoplasts were used immediately for transfection.

Protoplast transfection

High concentration (1 µg/µL) plasmid DNA was extracted via maxi-prep (Promega SV Wizard).

Lower quality plasmid DNA (e.g. from miniprep kits) typically resulted in low transfection rates.

The *Avr*, *R* and *luciferase* genes were expressed under control of maize ubiquitin promoter

(*UBI*), and 10 µg of each *Avr* and *R* plasmid and 20 µg of *luciferase* reporter plasmid were used

for each replicate. Due to high concentration and purity, plasmids were heated at 65^{o}C for 10 min

before use to reduce viscosity and ensure consistent measurement and delivery into cells. The

required plasmid DNA was added to 14 mL round-bottom culture tubes (Thermofisher,

#150268). 300 uL of protoplasts and 300 µL of PEG solution (PEG 4000 (40% w/v), Mannitol

(0.2 M), $CaCl_2$ (100 mM)) was then added and mixed immediately by gently flicking/rotating the

tube three times. Samples were incubated for 15 minutes, and the reaction halted with 900 µL

W5. Protoplasts were then collected by 100 x g centrifugation for 2 min, resuspended in 500 µL

modified W5 (W5 + 1 mM glucose) and transferred to a transparent, sterile 12 well culture plate

(Costar, #3513) whose wells were coated with 10% BSA to prevent sticking. Incubation was

carried out in light conditions, approximately 100 µmol/m^2/s. Wide bore or cut tips were used

throughout for handling protoplasts.

Protoplast Quantification

After the required incubation time (3, 4, 5, 6, and 18-hours for time course experiment, 4-18 hrs

for *pDefense-ELuc* reporters) protoplasts were prepared for luminescence measurement. 200 µL

of 1x cell lysis buffer (Promega #E3971) was added and samples vortexed and incubated for 15

min. Samples were centrifuged again to collect cell debris and 50 µL of lysate was added to an

opaque, flat-bottom, white, solid polystyrene 96 well plate (e.g. Corning® 96 well NBS™

Microplate #CLS3600). Plate was read for total luminescence (Tecan infinite 200Pro, 1000 ms

integration 0 ms settle) without addition of substrate, then 50 µL Steady-Glo (Promega #E2520)

Luciferin substrate was added, and the plate was read again.

Protoplast RNAseq sampling

Protoplasts were prepared as described above, using 315,00 protoplasts in 900 µL MMG per

replicate, with 4 replicates for each treatment group The *UBI-fLUC* reporter was excluded.

Additional replicates of each condition included the reporter to confirm cell death after 16hrs

(Data not shown). At 4 hrs post transfection the protoplasts were pelleted, snap-frozen, and

stored at -80 until RNA isolation. Samples from each cultivar were prepared and processed in

batches. Qiagen RNeasy Plant Mini Kit was used as directed, (#74904) omitting the first

mechanical cell lysis step as freezing and addition of RLT buffer was sufficient to lyse cells.

Qiagen RNase-Free DNase Set (#79254) was used as directed. RNA was checked for absence of

DNA contamination by PCR and gel electrophoresis and quantitated using Qubit and Nanodrop.

Total RNA Illumina sequencing was performed on the NovaSeq platform, in a 2x150bp paired-

end configuration, with poly A selection & strand specific library preparation for the 36 samples.

The resulting sequence data is deposited in the NCBI Sequence Read Archive (SRA) under

BioProject number PRJNA957082.

RNAseq Analysis

Transcript abundance was quantified with kallisto (Bray et al., 2016) using the Ensembl *Triticum*

aestivum transcriptome (IWGSC, INSDC Assembly GCA_900519105.1, Jul 2018) plus cDNA

sequences for *AvrSr50, Sr50*, *AvrSr35* and *Sr35* as reference. DESeq2 (Love et al., 2014) was

used to test for differential expression. We defined upregulation at a 4-fold change or greater.

GaboSr50 with AvrSr50 was compared against the four control treatment groups;

Gabo_AvrSr50, Gabo_EV, GaboSr50_AvrSr35 and GaboSr50_EV. Then, the intersect of the

upregulated genes was taken, for a total of 272 genes. For Fielder, Fielder_Sr35_AvrSr35 was

compared to the 3 control treatment groups; Fielder_AvrSr35, FielderSr35_AvrSr50,

Fielder_Sr35. Again, taking the intersect of the upregulated genes returned 98 upregulated genes.

66 genes were commonly upregulated between both lists (Zenodo Supplemental Data S5). For

GO terms analysis, we repeated the comparisons above with lower upregulation of 2-fold change

and used the gProfiler webtool and Ensembl *Triticum aestivum* annotation, running the g:GOst

program (Raudvere et al., 2019). The recommended g:SCS multiple testing correction algorithm

was used.

pDefense Promoter Selection

Expression in log fold change and transcripts per million (TPM) was visualized for the 66

commonly upregulated genes, present in both 2-fold upregulated lists used for GO analysis.

Thirteen genes were selected based on their overall expression and upregulation in *Avr/R*

treatment groups. The entire gene and region were compared in Geneious and 600-800 bp

upstream of the 13 genes was selected, including any annotated 5'UTR. These promoter

sequences then were synthesized with 5'and 3' 4bp MoClo promoter overhangs flanked by BsaI

recognition sites for Golden Gate cloning as promoter+5'UTR standard parts (Weber et al.,

2011). Any BpiI and BsaI enzyme recognition sites within the promoter sequence were removed

prior to synthesis. Promoters that could not be synthesized were cloned by PCR, using primers

(Zenodo Supplemental Data S6) with BpiI recognition sites and 4bp overhang to facilitate

assembly into universal level 0 vector *pAGM9121* (Addgene Plasmid #51833). Any BsaI

recognition sites were removed as per Grutzner & Marillionnet, (2020).

pDefense Plasmid Constructs

412   Plasmids were designed in Geneious. The plasmid constructs were designed according to the

413   MoClo syntax, assembled by Golden Gate cloning and verified by whole plasmid sequencing.

414   Supplemental Data (Zenodo Supplemental Data S6) and Plasmid maps Zenodo (DOI:

415   10.5281/zenodo.7844465, link: https://zenodo.org/record/7844465) are available, with all

416   generated plasmids deposited on Addgene.

Wilson et al., 2023

Ratiometric Assay

All isolation and transfection steps were as above aside from the substitution of the fLUC

reporter for the reporter (*UBI-pDefense)* and normalization (*UBI-pRedf*) plasmids. These require

the same substrate for activation, but express luciferase with different emission spectra, read by

specific green and red filters in the luminescence plate reader.A *UBI-ELuc* construct was also

generated to calibrate normalization and deconvolution calculations, necessary to resolve the

overlap in emission spectra of the two luciferase reporters (González-Grandío et al., 2021).

These calibration experiments were performed separately three times and with three replicates

per measurement, according to the instructions for Promega Chroma-Glo assay (Chroma-

Glo(TM) Luciferase Assay System Technical Manual, TM062). The plate reader was

programmed to measure total luciferase, then with green and red filters, with 1000 ms integration

time and 0 ms settle time, measured well-wise. A template spreadsheet for performing

calculations and a Python script for automating deconvolution and plotting are recorded on

protocols.io (https://dx.doi.org/10.17504/protocols.io.q26g7r3zkvwz/v1).

Cultivar testing

Protoplasts were isolated from nine different wheat cultivars (GaboSr50, Coorong, C6969,

Clement, Compair, HeinesPeko, K1483, Avocet and Fielder), using the method described above.

All were grown in the same conditions as above for 8 days. Where the cultivar carried an *R* gene

(GaboSr50, Coorong, C6969) the recognized *Avr* (*AvrSr50*, *AvrSr27* and *AvrSr35* respectively)

was transfected alone, with an avr (unrecognized by R protein) as control. In the other six

cultivars both the *Avr* and *R* gene were co-transfected. Normalized luminescence was calculated

from the ratio of deconvoluted *pDefense-Eluc* to *UBI-pRedf*.

Positive selection/pool screening

Protoplasts were prepared from GaboSr50 wheat leaves and transfected as above. Replicates

which contained the *Avr* pools used 10 µg of each plasmid mixed prior to transfection, while

replicates without pooled *Avrs* contained at total of 10 µg of *UBI-AvrSr50*. Measurements used

the ratiometric reporters and measurements, with *pDefense14-ELuc* as reporter.

Data availability

All supplemental data is available on Zenodo (DOI: 10.5281/zenodo.7880015, link: https://doi.org/10.5281/zenodo.7880015. All plasmids generated will be made available from Addgene, with plasmid maps deposited on Zenodo (DOI: 10.5281/zenodo.7844465, link: https://doi.org/10.5281/zenodo.7844465). Code used to perform RNAseq analysis is deposited on Github (https://github.com/ritatam/wheat-protoplast-RNAseq-DESeq2) and all RNAseq data is available on NCBI Sequence Read Archive (SRA) under BioProject number PRJNA957082.

Acknowledgements

We thank Prof. Robert Park for providing seeds of cv. C6969, A/Prof. Sambasivam Periyannan for providing seeds of cv. Fielder, cv. Gabo, cv. GaboSr50, and cv. Coorong, Dr. Michael Ayliffe and Dr. Peter Dodds for providing *pUBI-AvrSr35*, *pUBI-Sr35*, *pUBI-AvrSr50*, and *pUBI-Sr50*. B.S. acknowledges the financial support by the Australian Research Council via a Future Fellowship (FT180100024). This work was supported by computational resources provided by the Australian Government through the National Computational Infrastructure (NCI) under the ANU Merit Allocation Scheme.

Author Contributions

S. W., B. D., J. P. R., and B. S. conceived the project. S. W., B. D., R. T., L. M., N. H., and E. C. performed the experiments, S. W., B. D., and B. S. wrote the manuscript. R. T., E. C., and J. P. R. provided extensive feedback on the manuscript. B. S. secured funding. All authors read and approved the manuscript.

References

Aarts, N., Metz, M., Holub, E., Staskawicz, B.J., Daniels, M.J., and Parker, J.E. (1998). Different requirements for EDS1 and NDR1 by disease resistance genes define at least two R gene-mediated signaling pathways in Arabidopsis. *Proceedings of the National Academy of Sciences* **95**: 10306–10311.

Arndell, T. et al. (2023). Pooled effector library screening in protoplasts rapidly identifies novel
Avr genes.: 2023.04.28.538616.

Bi, G. et al. (2021). The ZAR1 resistosome is a calcium-permeable channel triggering plant
immune signaling. *Cell* **184**: 3528-3541.e12.

Boudsocq, M., Willmann, M.R., McCormack, M., Lee, H., Shan, L., He, P., Bush, J.,
Cheng, S.-H., and Sheen, J. (2010). Differential innate immune signalling via Ca2+
sensor protein kinases. *Nature* **464**: 418–422.

Bourras, S. et al. (2019). The AvrPm3-Pm3 effector-NLR interactions control both race-specific
resistance and host-specificity of cereal mildews on wheat. *Nat Commun* **10**: 2292.

Bouton, C., King, R.C., Chen, H., Azhakanandam, K., Bieri, S., Hammond-Kosack, K.E.,
and Kanyuka, K. (2018). Foxtail mosaic virus: A Viral Vector for Protein Expression in
Cereals. *Plant Physiology* **177**: 1352–1367.

Bray, N.L., Pimentel, H., Melsted, P., and Pachter, L. (2016). Near-optimal probabilistic
RNA-seq quantification. *Nat Biotechnol* **34**: 525–527.

Century, K.S., Holub, E.B., and Staskawicz, B.J. (1995). NDR1, a locus of Arabidopsis
thaliana that is required for disease resistance to both a bacterial and a fungal pathogen.
Proceedings of the National Academy of Sciences **92**: 6597–6601.

Chen, J. et al. (2017). Loss of *AvrSr50* by somatic exchange in stem rust leads to virulence for
Sr50 resistance in wheat. *Science* **358**: 1607–1610.

Christensen, A.H., Sharrock, R.A., and Quail, P.H. (1992). Maize polyubiquitin genes:
structure, thermal perturbation of expression and transcript splicing, and promoter
activity following transfer to protoplasts by electroporation. *Plant Mol Biol* **18**: 675–689.

Coppinger, P., Repetti, P.P., Day, B., Dahlbeck, D., Mehlert, A., and Staskawicz, B.J.
(2004). Overexpression of the plasma membrane-localized NDR1 protein results in
enhanced bacterial disease resistance in *Arabidopsis thaliana*. *The Plant Journal* **40**:
225–237.

Dagvadorj, B., Outram, M.A., Williams, S.J., and Solomon, P.S. (2022). The necrotrophic
effector ToxA from *Parastagonospora nodorum* interacts with wheat NHL proteins to
facilitate Tsn1-mediated necrosis. *The Plant Journal* **110**: 407–418.

Dodds, P.N. and Rathjen, J.P. (2010). Plant immunity: towards an integrated view of plant–
pathogen interactions. *Nat Rev Genet* **11**: 539–548.

Dracatos, P.M., Zhang, P., Park, R.F., McIntosh, R.A., and Wellings, C.R. (2016).
Complementary resistance genes in wheat selection "Avocet R" confer resistance to
stripe rust. *Theor Appl Genet* **129**: 65–76.

Förderer, A. et al. (2022). A wheat resistosome defines common principles of immune receptor
channels. *Nature* **610**: 532–539.

González-Grandío, E., Demirer, G.S., Ma, W., Brady, S., and Landry, M.P. (2021). A
Ratiometric Dual Color Luciferase Reporter for Fast Characterization of Transcriptional
Regulatory Elements in Plants. *ACS Synth. Biol*. **10**: 2763–2766.

Hewitt, T. et al. (2021). A highly differentiated region of wheat chromosome 7AL encodes a
Pm1a immune receptor that recognizes its corresponding AvrPm1a effector from
Blumeria graminis. *New Phytol* **229**: 2812–2826.

Jacob, P. et al. (2021). Plant "helper" immune receptors are Ca^{2+}-permeable nonselective
cation channels. *Science* **373**: 420–425.

Jensen, C. and Saunders, D.G.O. (2023). Magnaporthe oryzae pathotype Triticum (MoT) can
act as a heterologous expression system for fungal effectors with high transcript
abundance in wheat. *Sci Rep* **13**: 108.

Laflamme, B., Dillon, M.M., Martel, A., Almeida, R.N.D., Desveaux, D., and Guttman, D.S.
(2020). The pan-genome effector-triggered immunity landscape of a host-pathogen
interaction. *Science* **367**: 763–768.

Lee, W.-S., Hammond-Kosack, K.E., and Kanyuka, K. (2012). Barley Stripe Mosaic Virus-
Mediated Tools for Investigating Gene Function in Cereal Plants and Their Pathogens:
Virus-Induced Gene Silencing, Host-Mediated Gene Silencing, and Virus-Mediated
Overexpression of Heterologous Protein. *Plant Physiology* **160**: 582–590.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biology* **15**: 550.

Luo, M. et al. (2021). A five-transgene cassette confers broad-spectrum resistance to a fungal
rust pathogen in wheat. *Nat Biotechnol* **39**: 561–566.

Mago, R. et al. (2015). The wheat *Sr50* gene reveals rich diversity at a cereal disease resistance
locus. *Nat Plants* **1**: 15186.

McIntosh, R., Dyck, P., The, T., Cusick, J., and Milne, D. (1984). Cytogenetical studies in
wheat XIII. *Sr35* - a third gene from Triticum monococcum for resistance to Puccinia
graminis tritici. *Zeitschrift für Pflazenzüchtung* **92**: 1–14.

McIntosh, R., Wellings, C., and Park, R. (1995). Wheat Rusts: An Atlas of Resistance Genes.
In (CSIRO Publishing).

Meile, L., Croll, D., Brunner, P.C., Plissonneau, C., Hartmann, F.E., McDonald, B.A., and
Sánchez-Vallet, A. (2018). A fungal avirulence factor encoded in a highly plastic
genomic region triggers partial resistance to septoria tritici blotch. *New Phytologist* **219**:
1048–1061.

Müller, M.C. et al. (2022). Ancient variation of the *AvrPm17* gene in powdery mildew limits
the effectiveness of the introgressed rye *Pm17* resistance gene in wheat. *Proceedings of*
the National Academy of Sciences **119**: e2108808119.

Ortiz, D., Chen, J., Outram, M.A., Saur, I.M.L., Upadhyaya, N.M., Mago, R., Ericsson,
D.J., Cesari, S., Chen, C., Williams, S.J., and Dodds, P.N. (2022). The stem rust
effector protein AvrSr50 escapes Sr50 recognition by a substitution in a single surface-
exposed residue. *New Phytol* **234**: 592–606.

Praz, C.R. et al. (2017). *AvrPm2* encodes an RNase-like avirulence effector which is conserved
in the two different specialized forms of wheat and rye powdery mildew fungus. *New*
Phytol **213**: 1301–1314.

Raudvere, U., Kolberg, L., Kuzmin, I., Arak, T., Adler, P., Peterson, H., and Vilo, J. (2019).
g:Profiler: a web server for functional enrichment analysis and conversions of gene lists
(2019 update). *Nucleic Acids Research* **47**: W191–W198.

Saintenac, C., Zhang, W., Salcedo, A., Rouse, M.N., Trick, H.N., Akhunov, E., and
Dubcovsky, J. (2013). Identification of wheat gene *Sr35* that confers resistance to Ug99
stem rust race group. *Science* **341**: 783–786.

Salcedo, A. et al. (2017). Variation in the *AvrSr35* gene determines *Sr35* resistance against
wheat stem rust race Ug99. *Science* **358**: 1604–1606.

Sarrion-Perdigones, A., Chang, L., Gonzalez, Y., Gallego-Flores, T., Young, D.W., and
Venken, K.J.T. (2019). Examining multiple cellular pathways at once using multiplex
hextuple luciferase assaying. *Nat Commun* **10**: 5710.

Saur, I.M.L., Bauer, S., Lu, X., and Schulze-Lefert, P. (2019). A cell death assay in barley
and wheat protoplasts for identification and validation of matching pathogen AVR
effector and plant NLR immune receptors. *Plant Methods* **15**: 118.

Savary, S., Willocquet, L., Pethybridge, S.J., Esker, P., McRoberts, N., and Nelson, A.
(2019). The global burden of pathogens and pests on major food crops. *Nat Ecol Evol* **3**:
430–439.

Selote, D., Shine, M.B., Robin, G.P., and Kachroo, A. (2014). Soybean NDR1-like proteins
bind pathogen effectors and regulate resistance signaling. *New Phytologist* **202**: 485–498.

Upadhyaya, N.M. et al. (2021). Genomics accelerated isolation of a new stem rust avirulence
gene–wheat resistance gene pair. *Nat. Plants* **7**: 1220–1228.

Weber, E., Gruetzner, R., Werner, S., Engler, C., and Marillonnet, S. (2011). Assembly of
Designer TAL Effectors by Golden Gate Cloning. *PLOS ONE* **6**: e19722.

Wulff, B.B. and Krattinger, S.G. (2022). The long road to engineering durable disease
resistance in wheat. *Current Opinion in Biotechnology* **73**: 270–275.

Zheng, M.S., Takahashi, H., Miyazaki, A., Hamamoto, H., Shah, J., Yamaguchi, I., and
Kusano, T. (2004). Up-regulation of Arabidopsis thaliana *NHL10* in the hypersensitive
response to *Cucumber mosaic virus* infection and in senescing leaves is controlled by
signalling pathways that differ in salicylate involvement. *Planta* **218**: 740–750.

Zhong, Z. et al. (2017). A small secreted protein in *Zymoseptoria tritici* is responsible for
avirulence on wheat cultivars carrying the *Stb6* resistance gene. *New Phytologist* **214**:
619–631.

Zipfel, C., Robatzek, S., Navarro, L., Oakeley, E.J., Jones, J.D.G., Felix, G., and Boller, T.
(2004). Bacterial disease resistance in *Arabidopsis* through flagellin perception. *Nature*
428: 764–767.

Figure Legends

Figure 1: Cell death and differential gene expression in *Avr/R* interactions. **A-B**, Luciferase activity

was measured at different hours of post-transfection (hpt) in **(A)** cv. Fielder protoplasts

expressing *luciferase* alone, and *luciferase* with either *Sr35, AvrSr35,* or *Sr35* plus *AvrSr35* and

in **(B)** cv. GaboSr50 protoplasts expressing *luciferase* alone, and *luciferase* with either *AvrSr35*

or *AvrSr50.* **(C)** Shows the luminescence of the Avr/R interaction treatment groups from cv.

Fielder and cv. GaboSr50 alone when compared to the no plasmid transfection control replicates

(red line) which is the average of nine replicate across the six timepoints.

Figure 2: Co-expression of *Avr/R* gene pairs triggers defense signaling in wheat protoplasts.

(A) Heat map shows expression of *Avr* and *R* genes by normalized gene count. Each column

represents a replicate and the treatment group is indicated by a number as indicated in the legend.

R or *Avr* genes are expressed in their respective treatment groups only. This includes

endogenously expressed *Sr50* in cv. GaboSr50. **(B-D)** shows principal component analysis

(PCA) of the RNAseq analysis; all 36 samples **(B)**, cv. Fielder **(C)** and cv. Gabo/GaboSr50 **(D)**.

(E) GO terms of genes that are upregulated 2 fold in *Avr/R* treatment groups. A full list of

upregulated genes with overrepresented GO terms is recorded in supplemental data (Zenodo

Supplemental Data S4). **(F)** shows a heatmap of expression of thirteen candidate reporter genes

(D1-7, D10-12, D14-16) that are commonly upregulated by *Sr50/AvrSr50* and *Sr35/AvrSr35* in

cv. GaboSr50 or cv. Fielder, respectively. Each square represents the mean of four replicates for

treatment group.

Figure 3: Positive pDefense reporters detect defense signaling triggered by three distinct

***Avr/R* pairs in wheat protoplasts expressing the *R* gene endogenously.** Each panel of each

row shows the normalized luminescence of the positive defense reporter pDefense14 **(A)**,

pDefense15 **(B)**, or pDefense2 **(C)**. From left to right, protoplasts from wheat cultivars C6969

carrying *Sr35*, GaboSr50 carrying *Sr50*, and Coorong carrying *Sr27* were transfected with

AvrSr35, *AvrSr50* and *AvrSr27* constructs, respectively, and each wheat cv. was transfected with

avr (unrecognized *Avr*) construct and empty vector (EV) as indicated in the column headers and

x-axis labels. *UBI-pRedf* was used for normalization of all experiments.

Figure 4: The positive defense reporter *pDefence14* is broadly applicable to all tested wheat

cultivars. Each panel of each row shows the normalized luminescence of the positive defense

reporter *pDefense14*. Row **(A)**: Protoplast isolated from three wheat cultivars carrying a

described *R* gene, were transfected with its cognate *Avr* or controls. Rows **(B-C)**: Protoplasts

isolated from various wheat cultivars were transfected with both *R* and *Avr* constructs. The

cultivars Clement, Compair, HeinesPeko, and Kenya W1483 were tested with *AvrSr50/Sr50*.

Avocet and Fielder were tested with *AvrSr50/Sr50* and *AvrSr27/Sr27*, respectively. An *avr*

(unrecognised by R protein) was used as a control and *UBI-pRedf* was used for normalization of

all experiments.

Figure 5: The positive defense reporter enables multiplexed *Avr* screening. Each panel in **(A)**

shows the normalized luminescence of the positive defense reporter *pDefense14*. Each panel in

(B) shows the red luminescence of *UBI-pRedf.* A pool of ten *Avr* candidates, with and without

AvrSr50 or *AvrSr50* alone was transfected into protoplasts isolated from cultivars GaboSr50 (left

column) and Gabo (right column). One-way analysis of variance (ANOVA) and post hoc

Tukey's HSD test was used to assess differences among normalized luminescence or red

luminescence of treatment groups. Treatments marked with differing lowercase letters show

statistically significant difference (P <0.05).

Supplemental Material Legends

Supplemental Material S1: Luminescence data for Figure 1, protoplast time course experiment.

Supplemental Material S2: List of samples and treatment groups for RNAseq experiment.

Supplemental Material S3: Lists of Gabo/GaboSr50, Fielder and all cultivar genes upregulated

by 2-fold.

Supplemental Material S4: GO terms analysis Query tables and g:profiler results tables

Supplemental Material S5: Lists of Gabo/GaboSr50, Fielder and all cultivar genes upregulated

by 4-fold.

Supplemental Material S6: List of primers used.

Supplemental Material S7: List of plasmids used.

Supplemental Material S8: Luminescence data used for generating Figures 3, 4 and 5.