Title: Multiplexed effector screening for recognition by endogenous resistance genes using

positive defense reporters in wheat protoplasts

Salome Wilson*, Bayantes Dagvadorj*, Rita Tam, Lydia Murphy, Sven Schulz-Kroenert, Nigel

Heng, Emma Crean[$], Julian Greenwood, John P. Rathjen, and Benjamin Schwessinger

Research School of Biology, The Australian National University, Canberra, ACT, 2601,

Australia

*These authors contributed equally to this work

[$]Current address: Technical University of Munich, TUM Campus Straubing for Biotechnology

and Sustainability, Schulgasse 16, 94315 Straubing, Germany

Corresponding author: Benjamin Schwessinger

Email: benjamin.schwessinger@anu.edu.au

Short Title:

Defense signaling in wheat protoplasts

Summary

• Plant resistance (*R*) and pathogen avirulence (*Avr*) gene interactions play a vital role in

pathogen resistance. Efficient molecular screening tools for crops lack far behind their

model organism counterparts, yet they are essential to rapidly identify agriculturally

important molecular interactions that trigger host resistance.

• Here, we have developed a novel wheat protoplast assay that enables efficient screening of

Avr/R interactions at scale. Our assay allows access to the extensive gene pool of

phenotypically described *R* genes because it does not require the overexpression of cloned *R*

genes. It is suitable for multiplexed *Avr* screening, with interactions tested in pools of up to

fifty *Avr* candidates.

• We identified Avr/R-induced defense genes to create promoter-luciferase reporter. Then, we

combined this with a dual-color ratiometric reporter system that normalizes read-outs

accounting for experimental variability and Avr/R-induced cell-death. Moreover, we

introduced a self-replicative plasmid reducing the amount of plasmid used in the assay.

• Our assay increases the throughput of *Avr* candidate screening, accelerating the study of

cellular defense signaling and resistance gene identification in wheat. We anticipate that our

assay will significantly accelerate *Avr* identification for many wheat pathogens, leading to

improved genome-guided pathogen surveillance and breeding of disease-resistant crops.

Introduction

Plant pathogenic fungi cause severe damage to cereal crops. These pathogens secrete hundreds of

proteinaceous molecules called effectors during infection which manipulate plant processes to

facilitate disease (Dodds & Rathjen, 2010). Some effectors can be recognized by resistance (R)

proteins and are therefore also called avirulence (Avr) proteins. Avr recognition triggers a suite

of defense-related host responses to fight off pathogen infection including the hypersensitive

response (HR) that leads to cell-death at the site of infection and halts pathogen growth

(Laflamme *et al.*, 2020). The deployment of suitable *R* genes is a major breeding objective to

reduce the negative impact of pathogens on crop production including in wheat. Recent years

have made significant progress in identifying and cloning a wide variety of *R* genes in wheat

(Wulff & Krattinger, 2022). Yet the identification of the corresponding *Avr* genes is lacking far

behind. For example, there are only four confirmed *Blumeria graminis* s. str. (formerly known as

Blumeria graminis f. sp. *tritici*) *Avr* genes (Praz *et al.*, 2017; Bourras *et al.*, 2019; Hewitt *et al.*,

2021; Müller *et al.*, 2022), three for *Zymoseptoria tritici* (Zhong *et al.*, 2017; Meile *et al.*, 2018;

Amezrou *et al.*, 2023), and five for *Puccinia graminis* f. sp. *tritici* (*Pgt*) (Chen *et al.*, 2017;

Salcedo *et al.*, 2017; Upadhyaya *et al.*, 2021; Arndell *et al.*, 2023). There are no confirmed *Avr*

genes for *Puccinia triticina* and *Puccunia striiformis* f. sp. *tritici* despite both belonging to the

top five wheat pathogens globally (Savary *et al.*, 2019). This lack of confirmed *Avr* genes in

wheat pathogens and especially rust fungi is confounded by the absence of an efficient screening

tool for *Avr* recognition in wheat. Most current methods require cloned *Avr* and *R* gene
candidates that retain function when expressed heterologously and exhibit HR as general read-
out for Avr/R interactions. For example, the model plant *Nicotiana benthamiana* is used to
validate Avr/R interactions as it produces visible HR upon transient expression of *Avr/R* genes
through agroinfiltration. Virus-mediated overexpression (VOX) can be used to screen *Avr*
candidates in wheat cultivars carrying the corresponding *R* gene as the Avr recognition will lead
to visible HR symptoms on the infected leaves (Lee *et al.*, 2012; Bouton *et al.*, 2018). However,
the method is only applicable to small proteins (Lee *et al.*, 2012) and used viruses are not
accessible globally due to biosecurity regulations (Bouton *et al.*, 2018). Recently, HR-based
assays in wheat protoplasts have emerged as a potential alternative to test Avr/R interactions
(Saur *et al.*, 2019). In this transient assay, wheat protoplasts are co-transfected with *Avr/R* genes
along with a *luciferase* reporter gene. Avr/R interaction triggers cell-death causing significant
decreases in luciferase enzyme activity. This assay has been used to validate *Avr* candidates from
wheat rust pathogens (Saur *et al.*, 2019; Luo *et al.*, 2021; Upadhyaya *et al.*, 2021; Ortiz *et al.*,
2022). However, the HR-based wheat protoplast assay in its current form has several limitations
that make it impractical for high-throughput *Avr/R* screening. One such limitation is that it
requires the transient overexpression of both the *Avr* and *R* genes from plasmids (Saur *et al.*,
2019). This prevents screening against the large number of *R* genes which have been
phenotypically characterized but not yet been cloned (McIntosh *et al.*, 1995). Also, it is well
known that luminescence is highly variable between biological replicates and experiments (Saur
et al., 2019; González-Grandío *et al.*, 2021) which can make the results difficult to interpret for
high-throughput screening purposes. The assay in its current form relies on the reduction of
luciferase activity as read out which is unspecific and cannot be differentiated from cell-death
that is unrelated to HR. The assay in its current form also requires the preparation of large
quantities of target gene plasmids for the protoplast transfection to achieve observable Avr/R-
triggered cell-death (Saur *et al.*, 2019; Upadhyaya *et al.*, 2021; Ortiz *et al.*, 2022). These
limitations encouraged us to develop a novel assay in wheat protoplast that relies on a positive
defense specific read-out and accounts for experimental variability via normalization, which is
common practice for protoplast report assays (González-Grandío *et al.*, 2021). Our new assay
allows the research community to access the whole wheat gene pool to screen for Avr/R
interactions and enables the investigation of downstream defense signaling directly in wheat.

Results

Profiling temporal HR onset induced by Avr/R interactions in wheat protoplasts

Our end goal was to identify defense signaling responsive promoter elements downstream of
Avr/R recognition in wheat protoplast before or independent of the onset of HR. Hence, we first
sought to characterize the time of HR onset in wheat protoplasts following Avr/R recognition to
identify a suitable time point for RNAseq expression analyses. We used two published *Avr/R*
gene pairs, *AvrSr35/Sr35* (Saintenac *et al.*, 2013; Salcedo *et al.*, 2017) and *AvrSr50/Sr50* (Mago
et al., 2015; Chen *et al.*, 2017), from wheat stem rust *P. graminis* f. sp. *tritici* for initial method
development. As previously described (Saur *et al.*, 2019), we used luminescence as a proxy for
cell viability after overexpression of the *luciferase* reporter gene in conjunction with each *Avr/R*
combination and corresponding control. When expressed from a plasmid, *Avr* and *R* genes were
under control of maize ubiquitin promoter (*UBI*) (Christensen *et al.*, 1992). We used the wheat
cultivar (cv.) Fielder for the analysis of *AvrSr35/Sr35* and expressed both genes from a plasmid.
For *AvrSr50/Sr50*, we made use of the introgression of *Sr50* into cv. Gabo (GaboSr50) and
generated protoplast from cv. Gabo and cv. GaboSr50 (Jensen & Saunders, 2023). We only
expressed *AvrSr50* from a plasmid. We also included the controls, the reporter alone and the
reporter with *Avr* or *R* alone (Zenodo Supplemental Data S1). We measured luminescence at 3,
4, 5, 6, and 18 h post-transfection (hpt). We observed a gradual increase in luminescence
measurements in protoplast expressing luciferase alone or when expressed together with an *Avr*
or *R* gene only (Figure 1, A and B). The luciferase activity was strongest at 18 hpt in all
treatment groups except for the ones expressing *Avr/R* pairs. In cv. Fielder expressing
AvrSr35/*Sr35* and in cv. GaboSr50 expressing *AvrSr50*, luminescence measurements only
increased slightly early on during the time course when compared to mock treatments of
protoplasts transfected with no DNA (red line, Figure 1 C). At about 4 to 5 hpt we observed a
divergence of luciferase activity from control treatment groups when compared with the Avr/R
interaction treatment groups (Figure 1A and B). The lack of continuous increase of luciferase
activity specifically in protoplast expressing *Avr*/*R* pairs suggests that their co-expression leads
to cell-death in wheat protoplasts soon after transfection.

Transfected protoplasts are suitable for replicable transcriptional analysis

We concluded that 4 hpt is a suitable time point to capture *Avr/R*-dependent transcriptional
changes before strong HR induction based on our time course analysis (Figure 1). We repeated
the experimental setup while omitting the reporter plasmid and collected protoplasts at 4 hpt for
RNA sequencing (Figure 2 A, Zenodo Supplemental Data S2). We included four biological
replicates per treatment group and sequenced each sample on the Illumina platform generating
>20 million 150 bp paired-end reads per replicate. We used kallisto (Bray *et al.*, 2016) for
transcript quantification and DESeq2 (Love *et al.*, 2014) for downstream differential expression
analysis. As a first quality control step we quantified the expression of all *Avr* and *R* genes to
investigate their specific expression pattern at 4 hpt. This analysis clearly showed that all genes
are correctly expressed and at similar levels, including in treatment groups that co-express *Avr/R*
gene pairs (Figure 2 A). In addition, it revealed that the endogenous expression of *Sr50* in cv.
GaboSr50 is much lower level than the overexpressed of *Sr35* in cv. Fielder from a plasmid
under the *UBI* promoter. Next, we performed principal component analysis (PCA) of all sample
replicates combined. This revealed that the factor "wheat cultivar" had the strongest impact on
the observed variation in transcript abundance and accounted for close to 90% of all variation on
the first principal component (PC1), separating the two Gabo cultivars from cv. Fielder and
further separating cv. Gabo from cv. GaboSr50 along principal component two (PC2) (Figure 2
B). Consequently, we analyzed each cultivar separately. The wheat cultivar specific PCA
analysis showed a clear impact of Avr/R interaction on transcript abundance variation for both
AvrSr35/Sr35 and *AvrSr50/Sr50* (Figure 2 C and D). This impact was replicable because all four
biological replicates clustered closely together and clearly separated treatment groups including
Avr/R interactions sample types from their controls. This is consistent with the cell-death results
seen in Figure 1 A and B. These results suggest that these data could be used to identify reporter
genes specifically expressed downstream of Avr/R interactions in wheat protoplasts.

***Avr/R* recognition in wheat protoplasts leads to defense signaling**

Next, we investigated if the observed gene expression changes in *Avr/R* expressing protoplasts
overlapped with gene ontology (GO) terms commonly associated with defense signaling
pathways in wheat and other plant species. We identified all genes that are specifically
upregulated in response to Avr/R interactions in cv. Gabo/Gabo50 and cv. Fielder above a 2-fold

change threshold. We identified 309 and 922 genes in *AvrSr35/Sr35* and *AvrSr50/Sr50*

expressing protoplasts, respectively, when compared to all their control treatment groups.

Complete gene lists are available on Zenodo as Supplemental Data S3. We performed a

statistical enrichment analysis using these gene lists to test for overrepresentation of defense

signaling associated GO terms. Our results revealed significant enrichment for GO terms related

to cellular components of cell periphery (GO:0071944) and plasma membrane (GO:0005886) in

both cultivars (Figure 2 E). We identified defense response to other organisms (GO:0098542),

response to other organisms (GO:0051707), and biological process involved in interspecies

interaction between organisms (GO:0044419) when looking at cv. GaboSr50 alone (Zenodo

Supplemental Data S4). We found molecular functions GO terms being enriched for calcium ion

binding (GO:0005509), cation binding (GO:0043169), and calcium ion transmembrane

transporter activity (GO:0015085) in both cultivars (Figure 2 E). Our GO terms analysis suggests

that *Avr/R* expressing protoplasts undergo significant transcriptional reprogramming consistent

with calcium-mediated defense signaling before the onset of strong cell-death.

We were then interested in genes that are commonly upregulated by the Avr/R interaction in

both cv. GaboSr50 and cv. Fielder. We identified 66 genes commonly upregulated in both

cultivars, (Zenodo Supplemental Data S5) of which only 16 genes were associated with three

different GO terms, all related to the molecular function of ion binding (Zenodo Supplemental

Data S4). Our results showed that several of the identified genes have been previously implicated

in plant defense signaling including *TaNHL10* (*TRAESCS3D02G368800*) (Century *et al.*, 1995;

Aarts *et al.*, 1998; Dagvadorj *et al.*, 2022).

In summary, we observed the upregulation of genes involved in defense signaling in wheat

protoplasts expressing *Avr/R* pairs. The promoters of commonly upregulated genes, by two *Avr/R*

pairs and in two wheat cultivars (Figure 2F), are promising candidates to generate Avr/R-

inducible reporters.

Identification of positive defense reporters in wheat protoplasts

We further investigated the expression patterns of the 66 commonly upregulated genes to select a

subset whose promoters we could test as positive defense signaling reporters. We selected 13

promoters (D1-D7, D10-D12, and D14-D16) based on their genes' expression patterns in both

cultivars (Figure 2 F). To further evaluate these candidates, we cloned or synthesized their

promoters, consisting of 600-800 bp upstream of their respective genes' start codons. We

accounted for the impact of Avr/R-induced HR upon luciferase activity (Figure 1) by adapting a

dual-color green/red luciferase reporter system (González-Grandío *et al.*, 2021). This ratiometric

system uses green- and red-shifted luciferases (Sarrion-Perdigones *et al.*, 2019), where green-

shifted luciferase is fused to a conditionally responsive promoter and red-shifted luciferase

serves as a constitutively expressed reference for internal normalization, thus greatly reducing

intra experimental variation.

To test the activity of our thirteen candidate defense promoters, we generated promoter fusion

constructs driving the expression of green-emitting luciferase (E-Luc). These plasmids are

named *pDefense* (*pD[1-7, 10-12, 14-16]*), respectively. We co-transfected these plasmids along

with internal normalization plasmid *UBI-pRedf* and *UBI-AvrSr50* (Supplemental Figure 1) into

protoplasts isolated from cv. GaboSr50 and Gabo. For the initial activity screening, we

performed the test with two technical replicates. Ten of these promoters did not result in specific

and reproducible promoter activation downstream of AvrSr50/Sr50 interactions when compared

to their respective controls. In contrast, the three promoters D2, D14, and D15 showed a strong

increase in normalized ratiometric luminescence specifically downstream of AvrSr50/Sr50

interactions (Supplemental Figure 2). Figure 3 shows an example of the specific activation of

D14 downstream of AvrSr50/Sr50 in cv. GaboSr50 when compared to controls not expressing

the *R* gene or the matching *Avr*. These results indicate that our new ratiometric reporter system is

able to detect specific defense activation downstream the Avr/R interactions when the *R* gene is

expressed endogenously.

The wheat dwarf virus-derived replicons-based system further increases the sensitivity of

the positive defense reporter assay

The current wheat protoplast assay requires a large amount of DNA material for transfection to

obtain observable cell-death. This makes its application prohibitive for large scale screening of

many candidates or variants. We tested a replicon-based system using a disarmed version of the

wheat dwarf virus (WDV) in our expression vector backbone (Gil-Humanes *et al.*, 2017). We

reasoned that this would generate self-replicating plasmids in wheat protoplasts and thereby

reduce the amount of plasmids required to detect defense activation in our new ratiometric wheat

protoplast assay. We assembled Golden Gate compatible WDV vector from Gil-Humanes et al.

(2017) and denoted it as *pWDV1* (see Material and Methods). We expressed *AvrSr50* under

ubiquitin promoter using *pWDV1* (*pWDV1-AvrSr50*) and used it to generate dose response

curves using different amounts (0.1, 0.5, 1, 5 and 15 µg) of plasmid DNA for transfection. We

measured induction of D14 induction (normalized luminescence) and cell viability (red

luminescence) and compared it to the dose response of the standard non-viral plasmid

pICH4772-AvrSr50 (Figure 4). We observed specific D14 promoter induction in all samples of

cv. GaboSr50 protoplasts transfected with *pICH4772-AvrSr50* and *pWDV1-AvrSr50* when

compared to controls in cv. GaboSr50 transfected with *pWDV1-AvrSr35* (1 and 15 µg) and cv.

Gabo transfected with *pWDV1-AvrSr50* (1 µg) or *pWDV1-AvrSr35* (1 µg). We also observed that

the promoter induction increases relative to the amount of *pICH4772-AvrSr50* used. At the same

time, we were still able to observe induction of the defense reporter D14 even in the absence of

cell-death at 0.1, 0.5 and 1 µg plasmid used. Using *pWDV1-AvrSr50* further improved the

sensitivity of the assay as it increased defense reporter D14 activation at 0.1 and 0.5 µg plasmid

and decreased cell viability when compared to the same amounts of plasmid for *pICH4772-*

AvrSr50. Altogether, this result indicates that the ratiometric dual-color luciferase system

combined with *pDefense* reporter and *pWDV1* is highly sensitive and specific requiring much

less plasmid to detect Avr/R interactions than a single luciferase-based wheat protoplast assays.

Positive defense reporters are highly sensitive to detect intracellular Avr/R interactions in

wheat protoplast expressing *R* genes endogenously

Next, we explored if the defense reporters require overexpression of the *R* gene and if they are

applicable to additional Avr/R interactions. We tested the induction of D14 expressing *AvrSr27,*

AvrSr35, and *AvrSr50* in *pWDV1* using 1 µg plasmid, instead of the previously used 15 µg. We

transfected cv. Coorong, C6969, and GaboSr50, with *pWDV1* encoding the recognized *Avr*, an

unrecognized or non-matching *avr*, or the empty vector control (Figure 5). Wheat cv. Coorong

carries *Sr27* (Upadhyaya *et al.*, 2021), and cv. C6969 (*Triticum aestivum*) carries *Sr35* which

was introgressed from *Triticum monococcum* (McIntosh *et al.*, 1984). We observed a strong

induction of D14 specifically downstream of AvrSr27/Sr27, AvrSr35/Sr35, and AvrSr50/Sr50

interactions but not in any of the controls (Figure 5 A-C), validating the D14 defense promoter as

well as *pWDV1* applicability in wheat protoplast assay. We decided to focus on D14 and switch

all *Avr* expressions in *pWDV1* for the remainder of the study but to keep the reporter genes in the

non-viral vector. We further tested the D14 promoter with two newly reported stem rust Avr

proteins AvrSr13 and AvrSr22 (Arndell *et al.*, 2023) by transfecting them in wheat cv. Line S

and W3534 (Park, 2016), respectively (Figure 5 D and E). Similar to the other *Pgt* AvrSr/Sr

interactions, we observed strong specific induction of D14 from AvrSr13/Sr13 and AvrSr22/Sr22

combinations.

To assess the D14 promoter specificity to Avr/R interaction, we exploited the natural and

mutational variation of *AvrSr50* alleles that mediate the recognition by *Sr50* (Ortiz *et al.*, 2022).

We tested three AvrSr50 variants displaying varying interaction strength with Sr50; avrsr50-A1

which carries a single amino acid substitution at Q121K and is not recognized by Sr50, avrsr50-

B6 which is found in *Pgt* race QCMJC and is not recognized by Sr50 (Chen *et al.*, 2017) and

AvrSr50-C, which is found in *Pgt* race RKQQCs and is recognized by Sr50 (Chen *et al.*, 2017)

in cv. GaboSr50 (Supplemental Figure 3). The previously used *AvrSr50*, *AvrSr35* and empty

vector served as controls. Using our assay, we confirmed previous results that the specific single

amino acid substitutions in avrsr50-A1 and avrsr50-B6 were sufficient to abolish recognition by

Sr50, while changes to six different amino acids in AvrSr50-C only partially abrogated

recognition by Sr50 (Supplemental Figure 3). This demonstrates that the D14 promoter

successfully detects specific Avr/R interaction in allelic variants of Avr proteins such as

AvrSr50.

Next, to show the versatility of D14 promoter, we tested AvrPm2 from *Blumeria graminis* s. str.

(Praz *et al.*, 2017) (Figure 5F). We observed specific D14 induction compared to the controls

when we transfected *AvrPm2* into wheat cv. Federation4/Ulka, which carries *Pm2* (Sánchez-

Martín *et al.*, 2021).

These results support that the D14 promoter in combination with *pWDV1* is broadly applicable as

a defense reporter in wheat independent of the wheat cultivar and the specific *Avr/R* combination

used. This suggests that our protoplast assay is applicable to testing wheat cultivars that carry

endogenous *R* genes and does not require a cloned *R* gene.

Positive defense reporter D14 is able to detect extracellular AvrStb6/Stb6 interaction in

wheat protoplast

We were interested in the induction of D14 reporter by extracellular Avr/R interactions. For this

purpose, we tested AvrStb6 from *Zymoseptoria tritici* (Zhong *et al.*, 2017), which is recognized

by *Stb6* encoding a wall-associated receptor kinase-like protein (Saintenac *et al.*, 2018). We

decided to express AvrStb6 including its signal peptide (SP) allowing for AvrStb6 accumulation

outside the cells to allow for extracellular recognition of AvrStb6 by Stb6. We included the non-

secreted form of AvrStb6 without SP as negative control. We observed significant D14 induction

compared to the controls when we transfected secreted-AvrStb6 in protoplasts derived from

wheat cv. Chinese Spring which carries *Stb6* (Supplemental Figure 4) (Saintenac *et al.*, 2018).

The induction of D14 was specific to wheat lines carrying *Stb6* because we did not observe it in

wheat cv. Heines Kolben which lacks *Stb6* based on our PCR screening (Supplemental Figure 4).

This result indicates that our defense reporter assay is able to detect extracellular Avr/R

interactions and validates versatility of the assay among Avr proteins from different wheat

pathogens.

285   Positive defense reporters are broadly applicable for Avr/R interaction screening in many

286   wheat cultivars

We aimed to illustrate the broad applicability of our new assay in a wide range of wheat cultivars

as current cell-death based assays were mostly conducted in cv. Fielder (Saur *et al.*, 2019, p. 5;

Luo *et al.*, 2021; Upadhyaya *et al.*, 2021; Ortiz *et al.*, 2022; Arndell *et al.*, 2023). We tested the

defense signaling reporter D14 for its inducibility by three *Avr/R* combinations in protoplasts of

nine different wheat cultivars. We tested cultivars that carry *R* genes (cv. GaboSr50, cv. Coorong

and cv. C6969) with their corresponding *Avr* construct *(AvrSr50, AvrSr27* and *AvrSr35,*

respectively)*. The cultivars Clement, Compair, Avocet Yr8, (Dracatos *et al.*, 2016) and Kenya

W1483 (McIntosh *et al.*, 1995) were tested with overexpression of *AvrSr50/Sr50* combination.

The cultivars HeinesPeko (Dracatos *et al.*, 2016) and Fielder were tested with *AvrSr27/Sr27* and

AvrSr35/Sr35 combinations, respectively. We included a negative control in all cultivars by

expressing an unrecognized (non-matching to R protein) *avr, Avr* or *R* gene alone (Figure 6). In

all cultivars we tested, a consistent robust induction of normalized luminescence of *pD14* was

observed in protoplasts expressing cognate *Avr/R*-pair compared to the negative controls. The

relative normalized luminescence and its induction in the *Avr/R* treatment group varied between

the different cultivars. These results indicate that the positive defense reporter assay enables the

detection of Avr/R interactions in many wheat cultivars.

The positive defense reporter assay enables multiplexed screening of *Avr* candidates in bulk

We aimed to test if our assay could be used for multiplexed screening of Avr/R interactions

using AvrSr50/Sr50 as a proof of concept. We co-transfected a pool of twenty-four *Avr*

candidates (Methods for details) into protoplasts isolated from cv. GaboSr50 and cv. Gabo in the

presence or absence of *AvrsSr50*. The pools include 1 µg of each *pWDV1-Avr*. Each replicate

included *pD14* reporter and *UBI-pRedf* co-transfection to detect *Avr/R*-triggered defense

promoter induction and normalized luminescence measurements. In cv. GaboSr50 protoplasts,

AvrSr50 alone and the pooled mixes containing *AvrSr50* induced a high level of normalized

luminescence of *pD14* when compared to the pooled mix without *AvrSr50* and all other

treatment groups including controls in cv. Gabo protoplasts (Figure 7). We observed a clear

specific induction of the D14 promoter only in the presence of *AvrSr50* in cv. GaboSr50 when

provided as pairs or as multiplexed mixtures with no significant statistical difference between the

single or multiplexed treatment groups (Figure 7). Overall, these results show that our positive

defense reporter protoplast assay can be used to screen a pool of *Avr* candidates against a wheat

cultivar endogenously expressing the recognizing *R* gene, which then can be followed up to

identify the specific positive *Avr/R* interaction.

Discussion

Here, we describe an improved wheat protoplast assay to screen avirulence effector candidates

and study cellular defense signaling in wheat. We identify Avr/R-induced promoters and use

them to generate reporter constructs that produce a positive readout during Avr/R interaction. We

also reduce variability of luminescence measurement by adapting a ratiometric dual-color

luciferase reporter system (Sarrion-Perdigones *et al.*, 2019) and enable assays with candidate *Avr*

plasmids prepared from simple minipreps. We show the broad applicability of the positive

defense reporters by testing seven *Avr/R* pairs from three different wheat pathogens on thirteen

different wheat cultivars. The improved wheat protoplast assay is highly sensitive because it

allows for multiplexed screening of *Avr* candidates in wheat cultivars expressing *R* genes
endogenously.

This approach has several advantages over previous methods applied to identify Avr/R
interactions in wheat. First, it does not require a cloned and overexpressed *R* gene to test the
interaction. We demonstrate that the positive defense reporter assay is suitable to test interactions
directly in a broad range of wheat cultivars encoding specific *R* genes in their genome. This
creates opportunities to screen *Avr* candidates directly on resistant wheat cultivars, which has
been not easily possible with the cell-death based protoplast assays due to the inherent assay
variability without normalization (Saur *et al.*, 2019). Second, combining generic Avr/R-inducible
promoters with the dual-reporter system generates normalized positive signals, which enables a
much more sensitive screening approach by greatly reducing variability between replicates.
Third, the assay uses 10-20 times less plasmid per *Avr* and screens multiple candidates in a
single transfection, simplifying plasmid preparation and saving time and resources. Lastly, our
new wheat protoplast reporter assay enables us to study a wide range of plant defense signaling
directly in wheat. This is highlighted by our RNAseq analysis of defense signaling in wheat
protoplasts. Our results show that genes responsible for calcium ion binding, cation binding and
calcium ion transmembrane transporter activity along with other defense-related genes are
specifically upregulated in response to both AvrSr35/Sr35 and AvrSr50/Sr50 interactions in two
wheat cultivars. In line with our findings, there have been strong indications that plant defense
signaling triggered by Avr/R interactions involves plasma membrane reorganization including
formation of ion channels that regulate plant cell-death and disease resistance. Recent studies of
Arabidopsis ZAR1 (Bi *et al.*, 2021) and wheat Sr35 (Förderer *et al.*, 2022) showed the activated
ZAR1 and Sr35 proteins forming pentameric structures, so-called resistosomes, that appear to
function as non-selective calcium channels responsible for Ca^{2+} uptake, leading to cell-death.
Similarly, in *Arabidopsis*, an activated NRG1.1 protein, known as a downstream signaling
component of plant *R* proteins, composes a Ca^{2+} permeable non-selective cation channel enabled
by oligomerization, which controls cell Ca^{2+} influx and induces plant cell-death (Jacob *et al.*,
2021). Therefore, our findings support that the channel activity, especially Ca^{2+} regulating
channels, are conserved in broader Avr/R-induced downstream signaling.

Interestingly, one of our marker genes (the gene of promoter D14) encodes a predicted wheat

NDR1/HIN1-like 10 (NHL10) protein. The NHL proteins that are grouped in a large gene family

in *Arabidopsis* are involved in plant defense, such as disease resistance (Century *et al.*, 1995;

Aarts *et al.*, 1998) and pathogen sensing (Zheng *et al.*, 2004; Zipfel *et al.*, 2004; Boudsocq *et al.*,

2010). The NHL proteins have one or two transmembrane domain(s) (Zheng *et al.*, 2004) and

several of them were shown to localize to the plasma membrane (Coppinger *et al.*, 2004; Selote

et al., 2014; Dagvadorj *et al.*, 2022), which is thought to be crucial for their functioning in

disease resistance pathways. Moreover, *Arabidopsis NHL10* expression, which is highly induced

by flg22, is reported to be involved in Ca^{2+} signaling through the activation of calcium-

dependent protein kinases (Boudsocq *et al.*, 2010). In a recent study, wheat blotch disease

effector ToxA was shown to interact with a wheat NHL protein, TaNHL10, to promote cell-

death mediated by intracellular resistance like protein Tsn1 (Dagvadorj *et al.*, 2022). Future

studies will benefit from our defense specific promoters, in particular D14, to investigate

downstream defense signaling components in wheat.

Effectors are known to manipulate plant processes or suppress plant defense (Win *et al.*, 2012)

which could present challenges for screening large effector pools of unknown function. Some

effectors may interfere with the defense signaling process resulting in inhibition of the defense

promoter induction, which could prevent identifying a potential *Avr* candidate in the pool. This

problem can be solved by including a known *Avr* as positive control and transfecting the pool

into wheat cv. expressing cognate *R* gene of the known *Avr*. This positive control would reveal

whether the pool having *Avr* candidate(s) interferes with the defense promoter induction. In

addition, a split-pool approach that screens a limited number of subpools of *Avr* candidates with

partial overlaps can further reduce the risk of specific suppression of a recognized *Avr* by another

effector.

Future studies will further investigate the capabilities of our positive defense reporter based Avr

screening assay. We have performed screening of twenty-five different *Avr* candidates in this

study and we anticipate that the high sensitivity of the assay combined with self-replicative

pWDV1 plasmid can successfully screen far greater numbers of *Avr* candidates in a single wheat

protoplast transfection. Moreover, we anticipate that the *pWDV1* plasmid will simplify transient

gene overexpression in wheat protoplasts because it only requires miniprep plasmid samples

(Supplemental Figure 5), avoiding labor intensive and expensive maxiprep plasmid purification.

This will facilitate rapid functional gene and variant analysis directly in wheat as demonstrated

for *AvrSr50* variants in our study.

Our assay can be combined with a recent report that describes highly multiplexed avirulence

effector recognition screening using cell-death as a negative enrichment method (Arndell *et al.*,

2023). Multiplexed effector libraries (n>600) of *Pgt* were co-transfected into cv. Fielder

protoplasts overexpressing specific *Sr* genes from plasmids. The author used the overexpression

of YFP (Yellow Fluorescent Protein) as a cell viability marker to separate alive from dead

protoplasts via flow cytometry. This allowed for negative enrichment analysis of expressed

effectors when comparing live protoplasts before and after flow cytometry via RNAseq analysis.

The author could identify new matching *Avr* genes for two out of five *Sr* genes including

AvrSr13/Sr13 and *AvrSr22/Sr22* but not for *Sr21*, *Sr26*, and *Sr61*. Such large scale avirulence

effector screening approaches will be much more efficient when combined with our positive

reporter assays that will allow researchers to first identify avirulence effector pools that are

recognized by a specific wheat cultivar. Further refinement of the reported negative enrichment

screen will likely enable screening in wheat cultivars expressing *R* genes endogenously, similar

to our assay. Application of both multiplexed avirulence effector screening approaches promises

to usher in a transformative era of avirulence gene identification in economically important

pathogens of wheat and other crop species.

Methods

Protoplast Isolation

Protoplast isolation and plant growth protocol are described here:

https://dx.doi.org/10.17504/protocols.io.q26g7r3zkvwz/v1. Wheat plants of cv. Fielder, Gabo

and GaboSr50 were grown for 7-9 days in a growth cabinet with 150 µmol/m^2/s light intensity

and at 21 ^{o}C with 16 hr day length. Protoplasts were isolated by making a shallow cut across the

leaf epidermis and peeling back the epidermal layer to expose the mesophyll layer beneath. The

cut segments were placed, peeled side down in 0.6 M Mannitol for 10 mins, then placed into a 7

cm diameter petri dish containing filtered enzyme solution (MES-pH5.7 (20 mM), Mannitol (0.6

M), KCl (10 mM), Cellulase (Yakult R-10) 1.5% w/v, Macerozyme (Yakult R-10) 0.75% w/v,

$CaCl_2$(10 mM). BSA 0.1% w/v). The dish was wrapped in aluminum foil to shield it from light

and placed on an orbital shaker, horizontally rotating at 60 RPM, for three hours. Protoplasts

were collected by gentle centrifugation at 100 x g for 3 mins in a 30 mL round-bottomed tube,

then the enzyme solution was removed and the pellet was washed in W5 solution (MES-pH5.7 (2

mM), KCl (5 mM), $CaCl_2$ (125 mM), NaCl (154 mM). Five µL of concentrated protoplasts were

removed and used to quantify cells and check quality using a hemocytometer and light

microscope. Protoplasts were washed again with W5, this time left on ice for 45 minutes to

settle. W5 solution was removed, and protoplasts were resuspended to 300,000-350,000

protoplasts/mL using MMG (MES-pH5.7 (4 mM), Mannitol (0.4 mM), $MgCl_2$ (15 mM)).

Protoplasts were used immediately for transfection.

Protoplast transfection

High concentration (1 µg/µL) plasmid DNA was extracted via maxi-prep (Promega SV Wizard).

Lower quality plasmid DNA (e.g. from miniprep kits) typically resulted in low transfection rates.

The *Avr*, *R* and *luciferase* genes were expressed under control of maize ubiquitin promoter

(*UBI*), and 10 µg of each *Avr* and *R* plasmid and 20 µg of *luciferase* reporter plasmid were used

for each replicate. Due to high concentration and purity, plasmids were heated at $65^{\circ}C$ for 10 min

before use to reduce viscosity and ensure consistent measurement and delivery into cells. The

required plasmid DNA was added to 14 mL round-bottom culture tubes (Thermofisher,

#150268). 300 uL of protoplasts and 300 µL of PEG solution (PEG 4000 (40% w/v), Mannitol

(0.2 M), $CaCl_2$ (100 mM)) was then added and mixed immediately by gently flicking/rotating the

tube three times. Samples were incubated for 15 minutes, and the reaction halted with 900 µL

W5. Protoplasts were then collected by 100 x g centrifugation for 2 min, resuspended in 500 µL

modified W5 (W5 + 1 mM glucose) and transferred to a transparent, sterile 12 well culture plate

(Costar, #3513) whose wells were coated with 10% BSA to prevent sticking. Incubation was

carried out in light conditions, approximately 100 µmol/m^2/s. Wide bore or cut tips were used

throughout for handling protoplasts.

Protoplast Quantification

After the required incubation time (3, 4, 5, 6, and 18-hours for time course experiment, 4-18 hrs

for *pDefense-ELuc* reporters) protoplasts were prepared for luminescence measurement. Two-

hundred µL of 1x cell lysis buffer (Promega #E3971) was added and samples vortexed and

incubated for 15 min. Samples were centrifuged again to collect cell debris and 50 µL of lysate
was added to an opaque, flat-bottom, white, solid polystyrene 96 well plate (e.g. Corning® 96
well NBS™ Microplate #CLS3600). Plate was read for total luminescence (Tecan infinite
200Pro, 1000 ms integration 0 ms settle) without addition of substrate, then 50 µL Steady-Glo
(Promega #E2520) Luciferin substrate was added, and the plate was read again.

Protoplast RNAseq sampling

Protoplasts were prepared as described above, using 900 µL protoplast solution (350,000
cells/mL MMG) per replicate, with 4 replicates for each treatment group The *UBI-fLUC* reporter
was excluded. Additional replicates of each condition included the reporter to confirm cell-death
after 16hrs (Data not shown). At 4 hrs post transfection the protoplasts were pelleted, snap-
frozen, and stored at -80 until RNA isolation. Samples from each cultivar were prepared and
processed in batches. Qiagen RNeasy Plant Mini Kit was used as directed, (#74904) omitting the
first mechanical cell lysis step as freezing and addition of RLT buffer was sufficient to lyse cells.
Qiagen RNase-Free DNase Set (#79254) was used as directed. RNA was checked for absence of
DNA contamination by PCR and gel electrophoresis and quantitated using Qubit and Nanodrop.
Total RNA Illumina sequencing was performed on the NovaSeq platform, in a 2x150bp paired-
end configuration, with poly A selection & strand specific library preparation for the 36 samples.
The resulting sequence data is deposited in the NCBI Sequence Read Archive (SRA) under
BioProject number PRJNA957082.

RNAseq Analysis

Transcript abundance was quantified with kallisto (Bray *et al.*, 2016) using the Ensembl *Triticum*
aestivum transcriptome (IWGSC, INSDC Assembly GCA_900519105.1, Jul 2018) plus cDNA
sequences for *AvrSr50, Sr50*, *AvrSr35* and *Sr35* as reference. DESeq2 (Love *et al.*, 2014) was
used to test for differential expression. We defined upregulation at a 4-fold change or greater.
GaboSr50 with AvrSr50 was compared against the four control treatment groups;
Gabo_AvrSr50, Gabo_EV, GaboSr50_AvrSr35 and GaboSr50_EV. Then, the intersect of the
upregulated genes was taken, for a total of 272 genes. For Fielder, Fielder_Sr35_AvrSr35 was
compared to the 3 control treatment groups; Fielder_AvrSr35, FielderSr35_AvrSr50,
Fielder_Sr35. Again, taking the intersect of the upregulated genes returned 98 upregulated genes.
66 genes were commonly upregulated between both lists (Zenodo Supplemental Data S5). For

GO terms analysis, we repeated the comparisons above with lower upregulation of 2-fold change

and used the gProfiler webtool and Ensembl *Triticum aestivum* annotation, running the g:GOst

program (Raudvere *et al.*, 2019). The recommended g:SCS multiple testing correction algorithm

was used.

pDefense **Promoter Selection**

Expression in log fold change and transcripts per million (TPM) was visualized for the 66

commonly upregulated genes, present in both 2-fold upregulated lists used for GO analysis.

Thirteen genes were selected based on their overall expression and upregulation in *Avr/R*

treatment groups. The entire gene and region were compared in Geneious and 600-800 bp

upstream of the 13 genes was selected, including any annotated 5'UTR. These promoter

sequences then were synthesized with 5'and 3' 4bp MoClo promoter overhangs flanked by BsaI

recognition sites for Golden Gate cloning as promoter+5'UTR standard parts (Weber *et al.*,

2011). Any BpiI and BsaI enzyme recognition sites within the promoter sequence were removed

prior to synthesis. Promoters that could not be synthesized were cloned by PCR, using primers

(Zenodo Supplemental Data S6) with BpiI recognition sites and 4bp overhang to facilitate

assembly into universal level 0 vector *pAGM9121* (Addgene Plasmid #51833). Any BsaI

recognition sites were removed as per Grutzner & Marillionnet, (2020).

pDefense **Plasmid Constructs**

Plasmids were designed in Geneious. The plasmid constructs were designed according to the

MoClo syntax, assembled by Golden Gate cloning and verified by whole plasmid sequencing.

Supplemental Data (Zenodo Supplemental Data S6) and plasmid maps (Zenodo DOI:

10.5281/zenodo.7844465, link: https://zenodo.org/record/7844465) are available, with all

generated plasmids deposited on Addgene.

pWDV1 **vector**

The wheat dwarf virus genomic sequence from the isolate Enkoping1 (GenBank: AJ311031.1)

was modified to remove the V1 and V2 genes which encode the viral coat and transfer proteins,

respectively. In addition, one and two BsaI type II restriction sites were mutated to enable

downstream golden gate cloning steps. All mutations were in the Rep and RepA open reading

frames and were silent, with the exception that the mutation generated at amino acid position 263

for RepA results in a Serine to Threonine substitution at amino acid position 235 of the Rep

protein which is encoded by an alternately spliced transcript from the same Rep/RepA open

reading frame. Two BsaI sites were inserted between the Long and short intergenic regions in

place of the removed V1 and V2 genes. The BsaI sites were designed to generate a GGAG

overhang downstream of the long intergenic region and a CGCT overhang upstream of the short

intergenic region to facilitate level 1 golden gate assemblies as described previously (Engler *et*

al., 2014). BsmBI sites were also included at either end of the modified viral genome sequence

to facilitate golden gate cloning into the pYTK001 vector (Lee *et al.*, 2015) which was chosen

for its small size, ease of cloning and to facilitate selection and plasmid replication in bacteria.

The modified wheat dwarf virus sequence, called WDV_1, was synthesised as a 'gene' fragment

by twist bioscience and directly incorporated into the pYTK001 vector by golden gate cloning

using BsmBI. We sequence verified plasmids via whole plasmid sequencing. We named the

resulting plasmid *pWDV1*. We used *pWDV1* for subsequent cloning of genes by golden gate

cloning using the BsaI entry site.

Ratiometric Assay

All isolation and transfection steps were as above aside from the substitution of the fLUC

reporter for the reporter (*UBI-pDefense)* and normalization (*UBI-pRedf*) plasmids. These require

the same substrate for activation, but express luciferase with different emission spectra, read by

specific green and red filters in the luminescence plate reader.A *UBI-ELuc* construct was also

generated to calibrate normalization and deconvolution calculations, necessary to resolve the

overlap in emission spectra of the two luciferase reporters (González-Grandío *et al.*, 2021).

These calibration experiments were performed separately three times and with three replicates

per measurement, according to the instructions for Promega Chroma-Glo assay (Chroma-

Glo(TM) Luciferase Assay System Technical Manual, TM062). The plate reader was

programmed to measure total luciferase, then with green and red filters, with 1000 ms integration

time and 0 ms settle time, measured well-wise. Template spreadsheets for performing

calculations and a Python script for automating deconvolution and plotting are recorded on

protocols.io (https://dx.doi.org/10.17504/protocols.io.q26g7r3zkvwz/v1).

Cultivar testing

Protoplasts were isolated from nine different wheat cultivars (GaboSr50, Coorong, C6969,

Clement, Compair, HeinesPeko, KenyaW1483, AvocetYr8 and Fielder), using the method

described above. All were grown in the same conditions as above for 8 days. Where the cultivar

carried an *R* gene (GaboSr50, Coorong, C6969) the recognized *Avr* (*AvrSr50*, *AvrSr27* and

AvrSr35 respectively) was transfected alone, with an avr (unrecognized by R protein) as control.

In the other six cultivars both the *Avr* and *R* gene were co-transfected. Normalized luminescence

was calculated from the ratio of deconvoluted *pDefense-Eluc* to *UBI-pRedf.*

Multiplex screening

Protoplasts were prepared from cv. GaboSr50 and cv. Gabo wheat leaves and transfected as

above. Measurements used the ratiometric reporters and measurements, with *pDefense14-ELuc*

as reporter and *UBI-pRedf* for normalisation. The pool of twenty-five *Avr* candidates including

AvrSr50, AvrSr35, AvrSr27, AvrSr22, AvrSr13, AvrStb6, Avr3D1, AvrStb9, AvrPm1a, AvrPm2,

AvrPm3d3, AvrPm17, AvrSr50-A1, AvrSr50-B6, and other eleven wheat rust *Avr* candidates used

in our studies, including *Pst_104E_10266, Pst_M28_10266, Pst_104E_17331, Pst_M28_17331,*

Pst_104E_25466, Pst_M28_25466, Pttg_05697, Pttg_27353 and three alternatively spliced

version of *Pst_104E_25466.* The pool excluding the *AvrSr50* acted as negative controls, and 1

µg of *pWDV1-AvrSr50* alone acted as a positive control. Data processing and statistics was

carried out as above.

Data availability

All supplemental data is available on Zenodo (DOI: 10.5281/zenodo.7844446., link:

https://doi.org/10.5281/zenodo.7844446). All plasmids generated are available from Addgene,

with plasmid maps deposited on Zenodo (DOI: 10.5281/zenodo.7844464, link:

https://doi.org/10.5281/zenodo.7844464). Code used to perform RNAseq analysis is deposited

on Github (https://github.com/ritatam/wheat-protoplast-RNAseq-DESeq2) and all RNAseq data

is available on NCBI Sequence Read Archive (SRA) under BioProject number PRJNA957082.

Acknowledgements

We thank Prof. Robert Park and Prof. Robert McIntosh for providing seeds of cv. C6969, A/Prof.
Sambasivam Periyannan for providing seeds of cv. Fielder, cv. Gabo, cv. GaboSr50, and cv.
Coorong, A/Prof. Peng Zhang for providing seeds of Federation*4/Ulka, Dr. Michael Ayliffe and
Dr. Peter Dodds for providing *pUBI-AvrSr35*, *pUBI-Sr35*, *pUBI-AvrSr50*, and *pUBI-Sr50*. We
are grateful to Dr. Cecile Lorrain for useful suggestions on AvrStb6/Stb6 interaction. We thank
Jasper Lee for plant maintenance. B.S. acknowledges the financial support by the Australian
Research Council via a Future Fellowship (FT180100024). This work was supported by
computational resources provided by the Australian Government through the National
Computational Infrastructure (NCI) under the ANU Merit Allocation Scheme.

Author Contributions

572 S. W., B. D., J. P. R., and B. S. conceived the project. S. W., B. D., R. T., L. M., N. H., S. S. K.,
573 E. C., and J. G. performed the experiments, S. W., B. D., and B. S. wrote the manuscript. R. T.,
574 E. C., and J. P. R. provided extensive feedback on the manuscript. B. S. secured funding. All
575 authors read and approved the manuscript.

References

Aarts N, Metz M, Holub E, Staskawicz BJ, Daniels MJ, Parker JE. **1998**. Different
requirements for *EDS1* and *NDR1* by disease resistance genes define at least two *R* gene-
mediated signaling pathways in *Arabidopsis*. *Proceedings of the National Academy of Sciences*
95: 10306–10311.

Amezrou R, Audéon C, Compain J, Gélisse S, Ducasse A, Saintenac C, Lapalu N, Louet C,
Orford S, Croll D, *et al.* 2023. A secreted protease-like protein in *Zymoseptoria tritici* is
responsible for avirulence on *Stb9* resistance gene in wheat. *PLOS Pathogens* **19**: e1011376.

Arndell T, Chen J, Sperschneider J, Updhyaya NM, Blundell C, Niesner N, Wang A, Swain
S, Luo M, Ayliffe MA, *et al.* 2023. Pooled effector library screening in protoplasts rapidly
identifies novel *Avr* genes. *bioRxiv*: 2023.04.28.538616.

Bi G, Su M, Li N, Liang Y, Dang S, Xu J, Hu M, Wang J, Zou M, Deng Y, *et al.* 2021. The
ZAR1 resistosome is a calcium-permeable channel triggering plant immune signaling. *Cell* **184**:
3528-3541.e12.

Boudsocq M, Willmann MR, McCormack M, Lee H, Shan L, He P, Bush J, Cheng S-H,
Sheen J. **2010**. Differential innate immune signalling via Ca2+ sensor protein kinases. *Nature*
464: 418–422.

Bourras S, Kunz L, Xue M, Praz CR, Müller MC, Kälin C, Schläfli M, Ackermann P, Flückiger S, Parlange F, *et al.* **2019**. The *AvrPm3-Pm3* effector-NLR interactions control both race-specific resistance and host-specificity of cereal mildews on wheat. *Nature Communications* **10**: 2292.

Bouton C, King RC, Chen H, Azhakanandam K, Bieri S, Hammond-Kosack KE, Kanyuka K. **2018**. *Foxtail mosaic virus*: A Viral Vector for Protein Expression in Cereals. *Plant Physiology* **177**: 1352–1367.

Bray NL, Pimentel H, Melsted P, Pachter L. **2016**. Near-optimal probabilistic RNA-seq quantification. *Nature Biotechnology* **34**: 525–527.

Century KS, Holub EB, Staskawicz BJ. **1995**. *NDR1*, a locus of *Arabidopsis thaliana* that is required for disease resistance to both a bacterial and a fungal pathogen. *Proceedings of the National Academy of Sciences* **92**: 6597–6601.

Chen J, Upadhyaya NM, Ortiz D, Sperschneider J, Li F, Bouton C, Breen S, Dong C, Xu B, Zhang X, *et al.* **2017**. Loss of *AvrSr50* by somatic exchange in stem rust leads to virulence for *Sr50* resistance in wheat. *Science* **358**: 1607–1610.

Christensen AH, Sharrock RA, Quail PH. **1992**. Maize polyubiquitin genes: structure, thermal perturbation of expression and transcript splicing, and promoter activity following transfer to protoplasts by electroporation. *Plant Molecular Biology* **18**: 675–689.

Coppinger P, Repetti PP, Day B, Dahlbeck D, Mehlert A, Staskawicz BJ. **2004**. Overexpression of the plasma membrane-localized NDR1 protein results in enhanced bacterial disease resistance in *Arabidopsis thaliana*. *The Plant Journal* **40**: 225–237.

Dagvadorj B, Outram MA, Williams SJ, Solomon PS. **2022**. The necrotrophic effector ToxA from *Parastagonospora nodorum* interacts with wheat NHL proteins to facilitate *Tsn1*-mediated necrosis. *The Plant Journal* **110**: 407–418.

Dodds PN, Rathjen JP. **2010**. Plant immunity: towards an integrated view of plant–pathogen interactions. *Nature Reviews Genetics* **11**: 539–548.

Dracatos PM, Zhang P, Park RF, McIntosh RA, Wellings CR. **2016**. Complementary resistance genes in wheat selection 'Avocet R' confer resistance to stripe rust. *TAG. Theoretical and applied genetics. Theoretische und angewandte Genetik* **129**: 65–76.

Engler C, Youles M, Gruetzner R, Ehnert T-M, Werner S, Jones JDG, Patron NJ, Marillonnet S. **2014**. A Golden Gate Modular Cloning Toolbox for Plants. *ACS Synthetic Biology* **3**: 839–843.

Förderer A, Li E, Lawson AW, Deng Y, Sun Y, Logemann E, Zhang X, Wen J, Han Z, Chang J, *et al.* **2022**. A wheat resistosome defines common principles of immune receptor channels. *Nature* **610**: 532–539.

Gil-Humanes J, Wang Y, Liang Z, Shan Q, Ozuna CV, Sánchez-León S, Baltes NJ, Starker C, Barro F, Gao C, et al. 2017. High-efficiency gene targeting in hexaploid wheat using DNA replicons and CRISPR/Cas9. *The Plant Journal* **89**: 1251–1262.

González-Grandío E, Demirer GS, Ma W, Brady S, Landry MP. 2021. A Ratiometric Dual Color Luciferase Reporter for Fast Characterization of Transcriptional Regulatory Elements in Plants. *ACS Synthetic Biology* **10**: 2763–2766.

Hewitt T, Müller MC, Molnár I, Mascher M, Holušová K, Šimková H, Kunz L, Zhang J, Li J, Bhatt D, et al. 2021. A highly differentiated region of wheat chromosome 7AL encodes a *Pm1a* immune receptor that recognizes its corresponding *AvrPm1a* effector from Blumeria graminis. *The New Phytologist* **229**: 2812–2826.

Jacob P, Kim NH, Wu F, El-Kasmi F, Chi Y, Walton WG, Furzer OJ, Lietzan AD, Sunil S, Kempthorn K, et al. 2021. Plant "helper" immune receptors are Ca^{2+}-permeable nonselective cation channels. *Science* **373**: 420–425.

Jensen C, Saunders DGO. 2023. *Magnaporthe oryzae* pathotype *Triticum* (MoT) can act as a heterologous expression system for fungal effectors with high transcript abundance in wheat. *Scientific Reports* **13**: 108.

Laflamme B, Dillon MM, Martel A, Almeida RND, Desveaux D, Guttman DS. 2020. The pan-genome effector-triggered immunity landscape of a host-pathogen interaction. *Science* **367**: 763–768.

Lee ME, DeLoache WC, Cervantes B, Dueber JE. 2015. A Highly Characterized Yeast Toolkit for Modular, Multipart Assembly. *ACS synthetic biology* **4**: 975–986.

Lee W-S, Hammond-Kosack KE, Kanyuka K. 2012. *Barley Stripe Mosaic Virus*-Mediated Tools for Investigating Gene Function in Cereal Plants and Their Pathogens: Virus-Induced Gene Silencing, Host-Mediated Gene Silencing, and Virus-Mediated Overexpression of Heterologous Protein. *Plant Physiology* **160**: 582–590.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology* **15**: 550.

Luo M, Xie L, Chakraborty S, Wang A, Matny O, Jugovich M, Kolmer JA, Richardson T, Bhatt D, Hoque M, et al. 2021. A five-transgene cassette confers broad-spectrum resistance to a fungal rust pathogen in wheat. *Nature Biotechnology* **39**: 561–566.

Mago R, Zhang P, Vautrin S, Šimková H, Bansal U, Luo M-C, Rouse M, Karaoglu H, Periyannan S, Kolmer J, et al. 2015. The wheat *Sr50* gene reveals rich diversity at a cereal disease resistance locus. *Nature Plants* **1**: 15186.

McIntosh R, Dyck P, The T, Cusick J, Milne D. 1984. Cytogenetical studies in wheat XIII. *Sr35* - a third gene from *Triticum monococcum* for resistance to *Puccinia graminis tritici*. *Zeitschrift für Pflazenzüchtung* **92**: 1–14.

McIntosh R, Wellings C, Park R. **1995**. Wheat Rusts: An Atlas of Resistance Genes. In:
CSIRO Publishing.

Meile L, Croll D, Brunner PC, Plissonneau C, Hartmann FE, McDonald BA, Sánchez-
Vallet A. **2018**. A fungal avirulence factor encoded in a highly plastic genomic region triggers
partial resistance to septoria tritici blotch. *New Phytologist* **219**: 1048–1061.

Müller MC, Kunz L, Schudel S, Lawson AW, Kammerecker S, Isaksson J, Wyler M, Graf
J, Sotiropoulos AG, Praz CR, *et al*. 2022. Ancient variation of the *AvrPm17* gene in powdery
mildew limits the effectiveness of the introgressed rye *Pm17* resistance gene in wheat.
Proceedings of the National Academy of Sciences **119**: e2108808119.

Ortiz D, Chen J, Outram MA, Saur IML, Upadhyaya NM, Mago R, Ericsson DJ, Cesari S,
Chen C, Williams SJ, *et al*. 2022. The stem rust effector protein AvrSr50 escapes Sr50
recognition by a substitution in a single surface-exposed residue. *The New Phytologist* **234**: 592–
606.

Park PR. **2016**. The wheat stem rust pathogen in Australia- pathogenic variation and pathotype
designation. *Cereal Rust Report 2016* **14**: 1–4.

Praz CR, Bourras S, Zeng F, Sánchez-Martín J, Menardo F, Xue M, Yang L, Roffler S,
Böni R, Herren G, *et al*. 2017. *AvrPm2* encodes an RNase-like avirulence effector which is
conserved in the two different specialized forms of wheat and rye powdery mildew fungus. *The*
New Phytologist **213**: 1301–1314.

Raudvere U, Kolberg L, Kuzmin I, Arak T, Adler P, Peterson H, Vilo J. **2019**. g:Profiler: a
web server for functional enrichment analysis and conversions of gene lists (2019 update).
Nucleic Acids Research **47**: W191–W198.

Saintenac C, Lee W-S, Cambon F, Rudd JJ, King RC, Marande W, Powers SJ, Bergès H,
Phillips AL, Uauy C, *et al*. 2018. Wheat receptor-kinase-like protein Stb6 controls gene-for-
gene resistance to fungal pathogen *Zymoseptoria tritici*. *Nature Genetics* **50**: 368–374.

Saintenac C, Zhang W, Salcedo A, Rouse MN, Trick HN, Akhunov E, Dubcovsky J. **2013**.
Identification of wheat gene *Sr35* that confers resistance to Ug99 stem rust race group. *Science*
(New York, N.Y.) **341**: 783–786.

Salcedo A, Rutter W, Wang S, Akhunova A, Bolus S, Chao S, Anderson N, De Soto MF,
Rouse M, Szabo L, *et al*. 2017. Variation in the *AvrSr35* gene determines *Sr35* resistance
against wheat stem rust race Ug99. *Science* **358**: 1604–1606.

Sánchez-Martín J, Widrig V, Herren G, Wicker T, Zbinden H, Gronnier J, Spörri L, Praz
CR, Heuberger M, Kolodziej MC, *et al*. 2021. Wheat *Pm4* resistance to powdery mildew is
controlled by alternative splice variants encoding chimeric proteins. *Nature Plants* **7**: 327–341.

Sarrion-Perdigones A, Chang L, Gonzalez Y, Gallego-Flores T, Young DW, Venken KJT.
2019. Examining multiple cellular pathways at once using multiplex hextuple luciferase
assaying. *Nature Communications* **10**: 5710.

Saur IML, Bauer S, Lu X, Schulze-Lefert P. **2019**. A cell death assay in barley and wheat
protoplasts for identification and validation of matching pathogen AVR effector and plant NLR
immune receptors. *Plant Methods* **15**: 118.

Savary S, Willocquet L, Pethybridge SJ, Esker P, McRoberts N, Nelson A. **2019**. The global
burden of pathogens and pests on major food crops. *Nature Ecology & Evolution* **3**: 430–439.

Selote D, Shine MB, Robin GP, Kachroo A. **2014**. Soybean NDR1-like proteins bind pathogen
effectors and regulate resistance signaling. *New Phytologist* **202**: 485–498.

Upadhyaya NM, Mago R, Panwar V, Hewitt T, Luo M, Chen J, Sperschneider J, Nguyen-
Phuc H, Wang A, Ortiz D, *et al*. **2021**. Genomics accelerated isolation of a new stem rust
avirulence gene–wheat resistance gene pair. *Nature Plants* **7**: 1220–1228.

Weber E, Gruetzner R, Werner S, Engler C, Marillonnet S. **2011**. Assembly of Designer
TAL Effectors by Golden Gate Cloning. *PLOS ONE* **6**: e19722.

Win J, Chaparro-Garcia A, Belhaj K, Saunders DGO, Yoshida K, Dong S, Schornack S,
Zipfel C, Robatzek S, Hogenhout SA, *et al*. **2012**. Effector Biology of Plant-Associated
Organisms: Concepts and Perspectives. *Cold Spring Harbor Symposia on Quantitative Biology*
77: 235–247.

Wulff BB, Krattinger SG. **2022**. The long road to engineering durable disease resistance in
wheat. *Current Opinion in Biotechnology* **73**: 270–275.

Zheng MS, Takahashi H, Miyazaki A, Hamamoto H, Shah J, Yamaguchi I, Kusano T.
2004. Up-regulation of *Arabidopsis thaliana NHL10* in the hypersensitive response to *Cucumber*
mosaic virus infection and in senescing leaves is controlled by signalling pathways that differ in
salicylate involvement. *Planta* **218**: 740–750.

Zhong Z, Marcel TC, Hartmann FE, Ma X, Plissonneau C, Zala M, Ducasse A, Confais J,
Compain J, Lapalu N, *et al*. **2017**. A small secreted protein in *Zymoseptoria tritici* is
responsible for avirulence on wheat cultivars carrying the *Stb6* resistance gene. *New Phytologist*
214: 619–631.

Zipfel C, Robatzek S, Navarro L, Oakeley EJ, Jones JDG, Felix G, Boller T. **2004**. Bacterial
disease resistance in *Arabidopsis* through flagellin perception. *Nature* **428**: 764–767.

Figure Legends

Figure 1: Rapid onset of cell-death downstream of *Avr/R* interactions in protoplasts. **A-C**,

Luciferase activity was measured at defined timepoints in hours post-transfection (hpt) in wheat

protoplasts. **(A)** Shows boxplots of cv. Fielder protoplasts expressing *luciferase* alone, and

luciferase with either *Sr35, AvrSr35,* or *Sr35* plus *AvrSr35*. **(B)** shows boxplots of cv. GaboSr50

protoplasts expressing *luciferase* alone, and *luciferase* with either *AvrSr35* or *AvrSr50.* **(C)**
shows the luminescence of the Avr/R interaction treatment groups from cv. Fielder and cv.
GaboSr50 alone when compared to the no plasmid transfection control replicates (red line)
which is the average of nine replicate across the six timepoints.

Figure 2: Co-expression of *Avr/R* gene pairs triggers defense signaling in wheat
protoplasts. (A) Heat map shows expression of *Avr* and *R* genes by normalized gene count.
Each column represents a replicate and the treatment group is indicated by a number as shown in
the legend. *R* or *Avr* genes are expressed in their respective treatment groups only. This includes
endogenously expressed *Sr50* in cv. GaboSr50. **(B-D)** show principal component analysis (PCA)
of the RNAseq analysis; all 36 samples **(B)**, cv. Fielder **(C)** and cv. Gabo/GaboSr50 **(D)**. Figure
legend numbers corresponds to **(A)**, while colors correspond to **(B), (C)** and **(D)**. **(E)** shows the
GO terms of genes that are upregulated 2 fold in *Avr/R* treatment groups. A full list of
upregulated genes with overrepresented GO terms is recorded in supplemental data (Zenodo
Supplemental Data S4). **(F)** shows a heatmap of the expression of thirteen candidate reporter
genes (D1-7, D10-12, D14-16) that are commonly upregulated by *Sr50/AvrSr50* and
Sr35/AvrSr35 in cv. GaboSr50 or cv. Fielder, respectively. Each square represents the mean of
four replicates for treatment group.

Figure 3: *pDefense14* is a positive defense reporter in wheat protoplasts. **(A)** Wheat
protoplasts transfected with *pDefense14* and *UBI-pRedF* constructs show an increase in
normalized luminescence in the *Avr/R* treatment, compared to treatments with *R* or *avr* only and
without *R* or *Avr*. Panel **(B)** shows corresponding non-normalized red-filtered luminescence for
the treatments in panel A. Treatments marked with common letters were not significantly
different (P >0.05) assessed with one-way analysis of variance (ANOVA) and post hoc Tukey's
HSD test.

Figure 4: *pWDV1* plasmid reduces amount of *Avr* plasmid required to detect defense
signaling using *pDefense* reporter. *AvrSr50* was cloned into *pWDV1* (*pWDV1-AvrSr50*) and
non-viral plasmid backbone (*pICH4772-UBI-AvrSr50)* and transfected in increasing amounts
(0.1, 0.5, 1, 5 and 15 µg of plasmid DNA) alongside *pDefense14* and *UBI-pRedf.* Reporters
remained in the non-replicative plasmid. Controls in cv. GaboSr50 were transfected with
pWDV1-AvrSr35 (1 and 15 µg) and cv. Gabo transfected with *pWDV1-AvrSr50* (1 µg) or

pWDV1-AvrSr35 (1 µg). One-way analysis of variance (ANOVA) and post hoc Tukey's HSD

test was used to assess differences within normalized luminescence **(A)** and red luminescence

(B) groups. Treatments marked with a common letter were not significantly different (P >0.05).

Figure 5: Positive *pDefense* reporter detects defense signaling triggered by seven distinct

***Avr/R* pairs in wheat protoplasts expressing the *R* gene endogenously.** Protoplasts from

wheat cultivars **(A)** cv. Coorong carrying *Sr27*, **(B)** cv. C6969 carrying *Sr35,* **(C)** cv. GaboSr50

carrying *Sr50,* **(D)** cv. Line S carrying *Sr13*, **(E)** cv. W3534 carrying *Sr22*, **(F)** cv.

Federation4/Ulka carrying *Pm2* and **(G)** cv. Chinese Spring carrying *Stb6* were transfected with

UBI-pDefense14 and *Avr* constructs, as well as an *avr* (unrecognized *Avr*) construct and empty

vector **(EV)** construct as controls. **(H)** had identical treatments to **(G)**, however cv. Heines

Kolben lacks *Stb6. UBI-pRedf* was used for normalization of all experiments. For **(H)** and **(G)**

AvrStb6 was expressed with (SP-*AvrStb6)* and without (*AvrStb6*) secretion peptide to allow for

extracellular recognition including the appropriate control. One-way analysis of variance

(ANOVA) and post hoc Tukey's HSD test was used to assess differences among normalized

luminescence for each cultivar. Treatments marked with a common letter were not significantly

different (P >0.05).

Figure 6: The positive defense reporter *pDefence14* is broadly applicable to all tested wheat

cultivars. Each panel shows the normalized luminescence of the positive defense reporter

pDefense14. Protoplasts isolated from various wheat cultivars were transfected with both *R* and

Avr constructs, aside from cultivars GaboSr50, C6969 and Coorong which carry *Sr50, Sr35* and

Sr27 respectively. The cultivars Corack, Clement, Kenya W1483, Avocet Yr8 and Compair,

tested with *AvrSr50/Sr50* and Heines Peko was tested with *AvrSr27/Sr27*. An *avr* (unrecognised

by R protein) was used as a control and *UBI-pRedf* was used for normalization of all

experiments. One-way analysis of variance (ANOVA) and post hoc Tukey's HSD test was used

to assess differences among means of normalized luminescence measurements for each cultivar.

Treatments marked with common letters were not significantly different (P >0.05).

Figure 7: The positive defense reporter assay enables multiplexed *Avr* screening. *Avr* pools

of twenty-five candidates were transfected into wheat protoplasts isolated from **(A)** cv.

GaboSr50 and **(B)** cv. Gabo, along with *pD14* reporter and *UBI-pRedf*. *Avr* candidates included

were; *AvrSr13, AvrSr22, AvrSr27, AvrSr35, AvrSr50, AvrStb6, Avr3D1, AvrStb9, AvrPm1,*

AvrPm2, AvrPm3 AvrPm17 and thirteen other wheat rust effector candidates. One µg of each

pWDV1-Avr was used, with the negative control excluding *AvrSr50* and positive control of 1 µg

of *AvrSr50* alone. For each cultivar, one-way analysis of variance (ANOVA) and post hoc

Tukey's HSD test was used to assess differences among means of normalized luminescence and

red luminescence measurements for each treatment group. Treatments marked with common

letters were not significantly different (P >0.05).

Supplemental Material Legends

Supplemental Figure 1: Diagram of plasmids used in protoplast assay. (A) *pDefense14*

consists of D14 promoter that drives expression of the green-shifted E-Luc cassette. *pDs 1-7, 10,*

11, 12, 15 and 16 were constructed with the same modular plasmid components, replacing the

relevant defence-induced promoter **(B)** UBI-*pRedf*, used for normalization, consists of a red-

shifted coding sequence driven by *UBI* promoter. **(C)** Example *Avr* candidate plasmid, with

pWDV1 self-replicating plasmid backbone.

Supplemental Figure 2: Preliminary successful screening of *pDefense2, -14* and *-15* defense

reporters. Normalized luminescence for *pDefense14,* (panel **A**)*, pDefense 15* (panel **B**) and

pDefense2 (panel **C**), in three wheat cultivars. Left to right, cv. C6969 which carries *Sr35*, cv.

GaboSr50 which carries *Sr50* and cv. Coorong, which carries *Sr27*. *Avr+R* treatments were

transfected with plasmid which expressed a recognized *Avr, avr+R* treatments expressed a non-

recognized *avr* and *R* alone expressed an empty vector. Reporters were normalized against *UBI-*

pRedf for each replicate. The three reporters produced an increase in normalized luminescence in

treatments with Avr/R interaction, across the three *Avr/R* pairs tested.

Supplemental Figure 3: *pDefence* reporters confirm *AvrSr50* variants evade recognition by

a single amino acid substitution. *AvrSr50* variants; avrsr50-A1 (Q121K), avrsr50-B6

(avrsr50QCMJC) and AvrSr50-C (AvrSr50RKQQC), from Ortiz et al., (2022) were synthesized

and cloned into *pWDV1* vectors. avrsr50-A1 had a single amino acid substitution (Q121K) that

prevented recognition by Sr50, similar to avrsr50-B6, while AvrSr50-C had six amino acid

changes but was still recognized by Sr50. One-way analysis of variance (ANOVA) and post hoc

Tukey's HSD test was used to assess differences among means of normalized and red-filtered

luminescence measurements for each treatment. Treatments marked with common letters were

not significantly different (P >0.05).

Supplemental Figure 4: The *Stb6* gene amplification profile in ten different wheat cultivars.
The *Stb6* (exp: 782 bp), and wheat Polymerase A1 gene (*TaPolA1*; exp: 500 bp) partial
amplifications were performed. PCR products of 3 μL were loaded on 1% agarose gel. M: 1 kb
Plus DNA ladder (NEB #N3200).

Supplemental Figure 5: *pDefense* reporter can detect lower purity *Avr* plasmid. In **(A)** both
maxi prepped (left) and mini prepped (right) *AvrSr35* plasmid was co-transfected with *R* and *avr*
plasmids, in protoplasts from cv. Fielder. **(B)** shows corresponding non-normalized red-filtered
luminescence. One-way analysis of variance (ANOVA) and post hoc Tukey's HSD test was used
to assess differences among means of normalized and red-filtered luminescence measurements
for each treatment. Treatments marked with common letters were not significantly different (P
>0.05).

Supplemental Material S1: Luminescence data for Figure 1, protoplast time course experiment.

Supplemental Material S2: List of samples and treatment groups for RNAseq experiment.

Supplemental Material S3: Lists of Gabo/GaboSr50, Fielder and all cultivar genes upregulated
by 2-fold.

Supplemental Material S4: GO terms analysis Query tables and g:profiler results tables

Supplemental Material S5: Lists of Gabo/GaboSr50, Fielder and all cultivar genes upregulated
4-fold.

Supplemental Material S6: List of primers used.

Supplemental Material S7: List of plasmids used.

Supplemental Material S8: Luminescence data used for generating Figures 3 -7.

Supplemental Material S9: Statistical output for Figures 3-7.

Supplemental Material S10: Data and statistical output for Supplemental Figures 2, 3 and 4.

A

Normalized count (color scale: 1–4)

AvrSr35
AvrSr50
Sr35
Sr50

1 2 3 4 5 6 7 8 9

B All Samples

PC2: 5% variance

PC1: 90% variance

C Fielder

PC2: 33% variance

PC1: 46% variance

D Gabo/GaboSr50

PC2: 17% variance

PC1: 77% variance

Treatment Group

- 1. Fielder+*AvrSr35*
- 2. Fielder+*Sr35*+*AvrSr35*
- 3. Fielder+*Sr35*+*AvrSr50*
- 4. Fielder+*Sr35*
- 5. Gabo+*AvrSr50*
- 6. Gabo+*EV*
- 7. GaboSr50+*AvrSr35*
- 8. GaboSr50+*AvrSr50*
- 9. GaboSr50+*EV*

E

calcium ion binding
ion binding
metal ion binding
cation binding
P-type calcium transporter activity
membrane
intrinsic component of membrane
integral component of membrane
cell periphery
plasma membrane

-p adj log10

Cultivar
- Gabo/GaboSr50
- Fielder

number of genes
15
45
75

F

Fielder   Gabo/GaboSr50

Log Normalized Count (color scale: 7–10)

D1
D2
D3
D4
D5
D6
D7
D10
D11
D12
D14
D15
D16

Avr + + − −   + + − − −
avr − − + +   − − − − −
R   + − + +   + − − + +