1   **Title: Multiplexed effector screening for recognition by endogenous resistance genes using**
2   **positive defense reporters in wheat protoplasts**
3
4   Salome Wilson*, Bayantes Dagvadorj*, Rita Tam, Lydia Murphy, Nigel Heng, Emma Crean[$],
5   Julian Greenwood, John P. Rathjen, and Benjamin Schwessinger
6
7   Research School of Biology, The Australian National University, Canberra, ACT, 2601,
8   Australia
9
10 *These authors contributed equally to this work
11 [$]Current address: Institute for Plant Sciences, University of Cologne, Cologne, Germany
12
13
14 **Corresponding author: Benjamin Schwessinger**
15 Email: benjamin.schwessinger@anu.edu.au
16
17
18 **Short Title:**
19 Defense signaling in wheat protoplasts
20
21

Abstract

A key goal in agriculture is to create pathogen-resistant crop species via breeding or
biotechnological approaches. Plant resistance (*R*) and pathogen avirulence (*Avr*) gene
interactions play a vital role in pathogen resistance. Efficient molecular screening tools for crops
lack far behind their model organism counterparts, yet they are essential to rapidly identify
agriculturally important molecular interactions that trigger host resistance. To fill this important
gap, we have developed a novel wheat protoplast assay that enables efficient screening of Avr/R
interactions at scale. Our assay allows access to the extensive gene pool of phenotypically
described *R* genes because it does not require the overexpression of cloned *R* genes. It is suitable
for multiplexed *Avr* screening, with interactions tested in pools of up to fifty *Avr* candidates. Our

assay is based on newly identified Avr/R-induced defense genes which we used to create

promoter-luciferase reporter constructs. We combined these reporter constructs with a synthetic

biology based dual-color ratiometric reporter system that normalizes read-outs accounting for

experimental variability and Avr/R-induced cell-death. In addition, we introduced a replicon-

based plasmid derived from the wheat dwarf virus, which self-replicates in wheat reducing the

amount of plasmid used in the assay. Our new assay increases the throughput of *Avr* candidate

screening, accelerating the study of cellular defense signaling and resistance gene identification

in wheat. We anticipate that the uptake of our assay by the community will significantly

accelerate *Avr* identification for many wheat pathogens, leading to improved genome-guided

pathogen surveillance and breeding of disease-resistant crops.

Introduction

Plant pathogenic fungi cause severe damage to cereal crops. These pathogens secrete hundreds of

proteinaceous molecules called effectors during infection which manipulate plant processes to

facilitate disease (Dodds and Rathjen, 2010). Some effectors can be recognized by resistance (R)

proteins and are therefore also called avirulence (Avr) proteins. Avr recognition triggers a suite

of defense-related host responses to fight off pathogen infection including the hypersensitive

response (HR) that leads to cell-death at the site of infection and halts pathogen growth

(Laflamme et al., 2020). The deployment of suitable *R* genes is a major breeding objective to

reduce the negative impact of pathogens on crop production including in wheat. Recent years

have made significant progress in identifying and cloning a wide variety of *R* genes in wheat

(Wulff and Krattinger, 2022). Yet the identification of the corresponding *Avr* genes is lacking far

behind. For example, there are only four confirmed *Blumeria graminis* f. sp. *tritici Avr* genes

(Praz et al., 2017; Bourras et al., 2019; Hewitt et al., 2021; Müller et al., 2022), two for

Zymoseptoria tritici (Zhong et al., 2017; Meile et al., 2018), and five for *Puccinia graminis* f. sp.

tritici (Chen et al., 2017; Salcedo et al., 2017; Upadhyaya et al., 2021; Arndell et al., 2023).

There are no confirmed *Avr* genes for *Puccinia triticina* and *Puccunia striiformis* f. sp. *tritici*

despite both belonging to the top five wheat pathogens globally (Savary et al., 2019). This lack

of confirmed *Avr* genes in wheat pathogen and especially rust fungi is confounded by the

absence of an efficient screening tool for *Avr* recognition in wheat. Most current methods require

to have *Avr* and *R* gene candidates cloned, express them heterologously, and rely on HR as

general read-out for Avr/R interactions. For example, the model plant *Nicotiana benthamiana* is

used to validate Avr/R interactions as it produces visible HR upon transient expression of *Avr/R*

genes through agroinfiltration. Virus-mediated overexpression (VOX) can be used to screen *Avr*

candidates in wheat cultivars carrying the corresponding *R* gene as the Avr recognition will lead

to visible HR symptoms on the infected leaves (Lee et al., 2012; Bouton et al., 2018). However,

the method is only applicable to small proteins (Lee et al., 2012) and used viruses are not

accessible globally due to biosecurity regulations (Bouton et al., 2018). Recently, HR-based

assays in wheat protoplasts have emerged as a potential alternative to test Avr/R interactions

(Saur et al., 2019). In this transient assay, wheat protoplasts are co-transfected with *Avr/R* genes

along with a *luciferase* reporter gene. Avr/R interaction triggers cell-death causing significant

decreases in luciferase enzyme activity. This assay has been used to validate *Avr* candidates from

wheat rust pathogens (Saur et al., 2019; Upadhyaya et al., 2021; Luo et al., 2021; Ortiz et al.,

2022). However, the HR-based wheat protoplast assay in its current form has several limitations

that make it impractical for high-throughput *Avr/R* screening. Firstly, it requires the transgenic

overexpression of both the *Avr* and *R* genes from plasmids (Saur et al., 2019). This prevents its

use when the *R* gene has not yet been cloned thus stopping access to large numbers of

phenotypically described but uncloned *R* genes (McIntosh et al., 1995). Moreover, it is well

known that luminescence is highly variable between biological replicates and experiments (Saur

et al., 2019; González-Grandío et al., 2021) which can make the results difficult to interpret for

high-throughput screening purposes. In addition, the assay in its current form relies on the

reduction of luciferase activity as read out which is unspecific and cannot be differentiated from

cell-death that is unrelated to HR. Lastly, the assay requires large amount of target gene plasmids

for the protoplast transfection to achieve observable Avr/R-triggered cell-death. These

limitations encouraged us to develop a novel assay in wheat protoplast that relies on a positive

defense specific read-out and accounts for experimental variability via normalization. Our new

assay allows the research community to access the whole wheat gene pool to screen for Avr/R

interactions and enables the investigation of downstream defense signaling directly in wheat.

Results

Cell-death triggered by Avr/R interactions in protoplasts is time sensitive

We first sought to characterize the time of HR onset in wheat protoplasts following *Avr/R*

recognition. We used the two published *Avr/R* gene pairs, *AvrSr35/Sr35* (Salcedo et al., 2017;

Saintenac et al., 2013) and *AvrSr50/Sr50* (Chen et al., 2017; Mago et al., 2015), from wheat stem

rust *P. graminis* f. sp. *tritici* for initial method development. We aimed to identify the optimal

time point for RNAseq analyses that captures transcriptional changes prior to the onset of HR

triggered by Avr/R interactions. As previously described (Saur et al., 2019), we used

luminescence as a proxy for cell viability after overexpression of the *luciferase* reporter gene in

conjunction with each *Avr/R* combination and corresponding control. We expressed all *Avr* and *R*

genes under control of maize ubiquitin promoter (*UBI*) (Christensen et al., 1992) when expressed

from a plasmid. We used the wheat cultivar (cv.) Fielder for the analysis of *AvrSr35/Sr35* and

expressed both genes from a plasmid. For *AvrSr50/Sr50*, we made use of the introgression of

Sr50 into cv. Gabo (GaboSr50) and generated protoplast from cv. Gabo and cv. GaboSr50

(Jensen and Saunders, 2023). We only expressed *AvrSr50* from a plasmid. We also included the

controls, the reporter alone and the reporter with *Avr* or *R* alone (Zenodo Supplemental Data S1).

We measured luminescence at 3, 4, 5, 6, and 18 h post-transfection (hpt). We observed a gradual

increase in luminescence measurements in protoplast expressing luciferase alone or when

expressed together with an *Avr* or *R* gene only (Figure 1, A and B). The luciferase activity was

strongest at 18 hpt in all treatment groups except for the ones expressing *Avr/R* pairs. In cv.

Fielder expressing *AvrSr35*/*Sr35* and in cv. GaboSr50 expressing *AvrSr50*, luminescence

measurements only increased slightly early on during the time course when compared to mock

treatments of protoplasts transfected with no DNA (Figure 1 C). At about 4 to 5 hpt we observed

a divergence of luciferase activity from control treatment groups when compare with the Avr/R

interaction treatment groups (Figure 1A and B). The lack of continuous increase of luciferase

activity specifically in protoplast expressing *Avr*/*R* pairs suggests that their co-expressing leads

to cell-death in wheat protoplasts soon after transfection.

Transfected protoplasts are suitable for replicable transcriptional analysis

We concluded that 4 hpt is a suitable time point to capture *Avr/R*-dependent transcriptional

changes before strong HR induction based on our time course analysis (Figure 1). We repeated

the experimental setup while omitting the reporter plasmid and collected protoplasts at 4 hpt for

RNA sequencing (Figure 2 A, Zenodo Supplemental Data S2). We included four biological

replicates per treatment group and sequenced each sample on the Illumina platform generating

>20 million 150 bp paired-end reads per replicate. We used kallisto (Bray et al., 2016) for

transcript quantification and DESeq2 (Love et al., 2014) for downstream differential expression

analysis. As a first quality control step we quantified the expression of all *Avr* and *R* genes to

investigate their specific expression pattern at 4 hpt. This analysis clearly showed that all genes

are correctly expressed and at similar levels, including in treatment groups that co-express *Avr/R*

gene pairs (Figure 2 A). In addition, it revealed that the endogenous expression of *Sr50* in cv.

GaboSr50 is much lower level than the overexpressed of *Sr35* in cv. Fielder from a plasmid

under the *UBI* promoter. Next, we performed principal component analysis (PCA) of all sample

replicates combined. This revealed that the factor "wheat cultivar" had the strongest impact on

the observed variation in transcript abundance and accounted for close to 95% of all variation on

PC1 and PC2 combined (Figure 2 B). Consequently, we analyzed each cultivar separately. The

wheat cultivar specific PCA analysis showed a clear impact of Avr/R interaction on transcript

abundance variation for both *AvrSr35/Sr35* and *AvrSr50/Sr50* (Figure 2 C and D). This impact

was replicable because all four biological replicates clustered closely together and clearly

separated treatment groups including Avr/R interactions sample types from their controls. This is

consistent with the cell-death results seen in Figure 1 A and B. Overall, these results suggest that

these data could be used to identify reporter genes specifically expressed downstream of Avr/R

interactions in wheat protoplasts.

Avr/R recognition in wheat protoplasts leads to defense signaling

Next, we investigated if the observed gene expression changes in *Avr/R* expressing protoplast

overlapped with gene ontology (GO) terms commonly associated with defense signaling

pathways in wheat and other plant species. We identified all genes that are specifically

upregulated in response to Avr/R interactions in cv. Gabo/Gabo50 and cv. Fielder above a 2-fold

change threshold. We identified 309 and 922 genes in *AvrSr35/Sr35* and *AvrSr50/Sr50*

expressing protoplasts, respectively, when compared to all their control treatment groups.

Complete gene lists are available on Zenodo as Supplemental Data S3. We performed a

statistical enrichment analysis using these gene lists to test for overrepresentation of defense

signaling associated GO terms. Our results revealed significant enrichment for GO terms related

Wilson et al., 2023

to cellular components of cell periphery (GO:0071944) and plasma membrane (GO:0005886) in

both cultivars (Figure 2 E). We identified defense response to other organisms (GO:0098542),

response to other organisms (GO:0051707), and biological process involved in interspecies

interaction between organisms (GO:0044419) when looking at cv. GaboSr50 alone (Zenodo

Supplemental Data S4). We found molecular functions GO terms being enriched for calcium ion

binding (GO:0005509), cation binding (GO:0043169), and calcium ion transmembrane

transporter activity (GO:0015085) in both cultivars (Figure 2 E). Our GO terms analysis overall

confirms that *Avr/R* expressing protoplasts undergo defense signaling before the onset of strong

cell-death.

We were then interested in genes that are commonly upregulated by the Avr/R interaction in

both cv. GaboSr50 and cv. Fielder. We identified 66 genes commonly upregulated in both

cultivars, (Zenodo Supplemental Data S5) of which only 16 genes were associated with three

different GO terms, all related to the molecular function of ion binding (Zenodo Supplemental

Data S4). Our results showed that several of the identified genes have been previously implicated

in plant defense signaling including TaNHL10 (TRAESCS3D02G368800) (Century et al., 1995;

Aarts et al., 1998; Dagvadorj et al., 2022).

In summary, we observed the upregulation of genes involved in defense signaling in wheat

protoplasts expressing *Avr/R* pairs. The promoters of commonly upregulated genes, by two *Avr/R*

pairs and in two wheat cultivars (Figure 2F), are promising candidates to generate *Avr/R*

inducible reporters.

Identification of positive defense reporters in wheat protoplasts

We further investigated the expression patterns of the 66 commonly upregulated genes to select a

subset whose promoters we could test as positive defense signaling reporters. We selected 13

promoters (D1-D7, D10-D12, and D14-D16) based on their genes' expression patterns in both

cultivars (Figure 2 F). To further evaluate these candidates, we cloned or synthesized their

promoters, consisting of 600-800 bp upstream of their respective genes' start codons. We

accounted for the impact of *Avr/R* induced HR upon luciferase activity (Figure 1) by adapting a

dual-color green/red luciferase reporter system (González-Grandío et al., 2021). This ratiometric

system uses green- and red-shifted luciferases (Sarrion-Perdigones et al., 2019), where green-

shifted luciferase is fused to a conditionally responsive promoter and red-shifted luciferase

serves as a constitutively expressed reference for internal normalization, thus greatly reducing

intra experimental variation.

To test the activity of our thirteen candidate defense promoters, we generated promoter fusion

constructs driving the expression of green-emitting luciferase (E-Luc). These plasmids are

named *pDefense* (*pD*[*1-7, 10-12, 14-16*]), respectively. We co-transfected these plasmids along

with internal normalization plasmid *UBI-pRedf* and *UBI-AvrSr50* into protoplasts isolated from

cv. GaboSr50 and Gabo. For the initial activity screening, we performed the test with two

technical replicates. Ten of these promoters did not result in specific and reproducible promoter

activation downstream of AvrSr50/Sr50 interactions when compared to their respective controls.

In contrast, the three promoters D2, D14, and D15 showed a strong increase in normalized

ratiometric luminescence specifically downstream of AvrSr50/Sr50 interactions (Supplemental

Figure 1). Figure 3 shows an example of the specific activation of D14 downstream of

AvrSr50/Sr50 in cv. GaboSr50 when compared to its control not expressing the *R* gene or the

matching *Avr*. These results indicate that our new ratiometric reporter system is able to detect

specific defense activation downstream the Avr/R interactions when the *R* gene is expressed

endogenously.

The wheat dwarf virus-derived replicons-based system further increases the sensitivity of
199

the positive defense reporter assay

The current wheat protoplast assay requires a large amount of DNA material for transfection to

obtain observable cell-death. This makes its application prohibitive for large scale screening of

many candidates or variants. We tested a replicon-based system using a disarmed version of the

wheat dwarf virus (WDV) in our expression vector backbone (Gil-Humanes et al., 2017). We

reasoned that this would generate self-replicating plasmids in wheat protoplasts and thereby

reduce the amount of plasmids required to detect defense activation in our new ratiometric wheat

protoplast assay. We assembled Golden Gate compatible WDV vector from Gil-Humanes et al.

(2017) and denoted it as *pWDV1* (see Material and Methods). We expressed *AvrSr50* under

ubiquitin promoter using *pWDV1* (*pWDV1-AvrSr50*) and used it to generate dose response

curves using different amounts (0.1, 0.5, 1, 5 and 15 µg) of plasmid DNA for transfection. We

measured induction of D14 induction (normalized luminescence) and cell viability (red

Wilson et al., 2023

luminescence) and compared it to the dose response of the standard non-viral plasmid
pICH4772-AvrSr50 (Figure 4). We observed specific D14 promoter induction in all samples of
cv. GaboSr50 protoplasts transfected with *pICH4772-AvrSr50* and *pWDV1-AvrSr50* when
compared to controls in cv. GaboSr50 transfected with *pWDV1-AvrSr35* (1 and 15 µg) and cv.
Gabo transfected with *pWDV1-AvrSr50* (1 µg) or *pWDV1-AvrSr35* (1 µg). We also observed that
the promoter induction increases relative to the amount of *pICH4772-AvrSr50* used. At the same
time, we were still able to observe induction of the defense reporter D14 even in the absence of
cell-death at 0.1, 0.5 and 1 µg plasmid used. Using *pWDV1-AvrSr50* further improved the
sensitivity of the assay as it increased defense reporter D14 activation at 0.1 and 0.5 µg plasmid
and decreased cell viability when compared to the same amounts of plasmid for *pICH4772-*
AvrSr50. Altogether, this result indicates that the ratiometric dual-color luciferase system
combined with *pDefense* reporter and *pWDV1* is highly sensitive and specific requiring much
less plasmid to detect Avr/R interactions than a single luciferase based wheat protoplast assays.

Positive defense reporters are highly sensitive to detect Avr/R interactions in wheat
protoplast expressing *R* genes endogenously

Next, we explored if the defense reporters require overexpression of the *R* gene and if they are
applicable to additional Avr/R interactions. We tested the induction of D2, D14 and D15
promoters expressing *AvrSr27, AvrSr35,* and *AvrSr50* under ubiquitin promoter in *pWDV1* using
1 µg plasmid in instead of the previously used 15 µg. We transfected cv. Coorong, C6969, and
GaboSr50, with *pWDV1* encoding the recognized *Avr*, an unrecognized *avr*, or the empty vector
control (Figure 5). Wheat cv. Coorong carries *Sr27* (Upadhyaya et al., 2021), and cv. C6969
(*Triticum aestivum*) carries *Sr35* which was introgressed from *Triticum monococcum* (McIntosh
et al., 1984). Out of three promoters, we observed a strong induction of D14 specifically
downstream of AvrSr27/Sr27, AvrSr35/Sr35, and AvrSr50/Sr50 interactions but not in any of the
controls (Figure 5), validating the D14 defense reporter as well as *pWDV1* applicability in wheat
protoplast assay. We decided to focus on D14 and switch all *Avr* expressions in *pWDV1* for the
remainder of the study but to keep the reporter genes in the non-viral vector. These results
support that the D14 promoter in combination with *pWDV1* is broadly applicable as a defense
reporter in wheat independent of the wheat cultivar and the specific *Avr/R* combination used. In

addition, this suggests that our protoplast assay is applicable to testing wheat cultivars that carry

endogenous *R* genes and does not require overexpression of a cloned *R* gene.

Positive defense reporters are broadly applicable for Avr/R interaction screening in many wheat cultivars

We aimed to illustrate the broad applicability of our new assay in a wide range of wheat cultivars

as current cell-death based assays were mostly conducted in cv. Fielder (Saur et al., 2019, 5;

Ortiz et al., 2022; Upadhyaya et al., 2021; Luo et al., 2021; Arndell et al., 2023). We tested the

defense signaling reporter D14 for its inducibility by three *Avr/R* combinations in protoplasts of

nine different wheat cultivars. We tested cultivars that carry *R* genes (cv. GaboSr50, cv. Coorong

and cv. C6969) with their corresponding *Avr* construct *(AvrSr50, AvrSr27* and *AvrSr35,*

respectively). The cultivars Clement, Compair, Avocet Yr8, (Dracatos et al., 2016) and Kenya

W1483 (McIntosh et al., 1995) were tested with overexpression of *AvrSr50/Sr50* combination.

The cultivars HeinesPeko (Dracatos et al., 2016), and Fielder were tested with *AvrSr27/Sr27* and

AvrSr35/Sr35 combinations, respectively. We include a negative control in all cultivars by

expressing the unrecognized *avr, Avr* or *R* gene alone (Figure 4). In all cultivars we tested, a

consistent robust induction of normalized luminescence of *pD14* was observed in protoplasts

expressing cognate *Avr/R*-pair compared to the negative controls. The relative normalized

luminescence and its induction in the *Avr/R* treatment group varied between the different

cultivars. These results indicate that the new wheat protoplast assay enables the detection of

Avr/R interactions in many wheat cultivars.

The positive defense reporter assay enables multiplexed screening of *Avr* candidates in bulk

To explore multiplexed screening of *Avr* candidates using the positive defense reporter assay, we

transfected a pool of exemplifying ten or fifty *Avr* candidates into protoplasts isolated from cv.

GaboSr50 and cv. Gabo. The pools include 1 µg of *pWDV1-AvrSr50*, and 4.5 µg of each

pWDV1-AvrSr35 and *pWDV1-AvrSr27* in the pool of ten, or 24.5 µg of each *pWDV1-AvrSr35*

and *pWDV1-AvrSr27* in the pool of fifty so that one µg of *Avr* plasmid represents one Avr

candidate. We used the same *Avr* pools excluding the *AvrSr50* as negative controls, and 1 µg of

pWDV1-AvrSr50 alone as a positive control. Each replicate included *pD14* reporter and *UBI-*

pRedf co-transfection to detect *Avr/R*-triggered defense promoter induction and normalized

luminescence measurements. In cv. GaboSr50 protoplasts, *AvrSr50* alone and the pooled mixes

containing *AvrSr50* induced a high level of normalized luminescence of *pD14* when compared to

the pooled mix without *AvrSr50* and all other treatment groups including controls in cv. Gabo

protoplasts (Figure 7). We observed a clear specific induction of the D14 promoter only in the

presence of *AvrSr50* in cv. GaboSr50 when provided as pairs or as multiplexed mixtures with no

statistical difference between the single or multiplexed treatment groups (Figure 7). Overall,

these results show that our positive defense reporter protoplast assay can be used to screen a pool

of *Avr* candidates against a wheat cultivar endogenously expressing the recognizing *R* gene.

Discussion

Here, we describe an improved wheat protoplast assay to screen avirulence effector candidates

and study cellular defense signaling in wheat. We identify Avr/R-induced promoters and use

them to generate reporter constructs that produce a positive readout during Avr/R interaction. We

also reduce variability of luminescence measurement by adapting a ratiometric dual-color

luciferase reporter system (Sarrion-Perdigones et al., 2019) and reduce the amount of required

plasmids significantly enabling assays from simple minipreps. We show the broad applicability

of the positive defense reporters by testing three *Avr/R* pairs from the wheat stem rust

pathosystem on nine different wheat cultivars. The improved wheat protoplast assay is highly

sensitive because it allows for multiplexed screening of *Avr* candidates in wheat cultivars

expressing *R* genes endogenously.

This approach has several advantages over previous methods applied to identify Avr/R

interactions. Firstly, it does not require a cloned and overexpressed *R* gene to test the interaction.

We demonstrate that the positive defense reporter assay is suitable to test interactions directly in

a broad range of wheat cultivars encoding specific *R* genes in their genome. This will open

opportunities to screen *Avr* candidates directly on wheat-resistant cultivars, which was

previously considered not amenable using the cell-death based wheat protoplast assays (Saur et

al., 2019). Secondly, combining (generic) *Avr/R*-inducible promoters with the dual-reporter

system allows for generating normalized positive signals, which enables a much more sensitive

screening approach by greatly reducing variability between replicates. Furthermore, the high

sensitivity of the assay allows to use 10-20 times lower amount of plasmid for transfections than

Wilson et al., 2023

the former wheat protoplast assays, making the assay cost-effective, and easy to handle and

prepare a large list of *Avr* candidates for screening. Thirdly, this approach unlocks screening

multiple *Avr* candidates in a single transfection, which saves time and resources. Lastly, our new

wheat protoplast reporter assay enables us to study a wide range of plant defense signaling

directly in wheat. This is highlighted by our RNAseq analysis of defense signaling in wheat

protoplasts. Our results show that genes responsible for calcium ion binding, cation binding and

calcium ion transmembrane transporter activity along with other defense-related genes are

specifically upregulated in response to both AvrSr35/Sr35 and AvrSr50/Sr50 interactions in two

wheat cultivars. In line with our findings, there have been strong indications that plant defense

signaling triggered by Avr/R interactions involves plasma membrane reorganization including

formation of ion channels that regulate plant cell-death and disease resistance. Recent studies of

Arabidopsis ZAR1 (Bi et al., 2021) and wheat Sr35 (Förderer et al., 2022) showed the activated

ZAR1 and Sr35 proteins forming pentameric structures, so-called resistosomes, that appear to

function as non-selective calcium channels responsible for Ca^{2+} uptake, leading to cell-death.

Similarly, in *Arabidopsis*, an activated NRG1.1 protein, known as a downstream signaling

component of plant *R* proteins, compose a Ca^{2+} permeable non-selective cation channel enabled

by oligomerization, which controls cell Ca^{2+} influx and induces plant cell-death (Jacob et al.,

2021). Therefore, our findings support that the channel activity, especially Ca^{2+} regulating

channels, are conserved in broader Avr/R-induced downstream signaling.

Interestingly, one of our marker genes (the gene of promoter D14) encodes a predicted wheat

NDR1/HIN1-like 10 (NHL10) protein. The NHL proteins that are grouped in a large gene family

in *Arabidopsis thaliana* involved in plant defense, such as disease resistance (Aarts et al., 1998;

Century et al., 1995) and pathogen sensing (Boudsocq et al., 2010; Zheng et al., 2004; Zipfel et

al., 2004). The NHL proteins have one or two transmembrane domain(s) (Zheng et al., 2004) and

several of them were shown to localize to the plasma membrane (Selote et al., 2014; Coppinger

et al., 2004; Dagvadorj et al., 2022), which is thought to be crucial for their functioning in

disease resistance pathways. Moreover, *Arabidopsis NHL10* expression, which is highly induced

by flg22, is reported to be involved in Ca^{2+} signaling through the activation of calcium-

dependent protein kinases (Boudsocq et al., 2010). In a recent study, wheat blotch disease

effector ToxA was shown to interact with a wheat NHL protein, TaNHL10, to promote cell-

death mediated by intracellular resistance like protein Tsn1 (Dagvadorj et al., 2022). Future

studies will benefit from our defense specific promoters, in particular D14, to investigate
downstream defense signaling components in wheat.

Future studies will determine the capability of our positive defense reporter based Avr screening
assay. Although we have performed exemplified test for ten and fifty *Avr* candidates screening in
this study, we anticipate that the high sensitivity of the assay combined with self-replicative
pWDV1 plasmid can allow much higher number of *Avr* candidate screening in one wheat
protoplast transfection. Moreover, we anticipate that the *pWDV1* plasmid will simplify
overexpression of transient gene expression in wheat requiring only miniprep plasmid samples
(Supplemental Figure 2), which facilitate functional gene and variant analysis directly in wheat
protoplasts.

Our assay can be combined with a recent report that describes highly multiplexed avirulence
effector recognition screening using cell-death as a negative enrichment method (Arndell et al.,
2023). Multiplexed effector libraries (n>600) of *Pgt* were co-transfected into cv. Fielder
protoplasts overexpressing specific *Sr* genes from plasmids. The author used the overexpression
of YFP (Yellow Fluorescent Protein) as a cell viability marker to separate live protoplast from
dead protoplast via flow cytometry. This allowed for negative enrichment analysis of expressed
effectors when comparing live protoplast before and after flow cytometry via RNAseq analysis.
The author could identify new matching *Avr* genes for two out of five *Sr* genes including
AvrSr13/Sr13 and *AvrSr22/Sr22* but not for *Sr21*, *Sr26*, and *Sr61*. Such large scale avirulence
effector screening approaches will be much more efficient when combined with our positive
reporter assays that will allow researchers to first identify avirulence effector pools that are
recognized by a specific wheat cultivar. In addition, further improvements of the reported
negative enrichment screen will likely enable screening in wheat cultivars expressing *R* genes
endogenously similar to our assay. Hence the combination of both multiplexed avirulence
effector screening approaches promises to usher in a new area of avirulence gene identification
of economically important pathogens of wheat and other crop species.

Methods

Protoplast Isolation

Protoplast isolation and plant growth protocol are described here:
https://dx.doi.org/10.17504/protocols.io.q26g7r3zkvwz/v1. Wheat plants of cv. Fielder, Gabo
and GaboSr50 were grown for 7-9 days in a growth cabinet with 150 µmol/m^2/s light intensity
and at 21 ^{o}C with 16 hr day length. Protoplasts were isolated by making a shallow cut across the
leaf epidermis and peeling back the epidermal layer to expose the mesophyll layer beneath. The
cut segments were placed, peeled side down in 0.6 M Mannitol for 10 mins, then placed into a 7
cm diameter petri dish containing filtered enzyme solution (MES-pH5.7 (20 mM), Mannitol (0.6
M), KCI (10 mM), Cellulase (Yakult R-10) 1.5% w/v, Macerozyme (Yakult R-10) 0.75% w/v,
CaCl$_2$(10 mM). BSA 0.1% w/v). The dish was wrapped in aluminum foil to shield it from light
and placed on an orbital shaker, horizontally rotating at 60 RPM, for three hours. Protoplasts
were collected by gentle centrifugation at 100 x g for 3 mins in a 30 mL round-bottomed tube,
then the enzyme solution was removed and the pellet was washed in W5 solution (MES-pH5.7 (2
mM), KCI (5 mM), CaCl$_2$ (125 mM), NaCl (154 mM). 5 µL of concentrated protoplasts were
removed and used to quantify cells and check quality using a hemocytometer and light
microscope. Protoplasts were washed again with W5, this time left on ice for 45 minutes to
settle. W5 solution was removed, and protoplasts were resuspended to 300,000-350,000
protoplasts/mL using MMG (MES-pH5.7 (4 mM), Mannitol (0.4 mM), MgCl$_2$ (15 mM)).
Protoplasts were used immediately for transfection.

Protoplast transfection

High concentration (1 µg/µL) plasmid DNA was extracted via maxi-prep (Promega SV Wizard).
Lower quality plasmid DNA (e.g. from miniprep kits) typically resulted in low transfection rates.
The *Avr*, *R* and *luciferase* genes were expressed under control of maize ubiquitin promoter
(*UBI*), and 10 µg of each *Avr* and *R* plasmid and 20 µg of *luciferase* reporter plasmid were used
for each replicate. Due to high concentration and purity, plasmids were heated at 65^{o}C for 10 min
before use to reduce viscosity and ensure consistent measurement and delivery into cells. The
required plasmid DNA was added to 14 mL round-bottom culture tubes (Thermofisher,
#150268). 300 uL of protoplasts and 300 µL of PEG solution (PEG 4000 (40% w/v), Mannitol
(0.2 M), CaCl$_2$ (100 mM)) was then added and mixed immediately by gently flicking/rotating the
tube three times. Samples were incubated for 15 minutes, and the reaction halted with 900 µL
W5. Protoplasts were then collected by 100 x g centrifugation for 2 min, resuspended in 500 µL

modified W5 (W5 + 1 mM glucose) and transferred to a transparent, sterile 12 well culture plate

(Costar, #3513) whose wells were coated with 10% BSA to prevent sticking. Incubation was

carried out in light conditions, approximately 100 µmol/m^2/s. Wide bore or cut tips were used

throughout for handling protoplasts.

Protoplast Quantification

After the required incubation time (3, 4, 5, 6, and 18-hours for time course experiment, 4-18 hrs

for *pDefense-ELuc* reporters) protoplasts were prepared for luminescence measurement. 200 µL

of 1x cell lysis buffer (Promega #E3971) was added and samples vortexed and incubated for 15

min. Samples were centrifuged again to collect cell debris and 50 µL of lysate was added to an

opaque, flat-bottom, white, solid polystyrene 96 well plate (e.g. Corning® 96 well NBS™

Microplate #CLS3600). Plate was read for total luminescence (Tecan infinite 200Pro, 1000 ms

integration 0 ms settle) without addition of substrate, then 50 µL Steady-Glo (Promega #E2520)

Luciferin substrate was added, and the plate was read again.

Protoplast RNAseq sampling

Protoplasts were prepared as described above, using 315,00 protoplasts in 900 µL MMG per

replicate, with 4 replicates for each treatment group The *UBI-fLUC* reporter was excluded.

Additional replicates of each condition included the reporter to confirm cell-death after 16hrs

(Data not shown). At 4 hrs post transfection the protoplasts were pelleted, snap-frozen, and

stored at -80 until RNA isolation. Samples from each cultivar were prepared and processed in

batches. Qiagen RNeasy Plant Mini Kit was used as directed, (#74904) omitting the first

mechanical cell lysis step as freezing and addition of RLT buffer was sufficient to lyse cells.

Qiagen RNase-Free DNase Set (#79254) was used as directed. RNA was checked for absence of

DNA contamination by PCR and gel electrophoresis and quantitated using Qubit and Nanodrop.

Total RNA Illumina sequencing was performed on the NovaSeq platform, in a 2x150bp paired-

end configuration, with poly A selection & strand specific library preparation for the 36 samples.

The resulting sequence data is deposited in the NCBI Sequence Read Archive (SRA) under

BioProject number PRJNA957082.

RNAseq Analysis

Transcript abundance was quantified with kallisto (Bray et al., 2016) using the Ensembl *Triticum*

aestivum transcriptome (IWGSC, INSDC Assembly GCA_900519105.1, Jul 2018) plus cDNA

sequences for *AvrSr50, Sr50, AvrSr35* and *Sr35* as reference. DESeq2 (Love et al., 2014) was

used to test for differential expression. We defined upregulation at a 4-fold change or greater.

GaboSr50 with AvrSr50 was compared against the four control treatment groups;

Gabo_AvrSr50, Gabo_EV, GaboSr50_AvrSr35 and GaboSr50_EV. Then, the intersect of the

upregulated genes was taken, for a total of 272 genes. For Fielder, Fielder_Sr35_AvrSr35 was

compared to the 3 control treatment groups; Fielder_AvrSr35, FielderSr35_AvrSr50,

Fielder_Sr35. Again, taking the intersect of the upregulated genes returned 98 upregulated genes.

66 genes were commonly upregulated between both lists (Zenodo Supplemental Data S5). For

GO terms analysis, we repeated the comparisons above with lower upregulation of 2-fold change

and used the gProfiler webtool and Ensembl *Triticum aestivum* annotation, running the g:GOst

program (Raudvere et al., 2019). The recommended g:SCS multiple testing correction algorithm

was used.

pDefense Promoter Selection

Expression in log fold change and transcripts per million (TPM) was visualized for the 66

commonly upregulated genes, present in both 2-fold upregulated lists used for GO analysis.

Thirteen genes were selected based on their overall expression and upregulation in *Avr/R*

treatment groups. The entire gene and region were compared in Geneious and 600-800 bp

upstream of the 13 genes was selected, including any annotated 5'UTR. These promoter

sequences then were synthesized with 5'and 3' 4bp MoClo promoter overhangs flanked by BsaI

recognition sites for Golden Gate cloning as promoter+5'UTR standard parts (Weber et al.,

2011). Any BpiI and BsaI enzyme recognition sites within the promoter sequence were removed

prior to synthesis. Promoters that could not be synthesized were cloned by PCR, using primers

(Zenodo Supplemental Data S6) with BpiI recognition sites and 4bp overhang to facilitate

assembly into universal level 0 vector *pAGM9121* (Addgene Plasmid #51833). Any BsaI

recognition sites were removed as per Grutzner & Marillionnet, (2020).

pDefense Plasmid Constructs

Plasmids were designed in Geneious. The plasmid constructs were designed according to the

MoClo syntax, assembled by Golden Gate cloning and verified by whole plasmid sequencing.

Supplemental Data (Zenodo Supplemental Data S6) and Plasmid maps Zenodo (DOI:

10.5281/zenodo.7844465, link: https://zenodo.org/record/7844465) are available, with all

generated plasmids deposited on Addgene.

pWDV1 vector

The wheat dwarf virus genomic sequence from the isolate Enkoping1 (GenBank: AJ311031.1)

was modified to remove the V1 and V2 genes which encode the viral coat and transfer proteins,

respectively. In addition, one and two BsaI type II restriction sites were mutated to enable

downstream golden gate cloning steps. All mutations were in the Rep and RepA open reading

frames and were silent, with the exception that the mutation generated at amino acid position 263

for RepA results in a Serine to Threonine substitution at amino acid position 235 of the Rep

protein which is encoded by an alternately spliced transcript from the same Rep/RepA open

reading frame. Two BsaI sites were inserted between the Long and short intergenic regions in

place of the removed V1 and V2 genes. The BsaI sites were designed to generate a GGAG

overhang downstream of the long intergenic region and a CGCT overhang upstream of the short

intergenic region to facilitate level 1 golden gate assemblies as described previously (Engler et

al., 2014). BsmBI sites were also included at either end of the modified viral genome sequence to

facilitate golden gate cloning into the pYTK001 vector (Lee et al., 2015) which was chosen for

its small size, ease of cloning and to facilitate selection and plasmid replication in bacteria. The

modified wheat dwarf virus sequence, called WDV_1, was synthesised as a 'gene' fragment by

twist bioscience and directly incorporated into the pYTK001 vector by golden gate cloning using

BsmBI. We sequence verified plasmids via whole plasmid sequencing. We named the resulting

plasmid *pWDV1*. We used *pWDV1* for subsequent cloning of genes by golden gate cloning using

the BsaI entry site.

Ratiometric Assay

All isolation and transfection steps were as above aside from the substitution of the fLUC

reporter for the reporter (*UBI-pDefense)* and normalization (*UBI-pRedf*) plasmids. These require

the same substrate for activation, but express luciferase with different emission spectra, read by

specific green and red filters in the luminescence plate reader.A *UBI-ELuc* construct was also

generated to calibrate normalization and deconvolution calculations, necessary to resolve the

overlap in emission spectra of the two luciferase reporters (González-Grandío et al., 2021).

These calibration experiments were performed separately three times and with three replicates
per measurement, according to the instructions for Promega Chroma-Glo assay (Chroma-
Glo(TM) Luciferase Assay System Technical Manual, TM062). The plate reader was
programmed to measure total luciferase, then with green and red filters, with 1000 ms integration
time and 0 ms settle time, measured well-wise. Template spreadsheets for performing
calculations and a Python script for automating deconvolution and plotting are recorded on
protocols.io (https://dx.doi.org/10.17504/protocols.io.q26g7r3zkvwz/v1).

Cultivar testing

Protoplasts were isolated from nine different wheat cultivars (GaboSr50, Coorong, C6969,
Clement, Compair, HeinesPeko, KenyaW1483, AvocetYr8 and Fielder), using the method
described above. All were grown in the same conditions as above for 8 days. Where the cultivar
carried an *R* gene (GaboSr50, Coorong, C6969) the recognized *Avr* (*AvrSr50*, *AvrSr27* and
AvrSr35 respectively) was transfected alone, with an avr (unrecognized by R protein) as control.
In the other six cultivars both the *Avr* and *R* gene were co-transfected. Normalized luminescence
was calculated from the ratio of deconvoluted *pDefense-Eluc* to *UBI-pRedf*.

Multiplex screening

493 Protoplasts were prepared from cv. GaboSr50 and cv. Gabo wheat leaves and transfected as
494 above. Measurements used the ratiometric reporters and measurements, with pDefense14-ELuc
495 as reporter and UBI-pRedf for normalisation. Avr pools excluding the AvrSr50 acted as negative
496 controls, and 1 µg of pWDV1-AvrSr50 alone acted as a positive control. Data processing and
497 statistics was carried out as above.

Data availability

All supplemental data is available on Zenodo (DOI: 10.5281/zenodo.7844446., link:
https://doi.org/10.5281/zenodo.7844446 All plasmids generated will be made available from
Addgene, with plasmid maps deposited on Zenodo (DOI: 10.5281/zenodo.7844464, link:
https://doi.org/10.5281/zenodo.7844464). Code used to perform RNAseq analysis is deposited
on Github (https://github.com/ritatam/wheat-protoplast-RNAseq-DESeq2) and all RNAseq data
is available on NCBI Sequence Read Archive (SRA) under BioProject number PRJNA957082.

Acknowledgements

We thank Prof. Robert Park for providing seeds of cv. C6969, A/Prof. Sambasivam Periyannan for providing seeds of cv. Fielder, cv. Gabo, cv. GaboSr50, and cv. Coorong, Dr. Michael Ayliffe and Dr. Peter Dodds for providing *pUBI-AvrSr35*, *pUBI-Sr35*, *pUBI-AvrSr50*, and *pUBI-Sr50*. B.S. acknowledges the financial support by the Australian Research Council via a Future Fellowship (FT180100024). This work was supported by computational resources provided by the Australian Government through the National Computational Infrastructure (NCI) under the ANU Merit Allocation Scheme.

Author Contributions

S. W., B. D., J. P. R., and B. S. conceived the project. S. W., B. D., R. T., L. M., N. H., E. C., and J. G. performed the experiments, S. W., B. D., and B. S. wrote the manuscript. R. T., E. C., and J. P. R. provided extensive feedback on the manuscript. B. S. secured funding. All authors read and approved the manuscript.

References

Aarts, N., Metz, M., Holub, E., Staskawicz, B.J., Daniels, M.J., and Parker, J.E. (1998). Different requirements for EDS1 and NDR1 by disease resistance genes define at least two R gene-mediated signaling pathways in Arabidopsis. Proceedings of the National Academy of Sciences **95**: 10306–10311.

Arndell, T. et al. (2023). Pooled effector library screening in protoplasts rapidly identifies novel Avr genes.: 2023.04.28.538616.

Bi, G. et al. (2021). The ZAR1 resistosome is a calcium-permeable channel triggering plant immune signaling. Cell **184**: 3528-3541.e12.

Boudsocq, M., Willmann, M.R., McCormack, M., Lee, H., Shan, L., He, P., Bush, J., Cheng, S.-H., and Sheen, J. (2010). Differential innate immune signalling via Ca2+ sensor protein kinases. Nature **464**: 418–422.

Bourras, S. et al. (2019). The AvrPm3-Pm3 effector-NLR interactions control both race-specific resistance and host-specificity of cereal mildews on wheat. Nat Commun **10**: 2292.

Bouton, C., King, R.C., Chen, H., Azhakanandam, K., Bieri, S., Hammond-Kosack, K.E.,
and Kanyuka, K. (2018). Foxtail mosaic virus: A Viral Vector for Protein Expression in
Cereals. Plant Physiology **177**: 1352–1367.

Bray, N.L., Pimentel, H., Melsted, P., and Pachter, L. (2016). Near-optimal probabilistic
RNA-seq quantification. Nat Biotechnol **34**: 525–527.

Century, K.S., Holub, E.B., and Staskawicz, B.J. (1995). NDR1, a locus of Arabidopsis
thaliana that is required for disease resistance to both a bacterial and a fungal pathogen.
Proceedings of the National Academy of Sciences **92**: 6597–6601.

Chen, J. et al. (2017). Loss of *AvrSr50* by somatic exchange in stem rust leads to virulence for
Sr50 resistance in wheat. Science **358**: 1607–1610.

Christensen, A.H., Sharrock, R.A., and Quail, P.H. (1992). Maize polyubiquitin genes:
structure, thermal perturbation of expression and transcript splicing, and promoter
activity following transfer to protoplasts by electroporation. Plant Mol Biol **18**: 675–689.

Coppinger, P., Repetti, P.P., Day, B., Dahlbeck, D., Mehlert, A., and Staskawicz, B.J.
(2004). Overexpression of the plasma membrane-localized NDR1 protein results in
enhanced bacterial disease resistance in Arabidopsis thaliana. The Plant Journal **40**: 225–
237.

Dagvadorj, B., Outram, M.A., Williams, S.J., and Solomon, P.S. (2022). The necrotrophic
effector ToxA from Parastagonospora nodorum interacts with wheat NHL proteins to
facilitate Tsn1-mediated necrosis. The Plant Journal **110**: 407–418.

Dodds, P.N. and Rathjen, J.P. (2010). Plant immunity: towards an integrated view of plant–
pathogen interactions. Nat Rev Genet **11**: 539–548.

Dracatos, P.M., Zhang, P., Park, R.F., McIntosh, R.A., and Wellings, C.R. (2016).
Complementary resistance genes in wheat selection "Avocet R" confer resistance to
stripe rust. Theor Appl Genet **129**: 65–76.

Engler, C., Youles, M., Gruetzner, R., Ehnert, T.-M., Werner, S., Jones, J.D.G., Patron,
N.J., and Marillonnet, S. (2014). A Golden Gate Modular Cloning Toolbox for Plants.
ACS Synth. Biol. **3**: 839–843.

Förderer, A. et al. (2022). A wheat resistosome defines common principles of immune receptor
channels. Nature **610**: 532–539.

Gil-Humanes, J., Wang, Y., Liang, Z., Shan, Q., Ozuna, C.V., Sánchez-León, S., Baltes,
N.J., Starker, C., Barro, F., Gao, C., and Voytas, D.F. (2017). High-efficiency gene
targeting in hexaploid wheat using DNA replicons and CRISPR/Cas9. The Plant Journal
89: 1251–1262.

González-Grandío, E., Demirer, G.S., Ma, W., Brady, S., and Landry, M.P. (2021). A
Ratiometric Dual Color Luciferase Reporter for Fast Characterization of Transcriptional
Regulatory Elements in Plants. ACS Synth. Biol. **10**: 2763–2766.

Hewitt, T. et al. (2021). A highly differentiated region of wheat chromosome 7AL encodes a
Pm1a immune receptor that recognizes its corresponding AvrPm1a effector from
Blumeria graminis. New Phytol **229**: 2812–2826.

Jacob, P. et al. (2021). Plant "helper" immune receptors are Ca $^{2+}$ -permeable nonselective
cation channels. Science **373**: 420–425.

Jensen, C. and Saunders, D.G.O. (2023). Magnaporthe oryzae pathotype Triticum (MoT) can
act as a heterologous expression system for fungal effectors with high transcript
abundance in wheat. Sci Rep **13**: 108.

Laflamme, B., Dillon, M.M., Martel, A., Almeida, R.N.D., Desveaux, D., and Guttman, D.S.
(2020). The pan-genome effector-triggered immunity landscape of a host-pathogen
interaction. Science **367**: 763–768.

Lee, M.E., DeLoache, W.C., Cervantes, B., and Dueber, J.E. (2015). A Highly Characterized
Yeast Toolkit for Modular, Multipart Assembly. ACS Synth Biol **4**: 975–986.

Lee, W.-S., Hammond-Kosack, K.E., and Kanyuka, K. (2012). Barley Stripe Mosaic Virus-
Mediated Tools for Investigating Gene Function in Cereal Plants and Their Pathogens:
Virus-Induced Gene Silencing, Host-Mediated Gene Silencing, and Virus-Mediated
Overexpression of Heterologous Protein. Plant Physiology **160**: 582–590.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biology **15**: 550.

Luo, M. et al. (2021). A five-transgene cassette confers broad-spectrum resistance to a fungal
rust pathogen in wheat. Nat Biotechnol **39**: 561–566.

Mago, R. et al. (2015). The wheat Sr50 gene reveals rich diversity at a cereal disease resistance
locus. Nat Plants **1**: 15186.

McIntosh, R., Dyck, P., The, T., Cusick, J., and Milne, D. (1984). Cytogenetical studies in
wheat XIII. Sr35 - a third gene from Triticum monococcum for resistance to Puccinia
graminis tritici. Zeitschrift für Pflazenzüchtung **92**: 1–14.

McIntosh, R., Wellings, C., and Park, R. (1995). Wheat Rusts: An Atlas of Resistance Genes.
In (CSIRO Publishing).

**Meile, L., Croll, D., Brunner, P.C., Plissonneau, C., Hartmann, F.E., McDonald, B.A., and
Sánchez-Vallet, A.** (2018). A fungal avirulence factor encoded in a highly plastic
genomic region triggers partial resistance to septoria tritici blotch. New Phytologist **219**:
1048–1061.

Müller, M.C. et al. (2022). Ancient variation of the AvrPm17 gene in powdery mildew limits
the effectiveness of the introgressed rye Pm17 resistance gene in wheat. Proceedings of
the National Academy of Sciences **119**: e2108808119.

**Ortiz, D., Chen, J., Outram, M.A., Saur, I.M.L., Upadhyaya, N.M., Mago, R., Ericsson,
D.J., Cesari, S., Chen, C., Williams, S.J., and Dodds, P.N.** (2022). The stem rust
effector protein AvrSr50 escapes Sr50 recognition by a substitution in a single surface-
exposed residue. New Phytol **234**: 592–606.

Praz, C.R. et al. (2017). AvrPm2 encodes an RNase-like avirulence effector which is conserved
in the two different specialized forms of wheat and rye powdery mildew fungus. New
Phytol **213**: 1301–1314.

Raudvere, U., Kolberg, L., Kuzmin, I., Arak, T., Adler, P., Peterson, H., and Vilo, J. (2019).
g:Profiler: a web server for functional enrichment analysis and conversions of gene lists
(2019 update). Nucleic Acids Research **47**: W191–W198.

**Saintenac, C., Zhang, W., Salcedo, A., Rouse, M.N., Trick, H.N., Akhunov, E., and
Dubcovsky, J.** (2013). Identification of wheat gene Sr35 that confers resistance to Ug99
stem rust race group. Science **341**: 783–786.

Salcedo, A. et al. (2017). Variation in the *AvrSr35* gene determines *Sr35* resistance against
wheat stem rust race Ug99. Science **358**: 1604–1606.

**Sarrion-Perdigones, A., Chang, L., Gonzalez, Y., Gallego-Flores, T., Young, D.W., and
Venken, K.J.T.** (2019). Examining multiple cellular pathways at once using multiplex
hextuple luciferase assaying. Nat Commun **10**: 5710.

Saur, I.M.L., Bauer, S., Lu, X., and Schulze-Lefert, P. (2019). A cell death assay in barley
and wheat protoplasts for identification and validation of matching pathogen AVR
effector and plant NLR immune receptors. Plant Methods **15**: 118.

Savary, S., Willocquet, L., Pethybridge, S.J., Esker, P., McRoberts, N., and Nelson, A.
(2019). The global burden of pathogens and pests on major food crops. Nat Ecol Evol **3**:
430–439.

Selote, D., Shine, M.B., Robin, G.P., and Kachroo, A. (2014). Soybean NDR1-like proteins
bind pathogen effectors and regulate resistance signaling. New Phytologist **202**: 485–498.

Upadhyaya, N.M. et al. (2021). Genomics accelerated isolation of a new stem rust avirulence
gene–wheat resistance gene pair. Nat. Plants **7**: 1220–1228.

Weber, E., Gruetzner, R., Werner, S., Engler, C., and Marillonnet, S. (2011). Assembly of
Designer TAL Effectors by Golden Gate Cloning. PLOS ONE **6**: e19722.

Wulff, B.B. and Krattinger, S.G. (2022). The long road to engineering durable disease
resistance in wheat. Current Opinion in Biotechnology **73**: 270–275.

Zheng, M.S., Takahashi, H., Miyazaki, A., Hamamoto, H., Shah, J., Yamaguchi, I., and
Kusano, T. (2004). Up-regulation of Arabidopsis thaliana NHL10 in the hypersensitive
response to Cucumber mosaic virus infection and in senescing leaves is controlled by
signalling pathways that differ in salicylate involvement. Planta **218**: 740–750.

Zhong, Z. et al. (2017). A small secreted protein in Zymoseptoria tritici is responsible for
avirulence on wheat cultivars carrying the Stb6 resistance gene. New Phytologist **214**:
619–631.

Zipfel, C., Robatzek, S., Navarro, L., Oakeley, E.J., Jones, J.D.G., Felix, G., and Boller, T.
(2004). Bacterial disease resistance in Arabidopsis through flagellin perception. Nature
428: 764–767.

Figure Legends

Figure 1: Rapid onset of cell-death downstream of *Avr/R* interactions in protoplasts. **A-C**,

Luciferase activity was measured at defined timepoints in hours post-transfection (hpt) in wheat

protoplasts. **(A)** Shows boxplots of cv. Fielder protoplasts expressing *luciferase* alone, and

luciferase with either *Sr35, AvrSr35,* or *Sr35* plus *AvrSr35*. **(B)** shows boxplots of cv. GaboSr50

protoplasts expressing *luciferase* alone, and *luciferase* with either *AvrSr35* or *AvrSr50.* **(C)**

shows the luminescence of the Avr/R interaction treatment groups from cv. Fielder and cv.

GaboSr50 alone when compared to the no plasmid transfection control replicates (red line)

which is the average of nine replicate across the six timepoints.

Figure 2: Co-expression of *Avr/R* gene pairs triggers defense signaling in wheat protoplasts.

(A) Heat map shows expression of *Avr* and *R* genes by normalized gene count. Each column

represents a replicate and the treatment group is indicated by a number as shown in the legend. *R*

or *Avr* genes are expressed in their respective treatment groups only. This includes endogenously

expressed *Sr50* in cv. GaboSr50. **(B-D)** show principal component analysis (PCA) of the

RNAseq analysis; all 36 samples **(B)**, cv. Fielder **(C)** and cv. Gabo/GaboSr50 **(D)**. **(E)** shows the

GO terms of genes that are upregulated 2 fold in *Avr/R* treatment groups. A full list of

upregulated genes with overrepresented GO terms is recorded in supplemental data (Zenodo

Supplemental Data S4). **(F)** shows a heatmap of the expression of thirteen candidate reporter

genes (D1-7, D10-12, D14-16) that are commonly upregulated by *Sr50/AvrSr50* and

Sr35/AvrSr35 in cv. GaboSr50 or cv. Fielder, respectively. Each square represents the mean of

four replicates for treatment group.

Figure 3: *pDefense14* is a positive defense reporter in wheat protoplasts. (**A**) Wheat

protoplasts transfected with *pDefense14* and *UBI-pRedF* constructs show an increase in

normalized luminescence in the *Avr/R* treatment, compared to treatments with *R* or *avr* only and

without *R* or *Avr*. Panel (**B**) shows corresponding non-normalized red-filtered luminescence for

the treatments in panel A. Treatments marked with common letters were not significantly

different (P >0.05) assessed with one-way analysis of variance (ANOVA) and post hoc Tukey's

HSD test.

Figure 4: pWDV1 plasmid reduces amount of *Avr* plasmid required to detect defense

signaling using *pDefense* reporter. *AvrSr50* was cloned into *pWDV1* (*pWDV1-AvrSr50*) and

non-viral plasmid backbone (*pICH4772-UBI-AvrSr50)* and transfected in increasing amounts

(0.1, 0.5, 1, 5 and 15 µg of plasmid DNA) alongside *pDefense14* and *UBI-pRedf*. Reporters

remained in the non-replicative plasmid. Controls in cv. GaboSr50 were transfected with

pWDV1-AvrSr35 (1 and 15 µg) and cv. Gabo transfected with *pWDV1-AvrSr50* (1 µg) or

pWDV1-AvrSr35 (1 µg). One-way analysis of variance (ANOVA) and post hoc Tukey's HSD

test was used to assess differences within normalized luminescence (A) and red luminescence

(B) groups. Treatments marked with a common letter were not significantly different (P >0.05).

Figure 5: Positive pDefense reporter detect defense signaling triggered by three distinct

***Avr/R* pairs in wheat protoplasts expressing the *R* gene endogenously.** From left to right,

protoplasts from wheat cultivars Coorong carrying *Sr27*, C6969 carrying *Sr35* and GaboSr50

carrying *Sr50* were transfected with *UBI-pDefense14* and *AvrSr27, AvrSr35* and *AvrSr50*

constructs, respectively, and each wheat cv. was transfected with *avr* (unrecognized *Avr*)

construct and empty vector (EV) as indicated in the column headers and x-axis labels. *UBI-pRedf*

was used for normalization of all experiments. One-way analysis of variance (ANOVA) and post

hoc Tukey's HSD test was used to assess differences among normalized luminescence for each

cultivar. Treatments marked with a common letter were not significantly different (P >0.05).

Figure 6: The positive defense reporter *pDefence14* is broadly applicable to all tested wheat

cultivars. Each panel shows the normalized luminescence of the positive defense reporter

pDefense14. Protoplasts isolated from various wheat cultivars were transfected with both *R* and

Avr constructs, aside from cultivars GaboSr50, C6969 and Coorong which carry *Sr50, Sr35* and

Sr27 respectively. The cultivars Corack, Clement, Kenya W1483, Avocet Yr8 and Compair,

tested with *AvrSr50/Sr50* and Heines Peko was tested with *AvrSr27/Sr27.*An *avr* (unrecognised

by R protein) was used as a control and *UBI-pRedf* was used for normalization of all

experiments. One-way analysis of variance (ANOVA) and post hoc Tukey's HSD test was used

to assess differences among means of normalized luminescence measurements for each cultivar.

Treatments marked with common letters were not significantly different (P >0.05).

Figure 7: The positive defense reporter enables multiplexed *Avr* screening. Each panel in **(A)**

shows the normalized luminescence of the positive defense reporter *pDefense14*. Each panel in

(**B**) shows the red luminescence of *UBI-pRedf.* Pools exemplifying ten or fifty *Avr* candidates

were transfected into protoplasts isolated from cv. GaboSr50 (left) and cv. Gabo (right). The

pools include 1 µg of *pWDV1-AvrSr50*, and 4.5 µg of each *pWDV1-AvrSr35* and *pWDV1-*

AvrSr27 in the pool of ten, or 24.5 µg of each *pWDV1-AvrSr35* and *pWDV1-AvrSr27* in the pool

of fifty so that one µg of *Avr* plasmid represents one Avr candidate. The same *Avr* pools

excluding the *AvrSr50* are used as negative controls, and 1 µg of *pWDV1-AvrSr50* alone for

positive control. All replicates were co-transfected with *pD14* reporter *UBI-pRedf* for

normalization. One-way analysis of variance (ANOVA) and post hoc Tukey's HSD test was

used to assess differences among means of normalized luminescence and red luminescence

measurements for each treatment group. Treatments marked with common letters were not

significantly different (P >0.05).

Supplemental Figure 1: Preliminary successful screening of *pDefense2, -14* and *-15* defense

reporters. Normalized luminescence for *pDefense14,* (panel **A**)*, pDefense 15* (panel **B**) and

pDefense2 (panel **C**), in three wheat cultivars. Left to right, cv. C6969 which carries *Sr35*, cv.

GaboSr50 which carries *Sr50* and cv. Coorong, which carries *Sr27*. Avr+R treatments were

transfected with plasmid which expressed a recognized Avr, avr+R treatments expressed a non-

recognized avr and R alone expressed an empty vector. Reporters were normalized against *UBI-*

pRedf for each replicate. The three reporters produced an increase in normalized luminescence in

treatments with Avr/R interaction, across the three Avr/R pairs tested.

Supplemental Figure 2: *pDefense* reporter can detect lower purity *Avr* plasmid. In (**A**) both

maxi prepped (left) and mini prepped (right) *AvrSr35* plasmid was co-transfected with *R* and *avr*

plasmids, in protoplasts from the cultivar Fielder. **(B)** shows corresponding non-normalized red-

filtered luminescence. One-way analysis of variance (ANOVA) and post hoc Tukey's HSD test

was used to assess differences among means of normalized and red-filtered luminescence

measurements for each treatment. Treatments marked with common letters were not significantly

different ($P > 0.05$).**Supplemental Material Legends**

Supplemental Material S1: Luminescence data for Figure 1, protoplast time course experiment.

Supplemental Material S2: List of samples and treatment groups for RNAseq experiment.

Supplemental Material S3: Lists of Gabo/GaboSr50, Fielder and all cultivar genes upregulated

by 2-fold.

Supplemental Material S4: GO terms analysis Query tables and g:profiler results tables

Supplemental Material S5: Lists of Gabo/GaboSr50, Fielder and all cultivar genes upregulated

by 4-fold.

Supplemental Material S6: List of primers used.

Supplemental Material S7: List of plasmids used.

Supplemental Material S8: Luminescence data used for generating Figures 3 -7.

Supplemental Material S9: Statistical output for Figures 3-7.

Supplemental Material S10: Data and statistical output for supplemental figures 1 and 2.

A

GaboSr50

Gabo

B

GaboSr50

Gabo