Modelling complex growth profiles of *Bacteroides fragilis* and *Escherichia coli* on various carbohydrates in an anaerobic environment

Zachary McGuire,[1,2,3,+] Sainitya Revuru,[2,3,+] Sheng Zhang,[4,+] Amanda Blankenberger,[3,+] Moiz Rasheed,[3] Jacob D. Hosen,[5] Guang Lin,[4,6,7] Mohit S. Verma[2,3,8 *]

[1]College of Pharmacy, Purdue University, West Lafayette, IN, USA, 47907

[2]Birck Nanotechnology Center, Purdue University, West Lafayette, IN, USA, 47907

[3]Department of Agricultural and Biological Engineering, Purdue University, West Lafayette, IN, USA, 47907

[4]Department of Mathematics, Purdue University, West Lafayette, IN, USA, 47907

[5]Forestry and Natural Resources, Purdue University, West Lafayette, IN, USA, 47907

[6]Department of Statistics, Purdue University, West Lafayette, IN, USA, 47907

[7]School of Mechanical Engineering, Purdue University, West Lafayette, IN, USA, 47907

[8]Weldon School of Biomedical Engineering, Purdue University, West Lafayette, IN, USA, 47907

* To whom correspondence may be addressed. E-mail: msverma@purdue.edu

[+]Equal contribution

Abstract

Previously published models for microbial growth focus only on either death or growth and are unable to account for differently shaped growth curves. Currently, there is no model capable of incorporating combinations of microbial growth trends. This study creates a bacterial growth model that incorporates growth, death, lag, and tail phases as well as applies this model to the growth trends of *Bacteroides fragilis* and *Escherichia coli* on 13 different carbohydrate substances. Growth trends were collected by measuring the optical densities over 72 hours for either *B. fragilis* or *E. coli* in a chemically defined media supplemented by a mono- or disaccharide. The Digital Environment to Enable Data-driven Science (DEEDS) platform was utilized to parse data and apply the developed model to obtain parameter values. *E. coli* was found to grow on the chemically defined media alone while *B. fragilis* was unable to grow on it alone. *E. coli* growth was led by 10 mM concentration of substrates while *B. fragilis growth* was substrate dependent. Bacterial death only occurred for *B. fragilis* but was found to be dependent on concentration for the two most significant substrates. A singular model was developed that does not require prior knowledge of metabolomics and is capable of incorporating a combination of growth and death trends.

1. Introduction

The most widely used models for bacterial growth predict sigmoidal-shaped curves, which do not represent the entirety of the growth data observed in a closed system (since these include a death phase that is often ignored in growth models). In order to quantify a growth curve and perform statistical analysis on the influence of environmental conditions (e.g., different carbon sources), we need a model that can fit both growth and death phases without segmenting the data. Here, we present such a model by introducing a death rate term to the logistic growth model and use this model to quantify growth curves with and without death phases. We use this model to obtain and evaluate growth and death parameters in the presence of 13 different carbon sources at three concentrations each for two different bacteria: *Bacteroides fragilis* and *Escherichia coli* as seen in the workflow demonstrated in Fig. 1.

Published models in predictive microbiology tend to focus on either growth or death depending on the intended use in food safety and quality. Growth models focus on studies in food hazards and spoilage[1], while death models focus on attempts to kill bacteria in efforts to improve shelf life and food safety[2]. The death phase of the bacterial growth curve that occurs after bacteria reach their carrying capacity has largely been ignored. Since most of the work in this field has applications in food microbiology, the death phase may have been ignored in modelling as food cannot be unspoiled. Furthermore, not all experimental bacterial growth data will show a death phase, depending on the length of the experiment and the environmental conditions. The stationary phase of bacterial growth can be maintained for extended periods of time. The amount of time spent in each phase of the bacterial growth curve is dependent on the species of the bacteria as well as the environment, so the length of the stationary phase or the time until rapid death will occur is difficult to predict *a priori*[3].

Many models for non-sigmoidal growth require additional inputs which may not be available

to the researcher or possible to obtain given the bacterium of interest. Published literature has the

following three limitations: i) cybernetic models require growth studies of individual substrates

in minimal media[4] (which prohibits growth of certain bacteria because they may not have the

necessary micronutrients in minimal media), ii) diauxic models for biphasic growth require

knowledge of enzyme kinetics[5], iii) many models are designed for specific trends observed in

growth, e.g., sigmoidal[6,7], biphasic[4,5], death[2,7], death then recovery growth[7]. A comparison of the

published models with the model in this study is shown in Table 1. [8] [9] [10] [11]

To our knowledge there are limited models that can be used to model any combination of the

trends listed above. A model is necessary that is versatile and can fit many different trends in

bacterial growth while providing meaningful parameters that can be statistically analyzed. Here,

we present a growth model that can fit variable growth profiles, both with and without a death

phase. We apply this model to several different trends observed in the growth of *Bacteroides*

fragilis and *Escherichia coli* on 13 different substrates to demonstrate the versatility of the model

in fitting many different plot shapes.

2. Materials and Methods

2.1 Bacterial strains and culture

The bacterial strains used in this study include *Bacteroides fragilis* (ATCC 43858) and

Escherichia coli (LF82 strain obtained from Professor Ramnik Xavier's Lab). Frozen secondary

stocks of the bacteria were prepared (1 mL of culture and 1 mL of 50% glycerol in a 2 mL

cryovial) and stored in -80 °C. *B. fragilis* is cultured in commercially available chopped meat

broth (Pre-Reduced II Hungate Cap, BD BBL™) and *E. coli* LF82 is cultured in Tryptic Soy

Broth (BD Biosciences). Both the bacteria are incubated at 37 °C and allowed to grow under

anaerobic conditions for two days with a gas composition of 4% H_2, 5% CO_2, and 91% N_2. The

cells are incubated in a Heated Vinyl Anaerobic Customized Type B Chamber with a hydrogen

sulfide removal column (Coy Laboratory Products Inc., Grass Lake, MI).

After incubating for two days, the cultures are tightly sealed and centrifuged at 10,000 rpm

for 5 minutes. The supernatant is discarded and the pellets are suspended in required volumes of

phosphate-buffered saline (PBS, 137mM NaCl, 2.7mM KCl, 10 mM Na_2HPO_4, 1.8mM KH_2PO_4,

pH 7.4) in anaerobic conditions to yield OD_{600} of ~0.5 as measured in a Corning® 96-well clear

flat bottom polystyrene microplate (Catalog #3370, Corning®, Corning, NY) by pipetting 250

µL and using a Biotek® Epoch™ 2 multi-well plate reader (Biotek Instruments, Winooski, VT).

2.2 Defined media preparation

In this study, the bacteria are allowed to grow in a modified chemically defined base medium

ZMB1[12] (Table S1) supplemented with different carbohydrates (Fig. 2). ZMB1 is chosen as the

base medium because it supports high cell density growth of the selected bacteria[12]. Detailed

modified ZMB1 preparation information can be found in Table S1.

Carbohydrate (Fig. S1) stocks are prepared in concentrations of 111 mM as exemplified in

Table S2. The solution is then filtered using a Steriflip® Vacuum-driven Filtration System

(Catalog # SCGP00525, EMD Millipore Corporation, Burlington, MA). 111 mM carbohydrate

stocks were then serially diluted using two 10x dilutions into sterile modified ZMB1 to create

solutions at concentrations of 11.1 mM and 1.11 mM. If necessary, the pH of the solution is

adjusted to a value of 7.0 using 10N NaOH.

2.3 Experimental design and analysis

The experiments are carried out in a Corning® 96-well clear flat bottom polystyrene

microplate (Catalog #3370, Corning®, Corning, NY) to observe the growth of *B. fragilis* and *E.*

coli LF82 in all 13 carbohydrates over time (72 hrs) separately. For every set of growth

condition, eight technical replicates are used. 225 µL of prepared media (modified ZMB1 and

carbohydrates) are added into the microtiter plates. 25 µL of the inoculum (bacteria in PBS) is

then transferred into the wells containing the prepared media and covered with 50 µL sterile

mineral oil (Catalog #S25439, Fisher Science Education, Nazareth, PA) to prevent evaporation .

This dilution yields final experimental concentrations of 100 mM, 10 mM, and 1 mM for each

condition. Two blanks (modified ZMB1 and PBS sans bacteria) and negative controls (bacteria

in PBS and ZMB1 sans carbohydrates) of the same quantities are used for comparison (Fig. S2).

Optical density of the microplates is measured at 600 nm (OD_{600}) over 72 hours at an interval

of 30 mins using Epoch 2 Microplate Spectrophotometer (BioTek Instruments, Inc., Winooski,

VT) with the Corning® 96-well clear flat bottom polystyrene microplate #3370. Bacterial growth

is estimated via accumulation of biomass by subtracting the average of the blanks from each

experimental condition from the average of the technical replicates for that same condition. The

reading process for multiple plates is automated using a BioStack™ 4 microplate stacker

(BioTek Instruments, Inc., Winooski, VT), which cycles the experimental 96-well microplates

and removes the lids at the specified interval as to not interfere with spectrophotometric values.

After the Epoch 2 Microplate Spectrophotometer is finished reading each plate, the BioStack™ 4

microplate stacker returns the lid to the microplate and places the microplate back into the

stacker to wait for the next reading interval to begin.

2.4 Computational Model

According to the logistic model of population growth, we have

$$\frac{dY(t)}{dt} = r \cdot N(t) \cdot (1 - \frac{N(t)}{K})$$ (1)

128 where Y is the population size, K is the carrying capacity, and r is the per capita growth rate. In

129 our setting, the carrying capacity decreases over time as the resource is consumed. We use time-

130 dependent carrying capacity as follows:

$$\frac{dY(t)}{dt} = r \cdot Y(t) \cdot \left(1 - \frac{Y(t)}{K(t)}\right)$$ (2)

131 Denote t' = argmax_t Y(t). Since $\frac{dY(t)}{dt}$=0 when t=t', we have

$$0 = r \cdot Y(t') \cdot \left(1 - \frac{Y(t')}{K(t')}\right)$$ (3)

132 or

$$K(t') = Y(t')$$ (4)

133 Suppose the carrying capacity decreases over time in the rate of d. In other words,

$$K(t) - K(t') = -d \cdot (t - t')$$ (5)

134 We have

$$K(t) = Y(t') - d \cdot (t - t')$$ (6)

135 Substitute Eq. (5) into Eq. (2), and we have

$$\frac{dY(t)}{dt} = r \cdot Y(t) \cdot \left(1 - \frac{Y(t)}{Y(t') - d \cdot (t - t')}\right)$$ (7)

136 Given each dataset, we approximate dY(t)/dt numerically, and then obtain the growth rate

137 and the carrying capacity decrease rate for the dataset by solving the following optimization

138 problem:

$$R, D = argmin\_\{r, d\} \left(\frac{dY(t)}{dt} = r \cdot N(t) \cdot (1 - \frac{Y(t)}{Y(t') - d \cdot (t - t')}) \right)^2 \qquad (8)$$

2.5 DEEDS Workflow

The platform Digital Environment for Enabling Data-Driven Science (DEEDS) was used to

analyze the data presented in this paper. The raw data from the Gen5™ software (BioTek

Instruments, Inc.) was uploaded here. Two codes were ran in series to generate the parameter

data presented in this paper.

The first Python code (parsing.py) uses two files for input: the Gen5™ software's Microsoft

Excel document and a separate Microsoft Excel document detailing the experimental conditions

of each well of each plate in an experiment. The code then parses, labels, and subtracts the

carbohydrate blanks for all experimental conditions. The code creates a separate .csv file for each

experimental condition on an experiment. See Supporting Information for the code.

The second Python code (modelFit.py) uses the parsed .csv files from the previous code to

analyze the experimental data used in the model presented above. See Supporting Information

for the code.

Since this work was initially completed, the DEEDS platform is no longer available or

managed. Thus, we are providing the python code in supporting information to help replicate the

results reported here.

3.   Results and Discussion

3.1 Selection of carbohydrates for growth curves

A total of 13 carbohydrates are chosen to observe the growth of *B. fragilis* and *E. coli* LF82

in this study (Fig. 2A and B). D-glucose, a simple sugar, is a common carbon source for most

bacteria. We wanted to analyze the growth of bacteria in both mono- and di-saccharides - 1) by

including isomers of D-glucose (D-fructose and D-galactose), 2) by removing and adding

functional groups (L-fucose, D-glucosamine, N-acetyl-D-glucosamine), 3) by including sugars

that are a resultant of different glycosidic linkages between two glucose molecules and finally, 4)

by including di-saccharides that are a resultant of linkages between two different

monosaccharides (Fig. S1).

3.2 Optical density growth curves for *B. fragilis*

The growth of *B. fragilis* in ZMB1 media with individual carbohydrate sources varies per

source and concentration of sugar. *B. fragilis* did not grow in ZMB1 without a sugar present.

Like *E. coli*, *B. fragilis* has been reported to utilize amino acids for growth but in studies with

media consisting of only salts, minerals, and amino acids there was only a small amount of initial

growth[13]. Most notably, *B. fragilis* demonstrated a lag phase on most growth curves. It has been

hypothesized previously that at CO_2 levels like the 5% used in these experiments is not high

enough for the bacteria to be able to grow[14]. So in the lag phase, the bacteria are creating more

CO_2 until a level is reached in which the bacteria are able to replicate[14].

B. fragilis had significant growth rates on all sugars except trehalose and cellobiose. The

bacteria has been previously identified by its inability to ferment trehalose[15]. Similarly, one of

the defining differences among other *Bacteroides* species is *B. fragilis'* inability to ferment

cellobiose[16]. Unlike *E. coli*, all *Bacteroides* species are capable of fermenting maltose[16]. Maltose

fermentation utilizes the RokA kinase which is also used to phosphorylate hexose and amino

sugars[17].

In *B. fragilis*, N-acetyl glucosamine is used more efficiently than glucose and previously

published results suggest similar findings[18]. We also found that neuraminic acid was utilized by

B. fragilis. The kinase responsible for the catabolism of both neuraminic acid and n-acetyl

glucosamine is RokA[17].This kinase along with the hexA kinase are the only glucose kinases in

the cell[17].

3.3 Optical density growth curves for *E. coli* LF82

The growth of *E.coli* in ZMB1 also varied with the different carbohydrate sources. It should

be noted that *E. coli* was able to grow in the modified ZMB1 without an added carbohydrate

source. *E. coli* is typically grown on media containing Bacto[TM] Tryptone which is a supplier of

amino acids for the bacteria[19,20]. *E. coli* has been known to grow on L-alanine as its sole carbon

source[21]. It is probable the bacteria were able to utilize the amino acids included in the modified

ZMB1.

E. coli demonstrated significant growth on all monosaccharides, but only showed limited

growth on some disaccharides. Cellobiose and maltose both contain two glucose monomers but

the bacteria had little to no growth. Other strains of *E. coli* have been able to utilize cellobiose

through mutations likely not found in the LF82 strain used[22]. *E. coli*'s ability to utilize maltose

such as in the K12 and O157:H7 strains rose from a mutant of the original strain[23,24]. When two

glucose monomers are connected in an α-glycosidic linkage as in trehalose, there is significant *E.*

coli growth. The K12 and LF82 strains are capable of fermenting trehalose[25]. Additionally, the

growth curves of the 10 mM and 100 mM differ likely due to trehalose being degraded to

glucose by different methods dependent on the osmolarity of the growth conditions[26].

In previously published studies, *E. coli* is able to utilize N-acetyl-D-glucosamine more

efficiently than glucosamine[27]. Glucosamine reports a higher growth rate in this study likely due

to a mutation that limits *E. coli* growth on N-acetyl-D-glucosamine[28]. This mutation is likely on

the glucosamine-6-phospahte deaminase (NagB) which when affected allosterically causes

differences in growth rates[27]. *E. coli* is also capable of utilizing neuraminic acid well due to the

presence of an extracellular mutarotase (NanM) in *E. coli* allowing neuraminic acid to be utilized

as a carbohydrate source[29].

3.4 Analysis of model parameters

The r and d model parameters show growth or death contribution, respectively, to the growth

curve by an undetermined mechanism (Fig. 4 and 5). In this model, the r parameter is defined the

same way as the logistic equation. The d parameter describes the speed of decline of the carrying

capacity of the population over time. When $d = 0$, the model becomes the logistic model.

The growth was analyzed using the product of the r and Y_{max} parameter (Fig. 4). For *E. coli*,

growth is led by the 10 mM concentration for most sugars except for fucose and cellobiose

which are led by the 100 mM concentration. The growth of *B. fragilis* on these sugars appears to

be dependent on the sugar and not the concentration. Fucose, lactose, and neuraminic acid appear

to be concentration dependent as they are the only ones with large differences in their growth

values between concentrations. The death rate was only looked at for *B. fragilis* in which most of

the sugars produce a death effect. There is a concentration dependence for the d parameters for

maltose and lactulose, which is not seen when analyzing their growth.

A similar experiment conducted by Desai *et al.* using *E. coli* showed no growth for N-acetyl

neuraminic acid or cellobiose while our experiments found a statistically significant growth rate

for these two sugars[30]. Similar trends as our findings are found for cellobiose, galactose, glucose,

and N-acetyl glucosamine. We found growth on neuraminic acid while Desai *et al.* did not. The

difference is likely because Desai *et al.* using a different strain of *E. coli* (HS). Other differences

such as Desai *et al.* reporting higher growth levels for glucosamine and fucose than fructose are

likely due to differences in carbohydrate usage among different strains.

The interactive web application Dashing Growth Curves was utilized for a preliminary

comparison of the growth parameters of the Modified Gompertz and Modified Logistic models

to our model parameters[31]. The product of r or maximum growth rate and Y_{max} was calculated for

the three models. The results are not comparable due to the high standard deviation of the

parameters obtained from using the web application (Table S4). The high standard deviations are

likely due to the association of blanks to samples not being automated. Future work will include

a more comprehensive comparison of the models.

4. Conclusions

This paper provides a model that can be used to predict a variety of different trends found in

bacterial growth. The parameters are used to indicate a contribution to growth whether positive

or negative. This model improves upon previous models in two key ways: i) it does not require

knowledge of the metabolic interactions for the trends showing several growth periods, and ii) a

single model allows characterization of trends in growth that are sigmoidal, non-sigmoidal,

death, or a combination. A commonly used model like the Gompertz equation is said to be

applicable for both death and growth but has a severe limitation of $N(t=0)$ not equaling $N(0)$

which may disturb curve fitting. While logistic and log-linear equations can only be applied to

growth or death, respectively.

Three limitations of this model include: i) it lacks lag phase as a parameter, ii) the parameters

do not indicate what is causing change in growth, and iii) r value is calculated for entire period.

For example, this model is unable to determine the cause for the large death values seen in *B.*

fragilis grown using maltose as the carbon source. The *B. fragilis* strain used in these

experiments has prophage integrated in the genome. Experiments were conducted to induce the

phage to replicate the death seen (Fig. S3). No difference was seen between the control and

induced phage studies implying phage were not the source of the sudden drop in turbidity.

Investigations on chemical changes that may be causing the death are part of future studies.

Calculating the r value for the entire growth period creates discrepancies when analyzing data

that may have the same instantaneous growth rate, but reach Y_{max} more slowly or rapidly thus

reporting a different r value. Further work includes adapting the model for several growth rates

to deconvolute data that exhibits more than one growth rate before reaching Y_{max}.

We anticipate that the use of models like the one developed here can provide quantification

to commonly observed trends in growths. This quantification can then guide design of growth

environments for microbes, for example, in bioreactors or in microbiome engineering.

5. Data Availability

267   The raw data files for this work will be available on Mendeley Data under the title

268   "Modelling complex growth profiles of *Bacteroides fragilis* and *Escherichia coli* on various

269   carbohydrates in an anaerobic environment."

6. Acknowledgements

272   We are grateful to Purdue University's Data Science Consulting Services to help with the

273   data analysis and development of the model. We are also thankful to Ann Christine Caitlin

274   and Steven Clark for helping us set up the analysis on DEEDS. The work was funded

275   through Purdue's internal funds.

7. References

277
1. Pérez-Rodríguez, F. & Valero, A. Predictive Microbiology in Foods. in *Predictive Microbiology in*
Foods (eds. Perez-Rodriguez, F. & Valero, A.) 1–10 (Springer, 2013). doi:10.1007/978-1-4614-5520-
2_1.
2. Xiong, R., Xie, G., Edmondson, A. E. & Sheard, M. A. A mathematical model for bacterial
inactivation. *International Journal of Food Microbiology* **46**, 45–55 (1999).
3. Navarro Llorens, J. M., Tormo, A. & Martínez-García, E. Stationary phase in gram-negative bacteria.
FEMS Microbiology Reviews **34**, 476–495 (2010).
4. Kompala, D. S., Ramkrishna, D., Jansen, N. B. & Tsao, G. T. Investigation of bacterial growth on
mixed substrates: Experimental evaluation of cybernetic models. *Biotechnology and Bioengineering*
28, 1044–1055 (1986).
5. Narang, A. & Pilyugin, S. S. Bacterial gene regulation in diauxic and non-diauxic growth. *Journal of*
Theoretical Biology **244**, 326–348 (2007).
6. Zwietering, M. H., Jongenburger, I., Rombouts, F. M. & Riet, K. van 't. Modeling of the Bacterial
Growth Curve. *Appl. Environ. Microbiol.* **56**, 1875–1881 (1990).
7. Pruitt, K. M. & Kamau, D. N. Mathematical models of bacterial growth, inhibition and death under
combined stress conditions. *Journal of Industrial Microbiology* **12**, 221–231 (1993).
8. Linton, R. H., Carter, W. H., Pierson, M. D. & Hackney, C. R. Use of a Modified Gompertz Equation
to Model Nonlinear Survival Curves for Listeria monocytogenes Scott A. *J Food Prot* **58**, 946–954
(1995).
9. Ball, C. O. & Olson, F. C. W. Sterilization in food technology. Theory, practice, and calculations.
Sterilization in food technology. Theory, practice, and calculations. (1957).
10. Geeraerd, A. H., Herremans, C. H. & Van Impe, J. F. Structural model requirements to describe
microbial inactivation during a mild heat treatment. *Int J Food Microbiol* **59**, 185–209 (2000).
11. Albert, I. & Mafart, P. A modified Weibull model for bacterial inactivation. *International Journal of*
Food Microbiology **100**, 197–211 (2005).
12. Zhang, G., Mills, D. A. & Block, D. E. Development of chemically defined media supporting high-
cell-density growth of lactococci, enterococci, and streptococci. *Appl. Environ. Microbiol.* **75**, 1080–
1087 (2009).
13. Rios-Covian, D. *et al.* Bacteroides fragilis metabolises exopolysaccharides produced by
bifidobacteria. *BMC Microbiol* **16**, 150 (2016).
14. Caspari, D. & Macy, J. M. The role of carbon dioxide in glucose metabolism of Bacteroides fragilis.
Arch. Microbiol. **135**, 16–24 (1983).

15. San Joaquin, V. H., Griffis, J. C., Lee, C. & Sears, C. L. Association of Bacteroides fragilis with
Childhood Diarrhea. *Scandinavian Journal of Infectious Diseases* **27**, 211–215 (1995).

16. Tannock, G. W. Characteristics of Bacteroides isolates from the cecum of conventional mice. *Appl*
Environ Microbiol **33**, 745–750 (1977).

17. Brigham, C. J. & Malamy, M. H. Characterization of the RokA and HexA Broad-Substrate-
Specificity Hexokinases from Bacteroides fragilis and Their Role in Hexose and N-
Acetylglucosamine Utilization. *J Bacteriol* **187**, 890–901 (2005).

18. Chen, H.-C., Chang, C.-C., Mau, W.-J. & Yen, L.-S. Evaluation of N-acetylchitooligosaccharides as
the main carbon sources for the growth of intestinal bacteria. *FEMS Microbiology Letters* **209**, 53–56
(2002).

19. Lessard, J. C. Chapter Eleven - Growth Media for E. coli. in *Methods in Enzymology* (ed. Lorsch, J.)
vol. 533 181–189 (Academic Press, 2013).

20. Davami, F., Eghbalpour, F., Nematollahi, L., Barkhordari, F. & Mahboudi, F. Effects of Peptone
Supplementation in Different Culture Media on Growth, Metabolic Pathway and Productivity of
CHO DG44 Cells; a New Insight into Amino Acid Profiles. *Iran Biomed J* **19**, 194–205 (2015).

21. Tong, M. *et al.* Gene Dispensability in Escherichia coli Grown in Thirty Different Carbon
Environments. *mBio* **11**, e02259-20 (2020).

22. Kachroo, A. H., Kancherla, A. K., Singh, N. S., Varshney, U. & Mahadevan, S. Mutations that alter
the regulation of the chb operon of Escherichia coli allow utilization of cellobiose. *Molecular*
Microbiology **66**, 1382–1395 (2007).

23. Jones, S. A. *et al.* Glycogen and Maltose Utilization by Escherichia coli O157:H7 in the Mouse
Intestine. *Infect Immun* **76**, 2531–2540 (2008).

24. Siniagina, M. N. *et al.* Diversity and Adaptations of Escherichia coli Strains: Exploring the Intestinal
Community in Crohn's Disease Patients and Healthy Individuals. *Microorganisms* **9**, 1299 (2021).

25. Doudoroff, M., Hassid, W. Z., Putman, E. W., Potter, A. L. & Lederberg, J. DIRECT UTILIZATION
OF MALTOSE BY ESCHERICHIA COLI. *Journal of Biological Chemistry* **179**, 921–934 (1949).

26. Elbein, A. D. New insights on trehalose: a multifunctional molecule. *Glycobiology* **13**, 17R – 27
(2003).

27. Álvarez-Añorve, L. I., Calcagno, M. L. & Plumbridge, J. Why Does Escherichia coli Grow More
Slowly on Glucosamine than on N-Acetylglucosamine? Effects of Enzyme Levels and Allosteric
Activation of GlcN6P Deaminase (NagB) on Growth Rates. *J Bacteriol* **187**, 2974–2982 (2005).

28. White, R. J. Control of amino sugar metabolism in Escherichia coli and isolation of mutants unable to
degrade amino sugars. *Biochem J* **106**, 847–858 (1968).

29. Almagro-Moreno, S. & Boyd, E. F. Bacterial catabolism of nonulosonic (sialic) acid and fitness in the

gut. *Gut Microbes* **1**, 45–50 (2010).

30. Desai, M. S. *et al.* A Dietary Fiber-Deprived Gut Microbiota Degrades the Colonic Mucus Barrier

and Enhances Pathogen Susceptibility. *Cell* **167**, 1339-1353.e21 (2016).

31. Reiter, M. A. & Vorholt, J. A. Dashing Growth Curves – a web application for rapid and interactive

analysis of microbial growth curves. 2022.12.16.520708 Preprint at

https://doi.org/10.1101/2022.12.16.520708 (2022).

Figure 1. Schematic of experimental setup including the logic for including values in death

parameter analysis. r is the instantaneous growth rate. d is the linear rate of decline of carrying

capacity.

Figure 2. Chemical structures of the monosaccharides and disaccharides used in this work.

Figure 3. Overall growth curves of *Bacteroides fragilis* (A) and *Escherichia* coli (B) for 13 sugars at three concentrations. Each dark

line represents the average OD_{600} of the replicates for an experimental run. The gray shading represents the standard deviation of the

OD_{600} of the replicates. The media control was included to show the growth in the absence of a sugar.

Figure 4. (A) Overall growth rate data of *Bacteroides fragilis*. Each dot represents the average value of a different experimental condition or the negative (media) control. The lines represent the 95% confidence interval of each condition. (B) Overall growth rate data of *Escherichia coli*.

369   The gray shaded area represents the data that was not significantly greater than the media

370   control. The numbers on the right are the number of replicates.

372 .

373
Figure 5. Death rate data of *Bacteroides fragilis*. Each dot represents the average value of a

different experimental condition or the negative (media) control. The lines represent the 95%

confidence interval of each condition. The numbers on the right are the number of replicates.

Table 1. Comparison of growth and death models with the model of this study. [a] $N =$

instantaneous population size, r = instantaneous growth rate, and K = carrying capacity. [b] λ = lag

time, μ_m = maximum specific growth rate, A = carrying capacity. [c] k = initial cell count, Δ =

decrease of cell count over time, d_{max} = maximum death rate, α = shoulder. [d] D = decimal

reduction value, N and N_0 = population at time t. [e] K_{max} = maximum death rate, N_{res} = residual

population, C_c = concentration of the critical component. [f] $\gamma(t)$ = bacterial concentration at time

t. p = curve shape (concavity, convexity, and shoulder). t = time, δ = first decimal reduction

concentration for the population not belonging to N_{res}, N(0) = initial cell concentration, N_{res} =

residual bacterial concentration. [g] r = instantaneous growth rate, Y = instantaneous population

size, Y_{max} = max population size, d = linear rate of decline of carrying capacity, t_{max} = time at

which Y_{max} is reached. In all cases, t = time.

Name	Model	Model type	Trends fit	Limitation(s)	Reference
Logistic	$\frac{dN}{dt} = rN(1 - \frac{N}{K})$	Growth	fits sigmoidal with no lag	No lag parameter, fits long lag phases poorly	
Modified Gompertz[b]	$y = A * e^{-e^{\frac{\mu_m e}{A}(\lambda - t) + 1}}$	Growth	Fits lag or no lag	N(t=0)≠N(0)	[6]Zwietering et al.
Negative Gompertz[c]	$y = k - \Delta e^{-e^{\frac{d_{max} e}{\Delta}(\alpha - t) + 1}}$	Death	Fits shoulder and tail	N(t=0)≠N(0)	[8]Linton et al.
Log-linear[d]	$log\,N = log\,N_0 - \frac{t}{D}$	Death	Fits linear	Cannot fit data with shoulder or tail	[9]Ball and Olson
Shoulder-tail[e]	$N(t) = (N(0) - N_{res}) * e^{-k_{max}*t} \frac{1 + C_c(0)}{1 + C_c(0)e^{-k_{max}*t}} + N_{res}$	Death	Fits linear, shoulder, and tail		[10]Geeraerd et al.
Modified Weibull[f]	$\gamma\left(\frac{t}{\delta}\right) = (N_0 - N_{res})10^{(-(\frac{t}{\delta})^p)} + N_{res}$	Death	Fits linear, shoulder, and tail	N_{res} parameter loses accuracy with extended experiment duration.	[11]Albert and Mafart
This study[g]	$\frac{dY}{dt} = r * Y \frac{1 - Y}{(Y_{max} - d) * (t - t_{max})}$	Growth & Death	Fits growth with or without lag, linear death	Fitting complex trends with several growth or death rates	

392