RESEARCH ARTICLE

Title: Sulfur oxidation and reduction are coupled to nitrogen fixation in the roots of a salt marsh

foundation plant species

Authors: Rolando, J.L.[a]; Kolton, M.[a,b]; Song, T.[a], Liu, Y.[a,c]; Pinamang, P.[a], Conrad, R.[a]; Morris,

J.T.[d], Konstantinidis, K.T.[a,e], Kostka, J.E. [a,f,g]

Affiliations:

[a]Georgia Institute of Technology, School of Biological Sciences, Atlanta, GA 30332, USA

[b]French Associates Institute for Agriculture and Biotechnology of Drylands, Ben-Gurion

University of the Negev, Beer Sheva, Israel

[c]The Pennsylvania State University, Department of Civil & Environmental Engineering,

University Park, PA, 16802

[d] Belle Baruch Institute for Marine & Coastal Sciences, University of South Carolina, Columbia,

SC, 29201

[e]Georgia Institute of Technology, School of Civil and Environmental Engineering, Atlanta, GA

30332, USA

[f]Georgia Institute of Technology, School of Earth and Atmospheric Sciences, Atlanta, GA 30332,

USA

[g]Center for Microbial Dynamics and Infection, Georgia Institute of Technology, Atlanta, GA

30332, USA

Corresponding author:

Kostka, JE. e-mail: joel.kostka@biology.gatech.edu

Abstract

Symbiotic root microbiota are crucial for plant growth as they assist their hosts in nutrient acquisition. In the roots of coastal marine plants, heterotrophic activity in the rhizosphere by sulfate-reducing microorganisms has been linked to nitrogen fixation. In this study, we recovered 239 high-quality metagenome-assembled genomes (MAGs) from a salt marsh dominated by the foundation plant *Spartina alterniflora*, including diazotrophic sulfate-reducing and sulfur-oxidizing bacteria thriving in the root compartment. Here we show for the first time that highly-abundant sulfur-oxidizing bacteria in the roots of a coastal macrophyte encode and highly express genes for nitrogen fixation (nifHDK). Further, we leveraged a *S. alterniflora* biomass gradient to gain a mechanistic understanding on how root-microbe interactions respond to abiotic stress from anoxia and elevated sulfide concentration. We observed that the roots of the stressed *S. alterniflora* phenotype exhibited the highest rates of nitrogen fixation and expression levels of both the oxidative and reductive forms of the dissimilatory sulfite reductase gene (dsrAB). Approximately 25% and 15% of all sulfur-oxidizing dsrA and nitrogen-fixing nifK transcripts, respectively, were associated with novel MAGs of the *Candidatus* Thiodiazotropha genus in the roots of the stressed *S. alterniflora* phenotype. We conclude that the rapid cycling of sulfur in the dynamic *S. alterniflora* root zone is coupled to nitrogen fixation during both reductive and oxidative sulfur reactions, and that the *S. alterniflora – Ca.* Thiodiazotropha symbiosis is an adaptive response to anoxic and sulfidic sediment conditions, whereby the plants benefit from reduced sulfide toxicity and potential nitrogen acquisition.

Introduction

The microbial communities closely associated with plant hosts (i.e., plant microbiota) have received substantial attention due to their potential role in plant nutrient acquisition, phytohormone synthesis, prevention of soil-borne disease, and detoxification of the rhizosphere and root environment (Bulgarelli et al., 2013; Trivedi et al., 2020). Most plant microbiome studies have been performed in terrestrial ecosystems with an emphasis on agricultural plants (Trivedi et al., 2020). Plant microbiota from vegetated coastal ecosystems (i.e., seagrass meadows, mangroves, and salt marshes) remain mostly understudied, even though they play a key role in global climate regulation and the cycling of major nutrients (Barbier et al., 2011; Duarte et al., 2013). Covering less than 3% of the global surface area occupied by forests, coastal vegetated ecosystems exhibit an equivalent rate of carbon sequestration at the planetary scale due to their excess in primary production over organic matter decomposition (Duarte et al., 2013). Further, vegetated estuarine ecosystems play a key role in water purification by reducing coastal nutrient pollution (Barbier et al., 2011). Despite their high ecological value, little is known about how plant-microbe interactions contribute to the functioning of coastal marine ecosystems, their resilience to climate change, and provisioning of ecosystem services.

Previous studies have shown that the root zone of coastal marine plants is a hotspot for the cycling of carbon, nitrogen, and sulfur (Gandy and Yoch, 1988; Welsh et al., 1996; Spivak and Reeve, 2015; Thomas et al., 2014; Crump et al., 2018; Kolton et al., 2020). In coastal Georgia, US, sulfate-reducing and sulfur-oxidizing bacteria were recently shown to be overrepresented in the core root microbiome of the salt marsh foundation plant *Spartina alterniflora*, comprised of persistent taxa across multiple plant individuals (Rolando et al., 2022). Sulfate reduction, an anaerobic respiration pathway, represents a dominant terminal electron-accepting process coupled to the breakdown of

organic matter in marine ecosystems (Kostka et al., 2002; Jørgensen et al., 2019). Evidence from experimental and field research indicates that rhizospheric sulfate-reducing bacteria assimilate photosynthates from *S. alterniflora*, while fueling nitrogen fixation (Gandy and Yoch, 1988; Spivak and Reeve, 2015). Similarly, rates of nitrogen fixation have been closely associated with organic matter degradation by sulfate reduction in seagrass meadows (McGlathery et al., 1998; Herbert, 1999; Nielsen et al., 2001; Welsh et al., 1996). Conversely, the oxidation of reduced sulfur compounds is a chemolithotrophic process requiring potent terminal electron acceptors such as oxygen, nitrate, or oxidized metals. In particular, sulfide is a known phytotoxin, and its belowground oxidation is considered a detoxifying reaction for plants inhabiting coastal marine ecosystems (Lamers et al., 2013). Based on amplicon sequencing of the 16S rRNA gene, sulfur oxidation coupled with nitrogen fixation was recently hypothesized as an important process for plant growth under sulfidic conditions (Rolando et al., 2022). Nitrogen is often the limiting nutrient for salt marsh plants, and stress from sulfide toxicity and anoxia further impairs root active acquisition of nitrogen (Mendelssohn and Morris, 2000). Close relatives of diazotrophic sulfur-oxidizing symbionts from the *Sedimenticolaceae* family (genus: *Candidatus* Thiodiazotropha) have been shown to inhabit the roots of seagrasses and marsh plants in coastal marine ecosystems (Martin et al., 2020a, Rolando et al., 2022). The *Ca.* Thiodiazotropha genus was first discovered as bacterial symbionts of lucinid clams, where they provide both fixed carbon and nitrogen to their animal host by sulfur-mediated chemolithoautotrophy (Petersen et al., 2017). Most studies assessing the ecology and function of sulfur chemosymbiosis have been performed within marine invertebrates hosts (Petersen et al., 2017; Osvatic et al., 2021, 2023; Lim et al., 2019a, 2019b). In contrast, there is still no evidence for the coupling of sulfur oxidation with nitrogen and/or carbon fixation in the roots of coastal marine plants because the genomes, metabolic potential, and activity

of root-associated sulfur-oxidizing bacteria are still lacking. Limited studies have looked into the composition and abundance of sulfur-oxidizing bacteria present in the roots of coastal marine plants. These studies mainly relied on microscopy- and DNA/RNA amplicon-based community analyses (Thomas et al., 2014; Martin et al., 2020a; Kolton et al., 2020; Rolando et al., 2022). Only one study utilized a metatranscriptomics approach to investigate the gene expression of seagrass root-associated microbiomes (Crump et al., 2018). However, genome-wide gene expression profiles of sulfur-oxidizing bacteria and the coupling of sulfur oxidation with other biogeochemical processes are still to be deciphered in the roots of coastal marine plants.

Ecosystem models and empirical evidence indicate that climate change is altering the hydrology, biogeochemistry, and plant community composition of coastal wetlands (Guimond et al., et al., 2020; Donnelly and Bertness, 2001). Thus, a mechanistic understanding of how environmental perturbations impact plant-microbe interactions will be critical to forecasting the resilience of salt marsh ecosystems to climate change. According to the stress-gradient hypothesis (SGH), the frequencies of facilitative and competitive interactions are inversely related along abiotic stress gradients, with facilitative interactions occurring more frequently under high abiotic stress than under benign conditions (Bertness and Callaway, 1994). This concept has mostly been explored in plant-plant interactions; whereas, limited studies assessing plant-microbe interactions have found that the SGH is consistent in certain plant phenological stages (David et al., 2020). Following the SGH, we hypothesize that beneficial plant-microbe interactions will be more common in stressed coastal marine ecosystems, where plants experience greater salinity, redox, and sulfide toxicity stress. *S. alterniflora*-dominated salt marsh ecosystems represent an ideal natural laboratory in which to study the effects of stress on plant-microbe interactions because steep gradients in plant productivity are formed within short distances (Mendelssohn and Morris, 2000). The productivity

gradient is the result of decreased soil redox potential, along with increased salinity and anoxia, extending from vegetated tidal creek banks towards the interior of the marsh (Figure 1). Higher primary productivity is reflected in tall *S. alterniflora* plants (> 80 cm) growing adjacent to tidal creek banks, while smaller plants (< 50 cm) inhabit the interior of the marsh. We hypothesize that the strength of plant-sulfur chemosymbiosis increases in stressed marshes where elevated sulfide oxidation rates are coupled to nitrogen fixation.

Thus, the objectives of the present study were: (i.), to demonstrate the coupling of sulfur oxidation with nitrogen fixation in the root environment of *S. alterniflora* and (ii.) to evaluate the effect of environmental stress on the assembly, activity, and ecological interactions of the *S. alterniflora* root microbiome.

2. Results

2.1 Genome-centric multi-omics approach reveals prokaryotic sulfur metabolism is coupled to nitrogen fixation in the roots of a foundation salt marsh plant

We studied the root microbiome of the salt marsh foundation species of the US Atlantic and Gulf of Mexico coastlines, *Spartina alterniflora*, at Sapelo Island, GA during the summers of 2018, 2019 and 2020. A combination of prokaryotic DNA and RNA quantification, shotgun metagenomics, metatranscriptomics, and rate measurements of two key pathways of the nitrogen cycle were performed along a stress gradient within three compartments: sediment, rhizosphere and root (Figure 1). The short and tall *S. alterniflora* phenotypes represent the end extremes of the plant biomass gradient formed by stress from salinity, sulfide toxicity and anoxia (Figure 1). Four biological replicates per compartment and *S. alterniflora* phenotype were sequenced for both

metagenomic and transcriptomic analysis (Further details about sequencing in Materials and

Methods and Supplementary Table S1).

Using a custom-designed bioinformatics pipeline, we binned 239 high-quality metagenome

assembled genomes (MAGs; > 50 Quality score). MAGs were dereplicated into 160

genomospecies (gsp, plural: gspp) by grouping them according to 95% average nucleotide identity

(ANI) (Figure 2). MAGs were assigned to 19 phyla. Almost half of the recovered genomes were

taxonomically affiliated with the *Proteobacteria* or *Desulfobacterota* phyla (Figure 2, further

MAGs taxonomic and statistical information in Supplementary Table S2). Taxonomic novelty was

assessed using GTDB-Tk v2.1.0 with the reference database R07-RS207 (Chaumeil et al., 2022).

Recovered MAGs included 1 gspp from a previously undescribed order, 10 gspp from undescribed

families, 52 gspp from undescribed genera, 94 gspp from undescribed species, and only 3 gspp

from previously described species. In order to assess the MAGs' genetic potential to perform

important biogeochemical functions in the salt marsh environment, we annotated their open

reading frames (ORFs) using the eggNOG database, and focused on predicted genes involved in

the biogeochemical cycling of carbon, nitrogen, and sulfur (Supplementary Table S3).

A large proportion of binned *Proteobacteria*, including members of the *Ca.* Thiodiazotropha

genus, contained genes for nitrogen fixation, carbon fixation through RuBisCO, as well as for

dissimilatory sulfur metabolism, and thiosulfate oxidation using the soxABXYZ complex (Figure

2). Most MAGs from the *Desulfobacterota* phylum presented genes for dissimilatory metabolism

of sulfur, including members of the *Desulfosarcinaceae* family (Figure 2). Because the

dissimilatory sulfite reductase enzyme (dsrAB) can function in either sulfur reduction or oxidation,

we performed a phylogenetic analysis to identify the dsrAB type from our MAGs. We aligned

recovered dsrAB genes to a reference alignment (Müller et al., 2015) and annotated the dsrAB

type based on placement in an approximately-maximum-likelihood phylogenetic tree

(Supplementary Figure S1). All dsrAB genes retrieved from *Proteobacteria* gspp (*Alpha-* and

Gammaproteobacteria gspp) were placed within the oxidative type; while dsrAB genes recovered

from *Acidobacteriota*, *Bacteroidota*, *Chloroflexota*, *Desulfobacterota*, *Gemmatimonadota*, and

Nitrospirota gspp clustered within the reductive bacterial dsrAB type (Supplementary Figures S1,

S2).

We measure rates of $^{15}N_2$ fixation under oxic and anoxic conditions by sediment, a rhizosphere-

root mix, and root samples. Rates were measured under oxic and anoxic conditions to capture the

oxygen fluctuation experienced by microorganisms in the root-zone of *S. alterniflora*. The two *S.*

alterniflora phenotypes exhibited significantly higher N fixation rates in their root tissue compared

to the sediment or rhizosphere-root mix under both aerobic and anaerobic conditions (Figure 3).

Root tissue from the stressed short *S. alterniflora* phenotype showed 1.9- and 5.1-times greater N

fixation rates than roots of the tall phenotype under aerobic and anaerobic conditions, respectively

(Figure 3). Conversely, sediment from the tall *S. alterniflora* phenotype presented greater potential

nitrification rates between one and two orders of magnitude than all other studied compartments

(Figure 3). In the short *S. alterniflora*, potential nitrification rates were an order of magnitude

greater in root tissue compared to bulk and rhizospheric sediment (Figure 3).

We used metagenomic and metatranscriptomic short reads to relate nitrogen fixation activity with

multi-omics functional information. Metagenomic and metatranscriptomic reads were functionally

annotated using the eggNOG database. We normalized the functional profile of each library to

account for differences in sequencing effort and genome size by dividing the count matrix against

the median abundance of 10 universal single-copy phylogenetic marker genes as in Salazar et al.

(2019). Statistical difference between the normalized gene expression of the nitrogenase gene nifK

and the oxidative and reductive dsrA types was calculated between *S. alterniflora* phenotypes

across all compartments. Since gene expression within metabolic pathways was highly correlated,

we used nifK and dsrA as marker genes (Supplementary Figure S3). Even though nitrogen fixation

rates were the greatest in the root compartment of the short *S. alterniflora* phenotype, no statistical

significance was found in the nitrogenase gene expression between the two *S. alterniflora*

phenotypes (Figure 3). Conversely, the root compartment of the short *S. alterniflora* phenotype

had the highest normalized transcript abundance of both reductive and oxidative dsrA gene types

(Figure 3). To infer which members of the *S. alterniflora* root microbiome contributed the most to

nitrogen fixation, as well as to test if nitrogen fixation was coupled to sulfur metabolism, we

mapped all short read transcripts annotated as nifK and dsrA back to our binned gspp. Sulfur-

oxidizing *Ca.* Thiodiazotropha gspp in the short phenotype of *S. alterniflora* contributed

approximately 25% and 15% of all oxidative dsrA and nifK transcripts, respectively (Figure 3). In

addition, the most active sulfur-oxidizing and sulfate-reducing gspp in roots of the short phenotype

(gsp 31 and gsp 68) had a positive correlation between nitrogen fixation and sulfur

oxidation/reduction gene expression, respectively (Supplementary Figure S4). In the tall

phenotype, *Desulfosarcinaceae* gsp 134 contributed the most transcripts of the reductive dsrA

gene. However, in general, we were not able to recover most of the nifK genes as part of the

MAGs; and thus, transcripts, presumably due to issues with contig binning. Nevertheless, using

short read analysis, we found that the majority of the nifK transcripts in the roots of the tall

phenotype were assigned to bacteria from the *Desulfobacterota* phylum, while in the short

phenotype most nifK transcripts were affiliated with bacteria from the *Gammaproteobacteria* class

or the *Desulfobacterota* phylum (Supplementary Figure S5). Similarly, abundant transcripts of

oxidative dsrA in the roots of the short phenotype were mostly affiliated with the

Gammaproteobacteria class, while the reductive dsrA affiliated mainly with the *Desulfobacterales*
and *Desulfovibrionales* orders in both phenotypes (Supplementary Figures S6 and S7).

2.2 Phylogeny, genetic potential and gene expression of microorganisms enriched in the *S. alterniflora* root microbiome

To assess the phylogenetic novelty and relation of *Sedimenticolaceae* sulfur-oxidizing MAGs from the present study in comparison to that of marine invertebrate chemosymbionts, we retrieved all publicly available *Sedimenticolaceae* genomes as reported by GTDB release R07-RS207, as well as all *Ca.* Thiodiazotropha spp. analyzed by Osvatic et al. (2023). A maximum-likelihood phylogenetic tree using a 400 universal marker genes database was performed in PhyloPhlAn. We found that *S. alterniflora* root symbionts assigned to the *Ca.* Thiodiazotropha genus and an unknown genus from the *Sedimenticolaceae* family formed a monophyletic clade with lucinid clam chemosymbionts (Figure 4). All recovered MAGs that were phylogenetically related to marine invertebrate chemosymbionts were highly abundant in the root compartment of the *S. alterniflora* short phenotype, where sulfidic conditions are found in the marsh environment (Figure 1). To conserve phylogenetic coherence, we propose that *S. alterniflora* root symbionts assigned to the *Sedimenticolaceae* family and that formed a monophyletic group with *Ca.* Thiodiazotropha spp. are also members of such genus (Figure 4). Out of the 7 recovered *Ca.* Thiodiazotropha gspp, 6 had genes for carbon fixation (RuBisCO), 5 had the complete or partial nifDKH nitrogenase genes for nitrogen fixation, and all of them harbor the complete or at least partial genes for dissimilatory sulfite reductase (oxidative dsrAB type), and sulfur oxidation by the SOX complex (soxABYXYZ). Furthermore, we calculated the gene expression profile of the most active *Ca.* Thiodiazotropha gspp (gsp 31 and gsp 33), finding that genes for sulfur oxidation through the oxidative dsrAB, soxABYXYZ complex, and cytochrome c oxidase cbb3-type genes, as well as

genes for carbon and nitrogen fixation (RuBisCO and nifKDH) were among the most highly

transcribed (Supplementary Tables S4 and S5). We also found that sulfate-reducing MAGs from

the *Desulfosarcinaceae* family were enriched in the root compartment of *S. alterniflora*

(Supplementary Figure S8). *Desulfosarcinaceae* gspp had contrasting preferences for root

colonization, with gsp 68 mostly enriched in the roots of the short phenotype, gsp 134

preferentially enriched in the root of the tall phenotype, and gsp 80 equally abundant in both

phenotypes (Supplementary Figure S8). Of the three *Desulfosarcinaceae* gspp, all contained the

complete or partial genes for nitrogen fixation, and only gsp 80 did not have the reductive dsrAB

gene. The most transcribed genes from both gsp 68 and gsp 134 were related to dissimilatory

sulfate reduction (dsrAB and aprAB, Supplementary Tables S6 and S7).

<u>2.3 Sediment and root microbiomes have contrasting functional profiles across a salt marsh</u>
<u>primary productivity gradient</u>

A decreasing trend in prokaryotic alpha divestity was observed from the sediment compartment to

the root tissue in both *S. alterniflora* phenotypes (Figure 5). Similarly, quantification of the 16S

rRNA gene by qPCR showed statistically lower prokaryotic abundance in the root tissue compared

to the sediment and rhizosphere compartments (Figure 5). A stronger barrier for prokaryotic root

colonization, evidenced by a steeper decrease in prokaryotic abundance in the rhizosphere-root

interface, was observed in the less stressed tall *S. alterniflora* phenotype (Figure 5). Conversely,

the root compartment of both *S. alterniflora* phenotypes presented a higher transcript copy number

of the prokaryotic 16S rRNA than their sediment counterparts, indicating greater prokaryotic

activity in the root zone (Figure 5).

PERMANOVA and principal coordinate analysis from metagenomic and metatranscriptomic

functional profiles indicated that the environmental stress gradient significantly affects the

microbiome's potential and expressed functional repertoire (Figure 5, Supplementary Table S8).

The community-level abundance and expression of terminal oxidase genes for aerobic respiration

were specific to compartment and *S. alterniflora* phenotype. The relative abundance of genes for

the canonical cytochrome c oxidase system (coxA) was greater in the tall *S. alterniflora* phenotype

in both metagenomic and metatranscriptomic samples, while cbb3-type cytochrome c genes

(ccoN) were mostly present and transcribed in the root tissue of the short *S. alterniflora* phenotype

(Figure 5). Genes related to the redox cycling and turnover of nitrogen were more abundant and

transcribed in the sediment of the tall *S. alterniflora* phenotype than all other assessed marsh

compartments [i.e., nitrification, denitrification, and dissimilatory nitrate reduction to ammonium

(DNRA) genes] (Figure 5). Conversely, genes for prokaryotic carbon fixation by RuBisCO and

nitrogen fixation had the greatest relative abundance in the root compartment, especially in those

from the short *S. alterniflora* phenotype (Figure 5). Nevertheless, relative gene expression for

nitrogen fixation was greater in the sediment from the short phenotype and contributed mostly by

cyanobacteria species (Supplementary Figure S5). Finally, greater redox cycling of sulfur in the

root compartment, especially in the short *S. alterniflora,* was evidenced by the abundance and

expression of sulfur dissimilatory metabolism genes (Figure 5).

2.4 Plant species from coastal marine ecosystems harbor unique root microbiomes

In order to understand how root microbiomes are assembled in coastal marine ecosystems

compared to better-studied terrestrial plants, we used publicly available sequences to generate and

curate a 16S rRNA gene amplicon database from root microbiomes comprising 2,911 amplicon

samples from 56 plant species (complete dataset in Supplementary Table S9). We grouped

amplicon samples into 4 broad ecosystem types: seabed, coastal wetlands, freshwater wetlands,

and other terrestrial ecosystems. Species exchange based on the Bray-Curtis dissimilarity matrix

was largely explained by ecosystem type, with coastal wetland and seabed ecosystems clustering

together in an NMDS ordination (Figure 6). PERMANOVA analysis revealed that ecosystem type

alone significantly explained 13.2% of the variation of the species exchange between analyzed

samples (Supplementary Table S10). Furthermore, to explore two functional guilds that may

explain the difference in microbiome assembly between ecosystem types, we assigned putative

sulfate reduction and sulfur oxidation functions to our taxonomy table as performed by Rolando

et al. (2022). We found that the root microbiomes of both seabed and coastal wetland ecosystems

were highly enriched in putative sulfate-reducing and sulfur-oxidizing bacteria (Figure 6).

Furthermore, we discovered that in both seabed and coastal wetland ecosystems, amplicons

showing high sequence identity to known sulfur-oxidizing bacteria with the capability for nitrogen

fixation (*Sedimenticolaceae* family: *Ca.* Thiodiazotropha genus) were highly abundant

(Supplementary Figure S9). Conversely, amplicons showing high sequence identity to sulfate

reducers from the *Desulfosarcinaceae* family, particularly those from the genus *Desulfatitalea*

were highly enriched in the roots of coastal wetland plants (Supplementary Figure S9). The

taxonomic identity of highly abundant ASVs in the root compartment of coastal marine plants

coincide with diazotrophic MAGs retrieved from Sapelo Island, GA, such as sulfur oxidizers of

the *Sedimenticolaceae* family, and sulfate reducers of the *Desulfosarcinaceae* family.

3. Discussion

<u>3.1 Nitrogen fixation is stimulated by rapid sulfur cycling in the roots of *S. alterniflora*</u>

Nitrogen fixation in belowground coastal vegetated ecosystems has been mainly associated with

heterotrophy, and particularly with sulfate reduction (Gandy and Yoch, 1988; McGlathery et al.,

1998; Herbert, 1999; Nielsen et al., 2001; Welsh et al., 1996). The linkage of sulfate reduction to

nitrogen fixation was proposed based on studies quantifying rates of diazotrophy with and without

sulfate reduction inhibition by molybdate. Here, we show for the first time that highly-abundant

sulfur-oxidizing bacteria in the roots of *S. alterniflora* also encode and highly express genes for

nitrogen fixation (Figure 2, Figure 3). Furthermore, the MAG with the highest nitrogenase gene

expression in our study was a novel sulfur-oxidizing bacterium from the *Ca.* Thiodiazotropha

genus that also showed high expression of sulfur oxidation genes (Figure 3, Supplementary Figure

S4). Our results suggest that when sulfate reduction was inhibited in previous studies, it disrupted

the redox cycling of sulfur by impeding the flow of reduced sulfur compounds that serve as an

energy source for sulfur-oxidizing bacteria. Thus, most likely previous inhibition studies not only

blocked nitrogen fixation from sulfate-reducing microorganism, but also significantly reduced

nitrogen fixation coupled to sulfur oxidation. Unlike sulfate reduction, there is no reliable method

for measuring sulfide oxidation rates. Thus, the abundance and expression of sulfur-oxidation

genes is one of the few ways to directly link this otherwise unmeasurable process with nitrogen

fixation. We propose that both dissimilatory sulfur reduction and oxidation, and more importantly,

the rapid redox cycling of sulfur, stimulate nitrogen fixation in the root environment of *S.*

alterniflora. Since coastal wetland plants and seagrasses assemble similar root microbiomes

(Figure 6), and macrophyte activity boosts both nitrogen fixation and the cycling of sulfur, this

may be a common phenomenon in the root zone of many coastal marine plants (Whiting et al.,

1986; McGlathery et al., 1998; Welsh et al., 1996).

Further evidence supporting the significance of the rapid cycling of sulfur coupled to nitrogen

fixation is that members of the *Gammaproteobacteria* class and *Desulfobacterota* phylum

contributed most of the nitrogenase and oxidative/reductive dsrAB transcripts in the root

compartment of *S. alterniflora*. Similar results have been obtained from DNA and RNA amplicon

studies in seagrass and other coastal wetland plant species, where nitrogen fixation genes and their

expression have been affiliated with microorganism closely related to sulfate-reducing and sulfur-

oxidizing bacteria (Thomas et al., 2014; Crump, et al. 2018; Kolton et al., 2020).

3.2 The root zone of coastal marine ecosystems is a hotspot of sulfur redox cycling

In most coastal marine ecosystems, water saturated sediments are depleted in oxygen within the

first millimeters depth (Jørgensen et al., 2019). Because coastal marine ecosystems are bathed in

seawater containing high sulfate concentrations (28 mM), sediment sulfate-reducing

microorganisms often perform the terminal step of organic matter decomposition (Jørgensen et al.,

2019). Unlike aerobic respiration, energy flow during sulfate reduction is decoupled from the

carbon cycle, with most of the free energy conserved in reduced sulfur compounds (Howarth,

1984). The chemically stored energy is subsequently released by biotic and abiotic oxidation

reactions at marsh oxic-anoxic interfaces coupled to the reduction of potent electron acceptors

(oxygen, nitrate, or oxidized metals) (Howarth, 1984). Surface sediments of vegetated coastal

marine ecosystems experience rapid re-oxidation of most reduced sulfur compounds. This process

is highly concentrated in the microaerophilic root zone, which serves as a hotspot for the reaction

(Holmer et al., 2002; Kristensen and Alongi, 2006; Koop-Jakobsen et al., 2018). Our results

showing that the root compartment of coastal marine ecosystems is highly enriched in

microorganisms with the capability for both sulfur reduction and oxidation provides further

evidence for rapid sulfur cycling in the root zone (Figure 6). Furthermore, to the best of our

knowledge, this is the first study to employ genome reconstruction from metagenomic sequencing

to verify the function of uncultivated sulfate-reducing and sulfur-oxidizing microorganisms living

in the roots of coastal wetland plants. Our findings also reveal that genes that catalyze oxidative

and reductive reactions in the sulfur cycle are highly transcribed in the root environment,

particularly in the stressed *S. alterniflora* phenotype that thrives in a sulfidic environment (up to

1.5 mM sulfide concentration in our studied transects; Rolando et al. 2022). This is consistent with

previous studies of salt marsh and seagrass ecosystems, in which genes of sulfur-oxidation

pathways were shown to be highly expressed in the root compartment of *S. alterniflora* and *Zostera*

seagrass spp., respectively (Thomas et al. 2014, Crump et al., 2018). Thus, we propose that in

contrast to terrestrial ecosystems, coastal marine plants rely on the rapid cycling of sulfur in their

root zone for the breakdown of organic matter and recycling of nutrients through sulfate reduction

along with the re-oxidation of terminal electron acceptors by sulfur oxidation. Furthermore, we

propose that rapid rates of both oxidative and reductive sulfur reactions represent a key mechanism

to sustain high rates of energetically-expensive nitrogen fixation in the root zone of *S. alterniflora*.

3.3 Phylogenetic and ecological relationship between marine invertebrates and *S. alterniflora*

chemosymbionts

Endosymbionts from the *Ca.* Thiodiazotropha genus were initially discovered living in symbiosis

with lucinid clams, where they fix carbon and serve as a source of carbon and nitrogen to the

animal host using energy gained from the oxidation of reduced forms of sulfur (Petersen et al.,

2017). Recent studies using fluorescence in situ hybridization (FISH) microscopy and SSU rRNA

gene metabarcoding have shown that *Ca.* Thiodiazotropha bacteria also inhabit the roots of a

diverse array of seagrass species along with that of the coastal cordgrass *S. alterniflora* (Martin et

al., 2020a; Rolando et al., 2022). Here, we show that close relatives of diazotrophic sulfur

chemosymbionts associated with lucinid clams were highly abundant and active in the roots of *S.*

alterniflora. To the best of our knowledge, this is the first study reporting genomes and gene

expression profiles from *Ca.* Thiodiazotropha retrieved from macrophyte root samples. We show

that our MAGs formed a monophyletic clade with those from previously described marine

invertebrate animals (Figure 4). Similar to what has been reported in their symbiosis with lucinid

clams, the *S. alterniflora* symbionts highly expressed genes for sulfur oxidation, and carbon and

nitrogen fixation (Supplementary Table S4, Supplementary Table S5; Petersen et al., 2017; Lim et

al., 2019a). Studies of lucinid clams have shown that the *Ca.* Thiodiazotropha symbionts are

horizontally transmitted (Petersen and Yuen, 2021). Furthermore, invertebrate colonization by *Ca.*

Thiodiazotropha symbionts is not restricted by either the host or symbiont species (Lim et al.,

2019a). The flexibility of this symbiotic relationship could explain the evolution of *Ca.*

Thiodiazotropha symbionts colonizing macrophyte roots. Our results show that all *Ca.*

Thiodiazotropha symbionts of *S. alterniflora* are distinct species from those previously found in

lucinid clam hosts (ANI < 82% for all pairwise comparisons). However, additional metagenomic

sequencing of both macrophyte plants and lucinid clams is needed to better interrogate if

chemosymbionts populations are restricted by host biology at the kingdom level (i.e., to either

plant or animal hosts). In contrast to what is observed in lucinid clams, where a single bacterial

species dominates the gill microbiome; in *S. alterniflora* and seagrass species, *Ca.* Thiodiazotropha

spp. do not completely outcompete other microbial species in the root compartment (Lim et al.,

2019a, 2019b). However, the symbionts comprise a large proportion of the microbial community,

and are amongst the most active species (Rolando et al., 2022; Supplementary Figure S9).

3.4 Sulfur chemosymbiosis as an adaptation to anoxia and sulfide toxicity in *S. alterniflora* plants

We propose that the symbiosis between *S. alterniflora* and sulfur-oxidizing bacteria represents a

key adaptation supporting the resilience of coastal salt marsh ecosystems to environmental

perturbations. For instance, intertidal wetland ecosystems are vulnerable to climate change because

they are located in a narrow elevation range determined by tidal amplitude (Morris et al., 2021).

The Intergovernmental Panel on Climate Change (IPCC) has projected an increase in sea level

between 0.38 m and 0.77 m by 2100 (IPCC-AR6, Fox-Kemper et al., 2021). Although coastal

388 wetland ecosystems are dynamic and adapt to sea level rise by increasing sediment accretion rates,

389 ecosystem models predict that a large area of present-day marsh will drown because of accelerated

390 sea level rise (Kirwan et al., 2016). Increased hydroperiods will impact the redox balance of

391 vegetated sediments, imposing more severe anoxia and physiological stress from sulfide toxicity

392 to wetland plants. Under this scenario, a symbiosis of coastal vegetated plants with sulfur-

393 oxidizing bacteria could alleviate sulfide stress while at the same time coupling it to carbon and

394 nitrogen fixation for potential plant uptake; however, the mechanism for carbon and/or nitrogen

395 transfer between sulfur-oxidizing symbionts and the host plants still remains elusive and requires

396 further research. Similar to Thomas et al. (2014), we showed the transcription of sulfur oxidation

397 genes was greater in the stressed *S. alterniflora* short phenotype. In seagrass ecosystems, Martin

398 et al. (2020b) also found that the seagrass root microbiome was more enriched in *Ca.*

399 *Thiodiazotropha* spp. under stress conditions. Thus, we suggest that the *S. alterniflora – Ca.*

400 *Thiodiazotropha* symbiosis is an adaptive interaction to anoxic soil conditions, whereby the host

401 plant responds to stress from elevated dissolved sulfide concentrations. Further, we show that the

402 short *S. alterniflora* phenotype, which harbors a greater nitrogenase transcript abundance of *Ca.*

403 *Thiodiazotropha* sulfur chemosymbionts, also had the greatest rates of N fixation of all assessed

404 marsh compartments. Our results support the SGH in which greater stress strengthens the

405 relationship between sulfur oxidizing bacteria and macrophyte plants. Stressed plants benefit from

406 the symbiotic relationship through reduced sulfide toxicity and the coupling of sulfide oxidation

407 to nitrogen fixation, with nitrogen likely transferred to the plant host. Conversely, the tall *S.*

408 *alterniflora* phenotype showed a faster internal cycling of nitrogen, which is consistent with

409 previous studies showing greater rates of nitrogen mineralization in this zone of the ecosystem

410 (Figure 5, Rolando et al., 2022).

Intriguingly, we also found high transcript expression of RuBisCO from *Ca.* Thiodiazotropha gspp

in the root compartment of the stressed short phenotype of *S. alterniflora* (Supplementary Table

S5). When assessing the natural abundance of ^{13}C ($\delta^{13}C$) in the roots of the short *S. alterniflora*,

we found that it was significantly depleted compared to the $\delta^{13}C$ from the tall phenotype

(Supplementary Figure S10). As a C4 plant, *S. alterniflora* first fixes carbon by the action of PEP

carboxylase, which has lower discrimination against ^{13}C than RuBisCO (Farquhar, 1983). Thus,

we could speculate that the higher ^{13}C discrimination in the root of the short *S. alterniflora*

phenotype is a consequence of sulfur chemoautotrophic carbon fixation by RuBisCO. Consistent

with our hypothesis, a previous study by Hwang and Morris (1992) showed that the roots of *S.*

alterniflora fix carbon through an undeciphered dark reaction. Further, unlike other plant species,

S. alterniflora assimilates sulfur as sulfide (Carlson and Forrest, 1982), and under mild sulfidic

conditions, sulfur oxidation in *S. alterniflora* roots is mostly biological and was previously

hypothesized to be coupled with energetic gain (Lee et al., 1999). Interestingly, low doses of

sulfide (< 1 mM) have been shown to stimulate the productivity of *S. alterniflora* (Mendelssohn

and Morris, 2000). However, since other plant physiological processes could also explain

differences in $\delta^{13}C$, such as differing rates of dark respiration, and leakiness from the bundle sheath

cells to the mesophyll cell (Farquhar, 1983), further investigations are required to quantify the

relevance of sulfur chemolithoautotrophy in *S. alterniflora* root biomass production.

4. Materials and methods

4.1 Study site, field sampling and sample processing

The field fraction of this study was carried out in the Georgia Coastal Ecosystem - Long Term

Ecological Research (GCE-LTER) site 6, located on Sapelo Island, GA (Lat: 31.389° N, Long:

81.277° W). Four ~100 m transects along the tall to short *Spartina alterniflora* gradient were

studied in July 2018, 2019, and 2020 (Figure 1, further site description in Rolando et al., 2022). A combination of multi-omics approaches and biogeochemical rate measurements were performed across different salt marsh compartments and phenotypes of *S. alterniflora*.

The microbiome of *S. alterniflora* was analyzed using shotgun metagenomics from three compartments: sediment, rhizosphere, and root. The tall and short *S. alterniflora* extremes of the four studied transects were sampled (n: 4 transects * 2 *S. alterniflora* phenotypes * 3 compartments = 24). Samples from transects 1 and 2 were collected in July 2018, while those from transects 3 and 4 were collected in July 2019. All sampling was performed at the 0-5 cm depth profile. Root-associated samples were washed two times with creek water in the field to remove coarse chunks of sediment attached to the plant. All samples were immediately flash-frozen in an ethanol and dry ice bath, and stored at -80ºC until DNA extraction.

Samples for metatranscriptomic analysis and biogeochemical rate measurements were collected in July 2020 only from transect 4 to reduce plant disturbance time before flash freezing and incubations, respectively. Five independent plants at least 3 meters apart were sampled with a shovel at the two *S. alterniflora* biomass extremes. The entire plant, including the undisturbed root system with sediment attached, was sampled up to at least 20 cm depth, transferred to a 5 gallons bucket, and immediately transported to the field lab. For each plant, a paired sample of sediment was collected. In the lab, roots were washed with creek water two times. After the first wash, roots with sediment attached were collected and defined as the rhizosphere + root compartment. Live, sediment-free roots were sampled after the second wash and defined as the root compartment. Top 5 cm from each sample were homogenized and used for total RNA extractions and measurements of nitrogen fixation and potential nitrification rates. Root samples for RNA extractions were washed in an epiphyte removal buffer, as in Simmons et al. (2018). Sediment and root samples for

RNA analysis were immediately flash-frozen in an ethanol dry ice bath, and stored at -80 ºC until

extraction. Due to limited sample amount, we did not prepare rhizosphere libraries for

metatranscriptomic analysis.

<u>4.2 Nitrogen fixation and potential nitrification rates</u>

Rates of N fixation were calculated in $^{15}N_2$ incubations in 14 ml serum vials as in Leppänen et al.

(2013). Rates were measured for all samples under both oxic and anoxic conditions. About 2 g,

and 1 g of wet weight was used for tracer gas incubations of sediment and root samples,

respectively. An additional subsample of 2 g was immediately oven-dried at 60 ºC for 72 hours to

be used as a pre-incubation control. In all vials containing root samples, 4 ml of autoclaved and

0.22 um filter-sterilized artificial seawater was added to avoid tissue desiccation. No artificial

seawater was added to sediment and rhizosphere samples since they were already water-saturated.

In anoxic incubations, vials were flushed with N_2 gas for 5 minutes. After sealing all incubation

vials, 2.8 ml of gas was removed and immediately replaced with 2.8 ml of $^{15}N_2$ (98% enriched,

Cambridge Isotope Laboratories Inc, USA). Vials were overpressured by adding 0.3 ml of air or

N_2 gas in oxic or anoxic incubations, respectively. The incubations were carried out in the dark

and at room temperature for 24 hours. At the end of the incubation, samples were oven dried at 60

ºC for 72 hours, and ground using a PowerGen high throughput homogenizer (Fisherbrand,

Pittsburgh, PA). Ground samples were sent to the University of Georgia Center for Applied Isotope

Studies (https://cais.uga.edu/) for carbon and nitrogen elemental analysis and ^{13}C and ^{15}N stable

isotope analysis. Elemental analysis was performed by the micro-Dumas method, while isotopic

analysis by isotope ratio mass spectrometry. Rates of ^{15}N incorporation were calculated per dry

weight basis (DW) as in Leppänen et al. (2013):

N uptake rate $\left(\text{nmol g}^{-1}\text{ DW h}^{-1}\right)=$

480
$$\frac{\frac{1}{100}\times\frac{\%N}{100}\times\left[\frac{\text{atom}\%_{sample}-\text{atom}\%_{control}}{MW(N_2)}\right]\times10^9\times\frac{100}{\text{atom}\%_{headspace}}}{\text{hours}}\qquad[1]$$

Where %N is the N percent concentration of the oven-dried sample, and $MW(N_2)$ is the molecular

weight of N_2 (28.013446).

Potential nitrification rates were measured as detailed by Dollhopf et al. (2005) in 5 replicates for

each tall and short *S. alterniflora* compartments: sediment, rhizosphere + root, and root.

Homogenized samples (5 g, 5 g, and 2 g for sediment, rhizosphere + root, and root compartment,

respectively) were placed in 250-ml Erlenmeyer flasks filled with 50 ml sterile artificial seawater.

Sodium chlorate and ammonium sulfate were amended to all flasks to final concentrations of 10

mM and 500 µM, respectively. Flasks were incubated in the dark at room temperature and orbital

shaking (115 rpm) for 24 hours. Subsamples for NO_2 analysis were taken at 0, 4, 8, 12, 18, and 24

hours. Timepoint samples were filtered with 0.2 µm syringe filters, and stored at -20 ºC until

analysis. Filtered NO_2 concentration was measured by spectrophotometry (García-Robledo et al.,

2014). Potential nitrification rates (N nmol g^{-1}_{DW} h^{-1}) were calculated as the slope of the linear

regression of NO_2 concentration over time.

4.3 Nucleic acid extractions and multi-omics library preparation

For DNA extractions, compartment separation of the rhizosphere and root microbiomes was

performed by sonication in an epiphyte removal buffer, as detailed in Simmons et al. (2018).

Extracellular dissolved or sediment-adsorbed DNA was removed from bulk sediment samples

according to the Lever et al. (2015) procedure. DNA extractions from all samples were performed

using the DNeasy PowerSoil kit (Qiagen, Valencia, CA) following the manufacturer's instructions.

Shotgun metagenome sequencing was performed on an Illumina NovaSeq 6000 S4 2x150 Illumina flow cell at the Georgia Tech Sequencing Core (Atlanta, GA).

RNA from 4 biological replicates of both the sediment and root compartments from the tall and short phenotypes of *S. alterniflora* was extracted using the ZymoBIOMICS RNA Miniprep (Zymo Research Corp) kit according to the manufacturer's protocol (n: 4 replicates * 2 *S. alterniflora* phenotypes * 2 compartments = 16 samples). Rigorous DNA digestion was done with the TURBO DNase kit (Invitrogen), and eukaryotic mRNA was removed by binding and discarding the eukaryotic mRNA polyA region to oligo d(T)25 magnetic beads (England Biolabs). Finally, rRNA from both plant and prokaryotic organisms was depleted using the QIAseq FastSelect -rRNA Plant, and -5S/16S/23S kits, respectively (Qiagen, Valencia, CA). Metatranscriptomic libraries were sequenced in two lanes of Illumina's NovaSeq 6000 System flow cell utilizing the NovaSeq Xp workflow (SE 120bp) at the Georgia Tech Sequencing Core (Atlanta, GA). We retrieved 203, and 415 Gpb of metagenomic and metatranscriptomic raw sequences, respectively; with a median sequencing effort of 7.8, and 25.3 Gbp per metagenomic and metatranscriptomic library (Supplementary Table S1). After quality control and *in silico* removal of host and rRNA reads, median sequencing effort decreased to 5.3 and 9 Gbp per metagenomic and metatranscriptomic library, respectively (Supplementary Table S1).

4.4 Gene and transcript quantification of the prokaryotic 16S rRNA gene

Prokaryotic abundance and a proxy of prokaryotic activity were measured by quantitative polymerase chain reaction (qPCR) and Reverse Transcription-qPCR (RT-qPCR) of the SSU rRNA gene, respectively. All samples used for metagenome and metatranscriptome analysis were quantified by qPCR and RT-qPCR, respectively. Samples were analyzed in triplicate using the StepOnePlus platform (Applied Biosystems, Foster City, CA, USA) and PowerUp SYBR Green

Master Mix (Applied Biosystems, Foster City, CA, USA). Reactions were performed in a final

volume of 20µl using the standard primer set for the prokaryotic SSU rRNA gene: 515F (5′-

GTGCCAGCMGCCGCGGTAA′) and 806R (5′-GGACTACHVGGGTWTCTAAT′) (Caporaso

et al., 2011, Rolando et al., 2022). To avoid plant plastid and mitochondrial DNA/cDNA

amplification from root samples, peptide nucleic acid PCR blockers were added to all qPCR and

RT-qPCR reactions at a concentration of 0.75 µM (Lundberg et al., 2013). Standard calibration

curves were performed using a 10-fold serial dilution (10^3 to 10^8 molecules) of standard pGEM-T

Easy plasmids (Promega, Madison, WI, USA) containing target sequences from *Escherichia coli*

K12. Melting curve analyses was used to check for PCR specificity. Prokaryotic gene and

transcript abundance of the SSU rRNA gene were calculated as gene and transcript copy number

g^{-1} of fresh weight, respectively.

4.5 Metagenomic and metatranscriptomic quality control

Metagenomic and metatranscriptomic raw reads were quality trimmed (quality phred score < 20),

and filtered for Illumina artifacts, PhiX, duplicates, optical duplicates, homopolymers, and

heteropolymers using JGI's BBTools toolkit (Bushnell, 2014). Reads shorter than 75 bp were

removed, and the quality of both metagenomic and metatranscriptomic libraries was assessed with

FastQC (Andrews, 2010). Remaining reads were mapped against the only publicly available *S.*

alterniflora genome (NCBI BioProject: PRJNA479677) with bowtie2 (Langmead and Salzberg,

2012), followed by removal of short reads that aligned to the *S. alterniflora* genome using samtools

(parameters: view -u -f 12 -F256, Li et al., 2009). In addition, rRNA reads from metatranscriptomic

samples were removed using sortMeRNA v.4.3.4 (Kopylova et al., 2012). Finally, DNA

contamination in metatranscriptomic samples was assessed as indicated by Johnston et al. (2019).

Short-read transcripts were mapped to all assembled contigs, and strand-specificity (consistency

in sense/antisense orientation) was calculated for all genes with more than 100 hits. All assessed

samples had a greater than 95% average strand-specificity, thus considered to be free of DNA

contamination (Supplementary Table S1). Filtered, quality-trimmed, and host-free reads were

utilized for subsequent analyses.

<u>4.6 Metagenomic and metatranscriptomic short reads functional analysis and nonpareil diversity</u>

Short reads from all metagenomic and metatranscriptomic samples were aligned against the

eggNOG protein database (release 5.0.2) using eggnog-mapper v.2.1.9 (Cantalapiedra et al., 2021).

The DIAMOND tabular outputs were filtered by retrieving only the best hit based on bitscore. Hits

with less than 30% identity, less than 30% match of the read length, or that did not match a

prokaryotic domain were removed from the analysis. Gene profiles for each metagenomic and

metatranscriptomic library were constructed by counting hits of predicted KEGG orthology. Due

to differences in sequencing effort and genome size between multi-omics libraries, the count

matrixes were normalized by dividing them by their median count of 10 universal single-copy

phylogenetic marker genes (K06942, K01889, K01887, K01875, K01883, K01869, K01873,

K01409, K03106, and K03110) as in Salazar et al. (2019). Functional gene and transcript profiles

were analyzed by an principal coordinate analysis utilizing the Bray-Curtis dissimilarity distance.

Further, for both metagenomic and metatranscriptomic profiles, multivariate variation of the Bray-

Curtis dissimilarity matrix was partitioned to compartment (sediment, rhizosphere, and root) and

S. alterniflora phenotype, based on a permutational multivariate analysis of variance

(PERMANOVA) with 999 permutations performed in vegan v. 2.5.7 (Oksanen et al., 2013). The

normalized relative abundance of selected terminal oxidases and genes/transcripts from the carbon,

nitrogen, and sulfur cycles were assessed by compartment and *S. alterniflora* phenotype.

After removing reads annotated as Eukaryotic by eggnog-mapper, we calculated nonpareil

diversity from all metagenomic samples using nonpareil v. 3.401 (Rodriguez-R et al., 2018b).

Nonpareil diversity is a metric estimated from the redundancy of whole genome sequencing reads,

and has been shown to be closely related to classic metrics of microbial alpha diversity like the

Shannon index.

<u>4.7 Recovery of metagenome assembled genomes (MAGs)</u>

The following binning approach was motivated by a recently proposed method of iteratively

subtracting reads mapping to MAGs, re-assembling, and re-binning metagenomic libraries to

increase the number of recovered genomes (Rodriguez-R et al., 2020). An initial assembly using

idba-ud v1.1.3 with pre-error-correction for highly uneven sequencing depth was performed for

all individual metagenomic libraries (default parameters), as well as co-assemblies grouping

libraries from the same *S. alterniflora* phenotype and compartment (--mink 40 --maxk 120 --step

20 --min_contig 300) (Peng et al., 2012). The resulting contigs from both individual and co-

assemblies were binned using three different algorithms with default options: MaxBin v.2.2.7,

MetaBAT v.2.15, and CONCOCT v1.1.0 (Alneberg et al., 2014; Wu et al., 2016; Kang et al.,

2019). Recovered bins were dereplicated with DAS Tool v1.1.2 (Sieber et al., 2018), and the output

was refined for putative contamination with MAGPurify v2.1.2 (Nayfach et al., 2019).

Completeness, contamination, and quality score (Completeness – 5*Contamination) were

calculated with MiGA v0.7 (Rodriguez-R et al., 2018a). Recovered MAGs with a quality score

less than 50 were considered low quality and discarded. A second round of assembly and binning

was performed after discarding short reads that mapped against high-quality (HQ) MAGs.

Metagenomic libraries were mapped against HQ MAGs using bowtie2 with default parameters.

Paired reads that mapped against HQ MAGs were discarded using samtools (parameters: view -F

2, Li et al., 2009). Filtered metagenomic libraries were co-assembled once again using idba-ud

v1.1.3 with pre-error-correction for highly uneven sequencing depth (parameters: --mink 40 --

maxk 120 --step 20 --min_contig 300) in four groups: i) tall *S. alterniflora* root, ii) short *S.*

alterniflora root, iii) tall *S. alterniflora* sediment and rhizosphere, and iv) short *S. alterniflora*

sediment and rhizosphere. Binning, MAGs' refinement, and quality control were performed as

explained for the first iteration. Finally, HQ MAGs from both iterations were grouped into

genomospecies (gspp, singular gsp) by clustering MAGs with ANI > 95% using the MiGA v0.7

derep_wf workflow (Rodriguez-R et al., 2018a). MAGs with the highest quality score were

selected as representatives of their gsp, and most downstream analyses were performed with them.

Genomospecies relative abundance was estimated for each metagenomic library as in Rodriguez-

R et al. (2020). Sequencing depth was calculated per position using bowtie2 with default

parameters (Langmead and Salzberg, 2012), and bedtools genomecov (parameters: -bga, Quinlan

and Hall, 2010). Bedtools output was truncated to keep only the central 80% values, and the mean

of all retained positions was calculated using BedGraph.tad.rb from the enveomics collection, a

metric defined as TAD_{80} (truncated average sequencing depth) (Rodriguez-R and Konstantinidis,

2016). Relative abundance of each gsp was calculated by dividing TAD_{80} by genome equivalents

estimated for each metagenomic library with MicrobeCensus (Nayfach and Pollard, 2015).

<u>4.8 MAGs taxonomy, phylogenetic analysis, and gene functional annotation and expression</u>

Taxonomic classification of all MAGs was performed by GTDB-Tk v2.1.0 using the reference

database GTDB R07-RS207 (Chaumeil et al., 2022). A maximum likelihood phylogenetic tree of

all gspp was constructed using a 400 universal marker genes database in PhyloPhlAn v3.0.58

(parameters: -d phylophlan, --msa mafft, --trim trimal, --map_dna diamond, --map_aa diamond, -

-tree1 iqtree, --tree2 raxml, --diversity high, --fast, Asnicar et al., 2020). The phylogenetic tree was

decorated and visualized in ggtree v2.0.4 (Yu et al., 2018).

Protein-encoding genes from all binned MAGs were predicted with Prodigal v.2.6.3 using default

parameters (Hyatt et al., 2010), and resulting amino acid sequences aligned against the eggNOG

database (release 5.0.2) using eggnog-mapper v2.1.9 (Cantalapiedra et al., 2021). The DIAMOND

tabular output was filtered by retrieving only the best hit based on bitscore, and removing hits with

less than 30% identity and/or less than 50% match length.

Metagenomic and metatranscriptomic short reads were mapped against functionally annotated

ORFs of all gspp using megablast v.2.10.1. Only hits with greater than 95% percent identity and

90% read alignment were retained. Percent contribution of genes and transcripts to the total

microbial community was assessed by dividing the number of hits from each gsp against the total

number of functionally annotated short reads (from section 4.7). When assessing the transcript

expression profile of a specific gsp, the number of mapped hits per annotated transcript was

normalized by dividing it by gene length (bp) and the median abundance of 10 universal single-

copy phylogenetic marker genes of the prokaryotic community (K06942, K01889, K01887,

K01875, K01883, K01869, K01873, K01409, K03106, and K03110).

Finally, since the phylogeny of the dsrAB gene allows to discriminate between the oxidative and

reductive dsrAB types, we aligned all recovered genes from our binned gspp to a reference

alignment (Müller et al., 2015) with Clustal Omega v.1.2.4 (Sievers et al., 2011), and built an

approximately-maximum-likelihood phylogenetic tree with FastTree v.2.1.11 (Price et al., 2010).

Gene type was inferred based on placement in the phylogenetic tree.

<u>4.9 Phylogenetic reconstruction of the *Sedimenticolaceae* family</u>

Publicly available genomes from all species from the *Sedimenticolaceae* family, according to

GTDB R07-RS207, and all *Candidatus* Thiodiazotropha genomes from the Osvatic et al. (2023)

study were retrieved from NCBI. A full list and characteristics of genomes used for this analysis

is found in Supplementary Table S11. A phylogenetic tree of all retrieved genomes and binned

genomes from this study was constructed in PhyloPhlAn v3.0.58 using a 400 universal marker

genes database (parameters: -d phylophlan, --msa mafft, --trim trimal, --map_dna diamond, --

map_aa diamond, --tree1 fasttree, --tree2 raxml, --diversity low, --accurate, Asnicar et al., 2020).

The phylogenetic tree was decorated and visualized using ggtree v2.0.4 (Yu et al., 2018).

<u>4.10 Analysis of root microbiomes from contrasting ecosystems</u>

Publicly available 16S rRNA gene amplicon datasets, generated from next-generation sequencing,

were used to characterize the community assembly of root microbiomes from coastal marine

ecosystems. Studies were selected based on google scholar queries using a combination of the

following keywords: "salt marsh", "coastal", "wetland", "mangrove", "seagrass", "seabed", "crop",

"bog", "plant", "root", "endosphere", "microbial community", "microbiome", "amplicon", "next-

generation sequencing", "16S rRNA", and "SSU rRNA", as well as based on the authors prior

knowledge. Only studies that collected environmental samples were included (i.e., no greenhouse

or plants grown on potting media were included). When available, paired soil/sediment and

rhizosphere samples were also retrieved. Twenty-two studies that met our requirements were

selected, collecting a total of 2911 amplicon samples, with 1182 of them being from the root

compartment across 56 different plant species. Selected plants were categorized into 4 different

ecosystem types: seabed, coastal wetlands, freshwater wetlands, and other terrestrial ecosystems.

A complete list of selected amplicon samples with accompanying metadata is available in

Supplementary Table S9.

Cutadapt v.3.7 was used to detect and remove primer sequences from the datasets (Martin, 2011).

Primer-free sequences were quality filtered using DADA2's filterAndTrim function [options:

truncLen=c(175,150), maxN=0, maxEE=c(2,2), truncQ=10, rm.phix=TRUE] (Callahan et al.,

2016). Trimmed reads were randomly subsampled to a maximum of 20,000 reads using seqtk v.1.3

(Li, 2012) and used as input for dada2 amplicon sequence variant (ASV) calling (Callahan et al.,

2016). Chimeras were removed using the removeBimeraDenovo function from the DADA2

package. Taxonomy was assigned to ASVs utilizing the Ribosomal Database Project (RDP) Naive

Bayesian Classifier (Wang et al., 2007) against the SILVA SSU rRNA reference alignment

[Release 138, (Quast et al., 2012)]. Sequences classified as chloroplast, mitochondrial, and

eukaryotic or that did not match any taxonomic phylum were excluded from the dataset. Samples

that had less than 5,000 reads were removed at this stage. In order to merge studies from different

sequence runs and primer sets, we grouped ASVs at the genus level before merging all studies into

a single dataset in phyloseq v.1.36 (McMurdie and Holmes, 2013).

Microbial community assembly of the root microbiome was analyzed by performing a non-metric

multidimensional scaling (NMDS) ordination utilizing the Bray-Curtis dissimilarity distance.

Multivariate variation of the Bray-Curtis dissimilarity matrix was partitioned to ecosystem type,

and compartment (soil/sediment, rhizosphere, and root), based on a PERMANOVA with 999

permutations performed in vegan v. 2.5.7 (Oksanen et al., 2013). Finally, putative function for

sulfate reduction and sulfur oxidation was assigned based on homology at the genus level as in

Rolando et al. (2022).

Data availability

Raw metagenomic and metatranscriptomic sequences are deposited in the BioProject database

(http://ncbi.nlm.nih.gov/bioproject) under accessions PRJNA703972 and PRJNA950121,

respectively. Metagenome-assembled genomes are found in the BioProject PRJNA703972.

Acompanying metadata and custom scripts used in the present study are publicly available in

Supplementary Information and in a Zenodo repository: https://doi.org/10.5281/zenodo.7883423

Acknowledgements

This work was supported in part by an institutional grant (NA18OAR4170084) to the Georgia Sea

Grant College Program from the National Sea Grant Office, National Oceanic and Atmospheric

Administration, US Department of Commerce, and by a grant from the National Science

Foundation (DEB 1754756). Any opinions, findings, and conclusions or recommendations

expressed in this material are those of the authors and do not necessarily reflect the views of the

National Science Foundation.

Author contributions

692 Conceptualization: J.L.R., M.K., J.T.M., and J.E.K. Field sampling and analysis: J.L.R., M.K.,

693 T.S., Y.L., and J.E.K. Bioinformatic analysis: J.L.R., M.K., P.P., R.C., and K.T.K. Funding

694 acquisition: J.E.K. Writing—original draft: J.L.R. with inputs from all authors.

Ethics declarations

696 The authors declare no competing interests.

References:

1. Alneberg, J., Bjarnason, B.S., De Bruijn, I., Schirmer, M., Quick, J., Ijaz, U.Z., Lahti, L.,
Loman, N.J., Andersson, A.F. and Quince, C., 2014. Binning metagenomic contigs by
coverage and composition. Nature methods, 11(11), pp.1144-1146.
2. Andrews, S., 2010. FastQC: a quality control tool for high throughput sequence data.
3. Asnicar, F., Thomas, A.M., Beghini, F., Mengoni, C., Manara, S., Manghi, P., Zhu, Q.,
Bolzan, M., Cumbo, F., May, U., Sanders, J.G., 2020. Precise phylogenetic analysis of
microbial isolates and genomes from metagenomes using PhyloPhlAn 3.0. Nature
communications, 11:1-10.

4. Barbier, E.B., Hacker, S.D., Kennedy, C., Koch, E.W., Stier, A.C. and Silliman, B.R., 2011. The value of estuarine and coastal ecosystem services. Ecological monographs, 81(2), pp.169-193.

5. Bertness, M.D. and Callaway, R., 1994. Positive interactions in communities. Trends in ecology & evolution, 9(5), pp.191-193.

6. Bulgarelli, D., Schlaeppi, K., Spaepen, S., Van Themaat, E.V.L., Schulze-Lefert, P., 2013. Structure and functions of the bacterial microbiota of plants. Annual review of plant biology, 64:807-838.

7. Bushnell, B., 2014. BBTools software package. URL http://sourceforge. net/projects/bbmap.

8. Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., Holmes, S.P., 2016. DADA2: High-resolution sample inference from Illumina amplicon data. Nature methods, 13:581-583.

9. Cantalapiedra, C.P., Hernández-Plaza, A., Letunic, I., Bork, P., Huerta-Cepas, J., 2021. eggNOG-mapper v2: functional annotation, orthology assignments, and domain prediction at the metagenomic scale. Molecular biology and evolution, 38:5825-5829.

10. Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Lozupone, C.A., Turnbaugh, P.J., Fierer, N., Knight, R., 2011. Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. Proceedings of the national academy of sciences, 108:4516-4522.

11. Carlson Jr, P.R., Forrest, J., 1982. Uptake of dissolved sulfide by Spartina alterniflora: evidence from natural sulfur isotope abundance ratios. Science, 216:633-635.

12. Chaumeil, P.A., Mussig, A.J., Hugenholtz, P., Parks, D.H., 2022. GTDB-Tk v2: memory friendly classification with the genome taxonomy database. Bioinformatics, 38:5315-5316.

13. Crump, B.C., Wojahn, J.M., Tomas, F., Mueller, R.S., 2018. Metatranscriptomics and amplicon sequencing reveal mutualisms in seagrass microbiomes. Frontiers in microbiology, 9:388.

14. David, A.S., Thapa-Magar, K.B., Menges, E.S., Searcy, C.A., Afkhami, M.E., 2020. Do plant–microbe interactions support the Stress Gradient Hypothesis?. Ecology, 101:e03081.

15. Dollhopf, S.L., Hyun, J.H., Smith, A.C., Adams, H.J., O'Brien, S., Kostka, J.E., 2005. Quantification of ammonia-oxidizing bacteria and factors controlling nitrification in salt marsh sediments. Applied and environmental microbiology, 71:240-246.

16. Donnelly, J.P., Bertness, M.D., 2001. Rapid shoreward encroachment of salt marsh cordgrass in response to accelerated sea-level rise. Proceedings of the National Academy of Sciences, 98:14218-14223.

17. Duarte, C.M., Losada, I.J., Hendriks, I.E., Mazarrasa, I., Marbà, N., 2013. The role of coastal plant communities for climate change mitigation and adaptation. Nature climate change, 3:961-968.

18. Farquhar, G.D., 1983. On the nature of carbon isotope discrimination in C4 species. Functional Plant Biology, 10:205-226.

19. Fox-Kemper, B., Hewitt, H.T., Xiao, C., Aðalgeirsdóttir, G., Drijfhout, S.S., et al. 2021. Ocean, cryosphere and sea level change. In Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, ed. V Masson-Delmotte, P Zhai, A Pirani, SL Connors, C Péan, et al., pp. 1211–362. Cambridge, UK: Cambridge Univ. Press

20. Gandy, E.L., Yoch, D.C., 1988. relationship between nitrogen-fixing sulfate reducers and
fermenters in salt marsh sediments and roots of *Spartina alterniflora*. Applied and
Environmental Microbiology, 54:2031-2036.
21. García-Robledo, E., Corzo, A., Papaspyrou, S., 2014. A fast and direct spectrophotometric
method for the sequential determination of nitrate and nitrite at low concentrations in small
volumes. Marine Chemistry, 162:30-36.
22. Guimond, J.A., Yu, X., Seyfferth, A.L., Michael, H.A., 2020. Using hydrological-
biogeochemical linkages to elucidate carbon dynamics in coastal marshes subject to
relative sea level rise. Water Resources Research, 56:e2019WR026302.
23. Herbert, R.A., 1999. Nitrogen cycling in coastal marine ecosystems. FEMS microbiology
reviews, 23:563-590.
24. Holmer, M., Gribsholt, B., Kristensen, E., 2002. Effects of sea level rise on growth of
Spartina anglica and oxygen dynamics in rhizosphere and salt marsh sediments. Marine
Ecology Progress Series, 225:197-204.
25. Howarth, R.W., 1984. The ecological significance of sulfur in the energy dynamics of salt
marsh and coastal marine sediments. Biogeochemistry, 1:5-27.
26. Hwang, Y.H. and Morris, J.T., 1992. Fixation of inorganic carbon from different sources
and its translocation in Spartina alterniflora Loisel. Aquatic botany, 43(2), pp.137-147.
27. Hyatt, D., Chen, G.L., LoCascio, P.F., Land, M.L., Larimer, F.W. and Hauser, L.J., 2010.
Prodigal: prokaryotic gene recognition and translation initiation site identification. BMC
bioinformatics, 11(1), p.119.
28. Johnston, E.R., Kim, M., Hatt, J.K., Phillips, J.R., Yao, Q., Song, Y., Hazen, T.C., Mayes,
M.A., Konstantinidis, K.T., 2019. Phosphate addition increases tropical forest soil
respiration primarily by deconstraining microbial population growth. Soil Biology and
Biochemistry, 130:43-54.
29. Jørgensen, B.B., Findlay, A.J., Pellerin, A., 2019. The biogeochemical sulfur cycle of
marine sediments. Frontiers in microbiology, 10:849.
30. Kang, D.D., Li, F., Kirton, E., Thomas, A., Egan, R., An, H. and Wang, Z., 2019. MetaBAT
2: an adaptive binning algorithm for robust and efficient genome reconstruction from
metagenome assemblies. PeerJ, 7, p.e7359.
31. Kirwan, M.L., Temmerman, S., Skeehan, E.E., Guntenspergen, G.R., Fagherazzi, S.,
2016. Overestimation of marsh vulnerability to sea level rise. Nature Climate Change,
6:253-260.
32. Kolton, M., Rolando, J.L. and Kostka, J.E., 2020. Elucidation of the rhizosphere
microbiome linked to Spartina alterniflora phenotype in a salt marsh on Skidaway Island,
Georgia, USA. FEMS Microbiology Ecology, 96(4), p.fiaa026.
33. Koop-Jakobsen, K., Mueller, P., Meier, R.J., Liebsch, G., Jensen, K., 2018. Plant-Sediment
Interactions in Salt Marshes–An Optode Imaging Study of O_2, pH, and CO_2 Gradients in
the Rhizosphere. Frontiers in plant science, 9:541.
34. Kopylova, E., Noé, L., Touzet, H., 2012. SortMeRNA: fast and accurate filtering of
ribosomal RNAs in metatranscriptomic data. Bioinformatics, 28:3211-3217.
35. Kostka, J.E., Gribsholt, B., Petrie, E., Dalton, D., Skelton, H.,Kristensen, E., 2002. The
rates and pathways of carbon oxidation in bioturbated saltmarsh sediments. Limnology
and Oceanography, 47:230-240.

36. Kristensen, E., Alongi, D.M., 2006. Control by fiddler crabs (*Uca vocans*) and plant roots (*Avicennia marina*) on carbon, iron, and sulfur biogeochemistry in mangrove sediment. Limnology and Oceanography, 51:1557-1571.

37. Lamers, L.P., Govers, L.L., Janssen, I.C., Geurts, J.J., Van der Welle, M.E., Van Katwijk, M.M., Van der Heide, T., Roelofs, J.G., Smolders, A.J., 2013. Sulfide as a soil phytotoxin—a review. Frontiers in plant science, 4:268.

38. Langmead, B., Salzberg S., 2012. Fast gapped-read alignment with Bowtie 2. Nature Methods, 9:357-359.

39. Lee, R.W., Kraus, D.W., Doeller, J.E., 1999. Oxidation of sulfide by *Spartina alterniflora* roots. Limnology and Oceanography, 44:1155-1159.

40. Leppänen, S.M., Salemaa, M., Smolander, A., Mäkipää, R., Tiirola, M., 2013. Nitrogen fixation and methanotrophy in forest mosses along a N deposition gradient. Environmental and Experimental Botany, 90:62-69.

41. Lever, M.A., Torti, A., Eickenbusch, P., Michaud, A.B., Šantl-Temkiv, T., Jørgensen, B.B., 2015. A modular method for the extraction of DNA and RNA, and the separation of DNA pools from diverse environmental sample types. Frontiers in Microbiology 6:476.

42. Li, H., 2012. seqtk Toolkit for processing sequences in FASTA/Q formats. GitHub.

43. Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin, R., 2009. The sequence alignment/map format and SAMtools. Bioinformatics, 25:2078-2079.

44. Lim, S.J., Alexander, L., Engel, A.S., Paterson, A.T., Anderson, L.C., Campbell, B.J., 2019a. Extensive thioautotrophic gill endosymbiont diversity within a single *Ctena orbiculata* (Bivalvia: Lucinidae) population and implications for defining host-symbiont specificity and species recognition. MSystems, 4:e00280-19.

45. Lim, S.J., Davis, B.G., Gill, D.E., Walton, J., Nachman, E., Engel, A.S., Anderson, L.C., Campbell, B.J., 2019b. Taxonomic and functional heterogeneity of the gill microbiome in a symbiotic coastal mangrove lucinid species. The ISME journal, 13:902-920.

46. Lundberg, D.S., Yourstone, S., Mieczkowski, P., Jones, C.D., Dangl, J.L., 2013. Practical innovations for high-throughput amplicon sequencing. Nature methods, 10:999-1002.

47. Martin, M., 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet. journal, 17:10-12.

48. Martin, B.C., Middleton, J.A., Fraser, M.W., Marshall, I.P., Scholz, V.V., Hausl, B. and Schmidt, H., 2020a. Cutting out the middle clam: lucinid endosymbiotic bacteria are also associated with seagrass roots worldwide. The ISME journal, 14:2901-2905.

49. Martin, B.C., Alarcon, M.S., Gleeson, D., Middleton, J.A., Fraser, M.W., Ryan, M.H., Holmer, M., Kendrick, G.A., Kilminster, K., 2020b. Root microbiomes as indicators of seagrass health. FEMS Microbiology Ecology, 96:fiz201.

50. McGlathery, K.J., Risgaard-Petersen, N., Christensen, P.B., 1998. Temporal and spatial variation in nitrogen fixation activity in the eelgrass *Zostera marina* rhizosphere. Marine Ecology Progress Series, 168:245-258.

51. McMurdie, P.J. and Holmes, S., 2013. phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. PloS one, 8:e61217.

52. Mendelssohn, I.A. and Morris, J.T., 2000. Eco-physiological controls on the primary productivity of *Spartina alterniflora*. Concepts and Controversies in Tidal Marsh Ecology, pp.59-80.

53. Morris, J. T., Cahoon, D. R., Callaway, J. C., Craft, C., Neubauer, S. C., Weston, N. B.,
2021. Marsh Equilibrium Theory: Implications for Responses to Rising Sea Level. In:
FitzGerald, D. M. and Hughes, Z. J. (eds) Salt Marshes: Function, Dynamics, and
Stresses. Cambridge: Cambridge University Press, pp. 157–177.
54. Müller, A.L., Kjeldsen, K.U., Rattei, T., Pester, M., Loy, A., 2015. Phylogenetic and
environmental diversity of DsrAB-type dissimilatory (bi) sulfite reductases. The ISME
journal, 9:1152-1165.
55. Nayfach, S., Pollard, K.S. 2015. Average genome size estimation improves comparative
metagenomics and sheds light on the functional ecology of the human microbiome.
Genome Biology, 16:51.
56. Nayfach, S., Shi, Z.J., Seshadri, R., Pollard, K.S. and Kyrpides, N.C., 2019. insights from
uncultivated genomes of the global human gut microbiome. Nature, 568:505-510.
57. Nielsen, L.B., Finster, K., Welsh, D.T., Donelly, A., Herbert, R.A., De Wit, R., Lomstein,
B.A., 2001. Sulphate reduction and nitrogen fixation rates associated with roots, rhizomes
and sediments from *Zostera noltii* and *Spartina maritima* meadows. Environmental
Microbiology, 3:63-71.
58. Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin,
P.R., O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H.,
2020. vegan: Community Ecology Package. R package version 2.5-7. https://CRAN.R-
project.org/package=vegan
59. Osvatic, J.T., Wilkins, L.G., Leibrecht, L., Leray, M., Zauner, S., Polzin, J., Camacho, Y.,
Gros, O., van Gils, J.A., Eisen, J.A., Petersen, J.M., 2021. Global biogeography of
chemosynthetic symbionts reveals both localized and globally distributed symbiont groups.
Proceedings of the National Academy of Sciences, 118:e2104378118.
60. Osvatic, J.T., Yuen, B., Kunert, M., Wilkins, L., Hausmann, B., Girguis, P., Lundin, K.,
Taylor, J., Jospin, G., Petersen, J.M., 2023. Gene loss and symbiont switching during
adaptation to the deep sea in a globally distributed symbiosis. The ISME Journal, 17:453-
466.
61. Peng, Y., Leung, H.C., Yiu, S.M., Chin, F.Y., 2012. IDBA-UD: a de novo assembler for
single-cell and metagenomic sequencing data with highly uneven depth. Bioinformatics,
28:1420-1428.
62. Petersen, J.M., Kemper, A., Gruber-Vodicka, H., Cardini, U., Van Der Geest, M.,
Kleiner, M., Bulgheresi, S., Mußmann, M., Herbold, C., Seah, B.K., Antony, C.P., 2017.
Chemosynthetic symbionts of marine invertebrate animals are capable of nitrogen
fixation. Nature microbiology, 2:16195.
63. Petersen, J.M., Yuen, B., 2021. The symbiotic "all-rounders": partnerships between
marine animals and chemosynthetic nitrogen-fixing bacteria. Applied and Environmental
Microbiology, 87:e02129-20.
64. Price, M.N., Dehal, P.S. and Arkin, A.P., 2010. FastTree 2–approximately maximum-
likelihood trees for large alignments. PloS one, 5:e9490.
65. Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., Glöckner,
F.O., 2012. The SILVA ribosomal RNA gene database project: improved data processing
and web-based tools. Nucleic acids research, 41:D590-D596.
66. Quinlan, A.R., Hall, I.M., 2010. BEDTools: a flexible suite of utilities for comparing
genomic features. Bioinformatics, 26:841-842.

67. Rodriguez-R, L.M., Konstantinidis, K.T., 2016. The enveomics collection: a toolbox for specialized analyses of microbial genomes and metagenomes. PeerJ Prepr 4: e1900v1.

68. Rodriguez-R, L.M., Gunturu, S., Harvey, W.T., Rosselló-Mora, R., Tiedje, J.M., Cole, J.R., Konstantinidis, K.T., 2018a. The Microbial Genomes Atlas (MiGA) webserver: taxonomic and gene diversity analysis of Archaea and Bacteria at the whole genome level. Nucleic acids research, 46:W282-W288.

69. Rodriguez-R, L. M., Gunturu, S., Tiedje, J. M., Cole, J. R., Konstantinidis, K. T., 2018b. Nonpareil 3: fast estimation of metagenomic coverage and sequence diversity. MSystems, 3:e00039-18.

70. Rodriguez-R, L.M., Tsementzi, D., Luo, C., Konstantinidis, K.T., 2020. Iterative subtractive binning of freshwater chronoseries metagenomes identifies over 400 novel species and their ecologic preferences. Environmental Microbiology, 22:3394-3412.

71. Rolando, J.L., Kolton, M., Song, T., Kostka, J.E., 2022. The core root microbiome of *Spartina alterniflora* is predominated by sulfur-oxidizing and sulfate-reducing bacteria in Georgia salt marshes, USA. Microbiome, 10:37.

72. Salazar, G., Paoli, L., Alberti, A., Huerta-Cepas, J., Ruscheweyh, H.J., Cuenca, M., Field, C.M., Coelho, L.P., Cruaud, C., Engelen, S., Gregory, A.C., 2019. Gene expression changes and community turnover differentially shape the global ocean metatranscriptome. Cell, 179:1068-1083.

73. Sieber, C.M., Probst, A.J., Sharrar, A., Thomas, B.C., Hess, M., Tringe, S.G., Banfield, J.F., 2018. Recovery of genomes from metagenomes via a dereplication, aggregation and scoring strategy. Nature microbiology, 3:836-843.

74. Sievers, F., Wilm, A., Dineen, D., Gibson, T.J., Karplus, K., Li, W., Lopez, R., McWilliam, H., Remmert, M., Söding, J., Thompson, J.D., 2011. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. Molecular systems biology, 7:539.

75. Simmons, T., Caddell, D.F., Deng, S., Coleman-Derr, D., 2018. Exploring the root microbiome: extracting bacterial community data from the soil, rhizosphere, and root endosphere. Journal of visualized experiments: JoVE, 135.

76. Spivak, A.C., Reeve, J., 2015. Rapid cycling of recently fixed carbon in a *Spartina alterniflora* system: a stable isotope tracer experiment. Biogeochemistry, 125:97-114.

77. Thomas, F., Giblin, A.E., Cardon, Z.G., Sievert, S.M., 2014. Rhizosphere heterogeneity shapes abundance and activity of sulfur-oxidizing bacteria in vegetated salt marsh sediments. Frontiers in microbiology, 5:309.

78. Trivedi, P., Leach, J.E., Tringe, S.G., Sa, T., Singh, B.K., 2020. Plant–microbiome interactions: from community assembly to plant health. Nature reviews microbiology, 18:607-621.

79. Wang, Q., Garrity, G.M., Tiedje, J.M., Cole, J.R., 2007. Naive Bayesian classifier for rapid assignment of rRNA sequences into the bacterial taxonomy. Applied and environmental microbiology, 73:5261-5267.

80. Welsh, D.T., Bourgues, S., de Wit, R., Herbert, R.A., 1996. Seasonal variations in nitrogen-fixation (acetylene reduction) and sulphate-reduction rates in the rhizosphere of *Zostera noltii*: nitrogen fixation by sulphate-reducing bacteria. Marine Biology, 125:619-628.

81. Whiting, G.J., Gandy, E.L., Yoch, D.C., 1986. Tight coupling of root-associated nitrogen fixation and plant photosynthesis in the salt marsh grass *Spartina alterniflora* and carbon

dioxide enhancement of nitrogenase activity. Applied and environmental microbiology, 52:108-113.

82. Wu, Y.W., Simmons, B.A., Singer, S.W., 2016. MaxBin 2.0: an automated binning algorithm to recover genomes from multiple metagenomic datasets. Bioinformatics, 32:605-607.

83. Yu, G., Lam, T.T.Y., Zhu, H., Guan, Y., 2018. Two methods for mapping and visualizing associated data on phylogeny using ggtree. Molecular biology and evolution, 35:3041-3043.

Figure 1. *Spartina alterniflora* biomass gradient as a natural laboratory. A gradient in *S.*
alterniflora aboveground biomass is commonly observed with tall plants growing at the levees
next to large tidal creeks and a short phenotype of *S. alterniflora* dominating the interior of the
marsh. Sediments from the tall *S. alterniflora* zone are characterized as a more oxidized
environment with higher levels of iron, coupled nitrification-denitrification as well as higher
rates of organic matter hydrolysis and mineralization. Conversely, sediments from the short
phenotype tend to be more chemically reduced, with higher rates of sulfate reduction, elevated
porewater salinity, and less bioturbation and tidal flushing. Roots from the short *S. alterniflora*
phenotype have been proposed to harbor sulfur-oxidizing bacteria (SOB) that benefit the plant by
detoxifying the root environment.

Figure 2. Phylogenetic reconstruction of 160 dereplicated high-quality metagenome assembled
genomes (MAGs, > 50 quality-score) binned from *Spartina alterniflora* sediment, rhizosphere,
and root samples. Outer rim shows the presence/absence of genes for carbon fixation
(RuBisCO), nitrogen fixation (nifKDH), thiosulfate oxidation (soxABXYZ), dissimilatory sulfite
reduction/oxidation (dsrAB), nitrification (amoCAB), denitrification, and dissimilatory nitrate
reduction to ammonium (DNRA).

Figure 3. Drivers and metagenome assembled genomes (MAGs) associated with rates of N fixation and nitrification in the salt marsh environment. Rates of nitrification and N fixation (under anoxic and oxic conditions) per microbiome compartment and *Spartina alterniflora* phenotype (a). Normalized transcript relative abundance of the nitrogenase gene (nifK), and the reductive and oxidative dsrA types per microbiome compartment and *Spartina alterniflora* phenotype (b). Percentage from the total nitrogenase (nifK) and reductive and oxidative types of the dsrA transcripts mapping to the most active MAGs (c). Statistical significance based on non-parametric pairwise Mann-Whitney tests. n.s = p-value > 0.5, * p-value < 0.5, ** p-value < 0.01.

Figure 4. Phylogenetic tree of the Sedimenticolaceae family. Taxonomic annotation of the *Candidatus* Thiodiazotropha genus was based on Osvatic et al. (2023). The diagram next to the species name recognizes genomes recovered as symbionts of eukaryotic organisms. The average relative abundance of the metagenome assembled genomes (MAGs) from the present study is shown in the adjacent panels per microbiome compartment and *Spartina alterniflora* phenotype. Relative abundance was calculated at the DNA-level based on average coverage per position in metagenomic libraries. Purple sulfur bacteria *Allochromatium vinosum* was used as an outgroup for the phylogenetic tree.

Figure 5. Prokaryotic abundance, functional diversity, and activity are determined by *Spartina*
alterniflora phenotype and microbiome compartment. Average ± standard error of the
metagenomic nonpareil diversity, 16S rRNA gene and transcript abundance as quantified by
qPCR and RT-qPCR, respectively (n = 4) (a). Principal coordinate analysis (PCoA) ordination
plot based on the Bray-Curtis dissimilatory index of functional profiles from KEGG orthology
annotations (KO level) of metagenome and metatranscriptome libraries (b). Gene and transcript
normalized relative abundance of prokaryotic terminal oxidases and selected enzymes of the
carbon, nitrogen, and sulfur cycles (c). Different letter indicate statistical difference based on
pairwise Mann-Whitney tests (p-value < 0.5).

1023

1024

1025

Figure 6: Roots from marine influenced ecosystems assemble a distinct microbial community
enriched by bacteria with known to conserve energy from sulfur metabolism. Non-metric
multidimensional scaling (NMDS) ordination plot based on the Bray-Curtis dissimilatory index
of root-associated prokaryotic communities at the genus level, colored by ecosystem type (a).
Relative abundance of putative sulfur-oxidizing (b) and sulfate-reducing root bacteria (c) by
ecosystem type. Prokaryotic communities were characterized by analyzing an SSU rRNA gene
amplicon dataset of 1182 samples assessing roots from 56 plant species. Different letter indicate
statistical difference based on pairwise Mann-Whitney tests (p-value < 0.5).

1034

1035

1036

1037

Supplementary Figure S1. dsrAB phylogenetic tree of genes retrieved from metagenome-assembled genomes (MAGs). Genes were aligned to a reference database and tree was constructed with FastTree v.2.1.11. Oxidative and reductive types of the dsrAB gene were inferred based on phylogenetic placement in the constructed tree. Archean *Pyrobaculum aerophilum* was used as an outgroup.

1048 **Supplementary Figure S2**. Phylogenetic tree of the oxidative dsrAB type from metagenome
1049 assembled genomes (MAGs) aligned to a reference alignment database (Müller et al., 2015).
1050 Archean *Pyrobaculum aerophilum* was used as an outgroup of the phylogenetic tree.

1051

1052   **Supplementary Figure S3**. Gene expression of the nitrogenase (nifD, nifK, and nifH) and
1053   dissimilatory sulfite reductase genes (dsrA, dsrB, and aprA) are highly correlated among their
1054   own metabolic pathways.

1055

1056

1057

1058

1059

1060

1061

1062

1063

1064

1065 **Supplementary Figure S4.** Relation between the gene expression of the *Ca.* Thiodiazotropha
1066 sp. (gsp 31) oxidative, and *Desulfosarcinaceae* sp. (gsp. 68) reductive dsrA gene and the
1067 nitrogenase gene nifK in *Spartina alterniflora* root samples. Gene expression measured as the
1068 percentage of transcripts mapping to each genomospecies from the totality of functionally
1069 annotated short read transcripts.

1070

1071

1072

1073

1074

1075

1076

1077

1078

1079

1080

1081

1082

1083

1084

1085 **Supplementary Figure S5**. Gene and transcript relative abundance of the nitrogenase gene
1086 (nifK) partitioned at the finest taxonomic level as predicted by eggnog-mapper per microbiome
1087 compartment and *Spartina alterniflora* phenotype.

1088

1089

1090

1091

1092

1093

1094

1095

1096

1097

1098

1099

1100

1101 **Supplementary Figure S6.** Gene and transcript relative abundance of the oxidative dsrA gene
1102 partitioned at the finest taxonomic level as predicted by eggnog-mapper per microbiome
1103 compartment and *Spartina alterniflora* phenotype.

1104

1105

1106

1107

1108

1109

1110

1111

1112

1113

1114

1115

1116

1117 **Supplementary Figure S7.** Gene and transcript relative abundance of the reductive dsrA gene
1118 partitioned at the finest taxonomic level as predicted by eggnog-mapper per microbiome
1119 compartment and *Spartina alterniflora* phenotype.

1120

1121

1122

1123

1124

1125

1126

1127

1128

1129

1130

1131

1132

1133   **Supplementary Figure S8.** Relative abundance of selected metagenome assembled genomes
1134   (MAGs) from the *Desulfosarcinaceae* family across *Spartina alterniflora* phenotype and
1135   sampled compartment. Relative abundance was calculated at the DNA-level based on average
1136   coverage per position in metagenomic libraries. Different letter indicate statistical difference
1137   based on pairwise Mann-Whitney tests (p-value < 0.5).

1138

1139

1140

1141

1142

1143

1144

1145

1146

1147

1148

1149

1150

1151

1152

1153 **Supplementary Figure S9.** Relative abundance of amplicon sequence variants (ASVs)
1154 classified as members of the *Desulfosarcinaceae* and *Sedimenticolaceae* families, and the
1155 *Desulfatitalea* and *Ca.* Thiodiazotropha genera across soil/sediment, rhizosphere, and root
1156 samples from a seabed, coastal wetland, freshwater wetland and terrestrial ecosystems dataset.

1157

1158

1159

1160

1161

1162

1163   **Supplementary Figure S10.** Root carbon ^{13}C isotopic natural abundance by *Spartina*

1164   *alterniflora* phenotype. ^{13}C natural abundance was expressed as the per mille (‰) deviation from

1165   the Pee Dee Belemnite standard (PDB) ^{13}C:^{12}C ratio (δ^{13}C).

1166

1167

1168

1169