Rapid effects of valproic acid on the fetal brain transcriptome: Implications for brain development and autism

Susan G. Dorsey[1], Evelina Mocci[1], Malcolm V. Lane[2] and Bruce K. Krueger[3]*

[1]Department of Pain and Translational Symptom Sciences
University of Maryland School of Nursing, Baltimore, MD 21201
[2]Translational Toxicology/Department of Epidemiology and Public Health
University of Maryland School of Medicine, Baltimore, MD 21201
[3]Departments of Physiology and Psychiatry
University of Maryland School of Medicine, Baltimore, MD 21201

*Corresponding author
BKK Orcid: 0000-0002-0088-1931

ABSTRACT:

There is an increased incidence of autism among the children of women who take the anti-epileptic, mood stabilizing drug, valproic acid (VPA) during pregnancy; moreover, exposure to VPA *in utero* causes autistic-like symptoms in rodents and non-human primates. Analysis of RNAseq data obtained from fetal mouse brains 3 hr after VPA administration revealed that VPA significantly [p(FDR) ≤ 0.025] increased or decreased the expression of approximately 7,300 genes. No significant sex differences in VPA-induced gene expression were observed. Expression of genes associated with neurodevelopmental disorders such as autism as well as neurogenesis, axon growth and synaptogenesis, GABAergic, glutaminergic and dopaminergic synaptic transmission, perineuronal nets, and circadian rhythms was dysregulated by VPA. Moreover, expression of 400 autism risk genes was significantly altered by VPA as was expression of 247 genes that have been reported to play fundamental roles in the development of the nervous system, but are not linked to autism by GWAS. The goal of this study was to identify mouse genes that are: **(a) significantly up- or down-regulated by VPA in the fetal brain** and **(b) known to be associated with autism and/or to play a role in embryonic neurodevelopmental processes, perturbation of which has the potential to alter brain connectivity in the postnatal and adult brain.** The set of genes meeting these criteria provides potential targets for future hypothesis-driven approaches to elucidating the proximal underlying causes of defective brain connectivity in neurodevelopmental disorders such as autism.

INTRODUCTION:

Autism is a neurodevelopmental disorder characterized by social interaction deficits including language, and repetitive, stereotyped behavior with restricted interests (American Psychiatric Association, 2022). Autistic individuals may also display an increased incidence of anxiety, intellectual disability and seizures. The reported incidence of autism has increased dramatically over the last two decades (Zeidan et al., 2022) with some estimates now as high as 1:44 (Maenner et al., 2021). The severity of autism varies widely, ranging from high-functioning with minimal disability, to severely afflicted, where aggressive, self-destructive repetitive behaviors pose a threat to the safety of the patient, caregivers, and family. The term "autism spectrum disorders" (ASDs) reflects this symptomatic variability. The neurological mechanisms underlying ASDs are not understood nor is it known whether autism is a single disorder or multiple disorders sharing common core features.

Twin studies have revealed that 60 to 88% of the cases are inherited (Hallmayer et al., 2011; Rosenberg et al., 2009), however, many cases have been linked to *in utero* exposure to environmental factors such as pharmaceuticals, air pollution, chemical insecticides, and maternal infection (Landrigan, 2010; Rosenberg et al., 2009; Karimi et al., 2017).

The diagnosis of autism typically is made at around two years of age when the child fails to meet normal milestones for social development and language. However, it has become increasingly evident that autism arises before birth, with infants failing to show normal attention to faces (Macari et al., 2021). Epidemiological studies indicate that the onset of autism occurs during fetal development; one such study found that folate supplementation for pregnant women reduced the incidence of autism, but only when administered between two weeks before and four weeks after conception (Surén et al., 2013). Exposure to a variety of environmental factors, notably the anti-epileptic, mood stabilizing drug, valproic acid (VPA; Depakote®) (Christensen et al., 2013) and the organophosphate insecticide, chlorpyrifos (Burke et al., 2017), increase the incidence of autism and other neurodevelopmental disorders in the offspring of women who are exposed during pregnancy. Another environmental cause of autism is maternal immune activation (MIA) in which pregnant women who have systemic bacterial or viral infections with a hyper-immune response have an increased incidence of autism in their children (Bilbo et al., 2018; Careaga et al., 2017; Zawadzka et al., 2021).

Studying the etiology of autism in humans is limited to epidemiological approaches implicating genetic variants (single nucleotide polymorphisms, copy number variants, and single-gene syndromic mutations). The Simons Foundation Autism Research Initiative (SFARI) has compiled a list of 1,115 human

genes (https://gene.sfari.org/database/human-gene/) for which there is evidence for association with autism based on genome-wide association studies (GWAS), referred to as the "SFARI List" below.

Hypothesis-driven experiments to determine cause and effect need to be done in animal models, which display a range of behaviors that are remarkably similar to the core symptoms of autism (Silverman et al., 2010; Lan et al., 2017; Nicolini and Fahnestock, 2018). These autistic-like behaviors have been reported in transgenic animals lacking syndromic ASD genes such as *Cntnap2* and *Shank2* (Peñagarikano and Geschwind, 2012; Eltokhi et al., 2018), supporting the conclusion that pathogenic mutations in these genes are causative for syndromic autism and leading to the widespread use of these transgenic mice as animal models (Bey and Jiang, 2014; Kazdoba et al., 2016). Systemic administration of VPA (Nicolini and Fahnestock, 2018), chlorpyrifos (Lan et al., 2017) or induction of MIA (Careaga et al., 2017) in genetically normal pregnant rodents also leads to increased autistic-like behaviors in their offspring. Unlike transgenic mouse models, these environmental toxicity models of autism provide the opportunity to control the timing of exposure of the fetus to the toxic stressor, as will be discussed below. Gene expression analyses of the brains of mice in the VPA and MIA models have been conducted using RNA microarray analysis or next-generation RNA sequencing (RNAseq). In the majority of those animal studies, the fetus was exposed to the drug during mid-gestation, but RNA expression was analyzed in adult animals, typically following behavioral testing (e.g., see Qian et al., 2018; Zhao et al., 2019; Matsuo et al., 2022; Zhang et al., 2018). This approach has the potential to explain behavior in terms of contemporaneous gene expression but does not inform as to the initial molecular events occurring at the time the environmental stressor is present.

There are two competing, but not mutually exclusive, theories about the role of differential gene expression in autism. In one, abnormal gene expression in the child or adult at the time of behavioral testing, is responsible for autistic-like behavior. The second posits that abnormal gene expression, due to genomic variants or *in utero* environmental stressors, interferes with one or more critical steps in the "program" controlling fetal brain development. Such errors could carry forward throughout life leading to a brain with subtle anatomical or connectivity defects that underlie the abnormal development of an autistic brain. This has been referred to as a developmental "signature" (Ben-Ari, 2008). The present study focused on the second hypothesis by examining altered gene expression in fetal brains when the environmental stressor (VPA) is still present; these genes need not be continuously dysregulated throughout life. Consequently, genes associated with critical steps in early brain development such as neurogenesis, neuron fate specification, axon and dendrite growth, and synaptogenesis are of particular interest.

The incidence of autism is about 4-times higher in males than in females (Ferri et al., 2018). Sex differences in neural function are generally considered to be mediated by sex hormones acting from late prenatal brain development through to the adult. Since the time of VPA administration in the present study (E12.5) is just prior to the maturation of gonads and production of sex hormones (Ross and Capel, 2005), any sex differences in gene expression observed are likely due to sexually-dimorphic gene expression rather than to the effects of sex hormones.

The complexity and variability of the behavioral symptoms of autism together with identification of over 1,100 ASD-associated gene variants make a mechanistic understanding of the causes of the disorder a daunting task. One approach to this problem is to compare the results of multiple studies using GWAS results from human patients and gene expression results from animal models of autism, looking for common differentially expressed genes. The subset of genes in common from both types of studies can provide a short(er) list of candidate genes that could be subjected to future hypothesis-driven studies to establish causal relationships between genes and ASD symptoms.

The goal of the present study was to address this question using the VPA mouse model in which pregnant mice receive a single i.p. injection of VPA at gestational day 12.5 (E12.5). Pharmacokinetic studies have demonstrated that injected VPA dissipates within about 3−5 hours due to metabolism of the drug by the maternal liver (Nau and Löscher, 1982; Nau, 1986a); VPA levels in the fetal brain follow a similar trajectory (Nau, 1986b; Dickinson et al., 1980). Fetal brains were processed for RNAseq three hours after VPA administration to the pregnant dam; male and female fetuses were analyzed separately. Using this protocol, we propose that VPA-induced molecular events occurring on E12.5 are both necessary and sufficient for the autistic-like behaviors in VPA-treated animals assessed 5−6 weeks later. Analysis of these data revealed that VPA induced a significant increase or decrease in the expression of approximately 7,300 genes, of which 400 are among the 1,115 genes on the SFARI List and at least 247 additional genes are known to play fundamental roles in the development and function of the nervous system.

METHODS:

Timed-pregnant C57BL6 mice were generated by the University of Maryland School of Medicine Division of Veterinary Resources. All animal procedures were approved by the Institutional Animal Care and Use Committee. VPA was obtained from Sigma.

One male was paired overnight with two females; the day of separation was designated E0.5. On E12.5 pregnant females received i.p injections of VPA (400mg/kg) in sterile saline or saline alone. 7 pregnant females received VPA and 7 received saline. Three hours after VPA administration, pregnant females were euthanized by cervical dislocation and decapitation. Fetuses were transferred to ice-cold saline, decapitated and the entire brain was removed to Trizol and disrupted for 60 sec with a Bead Beater using 0.2 mm beads. The 105 individual fetal brain samples were frozen on dry ice and stored at −80°C prior to RNA extraction and quality control (RIN = 10 for all samples).

Sex was determined by analyzing *Sry* RNA derived from the torso of each fetus by PCR. Males were identified by the presence of both *Sry* and *Gapdh* RNA determined; *Gapdh* but not *Sry* RNA was expressed in females (Konopko et al., 2017). One male and one female brain from each of the 7 pregnancies was processed for RNAseq by the University of Maryland Institute for Genome Sciences.

Libraries were prepared from 25ng of RNA using the NEB Ultra II Directional RNA kit. Samples were sequenced on an Illumina NovaSeq 6000 with a 150bp paired-end read configuration. The quality of sequences was evaluated by using FastQC (Wingett and Andrews, 2018). The alignment was performed using HiSat (version HISAT2-2.0.4) (Kim et al., 2019) and *Mus musculus* GRCm38 as reference genome and annotation (version 102). Aligned bam files were used to determine number of reads by gene using HTSeq (Anders et al., 2015). On average, we sequenced 187,000,000 reads. 96.6% of them properly mapped the reference: 87% mapped exons, 2.4% mapped introns and the remaining 10.6% mapped intergenic regions.

Differential gene expression between mice treated with VPA and saline controls was conducted using DESeq2 (Love et al., 2014), which models gene counts using the negative binomial distribution. P-values were corrected for false discovery rate (FDR) using the Benjamini-Hochberg procedure; $p_{FDR} \leq 0.025$ and $\log_2$-fold change ≥ 0.32 was used as the criterion for significance. Differentially expressed genes due to VPA treatment were tested for enrichment in SFARI autism risk genes (https://gene.sfari.org/database/human-gene/) using Fisher's Exact Test.

Significance of sex-differences in FC was analyzed by 2-way ANOVA using the Limma-Voom tool (Law et al., 2014).

RESULTS:

RNAseq. 19,721 individual genes were analyzed in fetal mouse brain by RNAseq. Male and female brains were analyzed separately. The raw data showing base counts (mean of male and female counts) for each gene in each of seven independent fetal brain samples of each sex, each from a different pregnancy, are shown in Supplementary Table I (S1). n = 7 fetal brains from 7 different pregnancies for each condition. Only genes with base counts ≥100 were analyzed further. Genes that increased or decreased in response to VPA by < 1.25-fold and > −1.25-fold and those with adjusted p-values (p_{FDR}) ≥ 0.025 were also filtered out; a gene was included if VPA increased or decreased its expression by ≤ 1.25-fold and ≥ −1.25-fold in only one sex. Shown in Supplementary Table 2 (S2) are the 6516 genes significantly affected by VPA (p_{FDR} ≤ 0.025) in both sexes, 498 in females only and 280 in males only (listed in three separate tabs in Table S2). These apparent sex differences were due to variability in the replicates of one of the sexes (i.e., p_{FDR} > 0.025).

Throughout this report, a positive fold-change (FC) indicates that VPA increased gene expression; a negative FC indicates that VPA decreased expression such that the FC is the ratio of control to VPA gene expression. For example, FC = −3.0 indicates that V/C = 0.33, a 67% reduction by VPA.

Curating the mouse genes dysregulated by VPA in the fetal brain. The 7294 genes that were significantly altered by VPA (Table S2) were further analyzed in two ways. First, they were merged with the genes on the SFARI List identifying 400 common genes (Table 1). It should be noted that inclusion on the SFARI List is based on GWAS; in many cases, a mechanistic role in neither brain development nor the etiology of autism is known.

Second, we identified 247 genes significantly altered by VPA not on the list of SFARI risk genes but having a documented role in brain development including neurogenesis, synaptogenesis, and axon or dendrite growth (Tables 2 and 5). As will be discussed below, although these latter genes have not been associated with autism in GWAS, it is plausible that changes (either positive or negative) in the expression of their gene products in the fetal brain might adversely affect the trajectory of brain development leaving a permanent "signature" (Ben-Ari, 2008) on brain structure and connectivity leading to autistic-like behavior in the adult.

Table 1

Gene	Counts	Male FC	Male p_FDR	Female FC	Female p_FDR
Abat	8490	1.58	2.4E-08	1.68	1.7E-10
Abca7	4344	2.59	3.7E-24	2.58	7.2E-24
Abce1	14484	-1.25	2.2E-03	-1.20	1.3E-02
Abl2	6428	-1.33	1.3E-03	-1.25	1.4E-02
Ache	892	1.73	1.1E-03	1.82	3.1E-04
Actb	107135	-1.36	2.4E-08	-1.48	4.5E-13
Actn4	15736	1.43	3.2E-05	1.37	2.6E-04
Acy1	1770	1.70	2.3E-17	1.68	1.2E-16
Ada	284	1.30	1.9E-02	1.35	7.7E-03
Adcy3	1057	-1.69	1.4E-08	-1.81	1.1E-10
Adk	3332	1.30	1.4E-02	1.30	1.4E-02
Adnp	233	-1.63	4.0E-07	-1.58	3.0E-06
Adora2a	296	-1.25	4.9E-02	-1.36	4.9E-03
Ago3	5603	1.60	3.5E-07	1.90	1.9E-12
Agtr2	720	-6.44	2.7E-04	-4.45	4.1E-03
Ahi1	10423	1.57	1.1E-04	1.60	5.6E-05
Ahnak	9132	-3.07	1.2E-07	-2.64	5.6E-06
Akap9	18776	-2.30	5.3E-09	-1.85	2.4E-05
Aldoa	19582	1.32	1.1E-03	1.25	1.2E-02
Ankrd17	2867	-1.72	7.9E-13	-1.66	2.6E-11
Anxa1	480	-4.00	4.4E-05	-3.16	8.3E-04
Ap1s2	5451	1.35	1.5E-03	1.33	2.7E-03
Apba2	6951	-1.57	2.4E-04	-1.59	1.3E-04
Aph1a	4926	1.43	2.2E-08	1.31	2.6E-05
Arf3	12109	1.27	5.6E-06	1.20	9.6E-04
Arhgef2	9820	-1.95	1.2E-15	-2.08	9.6E-19
Arid1b	5659	-2.13	1.2E-16	-2.17	2.7E-17
Arid2	11217	-1.62	2.8E-15	-1.59	4.7E-14
Arntl2	511	1.71	3.1E-06	2.17	1.2E-11
Ash1l	9264	-1.69	1.4E-05	-1.40	6.9E-03
Asxl3	5208	-1.75	1.5E-10	-1.57	2.7E-07
Atp10a	473	-1.89	3.3E-16	-1.89	4.2E-16
Atp6v0a2	6136	1.42	2.4E-14	1.40	2.6E-13
Atrx	29927	-1.80	2.5E-06	-1.44	5.0E-03
Bace1	6271	1.83	4.3E-22	1.90	2.0E-24
Baiap2l1	386	3.78	2.5E-09	3.63	8.1E-09
Bcas1	279	-1.78	1.5E-03	-1.52	2.4E-02
Bcl11a	10078	-1.59	6.6E-05	-1.70	5.2E-06
Bcorl1	4918	1.89	1.9E-10	1.80	5.5E-09
Brinp2	2925	2.82	2.7E-12	2.61	1.2E-10
Brsk2	7140	-1.74	2.6E-04	-1.86	4.1E-05
Brwd3	6365	1.42	2.2E-03	1.66	4.9E-06
Btaf1	10163	1.41	1.9E-02	1.62	6.9E-04
Cacna1b	6214	1.34	4.7E-02	1.42	1.7E-02
Cacna1c	3446	-1.47	3.1E-03	-1.38	1.5E-02
Cacna2d1	9393	-1.84	2.7E-08	-1.59	3.0E-05
Cacnb2	498	-2.08	9.5E-13	-1.80	1.2E-08
Cadm1	12690	1.25	1.3E-02	1.28	5.3E-03
Camk2a	4142	1.94	1.6E-07	1.84	1.4E-06
Caprin1	3184	1.25	1.4E-03	1.31	1.0E-04
Cask	10174	-1.30	4.7E-03	-1.24	2.3E-02
Casz1	1306	2.43	2.5E-08	3.00	3.2E-12

Gene	Counts	Male FC	Male p_FDR	Female FC	Female p_FDR
Cc2d1a	2014	-1.38	3.6E-03	-1.44	1.1E-03
Ccdc91	1888	-1.95	2.9E-49	-1.82	4.0E-40
Ccng1	5463	1.49	2.9E-08	1.58	1.4E-10
Cd276	3140	-2.13	5.0E-36	-2.18	2.4E-38
Cd38	232	-1.84	8.7E-05	-1.90	3.5E-05
Cdc42bpb	14607	1.34	2.8E-04	1.31	9.7E-04
Cdh11	12820	-2.40	2.1E-06	-2.37	3.3E-06
Cdk19	6534	1.17	1.4E-02	1.25	3.0E-04
Cdk5rap2	8016	-1.37	6.2E-04	-1.27	1.2E-02
Cdkl5	3056	1.71	5.9E-08	1.95	9.5E-12
Cdon	13736	-1.37	4.4E-04	-1.20	4.7E-02
Celf6	1504	1.51	1.1E-03	1.44	4.8E-03
Cep135	2358	-1.51	7.3E-09	-1.38	9.7E-06
Cep41	1580	-1.91	2.3E-44	-1.94	1.8E-46
Chd2	12282	1.21	2.5E-02	1.35	3.5E-04
Chd3	39510	-1.98	9.7E-21	-2.00	3.8E-21
Chd7	13012	-1.35	1.4E-02	-1.33	2.0E-02
Chrna7	407.96	1.42	1.9E-04	1.28	9.1E-03
Clcn4	10609	-1.57	1.3E-06	-1.48	3.5E-05
Cln8	2506	1.77	2.3E-31	1.68	4.8E-26
Cmpk2	785	1.26	2.5E-02	1.26	2.4E-02
Cnr1	6596	-2.09	6.8E-05	-1.84	1.2E-03
Cntnap2	11429	1.42	7.8E-04	1.26	3.6E-02
Cpeb4	8073	1.60	9.9E-05	1.56	2.5E-04
Cpsf7	11294	-1.50	5.6E-09	-1.61	5.6E-12
Cpt2	1271	1.38	4.1E-04	1.29	5.1E-03
Crebbp	5496	-1.93	1.7E-10	-2.06	2.0E-12
Csde1	31901	-1.98	6.1E-40	-2.01	3.0E-41
Csnk1e	14294	-1.44	1.2E-06	-1.54	6.5E-09
Csnk2a1	16382	-1.27	5.8E-08	-1.26	8.2E-08
Csnk2b	5717	1.39	1.0E-11	1.32	1.4E-08
Ctcf	9148	-2.34	1.8E-38	-2.56	6.6E-47
Ctr9	13165	1.54	1.2E-27	1.54	1.4E-27
Cul7	10139	-1.69	4.5E-14	-1.80	1.2E-17
Cx3cr1	687	-2.05	1.2E-17	-2.28	1.3E-22
Cyfip1	6506	-1.45	1.4E-06	-1.45	1.6E-06
Ddc	823	-1.30	1.8E-02	-1.35	6.1E-03
Ddx3x	52188	-1.58	5.5E-09	-1.63	4.1E-10
Deaf1	3240	-1.47	4.7E-07	-1.57	2.5E-09
Depdc5	6414	1.60	1.5E-25	1.66	5.1E-29
Dhcr7	4286	1.56	6.0E-08	1.38	1.2E-04
Dhx30	9045	-1.55	3.6E-20	-1.65	7.1E-26
Dip2c	3203	-2.00	2.1E-11	-1.94	1.9E-10
Disc1	150	-2.17	8.3E-13	-1.71	1.1E-06
Dixdc1	6724	1.41	1.4E-06	1.50	1.0E-08
Dmpk	2879	1.32	4.3E-03	1.26	2.2E-02
Dnah10	231	2.35	4.2E-06	2.49	8.6E-07
Dnah17	157	5.09	4.6E-23	5.86	7.3E-27
Dnmt3a	15809	-2.22	4.4E-38	-2.11	3.4E-33
Dock1	4066	-1.49	7.5E-04	-1.38	7.5E-03
Dock8	232	-2.26	2.3E-04	-2.08	1.0E-03
Dolk	1049	1.29	1.9E-04	1.22	5.4E-03

Gene	Counts	Male FC	Male p_FDR	Female FC	Female p_FDR
Dpp4	163	-1.87	5.5E-03	-1.69	2.3E-02
Dpysl2	11973	-1.40	2.7E-02	-1.70	2.8E-04
Dpysl3	81721	-1.46	4.0E-03	-1.42	8.2E-03
Drd2	191	1.98	7.8E-06	1.92	2.1E-05
Dusp15	477	2.05	1.4E-05	2.24	9.4E-07
Dyrk1a	7472	-1.44	2.6E-08	-1.54	2.1E-11
Ebf3	12814	-1.68	2.6E-04	-1.60	1.1E-03
Ecpas	9645	-1.62	2.0E-10	-1.50	1.6E-07
Efr3a	5605	1.74	1.0E-15	1.91	5.7E-21
Ehmt1	11001	-1.32	3.2E-07	-1.44	5.5E-12
Ep300	10587	-1.39	1.2E-05	-1.31	4.0E-04
Epc2	5022	-3.53	7.0E-57	-3.45	8.2E-55
Ephb2	25546	1.44	1.4E-03	1.29	2.8E-02
Erbin	14418	-2.02	2.6E-26	-1.88	3.6E-21
Erg	1097	-2.37	4.0E-09	-2.44	1.2E-09
Exoc3	8530	-1.26	7.8E-06	-1.23	4.7E-05
Exoc6	1031	-1.67	5.4E-12	-1.60	2.1E-10
Ext1	5543	1.28	3.9E-04	1.29	2.6E-04
Fabp5	12171	-1.45	1.2E-07	-1.46	7.7E-08
Fam98c	976	2.59	4.5E-43	2.47	5.2E-39
Fbxo11	13248	-1.37	3.5E-04	-1.41	1.1E-04
Fbxo33	1584	1.59	1.5E-17	1.51	6.6E-14
Fezf2	1417	-1.73	5.8E-03	-1.99	4.1E-04
Foxp2	6899	-1.49	2.2E-08	-1.58	1.6E-10
Frk	338	-3.44	2.6E-06	-3.42	3.2E-06
Fxn	615	-1.96	1.2E-17	-2.04	1.4E-19
Gabbr2	570	-2.47	7.9E-03	-2.25	1.9E-02
Gabra4	739	2.46	3.0E-08	2.63	2.6E-09
Galnt10	4324	1.96	2.5E-16	1.91	4.3E-15
Galnt14	565	-2.75	7.3E-21	-2.53	1.2E-17
Gas2	1719	-1.55	2.7E-04	-1.45	2.1E-03
Gbe1	851	-1.62	4.7E-07	-1.67	7.3E-08
Gnai1	4191	1.42	4.4E-03	1.43	3.8E-03
Gnb1l	483	1.57	2.1E-15	1.60	1.3E-16
Grb10	45862	-1.51	5.4E-03	-1.49	6.7E-03
Grid2ip	169	2.70	6.3E-24	3.40	1.4E-34
Grik4	520	-1.44	1.2E-03	-1.38	4.7E-03
Grik5	7313	-1.31	2.2E-02	-1.40	3.6E-03
Grin2a	270	-2.28	1.5E-08	-2.27	2.0E-08
Gtf2i	23958	-2.36	1.2E-34	-2.54	8.0E-41
Hcfc1	17725	-2.30	2.7E-29	-2.29	4.5E-29
Hdlbp	25696	-1.29	9.1E-03	-1.25	2.5E-02
Hecw2	1165	-2.41	4.0E-22	-2.08	1.7E-15
Hivep2*	2898	-1.46	1.3E-02	-1.32	8.0E-02
Hnrnpk	69304	-1.30	9.0E-05	-1.36	3.4E-06
Hnrnpu	3018	-1.59	3.3E-04	-1.59	3.4E-04
Homer1	5300	1.61	9.6E-20	1.63	7.6E-21
Ica1	848	-1.41	2.3E-04	-1.30	5.0E-03
Igf1	2781	-3.30	8.9E-04	-3.18	1.3E-03
Ikzf1	271	-1.58	1.5E-05	-1.59	1.2E-05
Inpp1	1129	-2.01	6.8E-32	-2.01	8.4E-32
Ints6	3398	-1.53	3.8E-20	-1.47	2.7E-16

Gene	Counts	Male FC	Male p_FDR	Female FC	Female p_FDR
Iqgap3	4143	1.40	1.9E-03	1.39	2.5E-03
Iqsec2	1627	1.47	8.2E-05	1.40	7.0E-04
Irf2bpl	5211	-1.30	2.7E-02	-1.52	3.4E-04
Itpr1	4428	1.77	6.2E-07	2.01	6.7E-10
Kansl1	8932	-1.43	1.0E-14	-1.47	3.6E-17
Kansl1l	2089	-1.80	1.7E-19	-1.61	4.1E-13
Kat2b	1923	-1.71	1.0E-09	-1.73	4.2E-10
Kat6b	7883	-2.68	1.9E-23	-2.43	3.8E-19
Katnal1	2730	-1.97	2.7E-24	-2.04	9.5E-27
Kcnb1	1903	1.44	4.3E-03	1.49	1.6E-03
Kcnc3	5043	5.14	3.4E-42	4.74	3.4E-38
Kcnj10	1277	2.72	9.7E-28	2.66	1.8E-26
Kcnq2*	10691	-1.57	1.6E-02	-1.42	6.4E-02
Kdm1b	5673	2.26	3.4E-130	2.28	4.1E-133
Kdm5c	12138	-1.40	1.1E-14	-1.53	3.7E-23
Kdm6b	8176	-1.30	4.8E-02	-1.51	1.4E-03
Kif13b	3411	1.29	4.0E-03	1.48	4.3E-06
Kif14	2071	-1.46	4.8E-04	-1.48	2.9E-04
Klf16	1425	2.00	1.5E-16	1.78	8.3E-12
Klf7	6968	-1.26	2.5E-02	-1.27	2.3E-02
Kmt2e	18204	-1.33	3.4E-04	-1.36	8.9E-05
Kmt5b	6543	-6.45	2.2E-62	-6.59	9.4E-64
Kptn	941	-2.00	4.4E-45	-2.16	2.4E-55
Lamb1	8515	-1.45	4.2E-03	-1.39	1.3E-02
Las1l	6968	1.40	4.1E-24	1.37	1.2E-21
Ldb1	15935	-1.30	6.7E-05	-1.40	3.0E-07
Ldlr	2768	-1.36	3.2E-02	-1.57	1.1E-03
Lin7b	298	6.34	4.5E-69	6.45	1.5E-69
Lnpk	5745	1.28	3.6E-03	1.46	3.3E-06
Lrfn2	233	-1.63	3.9E-02	-1.67	3.1E-02
Lrp2	2856	1.43	3.2E-02	1.64	2.8E-03
Lrrc1	2446	1.64	7.0E-12	1.59	9.9E-11
Lrrc4	888	-3.27	1.1E-12	-3.64	8.2E-15
Lztr1	7298	-1.32	3.1E-15	-1.33	6.8E-16
Lzts2	3586	-1.20	3.2E-02	-1.33	5.7E-04
Maoa	3840	1.36	9.8E-05	1.39	2.0E-05
Map1a	10875	1.62	2.0E-04	1.61	2.3E-04
Map4k1	311	-1.27	6.1E-03	-1.37	3.2E-04
Mapk8ip1	12031	1.45	1.9E-07	1.37	1.0E-05
Mark1	5757	-1.39	3.0E-03	-1.42	1.7E-03
Mast3	433	2.89	1.1E-05	2.18	1.5E-03
Mbd3	6414	-2.01	1.9E-13	-2.27	3.7E-18
Mbd4	2576	1.30	1.3E-05	1.33	1.8E-06
Mboat7	5265	1.93	3.0E-26	1.91	2.5E-25
Mecp2	11258	-1.59	2.3E-05	-1.54	7.6E-05
Megf10	3621	-1.96	6.1E-22	-1.80	9.4E-17
Meis2	15231	-1.49	9.6E-04	-1.39	7.0E-03
Mink1	6672	1.33	6.4E-04	1.25	9.3E-03
Mrtfb	3074	-1.45	6.7E-10	-1.39	5.4E-08
Mtf1	1803	-1.16	2.5E-02	-1.25	5.6E-04
Mtor	10534	1.48	7.4E-11	1.56	1.1E-13
Myh9	16822	1.30	3.5E-02	1.29	4.4E-02

Table 1 (cont)

Gene	Counts	Male FC	Male p_FDR	Female FC	Female p_FDR	Gene	Counts	Male FC	Male p_FDR	Female FC	Female p_FDR
Mylk	1785	-1.20	4.4E-02	-1.25	1.5E-02	Ppp1r9b*	13672	1.28	5.3E-03	1.17	9.4E-02
Myo1e	3165	3.35	1.4E-64	3.28	2.9E-62	Ppp2r1b	7788	1.54	2.2E-11	1.44	2.0E-08
Myo5a*	17678	-1.36	7.3E-03	-1.22	9.8E-02	Ppp2r5d	10594	1.79	5.1E-25	1.68	2.5E-20
Myo9b	12403	1.43	3.5E-08	1.52	5.1E-11	Prex1	6987	1.55	5.5E-08	1.62	2.4E-09
Nacc1	8224	-1.15	1.4E-01	-1.29	3.2E-03	Prkd1	1487	1.33	9.2E-03	1.33	8.9E-03
Nexmif	3129	-2.12	1.7E-03	-1.76	2.3E-02	Prr12*	7437	-1.31	6.4E-02	-1.48	4.9E-03
Nf1	8492	-1.34	1.2E-04	-1.27	1.9E-03	Ptbp2	26867	-1.53	5.5E-08	-1.56	1.3E-08
Nfe2l3	803	4.19	8.5E-61	3.97	1.5E-56	Ptchd1*	234	-1.83	1.6E-02	-1.65	5.1E-02
Nfia	6750	-1.71	1.4E-03	-1.83	3.0E-04	Ptdss1	5032	1.32	6.1E-07	1.38	8.6E-09
Nfib	18590	-2.04	5.5E-04	-2.04	5.8E-04	Ptpn4	9452	1.32	1.6E-02	1.60	3.4E-05
Nfix	5859	-2.03	4.0E-03	-2.11	2.3E-03	Prkd2	1619	1.69	3.1E-11	1.65	3.4E-10
Nipa2	4612	1.32	9.0E-08	1.36	3.8E-09	Pten	12976	1.37	3.9E-06	1.37	3.9E-06
Nkx2-2†	355	-1.14	6.4E-01	-1.77	1.6E-02	Ptk7	9855	1.29	1.8E-02	1.27	2.5E-02
Nr1d1	234	2.42	2.9E-08	2.09	4.4E-06	Ptpn11	16665	1.31	9.8E-16	1.27	8.7E-13
Nr2f1	15518	-1.63	9.4E-06	-1.93	1.2E-09	Ptprb	1124	-1.77	2.0E-08	-1.83	2.6E-09
Nr3c2	254	2.04	7.0E-04	2.10	4.3E-04	Pxdn	15773	-1.48	4.5E-05	-1.38	1.1E-03
Nr4a2*	2860	1.35	1.7E-02	1.26	7.4E-02	Rab11fip5	3422	2.08	6.9E-20	2.09	3.7E-20
Nsd1	20486	-1.74	2.9E-11	-1.58	4.4E-08	Rad21	24544	-1.30	9.2E-06	-1.37	7.5E-08
Nsd2	33101	-1.33	1.0E-04	-1.28	9.4E-04	Rai1	7722	-1.57	2.5E-07	-1.56	3.3E-07
Nsmce3	1585	1.44	1.8E-18	1.36	2.3E-13	Ranbp17	928	-1.59	3.6E-07	-1.48	1.9E-05
Ntrk2	1174	-1.42	1.6E-02	-1.44	1.2E-02	Rassf5	168	1.54	2.2E-02	1.66	6.4E-03
Nxf1	17752	1.27	4.8E-08	1.26	1.6E-07	Reep3	6903	-1.26	5.0E-10	-1.26	5.5E-10
Nxph1	911	-2.60	8.0E-07	-2.48	3.3E-06	Rfx3	5786	-1.47	5.0E-08	-1.45	2.2E-07
Ofd1	2004	-1.35	8.8E-09	-1.34	2.4E-08	Rfx4	2512	-1.73	7.1E-04	-1.78	3.4E-04
Ophn1*	3837	-1.28	1.2E-02	-1.18	1.1E-01	Rims2	3497	-1.38	1.1E-02	-1.48	1.7E-03
Patj	1326	-2.29	1.3E-31	-2.20	1.1E-28	Rims3	4156	1.95	2.0E-08	1.98	1.2E-08
Pcca	2377	1.50	4.9E-07	1.51	4.8E-07	Rnf135	168	-1.99	4.9E-08	-1.97	8.1E-08
Pccb	2484	-1.67	1.1E-47	-1.81	4.0E-63	Rorb*	880	1.72	3.9E-04	1.32	8.8E-02
Pcdh10	8761	1.47	4.4E-03	1.54	1.4E-03	Rps6ka2	5307	1.80	8.1E-20	1.80	8.6E-20
Pcdha11	113	-2.30	2.5E-04	-1.94	4.1E-03	Rps6ka3	11427	1.21	3.3E-03	1.29	3.3E-05
Pcdha3	113	-2.00	1.2E-03	-1.97	1.7E-03	Runx1t1	7184	-1.62	4.1E-03	-1.61	4.8E-03
Pcdha7	112	-1.90	1.3E-02	-2.01	6.8E-03	Samd11	170	-1.90	1.4E-04	-2.03	2.3E-05
Pdk2	1859	2.00	1.8E-19	2.02	3.9E-20	Sbf1	12325	1.57	4.1E-08	1.53	2.8E-07
Pdzd8	8176	1.44	5.7E-22	1.50	1.2E-26	Scaf4	5878	1.30	4.1E-04	1.26	1.8E-03
Per1	3226	1.95	1.1E-11	1.77	8.2E-09	Scn9a*	2003	-2.28	2.2E-02	-1.29	5.3E-01
Per2	1092	1.71	1.0E-05	1.68	2.2E-05	Sdc2	2896	-1.53	2.6E-02	-1.49	3.8E-02
Pex7	836	-2.40	3.4E-59	-2.40	1.2E-58	Sema5a*	8354	-1.30	1.9E-02	-1.16	2.1E-01
Phb	6881	1.29	2.3E-05	1.27	1.1E-04	Set	42386	-1.41	4.4E-04	-1.59	1.3E-06
Phf2	6735	-1.85	5.0E-08	-2.00	5.6E-10	Setbp1	9254	-3.28	1.7E-29	-3.22	1.2E-28
Phf14	11218	-1.74	2.7E-18	-1.77	2.0E-19	Setd1b	4177	-2.04	1.6E-07	-2.28	1.1E-09
Phip*	21455	-1.44	1.2E-03	-1.22	9.0E-02	Setdb2	1135	1.56	1.1E-07	1.63	5.2E-09
Pik3cg	194	-2.20	4.0E-09	-2.41	6.5E-11	Sgsm3	2524	1.76	8.5E-29	1.69	8.3E-25
Pitx1	157	-3.67	5.1E-03	-5.34	2.6E-04	Shank2	2071	-2.21	1.8E-05	-2.32	4.8E-06
Plxnb1	17385	-1.79	1.2E-08	-1.91	2.6E-10	Sik1	2688	1.25	1.0E-02	1.26	8.0E-03
Pnpla7	498	-1.27	2.0E-03	-1.28	1.4E-03	Sin3a	12950	1.69	3.1E-27	1.62	8.5E-23
Pogz	8242	-2.18	2.3E-29	-2.25	8.9E-32	Slc22a15	1103	-2.14	3.7E-39	-1.90	6.7E-28
Polr2a	20372	-1.35	4.3E-04	-1.38	1.2E-04	Slc25a12	5108	1.38	2.3E-10	1.35	6.1E-09
Pola2	4393	1.67	7.1E-08	1.57	3.1E-06	Slc25a39	4509	1.70	1.8E-14	1.57	9.8E-11
Pomgnt1	3228	-1.63	1.2E-31	-1.69	7.8E-37	Slc29a4	3649	1.55	1.5E-05	1.54	2.4E-05
Pomt1	2351	-1.30	2.8E-07	-1.29	7.1E-07	Slc38a10	11004	1.62	9.3E-17	1.52	6.0E-13
Ppm1d	2918	1.33	2.6E-08	1.23	7.7E-05	Slc6a8	5359	1.72	5.6E-29	1.62	4.3E-23
Ppp1r1b	299	3.35	1.3E-04	4.05	8.5E-06	Slc7a3	316	2.75	5.8E-38	2.79	4.9E-39

Gene	Counts	Male FC	Male p_{FDR}	Female FC	Female p_{FDR}
Slc7a5	5758	1.57	2.8E-06	1.36	2.2E-03
Slc7a7	293	-1.48	5.4E-05	-1.62	4.5E-07
Slc9a6	7768	1.31	7.2E-03	1.39	1.1E-03
Slitrk5	2183	-1.56	6.1E-09	-1.59	1.4E-09
Smarca2	10932	1.25	3.1E-02	1.28	2.0E-02
Smarca4	29774	-1.59	5.2E-06	-1.69	2.5E-07
Smarcc2	27817	-1.55	1.9E-08	-1.63	4.3E-10
Smc1a	18407	-1.37	2.0E-06	-1.37	1.7E-06
Snap25	6289	1.64	8.5E-05	1.69	3.2E-05
Snd1	10450	1.27	3.4E-04	1.22	3.4E-03
Snx14	3275	1.73	5.1E-49	1.81	1.9E-56
Son*	59465	-1.32	3.8E-03	-1.14	2.2E-01
Sox6	5597	-1.29	2.3E-02	-1.32	1.4E-02
Sparcl1	6073	-1.59	2.4E-04	-1.48	2.5E-03
Srcap	371	-3.68	8.6E-20	-3.26	1.9E-16
Ssrp1	22251	-1.31	7.7E-05	-1.43	8.9E-08
St7	2111	-1.57	5.2E-08	-1.50	1.5E-06
St8sia2	9960	-1.84	8.8E-05	-1.91	3.1E-05
Stag1	8175	-2.42	1.3E-51	-2.34	7.8E-48
Stx1a	2912	1.38	5.2E-03	1.36	7.0E-03
Stxbp5	4749	1.46	3.0E-06	1.50	4.8E-07
Syap1	4114	1.57	3.4E-09	1.54	1.7E-08
Sybu	2165	-1.37	4.5E-02	-1.44	1.7E-02
Syn2	1795	1.63	3.8E-02	1.78	1.3E-02
Syncrip	22048	-1.48	7.5E-10	-1.51	7.8E-11
Synj1	9528	1.46	1.1E-03	1.54	1.8E-04
Taf1	11862	1.26	3.9E-04	1.36	1.7E-06
Taf4	3798	1.37	4.0E-04	1.28	7.5E-03
Taf6	5567	1.28	8.3E-05	1.18	1.1E-02
Taok1	12512	-1.49	4.8E-06	-1.36	4.8E-04
Tbc1d31	3810	2.35	1.3E-34	2.40	3.0E-36
Tbc1d5	1509	-1.60	6.8E-26	-1.59	4.1E-25
Tbl1x	11400	-1.64	2.6E-32	-1.68	4.3E-35
Tbl1xr1	4323	-1.42	2.5E-07	-1.57	2.4E-11
Tbr1	3672	-2.68	2.6E-05	-2.85	7.8E-06
Tbx22	155	-4.84	3.4E-02	-8.10	4.5E-03
Tceal1	996	-1.82	6.2E-13	-1.66	1.7E-09
Tcf4	24203	-1.55	9.7E-07	-1.48	1.4E-05
Tcf7l2	10030	-1.73	1.7E-02	-2.47	5.4E-05
Tet2	6298	-1.41	3.8E-05	-1.30	2.0E-03
Tet3	11236	-1.39	1.7E-03	-1.50	8.5E-05
Thbs1	7121	-1.91	1.8E-02	-1.90	1.9E-02
Tlk2	6144	-1.42	1.2E-14	-1.44	8.3E-16
Tm9sf4	4866	-1.34	2.6E-21	-1.36	1.1E-23

Gene	Counts	Male FC	Male p_{FDR}	Female FC	Female p_{FDR}
Tmem39b	1217	-2.03	9.6E-28	-2.26	2.6E-36
Tnrc6b	8717	-1.41	1.8E-06	-1.32	1.3E-04
Tns2	1379	-3.34	4.5E-19	-3.35	3.7E-19
Top3b	3624	-1.49	1.6E-43	-1.49	2.3E-43
Traf7	9194	1.42	3.8E-11	1.32	1.8E-07
Trappc6b	2765	-1.19	1.6E-02	-1.27	6.3E-04
Trim33	11213	-1.29	6.9E-05	-1.24	9.5E-04
Trio*	14535	-1.30	1.8E-02	-1.19	1.4E-01
Trrap	17379	-1.49	2.6E-06	-1.36	3.2E-04
Tsc2	11894	1.25	9.6E-04	1.25	8.7E-04
Tshz3	1699	1.28	4.1E-02	1.43	2.1E-03
Tspan17	1100	2.74	6.1E-25	2.80	7.2E-26
Tspan4	996	1.40	1.0E-03	1.36	2.8E-03
Ttn	914	-15.26	1.8E-03	-8.61	1.6E-02
Tubgcp5	3696	-1.29	2.7E-04	-1.21	8.0E-03
Ube2h	11325	1.37	7.8E-08	1.29	1.7E-05
Ubr1	5748	-1.47	1.3E-06	-1.37	9.7E-05
Ubr5*	28114	-1.28	3.3E-03	-1.18	5.5E-02
Unc13a	10790	2.41	2.4E-07	2.38	4.1E-07
Unc79	2448	-2.25	1.4E-06	-2.15	5.9E-06
Upf2	7646	1.31	1.6E-13	1.39	5.9E-19
Usp15	8350	1.26	6.5E-04	1.37	1.2E-06
Usp2	685	3.08	1.5E-34	2.78	1.2E-28
Usp30	2549	1.25	4.6E-09	1.15	3.5E-04
Usp45	4065	1.41	1.6E-14	1.43	1.5E-15
Vdr	170	6.19	1.5E-10	6.71	2.3E-11
Vezf1*	398	-1.31	4.8E-02	-1.11	4.8E-01
Vwa7	164	2.50	6.6E-09	2.60	1.4E-09
Wac	11365	-1.34	1.4E-10	-1.38	6.3E-13
Wdfy3	12067	-2.06	5.2E-11	-1.89	8.4E-09
Wnt1	77	-2.22	1.4E-02	-2.23	1.4E-02
Xpc	3233	1.40	2.2E-17	1.41	1.2E-17
Xrcc6	3705	1.36	7.6E-09	1.32	2.1E-07
Yeats2	8743	-1.43	3.0E-18	-1.38	3.4E-15
Ywhae	73412	-1.26	1.0E-04	-1.36	1.2E-07
Ywhaz	43502	-1.35	1.8E-10	-1.44	9.7E-15
Zbtb16	881	1.86	2.2E-04	1.70	1.7E-03
Zbtb20	6592	-2.52	2.0E-14	-2.13	5.5E-10
Zc3h4	6993	-2.62	3.8E-24	-2.79	3.1E-27
Zfyve26	2417	-1.46	1.9E-09	-1.33	1.0E-05
Zmiz1	19376	-1.45	1.1E-03	-1.53	1.8E-04
Zmym2	16595	-1.41	2.1E-06	-1.28	6.7E-04
Zmynd11	13113	-1.87	7.8E-23	-1.83	1.8E-21
Zmynd8	7708	-1.94	1.1E-19	-1.96	3.1E-20

* Significant effect of VPA in one sex only

Table 1. 400 genes that have been reported to be linked to autism by GWAS (SFARI list) that are also significantly up- or downregulated by VPA (from Table S2) in the fetal mouse brain.

Table 2

Gene	Counts	Male FC	p_FDR	Female FC	p_FDR
Adra2a	747	-1.66	2.3E-09	-2.01	1.1E-16
Aif1*	113	-1.49	2.7E-04	-1.15	2.4E-01
Ak2	7116	1.41	2.2E-04	1.27	1.2E-02
Akap8	17422	-1.26	3.2E-05	-1.29	6.1E-06
Akap8l	8525	1.60	5.9E-12	1.61	2.6E-12
Akt1	15951	-1.40	5.5E-09	-1.45	4.7E-11
Akt2	4906	-2.03	2.9E-50	-2.08	5.6E-54
Amotl2	6480	-1.60	8.2E-11	-1.72	4.1E-14
Apbb1ip	142	-1.73	1.1E-04	-1.57	1.7E-03
Apc2	25582	-1.41	3.9E-02	-1.48	1.7E-02
Arc	170	1.64	1.7E-04	1.66	1.4E-04
Arntl	1203	2.17	8.8E-45	2.07	1.6E-39
Asxl1	9564	-1.39	7.4E-06	-1.42	1.8E-06
Bcl10	2338	1.68	2.6E-19	1.59	1.4E-15
Bcl11b	7112	-1.58	1.7E-02	-1.67	8.1E-03
Bdnf	423	2.57	9.7E-12	3.10	3.3E-16
Bmp2	291	-1.49	3.6E-02	-1.68	6.3E-03
Bola2	2477	1.59	2.6E-05	1.48	5.2E-04
Brca1	5043	1.43	4.0E-03	1.48	1.7E-03
Brd1	7073	-1.39	1.9E-04	-1.47	1.5E-05
Brd2	19021	-1.26	1.3E-04	-1.35	6.2E-07
Brwd1	13743	-1.37	5.2E-06	-1.29	2.3E-04
C1qtnf1	444	-1.45	2.7E-04	-1.43	4.5E-04
C2cd5	6100	-1.54	1.9E-20	-1.50	3.3E-18
Cacnb4*	2101	-1.68	3.9E-03	-1.37	9.8E-02
Calm1	52234	1.44	1.0E-23	1.44	2.4E-24
Camk1	1971	-1.61	5.9E-24	-1.60	5.7E-23
Camkk1	2077	2.66	7.2E-41	2.53	6.9E-37
Camkv	1980	2.50	3.4E-28	2.30	1.1E-23
Carm1	6783	-1.40	1.9E-06	-1.52	1.4E-09
Casq2	132	-14.04	5.9E-04	-7.48	1.1E-02
Cbln2	2008	-1.82	1.5E-05	-1.71	1.2E-04
Cbs	736	5.02	2.1E-38	5.08	7.7E-39
Cdh7*	13012	-1.35	8.4E-03	-1.24	6.8E-02
Cdh15*	136	-4.45	6.9E-03	-2.61	9.7E-02
Clic6	1509	-1.80	3.4E-04	-1.58	6.3E-03
Col6a6	212	-4.15	3.8E-04	-4.13	4.2E-04
Creb3	2520	1.42	4.5E-15	1.43	1.2E-15
Cry1	3590	2.57	1.2E-98	2.41	5.4E-86
Cry2	3881	2.01	2.8E-27	1.94	1.1E-24
Csf1r	1501	-1.39	8.8E-04	-1.46	1.2E-04
Csgalnact1	179	-4.40	2.5E-08	-3.40	5.4E-06
Cux1*	12750	-1.17	5.2E-02	-1.27	2.6E-03
Cxcl14	1869	-3.84	2.6E-04	-3.31	1.3E-03
Dab2	6440	-1.69	1.0E-02	-1.63	1.8E-02
Dcbld2	9176	1.74	2.6E-25	1.68	3.3E-22
Dcps	2121	-1.27	3.8E-03	-1.34	3.2E-04
Dhfr	4614	1.56	3.7E-04	1.38	1.1E-02
Dlg5	6522	-2.95	3.9E-61	-2.96	1.3E-61
Dlx1*	5903	-1.77	1.2E-01	-2.29	2.0E-02
Dnajc12	1458	4.40	5.0E-179	4.66	1.6E-192
Dop1b	4122	1.41	4.7E-03	1.47	1.5E-03

Gene	Counts	Male FC	p_FDR	Female FC	p_FDR
Draxin	7093	-1.46	1.0E-02	-1.48	8.0E-03
Dyrk1b	1249	-2.21	3.1E-12	-2.12	3.7E-11
Ebf1	6893	-1.54	6.9E-05	-1.34	9.0E-03
Efna3	1500	2.61	1.2E-19	2.86	3.1E-23
Egfr	1567	-1.89	1.1E-02	-1.69	3.9E-02
Egr1	400	1.74	2.8E-04	1.71	4.8E-04
Elavl4	13824	-1.58	1.1E-05	-1.45	4.3E-04
Emp2	457	-1.49	4.9E-02	-1.62	1.6E-02
Eomes	3938	-1.81	1.6E-02	-1.86	1.1E-02
Ets2	2942	-1.45	8.6E-03	-1.42	1.5E-02
Evx1*	102	-3.25	1.3E-02	-1.42	5.1E-01
Farp1	6684	-1.72	5.0E-11	-1.72	7.1E-11
Fbxl21	373	3.18	2.4E-56	3.23	2.4E-57
Fbxl3	4323	-1.77	5.5E-43	-1.70	2.2E-37
Fezf1*	950	-1.20	4.6E-01	-1.68	2.3E-02
Fgf2	295	-2.94	3.9E-15	-2.79	1.1E-13
Fgf17*	317	-1.64	1.2E-02	-1.45	6.85E-02
Fgfr1op2	6435	1.26	5.9E-05	1.31	3.4E-06
Flrt2	7693	-1.73	2.6E-08	-1.52	3.0E-05
Fos	108	1.92	1.0E-03	2.49	3.2E-06
Foxa1	549	-1.47	2.2E-02	-1.75	5.7E-04
Foxc1	1824	-1.79	6.9E-04	-2.13	7.9E-06
Foxo1	2595	1.84	2.8E-13	1.80	1.8E-12
Foxo3	4755	1.66	1.9E-11	1.70	2.6E-12
Foxo4	3776	1.37	1.8E-05	1.36	3.1E-05
Foxo6	990	1.56	1.2E-02	1.50	2.3E-02
Fzd5	608	2.35	1.0E-10	1.80	1.3E-05
Gabra1	591	-1.75	2.6E-03	-1.68	5.9E-03
Gad1	3462	-2.50	1.0E-04	-2.54	7.7E-05
Gad2	5532	-3.18	2.9E-08	-3.38	4.9E-09
Gas7	1662	-1.64	6.9E-03	-1.67	4.8E-03
Gbx1	311	-2.29	3.5E-03	-2.39	2.2E-03
Gch1	460	5.58	7.6E-65	5.25	1.8E-60
Gfpt2	1177	1.67	1.2E-07	1.78	2.5E-09
Gls2	319	4.06	2.3E-62	4.84	4.0E-75
Grb2	9573	1.44	5.9E-08	1.37	3.4E-06
Grm2	321	-1.39	2.8E-02	-1.49	7.1E-03
Grm3	394	-1.84	2.7E-03	-1.59	2.7E-02
Hapln4	275	27.66	3.9E-126	23.71	2.8E-119
Hes1	1561	-1.52	3.5E-06	-1.70	2.6E-09
Hnrnpab	47049	-1.53	2.7E-05	-1.75	1.5E-08
Hnrnph1	93936	-1.60	9.0E-14	-1.61	5.2E-14
Homer3	3874	-1.27	1.0E-02	-1.43	9.4E-05
Hrh3	823	2.58	5.1E-15	2.34	2.6E-12
Hsp1a	235	6.27	1.4E-31	5.59	6.7E-28
Hsp1b	412	5.93	3.1E-35	5.20	1.9E-30
Hsp2	996	6.34	1.0E-96	6.09	1.8E-92
Icam5	104	16.47	6.4E-39	16.73	3.0E-37
Isl1	3389	-1.54	1.4E-03	-1.45	6.9E-03
Itpka	175	17.28	1.1E-58	10.83	3.3E-44
JunB	206	2.19	5.8E-10	2.20	5.4E-10
Kbtbd2	4187	-1.24	1.2E-05	-1.31	2.0E-08

Table 2 (cont)

Gene	Counts	Male FC	Male pFDR	Female FC	Female pFDR	Gene	Counts	Male FC	Male pFDR	Female FC	Female pFDR
Kcnmb2*	329	-2.11	5.5E-04	-1.45	1.1E-01	Plk1	3447	-1.54	5.4E-03	-1.83	7.0E-05
Kif17	387	4.17	2.5E-27	4.83	1.1E-32	Pnisr	26071	-1.95	1.5E-24	-1.81	1.0E-19
Klf4	882	-1.56	1.3E-02	-1.59	9.9E-03	Ppm1j	282	7.24	1.4E-36	7.60	4.4E-38
Kras	16185	1.38	2.1E-07	1.32	6.2E-06	Ppp3cb	14492	-1.25	1.0E-02	-1.29	3.0E-03
Lamp5	587	-6.05	9.0E-04	-4.36	7.7E-03	Ppp3cc	516	-1.27	6.9E-03	-1.25	1.2E-02
Lhx8	610	-2.58	7.4E-03	-2.81	3.5E-03	Ptk2b	117	1.75	1.6E-03	1.68	3.7E-03
Lmx1a	1011	-1.57	1.5E-03	-1.81	1.9E-05	Ptpn1	7485	1.59	8.2E-10	1.45	1.3E-06
Lrp6*	10654	-1.26	1.0E-04	-1.14	3.4E-02	Ptpn5	4140	-1.59	1.2E-02	-1.49	3.7E-02
Lrrk1	709	-2.99	2.8E-13	-2.85	3.1E-12	Ptpro	3982	-1.70	1.0E-02	-1.67	1.4E-02
Lrrk2	647	-2.40	1.1E-08	-2.20	2.9E-07	Pttg1ip	4768	1.30	1.4E-04	1.28	3.9E-04
Maf	3212	-1.63	3.8E-06	-1.64	2.8E-06	Pura	2394	1.32	2.7E-05	1.35	5.1E-06
Mafb	999	-1.57	8.5E-05	-1.71	2.2E-06	Raf1	11657	-1.62	2.3E-24	-1.60	1.8E-23
Map2	68999	-1.59	2.1E-04	-1.41	7.8E-03	Rarb	1030	-2.09	2.6E-05	-2.27	3.0E-06
Map2k1	4241	1.70	2.3E-18	1.67	3.6E-17	Rcan1	3802	1.67	4.1E-27	1.59	4.6E-22
Mapk1	20963	1.33	3.2E-15	1.29	7.0E-13	Rcan1	3802	1.67	4.1E-27	1.59	4.6E-22
Mapk14	7022	1.55	4.6E-40	1.52	9.9E-36	Rmnd5b	2194	1.35	3.1E-10	1.32	7.3E-09
Marcks	64056	-2.10	3.1E-25	-2.26	2.7E-30	Rnd3	17191	-1.47	1.3E-07	-1.44	6.9E-07
Mcam	4295	1.79	4.6E-17	1.65	5.1E-13	Rnf19b	2770	1.38	7.2E-05	1.32	7.2E-04
Mef2d	6108	1.54	1.3E-05	1.47	1.1E-04	Robo3*	6624	-1.73	5.8E-03	-1.23	3.4E-01
Metrn	4092	2.39	5.0E-10	2.07	2.6E-07	Robo4	695	-2.11	1.5E-17	-2.21	1.1E-19
Mllt11	16031	-1.56	2.5E-03	-1.49	8.0E-03	Rtn4rl1	469	2.04	4.0E-09	2.01	7.6E-09
Ncoa3	5325	1.35	1.9E-06	1.38	3.7E-07	Runx1	381	-3.19	1.6E-04	-2.98	4.3E-04
Ndst3	2340	-1.64	9.5E-05	-1.47	3.0E-03	Runx2	782	-3.89	1.5E-05	-3.99	1.1E-05
Nell2*	9177	-1.22	5.5E-02	-1.29	1.1E-02	Rxrb	4358	-1.27	5.6E-06	-1.34	3.0E-08
Neurod1	2403	-1.43	1.4E-03	-1.58	3.5E-05	S1pr1	1963	-1.25	2.4E-06	-1.38	4.0E-12
Neurog1	475	-2.00	4.6E-03	-2.49	1.5E-04	Sema3a	2749	-2.24	2.5E-07	-1.77	3.5E-04
Neurog2*	6066	-1.37	1.4E-01	-1.66	1.3E-02	Setd4	686	-1.85	1.3E-19	-1.92	5.7E-22
Nfatc1	283	-1.52	3.3E-04	-1.63	2.2E-05	Sh3bgr	135	1.74	1.1E-02	1.97	1.9E-03
Nfatc2	689	-1.65	4.9E-03	-1.60	8.5E-03	Shc2	2467	-1.42	1.9E-05	-1.41	3.2E-05
Nfatc4	3812	-2.04	9.2E-09	-2.10	2.2E-09	Shc4	325	-1.69	1.4E-05	-1.50	1.0E-03
Nhej1	194	-2.15	1.1E-16	-2.40	3.4E-21	Shox2	3795	-1.95	6.0E-05	-2.26	8.6E-07
Nhlh1	2918	-1.63	8.1E-06	-1.62	1.1E-05	Sim2	224	-2.06	9.2E-03	-2.27	2.9E-03
Nhlh2	5362	-2.89	1.7E-16	-2.74	4.9E-15	Sirt1	3366	-1.49	5.2E-15	-1.48	1.1E-14
Nkx2-1*	2086	-1.74	1.2E-01	-2.46	8.6E-03	Slc17a6	5910	-2.41	2.9E-09	-2.42	2.4E-09
Nkx2-4*†	120	-1.30	2.9E-01	-2.42	8.7E-05	Slc17a7	847	1.63	1.3E-02	1.63	1.4E-02
Nprl3	1675	-1.37	3.3E-11	-1.29	1.2E-07	Slc17a8	404	-1.72	1.1E-03	-1.43	3.8E-02
Nras	15777	-1.36	4.5E-09	-1.42	1.2E-11	Slc17a9	221	5.87	1.2E-33	6.74	2.1E-38
Ntn4	1222	-1.88	2.1E-07	-2.09	1.1E-09	Slc35e1	5103	1.48	2.7E-16	1.42	3.2E-13
Odc1	24222	1.49	5.0E-06	1.43	5.3E-05	Slc8a1	4042	-2.30	2.5E-05	-1.79	4.2E-03
Ogt	39637	-1.51	1.5E-06	-1.30	2.8E-03	Slc8a2	3842	1.99	4.1E-15	2.04	4.2E-16
Olig1*	341	2.11	7.4E-05	1.49	4.5E-02	Slc9a3r1	2902	3.04	2.4E-31	2.94	1.9E-29
Otp	1318	-2.64	2.1E-08	-2.31	1.8E-06	Slit2	10361	1.31	2.6E-04	1.46	2.2E-07
P2rx7	210	-2.26	6.8E-06	-1.67	6.3E-03	Slitrk1	1872	-1.44	8.6E-03	-1.46	6.4E-03
Palm3	720	1.99	7.7E-06	1.89	4.2E-05	Slitrk4	1131	1.62	2.8E-05	1.72	2.3E-06
Pax7	1240	-1.74	3.4E-03	-1.68	7.0E-03	Smyd2	1864	-1.24	3.7E-04	-1.28	4.3E-05
Pdk1	4064	1.24	1.1E-02	1.33	6.5E-04	Sos1	9101	1.25	6.5E-03	1.29	2.0E-03
Per3	1173	1.85	8.7E-21	1.92	5.5E-23	Sox2*	10110	-1.21	3.3E-01	-1.50	2.4E-02
Pigh	1044	-1.39	5.5E-07	-1.38	8.7E-07	Sp1	8616	-1.26	7.3E-07	-1.20	7.9E-05
Pigp	2067	-1.62	3.5E-20	-1.60	1.9E-19	Sp2	1254	-1.49	2.2E-03	-1.63	1.5E-04
Pik3cb	2150	1.73	3.6E-22	1.77	1.2E-23	Sp3	9744	-1.51	3.1E-11	-1.49	7.0E-11
Pik3r4	3800	1.26	3.3E-05	1.24	1.7E-04	Sparc	25277	-1.37	2.6E-02	-1.40	1.8E-02
Plcg2	339	1.95	1.2E-07	1.74	1.6E-05	Src	5937	-2.49	4.8E-21	-2.70	7.6E-25

Table 2 (cont)

Gene	Counts	Male		Female	
		FC	p_FDR	FC	p_FDR
*Srebf1**	4265	1.38	5.8E-04	1.21	5.7E-02
Sst	803	-1.79	1.0E-02	-1.86	6.2E-03
Stat3	5850	2.72	1.2E-44	2.69	1.7E-43
Stc1	534	1.35	8.9E-03	1.35	9.0E-03
Stc2	494	2.83	2.1E-23	2.55	4.8E-19
Stx18	1175	-1.59	1.1E-09	-1.68	9.8E-12
Sumo3	6505	-1.39	4.8E-09	-1.51	3.1E-13
Tacc2	7124	-1.50	5.9E-06	-1.52	2.5E-06
Tctn1	4106	1.47	1.8E-08	1.41	6.7E-07
Tctn3	1997	1.94	6.3E-22	1.99	1.9E-23
Tfap2a	954	-1.85	2.8E-03	-1.71	1.1E-02
Tgfb2	2234	-2.52	4.1E-13	-2.34	2.9E-11
Tgfb3	1115	-2.03	1.7E-04	-1.96	4.0E-04
Th	476	-1.42	5.1E-03	-1.52	6.6E-04
Tlr3	167	-1.75	1.2E-10	-1.95	2.6E-14
Tmem108	1225	1.53	7.9E-06	1.37	1.4E-03
Tmem163	1496	-1.62	1.4E-05	-1.68	2.4E-06
Top3a	2357	1.71	1.3E-20	1.66	1.2E-18
Trib2	8489	-1.73	3.2E-07	-1.73	3.4E-07
Trnp1	298	1.45	3.8E-02	1.55	1.4E-02
Trpm4	1776	2.04	1.2E-15	2.11	5.0E-17
Tubb3	58234	-1.25	1.8E-02	-1.27	1.4E-02
Tug1	14599	-1.19	1.6E-04	-1.26	2.8E-07
Unc5b	3707	1.69	4.9E-08	1.41	4.4E-04
Unc5c	4631	-2.20	1.5E-06	-1.86	2.1E-04
*Usp34**	19191	-1.41	6.4E-03	-1.17	2.7E-01
Vegfc	373	-1.52	4.0E-06	-1.73	1.1E-09
Vegfd	111	-2.21	2.6E-04	-1.91	3.5E-03
Wdr4	3027	-1.43	2.4E-11	-1.47	9.8E-13
Wdr41	1513	-2.70	1.1E-116	-2.73	2.1E-118
Wnt10b	116	-3.73	2.4E-03	-3.29	6.8E-03
Wnt16	127	2.71	1.2E-03	3.13	2.0E-04
*Wnt3**	639	-1.19	1.3E-01	-1.37	5.3E-03
Wnt3a	207	-3.64	1.6E-12	-4.01	2.7E-14
*Wnt4**	470	-1.70	2.3E-02	-1.39	1.8E-01
Wnt7b	4194	-1.68	1.4E-06	-1.97	1.5E-10
Wsb1	31033	-1.42	6.3E-08	-1.35	2.8E-06
Zfp804a	553	-1.88	3.6E-03	-1.68	2.0E-02
Zmym4	12662	-1.53	7.8E-14	-1.45	5.7E-11

* Significant effect of VPA in one sex only

Table 2. 247 genes involved in brain development and neurodevelopmental disorders that are dysregulated in the fetal brain by VPA (from Table S2) that have **not** been reported to be as sociated with autism in GWAS.

No significant sexually-dimorphic effects of VPA. Fig. 1 is a plot of the $\log_2$ fold-change (FC) of males versus females for the 647 genes in Tables 1 and 2. The gap in the data reflects the ±1.25-fold FC cut-off. Deviations from the regression line (r^2=0.973) indicate possible sex differences; however, in no case was the effect of VPA found to be significantly different between males and females. 17 genes (magenta) were upregulated by > 5-fold or downregulated > 80% by VPA (average of males and females).

Gene synonyms/protein products:

Arc, Arg3.1

Arntl, Bmal1

Cacna1c, Cav1.2 Ca^{2+} channel

Cacnb4, Cav2.2 CaCh β-subunit

Calm1, calmodulin

Chrna7, α7 nAChR

Cnr1, CB1 eCB receptor

Dop1b, Dopey2

Eomes, Tbr2

Gad1, GAD67

Kcnc3, Kv3.3 K channel

Kcnmb2, BK Ch β-subunit

Map2k1, Mek

Pik3cb, PI3 kinase β subunit

Pik3cg, PI3 kinase γ subunit

Ppp1r1b, DARPP-32

Ptk2b, Pyk2

Ptpn5, STEP

Rcan1, Dscr1

Slc17a6/7/8, VGLUT2/1/3

Slc17a9, NaHX9

Slc7a5, large amino acid transporter

Slc8a1/2, NCX1/2

Syn2, synapsin II

DISCUSSION:

The overarching hypothesis for this research is that neurodevelopmental disorders involving cognitive function such as ASDs and intellectual disability (ID) are the result of abnormalities in the connections among multiple brain regions. Normal connectivity is established beginning during fetal brain development as the various brain regions become populated with specific classes of neurons that subsequently connect with other neurons to establish the complex neuronal networks that underlie cognition and behavior. Although these networks are refined during late pre- and postnatal brain development, it is likely that certain early prenatal developmental epochs are vulnerable to alterations in the neurodevelopmental "program" (Ben-Ari, 2008). Although it is by no means clear that the fetal rodent brain accurately recapitulates the autistic human brain during prenatal development, the VPA model enables experimental control of the timing of exposure to an environmental factor that causes abnormal behavior resembling autism. In the present paradigm, VPA is administered to the pregnant dam at E12.5, a time when multiple early neurodevelopmental events critical for brain organization and connectivity are occurring. These include proliferation of neural progenitors (NPs), determination of the neuronal fate of these NPs, migration of newborn neurons away from the proliferative ventricular zones, differentiation of NPs to establish a mature neuronal phenotype, extension and branching of axons and dendrites and establishment of synaptic connections.

More than 7,300 genes were significantly up- or downregulated in the E12.5 fetal brain 3 hr after exposure to VPA (Table S2). 400 of those genes have been linked to autism in GWAS (Table 1), although in many cases the mechanism of action is unknown. In addition, at least 247 of the genes dysregulated by VPA but not on the SFARI Autism Risk list (Table 2), are known to play a mechanistic role in mediating one or more of the neurodevelopmental processes discussed above. Therefore, the goal of this study was to identify genes that are:

a. significantly up- or down-regulated by VPA in the fetal mouse brain AND

b. known to be linked to autism in GWAS or to play a role in embryonic neurodevelopmental processes, perturbation of which has the potential to alter brain connectivity in the postnatal and adult brain.

The set of genes meeting these criteria would provide potential targets for future hypothesis-driven approaches to understanding the underlying proximal causes of defective brain connectivity in neurodevelopmental disorders such as autism.

It was not practical to vet each of the more than 7,300 VPA-dysregulated genes for a potential role in abnormal brain development. However, as described below, the 247 genes in Table 2 and some of the genes in Table 1 have been reported to play substantial roles in specific aspects of fetal brain development. Consequently, it is plausible that the up- or downregulation of one or more of those genes could contribute to the abnormal connectivity in subjects with neurodevelopmental disorders. The validity of this potential contribution could be tested in animals by determining the effect of manipulating the expression of individual (or combinations of) genes in the fetal brain on subsequent behavior.

A common subset of genes in GWAS and VPA-fetal mouse brain studies. Several GWAS have identified genes linked to autism (see SFARI Autism list). Here we tabulated the findings of four GWAS studies (Satterstrom et al., 2020; Yuen et al., 2017; Stessman et al., 2017; Ruzzo et al., 2019) and one report categorizing autism risk genes potentially involved in neurogenesis (Garcia-Forn et al., 2020). As shown in Table 3, many of the genes identified in these reports are also dysregulated by VPA in the fetal mouse brain (Tables 1 and 2). Genes discussed in the respective papers are listed in each column in Table 3; genes that are also dysregulated by VPA in the present study are shown in **bold**. Five VPA-regulated genes in Tables 1 and 2 that also appear in all five of the published reports are in **red**. (*Arid1b*, *Dyrk1a*, *Pogz*, *Pten*, *Tbr1*.) Six genes in Tables 1 and 2 that also appear in four of the five other studies are in **blue**. (*Adnp*, *Ash1l*, *Chd2*, *Kmt5b*, *Tcf7l2*, *Wac*). There were four genes (yellow) identified in all five previously published studies that were not affected by VPA in the fetal mouse brain. *Chd2*, which is listed in three of the four published studies, and *Chd3*, (50% reduced by VPA), are both SFARI genes and have been reported to have similar functions to *Chd8*. Of the 11 **red** and **blue** genes in Table 3, three (*Adnp*, *Dyrk1a*, *Pten*) appear in Table 4 as VPA-regulated, autism-associated genes involved with the structural stability of neurons (Lin et al., 2016).

VPA dysregulation of high-confidence (hc)ASD genes in mid-fetal layer 5/6 cortical projection neurons. (Willsey et al., 2013) identified 9 hcASD genes that are expressed in layer 5/6 projection neurons in the fetal human brain. Three of these genes (*Dyrk1a*, *Pogz*, *Tbr1*) are downregulated in fetal mouse brain following VPA administration to the pregnant dam (Table 1). Willsey et al. (2013) used the nine hcASD genes as seed genes for co-expression network analysis which revealed 10 probable ASD genes, of which two (*Bcl11a*, *Nfia*) were down-regulated and one (*Aph1a*) was up-regulated by VPA (Table 2). All of these genes are on the SFARI autism risk gene list (Table 1) and the three seed genes were identified in all 5 reports analyzed in this study (Table 3). These findings raise the possibility that dysregulation of one or more of these genes in developing layer 5/6 projection neurons in cortical plate of the fetal brain contributes to permanent connectivity defects underlying autistic-like behaviors.

Table 3

Satterstrom	Garcia-Forn	Yuen	Stessman	Ruzzo	
41%	50%	37%	41%	44%	
		ADCY3	ADCY5		
ADNP		ADNP	ADNP	ADNP	
		AFF2			
		AGAP2		AKAP9	
ANK2	ANK2	ANK2	ANK2	ANK2	*
ANKRD11		ANKRD11	ANKRD11		
AP2S1					
ARID1B	ARID1B	ARID1B	ARID1B	ARID1B	
ASH1L		ASH1L	ASH1L	ASH1L	
ASXL3		ASXL3	ASXL3		
BABRB3					
BCL11A	BCL11A			BCL11A	
BRIN2B				BTRC	
CACNA1E				C16orf13	
CELF4		CACNA2D3	CACNA2D3	CACNA2D3	
			CASK	CAPN12	
			CDC42BPB	CCSER1	
CHD2		CHD2	CHD2	CHD2	
CHD8	CHD8	CHD8	CHD8	CHD8	*
		CIC		CMPK2	
		CLASP1	COL4A3BP		
		CNOT3			
CREBBP					
CTNNB1	CTNNB1		CTNNB1	CTTNBP2	
		CUL3	CUL3	CUL3	
DEAF1	DDX3X	DIP2C	DDX3X	DDX3X	
DNMT3A		DNMT3A	DIP2A	DNMT3A	
DPYSL2			DLG4		
			DLGAP1		
DSCAM		DSCAM	DOCK8	DSCAM	
		DYNC1H1	DSCAM		
DYRK1A	DYRK1A	DYRK1A	DYRK1A	DYRK1A	
EIF3G	FMR1	FAM47A		ERBIN	
				ETFB	
				FAM98C	
FOXP1	FOXP1	FOXP1	FOXP1	FOXP1	*
FOXP2				GABRB3	
GFAP					
GIGYF1		GIGYF1	GIGYF2	GIGYF1	
GNAI1		GRIN2B	GRIN2B	GRIA1	
IRF2BPL		KDM6A	HIVEP3	GRIN2B	
KCNQ3			ILF2	ILF2	
KDM5B			ITPR1	INTS6	
KDM6B		KDM6B		KDM6B	
KIAA0232		KIAA2022	KATNAL2	KATNAL2	
		KMT2A	KMT2A	KDM5B	
KMT2C		KMT2C	KMT2C	KMT2C	
			KMT2E	KMT2E	
KMT5B		KMT5B	KMT5B	KMT5B	
LDB1			LAMC3	MFRP	
MAP1A		MECP2	MECP2	MLANA	
MBD5					
MED13L		MED13	MED13L		

Satterstrom	Garcia-Forn	Yuen	Stessman	Ruzzo	
MKX		MYO5A		MYO5A	
MYT1L		MYT1L	MYT1L	MYT1L	
NRXN1		NAA15	NAA15	NCKAP1	
NSD1		NLGN3	NCKAP1	NRXN1	
PAX5		NLGN4X	NRXN1	P2RX5	
PHF12		PAX5	PARD3B	PCM1	
PHF21A		PCDH11X	PAX5		
		PHF2	PHF2	PHF2	
		PHF3	PHIP		
				PLXNB1	
POGZ	POGZ	POGZ	POGZ	POGZ	
			PPM1D	PRKAR1B	
PPP2R5D			PPP2R5D		
PRR12					
PTEN	PTEN	PTEN	PTEN	PTEN	
RAI1	SCN2A	SCN2A	PTK7	RANBP17	
RFX3			PTPN11	RAPGEF4	
RORB			RELN	SCN2A	
SATB1			RIMS1		
			SCN2A		
SCN2A			SETBP1		
			SETD2		
SETD5			SETD5	SETD5	
SHANK2		SHANK2		SHANK2	
SHANK3	SHANK3	SHANK3		SHANK3	
SIN3A			SLC6A1	SLC6A1	
SKI		SLC6A1	SMC3	SMURF1	
SLC6A1					
SMARCC2		SMARCC2			
SPAST		SPAST	SPAST	SPAST	
SRPRA					
		SRSF11	SRCAP		
			SRGAP3		
STXBP1			STXBP1		
SYNGAP1	SYNGAP1	SYNGAP1	SYNGAP1	SYNGAP1	*
			TANC2		
TBL1XR1		TAF6	TBL1XR1		
TBR1	TBR1	TBR1	TBR1	TBR1	
TCF20					
TCF4	TCF4		TCF4		
TCF7L2		TCF7L2	TCF7L2	TCF7L2	
TLK2				TMEM39B	
TM9SF4		TNRC6B	TNRC6B	TNRC6B	
		UBN2	TRIO		
TRIP12			TRIP12	TRIP12	
VEZF1		UPF3B	UNC80	TSPAN4	
				UIMC1	
				USP45	
WAC		WAC	WAC	WAC	
ZYMND8		WDFY3	WDFY3	WDFY3	
			WDR45	ZNF559	
			ZC3H4		
			ZNF292		

*Not affected by VPA

Autism-associated genes and structural stability of neurons

Lin et al. (2016) tabulated genes linked to autism in GWAS that have also been reported to be involved in the structural stability of neurons.

These genes were further sorted into three categories, viz., "neurite outgrowth", "spine/synapse formation" and "synaptic plasticity". 61 genes were assigned to these three categories although some genes appeared in two or all three categories, resulting in 29 different genes among the three categories (Table 4). Shown in red **bold** are 11 of these genes (38%) that were significantly up- or downregulated by VPA (i.e., in Table 1). As discussed in Lin et al. (2016), dysregulation of one or more of these genes (which could be due to gene variants or fetal exposure to VPA) could alter the structure and function of synapses throughout brain development.

Examination of Tables 3 and 4 revealed that *Pten* and *Dyrk1a* are common to the Lin et al., (2016) study and to all five of the GWAS. It is suggested that these two genes may be good candidates for investigating the molecular basis for the proximal causes of autism.

Table 4

Neurites	Synapse	Plasticity
CTTNBP2	**ADNP**	**CDH(S)**
CDH(S)	**CDH(S)**	**CDKL5**
CDKL5	**CDKL5**	DLGAP2
CNTN	CNTN	FMR1
CNTNAP2	**CNTNAP2**	GRIK2
DYRK1A	CTTNBP2	**GRIK4**
ELMO1	DLGAP2	GRIN2B
FMR1	**DYRK1A**	**MECP2**
GRIN2B	ELMO1	NLGN
MECP2	FMR1	NRXN
MYO16	GRIK2	**PTEN**
NCAM2	**GRIK4**	SHANK3
NLGN	GRIN2B	**STXBP5**
NRXN	**MECP2**	SYNGAP1
NTRK2	NLGN	UBE3A
PCDH	NRXN	
PTEN	**NTRK2**	
SPRICKLE1	PCDH	
STXBP5	**PTEN**	
TSC1	SHANK3	
TSC2	SYNGAP1	
UBE3A	TSC1	
	TSC2	
	UBE3A	

Table 4. Genes linked to autism in GWAS that have also been reported to regulate "neurite outgrowth", "synapse/spine formation" or "synaptic plasticity" (Lin et al., 2016). Genes in red **bold** are also up- or down-regulated by VPA in the fetal brain (Table 1). Lin et al. (2016) did not distinguish among the multiple known isoforms of cadherins (*Cdh's*); *Cdh11* is downregulated 60% by VPA in the fetal brain (c.f., Table 1).

Biological roles of genes dysregulated by VPA. Table 5 (next page) shows the biological function of genes that were significantly dysregulated by VPA in the E12.5 fetal brain. One or more developmental errors caused by these changes could induce a "signature" of abnormal brain circuitry resulting in abnormal behavior in juveniles and adults. In this discussion, the change in gene expression induced by VPA at E12.5 (expressed as fold-change) is shown in parentheses (mean of males and females). Following Table 5, the rationale for inclusion of the genes in the 18 categories is provided together with references to the literature.

Table 5

Intracellular signaling	*Ak2* **(+1.3)**, *Akap8* (−1.3), *Akap8l* **(+1.6)**, *Akt1* (−1.4), *Akt2* (−2.1), *Col6a6* (−4.1), *Dyrk1a* (−1.5), *Dyrk1b* (−2.2), *Grb2* **(+1.4)**, *Hnrnpab* (−1.6), *Kras* **(+1.4)**, *Map2k1* **(+1.7)**, *Mapk1* **(+1.3)**, *mTor* **(+1.4)**, *Ndst3* (−1.6), *Pik3cb* **(+1.8)**, *Pik3cg* (−2.3), *Pik3r4* **(+1.3)**, *Plcg2* **(+1.8)**, *Pten* **(+1.4)**, *Raf1* (−1.6), *Rmnd5b* **(+1.3)**, *Rnf19b* **(+1.4)**, *Sos1* **(+1.3)**, *Stat3* **(+2.7)**, *Tsc2* (−1.3)
Calcium signaling	*Calm1* **(+1.4)**, `*Cacna1b* **(+1.4)**, *Cacna1c* (−1.4), *Cacna2d1* (−1.7), *Cacnb2* (−1.9), *Camk1* (−1.6), *Camk2a* **(+1.9)**, *Camkk1* **(+2.6)**, *Camkv* **(+2.4)**, *Ppp3cb* (−1.3), *PPP3cc* (−1.3)
Retinoic acid signaling	*Megf10* (−1.9), *Cntnap2,* **(+1.3)**, *Meis2* (−1.4), *Cbln2* (−1.8), *Rxrb* (−1.3), *Rarb* (−2.2)
Neurogenesis	*Cdkrrap2* (−1.3), *Ebf1* (−1.4), *Eomes (Tbr2)* (−1.8), *Fgf17* (−1.6), *Foxc1* (−1.9), *Lrrk1 (−2.9)*, *Neurog1* (−2.2), *Plk1* (−1.7), *Trnp1* **(+1.5)**
Neuronal fate specification	*Ebf3* (−1.6), *Neurod1* (−1.5), *Nkx2-2* (−1.8), *Nr2f1 (Coup-tf1),* (−1.8), *Olig1* **(+2.1)**, *Tbr1* (−2.8)
Axon growth/guidance	*Draxin* (−1.5), *Marcks* (−2.2), *Slitrk1* (−1.5), *Syn2* (+1.7)
Neuronal migration	*Apc2* (−1.7), *Mllt11* (−1.5), *Neurod1* (−1.5), *Nsd1* (−1.4), *Otp* (−2.5), *Ptk2b* (−1.7), *Robo4* (−2.2)
Synaptogenesis	*Cbln2* (−1.8), *Csde1* (−2.0), *Ddx3x* (−1.6), *Pogz* (−2.2), *Slc17a7* **(+1.6)**, *Slitrk5* (−1.6), *Slitrk4* **(+1.7)**, *Syncrip* (−1.4), *Top3b* (−1.5)
Dendrite development	*Cc2d1a* (−1.4), *Csde1* (−2.0), *Disc1* (−1.9), *Dlg5* (−3.0), *Gas7* (−1.7), *Mef2d* **(+1.5)**, *P2rx7* (−2.0), *Slitrk1* (−1.5), *Tmem163* (−1.7),
Postsynaptic cytoskeleton	*Homer1* **(+1.6)**, *Homer3* (−1.4), *Shank2* (−2.3)
Excitation-Inhibition	*Arid1b* (−2.1), *Camk2a* **(+1.9)**, *Cc2d1a* (−1.4), *Gad1* (−2.5), *Gad2* (−3.3), *Gabra1* (−1.7), *Gabra2* (−1.7), *Gabra4* **(+2.5)**, *Grin2a* (−2.3), *Hapln4* **(+25.7)**, *Kmt5b* (−6.5), *Maf* (−1.6), *Mafb* (−1.6), *Nf1* (−1.3), *Trpm4* **(+2.1)**
Dopaminergic system	*Adora2a* (−1.3), *Bdnf* **(+2.8)**, *C1qtnf1* (−1.4), *Csf1r* (−1.4), *Drd2* **(+2.0)**, *Emp2* (−1.6), *Foxa1* (−1.6), *Klf4* (−1.6), *Lmx1a* (−1.7), *Mcam* (+1.7), *Palm3* **(+1.9)**, *Ppp1r1b* **(+3.7)**, *Robo4* (−2.2), *Rtn4rl1* **(+2.0)**, *S1pr1* (−1.3), *Slc9a3r1* **(+3.0)**, *Smyd2* (−1.3), *Sparc* (−1.4), *Srebf1* **(+1.4)**, *Th* (−1.4), *Tmem108* **(+1.4)**, *Wnt7b* (−1.8),
Cholinergic system	*Chrna7* **(+1.4)**, *Gbx1* (−2.3), *Isl1* (−1.5), *Lhx8* (−2.7), *Nkx2-1* (−2.5*),
Endocannabinoid system	*Cnr1* (−2.0)
Autism and Down syndrome	*Brwd1* (−1.3), *Dyrk1a* (−1.5), *Cbs* **(+5.0)**, *Dop1b* **(+1.4)**, *Erg* (−2.4), *Ets2* (−1.4), *Pigp* (−1.6), *Setd4* (−1.9), *Shebgr* **(+1.8)**, *Sim2* (−2.2)
Circadian rhythms	*Arntl* **(+2.1)**, *Cry1* **(+2.5)**, *Cry 2* **(+2.0)**, *Fbxl21* **(+3.2)**, *Fbxl3* (−1.8), *Gabra4* **(+2.5)**, *Gad1* (−2.5), *Gabbr2* (−2.4), *Mef2d* **(+1.5)**, *Per1* **(+1.9)**, *Per2* **(+1.7)**
Neuroinflammation	*Dnajc12* **(+4.5)**, *Hsp1a* **(+5.9)**, *Hsp1b* **(+5.6)**, *Hsp2* **(+6.2)**, *Igf1* (−3.2), *Igf2* (−2.7), *Tgfb2* (−2.4), *Tgfb3* (−2.0), *Tgfbr2* (−2.5)
Epigenetic regulation	*Ash1l* (−1.5), *Brd1* (−1.4), *Brd2* (−1.3), *Ehmt1* (−1.4), *Kmt5b* (−6.5), *Mecp2* (−1.6), *Setd1b* (−2.2)

Table 5. The biological roles of genes that are dysregulated by VPA in the fetal brain and are associated with autism or prenatal neurodevelopmental processes (Tables 1 and 2). The number in parentheses shows the FC induced by VPA (mean of males and females); **red** text indicates that VPA increased gene expression. Note that some genes are included in more than one category.

Intracellular signaling. As summarized in Figure 2, the genes encoding 10 members of the canonical

Figure 2

signaling pathways downstream from receptor tyrosine kinase receptors are significantly up-regulated by VPA in the fetal brain (green); four are down-regulated (red). ***Pten* (+1.4)** (Phosphatase and tensin homolog), a negative regulator of the PI3K-AKT-mTor signaling pathway, has a particularly strong association with autism, appearing in all GWAS (Tables 1, 3) and associated with the structural stability of neurons (Table 4) (Lin et al., 2016).

Twelve additional genes associated with intracellular signaling are included in Table 5. Thus, abnormal signaling via these pathways is a potential mechanism by which VPA could interfere with the establishment of normal brain connectivity. Meta-analysis of GWAS and copy number variation studies of autism-related genes revealed that three signaling networks, regulating steroidogenesis, neurite outgrowth and excitatory synaptic function, were enriched (Poelmans et al., 2013). A-kinase anchoring proteins (Akaps) functionally integrate signaling cascades within and among these networks. ***Akap8* (−1.3)** and ***Akap8l* (+1.6)** was decreased and increased, respectively, in fetal mouse brains exposed to VPA. ***Dyrk1b* (−2.2)** (which encodes dual specificity tyrosine phosphorylation regulated kinase 1B) is an early mediator of double-stranded DNA break repair (Dong et al., 2020) and regulates hedgehog signaling through activation of the mTor/AKT pathway (Singh et al., 2017; Upadhyay et al., 2021; Singh and Lauth, 2017). *Dyrk1b* has been linked to metabolic syndrome and autism (Orenstein et al., 2022). The 55% reduction in *Dyrk1b* expression induced by VPA could contribute to autistic-like behavior by dysregulating hedgehog or mTor/AKT signaling. ***Ppp1r1b* (+3.7)**, encodes the dopamine- and cAMP-regulated neuronal phosphoprotein (DARPP-32). DARPP-32 amplifies and/or mediates many actions of cyclic AMP-dependent protein kinase at the plasma membrane and in the cytoplasm, with a broad spectrum of potential targets and functions (Yger and Girault, 2011).

Ca^{2+} signaling. Ca^{2+} is an intracellular messenger in a wide variety of cellular responses; calmodulin, encoded by ***Calm1* (+1.4)**, acts as the Ca^{2+}-sensor in many responses including Ca^{2+}-calmodulin dependent kinases [***Camk1* (−1.6)**, ***Camk2a* (+1.9)**, ***Camkk1* (+2.6)**, ***Camkv* (+2.4)**], calcineurin, [***Ppp3cb* (−1.3)** and ***PPP3cc* (−1.3)**]. Four genes encoding subunits of voltage-gated Ca^{2+} channels on the SFARI list (Liao and Li, 2020; Breitenkamp et al., 2014) (Table 1) were dysregulated in the fetal brain: ***Cacna1b* (+1.4)**, ***Cacna1c* (−1.4)**, ***Cacna2d1* (−1.7)** and ***Cacnb2* (−1.9)**. Like *Cacnb2*, ***Cacnb4* (−1.5)** is an auxiliary subunit, however, it is not linked to autism by GWAS (Table 1). *Cacna1c* is associated

with Timothy syndrome, an ASD characterized by ID as well as long QT syndrome, a frequently lethal defect in heart rhythm (Herold et al., 2023; Endres et al., 2020).

Retinoic acid signaling. Retinoic acid (RA) is a diffusible derivative of vitamin A that has been reported to regulate the fetal development of the prefrontal cortex and the dopaminergic system in the striatum; dysregulation of RA signaling in mice can lead to autistic-like behavior. The RA receptor **Rarb (−2.2)**, which encodes RARβ, is downregulated by more than 50% by VPA in the E12.5 mouse brain. In addition, four additional genes dysregulated by VPA **Megf10 (−1.9)**, **Cntnap2**, **(+1.3) Meis2 (−1.4)**, **Cbln2 (−1.8)]** are involved in RA signaling; three of these (*Megf10*, *Cntnap2*, *Meis2*) are on the SFARI list. RA-dependent developmental processes in the fetal brain include loss of GABAergic striatal projection neurons and interneurons (Chatzi et al., 2013), reduced expression of D2 dopamine receptors in the ventral striatum (Krezel et al., 1998) and reduced cortical neurogenesis and cortical plate thickness (Haushalter et al., 2017). Shibata et al. (2021) reported that genetic deletion of *Rarb* disrupts proper molecular patterning of prefrontal and motor areas, development of prefrontal cortex (PFC)-mediodorsal thalamus connectivity and development of intra-PFC dendritic spinogenesis. RA signaling occurs through the RA receptors, RXRG and RARB, and genetic deletions of **Rxrb (−1.3)** and *Rarb* demonstrated that these enzymes are involved in the proper molecular patterning of the prefrontal and motor cortices, the development of cortico-thalamic connectivity and cortical spinogenesis (Shibata et al., 2021). RA signaling may also regulate the development of the dopamine (DA) system (see below), (Krezel et al., 1998); deletion of *Rxrb* and *Rarb* resulted in locomotor defects related to dysfunction of the mesolimbic DA signaling pathway, as well as expression of D1 and D2 DA receptors in the ventral striatum. VPA exposure reduced expression of *Rarb* (55%) and *Rxrb* (23%) in the fetal mouse brain. Interestingly, VPA doubled the expression the D2 DA receptor, **Drd2 (−1.9)**. Thus, these findings suggest that VPA could reduce the strength of RA signaling in the fetal brain, particularly in the dopaminergic system, resulting in altered numbers and/or connectivity of cortical or striatal neurons.

Neurogenesis. Excitatory neurons in the mammalian cortex are generated from proliferating radial glia cells (RGs) which undergo mitosis at the ventricular surface to expand the pool of neuroblasts and consequently regulate the final number of neurons (Silbereis et al., 2016). Neurogenesis proceeds according to a "program" that is tightly-regulated by intercellular signals from a variety of brain cells as well as the meninges. Disturbances in this regulation, for example by environmental factors such as VPA or gene mutations that disrupt the expression of the genes encoding the relevant extracellular signals, have the potential to alter the configuration and neuronal population of the developing brain leading to permanent defects in connectivity that could lead to altered behavior.

As discussed above, RA plays a number of roles in the regulation of cortical neurogenesis (Haushalter et al., 2017; Shibata et al., 2021). The RA has been reported to be produced in the dorsal forebrain meninges and act at receptors on the end-feet of RGs (Siegenthaler et al., 2009). This group used a transgenic mouse with reduced expression of *Foxc1* **(−1.9)** leading to defects in forebrain meningeal formation and loss of neuron progenitor production due to failure of proliferating RGs to exit the cell cycle. This would be predicted to lead to a delay in the generation of postmitotic neurons and consequently, an expansion of the proliferating RG pool neuroprogenitor pool. *Neurog1* **(−2.2)** has been reported to play a role as a negative regulator of cortical neurogenesis (Han et al., 2018). *Eomes* **(*Tbr2*)** **(−1.8)** is also implicated in the regulation of cortical neurogenesis (Englund et al., 2005; Baala et al., 2007; Arnold et al., 2008; Elsen et al., 2013; Sessa et al., 2017). The conversion from symmetrical to asymmetrical mitoses associated with the generation of post-mitotic neurons is regulated by the *Plk1* **(−1.7)** − *Lrrk1* **(−2.9)** − *Cdk5rap2* **(−1.3)** cascade (Hanafusa et al., 2015).

Comparison of the mechanisms controlling neurogenesis in rodents and humans has revealed that gyrification of the human brain is driven by changes in the expression of regulatory genes and growth factors (Borrell and Götz, 2014) . Stahl et al. (2013) reported that, over the period E12 to E16, decreased expression of the DNA-associated protein, *Trnp1* **(+1.5)**, leads to radial expansion of the cortex and the appearance of gyri-like folding of the mouse brain (Stahl et al., 2013). In the present study, VPA induced a 50% increase in the expression of *Trnp1* in the E12.5 fetal mouse brain; this would be predicted to inhibit or delay the shift to radial growth of the cortex. Fibroblast growth factor 2 (FGF2) has also been reported to induce gyrification in the mouse brain, having an effect opposite to that of *Trnp1* (Rash et al., 2013). In the present study, VPA caused a 65% reduction on the expression of *Fgf2* **(−2.9)** (Table 2). This would also be predicted to bias cortical neurogenesis toward tangential and away from radial growth that would lead to gyrification. These findings together with the present results lead to a hypothesis that increased *Trnp1* and decreased *Fgf2* expression in the fetal brain at E12.5 contribute to VPA-induced, autism-like behavioral defects. This hypothesis could be tested by experimentally decreasing *Trnp1* and increasing *Fgf2* expression in the mouse fetal brain exposed to VPA; these maneuvers would be predicted to normalize behavior.

Neuronal fate specification. *Tbr1* **(−2.8)** encodes T-brain-1, a brain-specific T-box transcription factor that has been linked to autism in multiple GWAS (see Table 3). VPA reduced *Tbr1* expression by 63% in the fetal mouse brain (Table 1). *Tbr1* has been described as a "master regulator of ASDs" because it regulates the expression of multiple genes with links to autism (Chuang et al., 2015), thereby controlling brain wiring in ASDs (Huang and Hsueh, 2015; Notwell et al., 2016). *Tbr1* has been reported to regulate differentiation of the preplate and layer 6 of the fetal cortex (Hevner et al., 2001) as well as the connectivity of cortical layer 6 (Fazel Darbandi et al., 2018). Recently, different *Tbr1* mutations

found in individuals with ASD, intellectual disability and developmental delay had differing effects on cortical development (Co et al., 2022); one such mutation (K228E) caused significant upregulation of *Tbr1* but similar behavioral phenotypes to those of *Tbr1* KO mice (Yook et al., 2019). Thus, it appears that both positive and negative deviations from an optimal level of *Tbr1* expression are associated with an autism-like behavioral phenotype. Considering its links to autism in GWAS and its central role in regulating fetal brain development, VPA-induced changes in *Tbr1* expression in the fetal brain could contribute to the behavioral abnormalities observed in animals exposed to VPA *in utero*.

Nr2f1 (***Coup-tf1***) **(−1.8)** encodes an orphan nuclear receptor that has been implicated in a range of neurodevelopmental functions including coordinating cortical patterning, neurogenesis, and laminar fate (Armentano et al., 2007; Faedo et al., 2008; Tocco et al., 2021). *Nr2f1* also has been reported to control subtype and laminar identity of cortical interneurons derived from the medial ganglionic eminence (Hu et al., 2017) as well as regional dynamics of neuroprogenitors in the cortex, possibly being responsible for abnormal gyrification (Bertacchi et al., 2020). *Nr2f1*, together with *Nr1f2*, regulates cell migration in the basal forebrain (Tripodi et al., 2004) and is required for forebrain commissural projections (Armentano et al., 2006). As discussed below, Engeln et al. (2021) reported that *Nr2f1* is a common regulator for multiple genes involved in spine morphology of D1-MSNs in the striatum.

Neuronal differentiation and regional specification in the CNS are regulated by the *Ebf* family of genes including ***Ebf3*** **(−1.6)** (Garel et al., 1997). Variants of *EBF3* are associated with hypotonia, developmental delay, intellectual disability, and autism (Padhi et al., 2021; Tanaka et al., 2017). The *Olig* genes (*Olig1/2/3*), which encode members of the basic helix-loop-helix (bHLH) family of transcription factors, were originally identified as regulators of oligodendrocyte production in the CNS. More recently, the *Olig* genes have been found to regulate the specification and differentiation of neuronal phenotype during fetal brain development (Chakrabarti et al., 2010; Szu et al., 2021). ***Olig1*** **(+2.1, significant in females only)** expression was more than doubled in fetal brains exposed to VPA. The basic Helix-loop-Helix gene, ***Neurod1*** **(−1.5)**, regulates multiple steps in cortical development including neuronal fate specification, differentiation, and migration (Tutukova et al., 2021).

Axon growth/guidance. ***Marcks*** **(−2.2).** Myristoylated Alanine-Rich C-kinase Substrate (MARCKS) plays a wide variety of roles in brain development and function (El Amri et al., 2018; Stumpo et al., 1995). These may be due at least in part to the regulation of axon growth in which MARCKS mediates membrane targeting of plasmalemmal precursor vesicles during axon development (Xu et al., 2014). Although total deletion of *Marcks* in mice is embryonic lethal, the 55% reduction in *Marcks* expression induced by VPA may be less severe and cause subtle changes in brain connectivity that could lead to autistic-like behavior. ***Syn2*** **(+1.7)** (synapsin-2) has been reported to be required for normal axon

formation (Ferreira et al., 1998). ***Draxin (−1.5)*** (dorsal inhibitory axon guidance protein) is essential for the axon guidance underlying development of the corpus callosum and thalamocortical projections (Liu et al., 2018; Ahmed and Shinmyo, 2021; Morcom et al., 2021).

Neuronal migration. *Robo4* **(−2.2)** has been reported to regulate the radial migration of newborn neurons in the developing neocortex (Zheng et al., 2012; Gonda et al., 2020). *Robo4* expression was reduced by more than half in the fetal brain by VPA. Dysregulation of *Robo4* expression at E12.5 could affect the developing connectivity of the fetal brain leading to behavioral abnormalities in the adult. *Mllt11* **(−1.5)** regulates migration and neurite outgrowth of cortical projection neurons during development (Stanton-Turcotte et al., 2022). ***Ptk2b (−1.7)*** (*Pyk2*) is a non-receptor cell-adhesion kinase and scaffold protein which, if either overexpressed or knocked down, impairs cortical neuron migration by altering growth cone dynamics (Fan et al., 2018). *Ptk2b* is regulated by ***Ptpn5 (−1.5)*** (STEP), a tyrosine phosphatase (Chatterjee et al., 2020). None of these three genes has been linked to autism in GWAS but downregulation of *Ptk2b*, *Ptpn5* and/or *Mllt11a* by VPA has the potential to alter neuronal migration at a critical time when circuits are forming in the fetal brain. ***Otp (−2.5)*** encodes a homeodomain transcription factor that is associated with the development of the hypothalamus in vertebrates. *Otp* is necessary for the migration of diencephalic neurons to the amygdala (García-Moreno et al., 2010) and affects neuropeptide switching in oxytocin neurons (Wircer et al., 2017; Acampora et al., 1999; Ryu et al., 2007). A loss of function mutation in ***Apc2 (−1.7)*** causes Sotos syndrome which is characterized by ID and characteristic facial features; this is because *Apc2* is a downstream effector of ***Nsd1 (−1.4)*** in regulating the migration and laminar positioning of cortical neurons (Almuriekhi et al., 2015). Thus, VPA inhibits this critical developmental pathway at two steps.

Synaptogenesis. *Slc17a7* **(+1.6)**, which encodes the vesicular glutamate transporter VGLUT1, promotes the development of cortical presynaptic terminals (Berry et al., 2012). The VPA-induced increase in *Slc17a7* would be consistent with the abnormal acceleration or premature initiation of this process. ***Slitrk*s** are single-pass transmembrane proteins of the leucine-rich repeat (LRR) superfamily that have been proposed as candidate genes for neuropsychiatric disorders (Proenca et al., 2011). There are 6 members of this family in humans and mice, viz., *Slitrk1−6*. ***Slitrk1 (−1.5)*** has been shown to modulate neurite outgrowth (Aruga and Mikoshiba, 2003) and to promote the development of excitatory synapses (Beaubien et al., 2016), *Slitrk2* (not significantly affected by VPA) is on the SFARI list and has been linked to schizophrenia and X-linked neurodevelopmental disorders (El Chehadeh et al., 2022) and ***Slitrk5 (−1.6)*** deficiency impairs corticostriatal circuitry (Shmelkov et al., 2010) and synaptogenesis (Song et al., 2017). *Slitrk1* and *Slitrk5* expression in the fetal brain was reduced 33% and 37% by VPA; ***Slitrk4 (+1.7)*** expression was increased 67% by VPA (Table 2). ***Cbln2 (−1.8)***. The cerebellins (CBLNs) are secreted glycoproteins that link presynaptic neurexins with postsynaptic δ1 glutamate receptors

(GluDs) to form trans-synaptic adhesion complexes and promote the formation and stability of excitatory synapses (Seigneur et al., 2021). The formation of these complexes is deficient in the *Cbln2* knockout mouse (Tao et al., 2018) resulting in the destabilization of excitatory synapses. Consequently, the 45% reduction in *Cbln* expression induced by VPA in the fetal mouse brain would be predicted to compromise excitatory synaptogenesis. *Top3b* **(−1.5)** encodes an RNA topoisomerase that works with FMRP to promote synapse formation (Xu et al., 2013) and is a high-confidence gene on the SFARI list (Table 1). *Top* family members have been reported to facilitate transcription of long genes linked to autism (King et al., 2013). *Syncrip* **(−1.4)** (SFARI list) (also known as *Hnrnpg*) encodes the synaptotagmin-binding cytoplasmic RNA-interacting protein, which is a candidate gene for ASD and ID (Rauch et al., 2012; Lelieveld et al., 2016; Semino et al., 2021). Bannai et al. (2004) reported that *Syncrip* is a component of mRNA-containing granules in dendrites, possibly regulating local protein synthesis in developing dendritic spines.

Disruption of *POGZ* is associated with ID, ASDs (Stessman et al., 2016, 2017) and impaired cortical development (Matsumura et al., 2020). Markenscoff-Papadimitriou et al. (2021) reported that *Pogz* **(−2.2)** promotes chromatin accessibility and expression of clustered synaptic genes. The X-linked gene, *Ddx3x* **(−1.6),** encodes an RNA helicase that functions in corticogenesis and synaptogenesis (Lennox et al., 2020; (Hoye et al., 2022). Mutations in *DDX3X* account for approximately 2% of intellectual disability in females (DDX3X syndrome) (Boitnott et al., 2021; Weil et al., 2020). Knockdown of the RNA binding protein, *Csde1* **(−2.0)**, has been reported to abnormal dendritic spine morphology and synapse formation (Guo et al., 2019).

Hapln4 **(+25.7)**, which encodes an extracellular matrix protein that is a component of perineuronal nets (PNNs), is a selective regulator of the formation of GABAergic synapses between Purkinje and deep cerebellar nuclei neurons (Edamatsu et al., 2018). The cerebellum is one of many brain regions implicated in the etiology of autism (Mapelli et al., 2022). The massive, 25-fold increase in *Hapln4* induced by VPA and the expected alterations in PNN density could alter the connectivity of cerebellar circuitry leading to cognitive behavioral deficits.

Dendrite development. *P2rx7* **(−2.0)** (ionotrophic P2X7 receptor, which is activated by extracellular ATP) has been described as the "central hub of brain diseases" (Andrejew et al., 2020). Inhibition of P2X7 receptors ameliorated dendritic spine pathology under pathological conditions (in *Mecp2*-deficient mice) (Garré et al., 2020), while P2rx7 deficiency caused dendritic branching deficits on otherwise normal mice (Mut-Arbona et al., 2023). Interestingly, *Tmem163* **(−1.7)** is required for full function of P2X7 receptors (Salm et al., 2020). *Sema3a* **(−2.0)** regulates dendritic complexity via *Farp1* **(−1.7)** in an activity dependent manner (Cheadle and Biederer, 2014). *Farp1* links postsynaptic cytoskeletal

dynamics and transsynaptic organization to coordinate synaptic development (Cheadle and Biederer, 2012). In contrast, *Itpka* **(+14.1)**, which has been reported to regulate dendrite morphology, was over-expressed by 14-fold in fetal brain exposed to VPA (Windhorst et al., 2012). Thus, the reduction in expression of *P2rx7*, *Tmem163*, *Sema3a* and *Farp1* and the increase in *Itpka* induced by VPA may work synergistically to alter dendritic branching of developing neurons in the E12.5 brain. *Icam5* **(+16.6)** (intercellular adhesion molecule 5) is overexpressed in the brain of the *Fmr1* knockout mouse (which models fragile X syndrome, an ASD), leading to dendritic spine abnormalities. The massive increase in *Icam5* expression induced by VPA has the potential to severely disrupt the connectivity of the develop-ing brain leading to autism-like behavior. *Gas7* **(−1.7)** encodes a spine initiation factor that responds to neuronal activity; *Gas7* knockdown decreased spine density in hippocampal neurons (Khanal et al., 2023). *Dlg5* **(−3.0)** encodes a membrane-associated guanylate kinase (MAGUK) protein, which is re-quired for dendritic spine formation and synaptogenesis (Wang et al., 2014), raising the possibility that synaptic development and organization may be impacted by the 67% reduction *Dlg5* expression in fetal brain exposed to VPA at E12.5. *Disc1* **(−1.9)** (Disrupted in schizophrenia-1) has been implicated in schizophrenia. Normal expression levels of *Disc1* are required for microtubule function and depletion of *Disc1* or dominant negative *Disc1* constructs impairs neurite outgrowth *in vitro* and proper develop-ment of the cortex *in vivo* (Kamiya et al., 2005).

Postsynaptic cytoskeleton. The organization and assembly of glutamate receptors in the postsyn-aptic membrane are dependent on scaffold proteins including members of the HOMER and SHANK families. (Guang et al., 2018) *Homer1* **(+1.6)** and *Shank2* **(−2.3)** are associated with autism by GWAS (Table 1) and are responsible for rare syndromic cases of autism. *Homer3* **(−1.4)** is also dysregulated in the fetal brain. (Eltokhi et al., 2018; Ronesi et al., 2012; Zaslavsky et al., 2019; Yoon et al., 2021).

Excitation-inhibition. A widely discussed hypothesis is that autism is caused by an imbalance be-tween neuronal excitation and inhibition due to increased excitation, reduced inhibition, or both (Ru-benstein and Merzenich, 2003; Nelson and Valakh, 2015). Consequently, alterations in the develop-mental program in the fetal brain that result in too many excitatory (glutamatergic) or too few inhibitory (GABAergic) neurons (or synapses) could contribute to autistic behavior. Two related genes, *Maf* **(−1.6)** and *Mafb* **(−1.6)**, which are both downregulated by about 40% by VPA, have been reported to play redundant roles in the generation of interneurons from the fetal median ganglionic eminence resulting in decreased numbers of cortical somatostatin-releasing, inhibitory interneurons (Pai et al., 2019). *Gad1* **(−2.5)** and *Gad2* **(−3.3)** encode isoforms of the enzyme that synthesizes the inhibitory transmitter, GABA*; Gad1* and *Gad2* were reduced by 60% and 70%, respectively, however, reduced *Gad* expres-sion may be a consequence of a deficit in the number inhibitory GABAergic neurons. Taken together, these findings are consistent with VPA increasing net brain excitation by decreasing the number of

inhibitory interneurons generated during fetal brain development. *Nf1* (−1.3) encodes NF1, which is mutated in neurofibromatosis type 1, an inherited neurocutaneous disorder associated with neurodevelopmental disorders including autism. *Nf1* deletion results in the specific loss of parvalbumin-expressing inhibitory cortical interneurons (Angara et al., 2020). In the present study, *Nf1* was downregulated by about 25% in fetal mouse brains exposed to VPA. These reductions in gene expression may contribute to a decrease in inhibition in mice exposed to VPA *in utero* (Markram et al., 2008).

<u>Dopaminergic system</u>. Several lines of evidence link autistic symptoms with dysfunction in the mesencephalic DA system (Langen et al., 2014; Subramanian et al., 2017; DiCarlo and Wallace, 2022). The former can be explained by abnormal circuitry (Golden et al., 2018), notably in the indirect pathway downstream from D2-receptor expressing medium spiny neurons (MSNs). The subthalamic nucleus (STN), an excitatory nucleus in the indirect pathway downstream from the D2-MSNs, is a key factor in the normal functioning of the basal ganglia. *Foxa1* (−1.6), which is reduced by 38% by VPA, has been reported to be essential for the development and functional integrity of the STN (Ferri et al., 2007; Lee et al., 2010; Gasser et al., 2016). *Foxa1* and *Lmx1a* (−1.7), which are both reduced by about 40% in the E12.5 brain exposed to VPA (Table 2), have been shown to promote mesencephalic DA neuron development and diversity (Yan et al., 2011; Doucet-Beaupré et al., 2015; Lin et al., 2009; Deng et al., 2011; Bodea and Blaess, 2015). The basic helix-loop-helix transcription factor *Srebf1* (+1.4 significant in males only) has been shown to be both necessary and sufficient for midbrain dopaminergic neurogenesis (Toledo et al., 2020). Engeln et al. (2021) reported that transgenic mice with the BDNF receptor, trkB, knocked out specifically in D1 MSNs, displayed autism-like stereotypic behavior that was reversed by chemogenetic inhibition of D1-MSNs. Using RNAseq of ribosome-associated mRNA, the authors found that 26 genes were dysregulated in the transgenic mouse D1-MSNs. Of these 17 were significantly dysregulated by VPA in the present study [*Slc9a3r1* (+3.0), *Csf1r* (−1.4), *Sparc* (−1.4), *Mcam* (+1.7), *Robo4* (−2.2), *Emp2* (−1.6), *Klf4* (−1.6), *Rtn4rl1* (+2.0), *Bdnf* (+2.8), *C1qtnf1* (−1.4), *Tmem108* (+1.4), *Smyd2* (−1.3), *Wnt7b* (−1.8), *Palm3* (+1.9), *Adora2a* (−1.3), *Drd2* (+2.0), *S1pr1* (−1.3)]. As discussed above, another gene linked to the development of the DA system is *Nr2f1* (−1.8), which was downregulated by 45% by VPA. It is of interest that all three DA receptor subtypes (*Drd1*, *Drd2*, *Drd3*) are linked to autism by GWAS (SFARI List) and that partial antagonists of the D2 DA receptor (encoded by *Drd2*) have demonstrated utility in reducing stereotypic behaviors in human subjects (Mandic-Maravic et al., 2022); *Drd2* (+2.2) expression was doubled by VPA in the present study. Tyrosine hydroxylase encoded by the *Th* (−1.4) gene is the rate limiting step in the biosynthesis of catecholamines including DA; *Th* is downregulated by about 35% in fetal brain exposed to VPA. *Ppp1r1b* (+3.7), encodes the dopamine- and cAMP-regulated neuronal phosphoprotein (DARPP-32). DARPP-32 amplifies and/or mediates many actions of cyclic AMP-dependent protein kinase at the

plasma membrane and in the cytoplasm, with a broad spectrum of potential targets and functions within the dopaminergic system and throughout the brain (Yger and Girault, 2011).

Cholinergic system. Cholinergic projection neurons in the nucleus basalis and cholinergic interneurons in the striatum are specified in the embryonic forebrain beginning at about E10 in mouse (Allaway and Machold, 2017). This process is regulated by several transcription factors including *Lhx8* **(−2.7)**, *Isl1* **(−1.5)**, and *Gbx1* **(−2.3)** all of which are downregulated in response to VPA in the fetal mouse brain. In the developing forebrain, *Isl1-Lhx8* hexameric complexes promote cholinergic identity (Cho et al., 2014). Conditional deletion of *Isl1* depletes cholinergic projection- and interneurons resulting in abolition of cholinergic innervation of the cortex (Elshatory and Gan, 2008). *Nkx2-1* **(−2.5**, significant in females only), a homeodomain transcription factor, is required for the development of cholinergic septohippocampal neurons and large subsets of basal forebrain cholinergic neurons (Magno et al., 2017). The 33−67% decrease in the expression of these transcription factors induced in the fetal brain by VPA has the potential to alter the proper development of circuits critical for normal brain function. Indeed, the loss of these neurons due to lack of *Nkx2-1* causes alterations in hippocampal theta rhythms and learning and memory defects (Magno et al., 2017). In addition, *Nkx2-1* acting together with *Lhx6* (not affected by VPA), is necessary to specify pallidal projection neurons and forebrain interneurons in the medial ganglionic eminence (Sandberg et al., 2016).

Endocannabinoid system. *Cnr1* **(−2.0)**, encodes the primary endocannabinoid **(**eCB) receptor in the brain; *Cnr1* has been liked to autism in GWAS. eCBs have been reported to regulate a wide range of embryonic neurodevelopmental processes (Harkany et al., 2007; Gomes et al., 2020) including neural progenitor proliferation (Aguado et al., 2005), long-range axon patterning (Mulder et al., 2008), interneuron migration and morphogenesis (Berghuis et al., 2005), and neuronal fate specification (Harkany et al., 2008). A 50% decrease in expression of *Cnr1* would be predicted to alter the cellular responses to eCB and could disrupt normal fetal brain development (Chakrabarti et al., 2015).

Autism and Down syndrome (DS). While the cause of autism is unknown, DS is caused by triplication of human chromosome 21 (Hsa21), leading to the hypothesis that a 50% increase in gene dosage of one or more Hsa21 genes is the cause of DS symptoms. ASDs and DS share ID as a common feature; moreover, DS individuals are diagnosed with ASD at a higher frequency than the general population (Hepburn et al., 2008; Rachubinski et al., 2017). In fact, several genes triplicated in DS that have been liked to autism are up- or downregulated by VPA in the fetal brain; these include *Dyrk1a* **(−1.5)**, *Brwd1* **(−1.3)** and *Cbs* **(+5.0)** (Park and Chung, 2013; Duchon and Herault, 2016; Fulton et al., 2022; Marechal et al., 2019). ID in both ASDs and DS has been attributed to overexpression of *Cbs* (cystathionine beta synthase) (Conan et al., 2023; Tisato et al., 2021); dysregulation of *Cbs* and

dihydrofolate reductase (*Dhfr*, +1.5) contributes to inborn errors of amino acid metabolism and is linked to ASDs (Zigman et al., 2021; Hoxha et al., 2021). *Dyrk1a* (which encodes dual specificity tyrosine phosphorylation regulated kinase 1A) is believed to underlie intellectual disability in DS due to its increased gene dosage in DS (Duchon and Herault, 2016). In contrast, haploinsufficiency of *DYRK1A* results in a (non-DS) syndrome characterized by intellectual disability including impaired speech development, autism spectrum disorder including anxious and/or stereotypic behavior problems, and microcephaly (Levy et al., 2021; Dang et al., 2018).

Circadian rhythms. Eleven genes involved in generating circadian rhythms were significantly affected by VPA in the fetal mouse brain (Table 5). Two of these (*Per1* and *Per1*) are SFARI risk genes. While **Per1 (+1.9), Per2 +1.7), Cry1 (+2.5), Cry 2 (+2.0), Arntl (+2.0) and Mef2d (+1.5)** (Geoffray et al., 2016; Mohawk et al., 2019; Lorsung et al., 2021) were upregulated by VPA, *Clock* was not affected. **Fbxl3 (−1.8)** a negative regulator of *Cry1* and *Cry2*, is downregulated by VPA whereas **Fbxl21 (+3.2)**, which is up-regulated by VPA, down-regulates *Fbxl3* (Hirano et al., 2013). Thus, these VPA-induced changes lead to increased *Per* and *Cry* gene expression. **Arntl (Bmal1) (+2.1)**, together with *Clock*, activates rhythmic transcription of *Per* and *Cry* genes; knockout of *Arntl* (*Bmal1*) induces autistic-like behavior and cerebellar dysfunction which was ameliorated by mTORC1 inhibition (Liu et al., 2022).

Daily rhythms in the fetal brain are thought to be entrained by the mother, however, whether there is a functional *Per/Cry/Clock* cycle in the fetal brain has not been explored. It is possible that disrupting the expression of these genes could alter early embryonic brain development through a mechanism unrelated to sleep. For example, (Noda et al., 2019) reported that *in utero* knockdown of *Per3* caused abnormal positioning of cortical excitatory neurons as well as impaired axon extension and dendritic arbor formation.

If VPA-induced changes in circadian rhythm gene expression were to persist postnatally, they could cause disrupted sleep patterns, a common occurrence in autism. Indeed, exposure of fetal rats to VPA on E12.5 caused sleep disturbances that were similar to those reported in autistic children (Cusmano and Mong, 2014); moreover, a VPA-induced reduction in GAD-67 (*Gad1*) was also found in the juvenile rat brains. In the present study, **Gad1 (−2.5)** was reduced by 60% while RNAs encoding GABA-A and -B receptors subunits, encoded by **Gabra4 (+2.5)** and **Gabbr2 (−2.4),** were found to be upregulated 2.5-fold and downregulated 60% by VPA, respectively.

Neuroinflammation. Immune dysregulation is believed to play a key role in the pathogenesis of autism. Neuroinflammatory molecules such as insulin-like growth factors [**Igf1 (−3.2), Igf2 (−2.7)**], transforming growth factor-β isoforms [**Tgfb2 (−2.4), Tgfb3 (−2.0)**], and the TGF-β receptor, **Tgfbr2 (−2.5)**, are decreased in fetal brain in response to VPA exposure. Expression of several heat-shock proteins

[*Hsp1a* (+5.9)*, Hsp1b* (+5.6)*, Hsp2* (+6.2) and *Dnajc12* (+4.5)]] was dramatically increased by VPA exposure. This may be a response to immune dysregulation following the administration of a high dose of VPA. *DNAJC12* encodes a chaperone protein responsible for the proper folding of phenylalanine hydrolase. *DNAJC12* deficiency causes hyperphenylalaninemia leading symptoms ranging from mild autistic features or hyperactivity to severe intellectual disability (Blau et al., 2018; Anikster et al., 2017).

Epigenetic regulation of gene expression in the action of VPA. The mechanism(s) by which VPA alters gene expression in the fetal brain is not known; however, VPA is a Class I histone deacetylase (HDAC) inhibitor, suggesting that many of the changes reported here may involve increased histone acetylation (Bourin, 2020; Gurvich et al., 2004). Indeed, Konopko et al. (2017) reported that VPA increased the acetylation of several histones at the promoter regions of *Bdnf* 5' untranslated exons 1, 4, and 6 and that this was positively correlated with increased expression of exon 9, the protein coding region. It is also likely that many of the genes targeted by VPA are themselves epigenetic writers, erasers, or readers; moreover, there is well-established crosstalk among epigenetic marks (Berger, 2007), including between DNA CpG methylation and covalent histone modifications, which could explain why HDAC inhibition could both increase and decrease gene expression. **Setd1b (−2.2)** encodes a lysine-specific methyltransferase which has been linked to autism in GWAS studies (Table 1). De novo variants of *SET1B* are associated with intellectual disability, ASD and epilepsy (Hiraide et al., 2019; Roston et al., 2021). **Kmt5b (−6.5)** encodes the epigenetic writer, lysine methyl transferase 5B, which methylates lysine residues on histones, resulting in gene activation or silencing, depending on the histone and lysine residue. Loss of function mutations in *KMT5B* lead to developmental delay and ASD (Eliyahu et al., 2022) (SFARI List). *Kmt5b* expression is reduced by 85% in fetal mouse brains exposed to VPA (Table 1). **Mecp2 (−1.6)** is an X-linked gene that encodes the epigenetic reader, methyl cytosine binding protein 2. Mutations in *MECP2* cause Rett syndrome, an ASD that affects mostly girls (Chahrour and Zoghbi, 2007); *MECP2* is on the SFARI autism list. MeCP2 binds to methylated CpGs in DNA mediating gene repression (Ip et al., 2018, Gonzales and LaSalle, 2010; Guy et al., 2011). Either increased or decreased expression of MeCP2 can cause developmental brain disorders. MeCP2 binds to CpGs throughout the genome; reduced MeCP2 would be expected to alter the expression of many genes. One such gene is **Bdnf (+2.8)**. MeCP2 binds to CpGs in the 5'UTR of exon 4 of *Bdnf* leading to suppression of BDNF expression (Martinowich et al., 2003). Reduction of MeCP2 as observed with VPA (Table 1), would be predicted to decrease normal epigenetic suppression of CpG methylated genes. Although reduced *Mecp2* expression induced by VPA would be expected to increase BDNF expression, VPA did not alter CpG methylation in the regulatory regions of *Bdnf*; instead, increased *Bdnf* expression was due to increased histone acetylation and methylation (Konopko et al., 2017).

Paulsen et al. (2022) reported that haploinsufficiency in the chromatin remodelers, *Kmt5b, Arid1b* and *Chd8*, increased the numbers of inhibitory GABAergic neurons in organoid models of human cortex leading to reduced spontaneous circuit activity in the organoids. In the present study, **Kmt5b (−6.5)** and **Arid1b (−2.1)** were down regulated in the fetal brain by 85% and 50% respectively, in response to VPA. Although *Chd8* expression was unaffected by VPA, **Chd2**, **Chd3** and **Chd7**, which have similar chromatin remodeling functions (Marfella and Imbalzano, 2007), were reduced by 23%, 50% and 23% respectively.

Caveats:

1. The results reported here are restricted to VPA-induced changes in RNA levels; the extent to which these changes reflect commensurate changes in the expression levels of the proteins encoded by these genes is not known. Although in many cases, RNA and protein levels are correlated, the extent to which RNA levels of a given gene track its corresponding protein levels may vary.

2. These results represent VPA-induced changes in gene expression at a single point in time (3 hr after VPA administration on E12.5). Whether the changes are transient and reverse as the VPA levels dissipate, as has been reported for *Bdnf* (Almeida et al., 2014), or are persistent, is not known. It is also not known whether the same VPA induced changes would be observed after VPA administration on a different gestational day.

3. There are many genes for which expression is increased or decreased by ≥ 1.25-fold or ≤ −1.25-fold in response to VPA but the FDR-corrected p-value did not reach the p_{FDR} ≤ 0.025 criterion applied in this study (Supplementary Table S1). Higher statistical power would be needed to determine whether these changes are significant.

4. The ± 1.25 FC cutoff used here is arbitrary; it is possible that smaller changes in the expression of one or more critical genes have profound effects on fetal brain development and these would not be identified in this analysis.

5. Only a fraction of the approximately 7,300 genes whose expression is significantly affected by VPA were curated in this study. It is likely that other genes, in addition to those shown in Tables 1 and 2, could play a role in mediating the effects of VPA on fetal brain development.

Reproducibility:

The stimulation by VPA of *Bdnf* expression (Table 2) has been independently confirmed by quantitative RT-PCR (Almeida et al., 2014; Konopko et al., 2017).

By separately analyzing gene expression in male and female brains, the results of this study were effectively replicated with independent biological samples (7 of each sex ± VPA, each from a different pregnancy). With very few exceptions, male and female gene expression levels were similar (c.f., Figure 1) demonstrating reproducibility across independent samples.

SUMMARY:

The approach for this study can be described as "hypothesis generating" in that we began without any preconceived idea (hypothesis) as to the mechanism by which a brief, transient dose of VPA, administered to the pregnant mouse at E12.5, can cause abnormal, autistic-like behavior in her offspring many weeks later. The analysis identified approximately 7,300 genes, expression of which was significantly affected by VPA. This set of genes was narrowed down in two ways, viz., identifying those genes significantly affected by VPA that were a) also linked to autism by GWAS (SFARI List; Table 1) and b) not on the SFARI list (Table 2), but known to play a role in critical steps of early brain development such as neuroprogenitor proliferation and fate determination, axon and dendrite growth, and synaptogenesis. Interference with one or more of these steps in the fetal brain has the potential to interfere with the "program" directing brain development to create a persistent pathological "signature" that leads to abnormal neuronal circuitry in the adult, long after the increase in VPA in the fetal brain and the initial change in gene expression have dissipated. Of these 647 genes, about half have known mechanisms of action associated with brain development or function; none showed significant sexually-dimorphic effects of VPA. Several genes affected by VPA in the fetal mouse brain appear in multiple GWAS studies (c.f., Tables 3 and 4) including *Adnp*, *Arid1b*, *Ash1l*, *Cdh2*, *Cdk5l*, *Dyrk1a*, *Kmt5b*, *Mecp2*, *Nr2f1*, *Pogz*, *Pten*, *Tbr1*, *Tcf7l2*, and *Wac*. This "short list" is a potential starting point for future hypothesis-driven studies to determine whether dysregulation of one or more of these genes by VPA is a proximal cause of the behavioral abnormalities identified in the adult animals. Nevertheless, it remains possible that dysregulation of one or more of the other genes altered by VPA contributes to the proximal cause of the autistic-like behavior. This hypothesis could be tested by determining the effect of manipulating gene expression (singly or in combination) on VPA-induced behavioral abnormalities.

REFERENCES:

Acampora D, Postiglione MP, Avantaggiato V, Bonito M Di, Vaccarino FM, Michaud J, Simeone A (1999) Progressive impairment of developing neuroendocrine cell lineages in the hypothalamus of mice lacking the Orthopedia gene. Genes Dev 13:2787–2800 Available at: www.genesdev.org.

Aguado T, Monory K, Palazuelos J, Stella N, Cravatt B, Lutz B, Marsicano G, Kokaia Z, Guzmán M, Galve-Roperh I (2005) The endocannabinoid system drives neural progenitor proliferation. FASEB J 19:1704–1706.

Ahmed G, Shinmyo Y (2021) Multiple Functions of Draxin/Netrin-1 Signaling in the Development of Neural Circuits in the Spinal Cord and the Brain. Front Neuroanat 15:766911.

Allaway KC, Machold R (2017) Developmental specification of forebrain cholinergic neurons. Dev Biol 421:1–7.

Almuriekhi M, Shintani T, Fahiminiya S, Fujikawa A, Kuboyama K, Takeuchi Y, Nawaz Z, Nadaf J, Kamel H, Kitam AK, Samiha Z, Mahmoud L, Ben-Omran T, Majewski J, Noda M (2015) Loss-of-function mutation in APC2 causes Sotos syndrome features. Cell Rep 10:1585–1598.

American Psychiatric Association (2022) Diagnostic and Statistical Manual of Mental Disorders (5th ed., text rev.).

Anders S, Pyl PT, Huber W (2015) HTSeq-A Python framework to work with high-throughput sequencing data. Bioinformatics 31:166–169.

Andrejew R, Oliveira-Giacomelli Á, Ribeiro DE, Glaser T, Arnaud-Sampaio VF, Lameu C, Ulrich H (2020) The P2X7 Receptor: Central Hub of Brain Diseases. Front Mol Neurosci 13:Article 124.

Angara K, Ling-Lin Pai E, Bilinovich SM, Stafford AM, Nguyen JT, Li KX, Paul A, Rubenstein JL, Vogt D (2020) Nf1 deletion results in depletion of the Lhx6 transcription factor and a specific loss of parvalbumin + cortical interneurons. Proc Natl Acad Sci USA 117:6189–6195.

Anikster Y et al. (2017) Biallelic Mutations in DNAJC12 Cause Hyperphenylalaninemia, Dystonia, and Intellectual Disability. Am J Hum Genet 100:257–266.

Armentano M, Chou SJ, Srubek Tomassy G, Leingärtner A, O'Leary DDM, Studer M (2007) COUP-TFI regulates the balance of cortical patterning between frontal/motor and sensory areas. Nat Neurosci 10:1277–1286.

Armentano M, Filosa A, Andolfi G, Studer M (2006) COUP-TFI is required for the formation of commissural projections in the forebrain by regulating axonal growth. Development 133:4151–4162.

Arnold SJ, Huang GJ, Cheung AFP, Era T, Nishikawa SI, Bikoff EK, Molnár Z, Robertson EJ, Groszer M (2008) The T-box transcription factor Eomes/Tbr2 regulates neurogenesis in the cortical subventricular zone. Genes Dev 22:2479–2484.

Aruga J, Mikoshiba K (2003) Identification and characterization of Slitrk, a novel neuronal transmembrane protein family controlling neurite outgrowth. Mol Cell Neurosci 24:117–129.

Baala L, Briault S, Etchevers HC, Laumonnier F, Natiq A, Amiel J, Boddaert N, Picard C, Sbiti A, Asermouh A, Attié-Bitach T, Encha-Razavi F, Munnich A, Sefiani A, Lyonnet S (2007) Homozygous silencing of T-box transcription factor EOMES leads to microcephaly with polymicrogyria and corpus callosum agenesis. Nat Genet 39:454–456.

Bannai H, Fukatsu K, Mizutani A, Natsume T, Iemura SI, Ikegami T, Inoue T, Mikoshiba K (2004) An RNA-interacting protein, SYNCRIP (heterogeneous nuclear ribonuclear protein Q1/NSAP1) is a component of mRNA granule transported with inositol 1,4,5-trisphosphate receptor type 1 mRNA in neuronal dendrites. J Biol Chem 279:53427–53434.

Beaubien F, Raja R, Kennedy TE, Fournier AE, Cloutier JF (2016) Slitrk1 is localized to excitatory synapses and promotes their development. Sci Rep 6:27343.

Ben-Ari Y (2008) Neuro-archaeology: pre-symptomatic architecture and signature of neurological disorders. Trends Neurosci 31:626–636.

Berger SL (2007) The complex language of chromatin regulation during transcription. Nature 447:407–412.

Berghuis P, Dobszay MB, Wang X, Spano S, Ledda F, Sousa KM, Schulte G, Ernfors P, Mackie K, Paratcha G, Hurd YL, Harkany T (2005) Endocannabinoids regulate interneuron migration and morphogenesis by transactivating the TrkB receptor. Proc Natl Acad Sci USA 102:19115–19120.

Berry CT, Sceniak MP, Zhou L, Sabo SL (2012) Developmental Up-Regulation of Vesicular Glutamate Transporter-1 Promotes Neocortical Presynaptic Terminal Development. PLoS One 7:e50911.

Bertacchi M et al. (2020) NR2F1 regulates regional progenitor dynamics in the mouse neocortex and cortical gyrification in BBSOAS patients. EMBO J 39:e104163.

Bey AL, Jiang Y hui (2014) Overview of mouse models of autism spectrum disorders. Curr Protoc Pharmacol 2014:5.66.1-5.66.26.

Bilbo SD, Block CL, Bolton JL, Hanamsagar R, Tran PK (2018) Beyond infection - Maternal immune activation by environmental factors, microglial development, and relevance for autism spectrum disorders. Exp Neurol 299:241–251.

Blau N, Martinez A, Hoffmann GF, Thöny B (2018) DNAJC12 deficiency: A new strategy in the diagnosis of hyperphenylalaninemias. Mol Genet Metab 123:1–5.

Bodea GO, Blaess S (2015) Establishing diversity in the dopaminergic system. FEBS Lett 589:3773–3785.

Boitnott A, Garcia-Forn M, Ung DC, Niblo K, Mendonca D, Park Y, Flores M, Maxwell S, Ellegood J, Qiu LR, Grice DE, Lerch JP, Rasin MR, Buxbaum JD, Drapeau E, De Rubeis S (2021) Developmental and Behavioral Phenotypes in a Mouse Model of DDX3X Syndrome. Biol Psychiatry 90:742–755.

Borrell V, Götz M (2014) Role of radial glial cells in cerebral cortex folding. Curr Opin Neurobiol 27:39–46.

Bourin M (2020) Mechanism of Action of Valproic Acid and Its Derivatives. SOJ Pharm Pharm Sci 7:1–4.

Breitenkamp AFS, Matthes J, Nass RD, Sinzig J, Lehmkuhl G, Nürnberg P, Herzig S (2014) Rare mutations of CACNB2 found in autism spectrum disease-affected families alter calcium channel function. PLoS One 9.

Buddell T, Friedman V, Drozd CJ, Quinn CC (2019) An autism-causing calcium channel variant functions with selective autophagy to alter axon targeting and behavior. PLoS Genet 15.

Burke RD, Todd SW, Lumsden E, Mullins RJ, Mamczarz J, Fawcett WP, Gullapalli RP, Randall WR, Pereira EFR, Albuquerque EX (2017) Developmental neurotoxicity of the organophosphorus insecticide chlorpyrifos: from clinical findings to preclinical models and potential mechanisms. J Neurochem 142 Suppl 2:162–177.

Careaga M, Murai T, Bauman MD (2017) Maternal Immune Activation and Autism Spectrum Disorder: From Rodents to Nonhuman and Human Primates. Biol Psychiatry 81:391–401.

Chahrour M, Zoghbi HY (2007) The story of Rett syndrome: from clinic to neurobiology. Neuron 56:422–437.

Chakrabarti B, Persico A, Battista N, Maccarrone M (2015) Endocannabinoid Signaling in Autism. Neurotherapeutics 12:837–847.

Chakrabarti L, Best TK, Cramer NP, Carney RSE, Isaac JTR, Galdzicki Z, Haydar TF (2010) Olig1 and Olig2 triplication causes developmental brain defects in Down syndrome. Nat Neurosci 13:927–934.

Chatterjee M, Singh P, Xu J, Lombroso PJ, Kurup PK (2020) Inhibition of striatal-enriched protein tyrosine phosphatase (STEP) activity reverses behavioral deficits in a rodent model of autism. Behavioural Brain Research 391:112713.

Chatzi C, Cunningham TJ, Duester G (2013) Investigation of retinoic acid function during embryonic brain development using retinaldehyde-rescued Rdh10 knockout mice. Developmental Dynamics 242:1056–1065.

Cheadle L, Biederer T (2012) The novel synaptogenic protein farp1 links postsynaptic cytoskeletal dynamics and transsynaptic organization. J Cell Biol 199:985–1001.

Cheadle L, Biederer T (2014) Activity-dependent regulation of dendritic complexity by Semaphorin 3A through Farp1. J Neurosci 34:7999–8009.

Cho H-H, Cargnin F, Kim Y, Lee B, Kwon R-J, Nam H, Shen R, Barnes AP, Lee JW, Lee S, Lee S-K (2014) Isl1 directly controls a cholinergic neuronal identity in the developing forebrain and spinal cord by forming cell type-specific complexes. PLoS Genet 10:e1004280.

Christensen J, Grønborg TK, Sørensen MJ, Schendel D, Parner ET, Pedersen LH, Vestergaard M (2013) Prenatal Valproate Exposure and Risk of Autism Spectrum Disorders and Childhood Autism. JAMA 309:1696–1703.

Chuang HC, Huang TN, Hsueh YP (2015) T-Brain-1 - A Potential Master Regulator in Autism Spectrum Disorders. Autism Research 8:412–426.

Co M, Barnard RA, Jahncke JN, Grindstaff S, Fedorov LM, Adey AC, Wright KM, O'Roak BJ (2022) Shared and distinct functional effects of patient-specific Tbr1 mutations on cortical development . J Neurosci 42:7166-7181.

Conan P, Léon A, Caroff N, Rollet C, Chaïr L, Martin J, Bihel F, Mignen O, Voisset C, Friocourt G (2023) New insights into the regulation of Cystathionine beta synthase (CBS), an enzyme involved in intellectual deficiency in Down syndrome. Front Neurosci 16:1110163.

Cusmano DM, Mong JA (2014) In Utero exposure to valproic acid changes sleep in juvenile rats: A model for sleep disturbances in autism. Sleep 37:1489–1499.

Dang T, Duan WY, Yu B, Tong DL, Cheng C, Zhang YF, Wu W, Ye K, Zhang WX, Wu M, Wu BB, An Y, Qiu ZL, Wu BL (2018) Autism-associated Dyrk1a truncation mutants impair neuronal dendritic and spine growth and interfere with postnatal cortical development. Mol Psychiatry 23:747–758.

Deng Q, Andersson E, Hedlund E, Alekseenko Z, Coppola E, Panman L, Millonig JH, Brunet JF, Ericson J, Perlmann T (2011) Specific and integrated roles of Lmx1a, Lmx1b and Phox2a in ventral midbrain development. Development 138:3399–3408.

DiCarlo GE, Wallace MT (2022) Modeling dopamine dysfunction in autism spectrum disorder: From invertebrates to vertebrates. Neurosci Biobehav Rev 133:104494.

Dickinson RG, Lawyer, CH, Kaufman SN, Lynn RK, Gerber N, Novy MJ, Cook MJ (1980) Materno-fetal pharmacokinetics and fetal distribution of valproic acid in a pregnant rhesus monkey. Pediatric Pharmacol (New York) 1:71–83.

Dong C, West KL, Tan XY, Li J, Ishibashi T, Yu C, Sy SMH, Leung JWC, Huen MSY (2020) Screen identifies DYRK1B network as mediator of transcription repression on damaged chromatin. Proc Natl Acad Sci USA 117:17019–17030.

Doucet-Beaupré H, Ang SL, Lévesque M (2015) Cell fate determination, neuronal maintenance and disease state: The emerging role of transcription factors Lmx1a and Lmx1b. FEBS Lett 589:3727–3738.

Duchon A, Herault Y (2016) DYRK1A, a dosage-sensitive gene involved in neurodevelopmental disorders, Is a target for drug development in down syndrome. Front Behav Neurosci 10:104.

Edamatsu M, Miyano R, Fujikawa A, Fujii F, Hori T, Sakaba T, Oohashi T (2018) Hapln4/Bral2 is a selective regulator for formation and transmission of GABAergic synapses between Purkinje and deep cerebellar nuclei neurons. J Neurochem 147:748–763.

el Amri M, Fitzgerald U, Schlosser G (2018) MARCKS and MARCKS-like proteins in development and regeneration. J Biomed Sci 25:43.

el Chehadeh S et al. (2022) SLITRK2 variants associated with neurodevelopmental disorders impair excitatory synaptic function and cognition in mice. Nat Commun 13:4112.

Eliyahu A, Barel O, Greenbaum L, Zaks Hoffer G, Goldberg Y, Raas-Rothschild A, Singer A, Bar-Joseph I, Kunik V, Javasky E, Staretz-Chacham O, Pode-Shakked N, Bazak L, Ruhrman-Shahar N, Pras E, Frydman M, Shohat M, Pode-Shakked B (2022) Refining the Phenotypic Spectrum of KMT5B-Associated Developmental Delay. Front Pediatr 10:Article 844845.

Elsen GE, Hodge RD, Bedogni F, Daza RAM, Nelson BR, Shiba N, Reiner SL, Hevner RF (2013) The protomap is propagated to cortical plate neurons through an Eomes-dependent intermediate map. Proc Natl Acad Sci U S A 110:4081–4086.

Elshatory Y, Gan L (2008) The LIM-homeobox gene Islet-1 is required for the development of restricted forebrain cholinergic neurons. J Neurosci 28:3291–3297.

Eltokhi A, Rappold G, Sprengel R (2018) Distinct Phenotypes of Shank2 Mouse Models Reflect Neuropsychiatric Spectrum Disorders of Human Patients With SHANK2 Variants. Front Mol Neurosci 11:240.

Endres D, Decher N, Röhr I, Vowinkel K, Domschke K, Komlosi K, Tzschach A, Gläser B, Schiele MA, Runge K, Süß P, Schuchardt F, Nickel K, Stallmeyer B, Rinné S, Schulze-Bahr E, van Elst LT (2020) New CaV1.2 channelopathy with high-functioning autism, affective disorder, severe dental enamel defects, a short QT interval, and a novel cacna1c loss-of-function mutation. Int J Mol Sci 21:1–8.

Engeln M, Song Y, Chandra R, La A, Fox ME, Evans B, Turner MD, Thomas S, Francis TC, Hertzano R, Lobo MK (2021) Individual differences in stereotypy and neuron subtype translatome with TrkB deletion. Mol Psychiatry 26:1846–1859.

Englund C, Fink A, Lau C, Pham D, Daza RAM, Bulfone A, Kowalczyk T, Hevner RF (2005) Pax6, Tbr2, and Tbr1 are expressed sequentially by radial glia, intermediate progenitor cells, and postmitotic neurons in developing neocortex. J Neurosci 25:247–251.

Faedo A, Tomassy GS, Ruan Y, Teichmann H, Krauss S, Pleasure SJ, Tsai SY, Tsai MJ, Studer M, Rubenstein JLR (2008) COUP-TFI coordinates cortical patterning, neurogenesis, and laminar fate and modulates MAPK/ERK, AKT, and β-catenin signaling. Cerebral Cortex 18:2117–2131.

Fan L, Lu Y, Shen X, Shao H, Suo L, Wu Q (2018) Alpha protocadherins and Pyk2 kinase regulate cortical neuron migration and cytoskeletal dynamics via Rac1 GTPase and WAVE complex in mice. Elife 7:e35242.

Fazel Darbandi S, Robinson Schwartz SE, Qi Q, Catta-Preta R, Pai ELL, Mandell JD, Everitt A, Rubin A, Krasnoff RA, Katzman S, Tastad D, Nord AS, Willsey AJ, Chen B, State MW, Sohal VS, Rubenstein JLR (2018) Neonatal Tbr1 Dosage Controls Cortical Layer 6 Connectivity. Neuron 100:831-845.e7.

Ferreira A, Chin L-S, Lian L, Lanier LM, Kosik KS, Greengard P (1998) Distinct roles of synapsin I an synapsin II during neuronal development. Molecular Medicine 4:2228.

Ferri ALM, Lin W, Mavromatakis YE, Wang JC, Sasaki H, Whitsett JA, Ang SL (2007) Foxa1 and Foxa2 regulate multiple phases of midbrain dopaminergic neuron development in a dosage-dependent manner. Development 134:2761–2769.

Ferri SL, Abel T, Brodkin ES (2018) Sex Differences in Autism Spectrum Disorder: a Review. Curr Psychiatry Rep 20:9.

Fulton SL et al. (2022) Rescue of deficits by Brwd1 copy number restoration in the Ts65Dn mouse model of Down syndrome. Nat Commun 13:6384.

Garcia-Forn M, Boitnott A, Akpinar Z, de Rubeis S (2020) Linking autism risk genes to disruption of cortical development. Cells 9:1–24.

García-Moreno F, Pedraza M, Di Giovannantonio LG, Di Salvio M, López-Mascaraque L, Simeone A, De Carlos JA (2010) A neuronal migratory pathway crossing from diencephalon to telencephalon populates amygdala nuclei. Nat Neurosci 13:680–689.

Garel S, Marin F, Marin M, Genevie M-G, Matte G, Vesque C, Vincent A, Charnay P (1997) Family of Ebf/Olf-1-Related Genes Potentially Involved in Neuronal Differentiation and Regional Specification in the Central Nervous System. Dev Dyn 210:191–205.

Garré JM, Silva HM, Lafaille JJ, Yang G (2020) P2X7 receptor inhibition ameliorates dendritic spine pathology and social behavioral deficits in Rett syndrome mice. Nat Com 11:1784.

Gasser E, Johannssen HC, Rülicke T, Zeilhofer HU, Stoffel M (2016) Foxa1 is essential for development and functional integrity of the subthalamic nucleus. Sci Rep 6:38611.

Geoffray MM, Nicolas A, Speranza M, Georgieff N (2016) Are circadian rhythms new pathways to understand Autism Spectrum Disorder? J Physiol Paris 110:434–438.

Golden CE, Buxbaum JD, de Rubeis S (2018) Disrupted circuits in mouse models of autism spectrum disorder and intellectual disability. Curr Opin Neurobiol 48:106–112.

Gomes TM, Dias da Silva D, Carmo H, Carvalho F, Silva JP (2020) Epigenetics and the endocannabinoid system signaling: An intricate interplay modulating neurodevelopment. Pharmacol Res 162:105237.

Gonda Y, Namba T, Hanashima C (2020) Beyond Axon Guidance: Roles of Slit-Robo Signaling in Neocortical Formation. Front Cell Dev Biol 8:Article 607415.

Gonzales ML, LaSalle JM (2010) The role of MeCP2 in brain development and neurodevelopmental disorders. Curr Psychiatry Rep 12:127–134.

Guang S, Pang N, Deng X, Yang L, He F, Wu L, Chen C, Yin F, Peng J (2018) Synaptopathology involved in autism spectrum disorder. Front Cell Neurosci 12.

Guo H et al. (2019) Disruptive variants of CSDE1 associate with autism and interfere with neuronal development and synaptic transmission. Sci Adv 5:eaax2166.

Gurvich N, Tsygankova OM, Meinkoth JL, Klein PS (2004) Histone Deacetylase Is a Target of Valproic Acid-Mediated Cellular Differentiation. Cancer Res 64:1079–1086.

Guy J, Cheval H, Selfridge J, Bird A (2011) The role of MeCP2 in the brain. Annu Rev Cell Dev Biol 27:631–652.

Hallmayer J, Cleveland S, Torres A, Phillips J, Cohen B, Torigoe T, Miller J, Fedele A, Collins J, Smith K, Lotspeich L, Croen LA, Ozonoff S, Lajonchere C, Grether JK, Risch N (2011) Genetic heritability and shared environmental factors among twin pairs with autism. Arch Gen Psychiatry 68:1095–1102.

Han S, Dennis DJ, Balakrishnan A, Dixit R, Britz O, Zinyk D, Touahri Y, Olender T, Brand M, Guillemot F, Kurrasch D, Schuurmans C (2018) A non-canonical role for the proneural gene neurog1 as a negative regulator of neocortical neurogenesis. Development 145:dev157719.

Hanafusa H, Kedashiro S, Tezuka M, Funatsu M, Usami S, Toyoshima F, Matsumoto K (2015) PLK1-dependent activation of LRRK1 regulates spindle orientation by phosphorylating CDK5RAP2. Nat Cell Biol 17:1024–1035.

Harkany T, Guzmán M, Galve-Roperh I, Berghuis P, Devi LA, Mackie K (2007) The emerging functions of endocannabinoid signaling during CNS development. Trends Pharmacol Sci 28:83–92.

Harkany T, Keimpema E, Barabás K, Mulder J (2008) Endocannabinoid functions controlling neuronal specification during brain development. Mol Cell Endocrinol 286S:S84–S90.

Haushalter C, Asselin L, Fraulob V, Dollé P, Rhinn M (2017) Retinoic acid controls early neurogenesis in the developing mouse cerebral cortex. Dev Biol 430:129–141.

Hepburn S, Philofsky A, Fidler DJ, Rogers S (2008) Autism symptoms in toddlers with Down syndrome: A descriptive study. J Appl Res Intellect Disabilities 21:48–57.

Herold KG, Hussey JW, Dick IE (2023) CACNA1C-Related Channelopathies. In Handbook of Experimental Pharmacology, Springer Nature, Switzerland.

Hevner RF, Shi L, Justice N, Bulfone A, Goffinet AM, Campagnoni AT, Rubenstein JLR (2001) Tbr1 regulates differentiation of the preplate and layer 6. Neuron 29:353–366.

Hiraide T, Hattori A, Ieda D, Hori I, Saitoh S, Nakashima M, Saitsu H (2019) De novo variants in SETD1B cause intellectual disability, autism spectrum disorder, and epilepsy with myoclonic absences. Epilepsia Open 4:476–481.

Hirano A, Yumimoto K, Tsunematsu R, Matsumoto M, Oyama M, Kozuka-Hata H, Nakagawa T, Lanjakornsiripan D, Nakayama KI, Fukada Y (2013) FBXL21 regulates oscillation of the circadian clock through ubiquitination and stabilization of cryptochromes. Cell 152:1106–1118.

Hoxha B, Hoxha M, Zappacosta B, Domi E, Gervasoni J, Persichilli S, Malaj V (2021) Folic acid and autism: A systematic review of the current state of knowledge. Cells 10:1976.

Hoye ML, Calviello L, Poff AJ, Ejimogu N-E, Newman CR, Montgomery MD, Ou J, Floor SN, Silver DL (2022) Aberrant cortical development is driven by impaired cell cycle and translational control in a DDX3X syndrome model. Elife 11:e78203.

Hu JS, Vogt D, Lindtner S, Sandberg M, Silberberg SN, Rubenstein JLR (2017) Coup-TF1 and coup-TF2 control subtype and laminar identity of MGE-derived neocortical interneurons. Development 144:2837–2851.

Huang TN, Hsueh YP (2015) Brain-specific transcriptional regulator T-brain-1 controls brain wiring and neuronal activity in autism spectrum disorders. Front Neurosci 9:Article 406.

Ip JPK, Mellios N, Sur M (2018) Rett syndrome: Insights into genetic, molecular and circuit mechanisms. Nat Rev Neurosci 19:368–382.

Kamiya A, Kubo K, Tomoda T, Takaki M, Youn R, Ozeki Y, Sawamura N, Park U, Kudo C, Okawa M, Ross CA, Hatten ME, Nakajima K, Sawa A (2005) A schizophrenia-associated mutation of DISC1 perturbs cerebral cortex development. Nat Cell Biol 7:1167–1178.

Karimi P, Kamali E, Mousavi SM, Karahmadi M (2017) Environmental factors influencing the risk of autism. J Res Med Sci 22:27.

Kazdoba TM, Leach PT, Yang M, Silverman JL, Solomon M, Crawley JN (2016) Translational mouse models of autism: Advancing toward pharmacological therapeutics. Curr Top Behav Neurosci 28:1–52.

Khanal P, Boskovic Z, Lahti L, Ghimire A, Minkeviciene R, Opazo P, Hotulainen P (2023) Gas7 Is a Novel Dendritic Spine Initiation Factor. eNeuro 10:0344–22.

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL (2019) Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol 37:907–915.

King IF, Yandava CN, Mabb AM, Hsiao JS, Huang HS, Pearson BL, Calabrese JM, Starmer J, Parker JS, Magnuson T, Chamberlain SJ, Philpot BD, Zylka MJ (2013) Topoisomerases facilitate transcription of long genes linked to autism. Nature 501:58–62.

Krezel W, Ghyselinck N, Samad TA, Dupé V, Kastner P, Borrelli E, Chambon P (1998) Impaired locomotion and dopamine signaling in retinoid receptor mutant mice. Science 279:863–867.

Lan A, Kalimian M, Amram B, Kofman O (2017) Prenatal chlorpyrifos leads to autism-like deficits in C57Bl6/J mice. Environ Health 16:43.

Landrigan PJ (2010) What causes autism? Exploring the environmental contribution. Curr Opin Pediatr 22:219–225.

Langen M, Bos D, Noordermeer SDS, Nederveen H, van Engeland H, Durston S (2014) Changes in the development of striatum are involved in repetitive behavior in autism. Biol Psychiatry 76:405–411.

Law CW, Chen Y, Shi W, Smyth GK (2014) voom: precision weights unlock linear model analysis tools for RNA-seq read counts. Genome Biol 15:R29.

Lee HS, Bae EJ, Yi SH, Shim JW, Jo AY, Kang JS, Yoon EH, Rhee YH, Park CH, Koh HC, Kim HJ, Choi HS, Han JW, Lee YS, Kim J, Li JY, Brundin P, Lee SH (2010) Foxa2 and Nurr1 synergistically yield A9 nigral dopamine neurons exhibiting improved differentiation, function, and cell survival. Stem Cells 28:501–512.

Lelieveld SH et al. (2016) Meta-analysis of 2,104 trios provides support for 10 new genes for intellectual disability. Nat Neurosci 19:1194–1196.

Lennox AL et al. (2020) Pathogenic DDX3X Mutations Impair RNA Metabolism and Neurogenesis during Fetal Cortical Development. Neuron 106:404-420.e8.

Levy JA, LaFlamme CW, Tsaprailis G, Crynen G, Page DT (2021) Dyrk1a Mutations Cause Undergrowth of Cortical Pyramidal Neurons via Dysregulated Growth Factor Signaling. Biol Psychiatry 90:295–306.

Liao X, Li Y (2020) Genetic associations between voltage-gated calcium channels and autism spectrum disorder: A systematic review. Mol Brain 13:96.

Lin W, Metzakopian E, Mavromatakis YE, Gao N, Balaskas N, Sasaki H, Briscoe J, Whitsett JA, Goulding M, Kaestner KH, Ang SL (2009) Foxa1 and Foxa2 function both upstream of and cooperatively with Lmx1a and Lmx1b in a feedforward loop promoting mesodiencephalic dopaminergic neuron development. Dev Biol 333:386–396.

Lin YC, Frei JA, Kilander MBC, Shen W, Blatt GJ (2016) A subset of autism-associated genes regulate the structural stability of neurons. Front Cell Neurosci 10:Article 263.

Liu D, Nanclares C, Simbriger K, Fang K, Lorsung E, Le N, Amorim IS, Chalkiadaki K, Pathak SS, Li J, Gewirtz JC, Jin VX, Kofuji P, Araque A, Orr HT, Gkogkas CG, Cao R (2022) Autistic-like behavior and cerebellar dysfunction in Bmal1 mutant mice ameliorated by mTORC1 inhibition. Mol Psychiatry:s41380.

Liu Y, Bhowmick T, Liu Y, Gao X, Mertens HDT, Svergun DI, Xiao J, Zhang Y, Wang J huai, Meijers R (2018) Structural Basis for Draxin-Modulated Axon Guidance and Fasciculation by Netrin-1 through DCC. Neuron 97:1261-1267.e4.

Lorsung E, Karthikeyan R, Cao R (2021) Biological Timing and Neurodevelopmental Disorders: A Role for Circadian Dysfunction in Autism Spectrum Disorders. Front Neurosci 15:642745.

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol 15:550.

Macari S, Milgramm A, Reed J, Shic F, Powell KK, Macris D, Chawarska K (2021) Context-Specific Dyadic Attention Vulnerabilities During the First Year in Infants Later Developing Autism Spectrum Disorder. J Am Acad Child Adolesc Psychiatry 60:166–175.

Maenner MJ et al. (2021) Prevalence and Characteristics of Autism Spectrum Disorder Among Children Aged 8 Years — Autism and Developmental Disabilities Monitoring Network, 11 Sites, United States, 2018. MMWR Surveillance Summary 70:1–16.

Magno L, Barry C, Schmidt-Hieber C, Theodotou P, Häusser M, Kessaris N (2017) NKX2-1 Is Required in the Embryonic Septum for Cholinergic System Development, Learning, and Memory. Cell Rep 20:1572–1584.

Mandic-Maravic V, Grujicic R, Milutinovic L, Munjiza-Jovanovic A, Pejovic-Milovancevic M (2022) Dopamine in Autism Spectrum Disorders—Focus on D2/D3 Partial Agonists and Their Possible Use in Treatment. Front Psychiatry 12.

Mapelli L, Soda T, D'Angelo E, Prestori F (2022) The Cerebellar Involvement in Autism Spectrum Disorders: From the Social Brain to Mouse Models. Int J Mol Sci 23:3894.

Marechal D, Brault V, Leon A, Martin D, Pereira PL, Loaëc N, Birling MC, Friocourt G, Blondel M, Herault Y (2019) Cbs overdosage is necessary and sufficient to induce cognitive phenotypes in mouse models of Down syndrome and interacts genetically with Dyrk1a. Hum Mol Genet 28:1561–1577.

Marfella CGA, Imbalzano AN (2007) The Chd family of chromatin remodelers. Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis 618:30–40.

Markenscoff-Papadimitriou E, Binyameen F, Whalen S, Price J, Lim K, Ypsilanti AR, Catta-Preta R, Pai ELL, Mu X, Xu D, Pollard KS, Nord AS, State MW, Rubenstein JL (2021) Autism risk gene POGZ promotes chromatin accessibility and expression of clustered synaptic genes. Cell Rep 37:110089.

Markram K, Rinaldi T, la Mendola D, Sandi C, Markram H (2008) Abnormal fear conditioning and amygdala processing in an animal model of autism. Neuropsychopharmacol 33:901–912.

Martinowich K, Hattori D, Wu H, Fouse S, He F, Hu Y, Fan G, Sun YE (2003) DNA methylation-related chromatin remodeling in activity-dependent BDNF gene regulation. Science 302:890–893.

Matsumura K et al. (2020) Pathogenic POGZ mutation causes impaired cortical development and reversible autism-like phenotypes. Nat Com 11:859.

Matsuo K, Shinoda Y, Abolhassani N, Nakabeppu Y, Fukunaga K (2022) Transcriptome Analysis in Hippocampus of Rats Prenatally Exposed to Valproic Acid and Effects of Intranasal Treatment of Oxytocin. Front Psychiatry 13:Article 859198.

Mohawk JA, Cox KH, Sato M, Yoo S-H, Yanagisawa M, Olson EN, Takahashi JS (2019) Neuronal Myocyte-Specific Enhancer Factor 2D (MEF2D) Is Required for Normal Circadian and Sleep Behavior in Mice. The Journal of Neuroscience 39:7958–7967.

Morcom L, Edwards TJ, Rider E, Jones-Davis D, Lim JWC, Chen KS, Dean RJ, Bunt J, Ye Y, Gobius I, Suárez R, Mandelstam S, Sherr EH, Richards LJ (2021) Draxin regulates interhemispheric fissure remodelling to influence the extent of corpus callosum formation. Elife 10:e61618.

Mut-Arbona P, Huang L, Baranyi M, Tod P, Iring A, Calzaferri F, de Los Ríos C, Sperlágh B (2023) Dual role of the P2X7 receptor in dendritic outgrowth during physiological and pathological brain development. J Neurosci: 43:1125–1142

Nau H (1986a) Species Differences in Pharmacokinetics and Drug Teratogenesis. Environ Health Perspect 70:113–129.

Nau H (1986b) Transfer of valproic acid and its main active unsaturated metabolite to the gestational tissue: Correlation with neural tube defect formation in the mouse. Teratology 33:21–27.

Nau H, Löscher W (1982) Valproic acid: brain and plasma levels of the drug and its metabolites, anticonvulsant effects and gamma-aminobutyric acid (GABA) metabolism in the mouse. J Pharmacol Exp Ther 220:654–659.

Nelson SB, Valakh V (2015) Excitatory/Inhibitory Balance and Circuit Homeostasis in Autism Spectrum Disorders. Neuron 87:684–698.

Nicolini C, Fahnestock M (2018) The valproic acid-induced rodent model of autism. Exp Neurol 299:217–227.

Noda M, Iwamoto I, Tabata H, Yamagata T, Ito H, Nagata K ichi (2019) Role of Per3, a circadian clock gene, in embryonic development of mouse cerebral cortex. Sci Rep 9:5874.

Notwell JH, Heavner WE, Darbandi SF, Katzman S, McKenna WL, Ortiz-Londono CF, Tastad D, Eckler MJ, Rubenstei JLR, McConnell SK, Chen B, Bejerano G (2016) TBR1 regulates autism risk genes in the developing neocortex. Genome Res 26:1013–1022.

Orenstein N, Gofin Y, Shomron N, Ruhrman-Shahar N, Magal N, Hagari O, Azulay N, Bazak L, Goldberg Y, Basel-Salmon L (2022) DYRK1B haploinsufficiency in a family with metabolic syndrome and abnormal cognition. Clin Genet 101:265–266.

Pai ELL, Vogt D, Clemente-Perez A, McKinsey GL, Cho FS, Hu JS, Wimer M, Paul A, Fazel Darbandi S, Pla R, Nowakowski TJ, Goodrich L v., Paz JT, Rubenstein JLR (2019) Mafb and c-Maf Have Prenatal Compensatory and Postnatal Antagonistic Roles in Cortical Interneuron Fate and Function. Cell Rep 26:1157-1173.e5.

Park J, Chung KC (2013) New Perspectives of Dyrk1A Role in Neurogenesis and Neuropathologic Features of Down Syndrome. Exp Neurobiol 22:244–248.

Paulsen B et al. (2022) Autism genes converge on asynchronous development of shared neuron classes. Nature 602:268–273.

Peñagarikano O, Geschwind DH (2012) What does CNTNAP2 reveal about autism spectrum disorder? Trends Mol Med 18:156–163.

Poelmans G, Franke B, Pauls DL, Glennon JC, Buitelaar JK (2013) AKAPs integrate genetic findings for autism spectrum disorders. Transl Psychiatry 3:e270.

Proenca CC, Gao KP, Shmelkov S v., Rafii S, Lee FS (2011) Slitrks as emerging candidate genes involved in neuropsychiatric disorders. Trends Neurosci 34:143–153.

Qian Z, Zhang R, Zhou J, Sun S, Di Y, Ren W, Tian Y (2018) RNA-Seq data on prefrontal cortex in valproic acid model of autism and control rats. Data Brief 18:787–789.

Rachubinski AL, Hepburn S, Elias ER, Gardiner K, Shaikh TH (2017) The co-occurrence of Down syndrome and autism spectrum disorder: is it because of additional genetic variations? Prenat Diagn 37:31–36.

Rash BG, Tomasi S, Lim HD, Suh CY, Vaccarino FM (2013) Cortical gyrification induced by fibroblast growth factor 2 in the mouse brain. Journal of Neuroscience 33:10802–10814.

Rauch A et al. (2012) Range of genetic mutations associated with severe non-syndromic sporadic intellectual disability: An exome sequencing study. The Lancet 380:1674–1682.

Ronesi JA, Collins KA, Hays SA, Tsai NP, Guo W, Birnbaum SG, Hu JH, Worley PF, Gibson JR, Huber KM (2012) Disrupted Homer scaffolds mediate abnormal mGluR5 function in a mouse model of fragile X syndrome. Nat Neurosci 15:431–440.

Rosenberg RE, Law JK, Yenokyan G, Mcgready J, Kaufmann WE, Law PA (2009) Characteristics and Concordance of Autism Spectrum Disorders Among 277 Twin Pairs. Arch Pediatr Adolesc Med 163:907–914.

Ross AJ, Capel B (2005) Signaling at the crossroads of gonad development. Trends Endocrinol and Metab 16:19–25.

Roston A, Evans D, Gill H, McKinnon M, Isidor B, Cogné B, Mwenifumbo J, van Karnebeek C, An J, Jones SJM, Farrer M, Demos M, Connolly M, Gibson WT (2021) SETD1B -associated neurodevelopmental disorder. J Med Genet 58:196–204.

Rubenstein JLR, Merzenich MM (2003) Model of autism: increased ratio of excitation/ inhibition in key neural systems. Genes Brain Behav 2:255–267.

Ruzzo EK, Pérez-Cano L, Jung JY, Wang L kai, Kashef-Haghighi D, Hartl C, Singh C, Xu J, Hoekstra JN, Leventhal O, Leppä VM, Gandal MJ, Paskov K, Stockham N, Polioudakis D, Lowe JK, Prober DA, Geschwind DH, Wall DP (2019) Inherited and De Novo Genetic Risk for Autism Impacts Shared Networks. Cell 178:850-866.e26.

Ryu S, Mahler J, Acampora D, Holzschuh J, Erhardt S, Omodei D, Simeone A, Driever W (2007) Orthopedia Homeodomain Protein Is Essential for Diencephalic Dopaminergic Neuron Development. Current Biology 17:873–880.

Salm EJ, Dunn PJ, Shan L, Yamasaki M, Malewicz NM, Miyazaki T, Park J, Sumioka A, Hamer RRL, He WW, Morimoto-Tomita M, LaMotte RH, Tomita S (2020) TMEM163 Regulates ATP-Gated P2X Receptor and Behavior. Cell Rep 31.

Sandberg M, Flandin P, Silberberg S, Su-Feher L, Price JD, Hu JS, Kim C, Visel A, Nord AS, Rubenstein JLR (2016) Transcriptional Networks Controlled by NKX2-1 in the Development of Forebrain GABAergic Neurons. Neuron 91:1260–1275.

Satterstrom FK et al. (2020) Large-Scale Exome Sequencing Study Implicates Both Developmental and Functional Changes in the Neurobiol of Autism. Cell 180:568-584.e23.

Seigneur E, Wang J, Dai J, Polepalli J, Südhof TC (2021) Cerebellin-2 regulates a serotonergic dorsal raphe circuit that controls compulsive behaviors. Mol Psychiatry 26:7509–7521.

Semino F, Schröter J, Willemsen MH, Bast T, Biskup S, Beck-Woedl S, Brennenstuhl H, Schaaf CP, Kölker S, Hoffmann GF, Haack TB, Syrbe S (2021) Further evidence for *de novo* variants in *SYNCRIP* as the cause of a neurodevelopmental disorder. Hum Mutat 42:1094–1100.

Sessa A, Ciabatti E, Drechsel D, Massimino L, Colasante G, Giannelli S, Satoh T, Akira S, Guillemot F, Broccoli V (2017) The Tbr2 Molecular Network Controls Cortical Neuronal Differentiation Through Complementary Genetic and Epigenetic Pathways. Cerebral Cortex 27:3378–3396.

Shibata M, Pattabiraman K, Lorente-Galdos B, Andrijevic D, Kim SK, Kaur N, Muchnik SK, Xing X, Santpere G, Sousa AMM, Sestan N (2021) Regulation of prefrontal patterning and connectivity by retinoic acid. Nature 598:483–488.

Shmelkov S v, Hormigo A, Jing D, Proenca CC, Bath KG, Milde T, Shmelkov E, Kushner JS, Baljevic M, Dincheva I, Murphy AJ, Valenzuela DM, Gale NW, Yancopoulos GD, Ninan I, Lee FS, Rafii S (2010) Slitrk5 deficiency impairs corticostriatal circuitry and leads to obsessive-compulsive-like behaviors in mice. Nat Med 16:598–602.

Siegenthaler JA, Ashique AM, Zarbalis K, Patterson KP, Hecht JH, Kane MA, Folias AE, Choe Y, May SR, Kume T, Napoli JL, Peterson AS, Pleasure SJ (2009) Retinoic Acid from the Meninges Regulates Cortical Neuron Generation. Cell 139:597–609.

Silbereis JC, Pochareddy S, Zhu Y, Li M, Sestan N (2016) The Cellular and Molecular Landscapes of the Developing Human Central Nervous System. Neuron 89:248–268.

Silverman JL, Yang M, Lord C, Crawley JN (2010) Behavioural phenotyping assays for mouse models of autism. Nat Rev Neurosci 11:490–502.

Singh R, Kumar Dhanyamraju P, Lauth M (2017) Oncotarget 833 www.impactjournals.com/oncotarget DYRK1B blocks canonical and promotes non-canonical Hedgehog signaling through activation of the mTOR/AKT pathway. Oncotarget 8:833–845.

Singh R, Lauth M (2017) Emerging roles of DYRK kinases in embryogenesis and Hedgehog pathway control. J Dev Biol 5:13.

Song M, Mathews CA, Stewart SE, Shmelkov S v., Mezey JG, Rodriguez-Flores JL, Rasmussen SA, Britton JC, Oh YS, Walkup JT, Lee FS, Glatt CE (2017) Rare synaptogenesis-impairing mutations in SLITRK5 are associated with obsessive compulsive disorder. PLoS One 12.

Stahl R, Walcher T, de Juan Romero C, Pilz GA, Cappello S, Irmler M, Sanz-Aquela JM, Beckers J, Blum R, Borrell V, Götz M (2013) Trnp1 regulates expansion and folding of the mammalian cerebral cortex by control of radial glial fate. Cell 153:535–549.

Stanton-Turcotte D, Hsu K, Moore SA, Yamada M, Fawcett JP, Iulianella A (2022) Mllt11 regulates migration and neurite outgrowth of cortical projection neurons during development. J Neurosci 42:3931–3948.

Stessman HAF et al. (2016) Disruption of POGZ Is Associated with Intellectual Disability and Autism Spectrum Disorders. Am J Hum Genet 98:541–552.

Stessman HAF et al. (2017) Targeted sequencing identifies 91 neurodevelopmental-disorder risk genes with autism and developmental-disability biases. Nat Genet 49:515–526.

Stumpo DJ, Bockt CB, Tuttle JS, Blackshear PJ (1995) MARCKS deficiency in mice leads to abnormal brain development and perinatal death (myristoylated, alanine-rich C kinase substrate/protein kinase C/Macs gene). Dev Biol 92:944–948.

Subramanian K, Brandenburg C, Orsati F, Soghomonian JJ, Hussman JP, Blatt GJ (2017) Basal ganglia and autism – a translational perspective. Autism Research 10:1751–1775.

Surén P, Roth C, Bresnahan M, Haugen M, Hornig M, Hirtz D, Lie KK, Lipkin WI, Reichborn-Kjennerud T, Schjølberg S, George M, Smith D, Øyen A-S, Susser E, Stoltenberg C (2013) Association Between Maternal Use of Folic Acid Supplements and Risk of Autism Spectrum Disorders in Children. J Am Med Assoc 309:570–577.

Szu J, Wojcinski A, Jiang P, Kesari S (2021) Impact of the Olig Family on Neurodevelopmental Disorders. Front Neurosci 15:659601.

Tanaka AJ, Cho MT, Willaert R, Retterer K, Zarate YA, Bosanko K, Stefans V, Oishi K, Williamson A, Wilson GN, Basinger A, Barbaro-Dieber T, Ortega L, Sorrentino S, Gabriel MK, Anderson IJ, Sacoto MJG, Schnur RE, Chung WK (2017) De novo variants in EBF3 are associated with hypotonia, developmental delay, intellectual disability, and autism. Cold Spring Harb Mol Case Stud 3:a002097.

Tao W, Díaz-Alonso J, Sheng N, Nicoll RA (2018) Postsynaptic δ1 glutamate receptor assembles and maintains hippocampal synapses via Cbln2 and neurexin. Proc Natl Acad Sci USA 115:E5373–E5381.

Tisato V, Silva JA, Longo G, Gallo I, Singh A V., Milani D, Gemmati D (2021) Genetics and epigenetics of one-carbon metabolism pathway in autism spectrum disorder: A sex-specific brain epigenome? Genes (Basel) 12:782.

Tocco C, Bertacchi M, Studer M (2021) Structural and Functional Aspects of the Neurodevelopmental Gene NR2F1: From Animal Models to Human Pathology. Front Mol Neurosci 14:767965.

Toledo EM, Yang S, Gyllborg D, van Wijk KE, Sinha I, Varas-Godoy M, Grigsby CL, Lönnerberg P, Islam S, Steffensen KR, Linnarsson S, Arenas E (2020) Srebf1 Controls Midbrain Dopaminergic Neurogenesis. Cell Rep 31:107607.

Tripodi M, Filosa A, Armentano M, Studer M (2004) The COUP-TF nuclear receptors regulate cell migration in the mammalian basal forebrain. Development 131:6119–6129.

Tutukova S, Tarabykin V, Hernandez-Miranda LR (2021) The Role of Neurod Genes in Brain Development, Function, and Disease. Front Mol Neurosci 14:662774.

Wang SHJ, Celic I, Choi SY, Riccomagno M, Wang Q, Sun LO, Mitchell SP, Vasioukhin V, Huganir RL, Kolodkin AL (2014) Dlg5 regulates dendritic spine formation and synaptogenesis by controlling subcellular N-cadherin localization. 34:12745–12761.

Weil D, Piton A, Lessel D, Standart N (2020) Mutations in genes encoding regulators of mRNA decapping and translation initiation: Links to intellectual disability. Biochem Soc Trans 48:1199–1211.

Willsey AJ et al. (2013) Coexpression networks implicate human midfetal deep cortical projection neurons in the pathogenesis of autism. Cell 155:997–1007.

Windhorst S, Minge D, Bähring R, Hüser S, Schob C, Blechner C, Lin HY, Mayr GW, Kindler S (2012) Inositol-1,4,5-trisphosphate 3-kinase A regulates dendritic morphology and shapes synaptic Ca 2+ transients. Cell Signal 24:750–757.

Wingett SW, Andrews S (2018) Fastq screen: A tool for multi-genome mapping and quality control. F1000Res 7:1–14.

Wircer E, Blechman J, Borodovsky N, Tsoory M, Nunes AR, Oliveira RF, Levkowitz G (2017) Homeodomain protein Otp affects developmental neuropeptide switching in oxytocin neurons associated with a long-term effect on social behavior. Elife 6:e22170.

Xu D et al. (2013) Top3β is an RNA topoisomerase that works with fragile X syndrome protein to promote synapse formation. Nat Neurosci 16:1238–1247.

Xu XH, Deng CY, Liu Y, He M, Peng J, Wang T, Yuan L, Zheng ZS, Blackshear PJ, Luo ZG (2014) MARCKS regulates membrane targeting of Rab10 vesicles to promote axon development. Cell Res 24:576–594.

Yan CH, Levesque M, Claxton S, Johnson RL, Ang SL (2011) Lmx1a and Lmx1b function cooperatively to regulate proliferation, specification, and differentiation of midbrain dopaminergic progenitors. 31:12413–12425.

Yger M, Girault JA (2011) DARPP-32, jack of all trades...master of which? Front Behav Neurosci 5:Article 56.

Yook C, Kim K, Kim D, Kang H, Kim SG, Kim E, Kim SY (2019) A TBR1-K228E Mutation Induces Tbr1 Upregulation, Altered Cortical Distribution of Interneurons, Increased Inhibitory Synaptic Transmission, and Autistic-Like Behavioral Deficits in Mice. Front Mol Neurosci 12:241.

Yoon S, Piguel NH, Khalatyan N, Dionisio LE, Savas JN, Penzes P (2021) Homer1 promotes dendritic spine growth through ankyrin-G and its loss reshapes the synaptic proteome. Mol Psychiat 26:1775–1789.

Yuen RKC et al. (2017) Whole genome sequencing resource identifies 18 new candidate genes for autism spectrum disorder. Nat Neurosci 20:602–611.

Zaslavsky K et al. (2019) SHANK2 mutations associated with autism spectrum disorder cause hyperconnectivity of human neurons. Nat Neurosci 22:556–564.

Zawadzka A, Cieślik M, Adamczyk A (2021) The role of maternal immune activation in the pathogenesis of autism: A review of the evidence, proposed mechanisms and implications for treatment. Int J Mol Sci 22:11516.

Zeidan J, Fombonne E, Scorah J, Ibrahim A, Durkin MS, Saxena S, Yusuf A, Shih A, Elsabbagh M (2022) Global prevalence of autism: A systematic review update. Autism Res 15:778–790.

Zhang R, Zhou J, Ren J, Sun S, Di Y, Wang H, An X, Zhang K, Zhang J, Qian Z, Shi M, Qiao Y, Ren W, Tian Y (2018) Transcriptional and splicing dysregulation in the prefrontal cortex in valproic acid rat model of autism. Reproductive Toxicology 77:53–61.

Zhao H, Wang Q, Yan T, Zhang Y, Xu H juan, Yu H peng, Tu Z, Guo X, Jiang Y hui, Li X jiang, Zhou H, Zhang YQ (2019) Maternal valproic acid exposure leads to neurogenesis defects and autism-like behaviors in non-human primates. Transl Psychiatry 9:267.

Zheng W, Geng AQ, Li PF, Wang Y, Yuan XB (2012) Robo4 regulates the radial migration of newborn neurons in developing neocortex. Cerebral Cortex 22:2587–2601.

Zigman T, Petković Ramadža D, Šimić G, Barić I (2021) Inborn Errors of Metabolism Associated With Autism Spectrum Disorders: Approaches to Intervention. Front Neurosci 15:673600.