Distinct regions of the kinesin-5 C-terminal tail are essential for mitotic spindle midzone localization and sliding force

Zachary R. Gergely[1,2,3], Michele H. Jones[1,2,3], Bojun Zhou[1],
Cai Cash[1], J. Richard McIntosh[2], and Meredith D. Betterton[1,2]

[1] *Department of Physics, University of Colorado Boulder, Boulder CO 80309*

[2] *Department of Molecular, Cellular, and Developmental Biology,
University of Colorado Boulder, Boulder CO 80309 and*

[3] *Equal contribution*

Kinesin-5 motor proteins play essential roles during mitosis in most organisms. Their tetrameric structure and plus-end-directed motility allow them to bind to and move along antiparallel microtubules, thereby pushing spindle poles apart to assemble a bipolar spindle. Recent work has shown that the C-terminal tail is particularly important to kinesin-5 function: the tail affects motor domain structure, ATP hydrolysis, motility, clustering, and sliding force measured for purified motors, as well as motility, clustering, and spindle assembly in cells. Because previous work has focused on presence or absence of the entire tail, the functionally important regions of the tail remain to be identified. We have therefore characterized a series of kinesin-5/Cut7 tail truncation alleles in fission yeast. Partial truncation causes mitotic defects and temperature-sensitive growth, while further truncation that removes the conserved BimC motif is lethal. We compared the sliding force generated by *cut7* mutants using a kinesin-14 mutant background in which some microtubules detach from the spindle poles and are pushed into the nuclear envelope. These Cut7-driven protrusions decreased as more of the tail was truncated, and the most severe truncations produced no observable protrusions. Our observations suggest that the BimC motif and adjacent C-terminal amino acids are essential for sliding force, while the residues N-terminal to the BimC motif are more important for midzone localization. Spindle microtubule protrusions may be a useful tool to study sliding force generated by kinesin-5 motors.

Kinesin-5 motors are essential for mitosis in many organisms because they help assemble the bipolar mitotic spindle (Fig. 1) [1–5]. Kinesin-5 tetramers are formed by two dimers linked in antiparallel orientation through their stalks, so the motor heads are positioned at both ends of a bipolar complex [6–11]. When kinesin-5s crosslink antiparallel microtubules (MT), their plus-end-directed motility generates force that slides the microtubules apart [12–18]. This sliding force is important for initial separation of mitotic spindle poles, assembly of a bipolar spindle (Fig. 1A), and for spindle elongation in early anaphase B [19–21]. Loss of kinesin-5 function commonly leads to the formation of monopolar mitotic spindles [2, 4, 22], consistent with the role of kinesin-5 sliding force in separating spindle poles.

However, increasing evidence suggests that this model for how kinesin-5s separate spindle poles is incomplete. Kinesin-5s are not only plus-end-directed, but can also walk or be carried toward spindle MT minus ends. Fungal kinesin-5s have intrinsic bidirectional motility, [23–34], while in vertebrates kinesin-5 becomes cargo for a minus-end directed motor [35–39]. At least in part because of this minus-end-directed motility, kinesin-5s localize more strongly near spindle poles (near MT minus ends) than in the center of the spindle, where MT plus ends predominate [4, 5, 38, 40–42]. While the functional significance of kinesin-5 minus-end-directed motility and pole localization have remained unclear, previously work suggested that pole localization may position kinesin-5s for spindle elongation [32, 43]. More recently, we proposed that these mechanisms may sequester kinesin-5 near the poles to limit sliding force [34]. In this light, the mechanisms by which kinesin-5 directionality and localization are controlled may be important to its regulation.

Kinesin-5 monomers have a conserved organization with an N-terminal motor domain, a stalk containing oligomerization domains, and a C-terminal tail (Fig. 1B). While the greatest amino acid conservation is in the motor domain, the tail has conserved regions known as the BimC box and NIMA-family kinase box [5, 41, 44–46]. These include consensus sites for cyclin-dependent kinase 1 (Cdk) and NIMA-family kinase (NFK, Fig. 1B). Because kinesin-5s are antiparallel tetramers, the C-terminal tail domain of one dimer lies near the motor domain of the other dimer (Fig. 1A). This geometry has led to a recent focus on the possible role of tail-motor interactions in kinesin-5 motility. It has been observed that cancer cells can develop resistance to kinesin-5/Eg5 inhibitors via tail truncation[47], further supporting the significance of the tail.

The kinesin-5 tail regulates motor function and localization. For metazoan kinesin-5s, truncation of the C-terminal tail alters motility, MT crosslinking, and sliding force of purified motor [18, 48, 49]. However, because the tail domain is also critical for spindle localization [18], the importance of the tail for kinesin-5 force in the spindle cannot be assessed for these organisms. Closed mitosis in fungi appears to have advantages in confining kinesin-5 tail truncations in a smaller volume where they can maintain spindle localization. Deletion of the kinesin-5/CIN8 tail in budding yeast biases directionality toward MT plus ends [50] and is lethal if the gene for a second kinesin-5 allele (KIP1) is also deleted [14, 50]. In fission yeast, truncation of the kinesin-5/Cut7 tail is lethal and affects bipolar spindle assembly [34]. Notably however, *cut7* becomes inessential in fission-yeast lacking kinesin-14 [51–53]. We used this genetic background to show that tail truncation alters Cut7p velocity and rate of direction switching [34]. While the C-terminal tail is predicted to be disordered based on its primary structure [34], cryo-EM analysis of *Drosophila* KLP61F found that interactions between the tail and motor domains can at least partially fold the tail [18]. Tail-motor interactions also alter motor structure, causing slower ATP hydrolysis and speed of purified motor [18]. Further, it has recently been proposed that C-terminal tails of kinesin-5 can interact with motor domains of adjacent motors, promoting cluster formation that enhances sliding force [49] and could affect direction switching [30, 33]. However, the extent and mechanisms of kinesin-5 regulation by the C-terminal tail remains incompletely understood.

The fission yeast *Schizosaccharomyces pombe* is a useful model in which to study kinesin-5. Cut7p is the sole kinesin-5 motor in this organism, and it shares domain structure and spindle localization with other kinesin-5s (Fig. 1). The Cut7p motor-domain structure has been solved [56]. In fission yeast lacking kinesin-14/Pkl1 or its binding partners, Wdr8p and Msd1p, spindle MT minus ends can be pushed past the spindle-pole body (SPB) and distort the nuclear envelope outward [51–53]. These spindle MT minus-end protrusions can differ in length, dynamics and in the polarity of the MTs they contain [53]. Importantly, spindle MT minus-end protrusions depend on Cut7p, since they do not form in cells in which Cut7 is deleted or has a loss of function [52, 53]. Protrusions can therefore assay force generated by Cut7p.

RESULTS

Sequential truncation of the *cut7* C-terminal tail leads to growth and spindle defects

In previous work, we found that the *cut7* tail truncation lacking the C-terminal 96 amino acids (*cut7-988*) was lethal in an otherwise wild-type genetic background [34]. To study which portions of the *cut7* tail are most important for cell viability, spindle assembly, and and sliding force, we generated a series of sequential C-terminal truncation mutants as well as an internal deletion (Fig. 2A). Each *cut7* allele was tagged with green fluorescent protein (GFP) at its C-terminus and was the sole copy of *cut7* in the cell. In addition, the cells contained low-level-expressed mCherry-tubulin to label spindle MTs [34, 57, 58].

FIG. 1. Overview of kinesin-5 mitotic function and C-terminal tail. (A) Schematic of Cut7p and its role in bipolar spindle assembly. Left: Cut7p tetramers near the fission-yeast spindle-pole bodies (SPBs) of a monopolar spindle. Inset: Cut7p tetramer showing motor, stalk, and tail domains. Right: Cut7p tetramers on a bipolar spindle. Inset: Cut7p crosslinking antiparallel microtubules in the spindle midzone. (B) Upper: domain map of *cut7* showing the motor, stalk, and tail. Middle and lower: ClustalW sequence alignment of kinesin-5 C-terminal tails near the BimC box (middle) and NIMA-family kinase box (lower), adapted from Ref. [46]. Alignment includes *Schizosaccharomyces pombe cut7*, *Homo sapiens* Eg5/KIF11, *Mus musculus* Eg5, *Xenopus laevis* Eg5, *Drosophila melanogaster* KLP61F, *Aspergillus nidulans* BimC, *Caenorhabditis elegans* BMK-1, and *Saccharomyces cerevisiae* KIP1 and CIN8. Conserved residues are highlighted in purple, *S. pombe* temperature-sensitive lethal mutations are highlighted in cyan, and the Cdk consensus motif is boxed [5, 42, 44–46, 54, 55].

While truncation of small portions of the tail did not significantly affect cell growth, greater truncation caused temperature-sensitive growth defects and lethality. The *cut7-1032* truncation removed the final 53 amino acids from the tail, including the NIMA-family kinase box. Growth was similar for *cut7-1032* and full-length *cut7* cells (*cut7-FL* Fig. 2B). Cells with *cut7-1017* lacking the final 68 amino acids grew similarly to *cut7-FL* and *cut7-1032* at 25ºC, but showed a strong growth defect at 37ºC. The *cut7-1006* allele lacked 79 amino acids, including the conserved BimC motif (Fig. 2A). We were unable to obtain *cut7-1006* cells by transformation as the sole *cut7* allele in otherwise wild-type cells, suggesting it is lethal. The *cut7-988* mutant lacked the entire tail up to the predicted coiled-coil sequence, and was a lethal allele [34]. Finally we made an internal deletion of a.a. 989-1028, which lacked the BimC motif, but retained a C-terminal portion of the BimC box and the NIMA-family kinase box (Fig. 2A) [46]. Cells with *cut7-989-1028∆* showed a slight growth defect at 25ºC and were inviable at 37ºC (Fig. 2B). These results suggest that the BimC box and adjacent residues are particularly important for cell growth.

The viable truncation mutants all showed bipolar spindle assembly at 25ºC with spindle-localized Cut7-GFP, but differed in spindle length and elongation (Figs. 2C-E, S1). We imaged cells by confocal fluorescence microscopy, then quantified spindle length over time and aligned

traces by defining $t = 0$ as the time of anaphase onset (Methods, Fig. S1). We measured pre-anaphase spindle length, anaphase spindle elongation speed, and the time from anaphase onset until steady spindle elongation was reached (Fig. 2E-H, Methods). The *cut7-FL* and *cut7-1032* spindles appeared similar, with pre-anaphase spindle length of $\sim$2.5 μm, anaphase spindle elongation speed of $\sim$ 0.8 μm/min, and a $\sim$2-3 min time after anaphase onset to reach steady elongation (Fig. 2E-H). Cells containing *cut7-1017* and *cut7-989-1028Δ* had shorter pre-anaphase spindles (2 and 1.25 μm, Fig. 2F). While the steady anaphase elongation speed was similar in all strains (Fig. 2G), *cut7-1017* and *cut7-989-1028Δ* cells took longer to reach steady elongation (Fig. 2H). This suggests that Cut7p with the tail truncated at a.a. 1017 or with the internal deletion might have defects in sliding force needed to elongate the spindle.

Spindle imaging at restrictive temperature showed that temperature-sensitive growth defects correlated with spindle defects. All Cut7-GFP mutants localized to the spindle at 37°C (Fig. 2I,J). All *cut7-FL* and *cut7-1032* cells imaged assembled a bipolar spindle (Fig. 2K). By contrast, *cut7-1017* cells, which showed growth defects at 37°C, also showed spindle assembly delays in 10% of cells (Fig. 2K), suggesting that spindle defects might cause their temperature-sensitive growth problems. For *cut7-989-1028Δ* cells that were inviable at restrictive temperature, all cells observed showed spindle assembly failure with monopolar spindles (Fig. 2J,K). Of the viable mutants we studied, *cut7-989-1028Δ* cells with the BimC motif removed had the most severe phenotype.

The BimC motif is required for proper Cut7p spindle midzone localization

Previous work showed that *cut7-988* lacking the entire tail showed changes in Cut7p localization and motility [34]. Therefore, we asked whether such localization or motility changes might explain *cut7* truncation mutant phenotypes at permissive temperature. We used automated image analysis (Methods, [34, 59]) to create kymographs of Cut7-GFP intensity along the spindle axis, determine average intensity profiles, and compute both total Cut7-GFP spindle fluorescence and the ratio of intensity at the midzone to that at the poles (Fig. 3, Methods). Because Cut7-GFP shows enhanced localization at spindle poles [34, 40, 51, 60] (Fig. 3A), we grouped profiles of Cut7-GFP fluorescence by peak-to-peak distance and averaged (Fig. 3B-D). This allowed us to measure how Cut7-GFP fluorescence redistributes as spindle length changes during mitosis. Cut7-FL-GFP average fluorescence intensity suggested that as the spindle increased in length from 2 to 4 μm, the midzone intensity increased slightly (Fig. 3B,C, black lines). Once spindle length reached 7 μm, the midzone intensity decreased and pole intensity became asymmetric (Fig. 3D, Methods). This quantitative analysis was consistent with previous [34, 40, 51, 60] and current (Fig. 3A) qualitative results.

Kymographs of Cut7-GFP movement on the spindle appeared similar for all the viable truncation mutants (Fig. 3A,E-H). Therefore, at permissive temperature truncation mutants of *cut7* localize to and move on spindle MTs. This suggests that the spindle length and elongation defects we observed do not occur due to complete loss of Cut7p motility. However, upon inspection Cut7-989-1028Δ-GFP kymographs appeared to have lower overall intensity and lower relative midzone intensity. Therefore, we quantified subtle changes in localization for different Cut7-GFP alleles, by averaging intensity profiles over a range of spindle length (Fig. 3I-L, Methods). While the average intensity profiles appeared similar for the full-length motor and truncation mutants, Cut7-989-1028Δ-GFP showed lower intensity, both at the spindle midzone and at poles. To quantify fluorescence, we computed the total Cut7-GFP intensity on the spindle for varying spindle length (Fig. 3M). Cut7-989-1028Δ-GFP indeed showed lower total intensity than the other strains. Lower Cut7p expression, decreased stability, or a spindle localization defect in these cells could lead to lower amounts of motor on the spindle, which in turn could explain the shorter spindles and longer

time before steady anaphase elongation in this mutant (Fig. 2E-H).

Removal of the BimC motif also altered the balance of Cut7p at the spindle midzone compared to the poles. Recently, we proposed that spindle poles may serve as a "waiting room" where kinesin-5s are sequestered to limit sliding force in the midzone [34]. Since our observations of altered spindle length and steady anaphase elongation delays suggested alterations in sliding force in some mutants, we asked whether this might be caused by shifts in Cut7p distribution along the spindle. We therefore quantified the ratio of midzone-to-pole Cut7-GFP fluorescence (Methods). Because fission yeast contain overlapping antiparallel MTs throughout the interpolar spindle [61–63], we defined the spindle midzone as all spindle MTs between the poles. The average midzone-to-pole intensity ratio was low (<1) for short spindles, then increased (to ∼1.5) in early anaphase B, and finally decreased in late mitosis (Fig. 3N). However, Cut7-989-1028Δ-GFP showed the increased ratio for shorter spindles than the other strains, while the overall ratio was lower (<1) for all spindle lengths observed. We additionally analyzed the midzone-to-pole ratio as a function of time relative to anaphase onset, and found that the time of increase was similar in all strains (Fig. 3O). These results showed that total Cut7-989-1028Δ-GFP fluorescence was lower and the relative amount of midzone-localized motor was lower in anaphase. Both of these changes could contribute to reduced sliding force and spindle length and elongation defects in these cells.

Deletion of kinesin-14 motor genes rescues lethality of Cut7 tail truncation

Deletion of kinesin-14/Pkl1 suppresses the deleterious effects of cut7 loss-of-function alleles [64–67]. Since deletion of *pkl1* allowed study of the otherwise lethal *cut7-988* truncation alleles [34], we introduced *cut7* and our four truncation alleles (Fig. 2A) into the *pkl1Δ klp2Δ* background to compare their phenotypes. All of our strains were viable (Fig. 4A). While no growth defects occurred at 25°C for *cut7-1032* and *cut7-1017* (Fig. 4A, top), removal of the BimC motif in *cut7-1006* severely retarded growth. We were surprised that removal of the entire tail in *cut7-988* showed a lesser growth defect; however, it may act more like a *cut7Δ* allele which has been shown to grow relatively better than certain *cut7* mutant alleles in this background [65, 66]. At 37°C, *cut7-1032* and *cut7-1017* grew similarly to full length, whereas *cut7-1006* was lethal and *cut7-988* had a moderate growth defect (Fig. 4A, bottom). At 37°C , the truncations that grew well showed nearly all bipolar spindles in live-cell imaging, while *cut7-1006* and *cut7-988* had monopolar spindles (Fig. S2). The rescue of these previously lethal alleles allowed us to further assess the potential effects of the amino acids N-terminal to the BimC motif (Fig. 1B) on Cut7p localization and force.

Dynamic spindle MT minus-end protrusions serve as a read-out of sliding force produced by Cut7 mutants

Cut7p force generated at the midzone of the spindle is required for protrusion of spindle MT minus ends in fission yeast (Fig. 4B) lacking kinesin-14/Pkl1 (or members of its complex, Wdr8p and Msd1p) [51–53, 60]. This suggested that protrusions might vary for Cut7p truncation mutants that caused changes in spindle length. Therefore, we imaged protrusions for our truncation mutants in the *pkl1Δ klp2Δ* background.

We found that Cut7p produced protrusions and accumulated at protrusion tips. We observed protrusions at 25°C in cells with fluorescently labeled MTs and Cut7p, as seen previously [51, 52]. In cells with protrusions, Cut7-GFP localized both to the SPBs and at the tip of the protruding MTs (Fig. 4B, C). Time-lapse imaging allowed us to observe protrusion dynamics. Some protrusions grew, shortened, and then grew again in the same cell (Fig. 4D), while other cells showed protrusions

from both SPBs simultaneously (Fig. 4E). Because we used low-level tubulin labeling [57, 68] and protrusions contain only a small number of MTs [51], protrusion MTs were often too dim to be be visible in every image (Fig. 4D-H). Despite this, Cut7-GFP at protrusion tips was consistently bright enough to identify protrusions and their dynamics (Fig. 4C-H). Because protruding MTs typically have their minus-ends distal [51, 53], Cut7-GFP accumulation at protrusion tips appeared to arise from Cut7p minus-end-directed motility (Fig. 4B, lower). We noted that the protrusion tip often showed brighter GFP signal than the adjacent SPB (Fig. 4D-F). Because protrusions depend on Cut7p sliding force, this observation suggests that Cut7p is plus-end-directed on midzone MTs while simultaneously minus-end-directed on protruding MTs.

Cut7p tail truncations showed reduced frequency of protrusions, even for alleles with mild growth and spindle phenotypes. In the *pkl1+ klp2+* background, *cut7-1032* and *cut7-1017* alleles with shorter tail truncation showed growth, spindle length, and midzone localization similar to *cut7-FL* (Fig. 2, 3). These observations suggested that the mild truncations could generate force similarly to full-length protein. However, when we measured the fraction of cells with visible protrusions, we found a decrease in all truncation mutants. Cells containing *cut7-FL* showed protrusions in 78% of cells, but this dropped to 65% for *cut7-1032* and 21% for *cut7-1017* (Fig. 4F,G,I). The removal of the residues C-terminal to the BimC motif in *cut7-1017* therefore appeared to cause a sharp decrease in Cut7p sliding force. When the BimC motif was completely removed in *cut7-1006* and *cut7-988*, we did not observe any protrusions (Fig. 4H,I). Thus, Cut7p sliding force (as reflected in protrusion frequency) was correlated with the amount of C-terminal tail present.

Different portions of the Cut7p tail affect sliding force and midzone localization

We asked whether the changes in protrusion frequency in *cut7* trail truncations might be correlated with changes in spindle length or Cut7p localization. Therefore, we analyzed spindle length dynamics and Cut7-GFP intensity distribution for truncation mutants in the *pkl1Δ klp2Δ* background (Methods). The spindle sometimes shortened when protrusions formed, suggesting that formation of protrusions altered spindle force balance. Therefore, when measuring spindle length dynamics, we only analyzed cells without visible protrusions. Among these cells, the pre-anaphase spindle length was shorter in *cut7-1006* and *cut7-988* cells lacking the BimC motif (Figs. 5A, S3). Since these strains also did not show protrusions in cells we imaged (Fig. 4I), this result suggests that both shorter spindle length and loss of protrusions reflect decreased Cut7p sliding force.

We then asked whether this decreased force might be caused by lower overall Cut7p levels or a reduced amount of motor at the spindle midzone. Here we quantified all cells, with and without protrusions. Because protrusions complicated automated identification of spindle poles, we used manual tracking of SPB position in cells with visible protrusions (Methods). The total Cut7-GFP signal on the entire spindle was similar for all alleles, except for Cut7-988-GFP which showed higher total intensity (Fig. S4A). The Cut7-GFP midzone-to-pole ratio was similar for most alleles, with the exception of Cut7-988-GFP which showed a consistently lower ratio than the other strains (Fig. 5B, S4B-F). Therefore, Cut7-988-GFP showed enhanced pole localization, in agreement with previous observations [34].

Surprisingly, Cut7-GFP midzone localization appeared to be retained even in *cut7-1006* and *cut7-1017* cells in which protrusions occurred less often. To compare protrusions and localization, we averaged the Cut7-GFP midzone-to-pole ratio (Fig. 5B) for all spindles with length from 3 to 7 μm. This gave a single average midzone-to-pole ratio that we compared to the protrusion frequency as a function of truncation position (Fig. 5C). Neither protrusion frequency nor midzone-to-pole ratio changed linearly with the length of C-terminal tail retained; instead, both quantities changed nonlinearly (compare red and blue curves to black linear model in Fig. 5C). Relative midzone

localization was retained for all truncations except that at a.a. 988, and remained fairly high even for the truncation at a.a. 1006 that removed the BimC motif. We noted that the midzone-to-pole ratio was similar for *cut7-1006* and *cut7-FL* cells in the *pkl1Δ klp2Δ* background, and slightly higher for *cut7-1017* and *cut7-1032* cells for unknown reasons. On the other hand, the BimC motif was required for protrusions to occur, and protrusion frequency increased sharply as the truncation site increased from a.a. 1006 to 1032. Therefore, our results show that significant protrusion frequency requires both the BimC motif and adjacent C-terminal amino acids, a region that includes nine previously identified phosphorylated residues [69–72]. These results suggest that the essential tail region for force production is the a.a. 1006-1032 (the BimC motif and adjacent C-terminal residues), while for anaphase midzone localization a.a. 988-1006 (N-terminal to the BimC motif) are most important.

DISCUSSION

Our results show that the fission-yeast kinesin-5/Cut7 C-terminal tail contributes to spindle midzone localization and sliding force for proper spindle assembly and elongation. Sequential truncation of the tail caused growth defects, temperature-sensitive lethality due to a failure of spindle assembly, and lethality as more of the tail was truncated. The lethality of the *cut7* tail truncation is consistent with studies of kinesin-5/CIN8 in budding yeast [50]. In viable mutants, we observed that pre-anaphase spindle length decreased and the time to reach steady anaphase elongation increased, suggesting that partial tail truncation limits Cut7p sliding force, despite spindle localization. We built on the observation that fission yeast lacking kinesin-14/Pkl1 form spindle MT minus-end protrusions that distort the nuclear envelope, a phenotype dependent on *cut7* [51–53]. We found that as the C-terminal tail was truncated, the frequency of visible protrusions decreased. Since protrusions require Cut7p sliding force, fewer protrusions provide evidence for defects in sliding force in tail truncation mutants.

These results extend recent work on homologous kinesin-5 motors in reconstituted systems. Tail truncation alters ATPase activity, motility, and force of *Drosophila* and human kinesin-5s [18], showing the tail plays a role in tuning kinesin-5 force. C-terminal tails may bridge to adjacent motors, promoting cluster formation that could affect sliding force [49] via motor domain structural change upon tail binding [18]. A computational model of budding yeast CIN8 proposed an attractive interaction that promotes motor clustering [33], which could be mediated by tail-motor interactions. Optical trap force measurements of human Eg5 antiparallel MT sliding showed that tail truncation decreases sliding force by a factor of 8 [18]. We found that full *cut7* tail truncation causes a loss of protrusions, consistent with a loss of sliding force. Protrusions are not observed in *pkl1Δ* cells lacking *cut7* [53] or carrying a temperature-sensitive *cut7* loss of function [52], making our tail truncation results consistent with loss of all sliding force. However, because some threshold level of force is likely required to deform the nuclear envelope [63, 67, 73], we cannot conclude that Cut7p alleles that form no protrusions produce no sliding force.

Our data suggest that midzone localization and sliding force are affected by different portions of the Cut7p tail. As has been seen with other *cut7* tail mutants [41], all of our truncated Cut7p alleles localized to spindle MTs/SPBs, despite the loss of the tail-MT binding site [74]. Even with overall spindle binding unaltered, decreases in motor midzone localization could alter sliding force [34]. However, we found that relative midzone motor fluorescence remained relatively high for all mutants except the *cut7-988* full tail truncation. The localization, spindle length, and protrusion measurements together point to amino acids N-terminal to the BimC motif as most important for midzone localization (a.a. 988-1006), while the BimC motif and adjacent C-terminal amino acids are essential for sliding force (a.a. 1006-1032). The BimC motif (a.a. 1008-1015)

contains a conserved Cdk phosphorylation site [5, 41, 42, 44, 45, 55]. When the Cdk-site threonine is mutated in vertebrate kinesin-5s, motor-spindle MT binding is reduced [5, 42, 45]. While for Cut7p spindle localization is retained with loss of the BimC motif, sliding force is reduced. Nine sites phosphorylated in mitosis have been identified in the *cut7* tail between amino acids 1007 and 1028 [69–72]. This corresponds remarkably well with the region most important for protrusions. We therefore speculate that phosphoregulation at these sites contributes to Cut7p sliding force.

While we did not see significant phenotypes of truncating amino acids 1033-1085 C-terminal to the BimC box region, these residues are still significant in the context of the internal deletion. The *cut7-988* and *cut7-989-1028*Δ alleles both lack the BimC motif, but only the full tail truncation shows a lethal phenotype. This suggests that even with this important region removed, retaining the other 57 tail amino acids is beneficial. Similarly, the internal deletion shows higher midzone-to-pole ratio compared to the full tail truncation, suggesting that the presence of the entire tail contributes to Cut7p midzone localization.

Tail truncation affected pre-anaphase spindle length and protrusion frequency, consistent with alterations in Cut7p force early in mitosis. While the truncation mutants did cause delays in reaching steady anaphase elongation, the late-anaphase elongation speed was similar for the different mutants. This is consistent with previous work showing that kinesin-6/Klp9 is the primary spindle elongation motor in fission-yeast anaphase B [75, 76].

Our results give further evidence of Cut7p bidirectional motility in cells. Previous work found that Cut7p moves toward plus and minus ends of MTs both when purified [28–30] and on the spindle [34]. In reconstituted antiparallel MT overlaps, crowding on the MT lattice shifted Cut7p to plus-end-directed movement [30]. However, we observed most Cut7p localized near the SPBs for all of our mutants, consistent with minus-end-directed movement that concentrates the motor near spindle poles. This appears to be at odds with the idea that crowding regulates Cut7p plus-end-directed motility on spindle MTs. However, the limited resolution of our images does not rule out the possibility that motor crowding near the SPB may cause plus-end-directed movement where Cut7p concentration is highest.

Our results also show that Cut7p can have varying directional bias on different parts of the spindle. Formation of protrusions depends on Cut7p sliding force, due to plus-end-directed motility on antiparallel MTs in the spindle midzone. At the same time, Cut7p minus-end-directed movement causes the motor to accumulate at the ends of protrusion MTs. (We note that the high Cut7-GFP signal observed at protrusion tips is evidence that it accumulates by minus-end-directed movement: if the motor localized solely based on binding to minus ends, one would expect the tip fluorescence to be proportional to MT number. Instead, Cut7-GFP fluorescence is bright, often exceeding that of the SPB-localized signal.) Therefore, subsets of Cut7p move toward both plus and minus ends of MTs on a bipolar spindle simultaneously during mitosis. Cut7p therefore behaves similarly to budding yeast CIN8, for which simultaneous plus- and minus-end-directed movement was observed on anaphase spindles [24]. The directional bias may be affected by MT geometry: protrusions contain single or small bundles of parallel MTs [51], which may contribute to minus-end-directed Cut7p motility. Antiparallel MTs in the spindle midzone appear to favor plus-end-directed motility that drives sliding force. Therefore, our results are consistent with previous work showing that kinesin-5 crosslinking contributes to its plus-end bias [24, 77].

METHODS

Reagents and techniques

Oligonucleotide primers were purchased from Integrated DNA Technologies (Coralville, IA). Restriction enzymes and Phusion HF DNA polymerase were purchased from New England Biolabs (Ipswich, MA). DNA was prepared using Qiaprep Spin Miniprep Kit and polymerase chain reaction (PCR) products were purified using Qiaquick PCR Purification Kit, both from Qiagen (Germantown, MD). Fission yeast genomic DNA was prepared using YeaStar Genomic DNA Kit from Zymo Research (Irvine, CA). DNA sequencing was performed by Quintarabio (Hayward, CA). DNA concentration was determined using a Thermo Scientific Nanodrop 2000. Strains were verified by PCR and sequence analysis of genomic DNA. At least two different transformant strains were analyzed in experiments, and cells were cultured using standard techniques [78]. Growth phenotypes were analyzed after 3-4 days on rich media plates with 5-fold serial dilution starting with 5 microliters of culture at 0.1 optical density.

Strain construction for truncation mutants

The plasmid pFA6a-GFP(S65T)-kanmx6 (Addgene 39292) was amplified by PCR using oligonucleotide primers Cut7-Cterm-pFA6a-GFP-F and Cut7-Cterm-pFA6a-GFP-R. The truncation mutants were made using the same reverse primer as above and the following forward primers: for 1-1032: Cut7-1-1032-GFP-F, for 1-1017: Cut7-1-1017-GFP-F for 1-1006: Cut7-1-1007-GFP-F, for 1-988: Cut7-1-988-GFP-F. There is a six amino acid linker (RIPGLI) between Cut7 and GFP. The PCR products were used for transformation using a lithium acetate method [79] and G418 resistant colonies were characterized. Transformation of yMB1084 (containing *pkl1+ klp2+*) yielded strains yMB1149 (*cut7*), yMB1125 (*cut7-1032*, and yMB1142 (*cut7-1017*) and transformation of strain yMB1147 (containing *pkl1Δ klp2Δ*) yielded strains yMB1162 (*cut7*), yMB1210 (*cut7-1032*), yMB1207 (*cut7-1017*), yMB1238 (*cut7-1006*) and yMB1155 (*cut7-988*).

Strain construction for internal deletion mutant

Plasmid pMB64 was made in a Bluescript backbone (kind gift from Wayne Wahls) and contains the *cut7* coding sequence from Bgl2 (b.p. 1835) to the C-terminus (b.p. 3255) and downstream sequence through the *his1* gene. Synthetic *cut7* alleles (Genscript) were inserted into pMB64, from Bcl1 (b.p. 2356) to BsrG1 (b.p. 3361) containing either wild-type sequence with four silent restriction sites (pMB71-FL) or a *cut7* fragment containing an internal deletion, lacking amino acids 989-1028. These were cloned into a related plasmid containing a C-terminal GFP (pMB44-FL and pMB45-intdel respectively) and a fragment from these plasmids was transformed into yeast strain MB1091 (in which the downstream his1 gene had been replaced with ura4) and *his+ura-* colonies were characterized. This yielded strains yMB1199 (*cut7*) and yMB1168 (*cut7-989-1028Δ*). Yeast with a *cut7* allele containing silent restriction sites (yMB1199) grew the same and behaved similarly in all assays as those with an allele lacking silent restriction sites (yMB1149). Therefore yMB1149 was used as the FL control for experiments in kinesin-14+ background.

Live-cell imaging

All microscopy used live-cell preparation on a Nikon Eclipse Ti (Nikon USA, Melville, NY) spinning-disk confocal microscope as previously described [34, 63]. Microscope temperature was maintained with ± 0.1°C precision using a CherryTemp controller (Cherry Biotech, Rennes, France), and samples were placed on *Bandeiraea simplicifolia* lectin (Sigma-Aldrich, St. Louis, MO) coated coverslips pre-equilibrated to the appropriate temperature of either 25 or 37°C as described previously [34].

Kymographs

For each truncation mutant, we selected the mitotic cells that formed a bipolar spindle, based on the bright spots of Cut7-GFP localized at both spindle poles. We manually segmented an elliptical region to include our cell of interest; any region outside this segmentation was not considered. The position of both Cut7-GFP spots was tracked to sub-pixel accuracy for each mitotic frame. We constructed kymographs by interpolating the Cut7-GFP intensity along the line connecting the spots in each frame and subtracting the background intensity. The brighter spot was used as a fixed reference point plotted on the left side of the kymograph, while the other spot varied in position. In some cases the line along the spindle axis extends beyond the boundary of the cell, into the region that has been cropped. In this case the kymograph includes pixels that have been cropped and therefore have zero intensity, which are visible in the kymographs as completely black areas.

Cut7-GFP intensity analysis

To group profiles of Cut7-GFP intensity along the spindle, we identified the Cut7-GFP peaks near the SPBs and measured the distance between the peaks. We then extracted lines from the kymographs with the relevant peak separation. We included images with peaks separated by ± 0.1 μm around the desired peak separation. In average curves, the right peak typically showed lower intensity than the left peak. This occurred because we manually selected the brighter Cut7-GFP spot to be at $x = 0$. We computed total Cut7-GFP intensity by summing the intensity from each spindle intensity profile extracted from the kymographs.

We determined the Cut7-GFP midzone-to-pole ratio by first defining the pole and midzone. We summed pole intensity from a square box 0.75 μm on a side centered on the pole intensity peak. The midzone intensity is summed in a rectangular box of width 0.75 μm with long axis centered on the spindle axis that runs between the two poles, including all of the spindle except the pole boxes. The total Cut7-GFP intensity is the sum of the pole and midzone signals. The midzone-to-pole ratio is the total midzone Cut7-GFP intensity, divided by total pole intensity, computed for each frame.

Spindle length dynamics

Microscope images were processed using TAMiT [59], which identifies spindle endpoints in each frame. The length of the spindle is determined from these points and smoothed. First, we divide each trace into 30 time bins and analyzed the standard deviation of spindle length in each bin. Bins with high standard deviation (corresponding to fluctuations on the order of 0.75 μm) were

flagged as noise and discarded. The data were then smoothed with a rolling average with window of width ~3.5 min.

Figure 3, F-K: For each cut7-GFP mutant, we plotted the average cut7-GFP intensity distribution at each integer spindle length from 1 to 10 μm.

We manually identified anaphase onset for each curve by choosing the point where the spindle begins to grow beyond fluctuations around its pre-anaphase length. To produce average spindle length traces from different cells, each curve was aligned by its anaphase onset time and divided into evenly spaced bins of width 15 s. All points from each curve that fall within a given bin were then averaged. Average curves for each phenotype included at least 4 curves.

To determine the anaphase elongation rate, we first manually identified the time range of constant growth speed, then performed a linear fit on this region. The fit elongation speed was averaged from multiple cells. The average time of onset of steady anaphase elongation was determined by averaging the identified time from all cells. Uncertainty is computed and displayed as the standard error of the mean.

ACKNOWLEDGMENTS

We thank Daniel Steckhahn for helpful discussions and figure formatting, the University of Colorado Boulder Light Microscopy and Biocore Facilities, Pombase and the Wellcome Trust, and the National Bio-Resource Project Yeast, Japan for providing original strains. This work was funded by NIH grant R01GM124371.

[1] A. P. Enos and N. R. Morris, Mutation of a gene that encodes a kinesin-like protein blocks nuclear division in A. nidulans, Cell **60**, 1019 (1990).

[2] I. Hagan and M. Yanagida, Novel potential mitotic motor protein encoded by the fission yeast cut7+ gene, Nature **347**, 563 (1990).

[3] M. A. Hoyt, L. He, K. K. Loo, and W. S. Saunders, Two Saccharomyces cerevisiae kinesin-related gene products required for mitotic spindle assembly., The Journal of Cell Biology **118**, 109 (1992).

[4] K. E. Sawin, K. LeGuellec, M. Philippe, and T. J. Mitchison, Mitotic spindle organization by a plus-end-directed microtubule motor, Nature **359**, 540 (1992).

[5] A. Blangy, H. A. Lane, P. d'Hérin, M. Harper, M. Kress, and E. A. Nigg, Phosphorylation by p34cdc2 regulates spindle association of human Eg5, a kinesin-related motor essential for bipolar spindle formation in vivo, Cell **83**, 1159 (1995).

[6] D. G. Cole, W. M. Saxton, K. B. Sheehan, and J. M. Scholey, A "slow" homotetrameric kinesin-related motor protein purified from Drosophila embryos., Journal of Biological Chemistry **269**, 22913 (1994).

[7] A. S. Kashina, R. J. Baskin, D. G. Cole, K. P. Wedaman, W. M. Saxton, and J. M. Scholey, A bipolar kinesin, Nature **379**, 270 (1996).

[8] D. M. Gordon and D. M. Roof, The Kinesin-related Protein Kip1p of Saccharomyces cerevisiae Is Bipolar, Journal of Biological Chemistry **274**, 28779 (1999).

[9] S. Acar, D. B. Carlson, M. S. Budamagunta, V. Yarov-Yarovoy, J. J. Correia, M. R. Niñonuevo, W. Jia, L. Tao, J. A. Leary, J. C. Voss, J. E. Evans, and J. M. Scholey, The bipolar assembly domain of the mitotic motor kinesin-5, Nature Communications **4**, ncomms2348 (2013).

[10] J. E. Scholey, S. Nithianantham, J. M. Scholey, and J. Al-Bassam, Structural basis for the assembly of the mitotic motor Kinesin-5 into bipolar tetramers, eLife **3**, e02217 (2014).

[11] S. K. Singh, H. Pandey, J. Al-Bassam, and L. Gheber, Bidirectional motility of kinesin-5 motor proteins: Structural determinants, cumulative functions and physiological roles, Cellular and Molecular Life Sciences **75**, 1757 (2018).

[12] D. J. Sharp, K. L. McDonald, H. M. Brown, H. J. Matthies, C. Walczak, R. D. Vale, T. J. Mitchison, and

J. M. Scholey, The Bipolar Kinesin, KLP61F, Cross-links Microtubules within Interpolar Microtubule Bundles of Drosophila Embryonic Mitotic Spindles, The Journal of Cell Biology **144**, 125 (1999).

[13] L. C. Kapitein, E. J. G. Peterman, B. H. Kwok, J. H. Kim, T. M. Kapoor, and C. F. Schmidt, The bipolar mitotic kinesin Eg5 moves on both microtubules that it crosslinks, Nature **435**, 114 (2005).

[14] E. R. Hildebrandt, L. Gheber, T. Kingsbury, and M. A. Hoyt, Homotetrameric Form of Cin8p, a Saccharomyces cerevisiae Kinesin-5 Motor, Is Essential for Its in Vivo Function, Journal of Biological Chemistry **281**, 26004 (2006).

[15] L. Tao, A. Mogilner, G. Civelekoglu-Scholey, R. Wollman, J. Evans, H. Stahlberg, and J. M. Scholey, A Homotetrameric Kinesin-5, KLP61F, Bundles Microtubules and Antagonizes Ncd in Motility Assays, Current Biology **16**, 2293 (2006).

[16] S. M. J. L. van den Wildenberg, L. Tao, L. C. Kapitein, C. F. Schmidt, J. M. Scholey, and E. J. G. Peterman, The Homotetrameric Kinesin-5 KLP61F Preferentially Crosslinks Microtubules into Antiparallel Orientations, Current Biology **18**, 1860 (2008).

[17] Y. Shimamoto, S. Forth, and T. M. Kapoor, Measuring Pushing and Braking Forces Generated by Ensembles of Kinesin-5 Crosslinking Two Microtubules, Developmental Cell **34**, 669 (2015).

[18] T. Bodrug, E. M. Wilson-Kubalek, S. Nithianantham, A. F. Thompson, A. Alfieri, I. Gaska, J. Major, G. Debs, S. Inagaki, P. Gutierrez, L. Gheber, R. J. McKenney, C. V. Sindelar, R. Milligan, J. Stumpff, S. S. Rosenfeld, S. T. Forth, and J. Al-Bassam, The kinesin-5 tail domain directly modulates the mechanochemical cycle of the motor domain for anti-parallel microtubule sliding, eLife **9**, e51131 (2020).

[19] G. Goshima and J. M. Scholey, Control of Mitotic Spindle Length, Annual Review of Cell and Developmental Biology **26**, 21 (2010).

[20] B. J. Mann and P. Wadsworth, Kinesin-5 Regulation and Function in Mitosis, Trends in Cell Biology (2018).

[21] J. M. Scholey, G. Civelekoglu-Scholey, and I. Brust-Mascher, Anaphase B, Biology **5**, 51 (2016).

[22] T. U. Mayer, T. M. Kapoor, S. J. Haggarty, R. W. King, S. L. Schreiber, and T. J. Mitchison, Small Molecule Inhibitor of Mitotic Spindle Bipolarity Identified in a Phenotype-Based Screen, Science **286**, 971 (1999).

[23] J. Roostalu, C. Hentrich, P. Bieling, I. A. Telley, E. Schiebel, and T. Surrey, Directional Switching of the Kinesin Cin8 Through Motor Coupling, Science **332**, 94 (2011).

[24] A. Gerson-Gurwitz, C. Thiede, N. Movshovich, V. Fridman, M. Podolskaya, T. Danieli, S. Lakämper, D. R. Klopfenstein, C. F. Schmidt, and L. Gheber, Directionality of individual kinesin-5 Cin8 motors is modulated by loop 8, ionic strength and microtubule geometry, The EMBO Journal **30**, 4942 (2011).

[25] R. Avunie-Masala, N. Movshovich, Y. Nissenkorn, A. Gerson-Gurwitz, V. Fridman, M. Kõivomägi, M. Loog, M. A. Hoyt, A. Zaritsky, and L. Gheber, Phospho-regulation of kinesin-5 during anaphase spindle elongation, Journal of Cell Science **124**, 873 (2011).

[26] C. Thiede, V. Fridman, A. Gerson-Gurwitz, L. Gheber, and C. F. Schmidt, Regulation of bi-directional movement of single kinesin-5 Cin8 molecules, BioArchitecture **2**, 70 (2012).

[27] V. Fridman, A. Gerson-Gurwitz, O. Shapira, N. Movshovich, S. Lakämper, C. F. Schmidt, and L. Gheber, Kinesin-5 Kip1 is a bi-directional motor that stabilizes microtubules and tracks their plus-ends in vivo, J Cell Sci **126**, 4147 (2013).

[28] M. Edamatsu, Bidirectional motility of the fission yeast kinesin-5, Cut7, Biochemical and Biophysical Research Communications **446**, 231 (2014).

[29] M. Edamatsu, Molecular properties of the N-terminal extension of the fission yeast kinesin-5, Cut7., Genetics and molecular research : GMR **15** (2016).

[30] M. Britto, A. Goulet, S. Rizvi, O. von Loeffelholz, C. A. Moores, and R. A. Cross, Schizosaccharomyces pombe kinesin-5 switches direction using a steric blocking mechanism, Proceedings of the National Academy of Sciences , 201611581 (2016).

[31] O. Shapira and L. Gheber, Motile properties of the bi-directional kinesin-5 Cin8 are affected by phosphorylation in its motor domain, Scientific Reports **6**, 25597 (2016).

[32] O. Shapira, A. Goldstein, J. Al-Bassam, and L. Gheber, Possible physiological role for bi-directional motility and motor clustering of the mitotic kinesin-5 Cin8, J Cell Sci , jcs.195040 (2017).

[33] H. Pandey, E. Reithmann, A. Goldstein-Levitin, J. Al-Bassam, E. Frey, and L. Gheber, Drag-induced directionality switching of kinesin-5 Cin8 revealed by cluster-motility analysis, Science Advances **7**, eabc1687 (2021).

[34] Z. R. Gergely, S. Ansari, M. H. Jones, B. Zhou, C. Cash, R. McIntosh, and M. D. Betterton, Kinesin-

5/Cut7 moves bidirectionally on fission-yeast spindles with activity that increases in anaphase, Journal of Cell Science , jcs.260474 (2023).

[35] M. Uteng, C. Hentrich, K. Miura, P. Bieling, and T. Surrey, Poleward transport of Eg5 by dynein–dynactin in Xenopus laevis egg extract spindles, The Journal of Cell Biology **182**, 715 (2008).

[36] N. Ma, U. S. Tulu, N. P. Ferenz, C. Fagerstrom, A. Wilde, and P. Wadsworth, Poleward Transport of TPX2 in the Mammalian Mitotic Spindle Requires Dynein, Eg5, and Microtubule Flux, Molecular Biology of the Cell **21**, 979 (2010).

[37] N. Ma, J. Titus, A. Gable, J. L. Ross, and P. Wadsworth, TPX2 regulates the localization and activity of Eg5 in the mammalian mitotic spindle, J Cell Biol **195**, 87 (2011).

[38] A. Gable, M. Qiu, J. Titus, S. Balchand, N. P. Ferenz, N. Ma, E. S. Collins, C. Fagerstrom, J. L. Ross, G. Yang, and P. Wadsworth, Dynamic reorganization of Eg5 in the mammalian spindle throughout mitosis requires dynein and TPX2, Molecular Biology of the Cell **23**, 1254 (2012).

[39] S. K. Balchand, B. J. Mann, J. Titus, J. L. Ross, and P. Wadsworth, TPX2 Inhibits Eg5 by Interactions with Both Motor and Microtubule, Journal of Biological Chemistry **290**, 17367 (2015).

[40] I. Hagan and M. Yanagida, Kinesin-related cut 7 protein associates with mitotic and meiotic spindles in fission yeast, Nature **356**, 74 (1992).

[41] D. R. Drummond and I. M. Hagan, Mutations in the bimC box of Cut7 indicate divergence of regulation within the bimC family of kinesin related proteins, Journal of Cell Science **111**, 853 (1998).

[42] J. Cahu, A. Olichon, C. Hentrich, H. Schek, J. Drinjakovic, C. Zhang, A. Doherty-Kirby, G. Lajoie, and T. Surrey, Phosphorylation by Cdk1 Increases the Binding of Eg5 to Microtubules In Vitro and in Xenopus Egg Extract Spindles, PLOS ONE **3**, e3936 (2008).

[43] R. Blackwell, C. Edelmaier, O. Sweezy-Schindler, A. Lamson, Z. R. Gergely, E. O'Toole, A. Crapo, L. E. Hough, J. R. McIntosh, M. A. Glaser, and M. D. Betterton, Physical determinants of bipolar mitotic spindle assembly and stability in fission yeast, Science Advances **3**, e1601603 (2017).

[44] M. M. Heck, A. Pereira, P. Pesavento, Y. Yannoni, A. C. Spradling, and L. S. Goldstein, The kinesin-like protein KLP61F is essential for mitosis in Drosophila., The Journal of Cell Biology **123**, 665 (1993).

[45] K. E. Sawin and T. J. Mitchison, Mutations in the kinesin-like protein Eg5 disrupting localization to the mitotic spindle, Proceedings of the National Academy of Sciences **92**, 4289 (1995).

[46] J. Rapley, M. Nicolàs, A. Groen, L. Regué, M. T. Bertran, C. Caelles, J. Avruch, and J. Roig, The NIMA-family kinase Nek6 phosphorylates the kinesin Eg5 at a novel site necessary for mitotic spindle formation, Journal of Cell Science **121**, 3912 (2008).

[47] E. Saeki, S. Yasuhira, M. Shibazaki, H. Tada, M. Doita, T. Masuda, and C. Maesawa, Involvement of C-terminal truncation mutation of kinesin-5 in resistance to kinesin-5 inhibitor, PLOS ONE **13**, e0209296 (2018).

[48] J. S. Weinger, M. Qiu, G. Yang, and T. M. Kapoor, A nonmotor microtubule binding site in kinesin-5 is required for filament crosslinking and sliding, Current Biology **21**, 154 (2011).

[49] S. Nithianantham, M. K. Iwanski, I. Gaska, H. Pandey, T. Bodrug, S. Inagaki, J. Major, G. J. Brouhard, L. Gheber, S. S. Rosenfeld, S. Forth, A. G. Hendricks, and J. Al-Bassam, The mechanochemical origins of the microtubule sliding motility within the kinesin-5 domain organization (2021).

[50] A. Düselder, V. Fridman, C. Thiede, A. Wiesbaum, A. Goldstein, D. R. Klopfenstein, O. Zaitseva, M. E. Janson, L. Gheber, and C. F. Schmidt, Deletion of the Tail Domain of the Kinesin-5 Cin8 Affects Its Directionality, Journal of Biological Chemistry **290**, 16841 (2015).

[51] M. Toya, M. Sato, U. Haselmann, K. Asakawa, D. Brunner, C. Antony, and T. Toda, γ-Tubulin complex-mediated anchoring of spindle microtubules to spindle-pole bodies requires Msd1 in fission yeast, Nature Cell Biology **9**, 646 (2007).

[52] M. Yukawa, C. Ikebe, and T. Toda, The Msd1–Wdr8–Pkl1 complex anchors microtubule minus ends to fission yeast spindle pole bodies, J Cell Biol **209**, 549 (2015).

[53] V. Syrovatkina and P. T. Tran, Loss of kinesin-14 results in aneuploidy via kinesin-5-dependent microtubule protrusions leading to chromosome cut, Nature Communications **6**, 7322 (2015).

[54] A. S. Rodriguez, A. N. Killilea, J. Batac, J. Filopei, D. Simeonov, I. Lin, and J. L. Paluh, Protein complexes at the microtubule organizing center regulate bipolar spindle assembly, Cell Cycle **7**, 1246 (2008).

[55] T. Akera, Y. Goto, M. Sato, M. Yamamoto, and Y. Watanabe, Mad1 promotes chromosome congression by anchoring a kinesin motor to the kinetochore, Nature Cell Biology **17**, 1124 (2015).

[56] O. von Loeffelholz, A. Peña, D. R. Drummond, R. Cross, and C. A. Moores, Cryo-EM Structure (4.5-Å)

of Yeast Kinesin-5–Microtubule Complex Reveals a Distinct Binding Footprint and Mechanism of Drug Resistance, Journal of Molecular Biology **431**, 864 (2019).

[57] H. A. Snaith, A. Anders, I. Samejima, and K. E. Sawin, Chapter 9 - New and Old Reagents for Fluorescent Protein Tagging of Microtubules in Fission Yeast: Experimental and Critical Evaluation, in *Methods in Cell Biology*, Vol. Volume 97, edited by Lynne Cassimeris and Phong Tran (Academic Press, 2010) pp. 147–172.

[58] R. Blackwell, O. Sweezy-Schindler, C. Baldwin, L. E. Hough, M. A. Glaser, and M. D. Betterton, Microscopic origins of anisotropic active stress in motor-driven nematic liquid crystals, Soft Matter **12**, 2676 (2016).

[59] S. Ansari, Z. R. Gergely, P. Flynn, G. Li, J. K. Moore, and M. D. Betterton, Quantifying yeast microtubules and spindles using the Toolkit for Automated Microtubule Tracking (TAMiT) (2023).

[60] M. Yukawa, Y. Teratani, and T. Toda, How Essential Kinesin-5 Becomes Non-Essential in Fission Yeast: Force Balance and Microtubule Dynamics Matter, Cells **9**, 1154 (2020).

[61] R. Ding, K. L. McDonald, and J. R. McIntosh, Three-dimensional reconstruction and analysis of mitotic spindles from the yeast, Schizosaccharomyces pombe., The Journal of Cell Biology **120**, 141 (1993).

[62] J. J. Ward, H. Roque, C. Antony, and F. Nédélec, Mechanical design principles of a mitotic spindle, eLife **3**, e03398 (2015).

[63] C. Edelmaier, A. R. Lamson, Z. R. Gergely, S. Ansari, R. Blackwell, J. R. McIntosh, M. A. Glaser, and M. D. Betterton, Mechanisms of chromosome biorientation and bipolar spindle assembly analyzed by computational modeling, eLife **9**, e48787 (2020).

[64] A. L. Pidoux, M. LeDizet, and W. Z. Cande, Fission yeast pkl1 is a kinesin-related protein involved in mitotic spindle function., Molecular Biology of the Cell **7**, 1639 (1996).

[65] S. A. Rincon, A. Lamson, R. Blackwell, V. Syrovatkina, V. Fraisier, A. Paoletti, M. D. Betterton, and P. T. Tran, Kinesin-5-independent mitotic spindle assembly requires the antiparallel microtubule crosslinker Ase1 in fission yeast, Nature Communications **8**, 15286 (2017).

[66] M. Yukawa, Y. Yamada, T. Yamauchi, and T. Toda, Two spatially distinct Kinesin-14 Pkl1 and Klp2 generate collaborative inward forces against Kinesin-5 Cut7 in S. pombe, J Cell Sci , jcs.210740 (2018).

[67] A. R. Lamson, C. J. Edelmaier, M. A. Glaser, and M. D. Betterton, Theory of Cytoskeletal Reorganization during Cross-Linker-Mediated Mitotic Spindle Assembly, Biophysical Journal **116**, 1719 (2019).

[68] R. Blackwell, O. Sweezy-Schindler, C. Edelmaier, Z. R. Gergely, P. J. Flynn, S. Montes, A. Crapo, A. Doostan, J. R. McIntosh, M. A. Glaser, and M. D. Betterton, Contributions of Microtubule Dynamic Instability and Rotational Diffusion to Kinetochore Capture, Biophysical Journal **112**, 552 (2017).

[69] J. T. Wilson-Grady, J. Villén, and S. P. Gygi, Phosphoproteome Analysis of Fission Yeast, Journal of Proteome Research **7**, 1088 (2008).

[70] A. Koch, K. Krug, S. Pengelley, B. Macek, and S. Hauf, Mitotic Substrates of the Kinase Aurora with Roles in Chromatin Regulation Identified Through Quantitative Phosphoproteomics of Fission Yeast, Science Signaling **4**, rs6 (2011).

[71] A. N. Kettenbach, L. Deng, Y. Wu, S. Baldissard, M. E. Adamo, S. A. Gerber, and J. B. Moseley, Quantitative Phosphoproteomics Reveals Pathways for Coordination of Cell Growth and Division by the Conserved Fission Yeast Kinase Pom1, Molecular & Cellular Proteomics **14**, 1275 (2015).

[72] M. P. Swaffer, A. W. Jones, H. R. Flynn, A. P. Snijders, and P. Nurse, Quantitative Phosphoproteomics Reveals the Signaling Dynamics of Cell-Cycle Kinases in the Fission Yeast Schizosaccharomyces pombe, Cell Reports **24**, 503 (2018).

[73] I. Derényi, F. Jülicher, and J. Prost, Formation and Interaction of Membrane Tubes, Physical Review Letters **88**, 238101 (2002).

[74] Z. T. Olmsted, A. G. Colliver, T. D. Riehlman, and J. L. Paluh, Kinesin-14 and kinesin-5 antagonistically regulate microtubule nucleation by γ-TuRC in yeast and human cells, Nature Communications **5**, 5339 (2014).

[75] C. Fu, J. J. Ward, I. Loiodice, G. Velve-Casquillas, F. J. Nedelec, and P. T. Tran, Phospho-Regulated Interaction between Kinesin-6 Klp9p and Microtubule Bundler Ase1p Promotes Spindle Elongation, Developmental Cell **17**, 257 (2009).

[76] M. Yukawa, M. Okazaki, Y. Teratani, K. Furuta, and T. Toda, Kinesin-6 Klp9 plays motor-dependent and -independent roles in collaboration with Kinesin-5 Cut7 and the microtubule crosslinker Ase1 in fission yeast, Scientific Reports **9**, 7336 (2019).

[77] L. C. Kapitein, M. E. Janson, S. M. J. L. van den Wildenberg, C. C. Hoogenraad, C. F. Schmidt, and

E. J. G. Peterman, Microtubule-Driven Multimerization Recruits ase1p onto Overlapping Microtubules, Current Biology **18**, 1713 (2008).

[78] S. Moreno, A. Klar, and P. Nurse, Molecular genetic analysis of fission yeast Schizosaccharomyces pombe, in *Guide to Yeast Genetics and Molecular Biology*, Vol. Volume 194 (Academic Press, 1991) pp. 795–823.

[79] K. Okazaki, N. Okazaki, K. Kume, S. Jinno, K. Tanaka, and H. Okayama, High-frequency transformation method and library transducing vectors for cloning mammalian cDNAs by trans-complementation of Schizosaccharomyces pombe, Nucleic Acids Research **18**, 6485 (1990).

FIG. 2. Sequential truncation of the *cut7* C-terminal tail compromises cell viability and spindle assembly and function. (A) Schematic of truncation mutants aligned with the BimC box region amino acid sequence (upper), showing conserved residues (purple), temperature-sensitive lethal mutations (cyan) and truncation sites (red triangles). Each allele is labeled with amino acids retained (gray bar and numbers), C-terminal GFP (green bar), and viability (left). (B) Serial dilution assays of cells grown at 25°C (upper) and 37°C (lower). All viable strains grew at 25 °C, and *cut7-1017* and *cut7-989-1028Δ* showed temperature-sensitive growth inhibition or lethality. (C-D) Fluorescence images showing mCherry-atb2 spindle MTs (red, left), Cut7-GFP (green, center), and merge (right) for cells grown and imaged at 25°C and shown as maximum-intensity projections of confocal sections. (C) Cut7-1032-GFP localized near SPBs and was visible on the interpolar spindle. (D) Cut7-989-1028Δ-GFP localized near SPBs and appeared dimmer at the spindle midzone. Scalebar, 1 μm. (E) Spindle length dynamics. Spindle length as a function of time was averaged for 19 *cut7*, 9 *cut7-1032*, 11 *cut7-1017*, and 14 *cut7-989-1028Δ* cells. We defined $t = 0$ as the time of anaphase onset to align the curves (dashed vertical line, Methods). Time at which steady anaphase elongation speed was reached marked on each curve by the vertical bar. (F) Pre-anaphase spindle length, (G) anaphase spindle elongation speed, and (H) the time between anaphase onset and start of steady elongation. Error bars in E-H are standard error of the mean. (I-J) Fluorescence images showing mCherry-atb2 spindle MTs (red, left), Cut7-GFP (green, center), and merge (right) for cells grown and imaged at 37°C and shown as maximum-intensity projections of confocal sections. (I) Cut7-1032-GFP localizes similarly to Cut7-GFP on the bipolar spindle. (J) Cut7-989-1028Δ-GFP localizes near the center of the monopolar spindle due to a failure of SPB bipolar spindle assembly. Scalebar, 1 μm. (K) Fraction of cells grown at 37°C that formed bipolar spindles (green), monopolar spindles that became bipolar (yellow), or persistent monopolar spindles (red) for 21 *cut7*, 17 *cut7-1032*, 21 *cut7-1017*, and 21 *cut7-989-1028Δ* cells.

FIG. 3. Kymographs of Cut7-GFP movement, intensity distribution along the spindle, and midzone-to-pole intensity ratio. (A) Kymograph of Cut7-GFP movement on a bipolar spindle. Bright regions are SPB-localized motor and dim streaks show midzone-localized moving clusters of motor. Scalebars, 1 μm horizontal, 60 sec vertical. (B-D) Cut7-GFP intensity along the spindle axis for spindle length of 2 μm (B), 4 μm (C) and 7 μm (D). Each colored line is from a separate image, and the thick black curve is the average. Data from 202 images (B), 45 images (C), and 28 images (D), from 29 cells. (E-H) Kymographs of Cut7-FL-GFP (E), Cut7-1032-GFP (F), Cut7-1017-GFP (G), and Cut7-989-1028Δ-GFP (H). (I-L) Average Cut7-GFP intensity along the spindle axis for Cut7-FL-GFP (I), Cut7-1032-GFP (J), Cut7-1017-GFP (K), and Cut7-989-1028Δ-GFP (L). (M) Total Cut7-GFP intensity on the spindle as a function of spindle length. (N) Cut7-GFP midzone-to-pole ratio as a function of spindle length. (O) Cut7-GFP midzone-to-pole ratio as a function of time, where $t = 0$ is anaphase onset (dashed vertical line). Intensity data from 29 *cut7*, 34 *cut7-1032*, 29 *cut7-1017*, and 30 *cut7-989-1028Δ* cells. Error bars in M and N are computed as the standard error of the mean assuming independent intensity measurements for each images at each integer spindle length. Error bars in O are computed as the standard error of the mean assuming independent intensity measurements are separated by 1 minute time intervals.

FIG. 4. Kinesin-14 deletion rescued lethal *cut7* tail truncation mutants and led to spindle MT minus-end protrusions. (A) Serial dilution assay of truncation mutants in the *pkl1Δ, klp2Δ* background 25°C (upper) and 37°C (lower). Cells with *cut7-1006* and *cut7-988* are viable in this background but show temperature-sensitive growth inhibition or lethality. (B) Schematic of protrusions and Cut7-GFP accumulation. Upper, MT with its minus-end detached from the SPB protrudes beyond the SPB and distorts the nuclear envelope. (Lower), Cut7-GFP near SPBs moves toward minus ends of protruding MT and collects at the protrusion tip. (C) Fluorescence image showing mCherry-atb2 spindle and protrusion MTs (red, left), Cut7-GFP (green, center), and merge (right) for cell grown and imaged at 25°C and shown as maximum-intensity projections of confocal sections. (D-H) Time-lapse imaging of spindle and protrusion dynamics showing mCherry-atb2 spindle and protrusion MTs (red, upper), Cut7-GFP (green, center), and merge (lower) for cells grown and imaged at 25°C and shown as maximum-intensity projections of confocal sections. (D-F) For *cut7-FL* and *cut7-1032* cells, protrusions are frequent and dynamic. (G) For *cut7-1017* cells protrusions are less frequent. (H) For *cut7-988* cells protrusions were not visible. Scalebar, 1 μm. Time between images 82.4s (D), 82.4s (E), 27.4s (F), 36.8s (G), and 35.8s (H). (I) Percentage of cells with visible protrusions decreases as more of the tail is truncated. Data from 75 *cut7*, 43 *cut7-1032*, 39 *cut7-1017*, and 39 *cut7-1006* and 39 *cut7-988* cells. Error bars are are estimated from Poisson counting statistics.

FIG. 5. Spindle length, Cut7-GFP midzone-to-pole intensity ratio, and protrusion frequency vary nonlinearly with tail truncation in the kinesin-14 *pkl1Δ klp2Δ* background. (A) Pre-anaphase spindle length decreases for the most severe tail truncation alleles *cut7-1006* and *cut7-988*. Data from 14 *cut7*, 13 *cut7-1032*, 11 *cut7-1017*, 6 *cut7-1006*, and 17 *cut7-988* cells that did not exhibit protrusions (Methods). Error bars are standard error of the mean. (B) Cut7-GFP midzone-to-pole intensity ratio as a function of spindle length shows similar trends as in the *pkl1+ klp2+* background but is consistently low for *cut7-988* cells. Intensity data from 44 *cut7*, 34 *cut7-1032*, 35 *cut7-1017*, 13 *cut7-1006*, and 40 *cut7-988* cells. (C) Protrusion frequency (red) and midzone-to-pole intensity ratio (dark blue, *pkl1Δ klp2Δ* background; light blue, *pkl1+ klp2+* background) vary nonlinearly with position of tail truncation (black line, linear model). Error bars of protrusion frequency are estimated from Poisson counting statistics, Error bars in B are computed as the standard error of the mean assuming independent intensity measurements for each images at each integer spindle length. Error bars in C are computed as the standard error of the mean assuming independent intensity measurements are separated by 1 minute time intervals. Lower, Cut7 tail with truncation sites marked with red triangles and phosphorylation sites with pink boxes. Inset: sequence and phosphorylated residues for a.a. 1006-1032.