Functional genomic analysis of non-canonical DNA regulatory elements of the

aryl hydrocarbon receptor

Tajhal D Patel^, Manjula Nakka*, Sandra L Grimm*#^, Cristian Coarfa*#^, Daniel A

Gorelick*#

*Center for Precision Environmental Health

#Department of Molecular & Cellular Biology

^Dan L Duncan Comprehensive Cancer Center

Baylor College of Medicine, Houston, Texas, USA

Authors for correspondence: CC (coarfa@bcm.edu), DAG (gorelick@bcm.edu)

ABSTRACT

The aryl hydrocarbon receptor (AHR) is a ligand-dependent transcription factor that

binds DNA and regulates genes in response to halogenated and polycyclic aromatic

hydrocarbons. AHR also regulates the development and function of the liver and the

immune system. In the canonical pathway, AHR binds a consensus DNA sequence,

termed the xenobiotic response element (XRE), recruits protein coregulators, and

regulates target gene expression. Emerging evidence suggests that AHR may regulate

gene expression via an additional pathway, by binding to a non-consensus DNA

sequence termed the non-consensus XRE (NC-XRE). The prevalence of NC-XRE

motifs in the genome is not known. Studies using chromatin immunoprecipitation and

reporter genes provide indirect evidence of AHR-NC-XRE interactions, but direct

evidence for an AHR-NCXRE interaction that regulates transcription in a natural

genomic context is lacking. Here, we analyzed AHR binding to NC-XRE DNA on a

genome-wide scale in mouse liver. We integrated ChIP-seq and RNA-seq data and

identified putative AHR target genes with NC-XRE motifs in regulatory regions. We also

performed functional genomics at a single locus, the mouse *Serpine1* gene. Deleting

NC-XRE motifs from the *Serpine1* promoter reduced the upregulation of *Serpine1* by

TCDD, an AHR ligand. We conclude that AHR upregulates *Serpine1* via NC-XRE DNA.

NC-XRE motifs are prevalent throughout regions of the genome where AHR binds.

Taken together, our results suggest that AHR regulates genes via NC-XRE motifs. Our

results will also improve our ability to identify AHR target genes and their physiologic

relevance.

INTRODUCTION

The toxic effects of secondhand cigarette smoke, fossil fuels, and byproducts of

industrial combustion are caused by halogenated and polycyclic aromatic hydrocarbons

such as TCDD. Understanding the signaling pathways by which such compounds cause

toxicity is crucial for predicting tolerable exposure levels and for reversing the effects of

adverse exposure. Aromatic hydrocarbons activate aryl hydrocarbon receptors (AHR),

ligand-dependent transcription factors that recruit protein coregulators to DNA and

directly regulate gene expression (Denison et al., 2011). AHR forms a dimer with aryl

hydrocarbon receptor nuclear translocator protein (ARNT1, also known as HIF1B) to

regulate transcription (Hankinson, 2005; Reyes et al., 1992). The AHR-ARNT1 complex

binds a consensus DNA sequence (GCGTG) termed the xenobiotic response element

(XRE) (Denison et al., 1988; Yao and Denison, 1992).

Recent evidence complicates these findings. AHR may regulate gene expression by

binding to non-consensus XRE sequences. A chromatin immunoprecipitation study

found that half of TCDD-AHR complexes in mouse liver were bound to DNA lacking a

consensus XRE (Dere et al., 2011). Exploring the genome more closely, Huang and

colleagues identified a non-consensus DNA sequence to which AHR binds (Huang and

Elferink, 2012). Chromatin immunoprecipitation, gel shift and reporter assays support

the hypothesis that AHR binds a non-consensus XRE DNA sequence (nucleotides

GGGA, referred to as NC-XRE motif) (Huang and Elferink, 2012; Jackson et al., 2014;

Joshi et al., 2015; Wilson et al., 2013). Such studies were limited because they did not

analyze AHR-NCXRE interactions on a genome-wide scale. Additionally, chromatin

immunoprecipitation identifies where AHR binds to DNA does not reveal whether

transcription actually occurs. Gel shift and reporter gene assays sometimes fail to reflect

what occurs in a natural genomic context. Direct evidence for an AHR-NCXRE

interaction that regulates transcription in a natural genomic context is lacking.

Here, we sought to analyze AHR binding to non-canonical DNA elements on a genome-

wide scale and, in the case of a single gene, directly test whether NC-XRE motifs are

required for AHR-dependent target gene expression.

RESULTS

Frequency of NC-XRE motifs

To determine the distribution of AHR binding to DNA following TCDD exposure, we

analyzed published AHR ChIP-seq data from livers of C57BL/6 male and female mice

(28 days old) following 2 hour exposure to 30 µg/kg TCDD (GEO accession

#GSE97634, GSE97636) (Fader et al., 2017; Fader et al., 2019). This data was

previously assayed for AHR binding to consensus XRE DNA [GCGTG] but was not

interrogated for AHR binding to non-consensus DNA such as NC-XRE [GGGA] or

RelBAHRE [GGGTGCAT]. We found that, on average, 30% of peaks (AHR binding to

DNA) contained a NC-XRE motif without a nearby XRE. 9% of peaks contained an XRE

motif without a nearby NC-XRE, while 51% of peaks contained an XRE and NC-XRE

(Fig 1A, C). Fewer than 1% of peaks contained RelBAHRE.

We wondered whether XRE and NC-XRE peaks were distributed equally among

promoter and enhancer DNA. We observed more peaks at promoter regions (within 3

kb of transcription start sites) containing XRE motifs alone or XRE+NC-XRE motifs

together compared to promoter regions containing NC-XRE alone (Fig 1B). This

suggests that AHR binding to XRE is stronger in promoter regions, whereas AHR

binding to NC-XRE is stronger in enhancer regions.

XRE, NC-XRE and RelBAHRE motifs are different lengths, which affects the frequency

with which they are found in the genome. We assessed the likelihood of finding XRE,

NC-XRE or RelBAHRE motifs in AHR ChIP peaks versus in a random region of DNA.

NC-XRE and XRE motifs are statistically significantly enriched in AHR ChIP peaks,

whereas RelBAHRE motifs are not (Fig 1D). Because of the low number of RelBAHRE

in AHR peaks, we focus on the XRE and NC-XRE motifs in this study.

Repeated NC-XRE motifs associated with AHR target genes

The preceding analysis examined presence of NC-XRE motifs in all regions where AHR

was bound to DNA. However, transcription factors like AHR may bind DNA without

affecting gene expression. Therefore, we sought to identify AHR ChIP-seq peaks

associated with AHR target genes. To identify AHR target genes, we integrated the AHR

ChIP-seq data with published RNA-seq data from mouse liver (Fader et al., 2017; Nault

et al., 2015; Nault et al., 2018). To maximize identification of AHR target genes, we

included RNA-seq data from mouse livers treated with a variety of doses of TCDD for

various durations of exposure (Table 1). We define AHR target genes as differentially

expressed genes (RNA-seq, fold difference in TCDD vs vehicle exceeding 1.5x to cover

both up- and downregulation, FDR<5%) containing AHR binding site (ChIP peak) within

10 kb of a gene body. The 10 kb parameter allows us to include promoter and gene

proximal regulatory DNA peaks, important because our analysis of ChIP-seq data

suggested that AHR binds NC-XRE at enhancers. Of the differentially expressed genes

in TCDD vs vehicle, 520 had AHR binding peaks at XRE, NC-XRE or XRE+NC-XRE

motifs within 10 kb of a gene body. 28 of these genes had AHR binding peaks at NC-

XRE but not at XRE or XRE+NC-XRE motifs.

The preceding analysis is for any single instance of XRE or NC-XRE DNA sequence.

For some transcription factors, repeated DNA binding sites produce a more robust

response (Klein-Hitpass et al., 1988). We sought to examine the association between

repeated runs of XRE or NCXRE motifs and target gene expression. Fig 2A shows the

relative frequency of AHR target genes with AHR peaks containing between 3 and 10

NC-XRE motifs separated by up to 50 basepairs (bp), compared to target genes found

near similar patterns of XRE peaks. 24% of AHR peaks contain 2 or more NC-XRE

motifs separated by 25 bp or less (Fig 2B). AHR peaks containing 5 NC-XRE motifs

separated by 25 bp or less were found proximal to 82 AHR target genes. Compared to

repeated XRE motifs, runs of 5 or more NC-XREs are enriched at AHR binding sites

proximal to transcription start sites (H3K4me, H3K27ac, Fig 2C). This suggests that

NCXREs may be more potent when they occur in tandem repeats. The top differentially

regulated genes with 5 or more NCXRE motifs are shown in Fig 2D. As validation, the

known AHR target gene *Serpine1* (aka *Pai1*) contains 5 NCXRE motifs in the putative

promoter, consistent with results that AHR upregulates *Serpine1* via NC-XRE DNA

(Huang and Elferink, 2012). AHR binding was also detected at NC-XRE DNA near

genes like *Nqo1*, a canonical AHR target gene thought to be regulated by XRE motifs

(Yeager et al., 2009). It is possible that NC-XRE DNA also contributes to the regulation

of NQO1 expression.

Motifs flanking XRE or NC-XRE sequence

Transcription factors bind regulatory DNA in a complex composed of multiple different

transcription factors. For example, AHR is associated with the transcription factor

estrogen receptor alpha at regulatory DNA (Beischlag and Perdew, 2005; Klinge et al.,

2000; Madak-Erdogan and Katzenellenbogen, 2012). Another transcription factor, ARNT,

is thought to be an obligate co-regulator of AHR (Lo and Matthews, 2012; Reyes et al.,

1992). Previous studies have focused on transcription factor binding at XRE motifs. We

identified known transcription factor motifs flanking NC-XRE motifs. We focused on AHR

ChIP peaks containing at least 2 NC-XRE motifs within 25 basepairs of each other. The

top 12 most frequently occurring known transcription factor motifs are shown in Figure

3. We found several motifs enriched in peaks containing both XRE and NC-XRE motifs.

As expected, the ARNT motif was enriched in both, supporting the previously known

association with XRE motifs and suggesting that the AHR-ARNT complex could be

associated with NC-XRE motifs. Several motifs were enriched flanking NC-XRE but not

XRE, such as androgen receptor binding site and SMAD3 site. This suggests that there

may be crosstalk between AHR and other signaling pathways, such as TGF-beta, at
NC-XRE motifs.

Testing NC-XRE function in a natural genomic context

We mined published RNA-seq and ChIP-seq data and identified candidate genes
upregulated by TCDD in mouse liver, where such genes contain an AHR ChIP peak
containing XRE or NCXRE (or both) within 10 kb of the gene body. We exposed mouse
Hepa1-6 liver cells to 10 nM TCDD for 2 hours and assayed expression of 8 candidate
genes identified from our analysis of mouse liver. Three genes were upregulated more
than 2-fold compared to vehicle in Hepa1-6 cells (Figure 4A). For one of these genes,
Serpine1, ChIP studies in mouse liver identified an AHR binding site ~150 bp upstream
of the transcription start site, which contains a run of 5 NC-XRE motifs (Fig 4B) (Huang
and Elferink, 2012). Indirect evidence suggests that AHR upregulates *Serpine1*
expression via NC-XRE motifs, but this has never been tested directly.

We deleted the 5x NC-XRE motif in the *Serpine1* promoter, generated clonal mutant cell
lines, and directly tested whether the 5x NC-XRE motif contributes to TCDD-dependent
Serpine1 expression. We transfected Hepa1-6 cells with Cas9 protein together with a
guide RNA targeting this NCXRE sequence. We picked single cells and established a
clonal cell line (clone 14) containing mutations in the target NCXRE (Fig 4B). Bulk
sequencing of clone 14 cells revealed a heterogenous population with 57% of the DNA
sequence containing a 24bp deletion of the 5x NC-XRE, 14% had a 2 bp deletion that
does not disrupt the NC-XRE motifs, and 28% of the DNA was wild-type. We then
derived additional single cell colonies from the original clone 14, denoted as clone 14-10
and clone 14-28. Bulk DNA sequencing of each of these clones revealed no wild-type
DNA. 95% of the DNA contained the expected 24bp deletion encompassing the 5x NC-
XRE, while 5% had the same 2bp deletion that does not disrupt the NC-XRE motifs.

We exposed mutant and wild-type cells to TCDD or vehicle for 2 hours and assayed
Serpine1 expression by RT-qPCR. In all clones, deletion of the NCXRE reduced

upregulation of *Serpine1* in response to TCDD (Fig 4C). To test whether AHR function
was normal, we also examined expression of *Cyp1a1*, a prototypical AHR target gene
thought to be regulated by XRE motifs. All cell lines exhibited increased *Cyp1a1*
expression following TCDD exposure, and there was no difference in mean *Cyp1a1*
expression levels between wild-type and any mutant cell line. Our results suggest that
AHR transcription factor function is normal in NC-XRE clone 14 mutant cell lines. We
conclude that AHR upregulates *Serpine1* via this NCXRE sequence 150 bp upstream of
the transcription start site.

DISCUSSION

We performed a genome-wide assessment of NC-XRE motifs and find they are
prevalent in regulatory regions associated with AHR target genes. We also provide
functional evidence that NC-XRE motifs in the *Serpine1* promoter are necessary for full
upregulation of *Serpine1* in response to TCDD.

There are few examples where a transcription factor binding site DNA was mutated in
an endogenous genomic context, allowing investigators to test whether a specific DNA
sequence is required for transcription factor activity. Carleton and colleagues used
catalytically inactive Cas9 to identify and block enhancer DNA bound by estrogen
receptor alpha (Carleton et al., 2017). Other labs have deleted hundreds or thousands
of basepairs in enhancer DNA to interrogate enhancer function (reviewed in (Lopes et
al., 2016)). These studies are fruitful but lack the resolution to identify specific DNA
sequence to which transcription factors bind. Here we used targeted, precise mutation
to demonstrate the necessity of repeated NC-XRE sequence to regulate an AHR target
gene.

However, we have only a basic understanding of the physiologic relevance of AHR
signaling at NC-XRE motifs. Only two target genes (*Serpine1* and *Cdkn1a*) are known
to be directly regulated by AHR at NC-XRE DNA, and these interactions were explored

in only one cell type, hepatocytes, in adulthood (Huang and Elferink, 2012; Jackson et

al., 2014). The degree to which other AHR target genes are regulated by non-

consensus DNA sequence, in additional tissues and cell types and at different stages of

organismal development, is not known.

ARNT1 is considered an obligate binding partner of AHR (Reyes et al., 1992), but these

studies focused on AHR-ARNT1 interactions at XRE DNA. Whether ARNT1 binds AHR

at NC-XRE DNA is not known. Studies suggest that under certain conditions AHR can

act independently of ARNT1 to upregulate gene expression. In mouse liver, AHR was

bound to NC-XRE in the promoter of the *Serpine1* gene following TCDD exposure, but

no ARNT1 was detected (Huang and Elferink, 2012). In a rat hepatoma cell line

containing a Serpine1:luciferase reporter construct, TCDD increased luciferase levels

but knockdown of ARNT1 had no effect (Huang and Elferink, 2012). These results

suggest that AHR may bind NC-XRE in the absence of ARNT1.

A limitation of the previous studies is that none considered ARNT2. ARNT2 was shown

to interact with AHR and XRE DNA *in vitro* (Hirose et al., 1996; Rowatt et al., 2003;

Tanguay et al., 2000), but to our knowledge there is absence of evidence whether

ARNT2 interacts with AHR *in vivo*. DNA binding, coregulator recruitment and target

gene expression could be different between AHR-ARNT1 and AHR-ARNT2

transcriptional complexes. One possibility is that ARNT1 preferentially associates with

AHR-XRE complexes, while ARNT2 preferentially associates with AHR-NC-XRE

complexes.

We appreciate that DNA regulatory elements may be incompletely conserved between

species. Showing that AHR binds NCXRE DNA to regulate gene expression in any

species is important foundational knowledge. While the specific number or configuration

of NCXRE sequence at a given promoter or enhancer may differ between strains of

animals or between species, it is extremely unlikely that AHR regulates gene expression

by binding to NCXRE in mice but not in humans. Considering that AHR binds identical

XRE DNA elements in zebrafish, mice and humans (Andreasen et al., 2002; Denison et

al., 1988; Tanguay et al., 1999; Yao and Denison, 1992), we argue that the same is likely

true for NCXRE.

METHODS

Re-analysis of published murine liver ChIP-Seq datasets

ChIP-Seq datasets of AHR in murine livers after 2hrs of TCDD treatment in male

(GSE97634) and female (GSE97636) mice were downloaded from NCBI Gene

Expression Omnibus (GEO) (Fader et al., 2017; Fader et al., 2019; Nault et al., 2016).

Data were trimmed using trim_galore, mapped to the mouse genome build UCSC

mm10 using bowtie2, and peaks were called using MACS2 using the provided controls

(Langmead and Salzberg, 2012; Zhang et al., 2008). Peaks were merged using

bedtoools across both sexes (Quinlan and Hall, 2010). Each peak was searched for

AHR motifs using an in-house script. To test the likelihood of finding AHR motifs

anywhere in the genome, random DNA peaks were generated matching the

chromosome and size distribution as previously described (Coarfa et al., 2020), then

searched for AHR motifs. Enrichment of AHR motifs compared to the random controls

was assessed using a Fisher's exact test, with significance achieved at p-value<0.05.

Venn diagrams of AHR motifs in the merged AHR peaks were derived using bedtools.

Genomic distribution of AHR peaks over genomic elements was inferred using bedtools

and visualized using GraphPad Prism.

Runs of AHR motifs, eg sequences of successive motifs within a maximum distance

from each other, were determined using an in-house Python script. Flanking sequences

were determined using the bedtools software. Enriched motifs within the flanking

sequences were determined using the HOMER software (Heinz et al., 2010).

Additional ChIP-Seq datasets of histone modifications in the mouse liver were

downloaded from NCBI GEO using Encode mouse liver from 8 week old mice:

GSM1000140, GSM769014 (Yue et al., 2014). Data were trimmed using trim_galore,

mapped to the mouse genome build UCSC mm10 using bowtie2 (Langmead and

Salzberg, 2012), and ChIP-Seq signal distribution over genomic regions was computed

using the HOMER software.

Re-analysis of published murine liver AhR RNA-Seq datasets and integration with

ChIP-seq datasets

Several RNA-seq datasets of mouse liver after treatment with TCDD at different doses

and for multiple timepoints were downloaded from NCBI GEO: GSE109863, GSE62902,

and GSE87519 (Fader et al., 2017; Nault et al., 2015; Nault et al., 2018)(Table 1). Data

was trimmed using trim_galore, then mapped onto the mouse genome build UCSC

mm10 using STAR (Dobin et al., 2013), and gene expression was quantified using

feature_counts (Liao et al., 2014). Differentially expressed genes between TCDD and

vehicle treatment were inferred using the EdgeR and RUVr R packages (Risso et al.,

2014; Robinson et al., 2010), with significance achieved at fold change exceeding 1.5x

and FDR-adjusted p-value<0.05. Genomic locations (represented as bed files) and

gene signatures were integrated using bedtools. To capture both promoter and proximal

enhancer effects, peaks were considered as associated with genes if the gene body

was within 10,000 basepairs of a peak.

Cell culture

The mouse hepatoma cell line Hepa 1–6 was obtained from ATCC (Manassas, VA)

(Darlington et al., 1980). Hepa 1-6 cells cells were cultured in 10 cm plates in

Dulbecco's Modified Eagle's medium (DMEM) supplemented with 1% (v/v)

Penicillin/Streptomycin and 10% (v/v) FBS in a humidified incubator (5% CO_2, 37°C).

CRISPR mutagenesis of Hepa 1-6 cells

We used CHOPCHOP (Labun et al., 2019) to design guide RNAs targeting the 5x NC-

XRE motif region 150bp upstream of transcription start in the mouse Serpine1 gene

(target sequence CAGCAAGTCACTGGGAGGGA<u>GGG</u>, PAM motif underlined).

Synthetic single guide RNA—modified with 2'-O-methyl at 3 first and last bases, and 3'-

phosphorothioate bonds between first 3 and last 2 bases—and purified SpCas9-2NLS

protein was purchased from Synthego (Redwood City, CA). We mixed guide RNA and

Cas9 (1.3:1 ratio) to form ribonucleoproteins according to Synthego's recommended

protocol. Guide RNA and Cas9 ribonucleoprotein complexes were transfected into

Hepa1-6 cells using Lipofectamine™ CRISPRMAX™ Cas9 Transfection reagent

(Thermo Fisher Scientific, Waltham, MA) according to the manufacturer's protocol.

100,000 Hepa-1-6 cells were mixed with the ribonucleoprotein-transfection solution and

split into two wells for genomic analysis and clonal expansion. Cells were incubated in a

humidified incubator (5% CO_2, 37 °C) for 24 hours. We then changed the media and

allowed the cells to incubate for another 3 days before genotyping or deriving single-cell

colonies for clonal expansion.

To derive colonies from single cells, we took 10 cells / mL suspension and aliquoted 100

µl into each well of the 96-well plate. The plates were checked daily and wells with one

cell were marked. Cells were monitored and allowed to grow for 1-2 weeks before

transfer to a 6 well plate for genotyping and further use.

Genotyping cell lines

Genomic DNA was extracted from cells using phenol-chloroform extraction. Cells were

lysed with DNA lysis buffer (10 mM Tris, 200 mM NaCl, 5 mM EDTA, 1% SDS, 0.4

mg/ml proteinase K). Equal volume of phenol:choloroform:isoamyl alcohol was added

for DNA extraction into upper phase followed by ethanol-sodium acetate precipitation of

DNA. We used PCR to amplify the region flanking the NC-XRE motifs using primers

5'- AAGCCAGGCCAACTTTTCCT and 5'- CGGTCCTCCTTCACAAAGCT. Amplicons

were then ligated into pCR4-TOPO vector using a TOPO TA cloning kit (catalogue

450030, ThermoFisherScientific, Waltham, MA) according to the manufacturer's

protocol. Plasmids were transfected into competent cells, plated and grown overnight.

10-20 bacterial colonies per plate were picked for miniprep DNA extraction and Sanger

sequencing using M13 forward primer. DNA sequence was aligned to wild-type using

MacVector software (MacVector Inc., Apex, NC). If 19 out of 20 colonies contained the

same mutant DNA sequence, and 1 colony contained wild-type DNA, we would

conclude that 95% of that cell line contained mutant DNA, and 5% wild-type. Clone 14

was selected based on the presence of a 24 bp deletion that encompassed the entire 5x

NC-XRE motif (see Figure 4).

Quantitative reverse-transcription PCR

70-80% confluent Hepa-1-6 wild-type and mutant cells were exposed to media

containing 10 nM TCDD or DMSO (vehicle) for 2 hours. To extract RNA, cells were

lysed in TRIzol followed by RNA extraction according to the manufacturer's protocol

using Direct-zol RNA Miniprep Kit, including the on-column DNase digestion (catalogue

11-331, Zymo Research, Irvine, CA). RNA concentration was measured using a

NanoDrop 2000 Spectrophotometer (ThermoFisherScientific, Waltham, MA). Purity of

the RNA (A260/A280) was ≥ 2 with the yield of 60 ng/µl or higher. We converted 1000

ng of the RNA into cDNA using iScript Reverse Transcription Supermix for RT-qPCR

(Biorad, Hercules, CA; cat no 1708841) according to the manufacturer's protocol (total

volume 20 µl, incubated for 5 min at 25°C, followed by incubation at 46°C for 20 min,

then 95°C for 1 min). For the qPCR reaction, 4 µl of 1:10 dilution of cDNA was mixed

with SsoAdvanced Universal SYBR Green Supermix (cat no 1725271, Bio-Rad,

Hercules, CA) and 500 nM each primer up to a total volume of 10 µl and amplified in a

CFX96 Real-Time System (Bio-Rad) with the following program: 95C for 30 sec, then

95C for 10 sec and 60C for 30 sec for 40 cycles. Primer pairs were either designed

using MacVector software or were previously published. Ct values were calculated

using CFX Maestro software version 4.1.2433.1219 (Bio-Rad). Target gene expression

was compared to reference gene expression using the $2^{-\Delta\Delta Ct}$ method. The results shown

were the average of three or more independent experiments. Statistical analysis was

performed using Prism version 9.4.2 (GraphPad Software, Boston, MA).

Primer sequences:

CYP1B1 gene (Liu et al., 2015)

Cyp1b1 FP: 5'-CCAGATCCCGCTGCTCTACA-3'

Cyp1b1 RP: 5'-TGGACTGTCTGCACTAAGGCTG-3'

AHRR (Aryl Hydrocarbon Receptor Repressor) gene (Bernshausen et al., 2006)

AHRR FP: 5'-GTTGGATCCTGTAGGGAGCA-3'

AHRR RP: 5'-AGTCCAGAGGCTCACGCTTA-3'

Mmp24os1 (BC029722) gene

BC029722 FP: 5'-CGCTTTCTAATCGCCTGCAC-3'

BC029722 RP: 5'-TGAGGAGATAAAAGCCAGGCC-3'

Cd36 gene (Niu et al., 2018)

CD36 FP: 5'-TTGTGGAGCTCAAAGACCTG-3'

CD36 RP: 5'-TGCAAGAAGCGGGATGTAGTC-3'

P21Cip1 gene (Jackson et al., 2014)

Cdkn1a FP: 5'-TGTCTTGCACTCTGGTGTCTGAG-3'

Cdkn1a RP: 5'-CAATCTGCGCTTGGAGTGATAG-3'

HROB gene

HROB FP: 5'-AAGAGGAGCTCTCAGAGGCA-3'

HROB RP: 5'-ATGGTGGATGCCCTGTCTTG-3'

IL1RN Gene (Isoda et al., 2005)

IL1RN FP: 5′-CTTTACCTTCATCCGCTCTGAGA-3′

IL1RN RP: 5′-TCTAGTGTTGTGCAGAGGAACCA-3′

MAF BZIP Transcription Factor F gene

MAFF FP: 5' ATGGCTGTGGATCCCTTATCT-3'

MAFF RP: 5' CATCAGCGCTTCATCCGACA-3'

UGT1A7C gene

UGT1A7C FP: 5'-GTCATCCAAAGACTCGGGCA-3'

UGT1A7C RP: 5'-GGGCATCATCACCATCGGAA-3'

18s rRNA (Turner et al., 2018)

m18s-FP: 5'-GTAACCCGTTGAACCCCATT-3'

m18s-RP: 5'-CCATCCAATCGGTAGTAGCG-3'

CYP1A1 gene (Jackson et al., 2014)

CYP1A1-FP: 5'-GCCTAACTCTTCCCTGGATGC-3'

CYP1A1-RP: 5'- GACATCACAGACAGCCTCATTGA -3'

Serpine1 gene (aka PAI-1) (Eren et al., 2014)

Serpine1-FP: 5′-ACGCCTGGTGCTGGTGAATGC-3′

Serpine1-RP: 5′-ACGGTGCTGCCATCAGACTTGTG-3′

ACKNOWLEDGMENTS

TDP, SLG and CC were partially supported by The Cancer Prevention Institute of Texas

(CPRIT) [RP210227, RP200504], NIH/NCI P30 shared resource grant [CA125123],

NIH/NIEHS center grants [P30 ES030285] and [P42 ES027725], and NIH/NIMHD

[P50MD015496]. DAG supported by NIEHS R01 ES026337.

420
421

FIGURE CAPTIONS

Figure 1. Frequency of NC-XRE motifs in DNA bound by AHR. (**A**) Following TCDD exposure, we analyzed mouse liver AHR ChIP-seq peaks for the presence of known AHR motifs: canonical xenobiotic response element (XRE), non-canonical xenobiotic response element (NC-XRE) or RelB AHR response element (RelBAHRE). Pie-chart shows the percent of AHR binding sites (AHR ChIP peaks) containing one or more of the motifs indicated. A majority of AHR peaks contain canonical and non-canonical xenobiotic response element sequences (XRE+NC-XRE). About a quarter of AHR peaks contain NC-XRE motifs alone. (**B**) Percent of AHR peaks containing XRE or NC-XRE sequences that are associated with promoters, enhancers or indicated genomic feature. A higher ratio of XRE+NC-XRE sites overlap with promoters compared to NC-XRE sites alone. (**C**) Venn diagram showing number of AHR peaks containing XRE, NCXRE, RelBAHRE alone or in combination. No AHR peaks contain RelBAHRE alone, two AHR peaks contain RelBAHRE + XRE. (**D**) Likelihood of finding XRE, NC-XRE or RelBAHRE motifs in AHR ChIP peaks versus in a random region of DNA. Table shows number of AHR peaks containing at least one XRE, NC-XRE or RelBAHRE motif. P value derived from Fisher's exact test.

Figure 2. Repeated NC-XRE motifs associated with AHR target genes. (**A**) Relative frequency of putative AHR target genes within 10kb of AHR ChIP peaks containing 3-10 NC-XRE motifs with 1-50 basepair (bp) spacers, compared to AHR peaks containing 3-10 XRE motifs with similar spacing. AHR ChIP peaks containing 5 NC-XRE motifs separated by 25 basepairs or less were found within 10 kb of 82 target genes. There are 4.9 fold more AHR target genes associated with AHR ChIP peaks containing 5 NCXRE motifs vs 5 XRE motifs separated by 25 bp or less. (**B**) Frequency of repeated runs of NCXRE motifs in AHR ChIP peaks. In 24% of AHR ChIP-seq peaks there are 2 or more NCXRE motifs separated by 25 basepairs or less. (**C**) Binding strength of H3K4me3 and H3K27ac at AHR peaks containing 5 or more NC-XRE or XRE motifs separated by 25 basepairs or less. Runs of NCXRE are enriched at AHR binding sites proximal to transcription start sites, as marked by H3K4me or H3K27ac ChIP peaks. (**D**) Examples of differentially expressed AHR target genes (fold change TCDD vs vehicle, mouse liver) containing AHR ChIP peaks with 5 or more NC-XRE motifs separated by 25 basepairs or less. We define AHR target genes as differentially expressed genes (RNA-seq, fold difference in TCDD vs vehicle > 1.5x, FDR<5%) containing AHR ChIP peak within 10 kb of a gene body.

Figure 3. Transcription factor motifs flanking XRE or NC-XRE motifs in AHR peaks.
Top 12 most frequently occuring known transcription factor motifs in the 50 basepairs flanking repeated NCXRE and XRE motifs (2 or more motifs within 25 basepairs of each other) in genomic regions where AHR binds. Highlighted motifs are enriched flanking NCXRE but not XRE. Significance defined as $p<10^{-11}$

Figure 4. Mutations in NCXRE DNA in *Serpine1* promoter reduced TCDD-dependent expression of *Serpine1*. (**A**) Hepa1-6 cells were exposed to 10 nM TCDD or vehicle for 2 hours, followed by RNA extraction and qPCR for the indicated genes. Gene expression normalized to vehicle, *18S rRNA* used as reference gene. Each circle represents a different biological replicate, horizontal black lines indicate average expression. Dotted line indicates 1x fold change. *Cyp1a1*, *Cyp1b1* and *Ahrr* were used as positive controls since they are known targets of TCDD-AHR. *Cdkn1a*, *Maff* and *Serpine1* exhibit more than 2x increase in expression following TCDD exposure. (**B**) Schematic of mouse *Serpine1* gene showing AHR ChIP peak upstream of transcription start site (TSS). An AHR peak 150 basepairs (bp) upstream of TSS contains NCXRE and no XRE sequences. We transfected Hepa1-6 cells with Cas9 plus guide RNA to generate mutations in these NCXRE motifs, picked single cells and derived a clonal population (clone 14) that completely lacks the NCXRE motifs. We then picked single cells from clone 14 and derived additional clonal populations, designated clone 14-10 and 14-28, with indentical mutations (deletion of the entire NCXRE motifs). (**C**) Hepa1-6 wild-type or mutant cells were exposed to 10 nM TCDD or vehicle for 2 hours, followed by RNA extraction and qPCR for *Serpine1* or *Cyp1a1*. Mutant cell lines lacking the entire NCXRE exhibit reduced increase in *Serpine1* following TCDD exposure compared to wild-type cells. Mutant cells showed similar increase in *Cyp1a1* following TCDD exposure as wild-type cells. One-tailed paired t test, * $p < 0.05$, ** $p < 0.01$, ns not significant ($p > 0.05$).

REFERENCES

Andreasen, E. A., Hahn, M. E., Heideman, W., Peterson, R. E. and Tanguay, R. L. (2002). The zebrafish (Danio rerio) aryl hydrocarbon receptor type 1 is a novel vertebrate receptor. *Mol. Pharmacol.* **62**, 234–249.

Beischlag, T. V. and Perdew, G. H. (2005). ER alpha-AHR-ARNT protein-protein interactions mediate estradiol-dependent transrepression of dioxin-inducible gene transcription. *J. Biol. Chem.* **280**, 21607–21611.

Bernshausen, T., Jux, B., Esser, C., Abel, J. and Fritsche, E. (2006). Tissue distribution and function of the Aryl hydrocarbon receptor repressor (AhRR) in C57BL/6 and Aryl hydrocarbon receptor deficient mice. *Arch. Toxicol.* **80**, 206–211.

Carleton, J. B., Berrett, K. C. and Gertz, J. (2017). Multiplex Enhancer Interference Reveals Collaborative Control of Gene Regulation by Estrogen Receptor α-Bound Enhancers. *Cell Syst.* **5**, 333-344.e5.

Coarfa, C., Grimm, S. L., Katz, T., Zhang, Y., Jangid, R. K., Walker, C. L., Moorthy, B. and Lingappan, K. (2020). Epigenetic response to hyperoxia in the neonatal lung is sexually dimorphic. *Redox Biol.* **37**, 101718.

Darlington, G. J., Bernhard, H. P., Miller, R. A. and Ruddle, F. H. (1980). Expression of liver phenotypes in cultured mouse hepatoma cells. *J Natl Cancer Inst* **64**, 809–819.

Denison, M. S., Fisher, J. M. and Whitlock, J. P. (1988). Inducible, receptor-dependent protein-DNA interactions at a dioxin-responsive transcriptional enhancer. *Proc Natl Acad Sci USA* **85**, 2528–2532.

Denison, M. S., Soshilov, A. A., He, G., DeGroot, D. E. and Zhao, B. (2011). Exactly the same but different: promiscuity and diversity in the molecular mechanisms of action of the aryl hydrocarbon (dioxin) receptor. *Toxicol. Sci.* **124**, 1–22.

Dere, E., Lo, R., Celius, T., Matthews, J. and Zacharewski, T. R. (2011). Integration of genome-wide computation DRE search, AhR ChIP-chip and gene expression analyses of TCDD-elicited responses in the mouse liver. *BMC Genomics* **12**, 365.

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M. and Gingeras, T. R. (2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21.

Eren, M., Boe, A. E., Murphy, S. B., Place, A. T., Nagpal, V., Morales-Nebreda, L., Urich, D., Quaggin, S. E., Budinger, G. R. S., Mutlu, G. M., et al. (2014). PAI-1-regulated extracellular proteolysis governs senescence and survival in Klotho mice. *Proc Natl Acad Sci USA* **111**, 7090–7095.

Fader, K. A., Nault, R., Kirby, M. P., Markous, G., Matthews, J. and Zacharewski, T. R. (2017). Convergence of hepcidin deficiency, systemic iron overloading, heme accumulation, and REV-ERBα/β activation in aryl hydrocarbon receptor-elicited hepatotoxicity. *Toxicol. Appl. Pharmacol.* **321**, 1–17.

Fader, K. A., Nault, R., Doskey, C. M., Fling, R. R. and Zacharewski, T. R. (2019). 2,3,7,8-Tetrachlorodibenzo-p-dioxin abolishes circadian regulation of hepatic metabolic activity in mice. *Sci. Rep.* **9**, 6514.

Hankinson, O. (2005). Role of coactivators in transcriptional activation by the aryl hydrocarbon receptor. *Arch. Biochem. Biophys.* **433**, 379–386.

Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y. C., Laslo, P., Cheng, J. X., Murre, C., Singh, H. and Glass, C. K. (2010). Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol. Cell* **38**, 576–589.

Hirose, K., Morita, M., Ema, M., Mimura, J., Hamada, H., Fujii, H., Saijo, Y., Gotoh, O., Sogawa, K. and Fujii-Kuriyama, Y. (1996). cDNA cloning and tissue-specific expression of a novel basic helix-loop-helix/PAS factor (Arnt2) with close sequence similarity to the aryl hydrocarbon receptor nuclear translocator (Arnt). *Mol. Cell. Biol.* **16**, 1706–1713.

Huang, G. and Elferink, C. J. (2012). A novel nonconsensus xenobiotic response element capable of mediating aryl hydrocarbon receptor-dependent gene expression. *Mol. Pharmacol.* **81**, 338–347.

Isoda, K., Sawada, S., Ayaori, M., Matsuki, T., Horai, R., Kagata, Y., Miyazaki, K., Kusuhara, M., Okazaki, M., Matsubara, O., et al. (2005). Deficiency of interleukin-1 receptor antagonist deteriorates fatty liver and cholesterol metabolism in hypercholesterolemic mice. *J. Biol. Chem.* **280**, 7002–7009.

Jackson, D. P., Li, H., Mitchell, K. A., Joshi, A. D. and Elferink, C. J. (2014). Ah receptor-mediated suppression of liver regeneration through NC-XRE-driven p21Cip1 expression. *Mol. Pharmacol.* **85**, 533–541.

Joshi, A. D., Mustafa, M. G., Lichti, C. F. and Elferink, C. J. (2015). Homocitrullination is a novel histone H1 epigenetic mark dependent on aryl hydrocarbon receptor recruitment of carbamoyl phosphate synthase 1. *J. Biol. Chem.* **290**, 27767–27778.

Klein-Hitpass, L., Kaling, M. and Ryffel, G. U. (1988). Synergism of closely adjacent estrogen-responsive elements increases their regulatory potential. *J. Mol. Biol.* **201**, 537–544.

Klinge, C. M., Kaur, K. and Swanson, H. I. (2000). The aryl hydrocarbon receptor
interacts with estrogen receptor alpha and orphan receptors COUP-TFI and
ERRalpha1. *Arch. Biochem. Biophys.* **373**, 163–174.

Labun, K., Montague, T. G., Krause, M., Torres Cleuren, Y. N., Tjeldnes, H. and
Valen, E. (2019). CHOPCHOP v3: expanding the CRISPR web toolbox beyond
genome editing. *Nucleic Acids Res.* **47**, W171–W174.

Langmead, B. and Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2.
Nat. Methods **9**, 357–359.

Liao, Y., Smyth, G. K. and Shi, W. (2014). featureCounts: an efficient general purpose
program for assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–
930.

Liu, X., Huang, T., Li, L., Tang, Y., Tian, Y., Wang, S. and Fan, C. (2015). CYP1B1
deficiency ameliorates obesity and glucose intolerance induced by high fat diet in
adult C57BL/6J mice. *Am. J. Transl. Res.* **7**, 761–771.

Lopes, R., Korkmaz, G. and Agami, R. (2016). Applying CRISPR-Cas9 tools to
identify and characterize transcriptional enhancers. *Nat. Rev. Mol. Cell Biol.* **17**, 597–
604.

Lo, R. and Matthews, J. (2012). High-resolution genome-wide mapping of AHR and
ARNT binding sites by ChIP-Seq. *Toxicol. Sci.* **130**, 349–361.

Madak-Erdogan, Z. and Katzenellenbogen, B. S. (2012). Aryl hydrocarbon receptor
modulation of estrogen receptor α-mediated gene regulation by a multimeric
chromatin complex involving the two receptors and the coregulator RIP140. *Toxicol.*
Sci. **125**, 401–411.

Nault, R., Fader, K. A. and Zacharewski, T. (2015). RNA-Seq versus oligonucleotide
array assessment of dose-dependent TCDD-elicited hepatic gene expression in mice.
BMC Genomics **16**, 373.

Nault, R., Fader, K. A., Kirby, M. P., Ahmed, S., Matthews, J., Jones, A. D., Lunt, S.
Y. and Zacharewski, T. R. (2016). Pyruvate Kinase Isoform Switching and Hepatic
Metabolic Reprogramming by the Environmental Contaminant 2,3,7,8-
Tetrachlorodibenzo-p-Dioxin. *Toxicol. Sci.* **149**, 358–371.

Nault, R., Doskey, C. M., Fader, K. A., Rockwell, C. E. and Zacharewski, T. (2018).
Comparison of Hepatic NRF2 and Aryl Hydrocarbon Receptor Binding in 2,3,7,8-
Tetrachlorodibenzo-p-dioxin-Treated Mice Demonstrates NRF2-Independent PKM2
Induction. *Mol. Pharmacol.* **94**, 876–884.

Niu, B., He, K., Li, P., Gong, J., Zhu, X., Ye, S., Ou, Z. and Ren, G. (2018). SIRT1
upregulation protects against liver injury induced by a HFD through inhibiting CD36
and the NF-κB pathway in mouse kupffer cells. *Mol. Med. Report.* **18**, 1609–1615.

Quinlan, A. R. and Hall, I. M. (2010). BEDTools: a flexible suite of utilities for
comparing genomic features. *Bioinformatics* **26**, 841–842.

Reyes, H., Reisz-Porszasz, S. and Hankinson, O. (1992). Identification of the Ah
receptor nuclear translocator protein (Arnt) as a component of the DNA binding form
of the Ah receptor. *Science* **256**, 1193–1195.

Risso, D., Ngai, J., Speed, T. P. and Dudoit, S. (2014). Normalization of RNA-seq
data using factor analysis of control genes or samples. *Nat. Biotechnol.* **32**, 896–902.

Robinson, M. D., McCarthy, D. J. and Smyth, G. K. (2010). edgeR: a Bioconductor
package for differential expression analysis of digital gene expression data.
Bioinformatics **26**, 139–140.

Rowatt, A. J., DePowell, J. J. and Powell, W. H. (2003). ARNT gene multiplicity in
amphibians: characterization of ARNT2 from the frog Xenopus laevis. *J. Exp. Zool. B*
Mol. Dev. Evol. **300**, 48–57.

Tanguay, R. L., Abnet, C. C., Heideman, W. and Peterson, R. E. (1999). Cloning and
characterization of the zebrafish (Danio rerio) aryl hydrocarbon receptor. *Biochim.*
Biophys. Acta **1444**, 35–48.

Tanguay, R. L., Andreasen, E., Heideman, W. and Peterson, R. E. (2000).
Identification and expression of alternatively spliced aryl hydrocarbon nuclear
translocator 2 (ARNT2) cDNAs from zebrafish with distinct functions. *Biochim.*
Biophys. Acta **1494**, 117–128.

Turner, N., Lim, X. Y., Toop, H. D., Osborne, B., Brandon, A. E., Taylor, E. N.,
Fiveash, C. E., Govindaraju, H., Teo, J. D., McEwen, H. P., et al. (2018). A
selective inhibitor of ceramide synthase 1 reveals a novel role in fat metabolism. *Nat.*
Commun. **9**, 3165.

Wilson, S. R., Joshi, A. D. and Elferink, C. J. (2013). The tumor suppressor Kruppel-
like factor 6 is a novel aryl hydrocarbon receptor DNA binding partner. *J. Pharmacol.*
Exp. Ther. **345**, 419–429.

Yao, E. F. and Denison, M. S. (1992). DNA sequence determinants for binding of
transformed Ah receptor to a dioxin-responsive enhancer. *Biochemistry* **31**, 5060–
5067.

Yeager, R. L., Reisman, S. A., Aleksunes, L. M. and Klaassen, C. D. (2009).
Introducing the "TCDD-inducible AhR-Nrf2 gene battery". *Toxicol. Sci.* **111**, 238–246.

Yue, F., Cheng, Y., Breschi, A., Vierstra, J., Wu, W., Ryba, T., Sandstrom, R., Ma,
Z., Davis, C., Pope, B. D., et al. (2014). A comparative encyclopedia of DNA
elements in the mouse genome. *Nature* **515**, 355–364.

Zhang, Y., Liu, T., Meyer, C. A., Eeckhoute, J., Johnson, D. S., Bernstein, B. E.,
Nusbaum, C., Myers, R. M., Brown, M., Li, W., et al. (2008). Model-based analysis
of ChIP-Seq (MACS). *Genome Biol.* **9**, R137.

GEO accession	PMID	Assay	Sex	Conditions
GSE97634	28213091, 31015483	ChIP-seq	male C57BL/6 liver	exposed to 30 ug/kg TCDD via oral gavage for 2 hours, n=5 mice
GSE97636	26582802	ChIP-seq	female C57BL/6 liver	exposed to 30 ug/kg TCDD via oral gavage for 2 hours, n=5 mice
GSE62902	25958198	RNA-seq	female C57BL/6 liver	exposed to 0.01, 0.03, 0.1, 0.3, 1, 3, 10, or 30 ug/kg TCDD or sesame oil vehicle via oral gavage, every 4 days for 28 days
GSE87519	28213091	RNA-seq	male C57BL/6 liver	exposed to 0.01, 0.03, 0.1, 0.3, 1, 3, 10, or 30 ug/kg TCDD or sesame oil vehicle via oral gavage, every 4 days for 28 days. At 7, 15, and 23d after the first dose, mice (vehicle and 30 µg/kg TCDD groups) were fasted for 6 hours (access to water but not food). At 22d after the first dose, oral glucose tolerance tests (OGTT) were performed (vehicle and 30 µg/kg TCDD groups; fasted for 6h). Briefly, at time 0 minutes (min) animals were orally gavaged with 2 g/kg glucose in a 25% solution and tail blood glucose was measured at 0, 5, 15, 25, 30, 60, and 120 min. At 26d after the first dose, mice (vehicle and 30 µg/kg TCDD groups) were transferred to Innocages lacking bedding for 2h, with access to water but not food.
GSE109863	29752288	RNA-seq	male C57BL/6 liver	On postnatal day 28 mice were orally gavaged with sesame oil vehicle or 30 μg/kg TCDD for 2, 4, 8, 12, 24, 72, and 168 hours

Table 1. Datasets analyzed in this manuscript

Figure 1. Frequency of NC-XRE motifs in DNA bound by AHR. (**A**) Following TCDD exposure, we analyzed mouse liver AHR ChIP-seq peaks for the presence of known AHR motifs: canonical xenobiotic response element (XRE), non-canonical xenobiotic response element (NC-XRE) or RelB AHR response element (RelBAHRE). Pie-chart shows the percent of AHR binding sites (AHR ChIP peaks) containing one or more of the motifs indicated. A majority of AHR peaks contain canonical and non-canonical xenobiotic response element sequences (XRE+NC-XRE). About a quarter of AHR peaks contain NC-XRE motifs alone. (**B**) Percent of AHR peaks containing XRE or NC-XRE sequences that are associated with promoters, enhancers or indicated genomic feature. A higher ratio of XRE+NC-XRE sites overlap with promoters compared to NC-XRE sites alone. (**C**) Venn diagram showing number of AHR peaks containing XRE, NCXRE, RelBAHRE alone or in combination. No AHR peaks contain RelBAHRE alone, two AHR peaks contain RelBAHRE + XRE. (**D**) Likelihood of finding XRE, NC-XRE or RelBAHRE motifs in AHR ChIP peaks versus in a random region of DNA. Table shows number of AHR peaks containing at least one XRE, NC-XRE or RelBAHRE motif. P value derived from Fisher's exact test.

Figure 2. Repeated NC-XRE motifs associated with AHR target genes. (**A**) Relative frequency of putative AHR target genes within 10kb of AHR ChIP peaks containing 3-10 NC-XRE motifs with 1-50 basepair (bp) spacers, compared to AHR peaks containing 3-10 XRE motifs with similar spacing. AHR ChIP peaks containing 5 NC-XRE motifs separated by 25 basepairs or less were found within 10 kb of 82 target genes. There are 4.9 fold more AHR target genes associated with AHR ChIP peaks containing 5 NCXRE motifs vs 5 XRE motifs separated by 25 bp or less. (**B**) Frequency of repeated runs of NCXRE motifs in AHR ChIP peaks. In 24% of AHR ChIP-seq peaks there are 2 or more NCXRE motifs separated by 25 basepairs or less. (**C**) Binding strength of H3K4me3 and H3K27ac at AHR peaks containing 5 or more NC-XRE or XRE motifs separated by 25 basepairs or less. Runs of NCXRE are enriched at AHR binding sites proximal to transcription start sites, as marked by H3K4me or H3K27ac ChIP peaks. (**D**) Examples of differentially expressed AHR target genes (fold change TCDD vs vehicle, mouse liver) containing AHR ChIP peaks with 5 or more NC-XRE motifs separated by 25 basepairs or less. We define AHR target genes as differentially expressed genes (RNA-seq, fold difference in TCDD vs vehicle > 1.5x, FDR<5%) containing AHR ChIP peak within 10 kb of a gene body.

Motif name	Motif sequence	Percent of AHR peaks with flanking motif & NCXRE	Percent of AHR peaks with flanking motif & XRE	NCXRE -log10(p value)	XRE -log10(p value)
Nr1a2	TRAGGTCA	50.62	46.52	215	71
AR half-site	CCAGGAACAG	44.61	0	27	not significant
RARa	TTGAMCTTTG	40.3	36.23	418	158
Arnt:Ahr	TBGCACGCAA	36.39	67.08	3704	5093
Tgif2	TGTCANYT	33.88	0	36	not significant
ERRa	CAAAGGTCAG	33.42	30.63	307	112
Smad3	TWGTCTGV	31.28	0	17	not significant
Hif1b	RTACGTGC	30.78	50.69	1036	1566
Tgif1	YTGWCADY	30.33	0	15	not significant
Klf14	RGKGGGCGK-GGC	29.25	32.36	25	39
Hic1	TGCCAGCB	27.12	0	26	not significant
COUP-TFII	AGRGGTCA	26.32	23.42	365	119

Figure 3. Transcription factor motifs flanking XRE or NC-XRE motifs in AHR peaks. Top 12 most frequently occuring known transcription factor motifs in the 50 basepairs flanking repeated NCXRE and XRE motifs (2 or more motifs within 25 basepairs of each other) in genomic regions where AHR binds. Highlighted motifs are enriched flanking NCXRE but not XRE. Significance defined as $p<10^{-11}$

A

B

C

Figure 4. Mutations in NCXRE DNA in *Serpine1* promoter reduced TCDD-dependent expression of *Serpine1*. (**A**) Hepa1-6 cells were exposed to 10 nM TCDD or vehicle for 2 hours, followed by RNA extraction and qPCR for the indicated genes. Gene expression normalized to vehicle, *18S rRNA* used as reference gene. Each circle represents a different biological replicate, horizontal black lines indicate average expression. Dotted line indicates 1x fold change. *Cyp1a1*, *Cyp1b1* and *Ahrr* were used as positive controls since they are known targets of TCDD-AHR. *Cdkn1a*, *Maff* and *Serpine1* exhibit more than 2x increase in expression following TCDD exposure. (**B**) Schematic of mouse *Serpine1* gene showing AHR ChIP peak upstream of transcription start site (TSS). An AHR peak 150 basepairs (bp) upstream of TSS contains NCXRE and no XRE sequences. We transfected Hepa1-6 cells with Cas9 plus guide RNA to generate mutations in these NCXRE motifs, picked single cells and derived a clonal population (clone 14) that completely lacks the NCXRE motifs. We then picked single cells from clone 14 and derived additional clonal populations, designated clone 14-10 and 14-28, with indentical mutations (deletion of the entire NCXRE motifs). (**C**) Hepa1-6 wild-type or mutant cells were exposed to 10 nM TCDD or vehicle for 2 hours, followed by RNA extraction and qPCR for *Serpine1* or *Cyp1a1*. Mutant cell lines lacking the entire NCXRE exhibit reduced increase in *Serpine1* following TCDD exposure compared to wild-type cells. Mutant cells showed similar increase in *Cyp1a1* following TCDD exposure as wild-type cells. One-tailed paired t test, * p < 0.05, ** p < 0.01, ns not significant (p > 0.05).