Title:

Boosting Live Malaria Vaccine with Cytomegalovirus Vector Can Prolong Immunity

through Innate and Adaptive Mechanisms

Authors: Komi Gbedande[1, 2 #a], Samad A. Ibitokou[1, #b], Monique L. Ong[3, 4], Mariapia A. Degli-

Esposti[3, 4], Michael G. Brown[5], Robin Stephens[1, 2, 6, *, #a]

Affiliation:

[1] Department of Internal Medicine, Division of Infectious Diseases, University of Texas Medical

Branch, Galveston, TX 77555-0435

[2] Center for Immunity and Inflammation, Rutgers New Jersey Medical School, Cancer Center,

205 S. Orange Avenue, Newark, NJ 07103

[3] Centre for Experimental Immunology, Lions Eye Institute; Nedlands, Western Australia,

Australia

[4] Infection and Immunity Program and Department of Microbiology, Biomedicine Discovery

Institute, Monash University; Clayton, Victoria, Australia

[5] Department of Medicine, Division of Nephrology, and the Beirne B. Carter Center for

Immunology Research, University of Virginia, Charlottesville, VA

[6] Department of Microbiology and Immunology, University of Texas Medical Branch,

Galveston, TX

[*] **Corresponding author**

[#a] Current Address: RS, KG: Center for Immunity and Inflammation, Rutgers New Jersey Medical

School, Cancer Center, 205 S. Orange Avenue, G-1230; Newark, NJ 07103;

#b Current Address SAI: Fate Therapeutics, 12278 Scripps Summit Drive, San Diego, CA 92131

Short Title: MCMV Boost Prolongs Malaria Vaccine Protection and Effector T cells

Abstract

Vaccines to persistent parasite infections have been challenging, and current iterations lack

long-term protection. Cytomegalovirus (CMV) chronic vaccine vectors drive protection against

SIV, tuberculosis and liver-stage malaria correlated with antigen-specific CD8 T cells with a Tem

phenotype. This phenotype is likely driven by a combination of antigen-specific and innate

adjuvanting effects of the vector, though these mechanisms are less well understood. Sterilizing

immunity from live *Plasmodium chabaudi* vaccination lasts less than 200 days. While *P.*

chabaudi-specific antibody levels remain stable after vaccination, the decay of parasite-specific T

cells correlates with loss of challenge protection. Therefore, we enlisted murine CMV as a booster

strategy to prolong T cell responses against malaria. To study induced T cell responses, we

included *P. chabaudi* MSP-1 epitope B5 (MCMV-B5). We found that MCMV vector alone

significantly protected against a challenge *P. chabaudi* infection 40-60 days later, and that

MCMV-B5 was able to make B5-specific Teff, in addition to previously-reported Tem, that

survive to the challenge timepoint. Used as a booster, MCMV-B5 prolonged protection from

heterologous infection beyond day 200, and increased B5 TCR Tg T cell numbers, including both

a highly-differentiated Tem phenotype and Teff, both previously reported to protect. B5 epitope

expression was responsible for maintenance of Th1 and Tfh B5 T cells. In addition, the MCMV

vector had adjuvant properties, contributing non-specifically through prolonged stimulation of

IFN-γ. *In vivo* neutralization of IFN-γ, but not IL-12 and IL-18, late in the course of MCMV, led

to loss of the adjuvant effect. Mechanistically, sustained IFN-γ from MCMV increased CD8α⁺

dendritic cell numbers, and led to increased IL-12 production upon *Plasmodium* challenge. In

addition, neutralization of IFN-γ before challenge reduced the polyclonal Teff response to

challenge. Our findings suggest that, as protective epitopes are defined, an MCMV vectored

booster can prolong protection through the innate effects of IFN-γ.

Authors' summary

Malaria represents a challenging target for vaccination. This is in part because of the

requirement for CD4 T cell immunity in addition to the standard B cell responses current vaccines

induce. However, human malaria vaccine approaches thus far have limited longevity of protection

due to decay of T cell responses. This includes the most advanced malaria vaccine, a virus-like

particle expressing one recombinant liver-stage antigen (RTS,S), and liver-stage parasites

attenuated by radiation (PfSPZ), as well as live vaccination with drug-treatment. Our work seeks

to prolong this protection using MCMV, a promising vaccine vector known to promote CD8 T cell

responses. We observed that boosting the live malaria vaccine with MCMV including a

Plasmodium antigen led to longer protection from *P. chabaudi* parasitemia, and can be used to

promote maintenance of antigen-specific CD4 T cells. In investigating the mechanisms of the

MCMV booster, we found that the cytokine IFN-γ is required for prolonged protection and

enhances priming of the innate immune system for prolonged protection from malaria. Our

research informs both the quest for a longer-lived malaria vaccine and that to understand

mechanisms of protection from persistent infection.

Introduction

Compared to acute and localized infections, persistent systemic infections are the most challenging

vaccines to develop [1, 2]. Despite impressive progress for malaria vaccine development, and

using emerging approaches including live or attenuated parasites, and recombinant protein

constructs targeting different stages in the parasite life cycle, none of the current vaccine

candidates provides both highly effective and long-lived immunity. Both natural parasite infection

and malaria vaccines generate immunity to parasitemia on re-infection that wanes quickly, and

this is shown to be due to a progressive loss of specific T cells [3, 4]. Key insights to decay of

protection from persistent systemic infections has come from the study of immunity to parasite

infections with *Leishmania* and *Plasmodium spp.,* which decays over time in the absence of

infection [3, 5, 6]. Protection from recurrent infection in nature is likely mediated by sustained

generation of otherwise short-lived Th1 Teff while infection persists [7, 8]. We and others have

shown that both short-lived effector T cells and long-lived effector memory T cells are generated

in persistent infection. It is unclear if long-lived memory T cells (Tmem) play an important role,

or could be used to improve protection from potentially persistent systemic infections. Persistent

parasite infection, including *Leishmania major* or *P. chabaudi,* generate recirculating Tcm that do

not protect as well as activated Teff or Tem, defined as previously divided $CD4^+CD62L^{lo}$ Tem/Teff

[9], or distinguished as $CD127^-$ Teff and $CD127^{hi}CD44^{hi}CD62L^{lo}CD27^-$ Tem [10]. In *P. chabaudi*

infection, we showed that inducing a pre-activated MSP1-specific B5 T cell receptor transgenic

(TCR Tg) T cell response, along with a B cell response to MSP-1 p19, before infection with *P.*

chabaudi, was an effective way to promote an effective reduction of the parasitemia of re-infection.

These data suggest an important role for effector T cells in protection. It should be noted that while

B5 T cells protect immunocompromised recipients from death, they do not eliminate parasite in

the absence of B cells [11]. By defining Teff as $CD127^-$, an early marker of T cell activation that

is transient and not present on Tem, we were able to determine that adoptively transferred Teff are

the most protective T cells in systemic, blood-stage *P. chabaudi* infection [8, 10]. However, short-

lived CD4 T cell immunity has been identified as the crucial variable resulting in short-term

protection from persistent *P. chabaudi* infection or vaccination [3, 12]. However, standard

vaccines do not generate durable Teff responses, leading to the problem at hand of what vaccine

vector to use to prolong protection, if the cell type that protects is short-lived.

Therefore, we investigated the potential of CMV to increase the duration of the T cell

response to effectively prolong vaccine-induced immunity to *Plasmodium* infection as a booster

vector. CMV has emerged as a promising long-lasting vaccine vector, for example, it provided

lasting immunity over a year against mucosal Simian Immunodeficiency virus (SIV) challenge in

Rhesus macaques [13]. CMV is a potent CD8 T cell-inducing vector with efficacy against SIV,

ebolavirus, tuberculosis, cancer and liver-stage malaria [14-18]. CMV infection can induce

sustained, or inflationary CD8 T cell responses with an effector memory phenotype. Th1-type

effector function is also generated [19-21], but the effect of persistent CMV on CD4 T cells is less

well understood. CMV also has non-specific effects suggesting a useful adjuvant effect of the

vector. For example, MCMV provides non-specific protection to *Listeria monocytogenes*, *Yersinia*

pestis and influenza virus, particularly in the latent phase of infection [22, 23]. Previous vaccine

studies all have used CMV in the priming phase of the vaccine response, and it has not been tested

as a vector for boosting existing responses, nor has it been studied extensively in the generation of

CD4 T cell responses. In addition, CMV generates effector/memory T cells, but the specific

phenotype has not been determined, and extending the duration of an effector response will be

critical to the fight against persistent infection.

In this study, we found that the MCMV vector boosted protection generated by a live

malaria vaccine against challenge infection with heterologous parasite. In addition to showing that

we can induce epitiope-specific effector T cells 40-60 days post-boost, we identified a significant

adjuvanting effect of boosting with MCMV. We showed that IFN-γ produced during the persistent

phase of MCMV increases CD8α$^+$ dendritic cell numbers. Upon challenge, MCMV boosted

animals had increased serum IL-12 and a larger CD4 polyclonal effector T cell response, which

were reduced by neutralization of IFN-γ before infection. This work informs our basic

understanding of the immunology and vaccinology of persistent infections, and suggests that

persistent stimulation of innate and adaptive mechanisms can promote persistence of protective

immunity.

Results

MCMV as a vaccine vector with adjuvant properties against *P. chabaudi*

In order to evaluate an MCMV vaccine vector and observe vector- and *Plasmodium*-

specific CD4 T cell phenotypes, we generated an MCMV vector carrying the *P. chabaudi* MSP-1

B5 CD4 T cell epitope (MCMV-B5) using bacterial artificial chromosome (BAC) recombineering.

The B5 epitope was inserted in the intermediate early 1 locus with a linker to facilitate processing,

as diagrammed in **S1A Fig**. As MCMV has been shown to induce more sustained CD4 T cell

numbers in BALB/c, than in C57BL/6 mice [24], we developed a BALB/c system. Although *P.*

chabaudi MSP-1 B5 CD4$^+$ T cells prevent immunodeficient animals from dying, B5 T cells alone

do not affect *P. chabaudi* parasitemia significantly unless they are pre-activated, and in the

presence of parasite-specific B cells [11]. Several aspects of the virology and immunology of the

two constructs were compared for rigor of the study. The kinetics and tissue tropism of MCMV-

B5 infection was confirmed to be the same as the parental virus, MCMV-BAC using plaque assay

(**S1B Fig**) and PCR (**S1C Fig**). The virus is largely cleared by day 30, however low recrudescence

can be seen at later timepoints. Proportions of polyclonal CD4$^+$CD127$^-$ Teff, and CD4$^+$

CD44^{hi}CD127^{hi} Tmem were similar for MCMV-BAC and MCMV-B5 at day 15 p.i. (**S1D Fig)**.

MCMV-specific CD4$^+$ T cells were identified using m78 peptide/I-A^d tetramers, using CLIP/I-A^d

as non-specific tetramer control, and MCMV-specific CD8$^+$ T cells were detected with the m45/D^d

-tetramer [20, 21]. Both MCMV-BAC and MCMV-B5 induced an MCMV m78-specific CD4 T

cell response at day 15 p.i. (**S1E Fig**). A large MCMV m45-specific CD8 T cell response to

MCMV-B5 was sustained near a million T cells per spleen from day 11 to 60 p.i. (**S1F Fig**).

MCMV is an excellent vaccine vector that showed early promise against liver-stage

malaria [16] and other infections fought by CD8 T cells. However, the mechanisms by which the

vector contributes to the adjuvanting protective effect are not well understood. To test for a

protective effect of the low chronic phase of MCMV infection on *P. chabaudi* challenge,

BALB/cByJ mice were infected with MCMV-B5 or MCMV-BAC or aged, and challenged 60 days

later with *P. chabaudi* AS, as shown schematically in **Fig 1A**. Groups given either MCMV-BAC

or MCMV-B5 had strongly reduced parasite growth upon *P. chabaudi* infection compared to a

primary infection age-matched control (**Fig 1B**). Pathology from malaria, as indicated by weight

loss, was also significantly inhibited by prior MCMV-BAC or MCMV-B5 infection (**Fig 1C**). This

strong, non-specific effect reduced parasitemia and pathology by about half, so not as effectively

as prior infection, or live vaccination [3].

MCMV stimulates and maintains *Plasmodium*-specific T cell responses

159  Using B5 TCR Tg T cells, we were able to distinguish the non-specific adjuvant effects

160 from epitope-specific effects on CD4$^+$ T cells [11]. To determine effects of the MCMV vector on

161 epitope-specific T cells, we adoptively transferred B5 Tg T cells (2 x 10^6) and infected the

162 recipients with MCMV-BAC or MCMV-B5, as illustrated in **Fig 2A**. The B5 CD4 T cell response

163 in the spleen 15 and 55 days post-MCMV infection was assessed by flow cytometry. The

164 proliferated B5 T cell (CTV$^-$ Thy1.2$^+$) number was specifically increased by MCMV-B5, which

165 leading to a specific increase that lasted until day 55 (**Fig 2B**). MCMV-BAC infected mice also

166 maintained proliferated CTV$^-$ B5 T cell numbers slightly above background to day 55. In order to

167 test the MCMV-driven generation of effector T cells, they were stained for the IL-7Rα (CD127),

168 which is downregulated on recently activated T cells. The majority of CTV$^-$ B5 TCR Tg T cells

169 on day 15 are CD127$^-$ Teff (**Fig 2C**). On day 55, the fraction of Tmem increases, but a significant

170 number of Teff also are present, as clearly seen in the flow cytometry plot of B5 T cells recovered

171 at this timepoint (**Fig 2D**). In addition, we observed that MCMV-B5, but not MCMV-BAC,

172 specifically promoted the differentiation of MCMV-B5 epitope-specific GC Tfh (CXCR5^{hi}PD-1^{hi},

173 **Fig 2E**).

174  We also detected some MCMV-specific CD4 T cell responses using MCMV m78 tetramer

175 at 11 and 60 days after MCMV-B5 infection (S2A **Fig**). MCMV-B5 infection induced MCMV

176 m78-specific CD4$^+$ T cell responses that persisted for two months (S2B **Fig**). Phenotypically,

177 MCMV vector-specific T cells were predominantly Teff (Tet$^+$ CD127$^-$) early in infection (day 11),

178 with re-upregulation of CD127 into a Tmem phenotype at day 60 (S2C **Fig**), similar to the B5-

179 specific response to MCMV-B5. Proliferation of transiently transferred B5 T cells in response to

180 MCMV-B5 was tested over various five-day windows. Congenic recipients (Thy1.1) were infected

181 with MCMV-BAC or MCMV-B5 or uninfected, and CTV-labelled CD4-purified B5 TCR Tg T

cells were adoptively transferred for the last five days of infection, testing presentation of B5

peptide between days 5-10, 25-30, or 55-60 of MCMV infection. Splenocytes were analyzed for

proliferation (%CTV-) of B5 T cells in each experimental window (**S2D Fig**). The MCMV-B5

vector, and not the MCMV-BAC control, led to strong specific proliferation of epitope-specific

B5 Tg T cells during acute infection (days 5-10) (**S2E Fig**). Despite the presence of B5 TCR Tg

Teff on day 55, there was no detectable proliferation in this assay at later timepoints. This may be

due to background proliferation, even in uninfected animals.

MCMV boosts long-term protection of live malaria vaccine

As the protection of malaria vaccines tested in mice and humans decays after one to two

years [25, 26], we sought to test a vaccine booster strategy to prolong immunity. *Plasmodium spp*.

infection followed by drug treatment is an effective vaccination protocol in C57BL/6 mice, and

also in human studies; however, this protection shows measurable decay by day 200 in mice [3,

27]. Using the protocol published by Freitas do Rosario et al, where mice are infected with *P.*

chabaudi AS followed by sub-curative dose of chloroquine each time parasitemia reaches 1%, we

tested sterilizing immunity with heterologous reinfection (**S3A Fig**). This experiment confirmed

that *P. chabaudi* infection induced sterilizing immunity for two months, and that a measurable

decay in protection was evident by day 200 in BALB/c mice (**S3B Fig**), as in B6. We also

confirmed that there was not a measurable decay in antibody concentration between 2 and 6.5

months p.i., as shown previously in B6 mice (**S3C Fig**), leading us to test a known T cell-response-

promoting vector to attempt to prolong T cell responses and protection through as yet unknown

adjuvant effects.

To test the potential of an MCMV-based booster vaccine, mice were vaccinated with live

P. chabaudi AS infection followed by sub-curative doses of chloroquine and then boosted with

MCMV-BAC or MCMV-B5 140 days later, as schematically depicted in **Fig 3A**. As vaccine-

induced protection decays by day 200, we tested MCMV boosters for protection from heterologous

re-infection with *P. chabaudi* AJ at day 200, which corresponds to 60 days after boosting. This

timing was chosen on the basis that neither immunity nor the T cell response to live vaccination

has completely decayed by day 120 [3, 12], and that the MCMV infection is below detectable by

day 60 [24], though we expected it to show prolonged stimulatory capacity at this timepoint [22].

Age- and sex-matched *Plasmodium*-vaccinated mice, with or without the MCMV-booster, were

challenged with heterologous *Plasmodium* infection. The booster prolonged live vaccine induced

protection, as evidenced by the data shown in **Fig 3B** that mice that were boosted with MCMV

had a lower parasitemia on challenge on day 200 post-vaccination (p.v., average of 0.06%, or

0.05% without or with the B5 epitope, respectively), on day 4 post-challenge (p.c.), than

vaccinated mice challenged without boosting (1.57% on day 4 p.c., and 0.5% on day 6 p.c.). The

boosting effect does not require the B5 epitope in a significant way, which is not surprising, B5

CD4 T cell epitope was included as a way to track specific T cells, not to protect as a single epitope.

Heterologous parasite-specific serum antibody was measured by ELISA at day 7 p.c. (**Fig 3C**). As

previously reported parasite-specific IgG responses remain stable up to day 200 post-live

vaccination [3], ELISA showed that levels of *P. chabaudi* AJ-specific serum IgG in the first week

of challenge were not altered by the MCMV booster (**Fig 3C**). The boost was specifically chosen

to promote T cell responses to support the ongoing antibody response generated by the live

vaccine. In summary, *Plasmodium* vaccination efficacy, but not serum antibody, fades with time,

and boosting with MCMV prolonged protection.

MCMV can boost parasite-specific T cell immunity induced by live malaria vaccine

CMV is a promising CD8 T cell-inducing vaccine vector and generates Tem and inflationary effector T cells [20, 28]. Less is known about CD4 T cell phenotypes in response to MCMV [20], and the potential for continuous generation of phenotypic effector CD4 T cells in the persistent phase is not well understood. In order to determine if MCMV-B5 boosted *Plasmodium*-specific CD4 T cell responses, B5 CD4 TCR Tg cells (Cell Trace Violet (CTV)-labelled) were adoptively transferred into congenic recipients (Thy1.1 BALB/cByJ), which were then immunized with live *P. chabaudi* vaccine, and boosted with MCMV-B5 or MCMV-BAC at day 140 p.v. (**Fig 4A**). B5 T cell effector and effector memory phenotype and cytokine responses were quantified at day 200 p.v.. Using MCMV-B5, as a booster to the live malaria vaccine specifically and significantly increased the numbers of divided B5 TCR Tg T cells ($CD4^+CTV^-Thy1.2^+$) present on day 200 p.v., while MCMV-BAC did not (**Fig 4B**). Therefore, more transferred B5 T cells survived to day 200 after MCMV-B5 boost. B5 TCR Tg cells are not detectable after 200 days in the absence of parasite [8]. The MCMV-B5 boost 60 days earlier specifically increased the numbers of both $CD127^-$ B5 Teff and $CD127^{hi}$ B5 Tmem present at day 200, while MCMV-BAC did not increase B5 T cell numbers (**Fig 4C**). Among Tmem subsets, the number of both $CD27^+$ and $CD27^-$ effector memory T cells ($CD44^{hi}CD127^{hi}CD62L^{lo}$, Tem) were significantly higher in MCMV-B5 boosted mice, while central memory ($CD62L^{hi}$ Tcm) were not increased by MCMV boost (**Fig 4D**). In addition, the MCMV-B5 booster specifically generated or maintained functional B5 T helper type 1-type cells, as shown by their production of IFN-γ (**Fig 4E**), TNF (**Fig 4F**) and IL-2 (**Fig 4G**) at day 200. This data show that the vector generated well-presented epitopes, and

could be used as a booster vaccine for T cells. In addition, the vector induced effector T cells for

a longer time than reported for acute stimulation, in addition to making Tem.

IFN-γ, but not IL-12 and IL-18, induced by MCMV, is associated with protection from *P.*

***chabaudi* infection**

As we had observed a protective effect by the MCMV-BAC, we took the opportunity to

understand mechanisms explaining the observed benefits of using this virus as a vaccine vector in

addition to the ability to present T cell epitopes and generate Teff. To determine if cytokines

induced by long-term MCMV infection are mechanistically responsible for promoting protection

from heterologous challenge, we used neutralizing antibodies to test the roles of IFN-γ, as well as

the combination of IL-12 and IL-18, in the period before challenge after 40 days of MCMV

infection. IFN-γ was neutralized starting on day 34 of MCMV-BAC infection, and before *P.*

chabaudi challenge, as shown schematically in **Fig 5A**. *In vivo* neutralization of IFN-γ after

MCMV, but before *P. chabaudi* infection was associated with a substantial loss of protection

against *P. chabaudi* challenge. The mice treated with anti-IFN-γ before infection had significantly

higher parasitemia compared to isotype-treated controls previously infected with MCMV (**Fig 5B**).

The level of parasitemia in mice receiving neutralizing antibody is similar to a first infection

suggesting that all or most of the reduction due to MCMV is due to IFN-γ. The effect of

neutralizing MCMV-induced IFN-γ on challenge is also reflected in increased pathology, as

measured by weight loss (**Fig 5C**), and hypothermia (**Fig 5D**). To determine if there is an effect of

MCMV persistence-induced IFN-γ on the T cell response to challenge, the polyclonal CD4$^+$ T cell

phenotype was examined on day 15 post-challenge with *P. chabaudi.* Surprisingly, there was a

significant reduction in responding CD4$^+$ Teff after pre-treatment with anti-IFN-γ (**Fig 5E**).

Evaluation of serum cytokines in MCMV infected animals showed IL-12p40 production

throughout MCMV infection (**Fig 5F)** that is increased upon *P. chabaudi* challenge (**Fig 5G**).

Strikingly, IL-12 p40 production on days 7, 8 and 15 p.c. as well as TNF, CXCL-9 and CXCL-10

on day 7 p.c. were significantly reduced by neutralization of IFN-γ for six days before challenge

while IL-10 level was increased at 7 days p.c. (**Fig 5H)**. We also measured other cytokines

including IL-6, IL-15, IL-18, IFN-α and IFN-β, which are known to be produced during MCMV

infection to see if they were affected by neutralization of IFN-γ but did not see significant

differences at these time points.

To investigate potential direct effects of neutralizing IFN-γ on challenge parasitemia due

to persisting antibody and determine the role of persistent IFN-γ induced by MCMV in protection,

we examined the effect of anti- IFN-γ clone H22 on *P. chabaudi* infection with or without an

earlier dose of MCMV (**S4A Fig**). We found that administration of neutralizing antibody up to 2

days before infection with *P. chabaudi* does not have a significant measurable effect on peak

parasitemia (**S4B Fig**), while it does affect parasitemia significantly, as above, if MCMV is given

before infection (**S4C Fig**). To determine if neutralizing levels of antibody remained in the serum

during challenge, we developed an assay to test induction of CXCL10, which is induced in

macrophages by IFN-γ. CXCL10 mRNA was measured after culture of RAW 264.7 cells with

serum from mice before challenge (d40, **S4D Fig**), and on day 7 post *P. chabaudi* challenge with

or without prior anti-IFN-γ administration (**S4E Fig**). We found no detectable change for the

relative CXCL10 mRNA level with Raw 264.7 cells cultured with d40 or d47 serum, though there

was a trend suggesting antibody was there. Adding recombinant IFN-γ (rIFN-γ) to the assay

allowed detection of neutralizing antibody. In this assay, a reduced level of CXCL10 mRNA was

detectable in the presence of serum from 2 days after anti-IFN-γ (**S4F Fig**) and day 7 *P. chabaudi*

challenge (**S4G Fig**). Therefore, though neutralizing antibody does remain in the system during *P.*

chabaudi challenge, we conclude that the large effect of blocking IFN-γ after MCMV, and before

challenge, is mostly due to inhibition of MCMV-induced not *P. chabaudi*-induced IFN-γ.

Therefore, we investigated the mechanisms of this strong effect.

As IL-12 and IL-18 can induce IFN-γ from NK cells or previously stimulated T cells, we

queried their role in induction of IFN-γ by MCMV, but did not find any effect of blocking IL-12

and IL-18 together in the time period before challenge on protection from *P. chabaudi* parasitemia

or pathology by the MCMV vector (**S5 Fig**).

To explore the effect of IFN-γ on persisting epitope-specific T cells and the innate response in the

context of late MCMV infection, we adoptively transferred CTV-labelled B5 TCR Tg T cells into

Thy1.1 mice and then infected them with MCMV-B5, as illustrated in **Fig 6A**. IFN-γ was

neutralized on days 44-48 of MCMV infection. Blocking IFN-γ induced by MCMV-B5 infection

did not change the number of CTV$^-$ B5 T cells at day 50 p.i. (**Fig 6B**). IFN-γ also did not affect

the phenotype, as measured by the fraction of both Teff and Tmem (**Fig 6C**), or representation of

Tmem subsets, as defined by CD27 and CD62L expression (**Fig 6D**). B5 T cells induced by

MCMV-B5 are mostly memory T cells, with a fraction of effector T cells that is larger than that

seen on day 60 of *P. chabaudi* infection [8]. A panel of innate cell markers showed that after IFN-

γ neutralization on day 50 post-MCMV infection, the number of CD8α$^-$ DCs (CD3$^-$

CD11c^{hi}MHCII^{hi}) remained constant while the number of CD8α$^+$ DCs showed a trend towards a

reduction (**Fig 6E**). The level of IL-12p40 in the serum on day 50 post-MCMV infection was

significantly reduced in the group that received IFN-γ neutralizing antibody compared to isotype

(**Fig 6F**). Splenic CD8α$^+$ DCs are the major producers of IL-12p40 in response to systemic stimuli

[29].

IFN-γ generated by the MCMV booster drives prolonged protection of live malaria vaccine

Given the ability of MCMV to prolong protection from challenge and to promote IL-12 in an IFN-γ dependent manner, we tested mechanisms of the prolonged protection. Mice were vaccinated and then boosted with MCMV at day 140, as in **Fig 4**, with adoptive transfer of B5 T cells to determine the effect on vector-driven T cells. B5 T cell recipients were immunized with live malaria vaccine and boosted with MCMV-B5 at day 140 p.v., as illustrated in **Fig 7A.** Half of these animals received neutralizing antibody to IFN-γ, or isotype control antibody, days 194-198 p.v. (days 54-58 after the MCMV-B5 booster). T cell and antigen-presenting cell phenotypes were assessed at day 200 after malaria immunization, in the absence of challenge. Live *P. chabaudi* vaccination plus MCMV-B5 booster tended to increase survival of B5 T cells compared to live vaccine alone, as above, though this effect did not reach significance in this experiment (**Fig 7B**). Neutralization of IFN-γ did not have an effect on specific malaria vaccine-driven, MCMV-B5 boosted B5 T cell numbers, or the composition of Tmem subsets (**Fig 7C**). These data indicate that IFN-γ does not promote T cell survival or Tem differentiation directly, after MCMV-B5 boost, supporting the specific effect of MCMV-B5 epitope on the specific T cell responses. Staining for innate cell types, CD11c^{hi} dendritic cells (DC) were found to be increased by MCMV boost (**Fig 7D**). Furthermore, *in vivo* neutralization of IFN-γ significantly reduced the DC numbers compared to isotype control (**Fig 7D**). This innate effect was seen most strikingly in the CD8α$^+$ DC (CD3$^-$ CD11c^{hi}MHCII^{hi}CD8α$^+$) population (**Fig 7E**). We also assessed NK cells, and monocyte derived subsets including monocytes, macrophages and inflammatory monocytes to see if their numbers or activation marker expression were affected by neutralization of IFN-γ, but did not see other significant differences **(S6 Fig)**. Blocking IFN-γ after vaccination and the MCMV booster caused

a trend towards a reduced serum level of IL-12p40, though this did not reach significance, as it did

in MCMV infection alone on day 50 (**Fig 7F**). To test the role of MCMV booster-induced IFN-γ

in prolonging vaccine-induced protection, mice were vaccinated and then boosted with MCMV-

BAC at day 140, as above in **Fig 3**, and as shown schematically in **Fig 7G**. The control groups

either had no live vaccine and no MCMV or no vaccine and then MCMV only. Prior to

heterologous challenge with *P. chabaudi* AJ at day 200, vaccinated mice with MCMV boost

received IFN-γ neutralizing antibody, or isotype control, at days 194-198 p.v. (54-58 days post-

MCMV booster). MCMV boosting live vaccination showed significant protection against weight

loss (**Fig 7H)** and parasitemia (**Fig 7I)** compared to the adjuvanting effect of MCMV or

unvaccinated control mice. Blocking IFN-γ before challenge significantly reduced the booster

effect of the MCMV-BAC on protection at day 200 (**Fig 7J**). This was measured as a significant

increase in parasitemia in the IFN-γ neutralized animals (0.4%), compared to the isotype group

(0.09%), on day 5 post-challenge. While this is clearly a non-specific effect, as there is no B5

epitope here, polyclonal T cell phenotyping on day 15 post-challenge again showed that blocking

IFN-γ before reinfection significantly reduced the fraction of Teff generated during *P. chabaudi*

challenge, and was associated with higher parasitemia, suggesting a role for antigen presentation

during challenge that was not tested here (**Fig 7K**). Serum levels of *P. chabaudi* AJ-specific IgG

increased after challenge, but there was no significant difference from neutralization of IFN-γ (**Fig**

7L). Collectively, these data suggest a novel innate pathway where viral vector-induced IFN-γ

drives an increase in CD8α+ DCs, which promote increased protective IL-12 and Teff upon

challenge. We have identified two novel mechanisms of adjuvanting supporting the promise of

CMV-based vaccines, and defined a promising line of research for prolonging protection from

Plasmodium infection. Testing the full capacity of this strategy would require the identification of

more protective T cell epitopes.

Discussion

Alternative high-efficacy approaches for all malaria vaccine continue to be developed, but

little engineering has been focused on increasing longevity. These data show significant promise

for use of chronic vaccination to prolong the current short half-life of vaccines against systemic

infections requiring CD4 T cell memory. In summary, we demonstrated an adjuvant effect

dependent on the MCMV vector-induced IFN-γ that enhances CD8α$^+$ dendritic cell numbers, and

IL-12 production upon challenge. In addition, we demonstrate two basic mechanisms for the

extension of protection provided by the chronic MCMV booster to a typical short-lived malaria

vaccine. The MCMV vaccine-expressing a malaria epitope prolongs survival of epitope-specific

CD4 Teff and Tem. Polyclonal effector T cell numbers also increase when challenge occurs in the

presence of MCMV-induced IFN-γ, suggesting that the underlying mechanism of protection from

challenge is an increased Th1 response to *Plasmodium* due to increased IL-12 downstream of

higher dendritic cell numbers at the time of challenge.

We observed a strong adjuvant effect of the MCMV vector itself on *P. chabaudi* infection.

The effect of MCMV alone reduced *P. chabaudi* challenge parasitemia by about one half. The

adjuvant effect alone also drove a significant fraction of the prolonged protection seen from using

the MCMV vector in a boosting strategy, as demonstrated by neutralizing the IFN-γ induced by

MCMV-BAC before challenge. MCMV and gamma herpesvirus 68 also have an adjuvant effect

against influenza virus and *Listeria monocytogenes*, but not West Nile virus [22, 23], indicating

some specificity. This specificity may be due to latency. Gamma Herpesvirus 68 with a point

mutation in ORF73, a gene required for latency, showed reduced protection from *Listeria*, and also

reduced IFN-γ levels and chronic macrophage activation. HSV-1 and Sindbis viruses, which do

not establish latency, did not have this protective effect [22]. MCMV-induced cross-protection

against influenza also lasts through early and established (5-12 weeks), but not long-standing (9

months) MCMV latency. Production of IFN-γ during the latent phase of infection has also been

implicated in MCMV-induced protection to *Listeria* and *Yersinia* [22], influenza; which was

abrogated in IFN-γ deficient mice [23]. Our work extends these findings mechanistically by

showing that neutralizing IFN-γ late in MCMV infection reduced the number of CD8α$^+$ dendritic

cells in the spleen, and reduced the amount of some cytokines and chemokines IL-12, CXCL-9

and CXCL-10 and the number of CD4 Teff, but increased IL-10 cytokine level on challenge. While

we did not specifically test for latency, our vectors showed similar kinetics and outcomes to

previous studies suggesting infection with our vectors would be latent in our mouse model by day

40 [24]. Mechanisms of this protection had not been specifically established.

Decay of immunity, and failure to maintain *Plasmodium*-specific CD4 T cells make the

development of an effective and long-lived malaria vaccine challenging. Both natural immunity to

the parasite, and vaccines, generate immunity that decays due to a progressive loss of malaria-

specific T cells limiting long-lived protection [3, 4, 30, 31]. To our knowledge, there are no current

vaccine formulations that drive long-lived CD4 T cell responses [2, 32]. Therefore, the potential

of CMV to induce a prolonged T cell response with the desired phenotype was used to extend

vaccine protection. In this study, a two-step vaccine strategy that first induces a long-lived

protective antibody response, and then boosts with an MCMV strategy was chosen for the potential

to extend effector/memory CD4 T cells responses over a longer period to prolong protection. While

the protective T cell epitopes are not well defined, using a parasite derived CD4 T cell epitope

allowed us to test this strategy in a pre-clinical setting. Boosting with the MCMV vector did

prolong the protection generated by the live infection and-drug cure malaria vaccine as evidenced

by reduced challenge parasitemia after boost. The data show that animals that received live malaria

vaccination became susceptible to heterologous re-infection by day 200 p.v., while the MCMV-

B5 boost maintained the protection seen at day 60 until day 200. All malaria vaccines tested in

human studies thus far, including the most advanced vaccine RTS,S, show insufficient longevity

of protection [25].

CMV induces a robust and long-lasting T cell response including both effector and memory

components [20, 33]. Hansen et al. showed recently that RhCMV-expressing *Plasmodium*

knowlesi antigens, generated a parasite antigen-specific immune response that delayed parasite

growth upon sporozoite challenge via generation of CD8 T cell responses [16]. However, whether

an MCMV vector could be used to boost pre-existing specificities has not been tested, and only a

few studies have investigated CD4 T cell responses to MCMV [34]. Here we show that the MCMV

vector can be used to specifically boost CD4 epitope-specific effector T cell responses, which is

one of the primary gaps in our current malaria vaccine arsenal. Both CD4 Teff and Tmem

phenotypes are generated in *P. chabaudi* infection, and both promote protection [8, 10]. Here, we

observed that MCMV-B5 increased epitope-specific T cells, and specifically generated GC-Tfh

and promoted maintenance of the Tem phenotype and effector functions (IFN-γ, TNF and IL-2)

past day 200. CD4 T cells are crucial for protective immunity to *P. chabaudi* as they produce

cytokines that both promote phagocytosis and B cell help [35]. Our previous findings revealed the

power of Teff in protection from persistent *P. chabaudi* with help from B cells. Adoptive transfer

of B5 effector CD4 T cells reduced the peak of *P. chabaudi* parasitemia by half [10]. While

memory T cells are less effective, CD27$^-$ TemL do reduce parasitemia, and are the most protective

memory phenotype. In this study, we showed that the MCMV-B5 vector specifically activated and

maintained both adoptively transferred epitope-specific T cells and MCMV-specific CD4$^+$ T cells

over 2 months. Interestingly, Teff were clearly present among the few m78-Tetramer MCMV-

specific CD4 T cells over 2 months, but did not reach significant levels compared to uninfected

controls. This may reflect the small number of cells detected, and so we used the B5 TCR Tg to

test for MCMV-induced proliferation at late time points, however this assay had high background.

Consistent with this, MCMV-B5 maintained both Teff and the potentially protective late effector

memory phenotype T cells generated by *P. chabaudi* infection. While we do not present evidence

here that the protection provided is due to increased survival of B5 T cells, their numbers are

boosted at a timepoint when live vaccine induced T cells and protection would otherwise decay,

though antibody levels remain constant. It should be noted that the B5 specificity is not especially

potent in protection, as the effect of this one specificity is only seen in immunocompromised

animals. CD8 T cells specific for some MCMV loci are given to inflationary activation, even after

15 weeks of MCMV infection [36]. The locus where B5 is expressed, IE1, can be a memory

inflation inducing locus for CD8 T cells [37]. CD4 T cell responses to MCMV are even less well

understood. Therefore, additional T cell epitope discovery in this field will aid in the long-term

goal of developing durable vaccine induced protection.

The MCMV vector increased CD8α$^+$ DCs and IL-12 both before challenge, and in response

to the parasite. Induction of IL-12 upon *P. chabaudi* challenge was reduced in the context of prior

neutralization of IFN-γ. Considering that CD8α$^+$ DCs are a key source of IL-12 production in many

infections [38], and also present antigen to T cells upon infection, the reduced IL-12 level in

response to *P. chabaudi* challenge after IFN-γ neutralization could be directly linked to the change

in Teff cell numbers seen. While IL-12 is more commonly associated with stimulating IFN-γ

production, IL-12 production by DC and macrophages can also be promoted by IFN-γ present in

the milieu, representing a positive feedback mechanism [39, 40]. Therefore, prolonged IFN-γ

appears to be maintaining CD8α^+ DCs, an IL-12 producing cell type, suggesting a novel

mechanism for this feedback. As IL-12 deficient mice have higher *P. chabaudi* parasitemia than

WT [41], or Ifng deficient animals [42], the reduced IL-12 response during challenge is likely to

explain the reduced control of parasites seen upon neutralization of IFN-γ. While more research

will be required to establish if the increase in dendritic cells late in MCMV infection is sufficient

to drive the increased IL-12 response or the increase in Teff upon challenge, the increase of CD8α^+

DCs supports this interpretation.

464   Prior work has shown that giving *P. chabaudi*-infected mice recombinant IFN-γ increased

the number of polyclonal CD4 Tem, and also prolonged heterologous immunity. In our system,

generation and maintenance of Tem is epitope specific, and not affected by blocking IFN-γ

generated during MCMV infection. Our previous work showed that both memory T cell

differentiation and prolonged infection contribute to maintenance of Tem numbers in *P. chabaudi*

infection [10]. It remains to be proven whether the MCMV-BAC driven increase in DCs, and/or

priming of them to respond/present antigen, leads directly to enhanced responsiveness of parasite

specific T cells/expansion and clearance of parasite by Th1-promoted macrophage activation and

Tfh-promoted antibody production. It is also possible that antigen presentation and Th1 response

are reduced with the presence of detectable anti-IFN-γ auto-antibody [43]. Exploring this

hypothesis to determine potential effects of anti-IFN-γ antibody, we found that administration of

neutralizing antibody up to 2 days before infection with *P. chabaudi* does not have a significant

effect on peak parasitemia, though there may be a trend towards increasing parasitemia. Indeed, in

Meding et al, where IFN-γ was neutralized during the infection, the change in parasitemia was

very small [42], similar to the effects of anti-IFN-γ seen on primary infection in our study. There

is a clear reduction of these IFN-γ-inducible chemokines (CXCL9 and CXCL10) in the serum

upon infection after anti-IFN-γ. This suggests that anti-IFN-γ directly or indirectly reduces their

production in response to the parasite. An indirect example would be that these chemokines can

be made by DCs, the reduction of IFN-γ induced chemokines could reflect the changes on DC

number that we have documented downstream of MCMV. Our results show that although

neutralizing antibody persists into the challenge phase, the largest effect of anti-IFN-γ after

boosting, requires MCMV-induced IFN-γ, and is not a direct effect on parasitemia.

Neutralization of IFN-γ led to a loss of the extended protection provided by MCMV

boosting and protection from *P. chabaudi* infection. Teff were reduced on challenge after IFN-γ

neutralization, which is particularly striking given the increase in parasite in anti-IFN-γ pre-treated

animals. IFN-γ production has been previously reported in MCMV and HCMV infection during

the period coinciding with latency [23]. Data from our laboratory and others establishes that IFN-

γ expression during persistent *P. chabaudi* infection, or recombinant IFN-γ also promote

protection from heterologous re-infection at day 200 [12, 44]. It was also reported that rIFN-γ

increased the number of polyclonal Tem at day 200 after *P. chabaudi* infection [44].

We have used basic knowledge of T cell differentiation in *P. chabaudi* infection to reverse

engineer a novel chronic vaccine strategy. We demonstrate a method and a mechanism to prolong

protection by a live malaria vaccine, which alone is short-lived, due to a loss of specific T cells.

Our work is novel as it indicates that MCMV works as a booster for CD4 T cell responses both by

maintaining specific effector and effector memory T cells and also as an adjuvant driving

protective IFN-γ production. The increase in dendritic cell numbers and increased IL-12 induced

by MCMV, together with and a trend towards increased Th1 cytokines upon challenge, suggest

that ultimately, a stronger T cell response to parasite explains the prolonged protection. This

strategy has the potential to resolve the major issue of short-lived protection generated by malaria

vaccines, as well as help understand how natural immunity could be prolonged. There are safety

issues to work out before CMV vectors could be used in humans. However, superinfection, the

ability to re-infect CMV carriers, and a varied T cell response are two benefits of this approach

[45, 46]. Further understanding of the mechanisms of this exciting new model could lead to more

rapidly translatable tools in the quest to prolong immunity to persistent or recurrent pathogens

without introducing another pathogen. This could be applicable to malaria, a disease that kills up

to half a million people a year; as well as other persistent or recurrent pathogens.

Materials and methods

Ethics statement

All experiments were carried out according to protocol approved by the University of

Texas Medical Branch Institutional Animal Care and Use Committee (IACUC).

Mice, Parasites and Immunization

All mice used in our study were maintained in the UTMB animal facility under specific

pathogen-free conditions with access to food and water *ad libitum*. Recipient Thy1.1 BALB/cByJ

were backcrossed to BALB/cJ (N4; The Jackson Laboratory, Bar Harbor, ME). B5 T cell receptor

transgenic (TCR Tg), which recognizes a peptide from merozoite surface protein-1 (MSP-1; 1157-

1171, ISVLKSRLLKRKKYI/I-E^d) was backcrossed to BALB/cJ (N10). B5 TCR Tg mice, and *P.*

chabaudi chabaudi AS stocks, were a kind gift from J. Langhorne (Francis Crick Institute, London,

UK). *P. chabaudi* AJ (MRA-740, a cloned line that is highly related but heterologous to AS [47,

48] was purchased from BEI Resources, formerly MR4. B5 TCR Tg mice were typed using

primers Vα2, 5′-gaacgttccagattccatgg-3′ and 5′-atggacaagatcctgacagcatcg-3′, and vβ8.1, 5'

cagagaccctcaggcggctgctcagg-3′ and 5′-atgggctccaggctgttctttgtggttttgattc-3′ [8]. Both male and

female recipients were used, however no sex differences in infection were noted. All adoptive

transfer experiments were done using age- and sex- matched recipient and donors.

For infection and drug cure vaccination, the protocol was performed as previously

documented in mice and humans [3, 27]. Six- to nine-week old mice were infected (10^6 *P.*

chabaudi) and treated with two subsequent noncurative doses of chloroquine (10 mg/kg body

weight/day; Sigma-Aldrich), when parasitemia reached 1-2% of erythrocytes. Parasites were

counted by light microscopy in thin blood smears stained with Giemsa (Sigma-Aldrich, St. Louis,

MO). Parasitemia was monitored by microscopic examination of thin blood smears from the tail.

Thin blood smears were stained with Giemsa (Ricca chemical company, Arlington, TX), and

parasites were counted by light microscopy.

Construction of MCMV and viral infection

A vaccine vector, MCMV-K181-B5, was generated as follows. MCMV-K181-Perth was

genetically modified to carry the MHC II-E^d-restricted B5 epitope (ISVLKSRLLKRKKYI) from

the *Plasmodium chabaudi* Merozoite Surface Protein MSP1 (1157–1171) in frame with the C-

terminus end of the MCMV IE1 gene (**Fig S1 A**). The B5 epitope was inserted into the MCMV-

K181 genome using bacterial artificial chromosome recombineering, as described previously [49].

The resulting virus was denoted MCMV-B5. Mice were infected by injecting $5x10^3$ PFU of

salivary gland-propagated MCMV-B5 virus or the parental virus, MCMV-K181-Perth (MCMV-

BAC). Viruses were injected i.p. in 0.2ml of PBS.

Virus quantification

MCMV-B5 or MCMV-BAC viral copy number of CMV/β-actin ratios were determined in

different organs in infected BALB/cJ mice. DNA was purified from tissue (liver, lung, and salivary

gland) using the Gentra Puregene Kit (Qiagen, Valencia, CA) following manufacturer's

instructions. All tissues were stored at -80°C in PBS for DNA, or Trizol (Thermo Fisher Scientific,

Carlsbad, CA) for RNA purification, until sample processing. qPCR was performed with 50 ng of

purified DNA. Specific primers for MCMV immediate early (IE1) gene (NCBI Accession No.

M11788) 5-tca gcc atc aac tct gct acc aac-3 and 5-gtg cta gat tgt atc tgg tgc tcc tc-3 and for murine

β-actin (Accession No. M12481) 5-gct gta ttc ccc tcc atc gtg -3 and 5-cac ggt tgg cct tag ggt tca-

3. Serial dilutions of a control plasmid were used to generate a standard curve for both MCMV-

IE1 and β-actin, as described previously, and qPCR of viral DNA has been shown to correlate

quantitatively with the plaque assay throughout the viral infection [50]. All sample measurements

were performed in triplicate. Real time PCR (VIIA, AB Biosystems, Life Technologies, Carlsbad,

CA).

Cell culture with serum, RNA isolation and real-time PCR analysis

RAW264.7 macrophage-like cell line was cultured in RPMI medium (Gibco)

supplemented with 10% (v/v) fetal bovine serum and antibiotics (100 U/ml penicillin and 100

lg/ml streptomycin), Sodium Pyruvate (100x, Gibco), 45% D-Glucose (100x, Sigma), L-glutamine

(100x, Corning) at atmosphere of 5% CO_2. RAW264.7 cells were incubated with sera form IFN-

γ neutralizing mice with or without recombinant IFN-γ for 24h. Total RNA was extracted with

Direct-Zol RNA reagents (Zymo research, R2025) as described by the manufacturer. cDNA was

synthesized from total RNA using the iScript cDNA Synthesis kit (Bio-Rad). Real-time PCR was

performed with cDNA using the SYBR Green Supermix (Bio-Rad). The β-actin gene was used for

normalization. Relative changes in gene expression were calculated using the comparative

threshold cycle (Ct) method. The following primers were used: CXCL-10 5'-

cctgctgggtctgagtggga-3' and 5'-gataggctcgcagggatgat-3'; β-actin 5-gctgtattcccctccatcgtg-3 and 5-

cacggttggccttagggttca-3.

Adoptive transfers and *in vivo* studies

MACS-purified CD4$^+$ T cells from young, uninfected B5 TCR-Tg mouse spleens (Miltenyi

San Diego, CA) were labelled with cell trace violet (CTV) and adoptively transferred (2×10^6) into

Thy1.1 congenic mice. For cytokine neutralization, three doses of anti-IFN-γ antibody (H22, 0.5

mg) or isotype control antibody (PIP, 0.5 mg); or a combination of anti-IL-12p40 (C17.8, rat

IgG2a, 300 µg) and anti-IL-18 antibody (YIGIF74–1G7, control Rat IgG2a, 200 µg) were given,

every other day. All *in vivo* antibodies were from BioXCell (Lebanon, NH).

Flow cytometry

Single-cell suspensions from spleens were prepared at the indicated time points after infection in

HBSS) with HEPES (Thermo Fisher Scientific, Waltham, MA) using gentleMACS (Miltenyi, San

Diego, CA) and incubated in RBC lysis buffer (eBioscience, San Diego, CA). Single cells were

stained extracellular in PBS, 2% FBS, and 0.1% sodium azide with antibodies. For T cells: anti-

CD127 (A7R34), anti-PD-1 (RMP1-30), anti-CD62L (MEL-14), CD4 (RM4-5), (anti-CD90.2

[30-H12], anti-CD44 [IM7], and anti-CD27 [LG-7F9]) (all from eBioscience/Invitrogen and

Biolegend, San Diego, CA). Phenotype of polyclonal CD4[+] T cell subsets and *Plasmodium* B5-

specific CD4[+] T cell subsets were identified with gating strategy as schematically depicted in **S7**

Fig. For follicular T cell staining with CXCR5, cells were stained in PBS + 0.5% BSA + 0.1%

sodium azide + 2% Normal Mouse Serum (NMS) and 2% FBS (Sigma, St. Louis, MO). Rat anti-

mouse purified CXCR5 (2G8, BDbioscience, San Jose, CA, 1 hr., 4°C) was followed by biotin-

conjugated AffiniPure Goat anti-rat (H+L, Jackson ImmunoResearch, West Grove, PA, 30 min,

4°C) followed by Streptavidin-eFluor 450. For antigen presenting cells were stained with anti-

Ly6C, anti-Ly6G (1A8), anti CD11b (M1/70), anti-CD11c (N418), DX5, MHCII/I-A/Eb

(M5/114.15.2). For intracellular cytokine staining, cells were cultured for 5 hours at 37°C, 5%

CO_2 in the presence of GolgiPlug (BD Bioscience) before extracellular staining and then fixed

with 2% paraformaldehyde overnight. Cells were then permeabilized using BD 10X Permeability

buffer diluted with water, and stained with anti-IFN-γ (XMG1.2), anti-TNF (MP6-XT22), anti-IL-

2 (JES6-5H4). Virus-specific CD4[+] T cells were identified using m78/I-A[d] tetramers, with peptide

(SQQKMTSLPMSVFYS) and CLIP/I-A[d] as tetramer control (PVSKMRMATPLLMQA) and

CD8[+] T cells with H2D[d] restricted m45-tetramer peptide 507-VGPALGRGL-515 [20, 21] (both

from NIH Tetramer Core Facility, Emory, Atlanta, GA). Tetramer staining was performed within

6 months of receiving the reagents at 4C for 1 hour before/after surface and intracellular staining.

Cells were acquired on a LSR II Fortessa using FACSDiva software (BD Biosciences, San Jose,

CA). Compensation was performed in FlowJo using single-stained splenocytes (with CD4 in all

colors). All data analysis was performed using the FlowJo software package (version 10, FlowJo,

BD San Jose, CA). Cell numbers were calculated as the % of recovered Tg cells/lymphocytes x

the number of lymphocytes, except for **Fig 6** and **Fig 7** which the recovered cells were counted by

inclusion of counting beads (Accu-Check, Molecular Probes, Thermo Fisher Scientific) in the

FACS sample according to manufacturer's instructions and calculated using the equation: Total

cell number = ((all cell event count / bead event count) x bead conc. (per µl) x (bead volume / cell

volume)) x volume of original sample. This formula only works if the whole volume of FACS

tube is collected.

Cytokines and antibody measurements

Cytokines and chemokines (CXCL10 (IP-10), IFN-γ, IFN-β, IFN-α, IL-10, IL-12p40, IL-

15, IL-18, IL-6, and TNF-α) were measured in the plasma by Legendplex according to

manufacturer's protocol (Biolegend, San Diego, CA). Plasma samples were collected at the

indicated days and were stored in -80°C until analysis. All data were acquired on an LSR II

Fortessa flow cytometer using FACSDiva software, version 8 (BD Biosciences, San Jose, CA).

The concentration of each analyte is determined with cytometric data based on a standard curve

using the LEGENDplex™ data analysis software (Biolegend, San Diego, CA). *P. chabaudi* AJ-

specific IgG was evaluated in plasma at indicated days by ELISA, as described previously [5].

Plates were coated with *P. chabaudi* AJ parasite lysate overnight at 4°C in PBS with 0.05%

Sodium Azide. Plates were blocked with 1% BSA, 0.3% Tween 20, 0.05% Sodium Azide and then

sera were incubated. Bound antibody was detected using Alkaline Phosphatase-conjugated goat

anti-mouse IgG (Southern Biotech, Birmingham, AL). The reaction was revealed with a p-

Nitrophenyl phosphate disodium salt (Sigma, USA) solution (1mg/ml) in diethanolamine buffer.

Plates were analyzed with an Omega plate reader (FLUOstar Omega BMG LABTECH Inc,

Durham, NC, USA).

Statistical analysis

All data were analyzed by ANOVA, followed by Tukey's or two-tailed Mann-Whitney test or two-tailed unpaired Student t test, where indicated. All data are presented as mean ± SEM. Statistical tests were performed using the statistical software package in Prism (version 9.2, GraphPad Software, USA): *$p < 0.05$, **$p \leq 0.01$, ***$p \leq 0.001$ and n.s., not significant.

Supporting information

Acknowledgments

We would like to thank Florentin Aussenac, Victor H. Carpio, Nadia D. Domingo, Lucinda Puebla-Clark and Peter Fleming for feedback and discussion. Thanks to Maria I. Giraldo Giraldo and Gregg Milligan for technical advice. We thank the NIH Tetramer Core Facility at Emory, Atlanta, GA for generating both the m78-Tetramer and m-45-Tetramer used in this study. This work was supported by the James W. McLaughlin Fellowship Fund (KG) and NIH R01AI135061, R01AI089953 (RS, KG).

Author Contributions

Conceptualization: KG, MGB, MAD, RS

Investigation: KG, SAI

Formal Analysis: KG, SAI, RS

Methodology and Resources: KG, SAI, MLO, MAD, MGB

Funding acquisition: KG, RS, MGB, MAD

Project Administration, Supervision: RS, MAD

Writing – Original Draft Preparation: KG

661 **Writing – Review & Editing:** RS, MGB, MAD

References

References

1.   Kennedy RB, Ovsyannikova IG, Palese P, Poland GA. Current Challenges in Vaccinology.

Front Immunol. 2020;11:1181. Epub 2020/07/17. doi: 10.3389/fimmu.2020.01181. PubMed

PMID: 32670279; PubMed Central PMCID: PMCPMC7329983.

2.   Pollard AJ, Bijker EM. A guide to vaccinology: from basic principles to new

developments. Nat Rev Immunol. 2021;21(2):83-100. Epub 2020/12/24. doi: 10.1038/s41577-

020-00479-7. PubMed PMID: 33353987; PubMed Central PMCID: PMCPMC7754704.

3.   Freitas do Rosario AP, Muxel SM, Rodriguez-Malaga SM, Sardinha LR, Zago CA,

Castillo-Mendez SI, et al. Gradual decline in malaria-specific memory T cell responses leads to

failure to maintain long-term protective immunity to *Plasmodium chabaudi* AS despite persistence

of B cell memory and circulating antibody. J Immunol. 2008;181(12):8344-55. PubMed PMID:

19050251.

4.   Palacpac NMQ, Horii T. Malaria vaccines: facing unknowns. F1000Res. 2020;9. Epub

2020/05/14. doi: 10.12688/f1000research.22143.1. PubMed PMID: 32399189; PubMed Central

PMCID: PMCPMC7194457.

5.   Achtman AH, Stephens R, Cadman ET, Harrison V, Langhorne J. Malaria-specific

antibody responses and parasite persistence after infection of mice with *Plasmodium chabaudi*

chabaudi. Parasite Immunol. 2007;29(9):435-44. doi: 10.1111/j.1365-3024.2007.00960.x.

PubMed PMID: 17727567.

6.   Scott P, Artis D, Uzonna J, Zaph C. The development of effector and memory T cells in

cutaneous leishmaniasis: the implications for vaccine development. Immunol Rev. 2004;201:318-

38. Epub 2004/09/14. doi: 10.1111/j.0105-2896.2004.00198.x. PubMed PMID: 15361250.

7.   Peters NC, Pagan AJ, Lawyer PG, Hand TW, Henrique Roma E, Stamper LW, et al.

Chronic parasitic infection maintains high frequencies of short-lived Ly6C+CD4+ effector T cells

that are required for protection against re-infection. PLoS Pathog. 2014;10(12):e1004538. Epub

2014/12/05. doi: 10.1371/journal.ppat.1004538. PubMed PMID: 25473946; PubMed Central

PMCID: PMCPMC4256462.

8.   Stephens R, Langhorne J. Effector memory Th1 CD4 T cells are maintained in a mouse

model of chronic malaria. PLoS Pathog. 2010;6(11):e1001208. Epub 2010/12/03. doi:

10.1371/journal.ppat.1001208.   PubMed   PMID: 21124875;   PubMed   Central   PMCID:

PMCPMC2991260.

9.   Zaph C, Uzonna J, Beverley SM, Scott P. Central memory T cells mediate long-term

immunity to Leishmania major in the absence of persistent parasites. Nat Med. 2004;10(10):1104-

10. Epub 2004/09/28. doi: 10.1038/nm1108. PubMed PMID: 15448686.

10.   Opata MM, Ibitokou SA, Carpio VH, Marshall KM, Dillon BE, Carl JC, et al. Protection

by and maintenance of CD4 effector memory and effector T cell subsets in persistent malaria

infection. PLoS Pathog. 2018;14(4):e1006960. doi: 10.1371/journal.ppat.1006960. PubMed

PMID: 29630679; PubMed Central PMCID: PMCPMC5908200.

11.   Stephens R, Albano FR, Quin S, Pascal BJ, Harrison V, Stockinger B, et al. Malaria-

specific transgenic CD4(+) T cells protect immunodeficient mice from lethal infection and

demonstrate requirement for a protective threshold of antibody production for parasite clearance.

Blood. 2005;106(5):1676-84. doi: 10.1182/blood-2004-10-4047. PubMed PMID: 15890689.

12.   Ibitokou SA, Gbedande K, Opata MM, Carpio VH, Marshall KM, Stephens R. Effects of

Low-Level Persistent Infection on Maintenance of Immunity by CD4 T Cell Subsets and Th1

Cytokines. Infect Immun. 2023;91(3):e0053122. Epub 2023/03/16. doi: 10.1128/iai.00531-22.

PubMed PMID: 36920200; PubMed Central PMCID: PMCPMC10016079.

13.   Hansen SG, Vieville C, Whizin N, Coyne-Johnson L, Siess DC, Drummond DD, et al.

Effector memory T cell responses are associated with protection of rhesus monkeys from mucosal

simian immunodeficiency virus challenge. Nat Med. 2009;15(3):293-9. Epub 2009/02/17. doi:

10.1038/nm.1935. PubMed PMID: 19219024; PubMed Central PMCID: PMC2720091.

14.   Beverley PC, Ruzsics Z, Hey A, Hutchings C, Boos S, Bolinger B, et al. A novel murine

cytomegalovirus vaccine vector protects against Mycobacterium tuberculosis. J Immunol.

2014;193(5):2306-16. doi: 10.4049/jimmunol.1302523. PubMed PMID: 25070842; PubMed

Central PMCID: PMCPMC4134927.

15.   Hansen SG, Ford JC, Lewis MS, Ventura AB, Hughes CM, Coyne-Johnson L, et al.

Profound early control of highly pathogenic SIV by an effector memory T-cell vaccine. Nature.

2011;473(7348):523-7. doi: 10.1038/nature10003. PubMed PMID: 21562493; PubMed Central

PMCID: PMCPMC3102768.

16.   Hansen SG, Womack J, Scholz I, Renner A, Edgel KA, Xu G, et al. Cytomegalovirus

vectors expressing *Plasmodium knowlesi* antigens induce immune responses that delay parasitemia

upon sporozoite challenge. PLoS One. 2019;14(1):e0210252. doi: 10.1371/journal.pone.0210252.

PubMed PMID: 30673723; PubMed Central PMCID: PMCPMC6343944 VIR Biotechnology

Inc., a company that may have a commercial interest in the results of this research and technology.

The potential individual and institutional conflicts of interest have been reviewed and managed by

OHSU. This does not alter our adherence to PLOS ONE policies on sharing data and materials.

17.   Tsuda Y, Parkins CJ, Caposio P, Feldmann F, Botto S, Ball S, et al. A cytomegalovirus-

based vaccine provides long-lasting protection against lethal Ebola virus challenge after a single

dose. Vaccine. 2015;33(19):2261-6. Epub 2015/03/31. doi: 10.1016/j.vaccine.2015.03.029.

PubMed PMID: 25820063; PubMed Central PMCID: PMCPMC4402448.

18.   Wilski NA, Snyder CM. From Vaccine Vector to Oncomodulation: Understanding the

Complex Interplay between CMV and Cancer. Vaccines (Basel). 2019;7(3). Epub 2019/07/22. doi:

10.3390/vaccines7030062.   PubMed   PMID: 31323930;   PubMed   Central   PMCID:

PMCPMC6789822.

19.   Snyder CM, Cho KS, Bonnett EL, Allan JE, Hill AB. Sustained CD8+ T cell memory

inflation after infection with a single-cycle cytomegalovirus. PLoS Pathog. 2011;7(10):e1002295.

doi: 10.1371/journal.ppat.1002295. PubMed PMID: 21998590; PubMed Central PMCID:

PMCPMC3188546.

20.   Snyder CM, Cho KS, Bonnett EL, van Dommelen S, Shellam GR, Hill AB. Memory

inflation during chronic viral infection is maintained by continuous production of short-lived,

functional T cells. Immunity. 2008;29(4):650-9. doi: 10.1016/j.immuni.2008.07.017. PubMed

PMID: 18957267; PubMed Central PMCID: PMCPMC2583440.

21.   Verma S, Weiskopf D, Gupta A, McDonald B, Peters B, Sette A, et al. Cytomegalovirus-

Specific CD4 T Cells Are Cytolytic and Mediate Vaccine Protection. J Virol. 2016;90(2):650-8.

Epub 2015/10/23. doi: 10.1128/JVI.02123-15. PubMed PMID: 26491148; PubMed Central

PMCID: PMCPMC4702662.

22.   Barton ES, White DW, Cathelyn JS, Brett-McClellan KA, Engle M, Diamond MS, et al.

Herpesvirus latency confers symbiotic protection from bacterial infection. Nature.

2007;447(7142):326-9. Epub 2007/05/18. doi: 10.1038/nature05762. PubMed PMID: 17507983.

23.   Furman D, Jojic V, Sharma S, Shen-Orr SS, Angel CJ, Onengut-Gumuscu S, et al.

Cytomegalovirus infection enhances the immune response to influenza. Sci Transl Med.

2015;7(281):281ra43. Epub 2015/04/04. doi: 10.1126/scitranslmed.aaa2293. PubMed PMID:

25834109; PubMed Central PMCID: PMCPMC4505610.

24.   Andrews DM, Estcourt MJ, Andoniou CE, Wikstrom ME, Khong A, Voigt V, et al. Innate

immunity defines the capacity of antiviral T cells to limit persistent infection. J Exp Med.

2010;207(6):1333-43. Epub 2010/06/02. doi: 10.1084/jem.20091193. PubMed PMID: 20513749;

PubMed Central PMCID: PMCPMC2882831.

25.   Campo JJ, Sacarlal J, Aponte JJ, Aide P, Nhabomba AJ, Dobano C, et al. Duration of

vaccine efficacy against malaria: 5th year of follow-up in children vaccinated with RTS,S/AS02

in   Mozambique.   Vaccine.   2014;32(19):2209-16.   Epub   2014/03/19.   doi:

10.1016/j.vaccine.2014.02.042. PubMed PMID: 24631081.

26.   Raja AI, Cai Y, Reiman JM, Groves P, Chakravarty S, McPhun V, et al. Chemically

Attenuated Blood-Stage *Plasmodium yoelii* Parasites Induce Long-Lived and Strain-Transcending

Protection. Infect Immun. 2016;84(8):2274-88. Epub 2016/06/02. doi: 10.1128/IAI.00157-16.

PubMed PMID: 27245410; PubMed Central PMCID: PMC4962623.

27.   Pombo DJ, Lawrence G, Hirunpetcharat C, Rzepczyk C, Bryden M, Cloonan N, et al.

Immunity to malaria after administration of ultra-low doses of red cells infected with *Plasmodium*

falciparum.   Lancet.   2002;360(9333):610-7.   Epub   2002/09/21.   doi: 10.1016/S0140-

6736(02)09784-2. PubMed PMID: 12241933.

28.   Holtappels R, Pahl-Seibert MF, Thomas D, Reddehase MJ. Enrichment of immediate-early

1 (m123/pp89) peptide-specific CD8 T cells in a pulmonary CD62L(lo) memory-effector cell pool

during latent murine cytomegalovirus infection of the lungs. J Virol. 2000;74(24):11495-503.

Epub 2000/11/23. doi: 10.1128/jvi.74.24.11495-11503.2000. PubMed PMID: 11090146; PubMed

Central PMCID: PMCPMC112429.

29.   Farrand KJ, Dickgreber N, Stoitzner P, Ronchese F, Petersen TR, Hermans IF. Langerin+

CD8alpha+ dendritic cells are critical for cross-priming and IL-12 production in response to

systemic   antigens.   J   Immunol.   2009;183(12):7732-42.   Epub   2009/11/20.   doi:

10.4049/jimmunol.0902707. PubMed PMID: 19923446.

30.   RTS. Efficacy and safety of RTS,S/AS01 malaria vaccine with or without a booster dose

in infants and children in Africa: final results of a phase 3, individually randomised, controlled

trial. Lancet. 2015;386(9988):31-45. Epub 2015/04/29. doi: 10.1016/S0140-6736(15)60721-8.

PubMed PMID: 25913272; PubMed Central PMCID: PMC5626001.

31.   Rts SCTP. Efficacy and safety of the RTS,S/AS01 malaria vaccine during 18 months after

vaccination: a phase 3 randomized, controlled trial in children and young infants at 11 African

sites. PLoS Med. 2014;11(7):e1001685. doi: 10.1371/journal.pmed.1001685. PubMed PMID:

25072396; PubMed Central PMCID: PMCPMC4114488.

32.   Draper SJ, Sack BK, King CR, Nielsen CM, Rayner JC, Higgins MK, et al. Malaria

Vaccines: Recent Advances and New Horizons. Cell Host Microbe. 2018;24(1):43-56. Epub

2018/07/13. doi: 10.1016/j.chom.2018.06.008. PubMed PMID: 30001524; PubMed Central

PMCID: PMCPMC6054918.

33.   Cicin-Sain L, Sylwester AW, Hagen SI, Siess DC, Currier N, Legasse AW, et al.

Cytomegalovirus-specific T cell immunity is maintained in immunosenescent rhesus macaques. J

Immunol. 2011;187(4):1722-32. doi: 10.4049/jimmunol.1100560. PubMed PMID: 21765018;

PubMed Central PMCID: PMCPMC3151292.

34.   Snyder CM, Loewendorf A, Bonnett EL, Croft M, Benedict CA, Hill AB. CD4+ T cell

help has an epitope-dependent impact on CD8+ T cell memory inflation during murine

cytomegalovirus   infection.   J   Immunol.   2009;183(6):3932-41.   Epub   2009/08/21.   doi:

10.4049/jimmunol.0900227. PubMed PMID: 19692644; PubMed Central PMCID: PMCPMC2766182.

PMCPMC2766182.

35.   Meding SJ, Langhorne J. CD4+ T cells and B cells are necessary for the transfer of

protective immunity to *Plasmodium chabaudi chabaudi*. Eur J Immunol. 1991;21(6):1433-8. doi:

10.1002/eji.1830210616. PubMed PMID: 1675172.

36.   Lemmermann NAW, Reddehase MJ. Direct Evidence for Viral Antigen Presentation

during Latent Cytomegalovirus Infection. Pathogens. 2021;10(6). Epub 2021/07/03. doi:

10.3390/pathogens10060731. PubMed PMID: 34200578; PubMed Central PMCID:

PMCPMC8229173.

37.   Simon CO, Holtappels R, Tervo HM, Bohm V, Daubner T, Oehrlein-Karpi SA, et al. CD8

T cells control cytomegalovirus latency by epitope-specific sensing of transcriptional reactivation.

J Virol. 2006;80(21):10436-56. Epub 2006/08/25. doi: 10.1128/JVI.01248-06. PubMed PMID:

16928768; PubMed Central PMCID: PMCPMC1641801.

38.   Mashayekhi M, Sandau MM, Dunay IR, Frickel EM, Khan A, Goldszmid RS, et al.

CD8alpha(+) dendritic cells are the critical source of interleukin-12 that controls acute infection

by *Toxoplasma gondii* tachyzoites. Immunity. 2011;35(2):249-59. Epub 2011/08/27. doi:

10.1016/j.immuni.2011.08.008. PubMed PMID: 21867928; PubMed Central PMCID:

PMCPMC3171793.

39.   Garris CS, Arlauckas SP, Kohler RH, Trefny MP, Garren S, Piot C, et al. Successful Anti-

PD-1 Cancer Immunotherapy Requires T Cell-Dendritic Cell Crosstalk Involving the Cytokines

IFN-gamma and IL-12. Immunity. 2018;49(6):1148-61 e7. Epub 2018/12/16. doi:

10.1016/j.immuni.2018.09.024. PubMed PMID: 30552023; PubMed Central PMCID:

PMCPMC6301092.

40.   Grohmann U, Belladonna ML, Vacca C, Bianchi R, Fallarino F, Orabona C, et al. Positive

regulatory role of IL-12 in macrophages and modulation by IFN-gamma. J Immunol.

2001;167(1):221-7. Epub 2001/06/22. doi: 10.4049/jimmunol.167.1.221. PubMed PMID:

11418652.

41.   Su Z, Stevenson MM. IL-12 is required for antibody-mediated protective immunity against

blood-stage *Plasmodium chabaudi* AS malaria infection in mice. J Immunol. 2002;168(3):1348-

55. Epub 2002/01/22. doi: 10.4049/jimmunol.168.3.1348. PubMed PMID: 11801675.

42.   Meding SJ, Cheng SC, Simon-Haarhaus B, Langhorne J. Role of gamma interferon during

infection with *Plasmodium chabaudi chabaudi*. Infect Immun. 1990;58(11):3671-8. Epub

1990/11/01. doi: 10.1128/iai.58.11.3671-3678.1990. PubMed PMID: 1977706; PubMed Central

PMCID: PMCPMC313713.

43.   Chen ZM, Yang XY, Li ZT, Guan WJ, Qiu Y, Li SQ, et al. Anti-Interferon-gamma

Autoantibodies Impair T-Lymphocyte Responses in Patients with Talaromyces marneffei

Infections. Infect Drug Resist. 2022;15:3381-93. Epub 2022/07/06. doi: 10.2147/IDR.S364388.

PubMed PMID: 35789796; PubMed Central PMCID: PMCPMC9250332.

44.   da Silva HB, de Salles EM, Panatieri RH, Boscardin SB, Rodriguez-Malaga SM, Alvarez

JM, et al. IFN-gamma-induced priming maintains long-term strain-transcending immunity against

blood-stage *Plasmodium chabaudi* malaria. J Immunol. 2013;191(10):5160-9. Epub 2013/10/18.

doi: 10.4049/jimmunol.1300462. PubMed PMID: 24133169.

45.   Caposio P, van den Worm S, Crawford L, Perez W, Kreklywich C, Gilbride RM, et al.

Characterization of a live-attenuated HCMV-based vaccine platform. Sci Rep. 2019;9(1):19236.

Epub 2019/12/19. doi: 10.1038/s41598-019-55508-w. PubMed PMID: 31848362; PubMed

Central PMCID: PMCPMC6917771.

46.   Liu J, Jaijyan DK, Tang Q, Zhu H. Promising Cytomegalovirus-Based Vaccine Vector

Induces Robust CD8(+) T-Cell Response. Int J Mol Sci. 2019;20(18). Epub 2019/09/13. doi:

10.3390/ijms20184457.   PubMed   PMID: 31510028;   PubMed   Central   PMCID:

PMCPMC6770317.

47.   Landau I, Killick-Kendrick R. Rodent plasmodia of the Republique Centrafricaine: the

sporogony and tissue stages of *Plasmodium chabaudi* and P. berghei yoelii. Trans R Soc Trop Med

Hyg. 1966;60(5):633-49. Epub 1966/01/01. doi: 10.1016/0035-9203(66)90010-1. PubMed PMID:

4163669.

48.   Martinelli A, Hunt P, Fawcett R, Cravo PV, Walliker D, Carter R. An AFLP-based genetic

linkage map of *Plasmodium chabaudi chabaudi*. Malar J. 2005;4:11. Epub 2005/02/15. doi:

10.1186/1475-2875-4-11.   PubMed   PMID: 15707493;   PubMed   Central   PMCID:

PMCPMC550669.

49.   Corbett AJ, Coudert JD, Forbes CA, Scalzo AA. Functional consequences of natural

sequence variation of murine cytomegalovirus m157 for Ly49 receptor specificity and NK cell

activation. J Immunol. 2011;186(3):1713-22. Epub 2010/12/29. doi: 10.4049/jimmunol.1003308.

PubMed PMID: 21187440.

50.   Wheat RL, Clark PY, Brown MG. Quantitative measurement of infectious murine

cytomegalovirus genomes in real-time PCR. J Virol Methods. 2003;112(1-2):107-13. PubMed

PMID: 12951218.

Figure Legends

Figure 1. Prior MCMV infection protects against *P. chabaudi* infection.

(A) Experimental schematic shows mice infected with MCMV-B5 or -BAC and challenged on

day 60 post MCMV infection with *P. chabaudi* AS. **(B)** Parasitemia and **(C)** weight loss were

determined after challenge. Data are representative of three independent experiments with 5

animals per group. Groups were compared using ANOVA followed by Tukey's post-test with

*, $P < 0.05$; **, $P < 0.01$. Mean shown with error bars representing SEM.

Figure 2. MCMV-B5 stimulates and maintains parasite-specific effector T cell responses.

(A) Experimental schematic shows MSP-1 specific B5 TCR Tg CD4 T cells (CTV$^-$Thy1.2$^+$) were

adoptively transferred into Thy1.1 mice that were then infected with MCMV-BAC or MCMV-B5.

Splenic T cells were gated on B5 TCR Tg (CD4$^+$CTV$^-$Thy1.2$^+$) recovered at days 15 and 55 post-

MCMV infection. **(B)** Plot, histogram and graphs show gating of proliferated B5 T cells. **(C)** Plots

and graphs show B5 Teff (CD4$^+$Thy1.2$^+$CTV$^-$, CD127$^-$CD44^{hi}) and Tmem (CD4$^+$Thy1.2$^+$CTV$^-$,

CD127^{hi}CD44^{hi}) **(D)** on day 15, or **(E)** day 55. Plots and graphs show B5 GC Tfh days 15 and 55

p.i.. Plots are concatenated from 3-5 animals per group. Data are representative of two experiments

with 3-5 mice in each group. Mean shown with error bars representing SEM and analyzed using

One-Way ANOVA followed by Tukey's post-test; *, $P < 0.05$; **, $P < 0.01$.

Fig 3. MCMV boosts long-term protection of live malaria vaccine.

(A) Experimental schematic shows infection and drug-cure live vaccine combined with MCMV

boosting strategy for protection to heterologous rechallenge. Mice were infected with *P. chabaudi*

AS and treated with two subsequent non-curative doses of chloroquine (CQ). Some groups

received MCMV-BAC or MCMV-B5 booster at day 140 (bottom timeline). All mice were

challenged with heterologous *P. chabaudi* AJ strain 200 days after live vaccination. Graph

showing **(B)** parasitemia of *P. chabaudi* AJ challengeand **(C)** serum level of *P. chabaudi* AJ-

specific IgG antibody at day 7 post-challenge. Data are representative of two independent

experiments with five mice per group showing mean, with error bars representing SEM and

analyzed using ANOVA followed by Tukey's post-test with **, $P < 0.01$; n.s., not significant.

Fig 4. MCMV-B5 promotes maintenance of *Plasmodium*-specific effector T cells. (A)

Experimental schematic shows live vaccine combined with MCMV boosting strategy with

adoptively transferred $CD4^+$ *Plasmodium* MSP1-specific T cells (B5 TCR Tg). $CD4^+$ $Thy1.2^+$

CTV^+ B5 T cells were adoptively transferred into Thy1.1 mice that were then vaccinated. Recipient

mice received MCMV-BAC or MCMV–B5 boosters at day 140, with T cell phenotyping at day

200 post-vaccination (p.v.). **(B)** Plot, histogram and graph show gating and quantification of

parasite-specific CTV^- B5 T cells recovered on d200. **(C)** Plots and graphs showing number of

Teff ($CD127^-$) and Tmem ($CD127^{hi}CD44^{hi}$) out of CTV^- B5 TCR Tg cells on day 200. **(D)** Plots

and graphs showing percentage and number of B5 memory T cell subsets gated on $CD44^{hi}CD127^{hi}$,

Tcm ($CD62L^+CD27^+$), Tem^{Early} (TemE, $CD62L^{lo}CD27^+$) or Tem^{Late} (TemL, $CD62L^{lo}CD27^-$).

Cytokines in B5 T cells were quantified and shown as histograms and graphs quantifying **(E)** IFN-

γ, **(F)** IL-2- and **(G)** TNF- positive CTV^- B5 T cells. Data are representative of two independent

experiments. Plots are concatenated from 5 animals per group, while graphs include individual

animals and show mean with error bars representing SEM. Groups were analyzed using One-Way

ANOVA followed by Tukey's post-test with *, $P < 0.05$; **, $P < 0.01$, ***, $P < 0.001$.

Fig 5. IFN-γ induced by MCMV leads to increased cytokines and protection from *P. chabaudi* infection.

(A) Experimental schematic shows mice infected with MCMV-BAC and anti-IFN-γ or isotype control was administered at indicated days (34, 36 and 38 p.i.) prior to challenge with *P. chabaudi* at day 40. Graphs show **(B)** percentage of parasitemia, **(C)** weight loss and **(D)** hypothermia upon *P. chabaudi* challenge. **(E)** Plot and graph show fraction of polyclonal effector T cells at day 15 p.c., isotype in white and anti-IFN-γ in black bars. Graphs show plasma IL-12 p40 levels **(F)** throughout MCMV infection and **(G)** after *P. chabaudi* challenge. **(H)** Graphs show chemokines and cytokines on day 7 of *P. chabaudi* challenge after MCMV-BAC and antibody treatment. Graphs include 5 animals per group and show mean with error bars representing SEM. Multiple comparisons Student *t* test and Unpaired or Mann Whitney were used for all except (F) which was analyzed with One-Way ANOVA. *, $P < 0.05$; **, $P < 0.01$, ***, $P < 0.001$; n.s., not significant.

Fig 6: IFN-γ induced by MCMV does not regulate MCMV driven T cell phenotype, but does induce IL-12 p40.

(A) Experimental schematic shows CTV-labelled B5 TCR Tg T cells were adoptively transfer into Thy1.1 (Balb/c) mice which were then infected with MCMV-B5 and received anti-IFN-γ antibody or isotype at indicated time points and spleen phenotyped for dendritic cells and B5 TCR Tg (Thy1.2$^+$) CD4$^+$ T cell at day 50 p.i.. **(B)** Plot, histogram and graph show gating and quantification of CD4$^+$Thy1.2$^+$CTV$^-$ B5 T cells. **(C)** Plots and summary bar graph show MCMV-B5-specific effector and memory T cell numbers. **(D)** Plot and summary bar graph show the number within each MCMV-B5-specific Tmem subset (Tcm, TemE and TemL) as defined by CD27 and CD62L

(E) Plot and histogram show DC and CD8α DC population gating and quantification forCD8α⁻

DC, CD8α⁺ DC and **(F)** serum IL-12p40 levels. Data shown are representative 5 mice per group

and mean shown with error bars representing SEM. The Student t test was used; *, $P < 0.05$.

Fig 7. IFN-γ induced by MCMV vector drives increase in CD8α⁺ dendritic cells, and boost of

long-term protection by live malaria vaccine.

(A) Experimental schematic shows B5 TCR Tg CD4 T cells labelled with cell trace violet (CTV)

were adoptively transferred into Thy1.1 mice that were then vaccinated with live malaria vaccine.

Recipient vaccinated mice received MCMV-B5 boosters at day 140 and then received anti-IFN-γ

antibody or isotype at indicated days (194, 196, 198) before spleen cell recovery for phenotyping

at day 200 p.v. **(B)** Plot, histogram and graph show gating and quantification of B5-epitope-

specific T cells (CTV⁻Thy1.2⁺) recovered on day 200. **(C)** Plot and summary bar graph show the

number within each vaccine and booster-B5-specific Tmem subset (Tcm, TemE and TemL) as

defined by CD27 and CD62L. * indicates significant difference between TemL subsets recovered

between groups. **(D)** Plots and graphs show gating and number of total DC population (CD3⁻

CD11cʰⁱMHCIIʰⁱ), and **(E)** CD8α⁻ or CD8α⁺ DC. Plots are concatenated from 5 animals per group.

(F) Graph of serum IL-12p40 level at day 200 p.v. before and after boosting. **(G)** Experimental

schematic for infection and drug-cure vaccine combined with MCMV boosting strategy for

protection to heterologous rechallenge. Age-matched mice were infected with *P. chabaudi* AS and

treated with two subsequent non-curative doses of chloroquine (CQ). Then they received MCMV

booster at day 140. Some mice received anti-IFN-γ antibody or isotype at indicated days (194,

196, 198) after *P. chabaudi* vaccination. All mice were reinfected with *P. chabaudi* AJ strain at

200 days p.v. Graphs showing **(H)** weight loss and **(I)** parasitemia after *P. chabaudi* AJ challenge

or infection. Comparisons of vaccinated or non-vaccinated groups are significantly different by

ANOVA and Tukey's multiple comparisons significance is not shown for the sake of simplicity

($p < 0.01$, for all time points for parasitemia and at day 9 p.c. for weight loss). **(J)** Graphs showing

parasitemia on day 5 *P. chabaudi* AJ challenge after IFN-γ neutralization (black bar) compared to

isotype (grey bar). **(K)** Graph of fraction of polyclonal effector T cells (CD127⁻) responding to

challenge (d15 p.c.). **(L) graph of** serum level of *P. chabaudi* AJ-specific IgG antibody at

indicated days (-5, 7 and 15) relative to challenge. Data shown represent 5 animals per group.

Mean shown with error bars representing SEM. Groups were analyzed using ANOVA followed

by Tukey's post-test for all except (J and K) which were analyzed with Student *t* test. *, $P < 0.05$;

**, $P < 0.01$, ***, $P < 0.001$; n.s., not significant.

S1 Fig. *Plasmodium chabaudi* MSP-1 B5 antigen-expressing MCMV vaccine vector.

(A) *P. chabaudi* merozoite surface protein MSP-1 (B5:1157–1171) B5 epitope is expressed as an

MCMV immediate early gene. B5 epitope was introduced into an MCMV-K181 bacterial artificial

chromosome (BAC) using recombineering. **(B)** The indicated organs were collected from MCMV

infected mice at different times post- MCMV infection, homogenized, and MCMV titres were

determined by plaque assay. The dashed line represents the limit of detection (L.O.D.) for each

organ. Mean ± SEM are plotted with three mice for each time point. **(C)** Time course of qPCR

assessment of MCMV infection in multiple organs. Mice were infected with MCMV-B5 or

MCMV-BAC and organs were collected at indicated times for vector quantification with n=6 per

time point. Copy number of CMV/β-actin (±SEM) ratio and L.O.D. of the assay are shown. L.O.D

was calculated using data from uninfected animal. (**D**) Mice were infected with MCMV-BAC or

MCMV-B5 and splenocytes were recovered 15 days later for MCMV-induced polyclonal T cell

phenotype. Plots show gating and graphs show numbers of polyclonal CD4⁺ T cells effector

(CD127⁻) and memory (CD44^{hi}CD127^{hi}) phenotype cells recovered after MCMV-B5 compared to

MCMV-BAC. **(E)** Enumeration of MCMV-specific m78-Tetramer-positive CD4 T cells upon

MCMV-B5 compared to MCMV-BAC from day 15 MCMV-infected mice. Plots represents

control CLIP tetramer or m-78-tetramer staining after MCMV-B5. Graph shows number of

MCMV-specific m78 Tet+ T cells (±SEM). **(F)** Enumeration of MCMV-specific m45-Tetramer-

positive CD8 T cells. Mice were infected with MCMV-B5 and splenocytes were recovered 11 and

60 days later for CD8 T cell quantification by staining with m45 tetramers. Plots and graph show

m45-Tet^{+} CD8^{+} T cells from uninfected and MCMV-B5 infected mice gating on vector specific T

cells (+/- SEM). Data shown are representative of two experiments with 3 or 5 mice each. Groups

were analyzed using One-Way ANOVA followed by Tukey's post-test with *, $P < 0.05$;

**, $P < 0.01$.

S2 Fig. Persistent MCMV-B5 infection induces vector-specific T cell response and may

promote continued B5 TCR Tg T cell proliferation.

(A) Experimental schematic shows mice were infected with MCMV-B5, and splenocytes were

recovered 11 or 60 days later to quantify MCMV m-78-specific Tetramer^{+} CD4 T cells. **(B)** Plots

and graph show uninfected and m78 tetramer-specific T cell numbers. Virus-specific CD4^{+} T cells

were identified using m78/I-A^{d} tetramers, with peptide (SQQKMTSLPMSVFYS) and CLIP/I-A^{d}

as tetramer control (PVSKMRMATPLLMQA). **(C)** Plots and graphs show the number of MCMV-

specific-Teff (CD44^{+}CD127⁻) and Tmem (CD44^{hi}CD127^{hi}) in the spleen gated on MCMV m78-

tetramer CD4^{+} T cell at day 11 and 60 p.i.. Data shown are representative of two experiments with

5 mice each and error bars representing SEM. **(D)** Experimental Schematic shows MCMV-BAC

or MCMV-B5 infection of different groups of age-matched Thy1.1 mice at indicated times (0, 30

and 50 days) followed by adoptive transfer of CTV$^+$ B5 TCR Tg CD4 T cells to all groups 5 days

before phenotyping, and 55 days after first infection. **(E)** Plots and graphs showing gating and

frequency of CTV-B5 T cells out of CD4 T cells, or number of proliferated B5 epitope-specific

CD4 T cells in response to MCMV-BAC or MCMV-B5 infection at different time points. Data are

representative of three independent experiments with 5 animals per group. Mean shown with error

bars representing SEM. Groups were analyzed using One-Way ANOVA followed by Tukey's

post-test with *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

1012

S3 Fig. Decay of *P. chabaudi* protection by days 200

(A) Experimental schematic shows age-matched mice were infected with *P. chabaudi* AS for

assessment of protection to heterologous challenge with *P. chabaudi* AJ 60 or 200 days later.

Graphs show **(B)** the percentage of parasitemia (iRBC/RBC) after heterologous challenge, and **(C)**

P. chabaudi AJ-specific IgG antibody serum level at day 7 post-challenge. Data are representative

of two independent experiments with five mice per group showing mean, with error bars

representing SEM. Groups were analyzed using One-Way ANOVA followed by Tukey's post-test

with *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

1021

S4 Fig: Evaluation of neutralizing effect of anti-IFN-γ antibody

(A) Experimental schematic shows mice were infected or not infected with MCMV-BAC to assess

the neutralizing effect of anti-IFN-γ naïve mice or in MCMV-induced protection to *P. chabaudi*.

Blocking of IFN-γ *in vivo* with anti-IFN-γ antibody or isotype control antibody was performed at

days 34, 36 and 38 prior to the challenge or infection with *P. chabaudi* at day 40 in both MCMV

infected and naïve mice. Graphs show percentage of parasitemia in **(B)** no MCMV mice and **(C)**

MCMV vaccinated mice. Relative CXCL-10 mRNA level normalized with β-actin from Raw

264.7 cells cultured for 24h with sera from: (**D**) 2 days after anti-IFN-γ administration, (**E**) day 7

post *P. chabaudi* challenge, (**F**) 2 days after anti-IFN-γ administration in presence of rIFN-γ

(25ng/ml), and (**G**) day 7 post *P. chabaudi* challenge administration in presence of rIFN-γ

(25ng/ml). Data shown represent 5 animals per group and mean shown with error bars representing

SEM. Groups were analyzed using Student *t* test (C, D and F) or One-Way ANOVA (B, E and G)

with *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

1035

S5 Fig. Combined IL-12 and IL-18 neutralization did not affect protection induced by

MCMV.

(A) Experimental schematic shows mice infected with MCMV-BAC and the combination of anti-

IL-12p40 and anti-IL-18 antibody or isotypes were given post MCMV infection. Graphs show

percentage of parasitemia **(B)**, weight loss **(C)**, and hypothermia **(D)** upon combined IL-12 and

IL-18 neutralization. Data shown represent 5 animals per group and mean shown with error bars

representing SEM and analyzed using multiple comparisons Student *t* test; n.s., not significant.

1043

S6 Fig. IFN-γ induced by MCMV vector does not change most antigen-presenting cells types

and their activation significantly.

(A) Experimental schematic shows mice infected with MCMV-B5 and received anti-IFN-γ

antibody or isotype at indicated time points. Splenic antigen-presenting cells types (monocytes,

macrophages and inflammatory monocytes) and NK cells were phenotyped at day 50 p.i.. **(B)**

Plots and graphs show gating and quantification of CD3$^-$CD11b$^-$Ly6C$^+$, CD3$^-$ CD11b$^+$Ly6C$^+$,

CD3$^-$ CD11b$^+$Ly6C$^-$ cells. (C) Histograms and graphs show MFI of CD3- CD11b+Ly6C-

expressing activation marker MHCII. (D) Histograms and graphs show MFI of CD3-

CD11b+Ly6C- expressing activation marker CD86. (**E**) Plots and graphs show gating and

quantification of CD3$^-$ DX5$^+$MHCII$^-$ NK cells.

(**F**) Experimental schematic shows Thy1.1 mice vaccinated with live malaria vaccine. Then

vaccinated mice received MCMV-B5 boosters at day 140 and then received anti-IFN-γ antibody

or isotype at indicated days (194, 196, 198) before spleen cell recovery for cells phenotyping at

day 200 p.v. **(G)** Plots and graphs show gating and quantification of CD3$^-$ CD11b$^-$Ly6C$^+$, CD3$^-$

CD11b$^+$Ly6C$^+$, CD3$^-$ CD11b$^+$Ly6C$^-$ cells. **(H)** Histograms and graphs show MFI of macrophages

(CD3$^-$CD11b$^+$Ly6C$^-$) expressing activation marker MHCII. **(I)** Histograms and graphs show MFI

of inflammatory monocytes (CD3$^-$ CD11b$^+$Ly6C$^+$) expressing activation marker MHCII. (**J**) Plots

and graphs show gating and quantification of CD3$^-$DX5$^+$MHCII$^-$ NK cells. Data shown are

representative 5 mice per group. Plots concatenated per group and mean shown with error bars

representing SEM. The Student *t* test or One-Way ANOVA were used.

1064

S7 Fig. An example of a gating strategy used to identify polyclonal CD4$^+$ T cells subsets and

B5-specific CD4$^+$ T cells subsets

(Top row) Plots show gating strategy for polyclonal CD4$^+$ T cell subsets Teff (CD4$^+$CD127$^-$

CD44^{hi}), Tmem (CD4$^+$CD127^{hi}CD44^{hi}); and memory T cell subsets gated on CD44^{hi}CD127^{hi} as

Tcm (CD62L$^+$CD27$^+$), Tem^{Early} (TemE, CD62L^{lo}CD27$^+$) or Tem^{Late} (TemL, CD62L^{lo}CD27$^-$).

(Bottom row) Plots show gating strategy for B5-specific T cells: Teff (CD4$^+$Thy1.2$^+$CTV$^-$,

CD127$^-$CD44^{hi}), Tmem (CD4$^+$Thy1.2$^+$CTV$^-$, CD127^{hi}CD44^{hi}) and B5 memory T cell subsets

gated on CD44^{hi}CD127^{hi}, Tcm (CD62L$^+$CD27$^+$), Tem^{Early} (TemE, CD62L^{lo}CD27$^+$) or Tem^{Late}

(TemL, CD62L^{lo}CD27$^-$).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7