1 **Sex or sex role – unravelling the evolutionary route of sex-specific senescence**
2

Freya A. Pappert[1,2,*], Arseny Dubin[1], Guillermo Torres[2], Olivia Roth[1,2]

[1]Marine Evolutionary Biology, Zoological Institute, Christian-Albrechts-Universität Kiel, Am Botanischen Garten 1-
9, 24118 Kiel, Germany
[2]Evolutionary Ecology of Marine Fishes, Helmholtz-Centre for Ocean Research Kiel (GEOMAR), 24105 Kiel,
Germany
[3]Institute of Clinical Molecular Biology (IKMB), Kiel University, Rosalind-Franklin-Straße 12, 24105 Kiel, Germany
*Corresponding author: Freya Adele Pappert, Marine Evolutionary Biology, Zoological Institute, Christian-
Albrechts-Universität Kiel, Am Botanischen Garten 1-9, 24118 Kiel, Germany. Telephone: +49 4318804224; E-
mail: fpappert@zoologie.uni-kiel.de
12
13

14 Keywords: life-history traits, sex-role reversal, parental investment, male pregnancy, Syngnathids, sex-biased gene
15 expression

16
Abstract

Sex-specific lifespan dimorphism is observed in many species, with females often outliving males. This is thought

to be due to sex-specific and unequal resource allocation between reproduction and maintenance. However, most

studies on ageing focus on animals with traditional sex roles, where females have the highest energy burden with

both egg production and parental care. This leaves a gap in our understanding of how resource allocation and

senescence are linked in species with non-traditional roles. To address this, we investigated the relationship

between sex (biological) and sex role (behavioural) on resource allocation trade-offs in somatic maintenance in the

broad-nosed pipefish *Syngnathus typhle*. This species is sex-role reversed, with males acting as the primary

caretaker for offspring through their unique male pregnancy, providing the opportunity to decouple the roles of sex

and sex role in resource allocation. Our results suggest that while some gene groups are influenced by sex roles

(e.g., genome maintenance and immune defence), others depend on biological sex and hormones (e.g., energy

and fat storage), with differences becoming more pronounced with age. These findings challenge traditional ageing

theories and improve our understanding of how sex-specific differences in ageing are shaped.

1. Introduction

Ageing is manifested as progressive physiological deterioration accompanied by decreasing or ceasing fertility that

correlates with an induced likelihood of mortality (Viña, Borrás, and Miquel 2007). Some species display

accelerated or rapid ageing, for instance, killifish with 13 weeks lifespan or pacific salmon senesce rapidly after

reproducing. Other species show slow or negligible senescence or have an extended lifespan such as rougheye

rockfish (up to 200 years) or the greenland shark (roughly 400 years) (Stenvinkel and Shiels 2019; Trifonova et al.

2018). However, differences in lifespan vary not only across species but also between sex (Jones et al. 2014).

Although the mechanistic pathways for ageing, also referred to as the "Hallmarks of Ageing", are thought to be

conserved (López-Otín et al. 2013), there is accumulating evidence that ageing leads to sex-specific phenotypes

and age-related diseases (Gubbels Bupp 2015; Vom Steeg and Klein 2016; Zuk 2009). For example, a study

looking at 101 mammal species, revealed that female lifespan is on average 18.6% longer than that of males (Lemaître et al. 2020). The resource allocation theory predicts that the higher sex-specific lifespan dimorphism observed in Mammalia, Insecta, Reptilia, and Actinopterygii compared to Arachnida, Aves, Chondrichthyes, and Amphibia is due to selection acting differently on males and females, resulting in shifted investments for growth and reproduction based on the reproductive roles of the sexes in the species (Xirocostas, Everingham, and Moles 2020). Males (defined as sperm producers) and females (egg providers) invest in different traits, behaviours and reproductive demands which lead to life history trade-offs and distinct mortality risks (Brooks and Garratt 2017; Janicke et al. 2016). On a similar line of thought Kirkwood's disposable soma theory states "longevity requires investments in somatic maintenance that reduces the resources available for reproduction" (Kirkwood and Rose 1991). Here, investment in reproduction is a traded-off with investment in somatic maintenance, thus reducing self-maintenance and determining lifespan.

Mammalian female reproductive requirements often include resource allocations for egg production, gestation, lactation and an overall higher parental investment as opposed to males' cheap sperm production, secondary sexual traits, and territory defence (Lemaître et al. 2020; Shaw and Shaw 2009). In such light, females may be more reproductively constrained by the time they devote to their offspring and will favour investing in longevity and lifespan, compared to males with a lower burden of parental care, allowing for multiple matings in a short period of time and thus focusing more on their reproductive success (Clutton-Brock and Isvaran 2007). This is, however, an example of stereotypical mammalian sex roles, where males compete for access to mates with their fitness being maximized by increasing mating rates, and where females are choosy with fitness being positively correlated with longevity (Bateman 1948; Rolff 2002; Williams 1957). In the animal kingdom, sex roles are not constrained and vary between species and even across an individual's lifetime based on different degrees of parental investment (Benvenuto et al. 2017; Janicke et al. 2016; Smith and Wootton 1995). For example, in the 85 teleost species that show parental care, the majority is dominated by paternal care, such as egg guarding, and oral brooding to full viviparity where the males will endure the largest amount of work (Smith and Wootton 1995; Wilson et al. 2001). Some of the most devoted male parents can be accredited to Syngnathids (seahorses, pipefishes and seadragons), which have evolved a range of brooding forms from simple attachment of eggs to the male tail to highly specialized male pregnancy with placenta-like structures analogous to those found in eutherian mammals, that provide embryos with nutrients, oxygen and immunological components (Beemelmanns and Roth 2016; Ripley and Foran 2009; Roth et al. 2011; Whittington and Friesen 2020; Wilson et al. 2001). Thus far experimental studies on parental investment have focused primarily on the mothers' fitness, linking it directly to the offspring's fitness, since females typically invest more not only in egg production but also in caring or nurturing (Wolf and Wade 2001). However, the range of sex role diversity in nature allows us to explore other avenues of resource allocation trade-offs and their effects on sex-specific senescence. Studying species with the varying spectrums of sex roles allows to test the

outcome of theories of ageing that were mainly developed for evolutionary consequences of conventional sex roles.

Unfortunately, there is scarce literature on sex-specific ageing in teleosts and fish in general (Stenvinkel and Shiels

2019). Much of the gerontology research on the mechanisms of ageing is biased toward a few species such as

roundworm (*Caenorhabditis elegans*), fruit fly (*Drosophila melanogaster*), mouse (*Mus musculus*) and rat (*Rattus

norvegicus domestica*) (Holtze et al. 2021; Mitchell et al. 2015). Studying the immense variety of species with sex

lifespan dimorphism would further our understanding of the evolutionarily conserved mechanisms of ageing and

improve the comparative approach between species (Holtze et al. 2021). Fish make up approximately half of all

existing vertebrate species (over 20,000) and not only have various forms of mating systems and parental

investments, but also have various forms of ageing from rapid, gradual, and negligible senescence (Holtze et al.

2021; Trifonova et al. 2018). They also make good comparative models with mammals as they have similar organs,

have an adaptive immune system and may suffer similar old age pathologies as senescing mammals (Froehlich et

al. 2013; Reznick, Ghalambor, and Nunney 2002). Nonetheless, while mammals and birds tend to have fairly

constant fecundity across age groups, fish fecundity increases with age as they have indeterminant growth and

thus can increase the number of eggs and offspring (Barneche, White, and Marshall 2018). This might cause

different life-history choices and change the way they evolved ageing compared to mammals, making it worth

investigating in fish.

In this study, we compared sex-biased gene expression between young and old broad-nosed pipefish *Syngnathus

typhle*, a sex-role reversed species with male pregnancy, providing the unique opportunity of decoupling the role

of the female sex (defined as the contribution of eggs) and pregnancy (body reshaping and energy allocation) (Roth

et al. 2011). Male pregnancy has evolved only once in vertebrates and from there on has diversified across

approximately 300 Syngnathidae species (Hamilton et al. 2017; Stiller et al. 2022). Syngnathids have also evolved

a gradient of mating systems including sex-role reversal; e.g., in broad-nosed pipefish, their brood pouch is size-

limited, and as the females produce on average more eggs than males can brood during the same period

(Sagebakken, Ahnesjö, and Kvarnemo 2016), males end up being the sex that is more reproductively constrained

(Mobley et al. 2018). Larger brood size and limited resource availability influence the cost of male care, making

embryo mortality inversely proportional to male condition (male mortality increases with a larger brood size)

(Whittington and Friesen 2020). In such role-reversed species, males will be the choosy sex whereas females will

be ornamented and actively court males (Mobley et al. 2018; Roth et al. 2011).

Using this unique species, we aimed to address two major gaps in our understanding of the evolutionary route to

sex-specific senescence. First, how does sex (defined as the contribution of eggs or sperm) or rather sex-role

(range of behaviours generally associated with being male or female) influence resource allocation trade-offs in

somatic maintenance? Research studying resource allocation theory and the effects of senescence on males and

females has been biased for animal models with conventional sex roles, where females generally have the highest

energy burden. Not allowing us to disentangle whether the evolution of a longer female lifespan is predominantly

due to sex hormones and sex-chromosomes, or due to the resource allocation trade-off in sex-role behaviours

such as parental investment. Second, if ageing theories are studied in animals where fertility declines with age to

various degrees between males and females, how does senescence affect somatic and germline maintenance in

teleost species where fertility increases proportionally with advancing age?

In this study, we used full transcriptome mRNA sequencing (RNA-Seq) to analyse gene expression in different

tissues (brain, head kidney and gonads) between young and old female and reproductively active (pregnant) and

non-reproductively active males. By identifying sex-biased gene expression and subsequent expression changes

throughout life and reproduction activity, we aimed to understand the importance of sex and sex role in determining

resource allocation in ageing. Has sex-role reversal as a consequence of increased male parental investment (i.e.,

male pregnancy) led to males investing more in longevity and females investing in reproduction? Alternatively, is

biological sex a more important feature for lifespan dimorphism? We hypothesized that sex-lifespan dimorphism is

more heavily influenced by sex-specific life-history traits, than by the biological sex. Our data revealed that while

sex roles may influence some gene groups and invert their expression when moving from conventional to sex-role

reversal, the activity of other gene groups may depend on the biological sex. This illuminates the importance of

disentangling sex and sex roles when we aim to understand sex-specific life-history strategies, sex-biased gene

expression and their consequences on longevity.

2. Methods

2.1 Sample collection

Broadnosed pipefish *Syngnathus typhle* were caught in the south-western Baltic Sea (54°39'N; 10°19'E) in spring.

They were brought to GEOMAR institute, where body mass and total body length were measured. The animals

were then euthanized with an overdose of MS-222 (500 mg/l; Sigma-Aldrich). Carcasses were dissected and

different tissues (brain, head kidney and gonads) were sampled and were immediately preserved in RNAlater.

Samples were kept at 4°C for three days, before transferring them to -20°C for long-term storage.

2.2 RNA isolation and Illumina sequencing

For our RNA-seq analysis, we extracted RNA from brain, head kidney, ovaries and testes, as gene expression is

known to differ between tissues. Gonads are frequently used in sex-biased gene expression studies, as sex organs

can reveal genes vital for sex-specific development or fertility regulation (Whittle and Extavour 2019). Also, the

brain has shown sex and age-specific gene expressions (Hale et al. 2018; Mayne et al. 2016). We additionally

used head kidney, which is unique to teleost fish but is analogous to the adrenal gland, producing both red blood

cells for oxygen transport and white blood cells for immune defence (Geven and Klaren 2017). Its clear role in

immune regulation makes it the perfect organ to focus on genes displaying sexual immune dimorphism. The RNA

extraction was performed using RNeasy Mini Kit (Qiagen, Venlo, Netherlands) according to the manufacturer's

protocol. As gene expression is known to differ between tissues, extraction yields were quantified using NanoDrop

ND-1000 spectral photometer (Peqlab, Erlangen, Germany), RNA was then stored at -80°C. Library preparation

(TruSeq stranded mRNA Kit by Illumina) and sequencing (RNA Seq NovaSeq6000 S4, 2x150bp, >17M reads per

sample) were carried out at the Institute of Clinical Molecular Biology (IKMB).

2.3 Transcriptome analysis

The resulting reads were quality-controlled using FastQC v.0.11.9 (Andrews 2010) and trimmed using Fastp

v.0.20.1 (Chen et al. 2018). Reads were aligned to a whole genome assembly of *Syngnathus typhle* (BioProject

ID: PRJNA947442) using STAR v.2.7.9a (Dobin et al. 2013). Transcript abundance was quantified with

TPMCalculator (Alvarez et al. 2019) in both raw read counts and transcripts per million (TPM), with counts

normalised for library size and gene length. We used Orthofinder v.2.4.0 (Emms and Kelly 2019) to assign

information to the annotated genes with these species lists (accession included): *Takifugu rubripes*

(rubripes.fTakRub1.2), *Syngnathus rostellatus* (GCF_901709675.1, fSynAcu1.2), *Syngnathus accus*

(GCF_901709675.1, GCF_901709675.1_fSynAcu1.2)*, Nerophis ophidion* (unpublished), *Oryzias latipes*

*(*ASM223467v1)*, Hippocampus erectus (*PRJNA613176)*, Hippocampus comes* (H_comes_QL1_v1,

GCA_001891065.1)*, Lepisosteus oculatus* (LepOcu1)*, Gasterosteus aculeatus* (BROADS1)*,* and *Danio rerio*

(GRCz11). Differential gene expression analysis of the RNA-Seq data from Illumina was done in Rstudio v.3.4.0

(R Core Team 2022), using the limma:voom package v.3.54.1 (Ritchie et al. 2015) on *S. accus* orthology

annotations. After scaling and normalizing with edgeR package v.3.14.0 (Robinson, McCarthy, and Smyth 2009),

we discarded any genes with less than 10 counts in at least 3 libraries. To look at the gene expression differences

between the organs we did a cluster analysis, detecting transcripts that were unique for each organ for subsequent

enrichment and functional analysis. The limma:voom pipeline was run separately for counts in brain, head kidney

and gonads, as the effects of other variables might be harder to detect with significant differences between tissues.

Within a single organ a multi-level analysis with categorical paired comparisons was performed, to investigate

differential gene expression for sex, age and pregnancy. This allowed to compute the contrasts between our single

variables of interest, whilst still considering the variable interactions within the linear model analysis of limma:voom.

2.4 Statistical analysis

Morphological data comparison between groups was analysed by a two-way ANOVA, with statistical significance

set at $P < 0.05$. To compare differentially expressed genes between the group variables (young and old females,

young and old reproductively active thus pregnant and non-reproductively active males), we did a Principal

Component Analysis (PCA) with the "regularized log transformation procedure" (rld) transformed expression values

to check for outliers and tissue variance (Figure S1). The few detected outliers were removed after further

consideration, and subsequently, samples clustered according to tissue type (Figure S1). To estimate differences

between factors such as sex, age and pregnancy, we separated the DEGs dataset for the three separate organs

(Figure 1D and Figure S3). To check if transcript expression varied significantly, we also applied a PERMANOVA

(Permutational Multivariate Analysis of Variance Using Distance Matrices) using function "adonis2" and method

"bray".

2.5 Orthology and enrichment analysis

For gene ontology analysis, we annotated the differentially expressed genes using a homology-based search of

Danio rerio with OrthoFinder. The overall number of DEGs ended up being reduced as it was not possible to find

orthologues for all transcripts. *Danio rerio* orthologues of DEGs were uploaded to g:Profiler (last access date

January 2023) (Raudvere et al. 2019) to examine the functional profiling of transcripts from the cluster analysis

based on tissue type, and the DEGs for young male (M.y.), young female (F.y.), old male (M.o.) and old female

(F.o). The thresholds used were: Organism "*Danio rerio* (Zebrafish)", Statistical domain scope set at "All known

genes", significance threshold "g:SCS threshold" at 0.05, numeric IDs treated as "ENTREZGENE_ACC", the data

sources were limited to GO biological processes and biological pathways the databases Kyoto Encyclopedia of

Genes and Genomes (KEGG) and Reactome (REAC) (Tables S1). A more targeted approach was used to

investigate the presence and expression pattern of immune-related genes. Following results from our gene

enrichment analysis in the head kidney, with terms such as innate immune system (REAC:R-DRE-168249) and

adaptive immune system (REAC:R-DRE-1280218) enriched, we searched in our DEGs dataset what immune

genes males and females had differentially expressed. As various genes in these pathways had numerous overlaps

with other non-immune related GO terms and some more unique immune transcripts had been missed out in the

GO analysis, we opted to use immunological keywords (see Table S4 for keywords) based on literature reviews

on the immune response in our whole dataset (Chaplin 2010; Yuan et al. 2014). The keywords that were found in

our dataset were grouped under umbrella terms as adaptive or innate immune response, with which we then

compared their differential expression for sex and age.

3. Results

3.1. Sample description and transcriptome analysis of differential gene expression in brain, gonads and head

kidney

In our study we used 10 females and 20 males; the latter included 10 pregnant and 10 not mated yet. All fish were

separated into old and young categories, based on morphological measurements (Figure S2). Body length and

weight are frequently used in fisheries to roughly determine age, as the vast majority of fish have indeterminate

growth with size and age being correlated (Barneche, White, and Marshall 2018). The average weight of the

sampled pipefish was 0.65 g (SD = 0.146) in young females (F.y.) and 0.6 g (SD = 0.082) in young males (M.y.),

whilst 1.71 g (SD = 0.191) in old females (F.o.) and 1.63 g (SD = 0.243) in old males (M.o.). The average total body

length was 12.3 cm (SD = 1.02) for F.y. and 11.56 cm (SD = 0.682) for M.y., 16.66 cm (SD = 0.764) in F.o. and

16.82 cm (SD = 1.41) in M.o. A two-way ANOVA confirmed that both total body length and weight were highly

significant for age (length $P < 2.8e-12$ and weight $P < 5.65e-15$), but not sex ($P > 0.05$).

RNA sequencing expression profiles resulted in a total number of 26072 transcripts with an average of 1M reads

per sample, of which 17200 remained after filtering out genes with less than 10 counts in at least 3 libraries. A

cluster analysis (Figure 1B) revealed groups of genes expressed mostly in either brain, head kidney or gonads

(Figure 1B). We then identified functional categories defining each organ of the pipefish based on clusters sharing

similar expression patterns (Table S1). Genes involved in neuronal pathways were mostly localized in brain, as

were immune response genes in head kidney (Figure 1C and Table S1). Gonads including both testes and ovaries

had mostly pathways involved in genome maintenance and proteostasis regulation (Figure 1B). Additionally, all

organs did have various metabolic related pathways enriched (Figure 1C). While there were strong differences

between organs, a PERMANOVA analysis did not reveal any significant effect of pregnancy in any of the single

tissues. Also, age did not quite reach significance in any of the tissues. However, sex was highly significant ($P <$

0.001) in both gonads and head kidney. Subsequent PCAs of each tissue confirmed these results (Figure S3 for

brain and gonads), but for the PCA of head kidney there was also some indication that PC2 may reflect the size

and thus age of the pipefish (Figure 1D).

Figure 1: A) A graphical representation of *Syngnathus typhle* divided per body compartment. Picture modified from (Schneider, Woltering, Adriaens, & Roth, 2022). B) DEGs organised in clusters according to their expression in different tissues with B for brain, HK for head kidney and G for gonads. C) Summary table of REAC annotated functional pathways from the group of clusters sharing similar expression patterns. The colours correlate with the regression line for each cluster, with yellow corresponding to brain tissue, green to head kidney and blue to gonads (testis and ovaries together). D) Principal component analysis in head kidney for all individuals. The colours represent sex with blue for males and pink for females, while the shapes correspond to age category, with a circle for old and a triangle for young.

3.2. Age-dependent increase in sex-biased gene expression in head kidney: Metabolic and cell maintenance

pathway analysis.

From the three distinct tissues, the differential gene expression analysis revealed that the head kidney had the

highest count of significant (adj.P < 0.05) DEGs between males and females, with sex-biased gene expression

differing even further when comparing young and old. The contrast M.y. vs. F.y. revealed 1919 DE transcripts, of

which 1077 were upregulated in F.o. and 842 in M.y., and for M.o. vs. F.o. 3828, of which were 2114 upregulated

in F.o. and 1714 in M.o. This suggests that sex-biased gene expression becomes more prevalent with age in

pipefish. We wanted to understand how many of the significant DEGs were sex-specific across all ages, and which

transcripts were biased only in young or old pipefish. We found that 569 transcripts were upregulated in males and

859 in females, irrespective of age (Figure 2A). On the other hand, 318 transcripts were only differentially expressed

for sex in young pipefish, of which 182 were upregulated in M.y. and 136 if F.y. (Figure 2B), and 2059 DEGs in old

pipefish, with 980 in M.o. and 1079 in F.o. (Figure 2C).

The following results of a functional enrichment analysis for both DEGs that were biased for sex, irrespective of

age, and for those who were uniquely expressed in either sex for young or old, are summarised in Figure 2D for

REAC and KEGG pathways (see supplements Table S1 for all results). There was a general enrichment in

categories related to different metabolic pathways (metabolism of RNA - REAC:R-DRE-8953854, metabolism of

lipids - REAC:R-DRE-556833, REAC:R-DRE-392499, fatty-acid metabolism - REAC:R-DRE-8978868) (Figure

2D). Both F.y. and F.o. had an overrepresentation of genes enriched for metabolic pathways involved in fat storage,

such as fatty acid metabolism (REAC:R-DRE-8978868) and lipid metabolism (REAC:R-DRE-556833), peroxisomal

lipid metabolism (REAC:R-DRE-390918) (Figure 2D). A closer look revealed upregulated genes such as

peroxisome proliferator-activated receptor gamma (*pparγ*) a nuclear receptor important in the regulation of fatty

acid metabolism balancing the storage and utilisation of fatty acids in the body (Rosen et al. 1999), or *adiponectin*

receptor 2 (*adipor2*) involved in the regulation of glucose (Table S2 and S3). F.o. differentially expressed genes

such as hormone-sensitive *lipase* (*lipea*) and *lipoprotein lipase* (*lpl*), that encode enzymes that break down fats and

regulate lipid metabolism.

Male pipefish had pathways such as metabolism of RNA (REAC:R-DRE-8953854), DNA metabolic process

(GO:0006259), translation (REAC:R-DRE-72766), and ribosome (KEGG:03010) that were enriched (Figure 2D).

M.y. and M.o. also both had an overrepresentation of genes involved in the non-sense mediated decay (NMD)

(REAC:R-DRE-2990846) pathway (Figure 2D). This important pathway maintains the integrity of the genetic

information by preventing the production of non-functional proteins, which is vital for correct intercellular

communication (Hug, Longman, and Cáceres 2015). However, M.o. additionally had pathways enriched for DNA

repair (REAC:R-DRE-73894), DNA double-strand break repair (REAC:R-DRE-5693532), and SUMOylation of DNA

damage response and repair (REAC:R-DRE-3108214) (Figure 2D). For example, Mo's expressed more *rad* genes

compared to any other group, such as *rad50*, *rad51b*, *rad52*, and *rad54* (Table S3). *Rad* genes are all involved in

DNA repair, specifically in the process of homologous recombination (Henning and Stürzbecher 2003; Jalan, Olsen,

and Powell 2019). M.o. also differentially expressed *ATM serine/threonine kinase* and *checkpoint kinase 2* (*chek2*),

both generally responsible for activating several downstream targets involved in repairing damaged DNA and

preventing the development of cancer (Maréchal and Zou 2013; Mustofa et al. 2020). Whilst we found that M.y.

differentially expressed *Sirtuin 2*, which is known to play a role in several processes from cell survival and DNA

repair, to energy metabolism (Wang et al. 2019). M.o. had *mTOR kinase* upregulated, which can lead to increased

cell growth and proliferation, but they also expressed *deptor*, which is a negative regulator of the mTOR signalling

pathway acting as a feedback inhibitor (Catena and Fanciulli 2017). The downstream targets of mTOR are

numerous (i.e., cell growth, metabolism, autophagy, translation initiation) and depend on sex hormones and

environmental energy availability (Efeyan, Comb, and Sabatini 2015). Nonetheless, females also had various cell

signalling pathways important for survival and maintenance, like FoxO signalling (KEGG:04068), with genes such

as *forkehad box O4* (*foxo4*) and *peroxisome proliferator-activated receptor gamma, coactivator 1 alpha* (*ppargc1a*)

homologous to *foxo1* upregulated in F.o.; or mTOR signalling (KEGG:04150), with express *lamtor1* and *lamtor2*

also upregulated in F.o., which act as scaffold-proteins by binding to mTOR complex1 promoting its activity (de

Araujo et al. 2017); and autophagy (REAC:R-DRE-9612973), important in eliminating damaged proteins, with F.o.

DE example *ATG5 autophagy related 5 homolog* (*atg5*) (Figure 2D).

Figure 2: Heatmaps of DEGs and table annotating pathways of the transcripts from the heatmap. A) Heatmap of the 399 filtered genes (adj.P.val < 0.05, logFC < -1 and logFC > 1) that were differentially expressed for sex in both young and old pipefish, of which 138 upregulated and 269 downregulated in males. B) Heatmap of 87 filtered (adj.P.val < 0.05, logFC < -1 and logFC > 1) transcripts that were uniquely differentially expressed for sex only in young pipefish (39 M.y. and 48 F.y.). C) Heatmap of 527 filtered (adj.P.val < 0.05, logFC < -1 and logFC > 1) transcripts that were uniquely expressed for sex only in old pipefish (348 Mo and 179 Fo). D) Summary table of the annotated pathways from the functional enrichment analysis of DEGs in head kidney for DEGs from the heatmap.

3.3. Sexual immune dimorphism in head kidney reveals old male-biased gene expression

Many aspects of immune regulation in vertebrates are fairly well conserved and most of the major immune genes exist in teleost fish but may have multiple copies (paralogues) of particular genes (Bjørgen and Koppang 2021; Zou and Secombes 2016). The GO enrichment analysis found immune-related pathways, such as innate immune system (REAC:R-DRE-168249) and adaptive immune system (REAC:R-DRE-1280218) for each group (Figure 2D). Overall, we found that males had more upregulated immune genes compared to female, in both young and old (Table S5). Young pipefish differentially expressed 43 significant genes, of which 25 were upregulated in M.y. and 18 were upregulated in F.y., while old pipefish differentially expressed 71, of which 48 were upregulated in M.o. and 23 in F.o. (Table S5).

We identified which innate and adaptive immune genes were upregulated in either M.y., F.y., M.o., or F.o. We

found the following anti-inflammatory markers expressed both in young and old males: *interleukin 13 receptor alpha*

1 (*il13ra1*); two copies of *interleukin-1 receptor type-1-like* (*il1rl1*), which acts as a decoy receptor for il1 by binding

to it but not transducing the signal (Zou and Secombes 2016), thus neutralizing its activity; *tumour necrosis factor*

alpha-induced protein 8-like 2b (*tnfaip8l2b*) downregulated the production of pro-inflammatory cytokines (NF-kB)

(Liu et al. 2021) (Table S5). As for young and old females, they expressed *interleukin 11 receptor alpha* (*il11ra*),

involved in the transduction of the signal of IL-11, an anti-inflammatory cytokine and interleukin 10 (*il10*), which

suppresses the production of pro-inflammatory cytokines such as TNF-alpha, IL-1, and IL-6 also inhibit the

activation of immune cells such as macrophages, which are important for the immunity against intracellular and

extracellular pathogens (Zou and Secombes 2016). Pro-inflammatory cytokines expressed in males were:

interleukin 12 receptor beta 2a (*il12rb2*), involved in the transduction of the signal of IL12 (Zou and Secombes

2016); *tumour necrosis factor receptor superfamily member 1B* (*tnfrsf1b*), involved in the transduction of TNF-alpha

signal (Baker and Reddy n.d.); and several *chemokine receptors* (*cxcr1* and *cscr4a*) (Table S5). They also

expressed multiple *neutrophil cytosolic factors (ncf1, ncf2* and *ncf4*), where NCF1 is considered a pro-inflammatory

protein that is involved in the production of reactive oxygen species important for killing pathogens (Holmdahl et al.

2016).

However, older males had additional pro-inflammatory elements expressed: *tumour necrosis factor receptor*

superfamily (*tnfrs*), *member 1a* and *21*; *interleukin 17 receptor C* (*il17rc*), which together with other IL-17 receptors

(*il17ra* and *il17rd*) are involved in the transduction of IL-17A and contribute to an inflammatory response (Zou and

Secombes 2016); toll-like receptors (TLR) were also only upregulated in old males, such as *tlr3* and *tlr18* (Table 1

and Table S5), the latter is a homolog of human *tlr1*, both important in initiating an immune response against viral

infections by recognising double-stranded RNA (Kawasaki and Kawai 2014; Li et al. 2017). In comparison, the pro-

inflammatory expression did not change all that much between young and old females, with them both expressing

il17ra1a, part of the il17 receptor complex described above and *interferon-gamma receptor 1-like* (*ifngr1l*) (Table

S5). Yet, F.y. did also have *interleukin 6 transducer* (*il6st*) upregulated, which is a transmembrane protein involved

in the signal transduction of cytokines belonging to the IL-6 family (Grayfer and Belosevic 2012).

F.o. had several variants of major histocompatibility complex (MHC) class 1 upregulated compared to males (Table

1), which are molecules present on the surface of all nucleated cells and bind to foreign antigens so that CD8+ T-

cells can recognize and eliminate the invading pathogens (Garcia et al. 1996). Nonetheless, M.o. in particular had

a variety of upregulated clusters of differentiation (CD) genes, such as *cd7*, c*d44b*, *cd209*, *cd81*, *cd247*, and *cd79a*

(Table 1). CD molecules encode surface proteins on white blood cells that are important in regulating the function

of immune cells (Tian et al. 2022). M.o. also had an increased biased expression for immunoglobulins (Ig) (Table

1), also known as antibodies that help identify and neutralize bacteria and viruses, they are divided into heavy and

light polypeptide chains (Jung et al. 2006). M.o. expressed IgD, two Ig heavy variable genes *ighv1-4* and *ighv4-3*

important in determining the antigen specificity of the Ig, two copies of Ig light 1 constant 3, and recombination

signal binding protein for immunoglobulin kappa J region-like (*rbpjl*) (Table 1). The latter plays a critical role in the

joining of the V, D, and J segments during the VDJ recombination process which is important in the generation of

a diverse set of antibodies (Webb et al. 2021).

	GENE	My	Fy	Mo	Fo
adaptive	*c2cd2l*		■		
	cd9b				■
	leng9				■
	mal2				■
	sema3bl				■
	mhc1uka				■
	mhc1uka.1				■
	tcta				■
	rag1	■			
	cd99l2	■			
	cd44b			■	
	CR847844			■	
	jak3			■	
	ighv1			■	
	ighv4			■	
	cd81a			■	
	igl1c3			■	
	blnk			■	
	bcap29			■	
	rbpjl			■	
	cd247l			■	
	swap70a			■	
	igl1c3.1			■	
	cd79a			■	
	nfat5b			■	
	ighd			■	
innate	*nfil3*		■		
	irak1		■		
	nfil3.1		■		
	il6st		■		
	irf6		■		
	tnfrsf9b				■
	ifrd1				■
	maea				■
	il17rd				■
	c8a				■
	traf2b	■			
	cxcr1.1	■			
	cxcr1.2	■			
	crlf3	■			
	tlr18			■	
	unc93b1			■	
	cxcl12a			■	
	ifih1			■	
	irf2bpl			■	
	mpeg1			■	
	cxcr3			■	
	irf8			■	
	il1fma			■	
	il17rc			■	
	tlr3			■	

	tnfrsf21			🟥	
	ccl25b			🟥	
	tnfrsf1a			🟥	

Table 1: The following table displays immune genes that were significantly differentially expressed in the head kidney for sex exclusively either in young or old pipefish. Boxes coloured in red indicate genes upregulated for either M.y., F.y., M.o. or F.o.

4. Discussion

Our main aim was to understand how male and female sex-role reversed pipefish differentiate in allocating their resources for reproduction or soma maintenance for age. We analysed gene expression in three different tissues of young, old, female and reproductively active (pregnant) and non-reproductively active males in the pipefish *Syngnathus typhle*. We identified distinct expression profiles in brain, head kidney and gonads, with most annotated pathways being unique to the organ type (Figure 1D). Pregnancy and age did not appear to strongly influence gene expression in any of the tissues assessed. Sex had the strongest effect on gonads and head kidney, and not at all on brain (Figure 1C). The analysis also revealed that age had an impact on the types of genes that were differentially expressed between males and females in the head kidney.

The gene ontology enrichment analysis comparing M.y., F.y., M.o. and F.o. revealed that females had many more upregulated transcripts relating to energy and lipid metabolism, whilst males had several upregulated pathways involved in the translation and genome maintenance process. Basic metabolic processes in males and females are similar in nature. Howbeit, different body composition (fat and muscle mass), sex hormones, sex-specific energy expenditure and age are all known to influence the effectiveness of how they metabolise nutrients and store energy (Krumsiek et al. 2015; Quirós Cognuck et al. 2020). Often a higher muscle mass correlates with a higher metabolic rate and in male mammals, sexual traits, such as body size or body colouration, are under stronger sexual selection (Halsey et al. 2022). Both F.y. and F.o. had enriched metabolic pathways involved in fat storage, such as fatty acid metabolism (REAC:R-DRE-8978868) and lipid metabolism (REAC:R-DRE-556833). Additionally, females had both *ppary* and *adiponectin* upregulated, the first a nuclear receptor important in the regulation of fatty acid metabolism which balances the storage and utilisation of fatty acids in the body (Rosen et al. 1999), found to be mostly higher expressed in the adipose tissue of females (Vidal-Puig et al. 1997), while the latter is involved in the regulation of glucose and was found to increase in older female mice (Gui, Silha, and Murphy 2004). These results match with research in conventional sex-role mammalian models (Figure 3), where females tend to have higher expression of genes involved in fat storage, consistent with the fact that they tend to store more body fat than males and are more efficient at using it as a source of energy (Shi, Seeley, and Clegg 2009). In mammals, increased lipid storage helps to ensure that enough energy is there to support herself and her offspring during development and provides a buffer against periods of food scarcity (Heldstab, van Schaik, and Isler 2017; Rödel et al. 2016). In the sex-role reversed broad-nosed pipefish this seems to remain a maternal trait and thus likely only

related to egg production, but not necessarily pregnancy. Furthermore, also sex-hormones play an important role

in metabolism and particularly oestrogen concentrations can alter how and where fat is stored in the body (Rubinow

2018). In *S. typhle* males produce testosterone and females produce oestrogen (Figure 3), despite the male

pregnancy and the sex-role reversal (Eens and Pinxten 2000; Mayer et al. 1993). Interestingly, a recent study in

medaka revealed that females turned into males upon a starvation period, the inhibition of the fatty acid synthesis

was suggested to have induced the sex reversal (Sakae et al. 2020). In *S. typhle* female would appear to rely

more on lipids as a source of energy for growth and egg production (Valdebenito, Gallegos, and Effer 2015), fat

deposits being possibly regulated by oestrogen. Male pipefish pregnancy is energetically costly, so they may be

utilizing other preferred sources of energy such as carbohydrates or proteins.

Male pipefish expressed genes from pathways involved in NMD, RNA, DNA metabolic process and ribosomes,

which could indicate changes in the rates and stability of translation and transcription processes in males compared

to females (Hug, Longman, and Cáceres 2015; López-Otín et al. 2013). This could be due to testosterones

promoting cell growth and more rapid division, which would possibly lead to a shorter lifespan (Ma et al. 2015; Park

et al. 2016). However, testosterone levels change with age and the levels usually drop during male pregnancy

(Eens and Pinxten 2000; Mayer et al. 1993; Scobell and Mackenzie 2011). Old male pipefish in particular had an

overrepresentation of pathways relating to DNA repair (Figure 2D). This may be due to exposure to environmental

damage or increased oxidative stress, with the cell trying to adapt and maintain genomic stability. From the

perspective of somatic maintenance, these results could be very beneficial, as an accumulation of DNA damage

over time is one of the major causes of ageing (López-Otín et al. 2013) and an efficient DNA repair system

mechanism is crucial in resisting oxidative stress and maintaining genomic integrity. This suggests a sex-role-

dependent selection for male pipefish to be investing more energy in protein and genome maintenance pathways

(Figure 3). Nonetheless, there were many more genes involved in autophagy that were upregulated in F.o. (Figure

2D). Autophagy is vital to recycling and/or degrading damaged proteins and organelles (López-Otín et al. 2013).

There is some research (primarily in rodents and humans) that indicate that there may be sex-specific differences

in the regulation of autophagy through sex hormones (Park et al. 2016). Testosterone has been shown to decrease

the expression of autophagy-related genes in men, while oestrogen increases them (Ma et al. 2015; Park et al.

2016). As previously mentioned, a sex-role reversal in *S. typhle* did not evolve with a reversal of sex hormones

(Mayer et al. 1993; Scobell and Mackenzie 2011). Thus, it would appear that sex, specifically sex hormones, may

be the determining factor for the regulation of autophagy. Moreover, it is important to note that the upregulation of

autophagy genes is beneficial to a certain degree, as it has been found to lead to cell death and cancer if

overexpressed (Finnegan et al. 2022).

Our differential gene expression analysis in head kidney also showed sex-specific immune gene expression, with

the differences between males and females becoming even stronger with age. Sex immune dimorphism is known

in diverse species from insects to lizards, birds and mammals, with an overall stronger innate and adaptive immune

response in females (Roved, Westerdahl, and Hasselquist, 2017; Klein and Flanagan 2016a). Theories suggest

that this could be a result of being either the heterogametic or homogametic sex, as the same sex chromosomes

help compensate for recessive mutations (Regan and Partridge 2013; Xirocostas, Everingham, and Moles 2020),

or sex-steroid hormones where, for example, oestrogens can act as antioxidants (Borrás et al. 2021), boost

telomere repair mechanism (Bayne et al. 2008), and testosterone can increase aggression providing an advantage

in the competition so by increasing the extrinsic hazard risk (Brooks and Garratt 2017) and additionally acting

immunosuppressive (Trigunaite, Dimo, and Jørgensen 2015). Generally, sex-dependent selection to optimise

energy investment in either soma maintenance or reproduction may have led to differences in resource allocation

in immune defence, metabolic requirements and ultimately mortality risk (Bronikowski et al. 2022; Kirkwood and

Rose 1991; Pennell, Galligan, and Fish 2012). From our results, we found a sex-specific differential expression of

immune genes, with males expressing more pro-inflammatory molecules, TLRs, CD genes, and immunoglobulins

(Table 1 and Table S5). These results are in contrast to studies in humans and other mammals (Figure 3), where

it has been shown that cell-mediated immunity seems to be more pronounced in females with both DC antigen-

presenting genes, TLRs and immunoglobulins proliferating at higher rates (Capone et al. 2018; Klein and Flanagan

2016). Furthermore, sex-hormone-specific research has shown that androgens typically induce an anti-

inflammatory response (Klein and Flanagan 2016; Taneja 2018), increasing also *il10* production, which conversely,

was significantly upregulated in female pipefish in our dataset, with also numerous pro-inflammatory markers being

upregulated in males (Table 1 and Table S5). Although female *S. typhle* did not have such a strong immune

variation with age, they did have MHC class 1 variants upregulated (Table 1). These molecules are a critical

component of the adaptive immune system and are responsible for presenting viral peptides, among others, to

CD8+ T cells for destruction (Garcia et al. 1996). This could mean that females are better equipped to mount a

cytotoxic T-cell response against infected cells, which could be the result of certain pathogens having a higher

prevalence in females compared to males. We know that sex-specific pathogens exist in other fish species due to

sex-specific behaviours, such as shoaling behaviour in female guppies' increased ectoparasite loads (Richards,

van Oosterhout, and Cable 2010). Preliminary in house data suggests that there is a difference in ectoparasite

prevalence between male and female pipefish and there is evidence to suggest that parent-specific

transgenerational immune priming does occur in broad-nosed pipefish (Beemelmanns and Roth 2016). Potential

future studies could investigate if parents are also priming their offspring to sex-specific pathogens.

The fact that males have overall more upregulated immune genes, particularly innate genes in older individuals,

might be the result of males requiring a more pronounced immune response to protect their health. In the sex-role

revered pipefish *S. typhle* males have a greater investment in the survival of their offspring and themselves, as

they are the sex that is reproductively constrained (Mobley et al. 2018). Also, males may have a higher exposure

and be more susceptible to microorganisms through females depositing their eggs in their pouch. Research has

shown that microbial diversity is highest in the pouch during pregnancy (Tanger et al. 2023) and embryos are highly

vulnerable to pathogenic attack (Wu et al. 2021). We also know from various animal systems that males and

females respond differently to microbes and other immune challenges and that numerous factors contribute to sex-

immune dimorphism from hormonal differences to behaviour-based pathogen exposure (Richards, van Oosterhout,

and Cable 2010; Rubinow 2018). Such results would indicate a sex pattern reversal of inflammatory response

compared with conventional sex-role species (Figure 3), thus making sex-role and the life-history trade-offs play a

more significant role here.

Figure 3: Here represented is a summary comparison of traits between a conventional sex-role mouse model and the broad-
nosed pipefish, which exhibits sex-role reversal. Specifically, we examine how these traits differ based on biological sex (pink for
females and blue for males). On the far right of the diagram, circular arrows indicate what life-history traits may be reversed
compared to the conventional model species.

In summary, sex-biased gene expression was strongest between old male and female *S. typhle* indicating the

importance of looking at sex traits in the context of senescence, specifically in species where fertility increasing

proportionally with age. Biological sex seems to be responsible for regulating fatty acid and lipid metabolism in

female pipefish, aligning with results from other species with conventional sex roles (Figure 3). Oppositely, our

results suggest that sex-roles may be leading male pipefish to allocate more energy to genome stability and repair,

and immune response as they are the sex that is time-constrained by the need to grow offspring in their pouch

(Figure 3). Similarly how for female mammals, having a higher parental time cost may result in them increasing

their reproductive time by upregulating their somatic maintenance and improving their health for the long haul.

Our study highlights the importance of considering the evolutionary context of senescence and the need to

investigate a variety of species with non-conventional sex behaviours to comprehensively understand the evolution

of sex lifespan dimorphism and sex-specific resource allocation. In future research, we aim to extend our studies

on sexual dimorphism beyond a single species of Syngnathids, given the diverse array of sex role changes present

within the family (Hamilton et al. 2017). By aggregating more datasets on sex-biased gene expression, we can gain

a better understanding of what determines sex and sex role in Syngnathids. Furthermore, future research that

involves comparing endocrine hormonal production and endocrinological manipulation could provide insights into

how sex is decoupled from sex roles, while also addressing the impact of hormones on gene expression and the

resulting phenotype.

5. Conclusion

Sex-specific lifespan dimorphism is found amongst many species and one prevalent hypothesis for this is that

males and females have distinct life-history strategies, which lead to unequal resource allocation between somatic

maintenance and reproduction, depending on how they acquire and optimise their fitness. In our study we analysed

gene expression of the sex-role reversed broad-nosed pipefish and found sex-specific differential gene expression

in head kidney varying with age. Specifically, females had many more upregulated transcripts relating to energy

and lipid metabolism, while males had many upregulated pathways involved in translation, genome maintenance

processes, and immune defence. Interestingly, while sex roles may influence some gene groups and reverse their

expression (e.g. genome maintenance and immune defence) when moving from conventional to sex-role reversal,

the activity of other gene groups may depend on biological sex and sex hormones (e.g., energy and fat storage).

This study underscores the importance of incorporating a wider range of species that exist along a spectrum of sex

roles, in order to enhance future research on sex-specific immune and lifespan-dimorphism, as well as the roles of

biological sex and sex soles in shaping them.

Ethics statement

Authors' contributions

F.A.P carried out the experiment, analysed the data, interpreted the results and wrote the article. G.T. helped

develop the DEGs limma:voom pipeline analysis. A.D. performed the orthology analysis and provided help with the

GO profiling. O.R. helped during the field work, supervised the project and supported the writing of the manuscript.

Acknowledgements

Special thanks to Dr. Ralf Schneider, who provided useful insights and feedback on the statistical analysis and

differential gene expression analysis.

Funding

This work was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) through

the Research Training Group for Translational Evolutionary Research (RTG 2501 TransEvo).

Conflict of interest declaration

The authors declare no conflicts of interest.

Data accessibility

Supplemental information include both additional figures (Figure S1-S3), the significant differentially expressed

genes in head kidney and the gene set enrichment analysis results as tables (Table S1-S5). The raw sequencing

data and metadata used in this study is available from the National Center for Biotechnology Information (NCBI)

Sequence Read Archive (SRA) under BioProject ID PRJNA943164 with submission ID SUB12940929. The whole

genome assembly of *Syngnathus typhle* can be found under BioProject ID PRJNA947442 with submission ID

SUB12974183.

References

Alvarez, Roberto Vera, Lorinc Sandor Pongor, Leonardo Mariño-Ramírez, and David Landsman. 2019.
"TPMCalculator: One-Step Software to Quantify MRNA Abundance of Genomic Features." *Bioinformatics*
35(11): 1960–62.

Andrews, S. 2010. " FastQC: A Quality Control Tool for High Throughput Sequence Data."
http://www.bioinformatics.babraham.ac.uk/projects/fastqc.

de Araujo, Mariana E G et al. 2017. 358 Science *Crystal Structure of the Human Lysosomal MTORC1 Scaffold*
Complex and Its Impact on Signaling.

Baker, Stacey J, and E Premkumar Reddy. *Modulation of Life and Death by the TNF Receptor Superfamily*.
http://www.stockton-press.co.uk/onc.

Barneche, Diego R, Craig R White, and Dustin J Marshall. 2018. "Fish Reproductive-Energy Output Increases
Disproportionately with Body Size." *Science* 645(May): 642–45.

Bateman, A. J. 1948. "Intra-Sexual Selection in Drosophila." *Heredity*.

Bayne, Sharyn et al. 2008. "Estrogen Deficiency Leads to Telomerase Inhibition, Telomere Shortening and
Reduced Cell Proliferation in the Adrenal Gland of Mice." *Cell Research*.

Beemelmanns, Anne, and Olivia Roth. 2016. "Bacteria-Type-Specific Biparental Immune Priming in the Pipefish
Syngnathus Typhle." *Ecology and Evolution* 6(18): 6735–57.

Benvenuto, C. et al. 2017. "Ecological and Evolutionary Consequences of Alternative Sex-Change Pathways in
Fish." *Scientific Reports* 7(1): 1–12.

Bjørgen, Håvard, and Erling Olaf Koppang. 2021. "Anatomy of Teleost Fish Immune Structures and Organs."
Immunogenetics 73(1): 53–63.

Borrás, C. et al. 2021. "Estrogen Replacement Therapy Induces Antioxidant and Longevity-Related Genes in
Women after Medically Induced Menopause." *Oxidative Medicine and Cellular Longevity* 2021.

Bronikowski, Anne M. et al. 2022. "Sex-Specific Aging in Animals: Perspective and Future Directions." *Aging Cell*
21(2): 1–25.

Brooks, Robert C., and Michael G. Garratt. 2017. "Life History Evolution, Reproduction, and the Origins of Sex-
Dependent Aging and Longevity." *Annals of the New York Academy of Sciences* 1389(1): 92–107.

Capone, Imerio et al. 2018. "Sexual Dimorphism Of Immune Responses: A New Perspective in Cancer
Immunotherapy." *Frontiers in Immunology* 9(MAR).

Catena, Valeria, and Maurizio Fanciulli. 2017. "Deptor: Not Only a MTOR Inhibitor." *Journal of Experimental and*
Clinical Cancer Research 36(1).

Chaplin, David D. 2010. "Overview of the Immune Response." *Journal of Allergy and Clinical Immunology* 125(2
SUPPL. 2).

Chen, Shifu, Yanqing Zhou, Yaru Chen, and Jia Gu. 2018. "Fastp: An Ultra-Fast All-in-One FASTQ
Preprocessor." In *Bioinformatics*, Oxford University Press, i884–90.

Clutton-Brock, T. H., and K. Isvaran. 2007. "Sex Differences in Ageing in Natural Populations of Vertebrates."
Proceedings of the Royal Society B: Biological Sciences.

Dobin, Alexander et al. 2013. "STAR: Ultrafast Universal RNA-Seq Aligner." *Bioinformatics* 29(1): 15–21.

Eens, Marcel, and Rianne Pinxten. 2000. 51 Behavioural Processes *Sex-Role Reversal in Vertebrates:*
Behavioural and Endocrinological Accounts. www.elsevier.com/locate/behavproc.

Efeyan, Alejo, William C. Comb, and David M. Sabatini. 2015. "Nutrient-Sensing Mechanisms and Pathways."
Nature 517(7534): 302–10.

Emms, David M., and Steven Kelly. 2019. "OrthoFinder: Phylogenetic Orthology Inference for Comparative
Genomics." *Genome Biology* 20(1): 1–14.

Finnegan, Ryan M., Ahmed M. Elshazly, Patricia v. Schoenlein, and David A. Gewirtz. 2022. "Therapeutic
Potential for Targeting Autophagy in ER+ Breast Cancer." *Cancers* 14(17).

Froehlich, Jacob M. et al. 2013. "Sarcopenia and Piscines: The Case for Indeterminate-Growing Fish as Unique
Genetic Model Organisms in Aging and Longevity Research." *Frontiers in Genetics* 4(AUG): 1–8.

Garcia, K. Christopher et al. 1996. "CD8 Enhances Formation of Stable T-Cell Receptor/MHC Class I Molecule
Complexes." *Nature* 384(6609): 577–81.

Geven, Edwin J.W., and Peter H.M. Klaren. 2017. "The Teleost Head Kidney: Integrating Thyroid and Immune
Signalling." *Developmental and Comparative Immunology* 66: 73–83.

Grayfer, Leon, and Miodrag Belosevic. 2012. "Cytokine Regulation of Teleost Inflammatory Responses." In *New*
Advances and Contributions to Fish Biology,.

Gubbels Bupp, Melanie R. 2015. "Sex, the Aging Immune System, and Chronic Disease." *Cellular Immunology*.

Gui, Yaoting, Josef v. Silha, and Liam J. Murphy. 2004. "Sexual Dimorphism and Regulation of Resistin,
Adiponectin, and Leptin Expression in the Mouse." *Obesity Research* 12(9): 1481–91.

Hale, Matthew C., Garrett J. McKinney, Frank P. Thrower, and Krista M. Nichols. 2018. "Evidence of Sex-Bias in
Gene Expression in the Brain Transcriptome of Two Populations of Rainbow Trout (Oncorhynchus Mykiss)
with Divergent Life Histories." *PLoS ONE* 13(2): 1–18.

Halsey, Lewis G. et al. 2022. "Variability in Energy Expenditure Is Much Greater in Males than Females." *Journal*
of Human Evolution 171.

Hamilton, Healy et al. 2017. "Molecular Phylogeny and Patterns of Diversification in Syngnathid Fishes."
Molecular Phylogenetics and Evolution 107: 388–403.

Heldstab, Sandra A., Carel P. van Schaik, and Karin Isler. 2017. "Getting Fat or Getting Help? How Female
Mammals Cope with Energetic Constraints on Reproduction." *Frontiers in Zoology* 14(1).

Henning, Wilhelm, and Horst-Werner Stürzbecher. 2003. "Homologous Recombination and Cell Cycle
Checkpoints: Rad51 in Tumour Progression and Therapy Resistance." *Toxicology* 193(1–2): 91–109.

Holmdahl, Rikard et al. 2016. "Ncf1 Polymorphism Reveals Oxidative Regulation of Autoimmune Chronic
Inflammation." *Immunological Reviews* 269(1): 228–47.

Holtze, Susanne et al. 2021. "Alternative Animal Models of Aging Research." *Frontiers in Molecular Biosciences* 8(May).

Hug, Nele, Dasa Longman, and Javier F. Cáceres. 2015. "Mechanism and Regulation of the Nonsense-Mediated Decay Pathway." *Nucleic Acids Research* 44(4): 1483–95.

Jalan, Manisha, Kyrie S. Olsen, and Simon N. Powell. 2019. "Emerging Roles of RAD52 in Genome Maintenance." *Cancers* 11(7).

Janicke, Tim, Ines K. Häderer, Marc J. Lajeunesse, and Nils Anthes. 2016. "Evolutionary Biology: Darwinian Sex Roles Confirmed across the Animal Kingdom." *Science Advances* 2(2).

Jones, Owen R. et al. 2014. "Diversity of Ageing across the Tree of Life." *Nature*.

Jung, David, Cosmas Giallourakis, Raul Mostoslavsky, and Frederick W. Alt. 2006. "Mechanism and Control of V(D)J Recombination at the Immunoglobulin Heavy Chain Locus." *Annual Review of Immunology* 24: 541–70.

Kawasaki, Takumi, and Taro Kawai. 2014. "Toll-like Receptor Signaling Pathways." *Frontiers in Immunology* 5(SEP).

Kirkwood, T. B.L., and M. R. Rose. 1991. "Evolution of Senescence: Late Survival Sacrificed for Reproduction." *Philos Trans R Soc Lond B Biol Sci.* Apr 29(332(1262)): 15–24.

Klein, Sabra L., and Katie L. Flanagan. 2016. "Sex Differences in Immune Responses." *Nature Reviews Immunology* 16(10): 626–38.

Krumsiek, Jan et al. 2015. "Gender-Specific Pathway Differences in the Human Serum Metabolome." *Metabolomics* 11(6): 1815–33.

Lemaître, Jean François et al. 2020. "Sex Differences in Adult Lifespan and Aging Rates of Mortality across Wild Mammals." *Proceedings of the National Academy of Sciences of the United States of America*.

Liu, Shuang Qing et al. 2021. "TNF-α-Induced Protein 8-like 2 Negatively Regulates the Immune Function of Dendritic Cells by Suppressing Autophagy via the TAK1/JNK Pathway in Septic Mice." *Cell Death and Disease* 12(11).

Li, Yajuan et al. 2017. "Pattern Recognition Receptors in Zebrafish Provide Functional and Evolutionary Insight into Innate Immune Signaling Pathways." *Cellular and Molecular Immunology* 14(1): 80–89.

López-Otín, Carlos et al. 2013. "The Hallmarks of Aging." *Cell*.

Maréchal, Alexandre, and Lee Zou. 2013. "DNA Damage Sensing by the ATM and ATR Kinases." *Cold Spring Harbor Perspectives in Biology* 5(9).

Mayer, Ian et al. 1993. "Plasma Levels of Sex Steroids in Three Species of Pipefish (Syngnathidae)." *Canadian Journal of Zoology*.

Ma, Yi et al. 2015. "Testosterone Regulates the Autophagic Clearance of Androgen Binding Protein in Rat Sertoli Cells." *Scientific Reports* 5.

Mayne, Benjamin T. et al. 2016. "Large Scale Gene Expression Meta-Analysis Reveals Tissue-Specific, Sex-Biased Gene Expression in Humans." *Frontiers in Genetics* 7(OCT): 1–14.

Mitchell, Sarah J., Morten Scheibye-Knudsen, Dan L. Longo, and Rafael de Cabo. 2015. "Animal Models of Aging Research: Implications for Human Aging and Age-Related Diseases." *Annual Review of Animal Biosciences*.

Mobley, Kenyon B. et al. 2018. "Female Ornamentation and the Fecundity Trade-off in a Sex-Role Reversed Pipefish." *Ecology and Evolution* 8(18): 9516–25.

Mustofa, Md Kawsar et al. 2020. "Roles of Chk2/CHEK2 in Guarding against Environmentally Induced DNA Damage and Replication-Stress." *Environmental and Molecular Mutagenesis* 61(7): 730–35.

Park, Jaekyoung, Hyejin Shin, Haengseok Song, and Hyunjung J. Lim. 2016. "Autophagic Regulation in Steroid Hormone-Responsive Systems." *Steroids* 115: 177–81.

Pennell, Leesa M., Carole L. Galligan, and Eleanor N. Fish. 2012. "Sex Affects Immunity." *Journal of Autoimmunity*.

Quirós Cognuck, Susana et al. 2020. "Sex Differences in Body Composition, Metabolism-Related Hormones, and Energy Homeostasis during Aging in Wistar Rats." *Physiological Reports* 8(20).

Raudvere, Uku et al. 2019. "G:Profiler: A Web Server for Functional Enrichment Analysis and Conversions of
Gene Lists (2019 Update)." *Nucleic Acids Research* 47(W1): W191–98.

R Core Team. 2022. "R: A Language and Environment for Statistical Computing."

Regan, Jennifer C., and Linda Partridge. 2013. "Gender and Longevity: Why Do Men Die Earlier than Women?
Comparative and Experimental Evidence." *Best Practice and Research: Clinical Endocrinology and
Metabolism* 27(4): 467–79.

Reznick, David N., Cameron Ghalambor, and Leonard Nunney. 2002. "The Evolution of Senescence in Fish."
Mechanisms of Ageing and Development 123: 773–89.

Richards, E. Loys, Cock van Oosterhout, and Joanne Cable. 2010. "Sex-Specific Differences in Shoaling Affect
Parasite Transmission in Guppies." *PLoS ONE* 5(10).

Ripley, Jennifer L., and Christy M. Foran. 2009. "Direct Evidence for Embryonic Uptake of Paternally-Derived
Nutrients in Two Pipefishes (Syngnathidae: Syngnathus Spp.)." *Journal of Comparative Physiology B:
Biochemical, Systemic, and Environmental Physiology*.

Ritchie, Matthew E. et al. 2015. "Limma Powers Differential Expression Analyses for RNA-Sequencing and
Microarray Studies." *Nucleic Acids Research* 43(7): e47.

Robinson, Mark D., Davis J. McCarthy, and Gordon K. Smyth. 2009. "EdgeR: A Bioconductor Package for
Differential Expression Analysis of Digital Gene Expression Data." *Bioinformatics* 26(1): 139–40.

Rödel, Heiko G., Teresa G. Valencak, Alexandra Handrek, and Raquel Monclús. 2016. "Paying the Energetic
Costs of Reproduction: Reliance on Postpartum Foraging and Stored Reserves." *Behavioral Ecology* 27(3):
748–56.

Rolff, Jens. 2002. "Bateman's Principle and Immunity." *Proceedings of the Royal Society of London. Series B:
Biological Sciences* 269(1493): 867–72.

Rosen, Evan D et al. 1999. 4 Molecular Cell *PPAR Functions*.

Roth, O., J. P. Scharsack, I. Keller, and T. B.H. Reusch. 2011. "Bateman's Principle and Immunity in a Sex-Role
Reversed Pipefish." *Journal of Evolutionary Biology*.

Rubinow, Katya B. 2018. "An Intracrine View of Sex Steroids, Immunity, and Metabolic Regulation." *Molecular
Metabolism* 15: 92–103.

Sagebakken, Gry, Ingrid Ahnesjö, and Charlotta Kvarnemo. 2016. "Costs and Benefits to Pregnant Male Pipefish
Caring for Broods of Different Sizes." *PLoS ONE*.

Sakae, Yuta et al. 2020. "Starvation Causes Female-to-Male Sex Reversal through Lipid Metabolism in the
Teleost Fish, Medaka (Olyzias Latipes)." *Biology Open* 9(4).

Schneider, R. F., Woltering, J. M., Adriaens, D., and Roth, O. (2022). A comparative analysis of the ontogeny of
syngnathids (pipefishes and seahorses) reveals how heterochrony contributed to their diversification.
Developmental Dynamics. https://doi.org/10.1002/DVDY.551

Scobell, S. K., and D. S. Mackenzie. 2011. "Reproductive Endocrinology of Syngnathidae." *Journal of Fish
Biology*.

Shaw, Laurence M.A., and Sebastian L.J. Shaw. 2009. "Menopause, Evolution and Changing Cultures."
Menopause International.

Shi, Haifei, Randy J. Seeley, and Deborah J. Clegg. 2009. "Sexual Differences in the Control of Energy
Homeostasis." *Frontiers in Neuroendocrinology* 30(3): 396–404.

Smith, Carl, and Robert J. Wootton. 1995. "The Costs of Parental Care in Teleost Fishes." *Reviews in Fish
Biology and Fisheries* 5(1): 7–22.

Vom Steeg, Landon G., and Sabra L. Klein. 2016. "SeXX Matters in Infectious Disease Pathogenesis." *PLoS
Pathogens*.

Stenvinkel, Peter, and Paul G. Shiels. 2019. "Long-Lived Animals with Negligible Senescence: Clues for Ageing
Research." *Biochemical Society Transactions* 47(4): 1157–64.

Stiller, Josefin et al. 2022. "Phylogenomic Analysis of Syngnathidae Reveals Novel Relationships, Origins of
Endemic Diversity and Variable Diversification Rates." *BMC Biology* 20(1): 1–21.

Taneja, Veena. 2018. "Sex Hormones Determine Immune Response." *Frontiers in Immunology* 9.

Tanger, Isabel S., Julia Stefanschitz, Olivia Roth, and Yannick Schwert. 2023. "The Source of Microbial
Transmission Influences Niche Colonization and Microbiome Development." *BioRxiv*.
https://doi.org/10.1101/2023.01.31.526426 (February 15, 2023).

Tian, Hong fei et al. 2022. "Cluster of Differentiation Antigens: Essential Roles in the Identification of Teleost Fish
T Lymphocytes." *Marine Life Science and Technology*.

Trifonova, Oxana P. et al. 2018. "Comparative Analysis of the Blood Plasma Metabolome of Negligible, Gradual
and Rapidly Ageing Fishes." *Fishes* 3(4): 46.

Trigunaite, Abhishek, Joana Dimo, and Trine N. Jørgensen. 2015. "Suppressive Effects of Androgens on the
Immune System." *Cellular Immunology* 294(2): 87–94. http://dx.doi.org/10.1016/j.cellimm.2015.02.004.

Valdebenito, Iván I., Patricia C. Gallegos, and Brian R. Effer. 2015. "Gamete Quality in Fish: Evaluation
Parameters and Determining Factors." *Zygote* 23(2): 177–97.

Vidal-Puig, Antonio J. et al. 1997. "Peroxisome Proliferator-Activated Receptor Gene Expression in Human
Tissues: Effects of Obesity, Weight Loss, and Regulation by Insulin and Glucocorticoids." *Journal of
Clinical Investigation* 99(10): 2416–22.

Viña, Jose, Consuelo Borrás, and Jaime Miquel. 2007. "Theories of Ageing." *IUBMB Life* 59(4–5): 249–54.

Wang, Yan et al. 2019. "SIRT2: Controversy and Multiple Roles in Disease and Physiology." *Ageing Research
Reviews* 55: 100961.

Webb, Louise M.C. et al. 2021. "Ageing Promotes Early T Follicular Helper Cell Differentiation by Modulating
Expression of RBPJ." *Aging Cell* 20(1).

Whittington, Camilla M, and Christopher R Friesen. 2020. "The Evolution and Physiology of Male Pregnancy in
Syngnathid Fishes." *Biological Reviews* 5.

Whittle, Carrie A., and Cassandra G. Extavour. 2019. "Selection Shapes Turnover and Magnitude of Sex-Biased
Expression in Drosophila Gonads." *BMC Evolutionary Biology* 19(1): 1–20.

Williams, George C. 1957. "Pleiotropy, Natural Selection, and the Evolution of Senescence." *Evolution*.

Wilson, A. B., A. Vincent, I. Ahnesjö, and Axel Meyer. 2001. "Male Pregnancy in Seahorses and Pipefishes
(Family Syngnathidae): Rapid Diversification of Paternal Brood Pouch Morphology Inferred from a
Molecular Phylogeny." *Journal of Heredity* 92(2): 159–66.

Wolf, J. B., and M. J. Wade. 2001. "On the Assignment of Fitness to Parents and Offspring: Whose Fitness Is It
and When Does It Matter?" *Journal of Evolutionary Biology*.

Wu, Yongli et al. 2021. "Molecular Characterization of TLR22 and Its Role in Immunological Modification of the
Brood Pouch of the Lined Seahorse, Hippocampus Erectus." *Aquaculture* 539: 736628.

Xirocostas, Zoe A., Susan E. Everingham, and Angela T. Moles. 2020. "The Sex with the Reduced Sex
Chromosome Dies Earlier: A Comparison across the Tree of Life." *Biology Letters*.

Yuan, Shaochun et al. 2014. "Comparative Immune Systems in Animals." *Annual Review of Animal Biosciences*
2: 235–58.

Zou, Jun, and Christopher J. Secombes. 2016. "The Function of Fish Cytokines." *Biology* 5(2).

Zuk, Marlene. 2009. "The Sicker Sex." *PLoS Pathogens*.