Direct translation of incoming retroviral genomes

Julia Köppke[1], Luise-Elektra Keller[1,2], Michelle Stuck[1,3], Nicolas D. Arnow[1], Norbert Bannert[1], Joerg Doellinger[4], and Oya Cingöz[1,*]

[1] Robert Koch Institute, Department of Infectious Diseases, Unit of Sexually Transmitted Bacterial Pathogens and HIV (FG18), Berlin, Germany

[2] Current address: Institute of Cardiovascular Regeneration, Goethe University Frankfurt, Germany

[3] Current address: Department of Chemistry, Heidelberg University, Heidelberg, Germany

[4] Robert Koch Institute, Center for Biological Threats and Special Pathogens, Proteomics and Spectroscopy (ZBS6), Berlin, Germany

* To whom correspondence should be addressed. Email: cingoezo@rki.de

Abstract

Viruses that carry a positive-sense, single-stranded (+ssRNA) RNA translate their genomes soon after entering the host cell to produce viral proteins, with the exception of retroviruses. A distinguishing feature of retroviruses is reverse transcription, where the +ssRNA genome serves as a template to synthesize a double-stranded DNA copy that subsequently integrates into the host genome. As retroviral RNAs are produced by the host cell transcriptional machinery and are largely indistinguishable from cellular mRNAs, we investigated the potential of incoming retroviral genomes to directly express proteins. Here we show through multiple, complementary methods that retroviral genomes are translated after entry. Our findings challenge the notion that retroviruses require reverse transcription to produce viral proteins. Synthesis of retroviral proteins in the absence of productive infection has significant implications for basic retrovirology, immune responses and gene therapy applications.

Introduction

All viruses, regardless of their nucleic acid type, composition or orientation, need to reach the mRNA stage for successful infection. Retroviruses carry two copies of a positive-sense, single-stranded RNA (+ssRNA) genome; however, they form a separate class from +ssRNA viruses in the Baltimore classification because their replication strategy involves reverse transcription of the ssRNA genome into a double-stranded (dsDNA) copy. Based on current knowledge most if not all +ssRNA viruses directly translate their RNA to synthesize viral proteins upon entry into host cells, with the exception of retroviruses, which undergo reverse transcription and degrade the original genomic RNA in the process. As retroviral genomes are produced by the host cell machinery, with a 5' cap and a 3' poly-A tail, we asked: Are incoming retroviral genomic RNAs also directly translated?

One small but notable difference between the full-length retroviral RNA packaged into the virions as the genome and the one that produces Gag and GagPol polyproteins is their transcription start sites (TSS). Full-length retroviral RNAs contain heterogenous TSS, which alters the RNA structure and impacts its dimerization and translation potential.[1, 2, 3] While 1G transcripts are primarily dimerized and selectively packaged into virions, 2G and 3G transcripts exist mainly as monomers and are enriched in polysomes.[1, 2] In addition, incompletely-spliced retroviral transcripts carry hypermethylated caps bound by NCBP3 instead of eIF4E, and translated in an mTOR-independent manner, although such hypermethylated caps were not detected in virion-packaged RNA.[4] As genomic RNA isolated from virions can be translated in *in vitro* systems to produce retroviral proteins,[5, 6, 7] the virion-packaged genomic RNA does not appear to be inherently untranslatable, even if it differs from the newly-produced full-length RNA that is translated to produce viral structural proteins.

Using multiple complementary approaches, including post-translational regulation of protein stability coupled to sensitive reporter assays, immunoprecipitation, polysome fractionation and SILAC-based mass spectrometry, we demonstrate here that incoming retroviral RNA genomes are translated shortly after cellular entry. This is a general process that occurs under a variety of conditions; with different viral genome lengths, cellular entry pathways and cell types. Extensive controls including nuclease treatments, checking for DNA or protein transfer, using fusion-defective viruses, and omission of various viral components (e.g. *env*, *gag*, packaging signal) confirm that the signal is truly due to direct *de novo* translation from incoming genomes. Capsid mutants that display altered stability and uncoating kinetics impact the translation of incoming RNAs by regulating the access of the encapsidated RNA to the translational machinery. The synthesis of retroviral proteins in the absence of reverse transcription has significant implications for basic retrovirology, immune responses during infection, and the use of retroviral vectors as RNA delivery vehicles.

Results

Post-translational control of protein stability minimizes virion-packaging and cellular delivery of reporter proteins

In the laboratory, retroviruses are typically produced by transfection of producer cells with plasmids encoding viral components. In case of viral genomes that carry a reporter gene, the reporter protein is also expressed in producer cells, which can get packaged into virions and delivered into recipient cells, yielding false positives.[8, 9, 10, 11, 12, 13] To minimize such producer cell background, which could mask the signal from *de novo* translated incoming retroviral RNAs, we employed a post-translational protein control system (ProteoTuner), where a destabilizing domain (DD) derived from a cellular gene with a very short half-life (FKBP12) is fused to the gene of interest, targeting it for rapid proteasomal degradation.[14] The unstable protein can be stabilized in a dose-dependent and reversible manner by adding a cell-permeable small molecule ligand called Shield1 that binds to the DD, allowing post-translational regulation of protein levels. We reasoned that producing viruses in the absence of the ligand would minimize reporter protein packaging into virions, whereas performing infections in the presence of the ligand would allow us to detect reporter expression from the incoming retroviral RNA genomes by stabilizing the reporter. As only two genomic RNAs per retrovirus particle are delivered into the cell upon entry, we selected the sensitive reporter nano-luciferase (Nluc) to assay the translation of incoming retroviral RNA genomes, which has superior sensitivity compared to other luciferase proteins (reviewed in [15]).

Nluc was cloned with or without the DD either under a CMV promoter in a minimal lentiviral vector to produce pLVX-(DD)-Nluc, or with deletions in the CMV promoter, IRES element and Neo resistance gene to generate pLVX-(DD)-Nluc-ΔCIN (Fig. 1A). Transfection of these constructs into 293T cells resulted in 30-45-fold less luciferase activity in case of DD-harboring constructs, while Shield1 treatment rescued the expression (Fig. 1B). The stabilization of DD-Nluc constructs by Shield1 was dose-dependent, whereas Nluc constructs without the responsive domain remained unaffected (Fig. 1C). To quantify the amount of reporter protein packaging into virions, we produced virions by transfecting cells with the lentiviral transfer vectors shown in Fig. 1A along with a packaging construct with or without an *env* plasmid (VSV-G) and assayed the virus particles themselves for luciferase activity. Virions produced by DD-Nluc vectors consistently yielded 120-270-fold lower signal compared to Nluc vectors for both constructs, while the presence or absence of an envelope protein (VSV-G) did not make a difference (Fig. 1D).

For HIV-1, the timing of the early events during infection are well-documented and cell type-

dependent. Reverse transcription can take anywhere from 6 - 48 hours, and integration follows about

5 hours after the completion of reverse transcription.[16, 17] As the constructs in Fig. 1A cannot produce

the reporter protein from full-length (unspliced) viral RNA, they require reverse transcription,

integration, transcription, splicing and translation to express luciferase. Accordingly, within the first

12 hours of infection, no change in signal was observed, indicating that the luciferase signal

represents Nluc protein that is transferred to and/or that remains associated with cells (Fig. 1E). The

presence or absence of an Env protein or the RT inhibitor nevirapine (NVP) had no effect on the

signal observed, although infection with the destabilized reporter viruses consistently yielded less

signal. Likewise, at 20 hours post-infection, there is still no expression from the provirus, but

expression did occur at 72 hours post-infection (hpi), which was reduced to background levels in the

absence of a functional Env or reverse transcription (Fig. 1F). These data show that post-translational

regulation of protein stability can markedly reduce passive protein packaging into virions and their

delivery into recipient cells, minimizing the background for assessment of incoming retroviral RNA

translation.

Incoming retroviral genomes are used as an mRNA for protein expression early after entry

To assess the direct translation potential of the full-length, incoming retroviral genomic RNA,

we cloned Nluc with or without the DD downstream of the packaging signal (Ψ; psi) in place of where

gag would normally be in a minimal lentiviral vector. The start codon of *gag* was mutated and any

additional elements that could affect translation efficiency (e.g. IRES, WPRE) were removed (Fig. 2A).

In line with previous results, transfection of DD-constructs resulted in ~180-fold decreased signal

compared to Nluc, which was completely rescued by Shield1 (S1) addition (Fig. 2B). We produced

virions and infected cells with serial dilutions of LV-DD-Nluc virus in the presence of NVP and S1, with

or without the translation inhibitor cycloheximide (CHX) to distinguish virion-packaged protein

delivery from *de novo* translation. Using viral doses as low as 0.0061 ng p24 per 100K cells, we

detected a CHX-sensitive signal (Fig. 2C). The signal to noise ratio (i.e. new protein synthesis vs.

packaged protein delivery) was highest between 0.165-4.5 ng p24 per 100K cells. Based on these

results, we opted to use 1-4 ng p24 in subsequent experiments.

Infection of cells with LV-Nluc virus resulted in an increase in the luciferase signal over time

despite the background of virion-packaged protein, which was diminished to background levels by

CHX treatment (Fig. 2D). RT inhibition had no effect on the luciferase signal within the first 8 hours,

indicating that the translation occurs independently of reverse transcription. We then performed an

infection with LV-DD-Nluc virus in the presence of NVP. The signal increased within the first 2 h, and

under S1-stabilized conditions increased further up to 6 h, whereas in the absence of S1, it decreased

after the initial hike due to the instability of the protein (Fig. 2E; red vs. blue lines). Translation

inhibition by CHX drastically reduced the signal in both cases, although increased stabilization of the

protein delivered passively by virions was also evident (Fig. 2E; green vs violet lines). Importantly,

when DD-Nluc is stabilized, there is a clear increase in reporter expression over time, which is

markedly reduced upon translation inhibition (Fig. 2E; red vs. violet lines). This difference is the result

of newly synthesized reporter proteins from the incoming retroviral genomic RNA in the absence of

reverse transcription. Similar results were observed regardless of whether the initial inoculum was

kept on the cells or whether the virus was removed after one hour, although the lack in continuous

virus uptake in the latter case was evident (Fig. 2F-G).

Expression from incoming retroviral genomes is due to *de novo* translation and occurs under

different conditions

Viral supernatants typically contain plasmids carried over from transfected producer cells.

Pre-treatment of our virus stocks with nucleases that degrade either DNA, RNA or both did not alter

reporter expression from the incoming viral genome after transduction, despite the enzymes being

functional, indicating that the observed signal is not due to nonspecific uptake of ambient DNA or

RNA (Fig. 3A-B). The presence of a functional viral envelope on particles was necessary, as viruses

pseudotyped with a VSV-G mutant defective in fusion activity (P127D) or those without an Env failed

to yield a signal above background (Fig. 3C). Similar results were observed with Gag-less "virus"

supernatants, when the packaging plasmid was omitted during virus production (Fig. 3D). To ensure

that our observations are not due to incomplete inhibition of RT, we confirmed these results with

viruses containing a catalytic mutant RT (Fig. 3E and Supp. Fig. 1A). To rule out endogenous reverse

transcripts already present in virions or the inadvertent packaging and delivery of plasmid DNA

fragments from producer cells as the source of the reporter signal, neither of which would be

susceptible to nuclease digestion, we transduced cells in the presence of transcription or translation

inhibitors (Actinomycin D [ActD] or CHX, respectively). Translation inhibition markedly reduced the

signal as shown previously (Fig. 2C-G), whereas transcription inhibition did not, indicating that the

observed signal is not due to DNA transfer, which would have required both of these processes (Fig.

3F). At the same concentrations both drugs inhibited expression from a transfected reporter plasmid,

validating their functionality (Fig. 3G). Although inhibition of reverse transcription is not required for

this process, we also confirmed these results during transduction in the presence of different RT

inhibitors, namely NVP, efavirenz (EFV) and tenofovir (TAF) used at 10 µM (Fig. 3H), which resulted in

similar expression levels as the untreated sample. When these RT inhibitors were used at higher

concentrations (100 µM) during infection, toxicity to cells was evident particularly in case of EFV,

where cells also lost the ability to express a transfected plasmid (Supp. Fig. 1B-C). Notably, the

translation of incoming retroviral genomes was observed with viral envelopes that use different entry

pathways (Fig. 3I) and in diverse cell lines and primary cells (Fig. 3J), indicating that this is a broad

process not restricted to a specific cell type or condition.

Immunoprecipitation, SILAC/MS and polysome fractionation confirm direct expression from

incoming HIV-1 genomes

The RNA genome of HIV-1 has an intricate secondary structure and several *cis*-acting RNA

elements with diverse functions in the replication cycle, which are not all present in the context of a

minimal vector.[18] To explore the contribution of such RNA elements to the process of incoming

retroviral genomic translation, we cloned DD-Nluc downstream of the packaging signal and a

mutated *gag* start codon in a near-full length HIV-1 construct based on NL4.3 that encodes firefly

luciferase in place of *nef* (Fig. 4A). Infection with NL43-DD-Nluc, produced by co-transfecting cells

with a packaging plasmid and a VSV-G *env*, yielded similar results to infection with minimal lentiviral

vectors in the presence of NVP (Fig. 4B vs. Fig 2E & 2G). Under S1-stabilized conditions the signal

increased over time, peaking at 4-6 hpi, whereas translation inhibition reduced the reporter activity

to baseline levels, confirming *de novo* protein synthesis (Fig. 4B; red vs. violet lines). Based on the

relatively short incubation times, we did not remove the virus inoculum, which explains why the

stabilized CHX condition shows a slight increase over time due to continuous reentry (violet line).

To evaluate the number of RNA copies delivered into cells, we performed transductions with

the parental NL43-Firefly virus in the presence or absence of NVP and quantified post-entry events at

24 and 48 hours. As infection with 4 ng p24 per 100K cells did not yield a detectable signal for RT

products above background, we performed infections using 10 times more virus than normally used,

which corresponds to a multiplicity of infection (MOI) of 1 based on the transducing units validated

by p24 staining. Under these conditions, we detected early RT products between 4.1 - 8.2 copies, late

RT products between 2.4 - 4.1 copies, and 2-LTR circles between 0.02 - 0.03 copies per haploid

genome (Supp. Fig. 2A).

As an alternative to reporter assays, we checked Gag protein production in the absence of

reverse transcription. Cells were challenged with VSV-G-pseudotyped NL43-Firefly virus (MOI = 1) in

the presence or absence of NVP and CHX. Gag production was assayed by immunoprecipitation (IP)

with an anti-Gag polyclonal antibody and western blot one day after infection to maximize the

accumulation of newly-synthesized viral proteins. In the absence of RT inhibition, Pr55-Gag was

produced at this time point, as expected. Although NVP treatment decreased this signal, viral protein

production was still evident and did not originate from incoming CA protein, as CHX treatment
abolished this signal (Fig. 4C).

To provide further evidence that authentic viral proteins are produced from the packaged
viral genome, we performed metabolic labeling via stable isotope labeling with amino acids in cell
culture followed by mass spectrometry (SILAC-MS). Briefly, cells grown in light medium were
transduced with VSV-G-pseudotyped NL43-Firefly virus also produced in light medium (MOI = 0.5). At
the time of transduction, cells were switched to heavy medium such that all newly produced proteins
would be heavy, whereas all pre-existing proteins light. After 18 hours, lysates were subjected to
immunoprecipitation with an anti-p24 antibody followed by mass spectrometry. Under conditions
where reverse transcription could take place, *de novo* viral Gag production was evident, as cells had
enough time to undergo the regular replication steps (Fig. 4D). Importantly, although the ratio of
heavy to light peptides that map to Gag were lower in case of RT inhibition, such peptides were still
detected, validating that incoming HIV-1 genomes are translated to produce viral proteins. Heavy
peptides corresponding to Gag were only detected under conditions where translation could take
place, confirming that these results truly are the result of *de novo* translation (Fig. 4D).

To validate the association of incoming HIV-1 RNA with polysomes - an indicator of active
translation - we also performed polysome fractionation on lysates from 293T cells after four hours of
infection with VSV-G-pseudotyped IIIB with a catalytic RT mutation (DD185/186AA; Supp. Fig. 1A).
Infected and uninfected cells yielded comparable polysome profiles with clearly separated peaks for
ribosomal subunits, whereas EDTA treatment completely disrupted polysomes (Fig. 4E). RT-qPCR on
fractions demonstrated the association of 10% and 72% of all HIV-1 RNA and the housekeeping gene
HPRT1 (hypoxanthine phosphoribosyltransferase-1 RNA) with polysomes, which was reduced to 1%
and 2% in EDTA-treated controls, respectively, indicating specific association of these RNAs with
polysomes (Fig. 4F). Taken together, these data provide multiple independent lines of evidence that
demonstrate the production viral proteins in the absence of reverse transcription.

Packaging signal and capsid stability mutations impact the course of incoming retroviral RNA
translation

The packaging signal (Ψ; psi) found in retroviral genomes is critical for the selective packaging
of the full-length viral RNA into budding virions (reviewed in [19]). Deletion of 38 nucleotides (750-787)
downstream of the Gag start codon in LV-DD-Nluc (ΔΨ) drastically reduced the amount of packaged
genomic viral RNA compared to WT (Fig. 5A), even though the expression from the ΔΨ construct by
transfection was comparable to WT with a 2.5-fold increase (Fig. 5B). Infection of cells with the two
viruses showed significantly reduced reporter signal in ΔΨ compared to WT, indicating that the

packaging of the viral genome into particles is essential for incoming viral RNA translation, as

expected (Fig. 5C).

Since the viral RNA genome is protected by the capsid, we reasoned that mutations affecting

capsid stability may alter incoming translation kinetics. To this end, we used two well-characterized

capsid mutants; the hypostable P38A mutant that loses its integrity early after entry, and the

hyperstable E45A mutant which does not easily dissociate. It was previously described that different

titration methods show considerable variation in predicting the transducing units of lentiviruses.[20]

We therefore normalized the viruses to each other by different methods: p24-CA amount by ELISA,

viral genomic RNA by RT-qPCR, virion-packaged Nluc by luciferase assay, or simply by using equal

volumes from viruses produced at the same time. Despite slight variations based on the

normalization method used (Fig 5D-E and Supp. Fig. 3A-B), collectively, infection with the P38A

mutant increased translation from incoming genomes compared to WT, whereas E45A mutant

decreased it (Fig. 5F). These data are consistent with unstable capsids allowing increased access to

the translational machinery, in contrast to hyperstable capsids shielding the viral RNA, linking capsid

stability and uncoating kinetics to the translation potential of incoming retroviral genomes.

We show here that the RNA genome packaged into retroviral particles can serve as an mRNA

for viral protein production independently of reverse transcription. This process occurs in the context

of both minimal and near full-length genomes, in the presence of RT inhibitors and catalytic mutants,

with Env proteins that use different entry pathways and in different cell types with different virus

amounts, suggesting that this is a general process. Using thorough controls that account for nucleic

acid contamination, passive DNA or protein delivery, transcription and translation inhibitors, fusion-

deficient envelopes and omission of various viral components, we confirm that the signal observed is

due to *de novo* translation. Using capsid mutant viruses, we further demonstrate that the translation

potential of the viral RNA is linked to capsid stability as the process of uncoating modulates the

access of the incoming RNA to ribosomes. In summary, we provide multiple lines of evidence that the

retroviral genome can serve as an mRNA early after cellular entry. Below we discuss potential

implications of our findings for basic retrovirology, host immune responses and RNA delivery

approaches.

Discussion

239   Early studies of avian and murine retroviruses reported the detection of full-length viral

240   genomes in polysome-containing pellets within 4 hpi.[21, 22, 23] A recent study also detected full-length

241   HIV-1 transcripts associated with polysomes at 8 hpi; although it is unclear whether the expression at

this time point is mediated by reverse-transcribed and integrated viral DNA that is then newly-transcribed, or from the incoming RNA genome itself, as the HIV-1 RT was functional in this case.[24] In support of our findings, a ribosome profiling study of HIV-1 infected cells detected ribosome-protected RNA fragments indicative of active translation in the *gag* coding region already within one hour of infection.[25] Based on the current knowledge regarding infection kinetics, as one hour is too short of a time period to complete reverse transcription and integration, the implication is that such expression is enabled by the direct translation of the RNA genome. Ribosome profiling studies in the absence of a functional RT at early time points following infection will be informative in identifying the specific regions of incoming retroviral genomes that are translated in different cell types.

In the field of gene therapy, recombinant retroviral vectors are very well-characterized and commonly used as tools for nucleic acid or protein delivery (reviewed in [26]). In a previous study using minimal gammaretroviral (MLV) vectors with primer binding site (PBS) mutations that cannot initiate reverse transcription, protein expression from a reporter gene was detected.[27] In other studies, modified lentiviral vectors containing 5' IRES or 3' WPRE insertions or major structural rearrangements of the genome (e.g. moving the U5-R regions further downstream towards the 3' end) were employed in order to enable direct translation from the packaged RNA.[28, 29] In all of these studies, however, protein or DNA transfer cannot be ruled out. In addition, to our knowledge, the production of actual retroviral proteins from a (near) full-length genome in the absence of reverse transcription has not been demonstrated.

A paradigm shift in retrovirology occurred with the finding that intact or near-intact HIV-1 capsids can be transported into the nucleus, where reverse transcription and uncoating is completed.[30, 31, 32, 33] How to reconcile the direct translation of incoming retroviral genomes surrounded by capsid and inaccessible to the translational machinery with the detection of intact core structures in the nucleus? In a given virus population not all particles are infectious or replication-competent. In fact, for animal viruses the particle-to-PFU (or particle to infectious unit; P/IU) ratio can vary greatly; from 1-2 for Semliki Forest Virus to as high as 10^7 for HIV-1, according to some estimates (reviewed in [34]). Such a high ratio highlights the presence of a large number of particles that may not proceed successfully to next steps of the replication cycle. We believe that incoming retroviral RNA translation occurs in case of particles that are able to enter the host cell, carry a packaged genomic RNA and start uncoating before reaching the nucleus; an idea that has been employed as a basis for the EURT entry/uncoating assay.[35] As direct translation of the genome is likely a dead-end for viral replication, it is reasonable to assume that virions that end up producing infectious progeny are the few ones that successfully make it to the nucleus or the nuclear pore intact, not those that are translated. In the stochastic environment of a viral population, some

genomes are translated, whereas some are reverse transcribed, just as some capsids fall apart before reaching the nucleus while some of them make it to the nucleus almost intact. As many of the assays used in this study are biochemical in nature, we cannot analyze the status of individual virus particles but rather the status of the population as a whole. Single-molecule live-imaging approaches to visualize incoming viral genomic RNAs and newly-synthesized proteins will be valuable in quantifying the percentage of particles that are translated after cellular entry.

The production of viral proteins in the absence of reverse transcription could have major consequences. Individuals who encounter HIV-1 but who do not get productively-infected, for instance due to pre-exposure prophylaxis (PrEP) usage, may still be able to process and present viral peptides to generate cell-mediated and/or humoral immune responses. An initial abortive infection may result in the recruitment and activation of T-cells, increasing the eligible target cell population locally for productive infection. During SIV infection of rhesus macaques, Gag- and Pol-specific CTL responses were identified very early (within 2h) after infection, whereas Env- or Nef-specific responses were not found until later, which was attributed to the ability of incoming viral proteins to be processed and presented.[36,37] Data presented here suggest that such responses may also occur due to *de novo* translation from the viral genome. In summary, our results challenge the notion that retroviruses require reverse transcription to produce viral proteins, warrant careful studies of immune responses during abortive infection and open up novel avenues for gene therapy and targeted vaccine approaches.

Methods

Cloning, constructs and virus production. As a basis for minimal lentiviral vectors the pLVX-IRES-Neo vector (Clontech) was used, which contains identical, full-length LTRs. DD-Nluc was synthesized as a gBlock (IDT) and cloned between XhoI and NotI sites of pLVX-IRES-Neo, either with or without the destabilizing domain to create CMV-(DD)Nluc. Similarly, (DD)Nluc was cloned between ClaI and MluI sites of pLVX-IRES-Neo to eliminate the CMV promoter, multiple cloning site, IRES element and Neo resistance gene to create (DD)Nluc-WPRE. LV-(DD)Nluc was created first by deleting the sequences between ClaI and KpnI sites (including CMV promoter, multiple cloning site, IRES element, Neo resistance gene and the WPRE sequence) in pLVX-IRES-Neo and religating the vector onto itself, then mutating the start codon of Gag, inserting the restriction sites BstBI and PacI after the packaging signal (Ψ) and finally cloning (DD)Nluc between these restriction sites. Packaging signal deletion (ΔΨ) was introduced into LV-DD-Nluc by overlap PCR resulting in a deletion between nucleotides 750-787 based on the pLVX-IRES-Neo vector sequence. NL43-Firefly is pNL4-3 e- r- FLuc (kindly provided by Ned Landau; NIH ARP-3418). NL43-DD-Nluc was created by mutating the Gag

start codon of NL43-Firefly and inserting the DD-Nluc sequence between PteI and SpeI sites. The

identity of all constructs was confirmed by restriction digest and sequencing.

Reporter viruses were produced by transfecting viral plasmids together with a plasmid

encoding VSV-G *env* (pCMV-VSV-G-myc, or the fusion-defective mutant P127D, kindly provided by

Wes Sundquist; Addgene #80054 and #80055), and in case of the minimal lentiviral vectors or NL43-

DD-Nluc, also with a packaging plasmid encoding HIV-1 gag, pol, tat and rev (psPAX2, kindly provided

by Didier Trono; Addgene #12260) or with the packaging plasmid pCD/NL-BH*deltavpu/RT- that

lacks RT activity due to the D110E mutation in the catalytic site (kindly provided by Jakob Reiser;

Addgene #136985). CA stability mutants P38A and E45A were kindly provided by Stephen Goff. The

full-length IIIB ΔRT construct carrying the DD185/186AA mutations in the catalytic domain of RT was

kindly provided by Michael Malim.[38] The HIV-1 *env* plasmid was kindly provided by Yiping Zhu

(University of Rochester). X-MLV *env* was a codon-optimized, V5-tagged, synthetic Xenotropic MLV-

Related Virus (XMRV) *env* in a pTH backbone, kindly provided by Alaa Ramadan.[39]

Virus stocks were produced by transfection of 293T cells with polyethylenimine (PEI),

followed by media change after one day and supernatant collection after two days. ΔEnv or ΔGag

viruses were produced by omitting the respective plasmids during the transfection step. Virus-

containing supernatants were filtered (0.45 μ), ultracentrifuged over a 20% sucrose cushion,

aliquoted and frozen at -80°C. Nuclease treatment of virus stocks was performed at room

temperature by DNase (Ambion), RNase (Qiagen) or benzonase (Millipore).

Cell culture, treatments, transductions. 293T and TZM-bl cells were maintained in DMEM

containing 9% FBS (Gibco) and 100 μg/ml Pen/Strep (Gibco). All suspension cell lines including THP-1,

U937, C8166, Jurkat, PM1 and A3.01 were maintained in RPMI with 9% FBS, 100 μg/ml Pen/Strep,

100 μg/ml Normocin (Invivogen). PBMCs of healthy donors were isolated from buffy coats from the

German Red Cross using standard Ficoll separation. Monocytes were selected by adhering PBMCs in

RPMI with 5% pooled AB human serum (Sigma), 1 mM HEPES (Gibco) and 24 μg/ml gentamicin

(Sigma) for several hours, followed by extensive washing to remove unbound cells. Monocyte-

derived macrophages were differentiated from primary monocytes by 50 ng/ml GM-CSF (R&D

Systems) treatment for 6-10 days. CD4+ T-cells were isolated from PBMCs by negative selection using

the MACS human CD4+ T-cell isolation kit (Miltenyi). Infections were performed by spinoculation at

1200 x g for 60-90 mins at 25°C. Unless indicated otherwise, the following concentrations were used

for treatments: NVP: 10-25 μM (Merck), CHX: 10-100 μg/ml (EMD/Millipore), ActD: 2 μg/ml (Sigma),

Shield1: 1.5-3 μM (Takara). With the exception of time-course experiments, Nluc activity was

measured after 4-6 hours post-infection, using the Nano-Glo luciferase assay system (Promega).

Polysome fractionation. Polysome fractionation was performed as previously with some

modifications.[40] Cells were split the day before such that they would reach a maximum of 60-80%

confluency on the day of the assay. At four hours post-infection with RT-deficient viruses or with WT

viruses in the presence of NVP treatment, cells were treated with 50 µg/ml CHX for 10 min at 37°C,

washed once with ice-cold PBS + CHX (50 µg/ml), collected by scraping in PBS+CHX, pelleted, and

lysed in 1X polysome lysis buffer containing 20 mM Tris-HCl pH 7.5, 150 mM KCl, 5 mM $MgCl_2$, 0.5 %

NP40, 1 mM DTT, 50 µg/ml cycloheximide, and protease inhibitors (Roche) on ice for 10 mins. Lysates

were passed through a 21-gauge needle 12 times, incubated on ice for another 5 mins and cleared by

spinning at 4°C, 13.000 rpm for 10 mins. Cleared lysates were then loaded on 15-45% linear sucrose

gradients (in 20 mM Tris-HCl pH 7.5, 150 mM KCl, 5 mM $MgCl_2$, 50 µg/ml CHX) prepared using a

BioComp gradient master in ultra-clear centrifuge tubes (Beckman Coulter) and centrifuged for 2:30

hours at 36.000 rpm in an SW41 rotor. Fractions (0.5 ml) were collected by a piston gradient

fractionator (Biocomp) with continuous UV absorbance recording at 254 nm. RNA from each fraction

was isolated by phenol-chloroform extraction and quantified by RT-qPCR.

qPCR, RT-qPCR. Viral genomic RNA was isolated from concentrated virus stocks with viral

RNA isolation kit (Qiagen). RNA was treated with Turbo DNase and inactivation beads (Ambion),

cDNA was synthesized using Superscript III (Invitrogen) or RevertAid RT (Thermo) with random

hexamers (Roche). qPCR was performed with the SensiFAST No-ROX Probe Master Mix (Bioline) on a

CFX96 qPCR machine (Bio-Rad) along with standards. Primer and probe sequences used in RT-qPCR

are listed in Table 1.

Target	F/R/P	Sequence	Reference
RT products	F	TGTGTGCCCGTCTGTTGTGT	[41]
	R	GAGTCCTGCGTCGAGAGATC	
	P	CAGTGGCGCCCGAACAGGGA	
2-LTR circles	F	AACTAGGGAACCCACTGCTTAAG	
	R	TCCACAGATCAAGGATATCTTGTC	
	P	ACACTACTTGAAGCACTCAAGGCAAGCTTT	
HIV-1 genome	F	TCTCGACGCAGGACTCG	[42]
	R	TACTGACGCTCTCGCACC	
	P	CTCTCTCCTTCTAGCCTC	
HPRT1 mRNA	F	TCTTTGCTGACCTGCTGGATT	
	R	TTATGTCCCCTGTTGACTGGT	
	P	AGTGATAGATCCATTCCTATGACTGT	

Table 1. List of primers and probes used in the study. F: forward, R: reverse, P: probe.

Immunoprecipitation, western blots, SILAC labeling. Immunoprecipitations were performed

as described with slight modifications.[43] Protein G Dynabeads (Invitrogen) were washed and coated

with anti-p55+p24+p17 antibody (Abcam) by rotating for 15 mins at 4°C. After washing off unbound

antibody, cleared cell lysates were added to the coated beads and incubated at 4°C with rotation for

1 hour. Beads were then separated by a magnet, washed three times and bound proteins were

released from the beads by boiling in the presence of a denaturing loading buffer. Released proteins

were analyzed by SDS-PAGE and western blot as previously described.[44] Briefly, cells were washed

with PBS, scraped, transferred to a tube and washed again with PBS. The pellet was lysed in NP40

lysis buffer (100 mM Tris, 30 mM NaCl, 0.5% NP-40) containing benzonase for 15-30 min on ice.

Lysates were cleared by centrifugation at 10,000 rpm for 5 min, supplemented with denaturing

loading buffer (Invitrogen) and run on an SDS-PAGE. Proteins were transferred to a PVDF membrane

(Millipore), blocked by blocking buffer (Rockland) and incubated with primary and IRdye-labeled

secondary antibodies (Licor). Blots were visualized on an Odyssey scanner (Licor). Anti-p24 antibody

used for HIV-1 capsid detection was AG3.0 (NIH ARP-4121)[45] and 183-H12-5C (NIH-ARP-3537).

Labeling and immunoprecipitation for mass spectrometry was performed using the SILAC

Protein Quantitation-Trypsin kit and MS-compatible Magnetic IP kit from Pierce (Thermo) according

to manufacturer's instructions. Cells were cultured in light medium with 10% dialyzed FBS and

transduced with viruses also produced in light medium. The virus used was VSV-G-pseudotyped,

NL4.3-Firefly reporter virus carrying 10 amino acids from the p6 region of SIVmac (pNL-luc3-SIVp6[17-

26]).[46] Infection was performed in the presence or absence of NVP (25 µM) or CHX (100 µg/ml) at an

MOI of 0.5 by spinoculation. The conditions were: Mock (no virus), virus (no drug), virus + NVP, and

virus + NVP + CHX. At the time of transduction, cells were switched to heavy medium. After

incubation for 18 hours, cells were washed extensively with PBS or PBS+CHX (50 µg/ml) to remove all

heavy media and lysed in IP-MS cell lysis buffer with protease inhibitors on ice. Immunoprecipitation

was performed with an anti-p24 antibody (183-H12-5C; NIH-ARP-3537) using Pierce Protein A/G

magnetic beads. Eluted proteins were analyzed by LC-MS. Samples were prepared and analyzed in

biological triplicates.

Liquid chromatography and mass spectrometry. Peptides were analyzed on an Evosep One

liquid chromatography system coupled online via the CaptiveSpray source to a timsTOF HT mass

spectrometer (Bruker Daltonics). Peptides were manually loaded onto Evotips Pure (Evosep) and

separated using the 30 samples per day (SPD) method on the respective performance column (15 cm

x 75 µm, 1.9 µm, Evosep). Column temperature was kept at 40°C using a column toaster (Bruker

Daltonics) and peptides were ionized using electrospray with a CaptiveSpray emitter (10 µm i.d.,

Bruker Daltonics) at a capillary voltage of 1400 V. The timsTOF HT was operated in ddaPASEF mode in

the m/z range of 100-1,700 and in the ion mobility (IM) range of $0.65 - 1.35$ Vs cm^{-2}.[47] Singly-charged

precursors were filtered out based on their m/z-ion mobility position. Precursor signals above 2,500

arbitrary units were selected for fragmentation using a target value of 20,000 arbitrary units and an

isolation window width of 2 Th below 700 Da and 3 Th above 700 Da. Afterwards, fragmented

precursors were dynamically excluded for 0.4 min. The collision energy was decreased as a function

of the IM from 59 eV at 1/K0 = 1.6 Vs cm^{-1} to 20 eV at 1/K0 = 0.6 Vs cm^{-1}. One cycle consisted of 10

PASEF ramps.

MS data analysis. The LC-IMS-MS/MS data were analyzed using FragPipe (version 20.0).[48]

Spectra were searched using MSFragger against the protein sequences of the human proteome

(UP000005640, UniProtKB) and of HIV-1 (NL4-3 e- r- Fluc [ARP-3418] with a modified SIV p6 between

aa 17-26) with a precursor and fragment mass tolerance of 20 ppm, strict trypsin specificity (Lysine:

K, Arginine: R) and allowing up to two missed cleavage sites. Cysteine carbamidomethylation was set

as a fixed modification and methionine oxidation, N-terminal acetylation of proteins as well as heavy

labeling of lysine and arginine (K + 8.014199 Da, R + 10.008269 Da) as variable modifications. Search

results were validated using Percolator with MSBooster enabled rescoring and converged to false

discovery rates of 1% on all levels. Proteins were quantified using IonQuant based on peptides

consistently identified in all replicates and requiring at least 2 peptides per protein.

p24 ELISA. p24-CA concentrations of viral stocks were determined by a homemade ELISA, as

previously described.[45] Briefly, 96-well plates were coated overnight at 4°C with the AG3.0 anti-p24

antibody diluted 1:100 in carbonate/bicarbonate buffer (Sigma). The next day, plates were washed

three times with wash buffer (PBS+0.05% Tween), blocked with PBS+2% milk powder at 37°C for 1 h

and washed again three times. Meanwhile, the viral supernatants were inactivated by incubating

with a final concentration of 0.2% Tween for 10 min at room temperature. Serial dilutions of viral

supernatants in dilution buffer (PBS+2% milk powder+0.05% Tween) were pipetted into the wells and

incubated at 37°C for 1 h. After three washes, the primary antibody in the form of pooled HIV+ serum

diluted 1:10000 was added to the wells and incubated at 37°C for 1 h. Following another three

washes, the secondary antibody anti-human IgG-HRP (Sigma) was added at 1:1000 dilution. After

three more washes, the substrate solution (12.5 ml phosphate/citrate buffer + 1 OPD tablet (5 mg;

Sigma) + 12 µl 30% H_2O_2 solution) was added and incubated at room temperature for 10 mins.

Reactions were stopped by adding 5% sulphuric acid (H_2SO_4). Absorbance was measured at 492 nm

and 620 nm.

Statistics

All statistics analyses were performed using GraphPad Prism 9. For each figure, the numbers

of biological replicates are as follows: Fig. 1B: n=7, Fig. 1C: n=6, Fig. 1D: n=3, Fig. 1E-F: n=2, Fig. 2B:

n=3, Fig. 2C: n=7, Fig. 2D-E: n=6, Fig. 2F-G: n=3, Fig. 3A: n=3-6, Fig. 3C-D: n=6, Fig. 3E: n=5, Fig. 3F:

n=6-11, Fig. 3G, n=3-6, Fig. 3H, n=6, Fig. 3I-J, n=3, Fig. 4B, n=5, Fig. 4D: n=3, Fig. 4F, n=4, Fig. 5A, n=5-

10, Fig. 5B: n=10, Fig. 5C, n=3, Fig. 5E, n=6, Fig. 5F, n=12. Supp. Fig. 1A: n=2-4, Supp. Fig. 1B-C: n=3,

Supp. Fig. 2A: n=2, Supp. Fig. 3A-B, n=3.

Acknowledgements

We thank members of the Cingöz Lab for helpful discussions, Christine Goffinet and her

group for valuable input and materials, as well as Igor Minia for help with polysome fractionation. We

thank Kornelia Gericke and Sabina Reichert for excellent laboratory management and our colleagues

in RKI-FG18. This work was funded by the priority program of the German Research Council (DFG)

awarded to OC and NB (SPP1923; innate sensing and restriction of retroviruses), intramural funds

from the Robert Koch Institute, as well as the KT Boost Fund awarded to OC by the Klaus Tschira

Foundation and the German Scholars Organization (GSO).

Data availability

444 The data that support the findings of this study are available from the corresponding author

445 upon reasonable request. Large datasets generated in this study have been deposited in the public

446 database PRIDE (https://www.ebi.ac.uk/pride/) under the accession number [N].

Figures and legends

Figure 1

Figure 1. Posttranslational regulation of protein stability reduces protein packaging into lentiviral particles and their delivery into recipient cells. (A) Schematic representation of the constructs generated, not drawn to scale. Nano-luciferase (Nluc) was inserted with or without a destabilizing domain (DD) downstream of a CMV promoter in a minimal lentiviral vector to generate pLVX-(DD)-Nluc (constructs I & II). The CMV promoter, IRES element and Neomycin resistance gene were deleted to generate pLVX-(DD)-ΔCIN (constructs III & IV). (B) 293T cells were transfected with the indicated constructs in the presence or absence of Shield1. Nluc activity was measured after 24 hours. (C) Transfection was performed as in B using different concentrations of Shield1. (D) Virus supernatants produced using the constructs in A with a packaging plasmid, with or without an Env (VSV-G) were assayed for virion-packaged Nluc activity. (E) Infection was performed with the viruses in D with or without NVP and assayed for Nluc activity at the indicated time points. (F) 293T cells were infected with pLVX-(DD)-Nluc carrying a VSV-G Env in the presence or absence of NVP, or with bald viruses without an Env. Nluc activity was measured at 20 hpi and 72 hpi. IRES: internal ribosomal entry site, WPRE: woodchuck hepatitis virus posttranscriptional regulatory element, PBS: primer binding site, psi (Ψ): packaging signal, RRE: rev-response element, PPT: polypurine tract, NVP: nevirapine. Data are presented as means +/- SD. Statistical analyses were performed by multiple Mann-Whitney tests using the False Discovery Rate (FDR) correction of Benjamini, Krieger and Yekutieli. ns: not significant.

Figure 2

Figure 2. Incoming minimal lentiviral RNA genomes are translated following entry into host cells.
(A) Schematic representation of the lentiviral constructs used; not drawn to scale. Nluc was cloned into a
minimal lentiviral vector downstream of the packaging signal (Ψ), with or without the destabilizing domain
(DD), and all other heterologous elements (CMV promoter, IRES, Neo resistance gene, WPRE) were removed.
(B) Cells were transfected with the constructs in A in the presence or absence of Shield1 (S1). Nluc reporter
activity was measured one day later. (C) Cells were infected with serial dilutions of LV-DD-Nluc in the presence
of S1 and NVP, with or without cycloheximide (CHX), followed by Nluc measurement. (D,F) Cells were infected
with VSV-G pseudotyped LV-Nluc virus in the presence or absence of NVP and/or CHX, where the virus was
either kept on the cells (D) or washed away (F). Luciferase activity was measured at the indicated time points.
(E,G) Infection was performed as in D and F, but with LV-DD-Nluc and in the presence of NVP, with or without
S1 and/or CHX. Data are presented as means +/- SD. Statistical analyses were performed using two-tailed
unpaired t-test with Welch's correction. ns: not significant.

Figure 3

Figure 3. Direct expression from incoming lentiviral RNA genomes is truly due to *de novo* translation and independent of reverse transcription.

Unless indicated otherwise, all infections were performed on 293T cells with LV-DD-Nluc virus in the presence of NVP and S1, with or without CHX, and assayed 4-6 hours later. (A) Equal amounts of virus stocks were treated with nucleases or left untreated prior to transduction of cells. (B) Different amounts of plasmid DNA (0.1, 1 or 10 µg) were treated with nucleases or left untreated and visualized on an agarose gel. (C) Cells were transduced with viruses pseudotyped with WT VSV-G, a fusion-deficient mutant of VSV-G (P127D) or without an Env. (D) Transduction with "viruses" produced in the presence or absence of a lentiviral packaging vector (WT vs. ΔGag, respectively). (E) Transduction with reporter virus that carries a catalytic RT mutation (D110E) in the presence of Shield1. (F) Transduction in the presence of S1 and NVP, with or without ActD or CHX. (G) Transfection of cells with a plasmid encoding CMV-driven Nluc in the presence of ActD or CHX. (H) Transduction in the presence of different RT inhibitors at 10 µM. (I) TZM-bl cells were transduced with viruses pseudotyped with HIV-1 Env, X-MLV Env or no Env. (J) The indicated cell types were transduced with reporter viruses.. Data are presented as means +/- SD. Statistical analyses were performed by one-way ANOVA with Tukey's test (A, D, F, G) or by unpaired t-test with Welch's correction (C, E, I). In case of multiple unpaired t-tests, the False Discovery Rate (FDR) correction of Benjamini, Krieger and Yekutieli were used (C, I). ns: not significant. NVP: Nevirapine, EFV: Efavirenz, TDF: Tenofovir Disoproxil Fumarate.

Figure 4. Incoming near full-length HIV-1 genomes are translated to produce viral proteins in the absence of
reverse transcription. (A) Schematic representation of the near full-length HIV-1 constructs, where 'x' denotes
mutated codons. DD-Nluc was cloned into the NL43-Firefly construct downstream of the packaging signal to
generate NL43-DD-Nluc. (B) 293T cells were infected with NL43-DD-Nluc with nevirapine (NVP) in the presence
or absence of Shield1 (S1) and/or CHX. Nluc activity was measured at the indicated time points. (C) Cells were
infected with VSV-G-pseudotyped NL43-Firefly with or without NVP or CHX. Lysates were collected one day
after infection, immunoprecipitated with a polyclonal anti-HIV-1 Gag antibody and probed for p24. (D) Cells
were labeled with heavy amino acids (SILAC) at the time of infection (MOI = 0.5) in the presence of the
indicated drugs. 18 hours later, Gag was immunoprecipitated from cell lysates and bound proteins were
analyzed by mass spectrometry. Data are represented as the heavy to light ratio of peptides that map to Gag.
(E) Representative polysome fractionation profiles from 293T cells infected (or not) with VSV-G pseudotyped
IIIB ΔRT (with a catalytic RT mutation DD185/186AA) at 4 hpi and lysed in the presence or absence of EDTA.
Lysates were run on a sucrose density gradient, then fractions were collected from top to bottom while
simultaneously measuring UV absorbance. (F) RNA was isolated from each fraction and the levels of HIV-1
genomic RNA and *HPRT1* were measured by RT-qPCR. The amount of each RNA is given as a percentage of total
RNA for that message. Data represent mean +/- SD. Statistical significance was determined by unpaired t-test
with Welch's correction, nd: not detected.

Figure 5

Figure 5. Packaging signal and capsid stability affect the translation of incoming RNA genomes.
(A) Viral RNA content of LV-DD-Nluc viruses produced with or without a deletion between nucleotides 750-787
in the packaging signal (ΔΨ; delta psi) measured by RT-qPCR. (B) 293T cells were transfected with the LV-DD-
Nluc WT or ΔΨ construct and subjected to luciferase assay one day later. (C) Cells were transduced with WT or
ΔΨ viruses; expression from the incoming viral genome was analyzed by luciferase assay at 4-6 hpi. (D) Western
blot of p24-CA in concentrated WT or the indicated capsid mutant virus stocks. (E) RT-qPCR for quantification of
viral RNA. (F) Cells were transduced with reporter viruses carrying WT or mutated CA. All transductions were
performed in 293T cells using LV-DD-Nluc viruses in the presence of both NVP and Shield1, with or without
CHX. Data represent mean +/- SD. Statistical significance was determined by unpaired t-test with Welch's
correction (C) or one-way ANOVA with Tukey's correction (F). ns: not significant.

     Supplementary Figures and Legends

Supp. Figure 1

Supplementary Figure 1. Loss of RT activity by catalytic mutations or the use of different RT inhibitors.
(A) PERT assay on RT catalytic mutant virus HIV-1 IIIB (DD185/186AA) used in polysome fractionation, or the LV-
DD-Nluc and NL43-DD-Nluc viruses produced with the RT catalytic mutant packaging vector (D110E) used in
reporter assays. (B) Luciferase assay on cells transduced with LV-DD-Nluc virus in the presence of 1-100 μM of
the RT inhibitors, with or without CHX. (C) Cells were transfected with a CMV-driven Nluc construct in the
presence of 100 μM RT inhibitors. Toxicity was measured by luciferase assay 6 hours after transfection. NVP:
Nevirapine, EFV: Efavirenz, TDF: Tenofovir disoproxil fumarate. Data represent mean +/- SD.
535

Supp. Figure 2

Supplementary Figure 2. Quantification of post-entry events after virus challenge.
(A) 293T cells were infected with NL43-Firefly virus (40 ng/100K cells) and DNA was isolated at 24 and 48 hours
post-infection. Early and late RT products, as well as 2-LTR circles were quantified by qPCR.

Supp. Figure 3

Supplementary Figure 3. Capsid stability mutations alter the translation potential of incoming retroviral
genomes.
(A) Cells were infected with LV-DD-Nluc viruses containing WT, P38A or E45A capsids in the presence of NVP
and Shield1, with or without CHX. Virus stocks were normalized based on either p24, viral RNA, virion-packaged
Nluc or simple volume. (B) Fold difference between CHX-treated and untreated samples based on the data in
(A). Data represent mean +/- SD.

References

1. Brown JD*, et al.* Structural basis for transcriptional start site control of HIV-1 RNA fate. *Science (New York, NY)* **368**, 413-417 (2020).

2. Kharytonchyk S*, et al.* Transcriptional start site heterogeneity modulates the structure and function of the HIV-1 genome. *PNAS* **113**, 13378-13383 (2016).

3. Rawson JMO, Nikolaitchik OA, Shakya S, Keele BF, Pathak VK, Hu WS. Transcription Start Site Heterogeneity and Preferential Packaging of Specific Full-Length RNA Species Are Conserved Features of Primate Lentiviruses. *Microbiol Spectr* **10**, e0105322 (2022).

4. Singh G, Seufzer B, Song Z, Zucko D, Heng X, Boris-Lawrie K. HIV-1 hypermethylated guanosine cap licenses specialized translation unaffected by mTOR. *PNAS* **119**, e2105153118 (2022).

5. Ghysdael J, Kettmann R, Burny A. Translation of bovine leukemia virus virion RNAs in heterologous protein-synthesizing systems. *Journal of virology* **29**, 1087-1098 (1979).

6. Kobayashi N, Yamamoto N, Koyanagi Y, Schneider J, Hunsmann G, Hatanaka M. Translation of HTLV (human T-cell leukemia virus) RNA in a nuclease-treated rabbit reticulocyte system. *Embo j* **3**, 321-325 (1984).

7. Philipson L, Andersson P, Olshevsky U, Weinberg R, Baltimore D, Gesteland R. Translation of MuLV and MSV RNAs in nuclease-treated reticulocyte extracts: Enhancement of the gag-pol polypeptide with yeast suppressor tRNA. *Cell* **13**, 189-199 (1978).

8. Nash KL, Lever AM. Green fluorescent protein: green cells do not always indicate gene expression. *Gene therapy* **11**, 882-883 (2004).

9. Liu ML, Winther BL, Kay MA. Pseudotransduction of hepatocytes by using concentrated pseudotyped vesicular stomatitis virus G glycoprotein (VSV-G)-Moloney murine leukemia virus-derived retrovirus vectors: comparison of VSV-G and amphotropic vectors for hepatic gene transfer. *Journal of virology* **70**, 2497-2502 (1996).

10. Geering B, Schmidt-Mende J, Federzoni E, Stoeckle C, Simon HU. Protein overexpression following lentiviral infection of primary mature neutrophils is due to pseudotransduction. *Journal of immunological methods* **373**, 209-218 (2011).

11. Gallardo HF, Tan C, Ory D, Sadelain M. Recombinant retroviruses pseudotyped with the vesicular stomatitis virus G glycoprotein mediate both stable gene transfer and pseudotransduction in human peripheral blood lymphocytes. *Blood* **90**, 952-957 (1997).

12. Kim JT*, et al.* Dendritic cell-targeted lentiviral vector immunization uses pseudotransduction and DNA-mediated STING and cGAS activation. *Sci Immunol* **2**, eaal1329 (2017).

13. Haas DL, Case SS, Crooks GM, Kohn DB. Critical Factors Influencing Stable Transduction of Human CD34+ Cells with HIV-1-Derived Lentiviral Vectors. *Molecular Therapy* **2**, 71-80 (2000).

14. Banaszynski LA, Chen L-c, Maynard-Smith LA, Ooi AGL, Wandless TJ. A Rapid, Reversible, and Tunable Method to Regulate Protein Function in Living Cells Using Synthetic Small Molecules. *Cell* **126**, 995-1004 (2006).

15. England CG, Ehlerding EB, Cai W. NanoLuc: A Small Luciferase Is Brightening Up the Field of Bioluminescence. *Bioconjugate Chemistry* **27**, 1175-1187 (2016).

16. Donahue DA, Sloan RD, Kuhl BD, Bar-Magen T, Schader SM, Wainberg MA. Stage-dependent inhibition of HIV-1 replication by antiretroviral drugs in cell culture. *Antimicrob Agents Chemother* **54**, 1047-1054 (2010).

17. Murray JM, Kelleher AD, Cooper DA. Timing of the components of the HIV life cycle in productively infected CD4+ T cells in a population of HIV-infected individuals. *Journal of virology* **85**, 10798-10805 (2011).

18. Watts JM*, et al.* Architecture and secondary structure of an entire HIV-1 RNA genome. *Nature* **460**, 711-716 (2009).

19. Rein A. RNA Packaging in HIV. *Trends in Microbiology* **27**, 715-723 (2019).

20. Geraerts M, Willems S, Baekelandt V, Debyser Z, Gijsbers R. Comparison of lentiviral vector titration methods. *BMC Biotechnology* **6**, 34 (2006).

21. Gallis BM, Eisenman RN, Diggelmann H. Synthesis of the precursor to avian RNA tumor virus internal structural proteins early after infection. *Virology* **74**, 302-313 (1976).

22. Shurtz R, Dolev S, Aboud M, Salzberg S. Viral genome RNA serves as messenger early in the infectious cycle of murine leukemia virus. *Journal of virology* **31**, 668-676 (1979).

23. Tobaly J, d'Auriol L, Yang WK, Peries J, Emanoil-Ravicovitch R. Murine retrovirus genome directs the synthesis of gag protein precursor early after infection. *Biochimie* **64**, 969-973 (1982).

24. Caliskan N KA, Buck S, Gribling A-S, Gilmer O, Pekarek L, Bohn P, Koch T, Mireisz CN-M, Schlosser A, Erhard F, Smyth R. The translational landscape of HIV-1 infected cells reveals novel gene regulatory principles. Preprint at https://www.researchsquare.com/article/rs-3352709/v3352701 (2023).

25. Labaronne E*, et al.* Extensive uORF translation from HIV-1 transcripts conditions DDX3 dependency for expression of main ORFs and elicits specific T cell immune responses in infected individuals. Preprint at https://www.biorxiv.org/content/10.1101/2022.1104.1129.489990v489991.full (2022).

26. Baum C, Schambach A, Bohne J, Galla M. Retrovirus Vectors: Toward the Plentivirus? *Molecular Therapy* **13**, 1050-1063 (2006).

27. Galla M, Will E, Kraunus J, Chen L, Baum C. Retroviral pseudotransduction for targeted cell manipulation. *Molecular cell* **16**, 309-315 (2004).

28. Mock U*, et al.* Novel lentiviral vectors with mutated reverse transcriptase for mRNA delivery of TALE nucleases. *Scientific Reports* **4**, 6409 (2014).

29. Counsell JR*, et al.* Re-structuring lentiviral vectors to express genomic RNA via cap-dependent translation. *Molecular Therapy* **20**, 357-365 (2021).

30. Zila V*, et al.* Cone-shaped HIV-1 capsids are transported through intact nuclear pores. *Cell* **184**, 1032-1046.e1018 (2021).

31. Burdick RC*, et al.* HIV-1 uncoats in the nucleus near sites of integration. *Proceedings of the National Academy of Sciences* **117**, 5486-5493 (2020).

32. Müller TG*, et al.* HIV-1 uncoating by release of viral cDNA from capsid-like structures in the nucleus of infected cells. *eLife* **10**, e64776 (2021).

33. Li C, Burdick RC, Nagashima K, Hu WS, Pathak VK. HIV-1 cores retain their integrity until minutes before uncoating in the nucleus. *Proceedings of the National Academy of Sciences of the United States of America* **118**, (2021).

34. Klasse PJ. Molecular determinants of the ratio of inert to infectious virus particles. *Prog Mol Biol Transl Sci* **129**, 285-326 (2015).

35. Da Silva Santos C, Tartour K, Cimarelli A. A Novel Entry/Uncoating Assay Reveals the Presence of at Least Two Species of Viral Capsids During Synchronized HIV-1 Infection. *PLOS Pathogens* **12**, e1005897 (2016).

36. Sacha JB*, et al.* Gag-specific CD8+ T lymphocytes recognize infected cells before AIDS-virus integration and viral protein expression. *J Immunol* **178**, 2746-2754 (2007).

37. Sacha JB*, et al.* Pol-specific CD8+ T cells recognize simian immunodeficiency virus-infected cells prior to Nef-mediated major histocompatibility complex class I downregulation. *J Virol* **81**, 11703-11712 (2007).

38. Bauby H*, et al.* HIV-1 Vpr Induces Widespread Transcriptomic Changes in CD4+ T Cells Early Postinfection. *mBio* **12**, e01369-01321 (2021).

39. Hong S*, et al.* Fibrils of prostatic acid phosphatase fragments boost infections with XMRV (xenotropic murine leukemia virus-related virus), a human retrovirus associated with prostate cancer. *Journal of virology* **83**, 6995-7003 (2009).

40. Cingöz O, Goff SP. Cyclin-dependent kinase activity is required for type I interferon production. *PNAS* **115**, E2950-e2959 (2018).

41. Mbisa JL, Delviks-Frankenberry KA, Thomas JA, Gorelick RJ, Pathak VK. Real-time PCR analysis of HIV-1 replication post-entry events. *Methods Mol Biol* **485**, 55-72 (2009).

42. Malnati MS*, et al.* A universal real-time PCR assay for the quantification of group-M HIV-1 proviral load. *Nature Protocols* **3**, 1240-1248 (2008).

43. Zhu Y, Wang GZ, Cingöz O, Goff SP. NP220 mediates silencing of unintegrated retroviral DNA. *Nature* **564**, 278-282 (2018).

44. Cingöz O, Arnow ND, Puig Torrents M, Bannert N. Vpx enhances innate immune responses independently of SAMHD1 during HIV-1 infection. *Retrovirology* **18**, 4 (2021).

45. Sanders-Beer BE, Eschricht M, Seifried J, Hirsch VM, Allan JS, Norley S. Characterization of a monoclonal anti-capsid antibody that cross-reacts with three major primate lentivirus lineages. *Virology* **422**, 402-412 (2012).

46. Sunseri N, O'Brien M, Bhardwaj N, Landau NR. Human Immunodeficiency Virus Type 1 Modified To Package Simian Immunodeficiency Virus Vpx Efficiently Infects Macrophages and Dendritic Cells. *Journal of virology* **85**, 6263-6274 (2011).

47. Meier F*, et al.* Online Parallel Accumulation-Serial Fragmentation (PASEF) with a Novel Trapped Ion Mobility Mass Spectrometer. *Mol Cell Proteomics* **17**, 2534-2545 (2018).

48. Yu F, Haynes SE, Teo GC, Avtonomov DM, Polasky DA, Nesvizhskii AI. Fast Quantitative Analysis of timsTOF PASEF Data with MSFragger and IonQuant. *Mol Cell Proteomics* **19**, 1575-1585 (2020).