RenSeq of *Silphium integrifolium* Reveals a Biotic Gradient for Disease Resistance

Kyle Keepers[1], Kelsey Peterson[1], Andrew Raduski[1†], Kathryn M. Turner[2], David Van Tassel[2], Kevin Smith[3], Alex Harkess[4], James D. Bever[5], Yaniv Brandvain[1]

[1]University of Minnesota, Department of Plant and Microbial Biology, 1500 Gortner Ave, St Paul, MN 55108

[2]The Land Institute, Crop Protection Ecology, 2440 E Water Well Rd, Salina, KS 67401

[3]University of Minnesota, Department of Agronomy and Plant Genetics, 1529 Lindig St, St Paul, MN 55108

[4]HudsonAlpha Institute for Biotechnology, 601 Genome Way, Huntsville, AL 35806

[5]University of Kansas, Department of Ecology and Evolutionary Biology, 1200 Sunnyside Ave, Lawrence, KS 66045

ABSTRACT

Intracellular plant defense against pathogens is mediated by a class of disease resistance genes known as NB-LRRs or NLRs (R genes). Many of the diseases these genes protect against are more prevalent in regions of higher rainfall, which provide better growth conditions for the pathogens. As such, we expect a higher selective pressure for the maintenance and proliferation of R genes in plants adapted to wetter conditions. In this study, we enriched libraries for R genes using RenSeq from baits primarily developed from the common sunflower (*Helianthus annuus*) reference genome. We sequenced the R gene libraries of *Silphium integrifolium* Michx, a perennial relative of sunflower, from 12 prairie remnants across a rainfall gradient in the Central Plains of the United States, with both Illumina short-read (n=99) and PacBio long-read (n=10) approaches. We found a positive relationship between the mean effective annual precipitation of a plant's source prairie remnant and the number of R genes in its genome, consistent with intensity of plant pathogen coevolution increasing with precipitation. We show that RenSeq can be applied to the study of non-model relatives of model organisms in the study of ecological hypotheses.

Keywords: *Silphium integrifolium*, R genes, RenSeq, common gardens, host-pathogen dynamics, prairies

INTRODUCTION

Silphium integrifolium Michx (commonly known as rosinweed or silflower) is a native perennial prairie species being domesticated for production as a forage and oilseed species (Vilela et al., 2018; Van Tassel et al., 2017). The plant's perennial nature allows it to provide novel ecosystem services in

agronomic production systems (Glover et al., 2010; Van Tassel et al., 2017) while producing oilseeds
with similar nutritional composition to annual sunflower (*Helianthus annuus* L. (Asterales: Asteraceae)
seeds (Reinert et al., 2018). However, *S. integrifolium* suffers infections from diverse pathogen
enemies across its native range. The stems, leaves, and flowers can be infected with multiple strains
(Turner, unpublished observations) of *Puccinia silphii* Schwein (1832), or *Silphium* rust (Turner et al.,
2018), *Colletotrichum silphii* Davis (1919) (Horst 2008) syn. and *Colletotrichum dematium* (Pers.)
Grove (Cybernome; Farr 1989) called leaf blotch hereafter. Recently, *Silphium* clear vein (SCV) leaf
and stem disease has been described and hypothesized to be caused by a virus based on the
identification of viral sequences in *Silphium integrifolium* that are similar to the Dahlia common Mosaic
Virus (DCMV) and Dahlia endogenous plant pararetroviral sequence (DvEPRS, formerly DMV-D10)
(Almeyda, 2014; Cassetta et al., in prep). Understanding how *S. integrifolium* defends against these
pathogens is crucial for progress in breeding programs and will elucidate evolutionary questions posed
in the species (Cassetta et al., in prep, Van Tassel et al, 2017). In this study, plants from 12 wild
populations of *S. integrifolium* were planted in a reciprocal transplant design to better understand
adaptation to disease pressures along a rainfall gradient. Phenotypic data revealed that these wild
populations have distinctive morphologies showed variable levels of tolerance to biotic stresses, such
as pests and pathogens (Peterson et al., in prep). Individuals with resistance to diseases of *S.*
integrifolium were visually identified in the study, and have been previously identified in breeding
populations (Turner et al., 2018), but the genetic basis of disease resistance has not been
investigated. Here, we look at the diversity of disease resistance genes in wild populations of *S.*
integrifolium to answer evolutionary questions regarding adaptation of wild populations to their
pathogen pressures and suggest applications for germplasm enhancement.
Environments with high precipitation are conducive to increased pathogen pressures (Clarkson
et al., 2014; Granke and Hausbeck 2010; Islam and Toyota, 2004; Magarey et al., 2005; Rowlandson
et al., 2015). Several studies have found patterns of resistance alleles consistent with pathogen
selection intensity increasing with precipitation (Wahl 1970, Abbott, Brown, and Burdon 1991, Dong et
al. 2009). We hypothesize that *S. integrifolium* populations evolved in areas with high precipitation will
have a more diverse complement of disease resistance genes to deal with this challenge. To test this
hypothesis, we collected seeds from *S. integrifolium* stands from four prairie remnants each in three
geographic regions in the Central Plains of the United States. The regions are situated along a
gradient of effective precipitation, or the amount of rainfall that isn't lost to potential evapotranspiration,
referred to in this study as the climate moisture index (CMI). We refer to these regions as "West", from
West-Central Kansas, "Central", from Eastern Kansas, and "East", from Central Illinois (Figure 1).
Prairie remnants in the West receive around -55 kg/m2/month of CMI per year, whereas sites in the
East receive closer to -15 kg/m2/month (Table 1). We note that plant defensive gene diversity
increased along this gradient in *Andropogon gerardii*, a native perennial grass that co-occurs with *S.*
integrifolium (Rouse et al. 2011) and soil pathogen diversity also increased along this same gradient
(Delavaux et al. 2021).

A major class of disease resistance genes in plants are the cytoplasmic nucleotide binding and
leucine-rich repeat (NLR) genes. (Caplan et al., 2008; Eitas and Dangl, 2010). These NLRs,
(hereafter, R genes), often contain an N-terminal domain that is either of a coiled-coil type, called
CNLs, or a toll-interleukin-like type (TNLs), whose divergence predates the split between monocots
and eudicots (Meyers et al., 1999), estimated to be about 160 Mya by TimeTree (Kumar et al., 2017).
R genes are highly diverse, numbering in the hundreds in many plant species (Jupe et al., 2012; Toda
et al., 2020). These genes evolve rapidly and are enriched in presence/absence variation even at the
intraspecies level, limiting the utility of a single reference genome to elucidate the diversity of R genes
within a species. This study employed R-gene enrichment sequencing (RenSeq; Jupe et al, 2013),
which uses DNA baits to reduce complexity of genomic DNA libraries. The baits we designed come
from the common sunflower, *Helianthus annuus* L., which is separated from *S. integrifolium* by
between 13.3 Ma (Mccann et al., 2018) and 19.3 Ma (Barreda et al., 2015). We sequenced the
RenSeq libraries from *S. integrifolium* plants across the rainfall gradient using both Illumina short-read
and PacBio long-read technologies. Using both simple linear models as well as mixed effect models,
we found a significant positive correlation between the effective precipitation of a plant's host prairie
and the number of R genes detected in both the Illumina and PacBio datasets. This study
demonstrates the economic utility of using RenSeq for the evaluation of ecological hypotheses, and
has broader implications for the sourcing of genetic diversity in wild populations for the development of
new crops.
95

METHODS
Plant Materials
Silphium integrifolium seeds were collected from 12 prairies in 2017 and 2018 (Figure 1; Table 1). The
prairie remnants collected spanned from central Kansas to central Illinois. The prairie remnants have
variable historical, geographic, and climatic characteristics described in Cassetta et al 2022, in prep.
From each of the 12 prairies, seeds were collected from approximately 40 mother plants randomly
chosen from throughout the prairie remnant. As *S. integrifolium* is an obligate outcrosser, seeds
collected from a single mother were a half sibling family.
S. integrifolium seeds from each prairie were placed on trays lined with moist germination paper
(Hoffman Manufacturing, Inc., Corvallis, OR, U.S.A.) in mid February, 2019, covered to maintain high
humidity, and stratified at 5 C for 3 weeks before returning the trays to room temperature. Individual
seeds were removed from the tray as soon as the emerging radicle was visible, and planted into
plastic plug trays (72 square cells per flat, Landmark Plastic, Akron, OH, U.S.A.) filled with BM6 peat-
pearlite medium (Berger, Saint-Modeste, Quebec, Canada). Plants were germinated at room
temperature together in a heated greenhouse at The Land Institute, Salina KS in 2019. A total of 96
genets comprising eight individuals from each prairie remnant from different mothers were selected
from the seedlings. Selection of the 96 individuals was randomized although the seedlings chosen had
to be moderately vigorous.

In early April of 2019, these seedlings were planted in 25-gallon fabric planters (Smart Pot,
Oklahoma City, OK, U.S.A.) filled with Pro-Mix BX (Premiere Horticulture, Rivière-du-Loup, Quebec,
Canada). The pots containing the seedlings were randomized and placed on greenhouse benches.
Pots were re-randomized several times during the season. Pots were sub-irrigated by an automated
ebb-and-flow system every 2 days for about 20 minutes. The irrigation frequency was increased as
plants grew and required more water. Natural light was supplemented automatically on cloudy days by
red and blue LED overhead lights, and these lights were used to extend natural day length to 14 hours
from April through December. Fertilizer was provided by 4 slow-release 7.5 g 15-8-11 (N-P-K) pellets
inserted into the root zone. Additional fertilizer pellets were inserted as necessary to maintain deep
green foliage. The plants grew rapidly but remained as vegetative rosettes. In late December 2019,
the foliage was trimmed off and the crowns were dug out of the containers. Crowns were placed in
plastic bags and vernalized in a dark, refrigerated room (5 C). Between February 05 and Feb 27,
2020, the crowns were removed from refrigeration and individually washed. Each crown (genet) was
divided into 4 pieces (ramets) with approximately equal numbers of dormant buds and roots, although
a few crowns were too small and only produced 2-3 pieces. The ramets were repotted into 10-gallon
fabric planters (Smart Pot) using Pro-Mix BX.
130

Tissue Collection
One ramet from each genet was grown in The Land Institute Greenhouse until young leaves had
emerged. On March 12, 2020, black plastic pots were placed over one clone of each of the 99 plants
emerging rosette and left for 48 hours to slightly etiolate the young leaves, which improves the quality
of extracted DNA in photosynthetic active tissue containing phenolic compounds (Dabe et al., 1993).
On March 14, 3-5 unexpanded leaves were excised from each crown. The collected tissue was placed
in a labeled envelope and immediately placed on dry ice in a cooler. Within 60 minutes of collection,
the sampled leaf tissue was placed in a lyophilizer and lyophilized for 48 hours.
139

Clone placement across the gradient and phenotyping
After young leaf tissue was collected from plants in the greenhouse, the plants remained in the
greenhouse under ideal conditions allowing them to grow rapidly. One cloned ramet from each of the
available genets was then moved to each of the following locations within the regions the seed was
sourced from: 1. breeding plots in Salina, KS at The Land Institute (West), 2. Prairie Park Nature
Center, Lawrence, KS (Central), 3. research plots at the ISU horticulture center in Normal, Illinois
(East), and 4. Land Institute Crop Protection Ecology greenhouse (backup site to conserve genets in a
disease and insect free state). The pots remained undisturbed at each site for two years, 2020 and
2021 except the Central site cloned plants, which, in May 2021, were moved from the Prairie Park
Nature Center to the Perennial Agriculture Project field station, approximately, 6 miles away. The
clones received occasional watering in each site to promote disease growth and avoid plant death due
to the above ground pots drying out.

In 2020 and 2021, disease scores and insect herbivory damage were recorded on each of the clones
in all 3 sites. Data was collected during their flowering period, between the months of July and
September. Diseases were recorded according to methods in Cassetta et al., in prep. Briefly, leaf rust
severity was quantified by visually assessing the total percentage of leaf area affected (0-100%) using
the rust scale described by Turner, et al. (2018). Similarly, stem rust severity was recorded as the total
percentage of the stems covered in rust (0-100%). Leaf blotch was quantified by assessing the total
percentage of leaf area affected (0-100%). Silphium clear vein was rated on a 0-10 scale based on
symptoms observed on the entire plant including: leaf distortion (narrow width, curled, toothed,
twisted, or pinched tips), and twisted meristem symptoms. Leaf herbivory scores were collected by
quantifying the percentage of total leaves with evidence of insect herbivore (mining, sucking, chewing)
damage (0-100%).
163

Climate Moisture Index
A BIOCLIM+ raster map of the continental U.S.A. containing 30-year climate moisture index averages
(1981-2010) was downloaded from the CHELSA (Climatologies at high resolution for the Earth's land
surface areas) database (Brun et al., 2022). The climate moisture index is calculated as the difference
between mean annual precipitation and mean potential evapotranspiration and, when negative,
represents the availability of water as a limiting resource in dry regions such as prairies.
170

Sequencing/RenSeq/QC
DNA was extracted from lyophilized leaf tissue using the CTAB method by the University of Minnesota
Genomics Center. DNA was then shipped on dry ice to Arbor Biosciences for sequencing, library
preparation, and RenSeq data acquisition. Extracted DNA was prepared for sequencing using Illumina
library prep kits. Sequencing libraries were enriched for R genes using the RenSeq protocol described
in Jupe et al., 2013 using Arbor's myBaits systems. In short, biotinylated single-stranded RNA oligo
baits were designed based on annotated R genes from *Helianthus annuus*, a relative of *Silphium*
integrifolium, within the same subtribe, estimated to be between 13.3 Ma (Mccann et al., 2018) and
19.3 Ma (Barreda et al., 2015) divergent. Hybridized oligos were pulled down using streptavidin-
coated magnetic beads. Libraries were then sequenced on an Illumina NovaSeq S4 sequencer using
paired-end 150bp reads by Arbor Biotechnologies in Cambridge, MA.
A subsample of 10 individuals was selected for a complementary R gene analysis using Pacific
Biosciences (PacBio) long-read sequencing. High-molecular-weight enriched DNA from three
individuals from each of the West, Central, and East prairies, as well as one individual derived from a
cultivated lineage, were prepared for long read sequencing and then sequenced on a Pacific
Biosciences Sequel II using circular consensus sequencing (CCS) by Arbor Biotechnologies.
A total of 99 Illumina libraries (96 from prairie remnants + 3 cultivated "elite" lineages) and 10 PacBio
libraries were sequenced. Both library types were filtered for sequence quality using Trimmomatic
v0.39 (Bolger et al., 2014) and then assessed using fastqc v0.11.7 (Andrews 2010). The Illumina

libraries were subsampled to a standardized assembly input size of 1Gb to avoid biasing R gene
counts by library input size. The PacBio libraries were not subsampled to a standard input size, but a
correlation between input size and total R gene count in this data type reveals no dependence of final
assembly size on initial input size (Figure S4; p=0.414). Six Illumina libraries failed to meet the input
threshold and were excluded from analysis. The libraries that met the input threshold were then *de*
novo assembled using SPAdes v3.15.3 (Bankevich et al., 2012) with -k 21,33,45,55,65,81.
196

Draft Genome Assembly
A chromosome-scale genomic assembly of an F1 *Silphium integrifolium* x *S. perfoliatum* is being
prepared at the time of writing. The *S. integrifolium* mother was a leaf blotch resistant, large-headed
selection from a breeding population that had undergone approximately 6 cycles of recurrent selection
for increased number of fertile ray florets ("feminization") per head and disease resistance. The
founders of the breeding population were sourced from wild central Kansas populations, i.e., located
within the geographic region we call "West" in this study. Flash-frozen tissue was sent to Arizona
Genomics Institute for high-molecular-weight DNA extraction, which was then prepared for PacBio
Sequel-II CCS sequencing at the Genome Sequencing Center at HudsonAlpha Institute for
Biotechnology. The libraries were sequenced at roughly 21.5X depth assuming a 10 Gb genome size.
Reads were assembled *de novo* using hifiasm assembler (Cheng et al., 2021) using default
parameters.
209

R Gene Annotations
We annotated the Illumina RenSeq libraries, PacBio RenSeq libraries, and the draft genome for their
R gene content. Each assembly was annotated for R genes using NLR-Annotator (Steuernagel et al.,
2020), which searches for common R gene peptide motifs in protein-coding sequence. The
annotations were performed with the "-a" flag, which outputs the amino acid sequence of the
nucleotide-binding domain (NBARC domain). We used the NBARC sequences of the draft genomic
assembly to infer a maximum likelihood tree in RAxML v.8.2.11 (Stamatakis 2006; Stamatakis 2014)
using the peptide substitution model "PROTGAMMAAUTO" with 100 rapid bootstraps.
The two broad, non-overlapping classes of R genes that we analyzed in this study, TNLs (containing a
toll-interleukin-like type N-terminal domain) and CNLs (containing a coiled-coil type N-terminal
domain), were identified from the annotations based on diagnostic motifs. To identify TNLs, we parsed
annotations for the motifs 13, 15, and 18 (as defined in Table S1 of Jupe et al., 2013). For CNLs, we
searched for motifs 2, 6, 16 and 17. These two sets of motifs occur only in their respective class of N-
terminal domain.
224

Principal Coordinates Analysis
To test for associations between R genes and the phenotypes collected on the cloned plants across
the gradient, we performed simple linear modeling as well as mixed effect modeling, which treats CMI

as a fixed effect, and source prairie as a random effect. We found one significant correlation between
R Genes of the various populations and leaf blotch. To interrogate which R genes were responsible for
the association between leaf blotch and R gene counts, we performed a census of which R gene
NBARC sequences occurred in which individuals (Table S5). The census was then used to generate a
principal coordinate analysis in the R package "vegan" (Oksanen 2015) against each of the five
phenotypic covariates included in the study. P-values were corrected for multiple tests using the
Benjamini & Hochberg method (1995).
235

Library Enrichment Calculations
The fasta files containing the annotated R genes for each Illumina library were concatenated into a
fasta containing every observed R gene from the Illumina dataset (hence, a "Pan-NLRome"). Reads
from each Illumina RenSeq library, as well as eleven 1X-depth WGS libraries from seven *S.*
integrifolium, two *S. perfoliatum*, one *S. laciniatum* and one *S. terebinthinaceum*, were mapped to the
Pan-NLRome using *bwa mem* (Li 2013). The fraction out of the total reads in the library that mapped
to the Pan-NLRome with a SAM CIGAR mapping score of between 148M-151M was calculated,
representing reads that mapped to an R gene in the Pan-NLRome with between 98.01%-100%
identity.
245

RESULTS
Reference Genome Assembly
The draft genome used in this study was 9.9 Gb in length and totalled 2,584 contigs. The longest
contig was 182 Mb long and the mean contig size was 3.8 Mb. The L50 (which is the number of
contigs, ordered longest-to-smallest, to comprise 50% of the assembly size) was 77, and the N50 (the
size of the contig that passes the 50% assembly size threshold) was 40.2 Mb.
The maximum likelihood tree constructed from the R genes in the draft assembly was
annotated according to peptide motifs that are diagnostic for either CNL or TNL identity (Figure 2). A
deep evolutionary divergence between the two broad classes is seen, although we observe isolated
instances of R genes falling incongruently within the wrong clade.
256

Library Enrichment
The libraries of seven *S. integrifolium* that were sequenced using WGS, and thus were unenriched for
R genes, contained a median of 1.78% of reads that mapped with high fidelity (98% identity or higher)
to contigs in the Pan-NLRome. Libraries that had been enriched for R genes using oligo baits
designed from *Helianthus annuus* contained a median of 63.4% of high fidelity reads mapping to R
genes, representing a 36-fold enrichment for R genes (Figure 3).
263

R Gene Counts by Geographic Region

The Illumina R gene assemblies ranged from 2.2 Mb to 51 Mb, and averaged 15 Mb in size. We

detected an average of 386.3, 410.0, and 414.0 R genes in plants sourced from the Western, Central,

and Eastern prairies, respectively (left facet of Figure 4A). The plants averaged 47.38 (West), 48.69

(Central), and 45.14 (East) TNL genes, and 166.1 (West), 171.3 (Central), and 175.3 (East) CNL

genes (Table S2).

The PacBio R gene assemblies ranged from 4.9 Mb to 13 Mb, but averaged 8 Mb. Within this

dataset, we only had three plants per region, so the inferential power was lower, but our confidence in

the contigs assembled was higher due to longer reads. We found an average of 355.0, 563.3, and

592.7 R genes in the Western, Central, and Eastern prairies, respectively (right facet of Figure 4A).

Within this dataset, we found an average of 54.00 (West), 81.67 (Central), and 81.33 (East) TNLs, and

121.3 (West), 158.7 (Central) and 159 (East) CNLs (Table S2). As compared to the Illumina data,

PacBio assemblies resulted in more "complete", as compared to "partial" genes for both intact and

pseudogenized R genes (Figure 4B), indicating that many of the R genes predicted as pseudogenes

in the Illumina data likely represent complete genes that are mis-categorized due to assembly

limitations.

Comparing to Reference Genome

The reference genome, which was sequenced at 21.5 X coverage using PacBio long reads, contained

a greater number of R genes in total (873; Table S2), of which 281 and 147 where annotated as CNLs

and TNLs, respectively. This compares to the 603 candidates found by NLR-Annotator in the closely

related sunflower genome (Toda et al., 2020). It contained a comparable percentage of complete R

genes relative to its total number of R genes (50.7%) as the enriched PacBio libraries (58.2%, 54.2%,

and 57.3% in the West, Central, and East, respectively). 837 R genes in the reference (as well as the

281 and 147 CNLs and TNLs, respectively) exceeded counts from both the lower-coverage, R-gene-

enriched PacBio assemblies or the Illumina short-read assemblies, below, suggesting either that the

reference had more R genes (which is plausible as it an interspecific F1) than the rest of the samples

and/or that some genes were missed in the enrichment process.

R Gene Counts Increase with Increasing Climate Moisture Index

We found a strong positive association between R gene count and CMI in both the Illumina and

PacBio datasets (Figure 5A,B; Table S3) from the simple linear model. We also found a significant

positive correlation between Illumina R gene counts and CMI using the mixed effect model (MSCMI =

9493, MSerror = 1059.6, F = 8.96. Degrees of freedom from Satterthwaite's method [Satterthwaite

1946]: dfCMI = 1, dferror = 9.73, p = 0.0139, Figure 5A; Table S4), on top of substantial variation

among source prairies. Specifically, the model predicted an additional 0.707 kg/m2/month (with a

standard error of 0.236 kg/m2/month) R genes for every additional unit of CMI. As average CMI

ranged from -57.3 units to -12.9 units, the model predicts that plants from prairies with the highest

annual CMI will have 31.4 more R genes than those from prairies with the least CMI. Because most

variation in CMI is associated with source region (West, Central, or East), we also modeled source region, rather than CMI as a fixed effect. This analysis revealed a similar, but nominally insignificant association between source region and the number of R genes (MSregion = 8753, MSerror = 4376.5, F = 4.13. Degrees of freedom from Satterthwaite's method: dfregion = 2, dferror = 8.82, p = 0.0542, see descriptive statistics in the section above and in Table S4).

Our PacBio sequencing recovered more R genes per individual and had relatively fewer R genes annotated as "partial" (relative to "complete") than the Illumina data (Table S2). Despite much lower power, we still see a significant association between CMI and R gene count in the PacBio data set (MSCMI = 92006, MSerror = 9961, F = 7.193, dfCMI = 1, dferror = 7, p = 0.0315, Figure 5B). Like the case for the Illumina data modeling R gene count from the PacBio data revealed a similarly consistent but nominally insignificant association (MSregion = 53074 MSerror = 10164, F = 4.96. Degrees of freedom dfregion = 2, dferror = 6, p = 0.054, see descriptive statistics in the section above and in Table S4).

CNLs and TNLs

We did not find a significant relationship between CMI and TNL counts (p = 0.695, Figure 5D, MSregion = 14.072, MSerror = 86.559, F = 0.1626. Degrees of freedom from Satterthwaite's method: dfregion = 1, dferror = 10.215), and only a marginal significance with CNL counts (Figure 5C; p=0.0634, MSregion = 651.19, MSerror = 150.22, F = 4.3349. Degrees of freedom from Satterthwaite's method: dfregion = 1, dferror = 10.22).

Due to having only one plant sampled per prairie, we report the results of a fixed-effect linear regression model for the PacBio dataset. Despite a much lower inferential power (n=9), we still found a positive correlation between CMI and the PacBio R genes (Figure 5B; p=0.019). There was a significant relationship between CMI and PacBio TNL counts (Figure S1B; p=0.018), but not CNLs (Figure S1A; p=0.14).

Correlations Between Phenotypes and R Gene Counts

There were no significant correlations between Illumina R gene counts and leaf rust severity (Figure 6B; p=0.63), rust response (Figure 6C; p=0.73), stem rust severity (Figure 6A; p=0.51), or insect leaf herbivory (not pictured; p=0.23). We found that R gene counts were significantly associated with leaf blotch scores (Figure 6D; p=0.0043). When source prairie was modeled as a random effect, leaf blotch scores were still found to significantly correlate with CMI (p=8.94e-5, slope=-0.0082).

Principal Coordinate Correlations Among Phenotypes

The PCoA analysis (Figure S5) of the unique NBARC sequences (Supplemental file "AllNBARCs.fasta"; Table S5) against the disease phenotypes yielded no significant candidates for genes responsible for disease resistance after correcting for multiple tests.

DISCUSSION

We enriched *Silphium integrifolium* libraries for R genes using baits developed from the related *Helianthus annuus* genome to characterize the diversity across a rainfall gradient. We found that the baits recovered a substantial portion of R genes in the reference. We found geographic associations between R gene count and rainfall. Here, we discuss the *Silphium* R gene content across the rainfall gradient, as well as the performance of the RenSeq enrichment method, and the biological implications of the geographic pattern we uncovered.

DISEASE RESISTANCE AND RAINFALL

R Gene Counts Correlate with Precipitation

The significant correlation between net precipitation rate, as estimated by average CMI, and the number of R genes in a plant's genome, detected in both the Illumina short-read dataset as well as the underpowered PacBio long-read libraries (n=9), suggests evidence of possible genomic adaptation for higher pathogen load. In fact, the steeper relationship of R gene number and CMI observed with the PacBio long-read libraries suggests that the Illumina dataset underestimates the strength of this relationship. The trend does not appear to be driven by either TNLs or CNLs, for which the results conflict between the Illumina and PacBio datasets. Genes of type TNL, or those containing a toll-interleukin-like receptor in their N-terminal domain, don't differ significantly according to CMI. A stronger understanding of the relationship between climate factors and the evolution of these two subtypes of R genes remains to be investigated.

Mixed Disease Phenotype Results

Despite the significant trend of the number of R genes increasing with CMI, leaf rust severity, leaf rust response, and stem rust severity did not significantly vary in the common gardens. These three traits are different phenotypic metrics to assess a specialist fungal pathogen, *Puccinia silphii*, commonly known as *Silphium* rust. These results are consistent with results from common gardens that found that eastern populations of *S. integrifolium* were not, on average, more resistant to *Silphium* rust (Cassetta in prep.). One possible explanation for why resistance to *Silphium* rust was not associated with variable R gene counts may lie in the concept of the 'disease triangle' reviewed by Velásquez et al., 2019. The clones in this study were placed in environments where *S. integrifolium* pathogens were present, and experienced conditions that would be ideal for pathogenic growth: warm and wet. Even with effective immunity represented by the R genes sequenced in the plant populations, the level of resistance conferred by the resistance genes might have been obscured by the pathogen spread. Disease resistance in these wild populations is highly likely to be quantitative, meaning that resistance is not an all-or-nothing response, but rather a distribution of potential responses ranging from minor to major (French et al., 2016). Further studies of the wild populations with sequenced R genes, with controlled inoculations of plants with pathogens, would elucidate associations between R genes that may confer various levels of response to pathogens.

Further, the evolutionary dynamics of R genes may be muddled by seemingly competing forces
of selection, with "arms race" dynamics expected to generate gradients in total host resistance, while
oscillatory "red-queen" dynamics may not (Bergelson et al., 2002). It is possible, for example, that
local co-evolution of *S. integrifolium* and *Silphium* rust has resulted in adaptation of the rust to its local
host population, rather than variation in overall resistance. This red-queen dynamic is supported by
observations from common gardens distributed along the gradient that showed much greater virulence
of *Silphium* rust on their populations from their local region than from other regions (Cassetta et al. in
prep). In the case of *Silphium* rust, R gene identity may be more important than R gene diversity.
We did observe that resistance allele diversity was associated with suppression of leaf blotch
(Fig. 6d). Our result affirms observations of a previous common garden study that found that eastern
populations of *S. integrifolium* were more resistant to *Silphium* blotch, as well as to clear vein virus
(Cassetta et al. in prep). These results are consistent with expectations of greater diversity of
resistance alleles contributing to greater resistance to pathogens, and are qualitatively consistent with
arms-race-type dynamics. In *S. integrifolium*, leaf blotch is caused by the generalist fungal pathogen
Colletotrichum dematium (Cybernome; Farr 1989), as well as its more specialized congener,
Colletotrichum silphii Davis (Horst 2008). It is possible that arms race dynamics are more likely
between hosts and generalist pathogens such as leaf blotch, while specialist pathogens such as the
Silphium rust are more likely to generate red-queen dynamics.
We note that it is possible that the observed relationship between resistance allele diversity
and resistance to blotch is spurious, as other heritable factors that covary with rainfall might contribute
to disease resistance, such as plant secondary chemicals. We note, however, that insect herbivory
may be an indicator of such overall resistance, and we did not observe any relationship between R
gene diversity and insect leaf herbivory. This result is consistent with R genes predominantly acting as
pathogen effector receptors, and not targeted toward resistance for insects (Chovelon et al., 2021).
403

A BROADER PATTERN OF R GENE EVOLUTION
R Gene Genealogy Recovers Expected Patterns
The maximum likelihood genealogy constructed from the NBARC domains of the R genes of the draft
genomic assembly recovers the deep evolutionary divergence between CNL-type and TNL-type R
genes that has been reported in many other studies (Meyers et al., 2003; Mun et al., 2017; Neupane
et al., 2018). The instances of individual R genes nesting incongruently within the wrong clade may be
attributable to N-domain switching via recombination, which, to the best of our reckoning, has yet to be
observed. More likely, it may be due to misidentification of upstream nucleotide sequence as
incidentally translating to peptide sequence that meets the identity threshold for one of the diagnostic
motifs that places those R genes in the wrong categorization.
414

PERFORMANCE OF RENSEQ FOR R GENE ENRICHMENT
Enrichment Using Distant Reference

The R gene libraries presented in this study were developed by enriching *Silphium integrifolium* DNA

for R genes using baits developed from *Helianthus annuus*, a model organism with well-developed

genomic resources. Despite an estimated divergence age of between 13.3 and 19.3 Ma, the baits

were able to enrich libraries to contain a median of 63% of reads originating from R genes,

representing a 36-fold enrichment over WGS. For comparison, a previous study by Andolfo et al.

(2014) found success enriching the *Solanum lycopersicum* (common tomato) with baits designed from

Solanum tuberosum (common potato), a congener estimated to be 6.7 Ma divergent by TimeTree

(Kumar et al., 2017). This study demonstrates the economical promise of RenSeq for studying the

immune systems of non-model plants under a variety of ecological and evolutionary pressures. It also

showcases the screening of crop wild relatives that are of agronomic interest, such as *S. integrifolium*,

for disease resistance genes that might enable more robust response to pathogens that pose a

challenge for the domestication of the plant.

The reference genome contained a much higher number of R genes in the draft genome

assembly compared to the enriched libraries (873 compared to ~400-600), which can be accounted

for by two factors. First, the draft genome is assembled from an F1 hybrid between two different

species, *S. integrifolium* and *S. perfoliatum*. While we expect overlap in many of the R genes due to

homology, our count is likely an overestimate of the true number contained within the haploid genome

of *S. integrifolium* due to the nature of R genes as a rapidly diversifying gene family. Second, RenSeq

enrichment likely captures a different (and smaller) subset of the R genes than the WGS PacBio

sequencing we employed for the draft assembly.

Caveats and Future Directions

While this study finds a genomic signature of adaptation for environmental pathogen load, it remains

to be determined which specific genes or cassette of genes are responsible. A complete, phased,

chromosome-level assembly of *S. integrifolium* will enable association studies to determine the loci of

pathogen resistance for incorporation into breeding programs. As substantial yield and crop loss is

attributed to pathogens each year and is recognized as an issue of increasing concern for farmers and

breeders (Savary et al., 2019), generating well annotated reserves of resistant plant material and

annotated libraries of R genes for *S. integrifolium* to draw upon will be important for the near and long

term use.

While more work is needed to identify functional consequences of individual R alleles, our

observation of strong correlations of R gene diversity with precipitation is consistent with these R

alleles conferring disease resistance. These results resonate with similar results from other systems

(Rouse et al 2011, Zhu et al 2013). Our direct test of this hypothesis yielded mixed results, as R gene

diversity was associated with resistance to Silphium blotch, but not Silphium rust. Further work should

aim to disentangle the impacts of R allelic identity from R gene diversity in conferring resistance to

pathogens.

CONCLUSION

We used RenSeq to sequence the R gene libraries of *S. integrifolium* plants from across a rainfall gradient to evaluate the hypothesis that CMI imposes a biotic stress whose evolutionary pressure should be evident within the genome. We detected a significant positive relationship between CMI and R gene count, supporting our hypothesis. We demonstrate that RenSeq can be used in the evaluation of ecological hypotheses and show promise for the economical interrogation of the specific R gene loci responsible for disease resistance.

ACKNOWLEDGEMENTS

We acknowledge support for this research from NSF Dimension of Biodiversity grant 1738041. This research was supported financially by the Perennial Agriculture Project, a joint project between The Land Institute and The Malone Family Land Preservation Foundation. We gratefully acknowledge financial support from the Foundation for Food and Agriculture Research (FFAR) Fellows program. We thank Eric Cassetta for assisting with all of the field work. We are thankful to Andrew Read and Peter Innes for assistance with code to generate the phylogeny and map figures, respectively.

COI

Authors declare no conflicts of interest.

AUTHOR CONTRIBUTIONS

KK assembled and annotated the R gene libraries, performed the analyses, generated the figures, and co-wrote the manuscript. KP performed the majority of the field work, common garden experiments, and phenotyping, and co-wrote the manuscript. KT created disease scales and led the recording of disease severity in the field. AH sequenced and assembled the draft genome of *S. integrifolium*. AR designed the RenSeq baits, developed the libraries, and performed the initial steps of bioinformatics. YB, KS, JB, KT and DVT conceived of the project and contributed to the writing.

DATA ACCESSIBILITY

Illumina short-read and PacBio long-read libraries are publicly available on the SRA database at Project PRJNA914292 under the project name "RenSeq of Silphium integrifolium samples across rainfall gradient."

CITATIONS

Almeyda, C. V., Eid, S. G., Saar, D., Samuitiene, M., & Pappu, H. R. (2014). Comparative analysis of endogenous plant pararetroviruses in cultivated and wild Dahlia spp. *Virus Genes*, *48*(1), 140-152.

Andrews, S. (2010). FastQC: A Quality Control Tool for High Throughput Sequence Data [Online]. Available online at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc/

Bankevich, A., Nurk, S., Antipov, D., Gurevich, A. A., Dvorkin, M., Kulikov, A. S., Lesin, V. M., Nikolenko, S. I., Pham, S., Prjibelski, A. D., Pyshkin, A. V., Sirotkin, A. V., Vyahhi, N., Tesler, G., Alekseyev, M. A., Pevzner, P. A. (2012). SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *Journal of Computational Biology*, 19(5), 455-477.

Barreda, V. D., Palazzesi, L., Tellería, M. C., Olivero, E. B., Raine, J. I., & Forest, F. (2015). Early evolution of the angiosperm clade Asteraceae in the Cretaceous of Antarctica. *Proceedings of the National Academy of Sciences*, *112*(35), 10989-10994.

Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B (Methodological)*, *57*(1), 289-300.

Bergelson, J., Kreitman, M., Stahl, E. A., and Tian, D. (2002). Evolutionary Dynamics of Plant R-Genes. Science, 292 (July), 2281–2289. https://doi.org/10.1126/science.1061337

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*, 30(15), 2114-2120.

Brun, P., Zimmermann, N. E., Hari, C., Pellissier, L., & Karger, D. N. (2022). Global climate-related predictors at kilometre resolution for the past and future. *Earth System Science Data Discussions*, 1-44.

Caplan, J., Padmanabhan, M., & Dinesh-Kumar, S. P. (2008). Plant NB-LRR immune receptors: from recognition to transcriptional reprogramming. *Cell Host & Microbe*, *3*(3), 126-135.

Cheng, H., Concepcion, G. T., Feng, X., Zhang, H., & Li, H. (2021). Haplotype-resolved de novo assembly using phased assembly graphs with hifiasm. *Nature Methods*, *18*(2), 170-175.

525 Chovelon, V., Feriche-Linares, R., Barreau, G., Chadoeuf, J., Callot, C., Gautier, V., Le Paslier, M. C.,
526 Berad, A., Faivre-Rampant, P., Lagnel, J., Boissot, N. (2021). Building a cluster of NLR genes
527 conferring resistance to pests and pathogens: the story of the Vat gene cluster in cucurbits.
528 *Horticulture Research*, *8*.
529

530 Clarkson, J. P., Fawcett, L., Anthony, S. G., & Young, C. (2014). A model for Sclerotinia sclerotiorum
531 infection and disease development in lettuce, based on the effects of temperature, relative humidity
532 and ascospore density. *PLoS One*, *9*(4), e94049.
533

534 Dabo, S. M., Mitchell, E. D., & Melcher, U. (1993). A method for the isolation of nuclear DNA from
535 cotton (Gossypium) leaves. *Analytical Biochemistry*, *210*(1), 34-38.
536

537 Delavaux, C. S., Schemanski, J. L., House, G. L., Tipton, A. G., Sikes, B., & Bever, J. D. (2021). Root
538 pathogen diversity and composition varies with climate in undisturbed grasslands, but less so in
539 anthropogenically disturbed grasslands. *The ISME journal*, *15*(1), 304-317.
540

541 Eitas, T. K., & Dangl, J. L. (2010). NB-LRR proteins: pairs, pieces, perception, partners, and pathways.
542 *Current Opinion in Plant Biology*, *13*(4), 472-477.
543

544 Farr, D. F., Bills, G. F., Chamuris, G. P., & Rossman, A. Y. (1989). *Fungi on Plants and Plant Products*
545 *in the United States*. APS press.
546

547 French, E., Kim, B. S., & Iyer-Pascuzzi, A. S. (2016). Mechanisms of quantitative disease resistance in
548 plants. In *Seminars in Cell & Developmental Biology* (Vol. 56, pp. 201-208). Academic Press.
549

550 Glover, J. D., Reganold, J. P., Bell, L. W., Borevitz J., Brummer, E. C., Buckler, E.S., Cox, C. M., Cox.,
551 T. S., Crews, T., E., Culman, W., Dehaan, R., Eriksson D., Gill, B., S., Holland, J., Hu, F., Hulke B., S.,
552 Ibrahim, A. M. H., Jackson W., Jones, S. S., Murray, S., C., Paterson, A. H., Ploschuk E., Sacks, E. J.,
553 Snapp, S., Tao, D., Van Tassel, D. L., Wade L. J., Wysean D. L., Xu, D. Y. (2010). Increased food and
554 ecosystem security via perennial grains. *Science*, *328*(5986), 1638-1639.
555

556 Granke, L. L., & Hausbeck, M. K. (2010). Effects of temperature, humidity, and wounding on
557 development of Phytophthora rot of cucumber fruit. *Plant Disease*, *94*(12), 1417-1424.
558

559 Horst, R. Westcott's Plant Disease Handbook; Springer: New York, NY, USA, 2008.
560

561 Islam, T. M., & Toyota, K. (2004). Effect of moisture conditions and pre-incubation at low temperature
562 on bacterial wilt of tomato caused by Ralstonia solanacearum. *Microbes and Environments*, *19*(3),
563 244-247.
564
565 Jupe, F., Pritchard, L., Etherington, G. J., MacKenzie, K., Cock, P. J., Wright, F., Sharma, S. K., Bolser,
566 D., Bryan, G. J., Jones, J. D. G., Hein, I. (2012). Identification and localisation of the NB-LRR gene
567 family within the potato genome. *BMC Genomics*, *13*(1), 1-14.
568
569 Jupe, F., Witek, K., Verweij, W., Śliwka, J., Pritchard, L., Etherington, G. J., Maclean, D., Cock, P. J.,
570 Leggett R. M., Bryan G. J., Cardle, L., Hein, I., Jones, J. D. (2013). Resistance gene enrichment
571 sequencing (RenSeq) enables reannotation of the NB-LRR gene family from sequenced plant
572 genomes and rapid mapping of resistance loci in segregating populations. *The Plant Journal*, *76*(3),
573 530-544.
574
575 Kumar, S., Stecher, G., Suleski, M., & Hedges, S. B. (2017). TimeTree: a resource for timelines,
576 timetrees, and divergence times. *Molecular biology and evolution*, *34*(7), 1812-1819.
577
578 Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv*
579 *preprint arXiv:1303.3997*.
580
581 Magarey, R. D., Sutton, T. B., & Thayer, C. L. (2005). A simple generic infection model for foliar fungal
582 plant pathogens. *Phytopathology*, *95*(1), 92-100.
583
584 Mccann, J., Jang, T. S., Macas, J., Schneeweiss, G. M., Matzke, N. J., Novák, P., Stuessy, T. F.,
585 Villaseñor, J. L., Weiss-Schneeweiss, H. (2018). Dating the species network: allopolyploidy and
586 repetitive DNA evolution in American daisies (Melampodium sect. Melampodium, Asteraceae).
587 *Systematic Biology*, *67*(6), 1010-1024.
588
589 Meyers, B. C., Dickerman, A. W., Michelmore, R. W., Sivaramakrishnan, S., Sobral, B. W., & Young, N.
590 D. (1999). Plant disease resistance genes encode members of an ancient and diverse protein family
591 within the nucleotide-binding superfamily. *The Plant Journal*, *20*(3), 317-332.
592
593 Meyers, B. C., Kozik, A., Griego, A., Kuang, H., & Michelmore, R. W. (2003). Genome-wide analysis of
594 NBS-LRR–encoding genes in Arabidopsis. *The Plant Cell*, *15*(4), 809-834.
595

596 Mun, J. H., Yu, H. J., Park, S., & Park, B. S. (2009). Genome-wide identification of NBS-encoding
597 resistance genes in Brassica rapa. *Molecular Genetics and Genomics*, *282*(6), 617-631.
598

599 Neupane, S., Andersen, E. J., Neupane, A., & Nepal, M. P. (2018). Genome-wide identification of
600 NBS-encoding resistance genes in sunflower (Helianthus annuus L.). *Genes*, *9*(8), 384.
601

602 Oksanen, J. (2015). Vegan: an introduction to ordination. *R Project*,
603 https://cran.r-project.org/web/packages/vegan/vignettes/intro-vegan.pdf.
604

605 Reinert, S., Money, K. L., Rockstad, G. B., Kane, N. C., Van Tassel, D. L., & Hulke, B. S. (2018). Two
606 contrasting laboratory methods improve Silphium integrifolium Michx. germination rate to
607 agronomically acceptable levels. *Euphytica*, *214*(9), 1-9.
608

609 Rouse, M. N., Saleh, A. A., Seck, A., Keeler, K. H., Travers, S. E., Hulbert, S. H., & Garrett, K. A.
610 (2011). Genomic and resistance gene homolog diversity of the dominant tallgrass prairie species
611 across the US Great Plains precipitation gradient. *PloS One*, *6*(4), e17641.
612

613 Rowlandson, T., Gleason, M., Sentelhas, P., Gillespie, T., Thomas, C., & Hornbuckle, B. (2015).
614 Reconsidering leaf wetness duration determination for plant disease management. *Plant Disease*,
615 *99*(3), 310-319.
616

617 Satterthwaite, F. E. (1946). An approximate distribution of estimates of variance components.
618 *Biometrics Bulletin*, *2*(6), 110-114.
619

620 Savary, S., Willocquet, L., Pethybridge, S. J., Esker, P., McRoberts, N., & Nelson, A. (2019). The
621 global burden of pathogens and pests on major food crops. *Nature Ecology & Evolution*, *3*(3), 430-
622 439.
623

624 Stamatakis, A. (2006). RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with
625 thousands of taxa and mixed models. *Bioinformatics*, *22*(21), 2688-2690.
626

627 Stamatakis, A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
628 phylogenies. *Bioinformatics*, *30*(9), 1312-1313.
629

Steuernagel, B., Witek, K., Krattinger, S. G., Ramirez-Gonzalez, R. H., Schoonbeek, H. J., Yu, G.,
Baggs, E., Witek, A. I., Yadav, I., Krasileva, K., V., Jones, J. D. G., Uauy, C., Keller, B., Ridout, C. J.,

Wulff, B. B. (2020). The NLR-Annotator tool enables annotation of the intracellular immune receptor
repertoire. *Plant Physiology*, 183(2), 468-482.

Toda, N., Rustenholz, C., Baud, A., Le Paslier, M. C., Amselem, J., Merdinoglu, D., & Faivre-Rampant,
P. (2020). NLGenomeSweeper: a tool for genome-wide NBS-LRR resistance gene identification.
Genes, *11*(3), 333.

Turner, M. K., Ravetta, D., & Van Tassel, D. (2018). Effect of Puccinia silphii on yield components and
leaf physiology in Silphium integrifolium: Lessons for the domestication of a perennial oilseed crop.
Sustainability, *10*(3), 696.

Van Tassel, D. L., Albrecht, K. A., Bever, J. D., Boe, A. A., Brandvain, Y., Crews, T. E., Gansberger, M.,
Gerstberger, P., González-Paleo, L., Hulke, B. S., Kane, N. C., Johnson, P. J., Pestsova, E. G.,
Picasso-Risso, V. D., Prasifka, J. R., Ravetta, D. A., Schlautman, B., Sheaffer, C. C., Smith, K. P.,
Speranza, P. R., Turner., M. K., Vilela, A. E., von Gehren, P., Wever, C. (2017). Accelerating silphium
domestication: an opportunity to develop new crop ideotypes and breeding strategies informed by
multiple disciplines. *Crop Science*, *57*(3), 1274-1284.

Velásquez, A. C., Castroverde, C. D. M., & He, S. Y. (2018). Plant–pathogen warfare under changing
climate conditions. *Current Biology*, *28*(10), R619-R634.

Vilela, A., González-Paleo, L., Turner, K., Peterson, K., Ravetta, D., Crews, T. E., & Van Tassel, D.
(2018). Progress and bottlenecks in the early domestication of the perennial oilseed Silphium
integrifolium, a sunflower substitute. *Sustainability*, *10*(3), 638.

Wieczorek, M. E., Jackson, S. E., & Schwarz, G. 2. (2018). Select attributes for NHDPlus version 2.1
reach catchments and modified network routed upstream watersheds for the conterminous United
States. *US Geological Survey. https://doi. org/10.5066/F7765D7V*.

Zhu, Q., Bennetzen, J. L., & Smith, S. M. (2013). Isolation and diversity analysis of resistance gene
homologues from switchgrass. *G3: Genes, Genomes, Genetics*, *3*(6), 1031-1042.

Figure 1: Prairie locations. Western prairies were distributed across Central Kansas. Central prairies
were located in Eastern Kansas, which has a much wetter climate than Central Kansas. Eastern
prairies were in Illinois, which has a similar level of precipitation to Eastern Kansas, but has higher
Climate Moisture Index (precipitation minus evapotranspiration), which allows more water to be
utilized by plants. Circles are filled according to the region-wise color scheme used throughout this
manuscript.

Figure 2: ML tree constructed from the peptide sequence of the nucleotide-binding domain (NB-ARC)
of 590 R genes in the genome of the draft *S. integrifolium* genome. Color bar represents the
probability that a sequence belongs to a TNL (green) or not (red).

Figure 3: All R genes among all samples in the dataset were concatenated into a single fasta file (Pan-
NLRome). The raw libraries from each sample were mapped to the Pan-NLRome. The ratios of the
number of mapped reads to the total read count in each library was calculated. A median of 63% of
reads in the enriched libraries were putatively sourced from R genes and 1.8% of reads in the un-
enriched WGS libraries. Note the log scale of the x-axis.

Figure 4: (A) Breakdown of R genes annotated by whether they contained a coiled-coil N-terminal
domain, a toll-interleukin-like N-terminal domain, or neither. Those that were not identifiable either
lacked a diagnostic N-terminal domain by their nature, or due to incomplete assembly. (B) R genes
broken down by their completeness and inferred functionality. Genes labeled "pseudo" are likely
pseudogenes due to the presence of internal stop codons. The higher number of "Complete" genes in
the PacBio assemblies compared to the Illumina libraries are likely due to more reliable assembly from
the long reads.

Figure 5: (A) Total R gene counts recovered from Illumina short-read libraries plotted against the
yearly CMI, or the amount of precipitation that is stored in the soil and not lost to evaporation. (B) Total
R gene counts recovered from the PacBio long-read libraries. (C) Counts of R genes from the short-
read libraries that were identified as type coiled-coil (CNLs). (D) Counts of R genes from the short-
read libraries of type toll-interleukin-like receptor (TNLs).

Table 1: Location information of the twelve pristine prairie remnants from which the *S. integrifolium*
plants were sourced. Climate Moisture Index is calculated as the difference between precipitation and
potential evapotranspiration, which is an indicator for the availability of moisture for plants to uptake.

Source	Prairie	Site Name	Latitude	Longitude	Climate Moisture Index (g/m²/month)
W1		W1	38.9908	-89.2300	-35.2
W2		W2	38.9908	-89.2300	-35.2
W3		W3	38.6316	-89.2866	-39.4
W4		W4	38.6366	-87.6729	-18.6
C1		Prairie Park	38.9031	-95.2122	-24.1
C2		Clog Prairie	38.7880	-95.2099	-25.1
C3		Kill Creek	38.9173	-94.5745	-24.4
C4		Lexington Lake	38.9664	-95.0102	-26.6
E1		Pellsville	40.4701	-87.5242	-12.8
E2		Prospect	40.4451	-86.8903	-12.6
E3		Loda	40.5273	-88.0765	-14.4
E4		Weston	40.7467	-88.6143	-20.4