14   **Disease resistance gene count increases with rainfall in _Silphium integrifolium_**
15

16   Kyle Keepers[1], Kelsey Peterson[1], Andrew Raduski[1†], Kathryn M. Turner[2], David Van Tassel[2], Kevin
17   Smith[3], Alex Harkess[4], James D. Bever[5], Yaniv Brandvain[1]
18
[1]University of Minnesota, Department of Plant and Microbial Biology, 1500 Gortner Ave,
St Paul, MN 55108
[2]The Land Institute, Crop Protection Ecology, 2440 E Water Well Rd, Salina, KS 67401
[3]University of Minnesota, Department of Agronomy and Plant Genetics, 1529 Lindig St,
St Paul, MN 55108
[4]HudsonAlpha Institute for Biotechnology, 601 Genome Way, Huntsville, AL 35806
[5]University of Kansas, Department of Ecology and Evolutionary Biology, 1200 Sunnyside
Ave, Lawrence, KS 66045

27
28

ABSTRACT

Intracellular plant defense against pathogens is mediated by a class of disease resistance

genes known as NB-LRRs or NLRs (R genes). Many of the diseases these genes protect

against are more prevalent in regions of higher rainfall, which provide better growth conditions

for the pathogens. As such, we expect a higher selective pressure for the maintenance and

proliferation of R genes in plants adapted to wetter conditions. In this study, we enriched

libraries for R genes using RenSeq from baits primarily developed from the common sunflower

(*Helianthus annuus*) reference genome. We sequenced the R gene libraries of *Silphium*

integrifolium Michx, a perennial relative of sunflower, from 12 prairie remnants across a rainfall

gradient in the Central Plains of the United States, with both Illumina short-read (n=99) and

PacBio long-read (n=10) approaches. We found a positive relationship between the mean

effective annual precipitation of a plant's source prairie remnant and the number of R genes in

its genome, consistent with intensity of plant pathogen coevolution increasing with precipitation.

We show that RenSeq can be applied to the study of ecological hypotheses in non-model

relatives of model organisms.

Keywords: *Silphium integrifolium*, R genes, RenSeq, common gardens, host-pathogen

dynamics, prairies

INTRODUCTION

Silphium integrifolium Michx (commonly known as rosinweed or silflower) is a native perennial

prairie species being domesticated for production as a forage and oilseed species (Vilela et al.,

2018; Van Tassel et al., 2017). The plant's perennial nature allows it to provide novel ecosystem

services in agronomic production systems (Glover et al., 2010; Van Tassel et al., 2017) while

producing oilseeds with similar nutritional composition to annual sunflower (*Helianthus annuus*

L. (Asterales: Asteraceae) seeds (Reinert et al., 2018). However, *S. integrifolium* suffers

infections from diverse pathogen enemies across its native range. The stems, leaves, and

flowers can be infected with multiple strains (Turner, unpublished observations) of *Puccinia*

silphii Schwein (1832), or *Silphium* rust (Turner et al., 2018), *Colletotrichum silphii* Davis (1919)

(Horst 2008) syn. and *Colletotrichum dematium* (Pers.) Grove (Cybernome; Farr 1989) called

leaf blotch hereafter. Recently, *Silphium* clear vein (SCV) leaf and stem disease has been

described and hypothesized to be caused by a virus based on the identification of viral

sequences in *Silphium integrifolium* that are similar to the Dahlia common Mosaic Virus (DCMV)

and Dahlia endogenous plant pararetroviral sequence (DvEPRS, formerly DMV-D10) (Almeyda,

2014; Cassetta et al., 2023). Understanding how *S. integrifolium* defends against these

pathogens is crucial for progress in breeding programs and will elucidate evolutionary questions

posed in the species (Cassetta et al., 2023, Van Tassel et al, 2017). In this study, plants from 12

wild populations of *S. integrifolium* were planted in a reciprocal transplant design to better

understand adaptation to disease pressures along a rainfall gradient. Phenotypic data revealed

that these wild populations have distinctive morphologies and showed variable levels of

tolerance to biotic stresses, such as pests and pathogens (Peterson et al., in prep). Individuals

with resistance to diseases of *S. integrifolium* were visually identified in the study, and have

been previously identified in breeding populations (Turner et al., 2018), but the genetic basis of

disease resistance has not been investigated. Here, we look at the diversity of disease

resistance genes in wild populations of *S. integrifolium* to answer evolutionary questions

regarding adaptation of wild populations to their pathogen pressures and suggest applications

for germplasm enhancement.

Environments with high precipitation are conducive to increased pathogen pressures

(Clarkson et al., 2014; Granke and Hausbeck 2010; Islam and Toyota, 2004; Magarey et al.,

2005; Rowlandson et al., 2015). Several studies have found patterns of resistance alleles

consistent with pathogen selection intensity increasing with precipitation (Wahl 1970, Abbott,

Brown, and Burdon 1991, Dong et al. 2009). We hypothesize that *S. integrifolium* populations

evolved in areas with high precipitation will have a more diverse complement of disease

resistance genes to deal with this challenge. To test this hypothesis, we collected seeds from *S.*

integrifolium stands from four prairie remnants each in three geographic regions in the Central

Plains of the United States. The regions are situated along a gradient of effective precipitation,

or the amount of rainfall that is not lost to potential evapotranspiration, referred to in this study

as the climate moisture index (CMI). We refer to these regions as "West", from West-Central

Kansas, "Central", from Eastern Kansas, and "East", from Central Illinois (Figure 1). Prairie

remnants in the West receive around -55 kg/m^2/month of CMI per year, whereas sites in the

East receive closer to -15 kg/m^2/month (Table 1). We note that plant defensive gene diversity

increased along this gradient in *Andropogon gerardii*, a native perennial grass that co-occurs

with *S. integrifolium* (Rouse et al. 2011) and soil pathogen diversity also increased along this

same gradient (Delavaux et al. 2021).

A major class of disease resistance genes in plants are the cytoplasmic nucleotide

binding and leucine-rich repeat (NLR) genes. (Caplan et al., 2008; Eitas and Dangl, 2010).

These NLRs, (hereafter, R genes), often contain an N-terminal domain that is either of a coiled-

coil type, called CNLs, or a toll-interleukin-like type (TNLs), whose divergence predates the split

between monocots and eudicots (Meyers et al., 1999), estimated to be about 160 Mya by

TimeTree (Kumar et al., 2017). R genes are highly diverse, numbering in the hundreds in many

plant species (Jupe et al., 2012; Toda et al., 2020). These genes evolve rapidly and are

enriched in presence/absence variation even at the intraspecies level, limiting the utility of a

single reference genome to elucidate the diversity of R genes within a species. This study

employed R-gene enrichment sequencing (RenSeq; Jupe et al, 2013), which uses DNA baits to

reduce complexity of genomic DNA libraries. The baits we designed come from the common

sunflower, *Helianthus annuus* L., which is separated from *S. integrifolium* by between 22

(Meireles et al., 2020) and 33 million years ago (Zhang et al., 2021) . We sequenced the

RenSeq libraries from *S. integrifolium* plants across the rainfall gradient using both Illumina

short-read and PacBio long-read technologies.

108   RenSeq is not a new technology – having been developed in 2013. It has been used to

address basic problems in plant immunity (Jupe et al., 2013) and applied questions in plant

breeding in numerous crop species and their wild relatives for crop improvement (e.g. Arora et

al., 2019) and to understand mechanisms of plant immunity (e.g. Witek et al., 2021). Despite the

utility of RenSeq in plant breeding projects, and in helping us better understand plant disease

resistance, this technology has not been used to address basic evolutionary and

biogeographical questions.

115   Using both simple linear models as well as mixed effect models, we found a significant

positive correlation between the effective precipitation of a plant's host prairie and the number of

R genes detected in both the Illumina and PacBio datasets. While this observation does not

itself prove the broader theory that disease resistance correlates with rainfall in plants, it is

certainly consistent with theoretical expectations, and should motivate additional evaluation of

these ideas in other taxa. Because we demonstrate the economic utility of the RenSeq

approach in non-model systems, our work not only points to data consistent with our hypothesis

but paves the way towards evaluating the generality of this finding.

METHODS

Plant Materials

Silphium integrifolium seeds were collected from 12 prairies in 2017 and 2018 (Figure 1; Table

1). The prairie remnants collected spanned from central Kansas to central Illinois. The prairie

remnants have variable historical, geographic, and climatic characteristics described in Cassetta

et al 2023. From each of the 12 prairies, seeds were collected from approximately 40 mother

plants randomly chosen from throughout the prairie remnant. As *S. integrifolium* is an obligate

outcrosser, seeds collected from a single mother were a half sibling family.

S. integrifolium seeds were germinated at room temperature together in a heated

greenhouse at The Land Institute, Salina KS in 2019. A total of 96 genets comprising eight

individuals from each prairie remnant from different mothers were selected from the seedlings.

Selection of the 96 individuals was randomized although the seedlings chosen had to be

moderately vigorous. The pots containing the seedlings were randomized and placed on

greenhouse benches. Pots were re-randomized several times during the season. The plants

grew rapidly but remained as vegetative rosettes. Each crown (genet) was divided into 4 pieces

(ramets) with approximately equal numbers of dormant buds and roots, although a few crowns

were too small and only produced 2-3 pieces. The ramets were repotted into 10-gallon fabric

planters (Smart Pot) using Pro-Mix BX.

Tissue Collection

One ramet from each genet was grown in The Land Institute Greenhouse until young leaves

had emerged. On March 12, 2020, black plastic pots were placed over one clone of each of the

99 plants emerging rosette and left for 48 hours to slightly etiolate the young leaves, which

improves the quality of extracted DNA in photosynthetic active tissue containing phenolic

compounds (Dabe et al., 1993). On March 14, 3-5 unexpanded leaves were excised from each

crown. The collected tissue was placed in a labeled envelope and immediately placed on dry ice

in a cooler. Within 60 minutes of collection, the sampled leaf tissue was placed in a lyophilizer

and lyophilized for 48 hours.

Clone placement across the gradient and phenotyping

After young leaf tissue was collected from plants in the greenhouse, the plants remained in the

greenhouse under ideal conditions allowing them to grow rapidly. One cloned ramet from each

of the available genets was then moved to each of the following locations within the regions the

seed was sourced from: 1. breeding plots in Salina, KS at The Land Institute (West), 2. Prairie

Park Nature Center, Lawrence, KS (Central), 3. research plots at the ISU horticulture center in

Normal, Illinois (East), and 4. Land Institute Crop Protection Ecology greenhouse (backup site to

conserve genets in a disease and insect free state). The pots remained undisturbed at each site

for two years, 2020 and 2021 except the Central site cloned plants, which, in May 2021, were

moved from the Prairie Park Nature Center to the Perennial Agriculture Project field station,

approximately, 6 miles away. The clones received occasional watering in each site to promote

disease growth and avoid plant death due to the above ground pots drying out.

In 2020 and 2021, disease scores and insect herbivory damage were recorded on each

of the clones in all 3 sites. Data was collected during their flowering period, between the months

of July and September. Diseases were recorded according to methods in Cassetta et al., 2023.

Briefly, leaf rust severity was quantified by visually assessing the total percentage of leaf area

affected (0-100%) using the rust scale described by Turner, et al. (2018). Similarly, stem rust

severity was recorded as the total percentage of the stems covered in rust (0-100%). Leaf

blotch was quantified by assessing the total percentage of leaf area affected (0-100%). Silphium

clear vein was rated on a 0-10 scale based on symptoms observed on the entire plant including:

leaf distortion (narrow width, curled, toothed, twisted, or pinched tips), and twisted meristem

symptoms. Leaf herbivory scores were collected by quantifying the percentage of total leaves

with evidence of insect herbivore (mining, sucking, chewing) damage (0-100%).

Climate Moisture Index

A BIOCLIM+ raster map of the continental U.S.A. containing 30-year climate moisture index

averages (1981-2010) was downloaded from the CHELSA (Climatologies at high resolution for

the Earth's land surface areas) database (Brun et al., 2022). The climate moisture index is

calculated as the difference between mean annual precipitation and mean potential

evapotranspiration and, when negative, represents the availability of water as a limiting

resource in dry regions such as prairies.

Sequencing/RenSeq/QC

DNA was extracted from lyophilized leaf tissue using the CTAB method by the University of

Minnesota Genomics Center. DNA was then shipped on dry ice to Arbor Biosciences for

sequencing, library preparation, and RenSeq data acquisition. Extracted DNA was prepared for

sequencing using Illumina library prep kits. Sequencing libraries were enriched for R genes

using the RenSeq protocol described in Jupe et al., 2013 using Arbor's myBaits systems. In

short, biotinylated single-stranded RNA oligo baits were designed based on annotated R genes

from *Helianthus annuus*, a relative of *Silphium integrifolium*, within the same subtribe, estimated

to be between 22.5 (Meireles et al., 2020) and 33.5 million years divergent (Zhang et al., 2022).

Hybridized oligos were pulled down using streptavidin-coated magnetic beads. Libraries were

then sequenced on an Illumina NovaSeq S4 sequencer using paired-end 150bp reads by Arbor

Biotechnologies in Cambridge, MA.

A subsample of 10 individuals was selected for a complementary R gene analysis using

Pacific Biosciences (PacBio) long-read sequencing. High-molecular-weight enriched DNA from

three individuals from each of the West, Central, and East prairies, as well as one individual

derived from a cultivated lineage, were prepared for long read sequencing and then sequenced

on a Pacific Biosciences Sequel II using circular consensus sequencing (CCS) by Arbor

Biotechnologies.

A total of 99 Illumina libraries (96 from prairie remnants + 3 cultivated "elite" lineages)

and 10 PacBio libraries were sequenced. Both library types were filtered for sequence quality

using Trimmomatic v0.39 (Bolger et al., 2014) and then assessed using fastqc v0.11.7

(Andrews 2010). The Illumina libraries were subsampled to a standardized assembly input size

of 1Gb to avoid biasing R gene counts by library input size. The PacBio libraries were not

subsampled to a standard input size, but a correlation between input size and total R gene

count in this data type reveals no dependence of final assembly size on initial input size (Figure

S3; p=0.414). Six Illumina libraries failed to meet the input threshold and were excluded from

analysis. The libraries that met the input threshold were then *de novo* assembled using SPAdes

v3.15.3 (Bankevich et al., 2012) with -k 21,33,45,55,65,81.

Draft Genome Assembly

A chromosome-scale genomic assembly of an F1 *Silphium integrifolium* x *S. perfoliatum* is

being prepared at the time of writing. The *S. integrifolium* mother was a leaf blotch resistant,

large-headed selection from a breeding population that had undergone approximately 6 cycles

of recurrent selection for increased number of fertile ray florets ("feminization") per head and

disease resistance. The founders of the breeding population were sourced from wild central

Kansas populations, i.e., located within the geographic region we call "West" in this study.

Flash-frozen tissue was sent to Arizona Genomics Institute for high-molecular-weight DNA

extraction, which was then prepared for PacBio Sequel-II CCS sequencing at the Genome

Sequencing Center at HudsonAlpha Institute for Biotechnology. The libraries were sequenced at

roughly 21.5X depth assuming a 10 Gb genome size. Reads were assembled *de novo* using

hifiasm assembler (Cheng et al., 2021) using default parameters.

R Gene Annotations

We annotated the Illumina RenSeq libraries, PacBio RenSeq libraries, and the draft genome for

their R gene content. Each assembly was annotated for R genes using NLR-Annotator

(Steuernagel et al., 2020), which searches for common R gene peptide motifs in protein-coding

sequence. The annotations were performed with the "-a" flag, which outputs the amino acid

sequence of the nucleotide-binding domain (NBARC domain). We used the NBARC sequences

of the draft genomic assembly to infer a maximum likelihood tree in RAxML v.8.2.11

(Stamatakis 2006; Stamatakis 2014) using the peptide substitution model

"PROTGAMMAAUTO" with 100 rapid bootstraps.

The two broad, non-overlapping classes of R genes that we analyzed in this study, TNLs

(containing a toll-interleukin-like type N-terminal domain) and CNLs (containing a coiled-coil

type N-terminal domain), were identified from the annotations based on diagnostic motifs. To

identify TNLs, we parsed annotations for the motifs 13, 15, and 18 (as defined in Table S1 of

Jupe et al., 2013). For CNLs, we searched for motifs 2, 6, 16 and 17. These two sets of motifs

occur only in their respective class of N-terminal domain.

Principal Coordinates Analysis

To test for associations between R genes and the phenotypes collected on the cloned

plants across the gradient, we performed simple linear modeling as well as mixed effect

modeling, which treats CMI as a fixed effect, and source prairie as a random effect. We found

one significant correlation between R Genes of the various populations and leaf blotch. To

interrogate which R genes were responsible for the association between leaf blotch and R gene

counts, we performed a census of which R gene NBARC sequences occurred in which

individuals (Table S5). The census was then used to generate a principal coordinate analysis in

the R package "vegan" (Oksanen 2015) against each of the five phenotypic covariates included

in the study. P-values were corrected for multiple tests using the Benjamini & Hochberg method

(1995).

Library Enrichment Calculations

The fasta files containing the annotated R genes for each Illumina library were concatenated

into a fasta containing every observed R gene from the Illumina dataset (hence, a "Pan-

NLRome"). Reads from each Illumina RenSeq library, as well as eleven 1X-depth WGS libraries

from seven *S. integrifolium*, two *S. perfoliatum*, one *S. laciniatum* and one *S. terebinthinaceum*,

were mapped to the Pan-NLRome using *bwa mem* (Li 2013). The fraction out of the total reads

in the library that mapped to the Pan-NLRome with a SAM CIGAR mapping score of between

148M-151M was calculated, representing reads that mapped to an R gene in the Pan-NLRome

with between 98.01%-100% identity.

RESULTS

Reference Genome Assembly

The draft genome used in this study was 9.9 Gb in length and totalled 2,584 contigs. The

longest contig was 182 Mb long and the mean contig size was 3.8 Mb. The L50 (which is the

number of contigs, ordered longest-to-smallest, to comprise 50% of the assembly size) was 77,

and the N50 (the size of the contig that passes the 50% assembly size threshold) was 40.2 Mb.

The maximum likelihood tree constructed from the R genes in the draft assembly was

annotated according to peptide motifs that are diagnostic for either CNL or TNL identity (Figure

2). A deep evolutionary divergence between the two broad classes is seen, although we

observe isolated instances of R genes falling incongruently within the wrong clade.

Library Enrichment

The libraries of seven *S. integrifolium* that were sequenced using WGS, and thus were

unenriched for R genes, contained a median of 1.78% of reads that mapped with high fidelity

(98% identity or higher) to contigs in the Pan-NLRome. Libraries that had been enriched for R

genes using oligo baits designed from *Helianthus annuus* contained a median of 63.4% of high

fidelity reads mapping to R genes, representing a 36-fold enrichment for R genes (Figure 3).

R Gene Counts by Geographic Region

The Illumina R gene assemblies ranged from 2.2 Mb to 51 Mb, and averaged 15 Mb in size. We

detected an average of 386.3, 410.0, and 414.0 R genes in plants sourced from the Western,

Central, and Eastern prairies, respectively (left facet of Figure 4A). The plants averaged 47.38

(West), 48.69 (Central), and 45.14 (East) TNL genes, and 166.1 (West), 171.3 (Central), and

175.3 (East) CNL genes (Table S2).

The PacBio R gene assemblies ranged from 4.9 Mb to 13 Mb, but averaged 8 Mb.

Within this dataset, we only had three plants per region, so the inferential power was lower, but

our confidence in the contigs assembled was higher due to longer reads. We found an average

of 355.0, 563.3, and 592.7 R genes in the Western, Central, and Eastern prairies, respectively

(right facet of Figure 4A). Within this dataset, we found an average of 54.00 (West), 81.67

(Central), and 81.33 (East) TNLs, and 121.3 (West), 158.7 (Central) and 159 (East) CNLs

(Table S2). As compared to the Illumina data, PacBio assemblies resulted in more "complete",

as compared to "partial" genes for both intact and pseudogenized R genes (Figure 4B),

indicating that many of the R genes predicted as pseudogenes in the Illumina data likely

represent complete genes that are mis-categorized due to assembly limitations.

Comparing to Reference Genome

The reference genome, which was sequenced at 21.5 X coverage using PacBio long reads,

contained a greater number of R genes in total (873; Table S2), of which 281 and 147 where

annotated as CNLs and TNLs, respectively. This compares to the 603 candidates found by

NLR-Annotator in the closely related sunflower genome (Toda et al., 2020). It contained a

comparable percentage of complete R genes relative to its total number of R genes (50.7%) as

the enriched PacBio libraries (58.2%, 54.2%, and 57.3% in the West, Central, and East,

respectively). 837 R genes in the reference (as well as the 281 and 147 CNLs and TNLs,

respectively) exceeded counts from both the lower-coverage, R-gene-enriched PacBio

assemblies or the Illumina short-read assemblies, below, suggesting either that the reference

had more R genes (which is plausible as it an interspecific F1) than the rest of the samples

and/or that some genes were missed in the enrichment process.

R Gene Counts Increase with Increasing Climate Moisture Index

We found a strong positive association between R gene count and CMI in both the Illumina and

PacBio datasets (Figure 5A,B; Table S3) from the simple linear model. We also found a

significant positive correlation between Illumina R gene counts and CMI using the mixed effect

model (MS_{CMI} = 9493, MS_{error} = 1059.6, F = 8.96. Degrees of freedom from Satterthwaite's

method [Satterthwaite 1946]: df_{CMI} = 1, df_{error} = 9.73, p = 0.0139, Figure 5A; Table S4), on top of

substantial variation among source prairies. Specifically, the model predicted an additional

0.707 kg/m^2/month (with a standard error of 0.236 kg/m^2/month) R genes for every additional

unit of CMI. As average CMI ranged from -57.3 units to -12.9 units, the model predicts that

plants from prairies with the highest annual CMI will have 31.4 more R genes than those from

prairies with the least CMI. Because most variation in CMI is associated with source region

(West, Central, or East), we also modeled source region, rather than CMI as a fixed effect. This

analysis revealed a similar, but nominally insignificant association between source region and

the number of R genes (MS_{region} = 8753, MS_{error} = 4376.5, F = 4.13. Degrees of freedom from

Satterthwaite's method: df_{region} = 2, df_{error} = 8.82, p = 0.0542, see descriptive statistics in the

section above and in Table S4).

Our PacBio sequencing recovered more R genes per individual and had relatively fewer

R genes annotated as "partial" (relative to "complete") than the Illumina data (Table S2). Despite

much lower power, we still see a significant association between CMI and R gene count in the

PacBio data set (MS_{CMI} = 92006, MS_{error} = 9961, F = 7.193, df_{CMI} = 1, df_{error} = 7, p = 0.0315,

Figure 5B). Like the case for the Illumina data modeling R gene count from the PacBio data

revealed a similarly consistent but nominally insignificant association (MS_{region} = 53074 MS_{error} =

10164, F = 4.96. Degrees of freedom df_{region} = 2, df_{error} = 6, p = 0.054, see descriptive statistics in

the section above and in Table S4).

CNLs and TNLs

We did not find a significant relationship between CMI and TNL counts (p = 0.695, Figure 5D,

MS_{region} = 14.072, MS_{error} = 86.559, F = 0.1626. Degrees of freedom from Satterthwaite's method:

df_{region} = 1, df_{error} = 10.215), and only a marginal significance with CNL counts (Figure 5C;

p=0.0634, MS_{region} = 651.19, MS_{error} = 150.22, F = 4.3349. Degrees of freedom from

Satterthwaite's method: df_{region} = 1, df_{error} = 10.22).

Due to having only one plant sampled per prairie, we report the results of a fixed-effect

linear regression model for the PacBio dataset. Despite a much lower inferential power (n=9),

we still found a positive correlation between CMI and the PacBio R genes (Figure 5B; p=0.019).

There was a significant relationship between CMI and PacBio TNL counts (Figure S1B;

p=0.018), but not CNLs (Figure S1A; p=0.14).

Correlations Between Phenotypes and R Gene Counts

There were no significant correlations between Illumina R gene counts and leaf rust severity

(Figure 6B; p=0.63), rust response (Figure 6C; p=0.73), stem rust severity (Figure 6A; p=0.51),

or insect leaf herbivory (not pictured; p=0.23). We found that R gene counts were significantly

associated with leaf blotch scores (Figure 6D; p=0.0043). When source prairie was modeled as

a random effect, leaf blotch scores were still found to significantly correlate with CMI (p=8.94e-

5, slope=-0.0082).

Principal Coordinate Correlations Among Phenotypes

The PCoA analysis (Figure S4) of the unique NBARC sequences (Supplemental file

"AllNBARCs.fasta"; Table S5) against the disease phenotypes yielded no significant candidates

for genes responsible for disease resistance after correcting for multiple tests.

DISCUSSION

We enriched *Silphium integrifolium* libraries for R genes using baits developed from the related

Helianthus annuus genome to characterize the diversity across a rainfall gradient. We found

that the baits recovered a substantial portion of R genes in the reference. We found geographic

associations between R gene count and rainfall. Here, we discuss the *Silphium* R gene content

across the rainfall gradient, as well as the performance of the RenSeq enrichment method, and

the biological implications of the geographic pattern we uncovered.

DISEASE RESISTANCE AND RAINFALL

R Gene Counts Correlate with Precipitation

The significant correlation between net precipitation rate, as estimated by average CMI, and the

number of R genes in a plant's genome, detected in both the Illumina short-read dataset as well

as the underpowered PacBio long-read libraries (n=9), suggests evidence of possible genomic

adaptation for higher pathogen load. In fact, the steeper relationship of R gene number and CMI

observed with the PacBio long-read libraries suggests that the Illumina dataset underestimates

the strength of this relationship. The trend does not appear to be driven by either TNLs or CNLs,

for which the results conflict between the Illumina and PacBio datasets. Genes of type TNL, or

those containing a toll-interleukin-like receptor in their N-terminal domain, don't differ

significantly according to CMI. A stronger understanding of the relationship between climate

factors and the evolution of these two subtypes of R genes remains to be investigated.

Mixed Disease Phenotype Results

Despite the significant trend of the number of R genes increasing with CMI, leaf rust severity,

leaf rust response, and stem rust severity did not significantly vary in the common gardens.

These three traits are different phenotypic metrics to assess a specialist fungal pathogen,

Puccinia silphii, commonly known as *Silphium* rust. These results are consistent with results

from common gardens that found that eastern populations of *S. integrifolium* were not, on

average, more resistant to *Silphium* rust (Cassetta et al., 2023). One possible explanation for

why resistance to *Silphium* rust was not associated with variable R gene counts may lie in the

concept of the 'disease triangle' reviewed by Velásquez et al., 2019. The clones in this study

were placed in environments where *S. integrifolium* pathogens were present, and experienced

conditions that would be ideal for pathogenic growth: warm and wet. Even with effective

immunity represented by the R genes sequenced in the plant populations, the level of

resistance conferred by the resistance genes might have been obscured by the pathogen

spread. Disease resistance in these wild populations is highly likely to be quantitative, meaning

that resistance is not an all-or-nothing response, but rather a distribution of potential responses

ranging from minor to major (French et al., 2016). Further studies of the wild populations with

sequenced R genes, with controlled inoculations of plants with pathogens, would elucidate

associations between R genes that may confer various levels of response to pathogens.

Further, the evolutionary dynamics of R genes may be muddled by seemingly competing

forces of selection, with "arms race" dynamics expected to generate gradients in total host

resistance, while oscillatory "red-queen" dynamics may not (Bergelson et al., 2002). It is

possible, for example, that local co-evolution of *S. integrifolium* and *Silphium* rust has resulted in

adaptation of the rust to its local host population, rather than variation in overall resistance. This

red-queen dynamic is supported by observations from common gardens distributed along the

gradient that showed much greater virulence of *Silphium* rust on their populations from their

local region than from other regions (Cassetta et al. 2023). In the case of *Silphium* rust, R gene

identity may be more important than R gene diversity.

We did observe that resistance allele diversity was associated with suppression of leaf

blotch (Fig. 6d). Our result affirms observations of a previous common garden study that found

that eastern populations of *S. integrifolium* were more resistant to *Silphium* blotch, as well as to

clear vein virus (Cassetta et al. 2023). These results are consistent with expectations of greater

diversity of resistance alleles contributing to greater resistance to pathogens, and are

qualitatively consistent with arms-race-type dynamics. In *S. integrifolium*, leaf blotch is caused

by the generalist fungal pathogen *Colletotrichum dematium* (Cybernome; Farr 1989), as well as

its more specialized congener, *Colletotrichum silphii* Davis (Horst 2008). It is possible that arms

race dynamics are more likely between hosts and generalist pathogens such as leaf blotch,

while specialist pathogens such as the Silphium rust are more likely to generate red-queen

dynamics.

We note that the observed relationship between resistance allele diversity and

resistance to blotch may be spurious, as other heritable factors that covary with rainfall might

contribute to disease resistance, such as plant secondary chemicals. We note, however, that

insect herbivory may be an indicator of such overall resistance, and we did not observe any

relationship between R gene diversity and insect leaf herbivory. This result is consistent with R

genes predominantly acting as pathogen effector receptors, and not targeted toward resistance

for insects (Chovelon et al., 2021).

A BROADER PATTERN OF R GENE EVOLUTION

R Gene Genealogy Recovers Expected Patterns

The maximum likelihood genealogy constructed from the NBARC domains of the R genes of the

draft genomic assembly recovers the deep evolutionary divergence between CNL-type and

TNL-type R genes that has been reported in many other studies (Meyers et al., 2003; Mun et

al., 2017; Neupane et al., 2018). The instances of individual R genes nesting incongruently

within the wrong clade may be attributable to N-domain switching via recombination, which, to

the best of our reckoning, has yet to be observed. More likely, it may be due to misidentification

of upstream nucleotide sequence as incidentally translating to peptide sequence that meets the

identity threshold for one of the diagnostic motifs that places those R genes in the wrong

categorization.

PERFORMANCE OF RENSEQ FOR R GENE ENRICHMENT

Enrichment Using Distant Reference

The R gene libraries presented in this study were developed by enriching *Silphium integrifolium*

DNA for R genes using baits developed from *Helianthus annuus*, a model organism with well-

developed genomic resources. Despite an estimated divergence age of between between 22.5

(Meireles et al., 2020) and 33.5 million years ago (Zhang et al., 2022), the baits were able to

enrich libraries to contain a median of 63% of reads originating from R genes, representing a

36-fold enrichment over WGS. For comparison, a previous study by Andolfo et al. (2014) found

success enriching the *Solanum lycopersicum* (common tomato) with baits designed from

Solanum tuberosum (common potato), a congener estimated to be 6.7 Ma divergent by

TimeTree (Kumar et al., 2017). This study demonstrates the economical promise of RenSeq for

studying the immune systems of non-model plants under a variety of ecological and

evolutionary pressures. It also showcases the screening of crop wild relatives that are of

agronomic interest, such as *S. integrifolium*, for disease resistance genes that might enable

more robust response to pathogens that pose a challenge for the domestication of the plant.

The reference genome contained a much higher number of R genes in the draft genome

assembly compared to the enriched libraries (873 compared to ~400-600), which can be

accounted for by two factors. First, the draft genome is assembled from an F1 hybrid between

two different species, *S. integrifolium* and *S. perfoliatum*. While we expect overlap in many of

the R genes due to homology, our count is likely an overestimate of the true number contained

within the haploid genome of *S. integrifolium* due to the nature of R genes as a rapidly

diversifying gene family. Second, RenSeq enrichment likely captures a different (and smaller)

subset of the R genes than the WGS PacBio sequencing we employed for the draft assembly.

Caveats and Future Directions

Consistent with the prediction that more rainfall results in more pathogens, which result in more

disease resistance – we find a positive correlation between R gene counts and precipitation.

However, the distance between our observed correlation and our causal motivation is quite

large. Here we suggest two routes to bridge this gap – independent replication and unraveling

the causal chain.

Correlation does not imply causation. However, repeated independent replication "does waggle

its eyebrows suggestively and gesture furtively while mouthing 'look over there'." (Munroe

2009). Thus, a key step in establishing that the correlation uncovered reflects our biological

hypothesis, rather than happenstance, would be to evaluate the generality of this pattern. Some

such evidence already exists – similar results have been seen in RFLP based studies of a few R

genes in big bluestem and switchgrass (Rouse et al 2011, Zhu et al 2013). Testing for this

pattern in other *Silphium* species represents a promising direction, as they would provide

evolutionarily independent replication, while covering a similar precipitation gradient, and could

use the same RenSeq baits developed here. Extending this study to more distant taxa would

provide further evidence supporting this hypothesis.

Functional studies of these R-genes would provide more evidence for our motivating

causal hypothesis. A complete, phased, chromosome-level assembly of *S. integrifolium* will both

allow for better assessment of whether these genes are functional, and enable association

studies to determine the loci of pathogen resistance for incorporation into breeding programs.

Additional evaluation of the hypothesis that the number of pathogens affecting Silphium (and/or

the variation in their ability to evade a specific R gene), as well as associating specific NLR

alleles to resistance to specific pathogens would allow for a more mechanistic understanding of

the association uncovered in this paper.

CONCLUSION

We used RenSeq to sequence the R gene libraries of *S. integrifolium* plants from across a

rainfall gradient to evaluate the hypothesis that CMI imposes a biotic stress whose evolutionary

pressure should be evident within the genome. We detected a significant positive relationship

between CMI and R gene count, supporting our hypothesis. We demonstrate that RenSeq can

be used in the evaluation of ecological hypotheses and show promise for the economical

interrogation of the specific R gene loci responsible for disease resistance.

FUNDING

This work was supported by a grant from the National Science Foundations' Dimensions in

Biodiversity Grant [Grant ID: 1738041].

DATA ACCESSIBILITY AND BENEFIT SHARING

The R gene libraries generated in this study are available on the NCBI SRA database at project

accession PRJNA914292.

AUTHOR CONTRIBUTIONS

KK assembled and annotated the R gene libraries, performed the analyses, generated the

figures, and co-wrote the manuscript. KP performed the majority of the field work, common

garden experiments, and phenotyping, and co-wrote the manuscript. KT created disease scales

and led the recording of disease severity in the field. AH sequenced and assembled the draft

genome of *S. integrifolium*. AR designed the RenSeq baits, developed the libraries, and

performed the initial steps of bioinformatics. YB, KS, JB, KT and DVT conceived of the project

and contributed to the writing.

ACKNOWLEDGEMENTS

We acknowledge support for this research from NSF Dimension of Biodiversity grant 1738041.

This research was supported financially by the Perennial Agriculture Project, a joint project

between The Land Institute and The Malone Family Land Preservation Foundation. We

gratefully acknowledge financial support from the Foundation for Food and Agriculture Research

(FFAR) Fellows program. We thank Eric Cassetta for assisting with all of the field work. We are

thankful to Andrew Read and Peter Innes for assistance with code to generate the phylogeny

and map figures, respectively.

COI

Authors declare no conflicts of interest.

CITATIONS

Almeyda, C. V., Eid, S. G., Saar, D., Samuitiene, M., & Pappu, H. R. (2014). Comparative

analysis of endogenous plant pararetroviruses in cultivated and wild Dahlia spp. *Virus Genes*,

48(1), 140-152.

Andrews, S. (2010). FastQC: A Quality Control Tool for High Throughput Sequence Data

[Online]. Available online at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc/

Arora, S., Steuernagel, B., Gaurav, K., Chandramohan, S., Long, Y., Matny, O., Johnson, R.,

Enk, J., Periyannan, S., Singh, N., Hatta, M. A., Athiyannan, N., Cheema, J., Yu, G., Kangara,

N., Ghosh, S., Szabo, L. J., Poland, J., Bariana, H., Jones, J. D. G., Bentley, A. R., Ayliffe, M.,

Olson, E., Xu, S. S., Steffenson, B. J., Lagudah, E. & Wulff, B.B.H. (2019). Resistance gene

cloning from a wild crop relative by sequence capture and association genetics. Nat Biotechnol

37, 139–143 (2019).

Bankevich, A., Nurk, S., Antipov, D., Gurevich, A. A., Dvorkin, M., Kulikov, A. S., Lesin, V. M.,

Nikolenko, S. I., Pham, S., Prjibelski, A. D., Pyshkin, A. V., Sirotkin, A. V., Vyahhi, N., Tesler, G.,

Alekseyev, M. A., Pevzner, P. A. (2012). SPAdes: a new genome assembly algorithm and its

applications to single-cell sequencing. *Journal of Computational Biology*, 19(5), 455-477.

Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: a practical and

powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B

(Methodological)*, *57*(1), 289-300.

Bergelson, J., Kreitman, M., Stahl, E. A., and Tian, D. (2002). Evolutionary Dynamics of

Plant R-Genes. Science, 292 (July), 2281–2289. https://doi.org/10.1126/science.1061337

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina

sequence data. *Bioinformatics*, 30(15), 2114-2120.

Brun, P., Zimmermann, N. E., Hari, C., Pellissier, L., & Karger, D. N. (2022). Global climate-

related predictors at kilometre resolution for the past and future. *Earth System Science Data

Discussions*, 1-44.

Caplan, J., Padmanabhan, M., & Dinesh-Kumar, S. P. (2008). Plant NB-LRR immune receptors:
from recognition to transcriptional reprogramming. *Cell Host & Microbe*, *3*(3), 126-135.

Cassetta, E., Peterson, K., Bever, J. D., Brandvain, Y., VanTassel, D., Lubin, T. K., ... & Turner,
K. (2023). Adaptation of pathogens to their local plant host, Silphium integrifolium, along a
precipitation gradient. *Ecosphere*, *14*(6), e4565.

Cheng, H., Concepcion, G. T., Feng, X., Zhang, H., & Li, H. (2021). Haplotype-resolved de novo
assembly using phased assembly graphs with hifiasm. *Nature Methods*, *18*(2), 170-175.

Chovelon, V., Feriche-Linares, R., Barreau, G., Chadoeuf, J., Callot, C., Gautier, V., Le Paslier,
M. C., Berad, A., Faivre-Rampant, P., Lagnel, J., Boissot, N. (2021). Building a cluster of NLR
genes conferring resistance to pests and pathogens: the story of the Vat gene cluster in
cucurbits. *Horticulture Research*, *8*.

Clarkson, J. P., Fawcett, L., Anthony, S. G., & Young, C. (2014). A model for Sclerotinia
sclerotiorum infection and disease development in lettuce, based on the effects of temperature,
relative humidity and ascospore density. *PLoS One*, *9*(4), e94049.

Dabo, S. M., Mitchell, E. D., & Melcher, U. (1993). A method for the isolation of nuclear DNA
from cotton (Gossypium) leaves. *Analytical Biochemistry*, *210*(1), 34-38.

Delavaux, C. S., Schemanski, J. L., House, G. L., Tipton, A. G., Sikes, B., & Bever, J. D. (2021).
Root pathogen diversity and composition varies with climate in undisturbed grasslands, but less
so in anthropogenically disturbed grasslands. *The ISME journal*, *15*(1), 304-317.

Eitas, T. K., & Dangl, J. L. (2010). NB-LRR proteins: pairs, pieces, perception, partners, and
pathways. *Current Opinion in Plant Biology*, *13*(4), 472-477.

Farr, D. F., Bills, G. F., Chamuris, G. P., & Rossman, A. Y. (1989). *Fungi on Plants and Plant
Products in the United States*. APS press.

French, E., Kim, B. S., & Iyer-Pascuzzi, A. S. (2016). Mechanisms of quantitative disease
resistance in plants. In *Seminars in Cell & Developmental Biology* (Vol. 56, pp. 201-208).
Academic Press.

Glover, J. D., Reganold, J. P., Bell, L. W., Borevitz J., Brummer, E. C., Buckler, E.S., Cox, C.
M., Cox., T. S., Crews, T., E., Culman, W., Dehaan, R., Eriksson D., Gill, B., S., Holland, J., Hu,
F., Hulke B., S., Ibrahim, A. M. H., Jackson W., Jones, S. S., Murray, S., C., Paterson, A. H.,
Ploschuk E., Sacks, E. J., Snapp, S., Tao, D., Van Tassel, D. L., Wade L. J., Wysean D. L., Xu,
D. Y. (2010). Increased food and ecosystem security via perennial grains. *Science*, *328*(5986),
1638-1639.

Granke, L. L., & Hausbeck, M. K. (2010). Effects of temperature, humidity, and wounding on
development of Phytophthora rot of cucumber fruit. *Plant Disease*, *94*(12), 1417-1424.

Horst, R. Westcott's Plant Disease Handbook; Springer: New York, NY, USA, 2008.

Islam, T. M., & Toyota, K. (2004). Effect of moisture conditions and pre-incubation at low
temperature on bacterial wilt of tomato caused by Ralstonia solanacearum. *Microbes and*
Environments, *19*(3), 244-247.

Jupe, F., Pritchard, L., Etherington, G. J., MacKenzie, K., Cock, P. J., Wright, F., Sharma, S. K.,
Bolser, D., Bryan, G. J., Jones, J. D. G., Hein, I. (2012). Identification and localisation of the NB-
LRR gene family within the potato genome. *BMC Genomics*, *13*(1), 1-14.

Jupe, F., Witek, K., Verweij, W., Śliwka, J., Pritchard, L., Etherington, G. J., Maclean, D., Cock,
P. J., Leggett R. M., Bryan G. J., Cardle, L., Hein, I., Jones, J. D. (2013). Resistance gene
enrichment sequencing (RenSeq) enables reannotation of the NB-LRR gene family from
sequenced plant genomes and rapid mapping of resistance loci in segregating populations. *The*
Plant Journal, *76*(3), 530-544.

Kumar, S., Stecher, G., Suleski, M., & Hedges, S. B. (2017). TimeTree: a resource for timelines,
timetrees, and divergence times. *Molecular biology and evolution*, *34*(7), 1812-1819.

Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv preprint arXiv:1303.3997.

Magarey, R. D., Sutton, T. B., & Thayer, C. L. (2005). A simple generic infection model for foliar
fungal plant pathogens. *Phytopathology*, *95*(1), 92-100.

Meireles, J. E., Cavender-Bares, J., Townsend, P. A., Ustin, S., Gamon, J. A., Schweiger, A. K.,
Schaepman, M. E., Asner, G. P., Martin, R. E., Singh, A., Schrodt, F., Chlus., A, & O'Meara, B.
C. (2020). Leaf reflectance spectra capture the evolutionary history of seed plants. *New*
Phytologist, *228*(2), 485-493.

Meyers, B. C., Dickerman, A. W., Michelmore, R. W., Sivaramakrishnan, S., Sobral, B. W., &
Young, N. D. (1999). Plant disease resistance genes encode members of an ancient and
diverse protein family within the nucleotide-binding superfamily. *The Plant Journal*, *20*(3), 317-
332.

Meyers, B. C., Kozik, A., Griego, A., Kuang, H., & Michelmore, R. W. (2003). Genome-wide
analysis of NBS-LRR–encoding genes in Arabidopsis. *The Plant Cell*, *15*(4), 809-834.

Mun, J. H., Yu, H. J., Park, S., & Park, B. S. (2009). Genome-wide identification of NBS-
encoding resistance genes in Brassica rapa. *Molecular Genetics and Genomics*, *282*(6), 617-
631.
651
652   Neupane, S., Andersen, E. J., Neupane, A., & Nepal, M. P. (2018). Genome-wide identification
653   of NBS-encoding resistance genes in sunflower (Helianthus annuus L.). *Genes*, *9*(8), 384.
654
655   Oksanen, J. (2015). Vegan: an introduction to ordination. *R Project*,
656   https://cran.r-project.org/web/packages/vegan/vignettes/intro-vegan.pdf.
657
658   Munroe, R. (2009, March 6). Correlation [Webcomic]. Retrieved from https://xkcd.com/552/
659
660   Reinert, S., Money, K. L., Rockstad, G. B., Kane, N. C., Van Tassel, D. L., & Hulke, B. S.
661   (2018). Two contrasting laboratory methods improve Silphium integrifolium Michx. germination
662   rate to agronomically acceptable levels. *Euphytica*, *214*(9), 1-9.
663
664   Rouse, M. N., Saleh, A. A., Seck, A., Keeler, K. H., Travers, S. E., Hulbert, S. H., & Garrett, K.
665   A. (2011). Genomic and resistance gene homolog diversity of the dominant tallgrass prairie
666   species across the US Great Plains precipitation gradient. *PloS One*, *6*(4), e17641.
667
668   Rowlandson, T., Gleason, M., Sentelhas, P., Gillespie, T., Thomas, C., & Hornbuckle, B. (2015).
669   Reconsidering leaf wetness duration determination for plant disease management. *Plant*
670   *Disease*, *99*(3), 310-319.
671
672   Satterthwaite, F. E. (1946). An approximate distribution of estimates of variance components.
673   *Biometrics Bulletin*, *2*(6), 110-114.
674
675   Savary, S., Willocquet, L., Pethybridge, S. J., Esker, P., McRoberts, N., & Nelson, A. (2019).
676   The global burden of pathogens and pests on major food crops. *Nature Ecology & Evolution*,
677   *3*(3), 430-439.
678
679   Stamatakis, A. (2006). RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with
680   thousands of taxa and mixed models. *Bioinformatics*, *22*(21), 2688-2690.
681
682   Stamatakis, A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-analysis of
683   large phylogenies. *Bioinformatics*, *30*(9), 1312-1313.
684
685   Steuernagel, B., Witek, K., Krattinger, S. G., Ramirez-Gonzalez, R. H., Schoonbeek, H. J., Yu,
686   G., Baggs, E., Witek, A. I., Yadav, I., Krasileva, K., V., Jones, J. D. G., Uauy, C., Keller, B.,
687   Ridout, C. J., Wulff, B. B. (2020). The NLR-Annotator tool enables annotation of the intracellular
688   immune receptor repertoire. *Plant Physiology*, 183(2), 468-482.
689
690   Toda, N., Rustenholz, C., Baud, A., Le Paslier, M. C., Amselem, J., Merdinoglu, D., & Faivre-
691   Rampant, P. (2020). NLGenomeSweeper: a tool for genome-wide NBS-LRR resistance gene
692   identification. *Genes*, *11*(3), 333.
693

694   Turner, M. K., Ravetta, D., & Van Tassel, D. (2018). Effect of Puccinia silphii on yield
695   components and leaf physiology in Silphium integrifolium: Lessons for the domestication of a
696   perennial oilseed crop. *Sustainability*, *10*(3), 696.
697
Van Tassel, D. L., Albrecht, K. A., Bever, J. D., Boe, A. A., Brandvain, Y., Crews, T. E.,
Gansberger, M., Gerstberger, P., González-Paleo, L., Hulke, B. S., Kane, N. C., Johnson, P. J.,
Pestsova, E. G., Picasso-Risso, V. D., Prasifka, J. R., Ravetta, D. A., Schlautman, B., Sheaffer,
C. C., Smith, K. P., Speranza, P. R., Turner., M. K., Vilela, A. E., von Gehren, P., Wever, C.
(2017). Accelerating silphium domestication: an opportunity to develop new crop ideotypes and
breeding strategies informed by multiple disciplines. *Crop Science*, *57*(3), 1274-1284.
704
705   Velásquez, A. C., Castroverde, C. D. M., & He, S. Y. (2018). Plant–pathogen warfare under
706   changing climate conditions. *Current Biology*, *28*(10), R619-R634.
707
708   Vilela, A., González-Paleo, L., Turner, K., Peterson, K., Ravetta, D., Crews, T. E., & Van Tassel,
709   D. (2018). Progress and bottlenecks in the early domestication of the perennial oilseed Silphium
710   integrifolium, a sunflower substitute. *Sustainability*, *10*(3), 638.
711
712   Wieczorek, M. E., Jackson, S. E., & Schwarz, G. 2. (2018). Select attributes for NHDPlus
713   version 2.1 reach catchments and modified network routed upstream watersheds for the
714   conterminous United States. *US Geological Survey. https://doi. org/10.5066/F7765D7V*.
715
716   Witek, K., Lin, X., Karki, H. S., Jupe, F., Witek, A. I., Steuernagel, B., Stam, R., van Oosterhout,
717   C., Fairhead, S., Heal, R., Cocker, J. M., Bhanvadia, S., Barrett, W., Wu, C. H., Adachi, H.,
718   Song, T., Kamoun, S., Vleeshouwers, V. G. A. A., Tomlinson, L., Wulff, B. B. H., and Jones, J.
719   D. G. (2021). A complex resistance locus in Solanum americanum recognizes a conserved
720   Phytophthora effector. Nature Plants 7:198-208.
721
722   Zhang, C., Huang, C. H., Liu, M., Hu, Y., Panero, J. L., Luebert, F., Gao, T., & Ma, H. (2021).
723   Phylotranscriptomic insights into Asteraceae diversity, polyploidy, and morphological innovation.
724   *Journal of Integrative Plant Biology*, *63*(7), 1273-1293.
725
726   Zhu, Q., Bennetzen, J. L., & Smith, S. M. (2013). Isolation and diversity analysis of resistance
727   gene homologues from switchgrass. *G3: Genes, Genomes, Genetics*, *3*(6), 1031-1042.