Sperm Toolbox – A selection of small molecules to study human spermatozoa

Franz S. Gruber[a*], Anthony Richardson [b*], Zoe C. Johnston [c*], Rachel Myles [c], Neil Norcross [b], David Day [b], Irene Georgiou[b], Laura Sesma Sanz [a], Caroline Wilson [b], Kevin D. Read [b], Sarah Martins da Silva [c] Christopher L. R. Barratt [c], Ian H. Gilbert [b], Jason R. Swedlow [a]

a. Divisions of Computational Biology and Molecular, Cellular and Developmental Biology and National Phenotypic Screening Centre, School of Life Sciences, University of Dundee, Dundee, DD1 5EH, UK

b. Drug Discovery Unit, Division of Biological Chemistry and Drug Discovery, School of Life Sciences, University of Dundee, Dundee, DD1 5EH, UK

c. Division of Systems Medicine, School of Medicine, Ninewells Hospital and Medical School, University of Dundee, Dundee, DD1 9SY, UK

Correspondence: jrswedlow@dundee.ac.uk

ORCID
Anthony Richardson: 0009-0005-3690-3409
Rachel Myles: 0009-0000-1667-5879
Irene Georgiou: 0000-0002-6042-9816
Laura Sesma Sanz: 0000-0001-7792-1631
Christopher L.R. Barratt: 0000-0003-0062-9979
Franz S. Gruber: 0000-0003-2008-8460
Zoe C. Johnston: 0000-0003-0904-7166
Caroline Wilson: 0000-0001-9940-0052
Neil R. Norcross: 0000-0001-8050-5217
Sarah Martins de Silva: 0000-0003-2579-4866
Jason R. Swedlow: 0000-0002-2198-1958
Ian H. Gilbert: 0000-0002-5238-1314
Kevin D. Read: 0000-0002-8536-0130

Abstract

Male contraceptive options and infertility treatments are limited. To help overcome these
limitations we established a small molecule library as a toolbox for assay development and screening
campaigns using human spermatozoa. We have profiled all compounds in our automated high-
throughput screening platform and provide the dataset, which allows direct side-by-side
comparisons of compound effects and assays using live human spermatozoa.

Main

One of the grand challenges for science and society is the development of novel contraceptives that
are safe and deliver control of reproduction to men and women all over the world. In parallel, a
deeper understanding of sperm maturation, motility, fertilization, and possible interventions for
male infertility is required to counter declining birth rates. Recently, the application of advanced
technologies has helped move the human fertility field forward, revealing new potential targets for
both contraception and fertility, and new compounds as contraceptive candidates (reviewed by [1-3]).
A key part of this effort is the development of mechanistic insight and drug candidates that affect
the behaviour and properties of human spermatozoa [4-7]. Towards this end, we have recently
developed an automated live human sperm phenotypic, target-agnostic screening platform that can
profile 1000s of compounds and identify modifiers of sperm function [8,9]. These effects can be either
positive or negative and therefore are starting points for either fertility treatments (including
medical assisted reproductive technologies) or contraceptive development. An advantage of using a
high-throughput platform is the ability to efficiently compare many compounds under the same
assay conditions, within a short amount of time.

Target-agnostic phenotypic screening can be enhanced by assaying libraries of compounds with
known targets. The resulting reference datasets can suggest mechanism(s) of action and target
identification for candidate compounds during a screening campaign. To facilitate our own, and
others, drug discovery efforts in contraception and infertility, we assembled a small library
containing 84 small molecules, which we have coined the Sperm Toolbox (STB). The idea of the STB
is in concordance with similar efforts e.g. the MMV Pandemic Response Box [10], the SGC Epigenetic
Chemical Probe Collection [11], or the Drug Repurposing Hub [12] that facilitate deeper understanding of
biology or pathology or repurpose approved drugs. We tested the STB compounds in multiple
functional sperm assays, as well as more general cytotoxicity and aqueous solubility assays (Figure
1A). The STB has been assembled to contain previously published compounds and compounds
detected in our own screening campaigns [8,9]. In addition, we included a few compounds which have
been inferred from patent applications or studies of proteins potentially relevant for sperm function
(Figure 1B). In constructing the STB, we have focussed on compounds reported in the literature that
affect human systems, but compounds tested in relevant animal models, cell-based, or biochemical
systems have also been included. Published analyses of the compounds selected in the STB have
indicated effects on crucial sperm functions e.g. motility, capacitation, acrosome reaction, Ca^{2+}
signalling, viability, or related to other processes (e.g. spermatogenesis, liquefaction, or ejaculation)
(Figure 1C). Compound annotations are included in the STB dataset and indicate that
Enzymes/Kinases, Receptors, channels and transporters are the most represented compound
classes. Using information from the Drug Repurposing Hub [12], we generated a protein-protein
interaction network of targets related to STB compounds (Figure 1E). This network consists of
multiple subclusters related to functions in spermatozoa (Figure 1F) and other biological processes,
functions, and pathways (Figure 1G).

Figure 1

(A) The Sperm Toolbox is a collection of 84 small molecules with published, observed or inferred effect on multiple biological processes related to the biogenesis or function of spermatozoa. Human sperm cells have been tested in various assays, which allows comparison of compounds under the same experimental conditions. (B) Summary of the sperm toolbox compound compositions. (C) Summary of annotated (i.e. published, observed, inferred) sperm function of Toolbox compounds. (D) Coarse classification of Sperm Toolbox target classes. (E) StringDB Protein-Protein interaction networks (Full network, confidence 0.7) of putative targets of Sperm toolbox compounds. Protein targets have been inferred from the repurposing hub. Network has been clustered using MCL clustering in Cytoscape. (F) Sperm related terms (GeneOntology, Reactome) within the network shown in (E). Marker size indicates false discovery rate (FDR). (G) Enriched terms within the biggest subclusters in (E). Redundant terms have been filtered using Cytoscape (cutoff 0.5). Color matches subcluster color in (E).

To compare the effects of STB compounds on specific sperm functions, we screened them in
multiple phenotypic functional assays. Each sperm function assay has been repeated in multiple
donor pools (see Material and Methods), where each pool consists of cells donated from at least 3
different donors to address donor to donor variability. Each compound has been tested at two
concentrations, 10 and 30 µM, and normalized to vehicle controls (DMSO). We have modified our
assays from the previously published workflow [8] by increasing incubation time and introducing
conditions that support capacitation, which is a sperm maturation process required for fertilization.
These changes expand the range of targets that we can explore in our assays. Our live-sperm-based
high-throughput assays use an automated microscope to record short timelapse movies and can run
in different modes. The first mode, denoted 'motility assay', utilises non-capacitating assay
conditions that model the state of spermatozoa after ejaculation. The second mode, denoted
'capacitation assay', utilises conditions which support capacitation (Extended Figure 1A). Both
modes allow sperm motility to be measured. In addition, the capacitation assay also scores
hyperactivation (i.e. distinct movement patterns concomitant with capacitation which are crucial for
fertilization)

The acrosome assays utilize a high-throughput flow cytometer to score for compounds inducing the
acrosome reaction, using peanut agglutinin conjugated to a fluorescent dye to label the inner
acrosomal membrane. As with the motility assays, we have developed two different screening
modes, named 'acrosome 1' and 'acrosome 2' (Extended Figure 1A). Both modes are capable of
measuring induction of the acrosome reaction. However, in the acrosome 2 assay, a mixture of
physiological agonists of the acrosome reaction is used [13]. This addition allowed measurement of
inhibition of acrosome reaction.

We also tested the effect of STB compounds on sperm viability using a complementary live-cell flow-
cytometer approach and propidium iodide, which does not permeate intact cell membranes
(Extended Figure 1B).

Finally, we also profiled each STB compound for general cytotoxicity in two cell-lines (HeLa and
HepG2). The HeLa-based assay uses the cell painting assay [14], while the HepG2 assay utilizes a
resazurin based viability read-out [15]. To ensure usability of STB compounds, we also measured
aqueous solubility. An overview of additional assays is given in Extended Figure 1C.

This multi-assay compound profiling dataset reveals a broader picture of the action of individual
compounds and helps find phenotypic similarities induced by compounds. Using unsupervised
clustering we observed clusters of compounds with a significant reduction of sperm motility and/or
increased amounts of acrosome reacted cells (e.g. Alexidine dihydrochloride, Tafenoquin or A23187)
(Figure 2A). Some of these compounds also show significant levels of cytotoxicity in HeLa and/or
HepG2 cells as well as decreased sperm viability suggesting that their effect on motility and
acrosome reaction is due to toxic effects on spermatozoa. However, other clusters show a
significantly increased acrosome reaction and hyperactive motility but do not show effects on sperm
viability or general cytotoxicity (e.g. LRRK2-IN-1, Clofarabine, Linsitinib, Trequinsin). One example,
LRRK2-IN-1, which enhances motility (progressive and hyperactivated), has been described as a
selective inhibitor of LRRK2 kinase, which is involved in Parkinson's disease [16]. This compound shares
some structural similarities with two kinase inhibitors in the STB, which decrease sperm motility:
Compound 19, described as a potent Testis-Specific Serine/Threonine Kinase 2 (TSSK2) inhibitor [17]
and CZC 25146, annotated as a potent LRRK2 inhibitor.

Interestingly, we observed that a number of compounds show weak or no effects in our assays
(Extended figure 2; grey datapoints). Given these compounds were sourced from publications,
patents, and screening systems that report activity in sperm, this highlights the need for
standardisation of conditions in phenotypic assays based on human sperm. The STB, and the use of
high-throughput assays allows the profiling of compounds using the same assay conditions (i.e.
buffer system, incubation time and screening concentration) on a large number of human
spermatozoa, thus providing reproducible, quantitative assay results. This combination also
facilitates the comparison of the effects of the STB compounds between different physiological
states (Extended Figure 1A&B), in a standardised, rapid, and quantitative manner that has not been
previously possible in the field of sperm biology. This standardized approach provides an excellent
starting point to establish control compounds and assay conditions for novel drug discovery
campaigns related to contraception and infertility. We provide the complete dataset (see
Supplement Information) as well as web interface for data analysis
(https://spermtoolbox.shinyapps.io/spermtoolbox/).

This work represents the first version of the STB. We aim to release annual updates with newly
published compounds modulating functions in spermatozoa. In our next release we aim to include
compounds that were published during the collection of the data for this first release e.g., a sAC
inhibitor TDI-11861 [4], a CatSper inhibitor RU1968 [18], a SLO3 inhibitor [19] or other contributions from
the research community. The STB is a reference for compounds that modulate sperm function and
can become an important resource for benchmarking new assays and new compounds. We aim to
use the STB as a reference in our screens and to define as many factors targeted by STB compounds
as possible that are critical for sperm function.

A Sperm Motility assays

Motility
Non-capacitating media

time with compounds

1 h recovery · 10 min · 3 h · 6 h

Capacitation
Capacitating media

30 min capacitation · 30 min with compounds

◆ Microscope read-out

B Sperm Acrosome assays

Acrosome 1
Capacitating media

3 h with compounds

Acrosome 2
Capacitating media

3 h with compounds · 1 h with agonist

Sperm Viability
Non-capacitating media

3 h with compounds

● Flow-cytometer read-out

C Additional assays

HeLa cytotoxicity

24 h with compounds · Microscope read-out

HepG2 cytotoxicity

70 h with compounds · Fluorometric read-out

Aqueous solubility

24 h with compounds · uHPLC read-out

Extended Figure 1
(A) Diagram showing sperm motility assays run on high-content microscope with compound incubation time/read-out time. Motility assays were run under non-capacitating conditions, with a 1 hour recovery phase after preparation of spermatozoa. Capacitation assays were run under capacitating conditions. Spermatozoa are capacitated for 30 min prior to compound incubation.
(B) Diagram showing assays run on high-throughput flow cytometer to measure acrosome status and sperm viability. Incubation time with compound, agonist and read-out times are indicated. Acrosome assays run under capacitating conditions. Sperm viability assay run under non-capacitating condition.
(C) Additional assays performed on sperm toolbox compounds.

Figure 2

(A) Heatmap showing standardized results (z-score) for every Toolbox compound screened at 30 µM in selected assays: Acrosome 1 (3h with compound then induction with acrosome reaction using an agonist mix), Motility Hyperactive (30 min capacitation then 30 min incubation with compound), Motility Progressive (3h incubation with compounds under non-capacitating conditions), Motility Total (3h incubation with compound under non-capacitating conditions)), Solubility (aqueous solubility of compounds after 24h incubation), Sperm viability (3h with compound then propidium iodide assay), Toxicity HeLa (24h compound incubation then cell painting assay), Toxicity HepG2 (70h incubation with compound then resaruzin based assay). Heatmap color indicates decrease (blue) or increase (red).

(B) UMAP plot showing groups of compounds with similar assay profile. Color indicates annotated sperm function.

(C) Structures and zoom-in of three examples related to LRRK2 kinase.

Extended Figure 2

Volcano plots of Acrosome, Capacitation and Motility assays. Upper panel indicates 10 µM, lower plot panels indicate 30 µM. Incubation times for assays are 3h with compound for Acrosome assay and Motility (non-capacitating) (PM = progressive motility), and 30 min for the Capacitation assay (HA = hyperactive motility, PM = progressive motility). Dotted line indicates significance levels of 0.05. Any point above dotted line is indicated in magenta. Buffer conditions are indicated below Plot title (capacitating vs. non-capacitating)

Material and Methods

Ethical approval

156 Written consent was obtained from each donor in accordance with under local ethical approval
157 (University of Dundee, SMED REC 20/45).

Sperm handling

Donated human spermatozoa with normal semen characteristics (concentration, motility; WHO
2010) from healthy volunteers with no known fertility problems were used in this study. Samples
were obtained by masturbation after ≥ 48 h of sexual abstinence, liquefied for 30 min at 37 °C, and
processed using density gradient centrifugation, to separate cells using 40/80% Percoll (Sigma
Aldrich, UK) fractions. Every experiment has been performed on pooled spermatozoa from ≥ 3
donors each.

For experiments in non-capacitating conditions, we used a buffer system slightly modified from
(Tardif et al., 2014) using Minimal Essential Medium Eagle (Sigma), supplemented with HEPES (1 M
solution, Gibco), sodium lactate (DL-solution, Sigma), sodium pyruvate (100 mM solution, Gibco) and
bovine serum albumin (7.5 % solution, Sigma).

For experiments in capacitating conditions, we used a buffer system slightly modified from the HTF from [13] to a final composition of 97.8 mM NaCl, 4.69 mM KCl, 0.2 mM $MgSO_4$, 0.37 mM KH_2PO_4, 2.04 mM $CaCl_2$, 0.33 mM Na-pyruvate, 12.4 mM Na-lactate, 2.78 mM glucose, 21 mM HEPES, 25 mM $NaHCO_3$ and 3 mg/mL BSA. All buffer components were supplied by Sigma-Aldrich unless otherwise stated and all buffer systems were adjusted to pH 7.4.

Compound handling

Compounds were purchased from vendors indicated in supplementary file x. 10 mM stock solutions in DMSO were generated. Quality control of each compound has been performed (HPLC-MS and NMR).

Compound plates were generated using 10 mM stock solutions of each STB compound in Echo 384-LDV plates (LP-0200). Prior to running experiments, assay ready plates have been generated using Echo acoustic dispensers (550/555). Each compound was screened at 10 and 30 µM final concentration. Plate types used for assays: Motility, HeLa cell painting and Sperm viability (PerkinElmer 384-well PhenoPlate), Acrosome (Greiner 384-PP V-bottom plates).

Motility assays

Motility assay (Non-capacitating):

Cells were rested for 1 h post density gradient centrifugation. 10 µL assay buffer was dispensed into assay ready plates, plates were put onto a shaker instrument for 30 seconds using 1,000 rpm, followed by a short pulse centrifugation. Plates were incubated for 15-30 mins at 37 °C prior to addition of 10 µL of pooled cells (at 1 M/mL concentration). Plates were then incubated in a Yokogawa CV7000 microscope set at 37 °C for 10 min prior to imaging. Imaging a 384-well plate requires ≤ 20 min using the following settings: 2 positions per well, 0.5 sec timelapse movies (3x binning; 11 ms exposure time, 22 ms interval time, 500 intervals). A second and a third timepoint was recorded after 3 h, and 6 h incubation at 37 °C. Data has been processed as described in Gruber et al., 2020.

Capacitation assay:

Cells were capacitated for 30 min at 37 °C and 5 % CO2. 10 µL assay buffer was dispensed into assay ready plates, plates were put onto a shaker instrument for 30 seconds using 1,000 rpm, followed by a short pulse centrifugation. Plates were incubated for 15-30 mins at 37 °C and 5% CO2 prior to addition of 10 µL of pooled cells (at 1 M/mL concentration). This was followed by a 20 min compound incubation step at 37 °C and 5 % CO2. Plates were then incubated in our CV7000 microscope set at 37 °C and 5% CO2 for 10 min prior to imaging. Imaging a 384-well plate requires ≤ 20 min using the following settings: 2 positions per well, 0.5 sec timelapse movies (3x binning; 11 ms exposure time, 22 ms interval time, 500 intervals). Data has been processed as described in Gruber et al., 2020.

Acrosome assays

In both experimental modes, 20 µL of pooled cells (at 1 M/mL concentration) were capacitated for 3 h in presence of compounds at 37 °C and 5% CO2. In mode a, cells were then stained with PNA-Alexa647 (ThermFisher, 1 mg/mL, final dilution 1:1000) and Hoechst dye for 30 min at 37 °C and 5% CO2, followed by 10 min fixation at RT by adding 20 µL of 4% paraformaldehyde (Sigma) in PBS. In mode b, a cocktail of progesterone, prostaglandin A and NH4Cl was added to all compound wells and incubated for an additional 1 h at 37 °C and 5% CO2 to induce acrosome reaction. This was followed by fixation and staining protocol as for mode a. In both modes fixatives and stains were removed and cells were resuspended in PBS, using a 405 platewasher. Plates were then measured on a Novocyte Advanteon high-throughput flow cytometer. Gating of data has been performed using the flow cytometer software to export counts of gated populations.

Sperm viability assay

Cells were incubated with compounds for a total of 3h at 37 °C. 20 min prior, propidium iodide was added (1.5 mM solution, Sigma, final dilution 1:2000). Plates were analysed using a Satorius iQue Screener with settings to allow for a sampling time of ≤30 min per 384-well plate. Gating of data has been performed using the flow cytometer software to export counts of gated populations.

HeLa cell painting

Cell were incubated with compounds for 24h. Cell painting was performed after Bray et al., 2016 with modifications suggested here (Cimini et al., bioRxiv 2022.07.13.499171; doi: https://doi.org/10.1101/2022.07.13.499171):

-   10 µL of MitoTracker (Life Technologies, M22426, 1 mM stock in DMSO as recommended by the vendor) is added to 50 µL of media (1:2000 final dilution)
-   Phalloidin-Alexa594 (Life Technologies, A12381, 66 µM stock in methanol as recommended by the vendor, 1:500 final dilution)
-   WGA-Alexa594 (Life Technologies, W11262, 5 mg/mL stock, 1:250 final dilution) was used
-   DAPI (5 mg/mL stock; 1:1000 final dilution) was used to stain DNA
-   20 µL of staining solution was used
-   PBS was used for washing steps

HepG2 cytotoxicity assay

HepG2 assay was performed as described previously [15].

Aqueous solubility

The aqueous solubility of test compounds was measured using UHPLC, as previously described in [20].

Data analysis

For cell painting imaging data, plates of images were imported into OMERO [21]. Cell segmentation and feature extraction were performed using CellProfiler 3 [22] and using instructions from [14] to normalize data. Timelapse data of motility assays was processed as previously described [8]. Acrosome flow cytometry data was processed using NovoCyte Express (PerkinElmer), to gate out background debris, select for single cells and calculate percentage values of acrosome positive events. Sperm viability flow cytometry data was processed using iQue Forecyt (Satorius) to gate out background debris, select for single cells and calculate percentage values of propidium iodide positive events. For all sperm functional assays, data was normalized to DMSO control wells. Each plate had 16 DMSO wells. A median value of all 16 wells was used to calculated percentage change ($pct\_ch = ((value / DMSO\_median) - 1) * 100$). For sperm viability we calculated fold-change ($fc = 1 - (value / DMSO\_median)$). For each compound we calculated a p-value using a Welch's t-test implemented in scipy (scipy.stats.ttest_ind) comparing a compounds effect to DMSO controls.

The following packages have been used for data analysis and visualization:

R: Tidyverse, plotly, pHeatmap

Python: Pandas, NumPy, SciPy, Seaborn, Matplotlib, UMAP

Acknowledgements

The authors acknowledge support by the Bill & Melinda Gates Foundation (INV-007117).

References

1. Norcross, N.R., Georgiou, I., Johnston, Z.C., Gruber, F.S., Swedlow, J.R., Read, K.D., Barratt, C.L., and Gilbert, I.H. (2022). Male contraceptive development: A medicinal chemistry perspective. Eur J Med Chem *243*, 114709. 10.1016/j.ejmech.2022.114709.

2. Anderson, D.J., and Johnston, D.S. (2023). A brief history and future prospects of contraception. Science *380*, 154-158. 10.1126/science.adf9341.

3. Service, C.A., Puri, D., Hsieh, T.C., and Patel, D.P. (2023). Emerging concepts in male contraception: a narrative review of novel, hormonal and non-hormonal options. Ther Adv Reprod Health *17*, 26334941221138323. 10.1177/26334941221138323.

4. Balbach, M., Rossetti, T., Ferreira, J., Ghanem, L., Ritagliati, C., Myers, R.W., Huggins, D.J., Steegborn, C., Miranda, I.C., Meinke, P.T., et al. (2023). On-demand male contraception via acute inhibition of soluble adenylyl cyclase. Nat Commun *14*, 637. 10.1038/s41467-023-36119-6.

5. Faber, E.B., Wang, N., John, K., Sun, L., Wong, H.L., Burban, D., Francis, R., Tian, D., Hong, K.H., Yang, A., et al. (2023). Screening through Lead Optimization of High Affinity, Allosteric Cyclin-Dependent Kinase 2 (CDK2) Inhibitors as Male Contraceptives That Reduce Sperm Counts in Mice. J Med Chem *66*, 1928-1940. 10.1021/acs.jmedchem.2c01731.

6. Carlson, E.J., Francis, R., Liu, Y., Li, P., Lyon, M., Santi, C.M., Hook, D.J., Hawkinson, J.E., and Georg, G.I. (2022). Discovery and Characterization of Multiple Classes of Human CatSper Blockers. ChemMedChem *17*, e202000499. 10.1002/cmdc.202000499.

7.　Hau, R.K., Tash, J.S., Georg, G.I., Wright, S.H., and Cherrington, N.J. (2022). Physiological
Characterization of the Transporter-Mediated Uptake of the Reversible Male Contraceptive
H2-Gamendazole Across the Blood-Testis Barrier. J Pharmacol Exp Ther *382*, 299-312.
10.1124/jpet.122.001195.

8.　Gruber, F.S., Johnston, Z.C., Barratt, C.L., and Andrews, P.D. (2020). A phenotypic screening
platform utilising human spermatozoa identifies compounds with contraceptive activity.
Elife *9*. 10.7554/eLife.51739.

9.　Gruber, F.S., Johnston, Z.C., Norcross, N.R., Georgiou, I., Wilson, C., Read, K.D., Gilbert, I.H.,
Swedlow, J.R., Martins da Silva, S., and Barratt, C.L.R. (2022). Compounds enhancing human
sperm motility identified using a high-throughput phenotypic screening platform. Hum
Reprod *37*, 466-475. 10.1093/humrep/deac007.

10.　Samby, K., Besson, D., Dutta, A., Patra, B., Doy, A., Glossop, P., Mills, J., Whitlock, G., Hooft
van Huijsduijnen, R., Monaco, A., et al. (2022). The Pandemic Response Box horizontal line
Accelerating Drug Discovery Efforts after Disease Outbreaks. ACS Infect Dis *8*, 713-720.
10.1021/acsinfecdis.1c00527.

11.　Brown, P.J., and Muller, S. (2015). Open access chemical probes for epigenetic targets.
Future Med Chem *7*, 1901-1917. 10.4155/fmc.15.127.

12.　Corsello, S.M., Bittker, J.A., Liu, Z., Gould, J., McCarren, P., Hirschman, J.E., Johnston, S.E.,
Vrcic, A., Wong, B., Khan, M., et al. (2017). The Drug Repurposing Hub: a next-generation
drug library and information resource. Nat Med *23*, 405-408. 10.1038/nm.4306.

13.　Prajapati, P., Kane, S., McBrinn, R.C., Dean, M.S., Martins da Silva, S.J., and Brown, S.G.
(2022). Elevated and Sustained Intracellular Calcium Signalling Is Necessary for Efficacious
Induction of the Human Sperm Acrosome Reaction. Int J Mol Sci *23*.
10.3390/ijms231911253.

14.　Bray, M.A., Singh, S., Han, H., Davis, C.T., Borgeson, B., Hartland, C., Kost-Alimova, M.,
Gustafsdottir, S.M., Gibson, C.C., and Carpenter, A.E. (2016). Cell Painting, a high-content
image-based assay for morphological profiling using multiplexed fluorescent dyes. Nat
Protoc *11*, 1757-1774. 10.1038/nprot.2016.105.

15.　Baragana, B., Hallyburton, I., Lee, M.C., Norcross, N.R., Grimaldi, R., Otto, T.D., Proto, W.R.,
Blagborough, A.M., Meister, S., Wirjanata, G., et al. (2015). A novel multiple-stage
antimalarial agent that inhibits protein synthesis. Nature *522*, 315-320.
10.1038/nature14451.

16.　Deng, X., Dzamko, N., Prescott, A., Davies, P., Liu, Q., Yang, Q., Lee, J.D., Patricelli, M.P.,
Nomanbhoy, T.K., Alessi, D.R., and Gray, N.S. (2011). Characterization of a selective inhibitor
of the Parkinson's disease kinase LRRK2. Nat Chem Biol *7*, 203-205. 10.1038/nchembio.538.

17.　Hawkinson, J.E., Sinville, R., Mudaliar, D., Shetty, J., Ward, T., Herr, J.C., and Georg, G.I.
(2017). Potent Pyrimidine and Pyrrolopyrimidine Inhibitors of Testis-Specific
Serine/Threonine Kinase 2 (TSSK2). ChemMedChem *12*, 1857-1865.
10.1002/cmdc.201700503.

18.　Schierling, T., Tosi, B., Eisenhardt, C., Reining, S., Daniliuc, C.G., Brenker, C., Strunker, T., and
Wunsch, B. (2023). Synthesis and Functional Characterization of Novel RU1968-Derived

328   CatSper Inhibitors with Reduced Stereochemical Complexity. ACS Pharmacol Transl Sci *6*,
329   115-127. 10.1021/acsptsci.2c00188.

330   19.   Lyon, M., Li, P., Ferreira, J.J., Lazarenko, R.M., Kharade, S.V., Kramer, M., McClenahan, S.J.,
331   Days, E., Bauer, J.A., Spitznagel, B.D., et al. (2023). A selective inhibitor of the sperm-specific
332   potassium channel SLO3 impairs human sperm function. Proc Natl Acad Sci U S A *120*,
333   e2212338120. 10.1073/pnas.2212338120.

334   20.   Smith, A., Wall, R.J., Patterson, S., Rowan, T., Rico Vidal, E., Stojanovski, L., Huggett, M.,
335   Hampton, S.E., Thomas, M.G., Corpas Lopez, V., et al. (2022). Repositioning of a
336   Diaminothiazole Series Confirmed to Target the Cyclin-Dependent Kinase CRK12 for Use in
337   the Treatment of African Animal Trypanosomiasis. J Med Chem *65*, 5606-5624.
338   10.1021/acs.jmedchem.1c02104.

339   21.   Allan, C., Burel, J.M., Moore, J., Blackburn, C., Linkert, M., Loynton, S., Macdonald, D.,
340   Moore, W.J., Neves, C., Patterson, A., et al. (2012). OMERO: flexible, model-driven data
341   management for experimental biology. Nat Methods *9*, 245-253. 10.1038/nmeth.1896.

342   22.   McQuin, C., Goodman, A., Chernyshev, V., Kamentsky, L., Cimini, B.A., Karhohs, K.W., Doan,
343   M., Ding, L., Rafelski, S.M., Thirstrup, D., et al. (2018). CellProfiler 3.0: Next-generation image
344   processing for biology. PLoS Biol *16*, e2005970. 10.1371/journal.pbio.2005970.