1

Genetic manipulation of candidate phyla radiation bacteria provides

functional insights into microbial dark matter

Yaxi Wang[1,†], Larry A. Gallagher[1,†], Pia A. Andrade[1], Andi Liu[1], Ian R. Humphreys[2,3], Serdar Turkarslan[4], Kevin J. Cutler[5], Mario L. Arrieta-Ortiz[4], Yaqiao Li[1,4], Matthew C. Radey[1], Qian Cong[6,7,8], David Baker[2,3,9], Nitin S. Baliga[4], S. Brook Peterson[1], and Joseph D. Mougous[1,9,10,*]

[1]Department of Microbiology, University of Washington, Seattle, WA 98109, USA
[2]Department of Biochemistry, University of Washington, Seattle, WA 98109, USA
[3]Institute for Protein Design
[4]Institute for Systems Biology, Seattle, WA 98109, USA
[5]Department of Physics, University of Washington, Seattle, WA 98195, USA
[6]Eugene McDermott Center for Human Growth and Development, University of Texas Southwestern Medical Center, Dallas, TX, USA.
[7]Department of Biophysics, University of Texas Southwestern Medical Center, Dallas, TX, USA.
[8]Harold C. Simmons Comprehensive Cancer Center, University of Texas Southwestern Medical Center, Dallas, TX, USA.
[9]Howard Hughes Medical Institute, University of Washington, Seattle, WA 98195, USA
[10]Microbial Interactions and Microbiome Center, University of Washington, Seattle, WA 98109, USA

† Authors contributed equally.

* To whom correspondence should be addressed:

    Email – mougous@uw.edu

Summary

The study of bacteria has yielded fundamental insights into cellular biology and physiology,

biotechnological advances and many therapeutics. Yet due to a lack of suitable tools, the

significant portion of bacterial diversity held within the candidate phyla radiation (CPR) remains

inaccessible to such pursuits. Here we show that CPR bacteria belonging to the phylum

Saccharibacteria exhibit natural competence. We exploit this property to develop methods for

their genetic manipulation, including the insertion of heterologous sequences and the

construction of targeted gene deletions. Imaging of fluorescent protein-labeled Saccharibacteria

provides high spatiotemporal resolution of phenomena accompanying epibiotic growth and a

transposon insertion sequencing genome-wide screen reveals the contribution of enigmatic

Saccharibacterial genes to growth on their Actinobacteria hosts. Finally, we leverage

metagenomic data to provide cutting-edge protein structure-based bioinformatic resources that

support the strain *Southlakia epibionticum* and its corresponding host, *Actinomyces israelii*, as a

model system for unlocking the molecular underpinnings of the epibiotic lifestyle.

Introduction

The vast majority of metagenomic DNA sequences obtained from microbial species-rich environmental sources is derived from Bacteria and Archaea that have not been cultivated. Conservative estimates suggest that these sequences, often referred to as microbial dark matter, represent organisms constituting approximately half of phylum level diversity within these domains[1]. Microbial dark matter holds great interest as a reservoir of biosynthetic pathways and enzymes with potential for biotechnological application[2]. In addition, understanding the functions of these genes is paramount to defining the molecular processes supporting a given ecosystem and for unraveling the physiology and cell biology of the organisms within[3].

The candidate phyla radiation (CPR) of bacteria are reported to contain well over 50 phyla, only four of which have cultivated representatives[4]. This large monophyletic lineage of phyla and superphyla, including the group currently known as Patescibacteria, is thought to represent 15-50% of total bacterial diversity and contributes disproportionately to microbial dark matter[5]. Common features of CPR bacteria are their small size (as little as 100-200 nm in width), reduced genomes (typically < 1 megabase), and limited metabolic capability[6]. This has led to the hypothesis that these organisms broadly share a requirement for host organisms to support their growth. Indeed, experiments show that most cultivated CPR bacteria attach to and proliferate on the surface of other bacteria – living as obligate epibionts[4 7-10]. Genomic analyses have yielded speculation regarding the molecular functions that support the epibiotic lifestyles of CPR bacteria[5,11]. However, owing to the phylogenetic distance separating CPR and well characterized bacteria, the function of much of their proteome cannot be predicted and a lack of genetic tools for these organisms has heretofore precluded the experimental investigation of genotype–phenotype relationships.

Among the CPR, members of the phylum Saccharibacteria, originally named Tm7, were

the first to be cultivated in the laboratory[8]. Saccharibacteria are found in a multitude of terrestrial

and marine environments, yet early interest in them stemmed from their widespread occurrence

in human oral microbiomes[12]. Archaeological findings show this association dates to before the

mesolithic period and recent work links Saccharibacteria to human oral health[12-14]. The growth of

Saccharibacteria relies on the co-cultivation of host bacteria belonging to the class

Actinomycetia with the phylum Actinomycetota, for which they exhibit strain level specificity[15-

17]. Employing panels of Actinomycetia strains for Saccharibacterial enrichment has facilitated

the isolation and sequencing dozens of strains[16,18,19]. Despite this progress, the phylum remains

poorly sampled, with many divergent clades uncultivated, and the extent of genetic diversity

unresolved.

The genome sequences of Saccharibacteria reveal common features that provide insight

into molecular functions underlying their cellular physiology and lifestyle. Akin to other CPR

bacteria, Saccharibacteria generally lack a respiratory chain, and pathways for the *de novo*

generation of amino acids, nucleotides, and fatty acids[6]. On the contrary, Saccharibacteria

universally possess a relative wealth of specialized secretory mechanisms including the type II

and IV secretion systems (T2SS, T4SS)[11]. These diverge significantly from related systems in

bacterial pathogens that deliver toxins and effectors to eukaryotic host cells; however, it has been

speculated that they function in an analogous fashion to support bacterial host co-option by

Saccharibacteria[11]. Saccharibacteria also possess type IV pili (T4P), which were implicated in

twitching motility and host adhesion through the use of a small molecule inhibitor of pilus

extrusion[17]. Though such genomic analyses and experiments provide fertile ground for the

formulation of hypotheses, progress toward a mechanistic understanding of the unique biology of

Saccharibacteria and CPR bacteria as a whole has been stymied by a lack of genetic tools[20].

Here, we discover that natural competence can be harnessed for genetic manipulation of

Saccharibacteria. With this capability in hand, we go on to use fluorescent protein expression to

conduct time-lapse microscopic analysis of the Saccharibacterial lifecycle and we perform

transposon mutagenesis to identify genes important for epibiotic growth. Our findings offer an

initial mechanistic glimpse into the cellular functions encoded in microbial dark matter.

Results

Isolation and characterization of Saccharibacteria strains

We collected, pooled, homogenized, and filtered saliva and dental plaque from

volunteers, and enriched this material for Saccharibacteria using the method described by Bor *et*

al[18]. This led to the isolation of two strains, which exhibit distinct host specificity (Figure 1A).

Phylogenetic analysis using complete 16S rRNA gene sequences placed the two strains within

human oral subclades of the G1 clade of Saccharibacteria (Figure 1B,C)[21]. Based on these

assignments, we named our strains *Candidatus Nanosynbacter lyticus* ML1 (*Nl*) and *Ca.*

Southlakia epibionticum ML1 (*Se*). Complete genome sequences of the two strains indicated

they bear features typical of Saccharibacteria, such as genes associated with specialized secretion

systems, cell surface appendages, and competence, coupled with a lack of genes required for a

multitude of biosynthetic pathways and an overall reduced genome size (Figure 1D,E).

Genetic manipulation of Saccharibacteria via natural transformation

We sought to develop methods for reverse genetic analyses within Saccharibacteria.

Genes encoding the core components of the *com* DNA uptake system are conserved across CPR

phyla, including the Saccharibacteria[5]. These consist of ComEC, the central membrane DNA

conduit, DprA, a catalyst of Rec-mediated recombination, and ComFC, which binds ssDNA and

links import to recombination[22-24]. Com proteins function in concert with type IV pili, which are

also widely distributed in Saccharibacteria, and CPR bacteria more generally[5,23]. Given the lack

of nucleotide biosynthetic capability in CPR bacteria, it has been proposed that these systems

facilitate nucleotide acquisition[5]. Although the presence of the Com system is largely not

predictive of DNA transformation in a laboratory setting[25], we sought to test whether exogenous

DNA could be exploited for genetically manipulating Saccharibacteria.

As a first step toward assessing feasibility of genetics in Saccharibacteria, we searched

for antibiotics with convenient resistance determinants that potently inhibit the growth of *Se*

without impacting that of its preferred host, *Actinomyces israelii* F0345 (*Ai*). These experiments

revealed that across a wide range of concentrations, the aminoglycoside hygromycin fulfills

these criteria (Figure S1). Next, we designed and generated a linear cassette containing the

hygromycin resistance gene (*hph*) codon optimized for *Se* flanked by the promoter and

terminator regions of the TM7x *tuf* gene (elongation factor Tu), an open reading frame (ORF)

predicted to be highly expressed (Figure 2A). For the insertion of this cassette, we selected an

intergenic region located between two convergently transcribed ORFs, SEML1_0215 and

SEML1_0216, hereafter referred to as neutral site 1 (NS1). To promote homologous

recombination, approximately 1000 bp on either side of the insertion site were added to the 5′

and 3′ ends of our cassette.

To transform *Se*, we incubated *Se–Ai* co-cultures with 2.0 ng/μL of our linear cassette for

six hours before initiating selection with hygromycin (Figure 2B). Naïve host was added

concomitantly in order to permit the outgrowth of successfully transformed *Se* cells. Reasoning

that transformation may be inefficient, we passaged these co-cultures twice, at 48 hour intervals,

with continued hygromycin selection and the addition of naïve host. Cultures were then diluted

and plated to obtain colonies, where were selected and propagated with naïve host without

selection before genotyping (Figure 2B). The latter step was included in order to bottleneck the

Se population and facilitate the isolation of clonal populations. Remarkably, each *Se*-infected

culture we tested – accounting for the majority of colonies selected – contained our synthetic

cassette inserted at the expected location (Figure 2C). Whole genome sequencing confirmed

these PCR results and it further showed that cassette integration occurred without introducing

off-target mutations.

Quantitative PCR (qPCR) analysis of total *Se*, transformed *Se*, and *Ai* at regular intervals

during our transformation procedure demonstrated that approximately 0.2% of Se contain the

integrated cassette by the conclusion of the initial incubation period (Figure 2D). Though *Se*

levels remain low through the second passage under selection with hygromycin, all surviving *Se*

cells bear the cassette at this time point. In the final passage, the population of *Se* continues to

maintain the cassette and expands markedly, far surpassing levels of the host (Figure S2A). In

the absence of hygromycin, similar quantities of initially transformed *Se* are observed; however,

this small proportion fails to expand despite overall robust growth of *Se*. We observed similar

transformation behavior using lengths of DNA with homology to the insertion site flanking

regions as short as 221 bp (Figure S2A) and with as little as 0.02 ng/μL (Figure S2B). Finally, to

probe the generality of our methods, we identified a predicted neutral site within *Nl* and

subjected this second Saccharibacterial strain to an analogous transformation protocol.

Genotyping of transformed populations indicated cassette insertion at the desired location also

occurred within this strain (Figure S2C,D).

The ability to introduce heterologous DNA into *Se* has numerous foreseeable

applications, one of which is the expression of reporter genes that allow *Se* to be distinguished

and studied within the context of co-culture with their hosts. To explore this possibility, we

designed and generated NS1 insertion cassettes containing genes encoding nanoluciferase,

mCherry and GFP under the control of the *tuf* promoter and upstream of *hph* driven by a second

predicted strong promoter of TM7x, that of *rpsJ* (Figure 3A,B). Using our transformation

protocol, we obtained clonal populations of hygromycin-resistant *Se* containing each of these

genes. Luminescence assays and fluorescence microscopy demonstrated robust activity of each

reporter gene (Figure 3A,B). We did not detect their activity in host cells, indicating the

feasibility of achieving specific manipulation of *Se* in the context of a co-culture.

The expression of fluorescent proteins within *Se* provided the opportunity to visualize

CPR bacterial growth with extended, time-lapse fluorescence imaging. Over the course of 20 hrs,

both *Se* and *Ai* populations – deposited from co-cultures at low *Se*:*Ai* ratio onto an agar substrate

containing growth media – showed clear evidence of expansion (Videos S1-10). Apparent T4P-

mediated motility of *Se* was also observed, as reported by Xie *et al*[17]. Our time-lapse imaging

further captured features of the CPR lifecycle at unprecedented spatiotemporal resolution. For

instance, we could distinguish productive (*Se* growth supporting) versus non-productive (*Se*

adhered without concomitant growth) interactions, and directly measure their respective impact

on individual host cells (Figure 3C-E). Additionally, we observed productively adhered mother

cells producing small swarmer cell progeny via repeated polar budding, and the differentiation of

a subset of these progeny into mother cells (Videos S1-10). Altogether, these findings show that

natural transformation can be exploited to genetically manipulate CPR bacteria in a directed

manner and open a window into the distinctive biology of this largely unexplored group of

organisms.

Tn-seq of *Se* provides insights into the epibiotic lifestyle

The ability to genetically manipulate Saccharibacteria enables myriad avenues of

investigation. As a first step toward genetic dissection of the *Se* epibiotic relationship with *Ai*, we

conducted transposon-insertion sequencing (Tn-seq) within *Se* during growth on *Ai*. To this end,

we performed *in vitro* Tn5-based transposition on purified *Se* genomic DNA, repaired gaps as

described by Manoil and colleagues, and used sequencing to confirm high frequency,

homogenous insertion across the genome. This DNA was then used to transform *Se*, with slight

modifications from our basic transformation protocol (see Methods). Most notably, we elected to

increase the scale of the experiment to account for our previously measured transformation

efficiency and thereby avoid population bottlenecking following the onset of selection. We

collected four samples for Tn-seq analysis, an initial sample following the transformation and

recovery period (T0), and three additional samples representing the population after 48 hr serial

passages with the addition of naïve host (T1-3). Measurements of *Se* and *Ai* levels at each of

these timepoints revealed a precipitous drop in *Se* levels at T1 that is not observed at a similar

timepoint in transformations targeting NS1, suggesting that the large majority of transposon

insertions in *Se* are deleterious (Figure S3). Our measurements also showed that, as expected, *Ai*

levels drop relative to those of *Se* at later passages, such that by T3, *Se* outnumbers *Ai* by

approximately 50-fold.

Preliminary analysis of our Tn-seq data indicated that the T0 population of *Se* contained a

large number of unique insertions (9,996) relative to subsequent timepoints (T1, 1709; T2, 1887;

T3, 1699) despite similar sequencing depth (Table S1, Figure 4A). Furthermore, application of

the TRANSIT algorithm to the T0 data identified a small number of essential genes relative to

our expectations (905 *Se* genes; 56 essential by Gumbel or HMM)[26]. Taken together with our

measurement of *Se* and *Ai* population levels, we interpret these findings as evidence that little

selection occurred prior to T0 sampling. Therefore, we compared subsequent timepoints to T0

using TRANSIT resampling in order to assess gene level fitness contributions throughout the

experiment. In total, this approach identified 214 genes that demonstrate statistically significant

(multiple-comparison adjusted P value < 0.05) fitness contributions across all timepoints, with

selection yielding greater numbers at each subsequent sampling (T1, 222; T2, 252; T3, 275)

(Table S1).

The transposon insertion density we observed in our T1-3 samples is below the value

needed to obtain optimal results in TRANSIT. Given this potential caveat, we elected to further

process our data with ALDEx2, a program widely used for differential abundance analysis of

compositional data[27]. Overlaying genes with significantly differential insertion abundance

determined by ALDEx2 with those we obtained using TRANSIT revealed a group of 79 genes

that we deem unambiguously critical for *Se* growth on *Ai* in the context of our experimental

conditions (Table S1 and Figure 4A). This group includes genes encoding proteins that comprise

core cellular machinery such as DNA polymerase, RNA polymerase and the ribosome.

Interestingly, the group also includes non-canonical essential genes that likely encode specialized

functions that support the epibiotic lifestyle of CPR bacteria; examples include genes within a

T4P operon and several genes lacking homologs in non-CPR bacteria (Figure 4B).

If our Tn-seq findings provide an accurate measure of gene-level fitness contributions,

then those identified as critical for *Se* fitness via TRANSIT or TRANSIT and ALDEx2 should be

difficult to inactivate by targeted mutagenesis. Accordingly, to validate these data, we selected

four genes identified as significant contributors to fitness in one or both analyses, along with four

control genes, and designed constructs to replace each with an *hph* expression cassette. The

constructs shared equivalent length flanking sequences to enable the direct comparison of their

behavior in our transformation protocol. We assessed the fitness impact of inactivating each gene

by quantifying *Se* populations following transformation and outgrowth under selection with

hygromycin. For the control genes not predicted to contribute to fitness, we achieved robust

levels of *Se* growth by this time point, comparable to that achieved when introducing the *hph*

expression cassette at NS1 (Figure 4C). In contrast, in transformations targeting each of the four

predicted essential genes examined, *Se* failed to proliferate, consistent with the inactivation of

these genes strongly impacting fitness. Together, these findings provide confirmation that our

Tn-seq analysis successfully identified the relative fitness contributions of *Se* genes during co-

culture with *Ai.*

We noted that genes encoding homologs of *com* system components ComEC, ComF and

DprA were not among those genes defined as contributing significantly to *Se* fitness in our Tn-

seq study. This finding suggests that this DNA competence machinery is not required for

acquiring nucleotides to support *Se* growth, as had been previously suggested[5]. To determine

whether the *com* system of *Se* instead functions to mediate natural transformation, we generated

an *Se* strain in which the *comEC* ORF is replaced by the *hph* expression cassette (*Se*

Δ*comEC*::*hph*). To assess whether this mutation affects *Se* transformation, we measured the

efficiency of inserting a second, unmarked cassette at NS1. At the conclusion of the

transformation protocol, insertion at the NS1 site was only detectable in the wild-type

background, supporting the hypothesis that the *com* system mediates natural transformation in

this species (Figure 4D,E).

Banfield and colleagues previously leveraged the large number of CPR genomes

available from metagenomic and traditional genome sequencing datasets (n=2,321) to define

CPR-enriched protein families[5]. To gain insight into which of these contribute to fitness during

Se growth on *Ai*, we identified *Se* proteins belonging to these families, and cross-referenced hits

against the gene lists obtained from our Tn-seq experiment following TRANSIT or TRANSIT

and ALDEx2 analysis (Figure S4). Interestingly, we found that the representation of CPR-

enriched protein families among proteins critical for *Se* survival on *Ai*, 22% and 19%,

respectively, is considerably higher than their proportion within the overall *Se* proteome (12%)

(Table S1 and Figure S4). As many of these protein families have no predicted function, nor

orthologs outside of CPR bacteria, their future study is likely to reveal unique features of these

organisms.

Development of resources for dissecting Saccharibacteria–host cell interactions

With available genetic tools and gene-level fitness data, *Se* could become a tractable

model for the study of Saccharibacteria. However, comprehensive genotype–phenotype

dissection of the interaction between *Se* and its host remains limited by lack of equivalent tools

and data for *Ai*. As a first step towards bridging this gap, we obtained a complete genome

sequence for the host strain we employed, *A. israelii* F0345 (Figure S5A). This strain was

isolated from the human oral cavity as part of the Human Microbiome Project, but a genome

sequence had not been previously deposited in a public database. Although genetic tools have

not been developed for this organism, gene inactivation via the electroporation-mediated

introduction of suicide plasmid or linear gene inactivation constructs has been achieved in

related species[28,29]. The availability of genetic information for this strain should facilitate

development of selectable markers and counter selection strategies.

Automated annotation of the *Se* genome sequence failed to predict functions for the

proteins encoded by many ORFs (278 of 855), including 14/79 belonging to the conservative list

of essential genes we identified by Tn-seq (Table S1). To improve functional predictions

associated with ORFs in the *Se* genome, we applied a battery of sequence and structure-based

computational tools. ProtNLM is a natural language model trained on UniProt to predict protein

names given a sequence[30]. Applying ProtNLM to the *Se* proteome yielded functional annotations

for 337 of 855 proteins (ProtNLM score ≥ 0.5). This sequence-based annotation was further

improved by mapping each protein to Pfam domains using hmmscan, detecting Pfam domains in

an additional 270 proteins (see Methods for cutoffs), 245 of which could be assigned

functions[31,32].

To complement our sequence-based annotations, we took advantage of recent advances in

protein structure prediction to conduct genome-wide structure-based homology analyses of the *Se*

proteome[33,34]. Generation of structural models using AlphaFold (AF) relies on evolutionary

information extracted from multiple sequence alignments (MSAs)[34]. For nearly 25% of the *Se*

proteome (220 proteins), our initial MSAs based on HHblits searches of UniRef and BFD (the

default databases used by AF) were too shallow for high confidence structure prediction (<500

sequences post filtering, Figure 5A)[35-37]. To improve the MSA depth for these proteins, we

implemented a hidden Markov model (HMM)-based approach to identify and align additional

homologs for each protein from multiple metagenomic datasets. This resulted in considerably

deeper MSAs (>500 sequences) for an additional 9% of the proteome (Figure 5B). Using the

highest depth MSA obtained for each protein, we obtained AF models for >99% of the predicted

Se proteome. For comparison, we also computed AF models using the original 220 MSAs that

contained <500 sequences and compared the model confidence metric obtained using these and

the improved depth MSAs. Particularly for low confidence models obtained using MSAs generated

with the default databases (average pLDDT<50), we found that the use of the deeper MSAs for

structural prediction resulted in substantial model confidence improvement (Figure 5C). The

structure model improvements enabled by using extensive metagenomic databases for MSA

generation were further underscored by structural homology search results obtained using

Foldseek (FS)[38]. For some proteins, the improved structural models led to the identification of

structural homologs for proteins that initially had no FS matches passing our cutoffs, while for

others, the changes in the overall predicted structure led to the identification of different and more

closely aligning top matches (Figure 5D,E, Figure S5B-D). In total, using FS we were able to

identify similar structures for 88% of modeled proteins (748/852).

Many of the AF models with similar structure identified for *Se* proteins are

uncharacterized (12%), providing little information regarding function. Moreover, unrelated

proteins may share similar structure due to convergence, and additional evidence is usually

needed to assess the homologous relationships. Domain Parser for AF Models (DPAM) is a

recently developed tool that parses structural domains from AF modeled structures, and

integrates both structure and sequence searches to map protein domains to Evolutionary

Classification of protein Domains (ECOD)[39,40]. Using DPAM, we detected homologous ECOD

domains for 80% of predicted *Se* proteins. In supplementary material accompanying this report,

we provide alignments for each mapped *Se* protein domain, along with links to corresponding

Protein Data Bank identifiers (PDB IDs) and ECOD classifications

(https://conglab.swmed.edu/ECOD_Se/Se_ECOD.html). In total, these predictions provide a

foundation for future use of the *Se–Ai* system as a model for understanding CPR bacteria.

Discussion

324   Scientists have been aware of CPR bacteria in environmental samples for many years, yet

our understanding of this group of organisms has lagged[4]. One major challenge is their apparent

strict requirement for host bacteria, the identity of which cannot currently be determined a priori,

thus adding significant complexity to CPR isolation[7,18,41,42]. It is foreseeable that methods

bringing to bear both experimental and computational approaches could provide predictions of

host–epibiont associations in the future. Here we addressed a second impediment for studying

CPR bacteria – a lack of genetic tools for their manipulation. Indeed, despite streamlined

methods for isolating Saccharibacteria on host Actinobacteria that have been employed by

several laboratories, this latter challenge has to-date prohibited a molecular dissection of their

epibiotic lifestyle[16,17,42]. Our discovery that natural transformation can facilitate targeted

mutagenesis in Saccharibacteria opens the door to direct interrogation of genotype–phenotype

relationships within these CPR bacteria; however, challenges remain. The most fundamental

hurdle is that factors required for the Saccharibacteria–host cell interaction are also required for

viability. In analogous situations, researchers overcome this using assorted conditional

inactivation strategies (e.g. modulated expression, temperature sensitivity, inducible

degradation). While it is conceivable that such methods could be implemented in

Saccharibacteria, a powerful approach in this system may involve exploiting host genetics to

identify gene interactions that suppress otherwise lethal mutations in the epibiont.

The small genome of Saccharibacteria stands in contrast to the apparent complexity of

their lifecycle. Xie *et al.* proposed a model for the epibiotic growth of Saccharibacteria that

consists of four stages: T4P-mediated infection, growth, bud formation and asymmetric

division[17]. Our time-lapse microscopy generally supports this model; however, it further revealed

facets of the Saccharibacteria–host interaction not previously captured. While the

aforementioned model implies uniform progression, our data suggest the lifecycle is more

complex. We find that only a small subset of *Se* achieve productive infections, and that after an

extended enlargement period, these cells rapidly bud a large number of highly motile swarmer

cell progeny (>10 in a 20 hr period observed in many instances). Quantitative analyses of our

data additionally permitted us to link productive and non-productive *Se* infections with

corresponding host cell outcomes. We observed a striking inverse relationship between *Se*

growth and that of *Ai*; cells productively infected with as few as one *Se* not only failed to divide,

but diminished in size, whereas those infected unproductively by multiple *Se* readily proliferated.

Our microscopy observations suggests a potential division of labor within *Se* populations,

wherein one subpopulation is devoted to reproduction, and a second, motile subpopulation

searches for new compatible host cells. Operating under the assumption that the

Saccharibacteria–Actinobacteria relationship is chiefly one of parasitism, which we maintain is

not fully resolved in the literature, then the relevant prevailing evolutionary theory predicts that

Se fitness is dependent on both the rate at which it causes new infections and the duration of

those infections[43]. The reproductive strategy of *Se* appears consistent with this framework;

release of progeny cells destined to infect other host cells tempers host cell burden and prolongs

the infection, while concurrently maximizing *Se* reproduction. A similar outcome could be

achieved if *Se* employed active mechanisms to block superinfection, reminiscent of those utilized

by phage; however, swarmer cell production has the added advantage of increasing *Se* dispersal

in the face of low host cell densities[44]. Interestingly, most nascent swarmer cells we observed do

not themselves establish a productive infection within our 20 hr observation period, despite close

proximity and apparent adherence to host cells. It may be that the time required to establish a

productive interaction varies, and many of these cells would go on to become productive. It is

also conceivable that infected host cells defend against secondary infections, perhaps even

inducing a defensive state in neighboring kin cells. Future studies coupling quantitative

microscopy with genetics to further dissect the *Se–Ai* interaction will no doubt address these

questions and shed light on this fascinating interphyla relationship.

Pili are a basal feature of CPR bacteria and our Tn-seq results highlight the critical role

they play for these organisms[7,17,45]. Among the 15 genes that are both enriched in CPR and that

we conservatively deemed essential in *Se*, six are predicted T4P components (Table S1). A prior

study utilizing quercetin, a small molecule inhibitor of pilus retraction, reported that T4P are

important for host attachment and twitching motility in a second Saccharibacterial species[17]. In

naturally competent bacteria, T4P work in concert with the *com* system to bind and mediate the

uptake of extracellular DNA[46,47]. Our observation that *Se* cells lacking *comEC* are viable

suggests that the essentiality of T4P is unrelated to its role in competence. *Se* lacks *de novo*

nucleotide synthesis pathways, thus the organism presumably possesses an essential DNA or

nucleotide uptake pathway that operates independently of the *com* system. Whether T4P

participate in such an alternative DNA uptake mechanism remains unknown.

The arginine deiminase system (ADS) is an ATP generating catabolic pathway prevalent

in mammalian-adapted Saccharibacteria species[48]. A prior study demonstrated that arginine

supplementation to *N. lyticus* TM7x–*A. odontolyticus* co-cultures permits acid neutralization via

ammonia production, supporting the viability of both strains during acid stress[48]. The authors of

this study put forth a model in which hosts lacking ADS benefit from and are specifically

permissive to colonization by Saccharibacteria containing the pathway. In *Se*, arginine deiminase

(ArcA) is a fusion of two proteins, ArcA and ornithine carbamoyltransferase (ArcB); *arcA* is

among our conservative list of genes deemed essential for *Se* growth. Inactivation of genes

encoding the remaining ADS components, carbamate kinase ArcC and arginine/ornithine

antiporter ArcDE*,* also yielded significant *Se* fitness defects by one or more metric (Table S1).

Interestingly, in contrast to the proposed model for ADS function in Saccharibacterial–host cell

interactions, we find intact homologs of each ADS gene in *Ai*. Taken together with our Tn-seq

data, our results suggest that in some co-culture pairs, the organisms may compete for arginine.

However, the mechanism by which the pathway influences Saccharibacteria–host cell dynamics

in vivo, including specific niches in the oral cavity, remains an open question[48].

We anticipate that the methods and genetic tools presented here will facilitate molecular

level characterization of Saccharibacteria and CPR bacteria more broadly. In this regard, one key

question is the extent to which natural competence is active and able to be similarly exploited

across the CPR. The ability to manipulate CPR bacteria outside of Saccharibacteria, particularly

those with phylogenetically distinct hosts and inhabiting diverse niches, should aid in elucidating

the core requirements of the epibiotic lifestyle. Regardless of the precise methods utilized,

genetic manipulation of CPR bacteria will open the door to phenotypic studies of the rich

reserves of microbial dark matter these organisms contain, potentially revealing unprecedented

biological mechanisms.

Acknowledgements

The authors wish to thank Jeffrey McLean, Simon Dove and Joshua Woodward for insightful comments on the manuscript, and members of the Mougous laboratory for helpful suggestions. This work was supported by grants from the NIH (R01AI128215 and R01AI141953 to N.S.B.), the National Science Foundation (MCB-2105570 to N.S.B. and S.T., and IIBR-2042948 and IOS-2050550 to N.S.B.), the Department of Defense, Defense Threat Reduction Agency (HDTRA1-21-1-0007 [DTRA J5B] to D.B.), the Bill and Melinda Gates Foundation (OPP1156262 to DB), and the Welch Foundation (I-2095-20220331). Q.C. is a Southwestern Medical Foundation endowed scholar. J.D.M. and D.B. are HHMI Investigators, and J.D.M. is supported by the Lynn M. and Michael D. Garvey Endowed Chair at the University of Washington.

Declaration of interests

The authors declare no competing interests.

Author contributions

L.A.G, A.L., S.B.P and J.D.M conceived the study. Y.W., L.A.G., A.L., I.R.H, S.T., K.J.C., Q.C., N.S.B., S.B.P. and J.D.M. designed the study. Y.W., L.A.G., P.A.A., A.L., and K.J.C. performed experiments. Y.W., L.A.G., I.R.H., S.T., K.J.C., M.L.A-O., Y.L., M.C.R., and Q.C. processed, analyzed and visualized the data. Y.W., L.A.G., I.R.H., S.B.P. and J.D.M wrote the manuscript. D.B., N.S.B., S.B.P. and J.D.M. provided supervision. Q.C., D.B., N.S.B. and J.D.M funded the study. All authors contributed to manuscript editing and support the conclusions.

Methods

Strains, media and growth conditions

Saccharibacteria strains used in this study include *Southlakia epibionticum* ML1 and *Nanosynbacter lyticus* ML1, both isolated in this study. Host bacteria strains used include *Actinomyces israelii* F0345, *A. odontolyticus* F0309, *A. urogenitalis* S6-C4, *Actinomyces sp.* F0386 and *Propionibacterium propionicum* F0230 (*Pp*)[49-51]. Host bacteria mono-cultures, *Se-Ai* co-cultures and *Nl-Pp* co-cultures were routinely grown statically under an atmosphere of air supplemented with 5% CO_2 at 37°C or anaerobically with shaking at 37°C using the GasPak EZ Anaerobe Container System with Indicator (BD 260626 and 260001) in TSY media (30 g/L tryptic soy and 5 g/L yeast extract) or TSYR media (TSY media supplemented with 10 mM arginine), or anaerobically on TSBY agar plates (30 g/L tryptic soy, 5 g/L yeast extract, 15 g/L agar, supplemented with 5% (v/v) horse blood). For selection of hygromycin-resistant Saccharibacteria, hygromycin was used at 150 µg/mL. Saccharibacteria-host co-cultures were stored at -80°C in TSY or TSYR supplemented with 10% (v/v) Dimethylsulfoxide (DMSO).

Isolation of Saccharibacteria strains *Se* ML1 and *Nl* ML1

Isolation of Saccharibacteria strains in co-culture with host bacteria was carried out as essentially as previously described[42,52]. Anonymous volunteers aged over 18 years provided oral samples for Saccharibacteria isolation. Supragingival plaque samples were collected with toothpicks and dispersed in 1 mL of Maximum Recovery Diluent (MRD; Peptone 1.0 g/L, Sodium Chloride 8.5 g/L, pH 7.0) buffer. 5 mL of saliva was collected by voluntary expectoration into sterile 50 mL conical tubes. All saliva samples and plaque samples were then pooled into 20 mL MRD. Pooled samples were then vigorously resuspended by vortexing and

filtered with a 0.22 μm filter, and the flowthrough was collected. Residual bacteria in the sample

tube and filter were washed once with 10 mL MRD, filtered again, and combined with the

previous flowthrough. Filtrates were collected by centrifuging at 60,000 rcf for 1 hr at 4°C.

Supernatant was removed, and pellets were resuspended in 1 mL MRD. The presence of

Saccharibacteria in these samples were confirmed by PCR. The collected Saccharibacteria were

then added to mono-cultures of a panel of five potential host species (*A. odontolyticus, A.*

urogenitalis, Actinomyces sp. F0386, *Ai,* and *Pp*) and the cultures were passaged every 24 or 48

hr. The presence of Saccharibacteria cells in the final cultures was confirmed by qPCR with

universal Saccharibacteria primers and microscopic imaging. The cultures were then streaked on

TSY or TSBY agar, isolated colonies of host bacteria were tested for Saccharibacteria by PCR,

and positive colonies were re-cultured in liquid medium and stored as clonal co-cultures.

To purify Saccharibacteria from co-cultures, the co-culture was centrifuged at 3,000 rcf

for 5 min at room temperature. Supernatant containing free-floating Saccharibacteria was

collected and filtered twice using a pre-sterilized 5 μm mixed cellulose esters (MCE) filter.

Filtrate was centrifuged at 15,000 rcf for 30 min at room temperature to pellet Saccharibacteria.

The resulting pellet was resuspended in TSYR supplemented with 10% (v/v) DMSO and stored

at -80°C.

Saccharibacteria host compatibility testing

Purified populations of *Se* and *Nl* for host compatibility testing were generated by

growing 100 ml co-cultures of each with their respective host strains according to the standard

protocol described above. Co-cultures were passed through a 0.45 μm SFCA filter to remove

host cells then spun at 80,000 rcf for 20 min to pellet Saccharibacteria. Supernatant was removed

and cell pellets were resuspended in 1 mL fresh medium. Purified Saccharibacteria cells were

then add to $OD_{600} = 0.2$ cultures of *Ai* and *Pp* (compatible host for *Nl*) at an MOI of 0.2. Cultures

were then incubated at 37°C under 5% CO_2 for up to 72 hr, and populations of *Se* and *Nl* were

monitored over time using qPCR.

Phylogenetic analysis of 16S rRNA gene sequences

The Saccharibacterial 16S rRNA phylogeny was generated by extracting Saccharibacteria

sequences in HOMD and reported in McLean *et al.* and aligning in SILVA, collecting nearest

neighbors (n=20, 89% cutoff)[11,53,54]. Unique sequences were then extracted in Geneious, the

resulting sequences were re-align using the SILVA SINA aligner, positions with 10% gaps were

masked, and a tree was generated using RAXML[55,56].

Whole genome sequencing

Genomic DNA of *Se* ML1, *Nl* ML1and *Ai* F0345 was isolated using the Wizard HMW

DNA Extraction kit (Promega). Sequencing was performed on Illumina iSeq and MiSeq and

Oxford Nanopore (ONT) MinION instruments after standard sequencing library preparation

protocols (Illumina and Oxford Nanopore). De-novo assemblies were generated using the

Trycyler pipeline[57]. Specifically, ONT long reads were filtered using Filtlong v0.2.1

(https://github.com/rrwick/Filtlong) with Illumina reference reads, using --keep_percent 95 to

retain approximately 95% of the reads. Long reads were then subsampled into 12 bins and

assembled into 12 assemblies using the Flye v2.9

(https://github.com/fenderglass/Flye/releases/tag/2.9. Accessed 4 October 2021), Raven v1.8.1,

Miniasm v0.3r179[58] and Minipolish v0.1.3

(https://github.com/rrwick/Minipolish/blob/main/miniasm_and_minipolish.sh) assemblers.

Assembly contigs were manually curated and then reconciled using Trycycler v0.5.2[57]. A

consensus assembly for each bacterium was generated and then polished with Illumina short

reads using Polypolish v0.5.0[57] and POLCA from MaSuRCA v4.0.9[59]. Initial annotations were

generated using PROKKA v1.14.5[60].

To sequence clonal transformed *Se* (see Results and below), genomic DNA was isolated

from frozen pellets of purified *Se* by re-suspending in Buffer PB (Qiagen) to a total volume of

500 μL, adding 250 μL of 0.1mm zirconia/silica beads (BioSpec Catalog # 11079101z), 250 μL

of 20% SDS, and 550 μL of phenol:chloroform:IAA (25:24:1) (Invitrogen Catalog #15593-031),

and bead-beating in a Mini-BeadBeater-16 (Biospec Model 607) with settings 3450 RPM, 115V,

10A, and ½ HP, for four, 30-second intervals, each followed by cooling on ice for 1 minute.

Purification of the DNA was performed by applying the aqueous phase directly to a DNeasy

Blood & Tissue Prep Kit (Qiagen) purification column and following the recommended protocol

for washing and elution. Sequencing was performed on an Illumina iSeq using standard library

preparation protocols (Illumina). Reads were mapped to the assembled *Se* ML1 genome using

minimap2 and variants were called using LoFreq v2.

Measuring hygromycin sensitivities of *Ai* and *Se*

Hygromycin sensitivity of *Ai* and *Se* was measured in liquid cocultures. Duplicate co-

cultures were initiated by mixing purified *Se* with *Ai* at an OD_{600} of 0.2 in TSY and at an

approximate cellular ratio of 1:2 (*Se*:*Ai*). The co-cultures were divided into multiple aliquots in

96-well culture plates and hygromycin was added to the final concentrations shown in Figure S1.

The plates were covered with Breath-Easy sealing membrane (Sigma Z380059) and incubated

without lids at 37°C with 5% CO_2. At 1 day and 3 days, the cells within individual wells were

pelleted (>15 min at >15,000 rcf) and stored at -20°C for later genomic DNA isolation and

qPCR-based quantification of *Se* and *Ai* (see below).

Design and generation of cassettes for heterologous gene expression in *Se* and *Nl*

Cassettes for heterologous gene expression in *Se* and *Nl* were designed by appending

promoter and terminator sequences from the TM7x genome[8] to the 5' and 3' ends, respectively,

of ORFs codon optimized for *Se*. The promoter and terminator elements were sourced from

TM7x rather than *Se* to reduce the likelihood of off-target integration at corresponding *Se* loci.

The TM7x promoters were chosen from genes expected to be highly and constitutively

expressed, *tuf* and *rpsJ*. *Se* codon usage was calculated using the Dynamic Codon Biaser

(DCB)[61] Heterologous ORFs were optimized to match relative codon usage frequencies found in

Se, but omitting codons with less than 10% usage. Heterologous genes utilized included *hph*

(hygromycin B phosphotransferase) from *Streptomyces hygroscopicus*, superfolder GFP

(www.fpbase.org/protein/superfolder-gfp/), mCherry2 (www.fpbase.org/protein/mcherry2/) and

NanoLuc Luciferase[62,63]. Table S2 reports the composition and complete sequences of the

designed cassettes. Cassettes were obtained as gBlocks from Integrated DNA Technologies, Inc

(IDT).

Linear fragments used for transformations were generated by adding *Se* or *Nl* genomic

sequences corresponding to the targeted insertion or allelic replacement sites to the left and right

sides of a heterologous gene expression cassette or of two cassettes joined together. Overlap

extension PCR was used to join gBlock cassettes and genomic fragments, the latter of which

were individually amplified from *Se* or *Nl* genomic DNA. In some cases, complete fragments

including genomic sequences were obtained as gBlocks from IDT. Fragments used for flank-

length tests were generated by amplification using larger fragments as templates followed by gel-

purification. Table S2 reports the composition and complete sequences of the fragments utilized

and primers used are listed in Table S3.

Genetic transformation of *Se* and *Nl* with targeted insertion and allelic replacement

constructs

To prepare *Se-Ai* or *Nl-Pp* co-cultures for transformation, 1-mL aliquots of a previously

frozen co-culture (see methods on co-culturing) were thawed on ice and added to 9 mL of TSY

supplemented with recently passaged *Ai* to a final OD_{600} of 0.2, incubated for 2 d at 37°C in 5%

CO_2, then enlarged by the addition of 10 volumes of TSY and incubated anaerobically using the

GasPak EZ Anaerobe Container System with Indicator (BD 260626 and 260001) at 37°C for 2

days with shaking at 160 rpm.

For transformation of *Se* or *Nl* with linear targeted insertion constructs, 0.22 - 0.3 mL

aliquots of prepared co-culture were incubated with transforming DNA for 6 h at 37°C in 5%

CO_2 in culture tubes. The mixtures were subsequently enlarged by the addition of TSYR to

approximately 5 mL and supplemented with recently passaged *Ai* (for *Se*) or *Pp* (for *Nl*) to a final

OD_{600} of approximately 0.06. This time point was designated "TF" (time zero after

transformation). After removal of a sample for later qPCR analysis, the cultures were divided

and one half was supplemented with hygromycin to 150 μg/ml. The cultures were then incubated

at 37°C in 5% CO_2 without agitation for 2-3 days. This time point was designated "P1" (end of

initial passage). Cultures were subsequently serially passaged as many as four times by five-fold

dilution into fresh TSYR supplemented with *Ai* (for *Se*) or *Pp* (for *Nl*) to a final OD_{600} of 0.06

and, for the hygromycin containing cultures, with additional hygromycin to 150 μg/ml. These

passages (designated "P2", "P3", etc.) were each incubated for 2-3 days at 37°C in 5% CO_2

without agitation. Cultures were sampled at each passage by removing and pelleting (>15 min at

>15,000 rcf) of 0.1 to 1 ml, and the pellets were stored at -20°C for later qPCR analysis.

qPCR assays

We employed qPCR for quantification of Saccharibacteria and host populations in

cocultures, as well as to monitor *Se* transformation. For these assays, genomic DNA was isolated

from frozen co-culture pellets using Instagene matrix (Bio-Rad) and quantitative PCR (qPCR)

was performed using a CFX Connect Real-Time PCR Detection System (Bio-Rad). To quantify

Se and *Ai*, amplification employed primers targeting *Se uvrB* or the 16S rRNA gene of *Ai* and

was performed in 20 μL reactions with 1X SsoAdvanced Universal SYBR Green Supermix (Bio-

Rad), 300 nM each primer and 4 μL template DNA. Thermocycling conditions were 95°C for 5

min followed by 35-40 cycles of 95°C for 20 s, 60°C for 30 s and a read of fluorescence. To

quantify transformed *Se* (insertions at NS1), amplification employed primer pairs targeting the

insertion sequence and sequence adjacent to NS1 but outside of the linear transformation

construct arms and was performed in 20 μL reactions with 1X Phusion HF Buffer, 0.2 mM

dNTPs, 250 nM each primer, 0.5X SYBR Green I (ThermoFisher), 4 μL template DNA and 0.4

units of Phusion High-Fidelity DNA Polymerase (NEB). Thermocycling conditions were 98°C

for 90 s followed by 35-40 cycles of 98°C for 15 s, 70°C for 20 s, 72°C for 90 s and a read of

fluorescence. Melt curves were performed following each amplification to verify product

homogeneity. Absolute target sequence abundance was determined by comparison to duplicate

standard curve reactions performed with each amplification. The standard curve templates were

generated by serial dilution of previously amplified and gel-purified products quantified by Qubit

(ThermoFisher). Primer sequences employed in qPCR reactions are provided in Table S2.

Isolation of isogenic mutant co-cultures

To obtain co-cultures with isogenic mutant (transformed) *Se*, transformation mixtures

grown for three passages under hygromycin selection were serially diluted, plated on TSBY agar

without antibiotic and incubated for 7 days anaerobically. Isolated colonies (representing *Ai*

potentially colonized with *Se*) were picked into 0.2 mL TSYR supplemented with *Ai* at a final

OD_{600} of 0.2 and incubated at 37C in 5% CO_2 for 2-4 days, then screened for the presence of *Se*

and for the mutant (genomic insertion) and wild-type (no insertion) alleles by PCR (e.g., Figure

2A). Cultures containing pure mutant *Se* were frozen and/or further propagated for purification

of the *Se* for WGS (see above).

Nanoluciferase assay

Nanoluciferase assays were performed using isogenic wild-type *Se-Ai* co-cultures and *Se*

NS1::*nluc-hph2-Ai* co-cultures grown statically under an atmosphere of air supplemented with

5% CO_2 at 37°C in TSY media for three passages as described above. To separate *Se* from host

cells, 20 mL of each co-cultures was passed through a 0.45 μM SCFM filter and the resulting

filtrate was spun at 80,000 rcf for 30 min to pellet *Se*, supernatant was removed, and the pelleted

Se was resuspended in 750 μL TSY. 100 μL of these purified *Se* cells or 100 μL of the

corresponding *Se-Ai* cocultures were mixed with 100 μL Nano-Glo Luciferase assay reagent

(50:1 mixture of substrate:buffer, Promega N1110) in a 96-well plate. Luminescence signal

indicative of nanoluciferase activity was detected using a Cytation 2 plate reader. Luminescence

signal was later normalized by *Se* abundance in co-cultures and filtrates as measured by qPCR

using *Se*-specific primers as described above.

Microscopy

Imaging was performed on a Nikon Eclipse Ti-E wide-field epi-fluorescence microscope,

equipped with a sCMOS camera (Hamamatsu) and X-cite LED for fluorescence imaging. We

imaged through a Nikon Plan Apo λ 60X 1.4 NA oil-immersion Ph3 objective. The microscope

was controlled by NIS-Elements 3.30.02. Isogenic *Se* NS1::*mcherry-hph2-Ai* co-cultures and *Se*

NS1::*sfgfp-hph2-Ai* co-cultures were grown statically under an atmosphere of air supplemented

with 5% CO_2 at 37°C in TSYR media as described above. Cell samples were spotted on a 3%

(w/v) agarose pad made with TSYR media supplemented with 0.4% glucose placed on a

microscope slide. The microscope chamber was heated to 37°C for time-lapse experiments.

Time-lapse images were aligned and segmented with Omnipose according to the phase contrast

channel and the published bact_phase_omni model[64]. Masks were manually linked and corrected

for segmentation errors in regions of *Ai* cell overlap using Napari. Regions corresponding

to *Se* were also removed to accurately track host *Ai* growth alone. For visualization purposes,

fluorescence intensity was gamma-corrected to normalize background levels on a frame-by-

frame basis while not distorting the *Se* signal. No bleaching correction was implemented. Figure

3C-E and Videos S1-10 were generated using Python and annotated in Adobe Premiere.

Mutagenesis of Sac1a genomic DNA and transposon mutant library generation

The transposon used for *in vitro* transposition (here named T36) was generated by

amplification of the *hph1* insertion cassette using the 5'-phosphorylated primers T36-ampF and

T36-ampR, which add required 19-bp Tn5 mosaic end sequences to each end of the amplicon,

followed by purification using the Qiagen PCR Purification Kit with elution in TE. *In vitro*

transposition was performed in multiple 45-μl reactions containing 1.8 μg *Se* genomic DNA, 290

ng purified T36, 1X EZ-Tn5 Reaction Buffer and 1.2 U EZ-Tn*5* Transposase (Biosearch

Technologies) with incubation for 2 hr at 37 °C. The reactions were stopped, DNA was ethanol

precipitated, and gaps were repaired as described[65]. Following gap repair, the DNA was purified

using Qiagen PCR Purification Kit with elution in water after the columns were washed twice

with Buffer PE.

Transformation of *Se* to generate a transposon mutant library was performed similarly to

the procedure described above for transformation with targeted insertion fragments, but at a

larger scale and with modifications. Specifically, the initial 6 hr incubation contained 4.0 μg of

in vitro transposon-mutagenized genomic DNA and 15 mL of *Ai-Se* co-culture. This co- culture

was enlargement by adding 135 mL of TSYR supplemented with *Ai*, then incubated for two

additional hours before hygromycin addition. Concomitantly with hygromycin addition, $MgCl_2$

(to 1 mM) and benzonase (Sigma) (to 25 U/mL) were added to degrade extracellular DNA, and

the cultures were supplemented with benzonase similarly approximately every 24 h during

passaging. After the initial 2-day growth passage (ending at time point T0), the culture was

serially passaged four additional times by dilution of 30 mL culture into 120 mL in TSYR

supplemented with *Ai* at OD_{600} of 0.067. These passages were each incubated for 2 days at 37°C

in 5% CO_2. Samples for Tn-seq analysis were taken at the end of the initial passage (60 mL, T0)

and at the end passages 1, 2 and 4 (80 mL, T1-T3). These samples were treated with additional

benzonase (50 U/mL) for 30 min, then EDTA was added to 15 mM and pellets were collected by

centrifugation for 30 min at 15,000 rcf and stored at -80°C.

Tn-seq library preparation and sequencing

Genomic DNA was isolated from frozen co-culture pellets using the bead-beating method

described above (section on whole genome sequencing). Tn-seq libraries were prepared by the

C-Tailing method as described[66]. The transposon-specific primers utilized are listed in Table S3.

Sequencing was performed in multiplex as 50-bp single-end reads on an Illumina MiniSeq with

25-40% PhiX spike-in.

Custom scripts (https://github.com/lg9/Tn-seq[65,66]) were used to process the Illumina

reads. In brief, reads were first filtered for those displaying transposon end sequence as their

initial bases (the sequencing primer was designed to anneal six bases from the end of the

transposon). These reads were then mapped to the *Se* genome after removing the transposon end

sequences. Reads per unique mapping position and orientation were tallied and read counts per

gene were calculated by summing reads from all unique sites within a given gene.

Se **gene fitness and essentiality analysis**

We first checked the saturation level (i.e., number of insertions detected) of the Tn-seq

library sampled along four time points (T0-T3). Because the T0 sample had the highest number

of unique insertions (9,996), we initially focused on this time point to identify essential genes in

the *Se* genome. Gene essentiality was determined based on the number of insertions detected in

each gene (using the GUMBEL method, part of the TRANSIT tool), or alternatively by taking

into account the read counts per insertion(s) in each gene (using the Hidden Markov Model

method, HMM, also part of TRANSIT)[26]. Default parameters were used for both GUMBEL and

HMM analyses. To complement these analyses and to identify genes whose disruption was

deleterious for *Se* in later time points, we used a comparative approach with respect to T0. The

comparative approach was necessary due to the significant reduction in the number of unique

insertions detected at T1-T3 (the average number of insertions for these time points was 1,765),

which suggested a strong selection pressure over time. Thus, the distribution of insertions at T1,

T2 or T3 were independently compared with T0. We detected genes associated with

differentially abundant insertions at T1-T3 vs. T0 using TRANSIT and ALDEx2[26,27]. For the

TRANSIT-based analysis, we used the Resampling method to identify differentially abundant

genes between the two time points being compared. Genes with TRANSIT-estimated adjusted p-

value < 0.05 were considered as differentially abundant (this gene set would include under- and

over-represented genes at the late time points with respect to T0). ALDEx2 takes into account

the compositional nature of sequencing data generated during selection assays (i.e., competitive

growth mutant assays). Given the lack of replicates (expected in the standard ALDEx2 pipeline)

in our data, we adapted the ALDEx2 workflow by using the aldex.clr function available in the

ALDEx2 R package to generate Dirichlet distributions of centered-log ratio (CLR) transformed

abundances for any pair of samples being compared (e.g., T1 vs. T0) [27]. We used 1,000 Monte

Carlo instances to generate the Dirichlet distributions. Then, we computed the median CLR

value per gene for each sample. Finally, we computed a delta score similar to ALDEx2 original

"dif.btw" by computing gene wise median difference between the two samples being compared

(e.g., T1-T0). Genes with delta scores < -1.5 were considered under-represented in the late time

points (with respect to T0), suggesting that insertions that disrupt those genes were deleterious

for Se. To define a meaningful threshold for our delta score and confirm the performance of our

ALDEx2-based analysis, we leveraged barcode sequencing data we previously generated for

competition growth assays of an antibiotic-treated Escherichia coli pooled single-gene deletion

library[67]. This benchmarking dataset included four time points (T0-T3), each one with four

replicates. For the T1 vs. T0 and T2 vs. T0 comparisons we applied our strategy for each

individual replicate and then evaluated the overlap with the original results obtained when

considering all four replicates. We found that indeed the set of genes with absolute delta score >

1.5 for single replicate analyses significantly overlap with the set of genes in the analysis that

included all replicates. Three out of the four single replicate analyses had hypergeometric test p-

values < 0.05 for T1 vs. T0 (two with p-values < 0.005). Moreover, all single replicate analyses

for T2 vs T0 had hypergeometric test p-values < 1E-202.

Transformations to validate gene essentiality predictions

To validate Tn-seq essentiality predictions, we designed constructs to replace target genes

with *hph*. Each construct contained 335-bp sequences targeting the desired site flanking the *hph*

expression cassette described above. 100 ng synthesize linear DNA or water were added to 300

μL wild-type *Se-Ai* co-cultures (4 biological replicates per condition) prepared as described

above. Transformation and subsequent passaging were performed as described above. At the end

of passage 3, 1 mL of each transformed co-culture was treated with 1 mM $MgCl_2$ and 25 U/mL

benzonase for 30 min at 37°C to remove extracellular DNA. 15 mM EDTA was then added to

stop the benzonase reaction, samples were centrifuged at 21,000 rcf for 1 hr and the resulting

pellets were stored at -80°C. *Se* abundance was then measured by qPCR as described above.

Generation and transformation efficiency of *Se* Δ*comEC::hph*

A clonal population of *Se* Δ*comEC::hph* was generated by replacing *comEC* with our9

hygromycin resistance cassette following the transformation and clonal mutant isolation

protrocols described above. To assess *Se* Δ*comEC::hph* transformability, 50 ng of synthesized

linear gene product for inserting unmarked *sfGFP* at NS1 (with 545 bp flanking regions) was

used to transform 150 μL isogenic wild-type or Δ*comEC Se-Ai* co-cultures (4 biological

replicates per condition). After 6.5 h of incubation between DNA and co-cultures at 37°C, the

entire transformation culture was treated with 1 mM $MgCl_2$ and 25 U/mL benzonase for 30 min

at 37°C to degrade extracellular DNA. 15 mM EDTA was added to stop the benzonase reaction

and the samples were centrifuged at 21,000 rcf for 1 hr and the resulting pellets were stored at -

80°C. We then used qPCR to measure the abundance of transformed and total *Se* using NS1-

sfGFP or *uvrB-* specific primers as described above.

CPR-enriched protein families analysis

To assess the protein family distribution of the *Se* and *Nl* genomes, we collected amino

acid sequences for 22,977 protein clusters (protein families) from Meheust *et al.*[5]. These protein

families were built through their occurrences across at least 5 distinct non-redundant or draft

CPR bacteria, non-CPR bacteria and Archaeal genomes. The presence or absence of any given

family in *Se* or *Nl* were determined through all-vs-all protein sequence search of every protein in

the genome against the database of protein family sequences. Protein sequence search was

performed by using the MMseqs2 (version: 14-7e284) algorithm with the following parameters:

module: easy-search; sensitivity (-s): 7.5; alignment coverage (-c): 0.5; greedy-best-hits: 1. The

resulting presence or absence matrix was combined with the core family matrix of 921 protein

families across 2890 genomes (collected from Meheust *et al.*[5]) for clustering analysis. A Jaccard

distance based on the presence or absence of core families across *Se*, *Nl* and 2890 other genomes

was calculated in R by using proxyC package (version: 0.3.3). Agglomerative hierarchical

clustering was performed by using cluster package (version: 2.1.4) in R with the "wand" method.

A hierarchical clustering heatmap was built using complexHeatmap package in R (version:

2.14.0) by plotting presence (black) or absence (white) of protein families (columns) in a given

genome (rows). Enrichment or depletion of core protein family clusters in *Se* and *Nl* was

performed using Fisher's exact test in R with input table of presence or absence of protein

families in the *Se* and *Nl* genomes. A threshold of Benjamini-Hochberg corrected p-values of <

5e-05 was used to determine the enrichment or depletion.

Sequence-based gene annotation of the *Se* genome

765   We used ProtNLM (https://github.com/google-research/google-

research/tree/master/protnlm) to predict the function annotations of proteins in the *Se* proteome

from sequence[30]. A prediction of score ≥0.5 corresponds to >70% accuracy for proteins without

close UniRef50 matches and ~95% accuracy for proteins with close matches. We thus

considered the 337 proteins with score ≥0.5 to have confident annotations (Table S1).

Additionally, we mapped each protein to Pfam domains (Nov, 2021 release v35.0[46]) using

hmmscan[31]. We filtered the hit Pfam domains by full sequence E-value ≤ 1e-3 and domain-based

CE-value ≤ 1e-5 or CI-value ≤ 1e-5, domain coverage ≥ 50% (coverage of aligned portion of

domain to full length domain) and selected the top hit for each aligned region.

Generation of multiple sequence alignments

776   To construct MSAs for each protein in the *Se* proteome, we initially used HHblits[35] to

search against UniRef (2022 release[37]) and BFD (2019 release[36]). For those MSA that contained

<500 sequences after this approach, we conducted extensive homology searches against

metagenome databases similar to Anishchenko *et al.*[68]. We first converted the initial HHblits (E-

value ≤ 1e-3) MSAs for each protein into a hidden Markov model (HMM) which we used as a

seed sequence profile to search against metagenomic and metatranscriptomic sequences from

JGI[69], MGnify[70], and UniRef with hmmsearch[31], respectively. Hits from JGI, MGnify, and

UniRef were gathered and aligned to the query protein by phmmer and jackhammer[31], to retain

sequences with E-value ≤ 1e-5 at each stage. We removed columns containing gaps in the query

sequence from the MSAs, which were then subjected to redundancy filtering at 95% sequence

identity and 50% sequence coverage. We selected the deepest MSA for each protein (from

HHblits against UniRef+BFD or phmmer/jackhmmer against metagenome) for downstream

analysis.

Structure-based annotations

Using MSAs described above, we computed AlphaFold (AF) models (model_1 without

structural template search, 10 recycles, and version 1.0 weights) for 852 of the 855 proteins (3

were excluded because of MSA depth or protein size)[34]. We then applied FoldSeek[38] against

AFdatabase50 (AF Protein Structure Database clustered at 50% sequence identity and 80%

coverage). We report the top hits in Table S2 based on E-value ≤ 0.00001, TM-alignment score

≥ 0.5 (which correlates well to fold-family level similarity[51]), and query coverage and target

coverage ≥ 50%. To classify proteins into evolutionary contexts based on structure and sequence

similarity, we used Domain Parser for AlphaFold Models (DPAM)[40], and the full results are

available at https://conglab.swmed.edu/ECOD_Se/Se_ECOD.html.

Nucleotide sequence accession numbers.

The complete genome sequences of *Se* ML1, *Nl* ML1and *Ai* F0345 have been deposited

in GenBank under BioProject PRJNA957798 with accession numbers SAMN34266291,

SAMN34266292 and SAMN34266293, respectively.

 Figure Legends

Figure 1. Phylogenetic placement and genome sequencing of newly isolated Saccharibacteria

strains *S. epibionticum* ML1 and *N. lyticus* ML1. (A) Maximum growth (fold change) achieved

by *S. epibionticum* ML1 and *N. lyticus* ML1 during co-culture with compatible host species *A.*

israelii and *Propionibacterium propionicum* respectively, and population change (growth or

death) detected at equivalent timepoints with an incompatible host. (B, C) 16S rRNA gene

sequence phylogeny of CPR bacteria (B) and Saccharibacteria (C) indicating placement of *S.*

epibionticum and *N. lyticus* ML1. (B) Is adapted from Castelle *et al.*[6] Lineages designations in

(C) derive from McLean *et al*[11]. (D, E) Overview of the genome sequences of *S. epibionticum*

ML1 and *N. lyticus* ML1.

Figure 2. Harnessing natural transformation to generate mutant Saccharibacteria. (A) Schematic

depicting the intergenic neutral site (blue, NS1) targeted for insertion of a hygomycin resistance

cassette (yellow) in the *Se* genome and the linear DNA fragment employed in transformation

experiments. Primer binding sites used for genotyping are indicated (sites 1-3). (B) Overview of

the *Se* transformation protocol. After incubation with linear DNA, *Se + Ai* co-cultures are

enlarged concomitant with hygromycin addition and serially passage with addition of naïve host

at each dilution to promote *Se* growth (grey box). Clonal transformed *Se* populations were

obtained by plating to isolate single colonies of *Ai* with accompanying *Se* cells, followed by

growth in liquid culture, with additional *Ai,* to promote *Se* population expansion. (C) PCR-based

genotyping of *Se* clones obtained following transformation according the protocol show in (B) in

the presence (right) or absence (left) of selection with hygromycin during the expansion and

passaging steps. Binding sites for primers targeting NS1 (1, 2) and *hph* (3) are shown in (A).

Positive control primers (*Se*) target a locus distant from NS1. (D) Growth (red) and percent of *Se*

transformed (grey) over the course of transformation protocol depicted in (B), in the presence

(squares) or absence (circles) of selection with hygromycin. See also Figures S2A, S2B.

Figure 3. Introduction of reporter genes via natural transformation enables tracking of the *S.*

epibionticum lifecycle. (A) Luminescence production from *Se*–*Ai* co-cultures (left) or co-culture

filtrates (right) in which *Se* contains a nanoluciferase expression cassette inserted at NS1 (shown

at bottom). (B) Fluorescence and phase contrast micrographs of *Se*–*Ai* co-cultures in which *Se*

carries an *mcherry* (top) or *sfgfp* (bottom) expression cassette inserted at NS1. (C) Snapshots

captured at the indicated time points from timelapse fluorescence and phase contrast microscopy

of GFP-expressing *Se* grown in co-culture with *Ai.* Arrows indicate *Se* cells exhibiting

productive (pink, purple) and non-productive (blue) interactions with *Ai* cells. (D) Omnipose-

generated segmentation of *Se* and *Ai* cells depicted in (C), at the start (left) and end (right) of the

20 hr growth period. (E) Growth of individual *Ai* cells as impacted by productive (light grey) or

non-productive *Se* cells (black, dark grey). Colors correspond to cell masks shown in (D). For the

full time course captured in (C-E) and additional examples of tracked *Se*–*Ai* growth, see Videos

S1-S10. Data in (A) represent mean ± s.d. Asterisks indicate statistically significant differences

(unpaired two-tailed student's t-test; *$p<0.05$).

Figure 4. Identification of genes important for fitness of *S. epibionticum* during co-culture with

Ai identified by Tn-seq. (A) Overview of normalized transposon insertion frequency across the

Se genome detected in input DNA used for mutagenesis (dark grey), and from samples collected

immediately following transformation (T0, dark blue) and subsequent outgrowth time points

(T1-T3, shades of blue). Genes encoding proteins belonging to CPR-enriched protein families

(light green) and those found to be significantly important for fitness across all time points and

by two different metrics (dark green). The location of the arginine deiminase system (ADS)

genes and two loci containing T4P genes (T4P$_1$, T4P$_2$) are indicated in the outer circle. (B)

Schematic depicting T4P$_1$ and flanking genes (separated by vertical lines) and the relative fitness

contribution of each gene as determined by the TRANSIT resampling algorithm (adjusted p-

values) and ALDEx2 (*delta* scores). Asterisks indicated genes found to be significant by both

metrics across all time points. Gene annotations were derived from Foldseek queries using AF

models generated for each gene product (see methods). (C) *Se* population levels detected in *Se–*

Ai cocultures following transformation with constructs designed to replace the indicated genes

with *hph.* Candidate essential genes tested were considered significant for *Se* fitness by

TRANSIT and ALDEx2 at a minimum of 2 of 3 time points (see Table S1). (D, E) Total *Se*

population (D) and proportion transformed (E) following transformation with an unmarked

cassette targeted to NS1 in the indicated strains of *Se*. See also Figures S3 and S4, Table S1.

Data in (C-E) represent mean ± s.d. Asterisks indicate statistically significant differences (C,

one-way ANOVA followed by Dunnett's compared to no DNA control; E, unpaired two-tailed

student's t-test; *$p < 0.05$, ns, not significant).

Figure 5. Inclusion of extensive metagenomic data in MSAs enables proteome-wide AF

modeling of *Se* protein structures. (A) Histograms depicting MSA depths obtained for *Se*

proteins using HHblits. (B) Maximum depths obtained for *Se* protein MSAs that initially

contained <500 sequences. Additional sequences were sourced from metagenomic sequence

databases and incorporated into MSAs using Jackhmmer or Phmmer (see Methods). (C)

Comparison of the AF confidence metric (pLDDT) determined using Hhblits or

Jackhmmer/Phmmer (Metagenome)-generated MSAs for *Se* proteins with initially shallow

MSAs (<500). (D and E) Example *Se* protein structure models and associated predicted

alignment matrices obtained using shallow (right) or metagenomic sequence-improved (left)

MSAs. *Se* proteins models (blue) are aligned to models from top FoldSeek (FS) hits (light grey,

A0A2H0BDY7 (D), A0A7W4EM10, metagenome and A0A660M3E3, shallow, (E)), when

available. Some structures are trimmed to highlight the alignment. See also Figure S5.

Figure S1. Hygromycin resistance phenotypes of *Se* and *Ai* enable use of *hph* as a selectable

marker in *Se*. Growth of *Ai* (top) and *Se* (bottom) in co-cultures containing the indicated

concentrations of hygromycin.

Figure S2. Optimization of the *Se* transformation protocol and transformation of a second

Saccharibacteria species. (A, B) Quantification of *Se, Ai* and transformed *Se* populations over the

course of our transformation protocol (See Figure 2B) with varying lengths (A) or concentrations

(B) of transforming DNA. Transformations were performed in the presence (top panels) and

absence (bottom panels) of selection for transformants with hygromycin. (C) Schematic

depicting the intergenic neutral site (blue, NS1) targeted for insertion of a hygomycin resistance

cassette (yellow) in the *N. lyticus* ML1 genome and the linear DNA fragment employed in

transformation experiments with this species. Primer binding sites used for genotyping are

indicated (sites 1-3). (D) Genotyping of *N. lyticus*–*P. propionicum* co-cultures transformed with

the linear DNA fragment depicted in (C) (right) or parallel negative control co-cultures with no

DNA added (left) at the end of passage 4 (see Methods) with primers targeting NS1 (1,2) or *hph*

(3). Positive control *N. lyticus* primers target a genomic locus distant from NS1.

Figure S3. Population dynamics of *Se* and *Ai* during transposon mutagenesis. Transformation co-

culture was diluted and passaged into fresh media, with the addition of fresh *Ai,* after collection

of the T0, T1 and T2 samples (see Methods).

Figure S4. Distribution of 921 core protein families across CPR and non-CPR bacterial genomes,

including *Se* and *N. lyticus* ML1. Columns represent core families (derived from Meheust *et al.*[5])

and rows represent individual genomes from the indicated bacterial groups. CPR enriched

protein families indicated at top (blue), and dendrogram at left represents clustering of bacterial

strains based on protein family content.

Figure S5. The *Ai* genome and structural models for the *Se* proteome generated using

metagenomic sequence enriched MSAs for use in future studies of the *Se-Ai* system. (A)

Overview of the genome sequence of *A. israelii* F0345. (B-D) Example *Se* protein structure

models and associated predicted alignment matrices obtained using shallow (right) or

metagenomic sequence-improved (left) MSAs. *Se* proteins models (blue) are aligned to models

for top FS hits (light grey, AF database50 numbers A0A8B1YQG7 (B), A0A660LZS9 (C) and

A0A563CGZ5 (D)), when available. Some structures are trimmed to highlight the alignment.

References

1.   Hug, L.A., Baker, B.J., Anantharaman, K., Brown, C.T., Probst, A.J., Castelle, C.J.,
Butterfield, C.N., Hernsdorf, A.W., Amano, Y., Ise, K., et al. (2016). A new view of the tree of
life. Nat Microbiol *1*, 16048. 10.1038/nmicrobiol.2016.48.

2.   Katz, M., Hover, B.M., and Brady, S.F. (2016). Culture-independent discovery of natural
products from soil metagenomes. J Ind Microbiol Biotechnol *43*, 129-141. 10.1007/s10295-015-
1706-6.

3.   Rinke, C., Schwientek, P., Sczyrba, A., Ivanova, N.N., Anderson, I.J., Cheng, J.F.,
Darling, A., Malfatti, S., Swan, B.K., Gies, E.A., et al. (2013). Insights into the phylogeny and
coding potential of microbial dark matter. Nature *499*, 431-437. 10.1038/nature12352.

4.   Ji, Y., Zhang, P., Zhou, S., Gao, P., Wang, B., and Jiang, J. (2022). Widespread but
Poorly Understood Bacteria: Candidate Phyla Radiation. Microorganisms *10*.
10.3390/microorganisms10112232.

5.   Meheust, R., Burstein, D., Castelle, C.J., and Banfield, J.F. (2019). The distinction of
CPR bacteria from other bacteria based on protein family content. Nat Commun *10*, 4173.
10.1038/s41467-019-12171-z.

6.   Castelle, C.J., Brown, C.T., Anantharaman, K., Probst, A.J., Huang, R.H., and Banfield,
J.F. (2018). Biosynthetic capacity, metabolic variety and unusual biology in the CPR and
DPANN radiations. Nat Rev Microbiol *16*, 629-645. 10.1038/s41579-018-0076-2.

7.   Batinovic, S., Rose, J.J.A., Ratcliffe, J., Seviour, R.J., and Petrovski, S. (2021).
Cocultivation of an ultrasmall environmental parasitic bacterium with lytic ability against
bacteria associated with wastewater foams. Nat Microbiol *6*, 703-711. 10.1038/s41564-021-
00892-1.

8.   He, X., McLean, J.S., Edlund, A., Yooseph, S., Hall, A.P., Liu, S.Y., Dorrestein, P.C.,
Esquenazi, E., Hunter, R.C., Cheng, G., et al. (2015). Cultivation of a human-associated TM7
phylotype reveals a reduced genome and epibiotic parasitic lifestyle. Proc Natl Acad Sci U S A
112, 244-249. 10.1073/pnas.1419038112.

9.   Kuroda, K., Yamamoto, K., Nakai, R., Hirakata, Y., Kubota, K., Nobu, M.K., and
Narihiro, T. (2022). Symbiosis between Candidatus Patescibacteria and Archaea Discovered in
Wastewater-Treating Bioreactors. MBio *13*, e0171122. 10.1128/mbio.01711-22.

10.   Yakimov, M.M., Merkel, A.Y., Gaisin, V.A., Pilhofer, M., Messina, E., Hallsworth, J.E.,
Klyukina, A.A., Tikhonova, E.N., and Gorlenko, V.M. (2022). Cultivation of a vampire:
'Candidatus Absconditicoccus praedator'. Environmental microbiology *24*, 30-49. 10.1111/1462-
2920.15823.

11.   McLean, J.S., Bor, B., Kerns, K.A., Liu, Q., To, T.T., Solden, L., Hendrickson, E.L.,
Wrighton, K., Shi, W., and He, X. (2020). Acquisition and Adaptation of Ultra-small Parasitic

Reduced Genome Bacteria to Mammalian Hosts. Cell Rep *32*, 107939.
10.1016/j.celrep.2020.107939.

12.   Bor, B., Bedree, J.K., Shi, W., McLean, J.S., and He, X. (2019). Saccharibacteria (TM7)
in the Human Oral Microbiome. J Dent Res *98*, 500-509. 10.1177/0022034519831671.

13.   Chipashvili, O., Utter, D.R., Bedree, J.K., Ma, Y., Schulte, F., Mascarin, G., Alayyoubi,
Y., Chouhan, D., Hardt, M., Bidlack, F., et al. (2021). Episymbiotic Saccharibacteria suppresses
gingival inflammation and bone loss in mice through host bacterial modulation. Cell Host
Microbe *29*, 1649-1662 e1647. 10.1016/j.chom.2021.09.009.

14.   Adler, C.J., Dobney, K., Weyrich, L.S., Kaidonis, J., Walker, A.W., Haak, W., Bradshaw,
C.J., Townsend, G., Soltysiak, A., Alt, K.W., et al. (2013). Sequencing ancient calcified dental
plaque shows changes in oral microbiota with dietary shifts of the Neolithic and Industrial
revolutions. Nature genetics *45*, 450-455, 455e451. 10.1038/ng.2536.

15.   Bor, B., McLean, J.S., Foster, K.R., Cen, L., To, T.T., Serrato-Guillen, A., Dewhirst,
F.E., Shi, W., and He, X. (2018). Rapid evolution of decreased host susceptibility drives a stable
relationship between ultrasmall parasite TM7x and its bacterial host. Proc Natl Acad Sci U S A
115, 12277-12282. 10.1073/pnas.1810625115.

16.   Nie, J., Utter, D.R., Kerns, K.A., Lamont, E.I., Hendrickson, E.L., Liu, J., Wu, T., He, X.,
McLean, J., and Bor, B. (2022). Strain-Level Variation and Diverse Host Bacterial Responses in
Episymbiotic Saccharibacteria. mSystems *7*, e0148821. 10.1128/msystems.01488-21.

17.   Xie, B., Wang, J., Nie, Y., Tian, J., Wang, Z., Chen, D., Hu, B., Wu, X.L., and Du, W.
(2022). Type IV pili trigger episymbiotic association of Saccharibacteria with its bacterial host.
Proc Natl Acad Sci U S A *119*, e2215990119. 10.1073/pnas.2215990119.

18.   Bor, B., Collins, A.J., Murugkar, P.P., Balasubramanian, S., To, T.T., Hendrickson, E.L.,
Bedree, J.K., Bidlack, F.B., Johnston, C.D., Shi, W., et al. (2020). Insights Obtained by
Culturing Saccharibacteria With Their Bacterial Hosts. J Dent Res *99*, 685-694.
10.1177/0022034520905792.

19.   Cross, K.L., Campbell, J.H., Balachandran, M., Campbell, A.G., Cooper, S.J., Griffen,
A., Heaton, M., Joshi, S., Klingeman, D., Leys, E., et al. (2019). Targeted isolation and
cultivation of uncultivated bacteria by reverse genomics. Nat Biotechnol *37*, 1314-1321.
10.1038/s41587-019-0260-6.

20.   He, X. (2023). Culture-based approaches to studying "microbial dark matter". Proc Natl
Acad Sci U S A *120*, e2219691120. 10.1073/pnas.2219691120.

21.   Stackenbrandt, E., and Goebel, B.M. (1994). Taxonomic Note: A Place for DNA-DNA
Reassociation and 16S rRNA Sequence Analysis in the Present Species Definition in
Bacteriology. Int. J. Syst. Evol. Microbiol. *44*, 846-849.

22.   Damke, P.P., Celma, L., Kondekar, S.M., Di Guilmi, A.M., Marsin, S., Depagne, J.,
Veaute, X., Legrand, P., Walbott, H., Vercruyssen, J., et al. (2022). ComFC mediates transport

and handling of single-stranded DNA during natural transformation. Nat Commun *13*, 1961.
10.1038/s41467-022-29494-z.

23.   Dubnau, D., and Blokesch, M. (2019). Mechanisms of DNA Uptake by Naturally
Competent Bacteria. Annu Rev Genet *53*, 217-237. 10.1146/annurev-genet-112618-043641.

24.   Sharma, D.K., Misra, H.S., Bihani, S.C., and Rajpurohit, Y.S. (2023). Biochemical
Properties and Roles of DprA Protein in Bacterial Natural Transformation, Virulence, and Pilin
Variation. J Bacteriol *205*, e0046522. 10.1128/jb.00465-22.

25.   Johnston, C., Martin, B., Fichant, G., Polard, P., and Claverys, J.P. (2014). Bacterial
transformation: distribution, shared mechanisms and divergent control. Nat Rev Microbiol *12*,
181-196. 10.1038/nrmicro3199.

26.   DeJesus, M.A., Ambadipudi, C., Baker, R., Sassetti, C., and Ioerger, T.R. (2015).
TRANSIT--A Software Tool for Himar1 TnSeq Analysis. PLoS computational biology *11*,
e1004401. 10.1371/journal.pcbi.1004401.

27.   Fernandes, A.D., Reid, J.N., Macklaim, J.M., McMurrough, T.A., Edgell, D.R., and
Gloor, G.B. (2014). Unifying the analysis of high-throughput sequencing datasets: characterizing
RNA-seq, 16S rRNA gene sequencing and selective growth experiments by compositional data
analysis. Microbiome *2*, 15. 10.1186/2049-2618-2-15.

28.   Bedree, J.K., Bor, B., Cen, L., Edlund, A., Lux, R., McLean, J.S., Shi, W., and He, X.
(2018). Quorum Sensing Modulates the Epibiotic-Parasitic Relationship Between Actinomyces
odontolyticus and Its Saccharibacteria epibiont, a Nanosynbacter lyticus Strain, TM7x. Front
Microbiol *9*, 2049. 10.3389/fmicb.2018.02049.

29.   Wu, C., and Ton-That, H. (2010). Allelic exchange in Actinomyces oris with mCherry
fluorescence counterselection. Applied and environmental microbiology *76*, 5987-5989.
10.1128/AEM.00811-10.

30.   Gane, A., Bileshi, M.L., Dohan, D., Speretta, E., Heliou, A., Meng-Papaxanthos, L.,
Zellner, H., Brevdo, E., Parikh, A., Martin, M.J., et al. (2022). ProtNLM: Model-based Natural
Language Protein
Annotation. Preprint.

31.   Eddy, S.R. (2011). Accelerated Profile HMM Searches. PLoS computational biology *7*,
e1002195. 10.1371/journal.pcbi.1002195.

32.   Mistry, J., Chuguransky, S., Williams, L., Qureshi, M., Salazar, G.A., Sonnhammer,
E.L.L., Tosatto, S.C.E., Paladin, L., Raj, S., Richardson, L.J., et al. (2021). Pfam: The protein
families database in 2021. Nucleic Acids Res *49*, D412-D419. 10.1093/nar/gkaa913.

33.   Baek, M., DiMaio, F., Anishchenko, I., Dauparas, J., Ovchinnikov, S., Lee, G.R., Wang,
J., Cong, Q., Kinch, L.N., Schaeffer, R.D., et al. (2021). Accurate prediction of protein structures
and interactions using a three-track neural network. Science *373*, 871-876.
10.1126/science.abj8754.

34.   Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,
Tunyasuvunakool, K., Bates, R., Zidek, A., Potapenko, A., et al. (2021). Highly accurate protein
structure prediction with AlphaFold. Nature *596*, 583-589. 10.1038/s41586-021-03819-2.

35.   Remmert, M., Biegert, A., Hauser, A., and Soding, J. (2011). HHblits: lightning-fast
iterative protein sequence searching by HMM-HMM alignment. Nature methods *9*, 173-175.
10.1038/nmeth.1818.

36.   Steinegger, M., Mirdita, M., and Soding, J. (2019). Protein-level assembly increases
protein sequence recovery from metagenomic samples manyfold. Nature methods *16*, 603-606.
10.1038/s41592-019-0437-4.

37.   Suzek, B.E., Huang, H., McGarvey, P., Mazumder, R., and Wu, C.H. (2007). UniRef:
comprehensive and non-redundant UniProt reference clusters. Bioinformatics *23*, 1282-1288.
10.1093/bioinformatics/btm098.

38.   van Kempen, M., Kim, S.S., Tumescheit, C., Mirdita, M., Le, J., Gilchrist, C.L.M.,
Soding, J., and Steinegger, M. (2023). Fast and accurate protein structure search with Foldseek.
bioRxiv.

39.   Cheng, H., Schaeffer, R.D., Liao, Y., Kinch, L.N., Pei, J., Shi, S., Kim, B.H., and
Grishin, N.V. (2014). ECOD: an evolutionary classification of protein domains. PLoS
computational biology *10*, e1003926. 10.1371/journal.pcbi.1003926.

40.   Zhang, J., Schaeffer, R.D., Durham, J., Cong, Q., and Grishin, N.V. (2023). DPAM: A
domain parser for AlphaFold models. Protein Sci *32*, e4548. 10.1002/pro.4548.

41.   Moreira, D., Zivanovic, Y., Lopez-Archilla, A.I., Iniesto, M., and Lopez-Garcia, P.
(2021). Reductive evolution and unique predatory mode in the CPR bacterium Vampirococcus
lugosii. Nat Commun *12*, 2454. 10.1038/s41467-021-22762-4.

42.   Murugkar, P.P., Collins, A.J., Chen, T., and Dewhirst, F.E. (2020). Isolation and
cultivation of candidate phyla radiation Saccharibacteria (TM7) bacteria in coculture with
bacterial hosts. J Oral Microbiol *12*, 1814666. 10.1080/20002297.2020.1814666.

43.   Cressler, C.E., Mc, L.D., Rozins, C., J, V.D.H., and Day, T. (2016). The adaptive
evolution of virulence: a review of theoretical predictions and empirical tests. Parasitology *143*,
915-930. 10.1017/S003118201500092X.

44.   Egido, J.E., Costa, A.R., Aparicio-Maldonado, C., Haas, P.J., and Brouns, S.J.J. (2022).
Mechanisms and clinical importance of bacteriophage resistance. FEMS Microbiol Rev *46*.
10.1093/femsre/fuab048.

45.   He, C., Keren, R., Whittaker, M.L., Farag, I.F., Doudna, J.A., Cate, J.H.D., and Banfield,
J.F. (2021). Genome-resolved metagenomics reveals site-specific diversity of episymbiotic CPR
bacteria and DPANN archaea in groundwater ecosystems. Nat Microbiol *6*, 354-365.
10.1038/s41564-020-00840-5.

46.   Ellison, C.K., Dalia, T.N., Vidal Ceballos, A., Wang, J.C., Biais, N., Brun, Y.V., and
Dalia, A.B. (2018). Retraction of DNA-bound type IV competence pili initiates DNA uptake
during natural transformation in Vibrio cholerae. Nat Microbiol *3*, 773-780. 10.1038/s41564-
018-0174-y.

47.   Hepp, C., and Maier, B. (2016). Kinetics of DNA uptake during transformation provide
evidence for a translocation ratchet mechanism. Proc Natl Acad Sci U S A *113*, 12467-12472.
10.1073/pnas.1608110113.

48.   Tian, J., Utter, D.R., Cen, L., Dong, P.T., Shi, W., Bor, B., Qin, M., McLean, J.S., and
He, X. (2022). Acquisition of the arginine deiminase system benefits epiparasitic
Saccharibacteria and their host bacteria in a mammalian niche environment. Proc Natl Acad Sci
U S A *119*. 10.1073/pnas.2114909119.

49.   Batty, I. (1958). Actinomyces odontolyticus, a new species of actinomycete regularly
isolated from deep carious dentine. J Pathol Bacteriol *75*, 455-459. 10.1002/path.1700750225.

50.   Dewhirst, F.E., Chen, T., Izard, J., Paster, B.J., Tanner, A.C., Yu, W.H., Lakshmanan, A.,
and Wade, W.G. (2010). The human oral microbiome. J Bacteriol *192*, 5002-5017.
10.1128/JB.00542-10.

51.   Nikolaitchouk, N., Hoyles, L., Falsen, E., Grainger, J.M., and Collins, M.D. (2000).
Characterization of Actinomyces isolates from samples from the human urogenital tract:
description of Actinomyces urogenitalis sp. nov. Int J Syst Evol Microbiol *50 Pt 4*, 1649-1654.
10.1099/00207713-50-4-1649.

52.   Collins, A.J., Murugkar, P.P., and Dewhirst, F.E. (2021). Establishing Stable Binary
Cultures of Symbiotic Saccharibacteria from the Oral Cavity. J Vis Exp. 10.3791/62484.

53.   Escapa, I.F., Chen, T., Huang, Y., Gajare, P., Dewhirst, F.E., and Lemon, K.P. (2018).
New Insights into Human Nostril Microbiome from the Expanded Human Oral Microbiome
Database (eHOMD): a Resource for the Microbiome of the Human Aerodigestive Tract.
mSystems *3*. 10.1128/mSystems.00187-18.

54.   Pruesse, E., Quast, C., Knittel, K., Fuchs, B.M., Ludwig, W., Peplies, J., and Glockner,
F.O. (2007). SILVA: a comprehensive online resource for quality checked and aligned ribosomal
RNA sequence data compatible with ARB. Nucleic Acids Res *35*, 7188-7196.
10.1093/nar/gkm864.

55.   Pruesse, E., Peplies, J., and Glockner, F.O. (2012). SINA: accurate high-throughput
multiple sequence alignment of ribosomal RNA genes. Bioinformatics *28*, 1823-1829.
10.1093/bioinformatics/bts252.

56.   Stamatakis, A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-
analysis of large phylogenies. Bioinformatics *30*, 1312–1313. 10.1093/bioinformatics/btu033.

57.   Wick, R.R., Judd, L.M., Cerdeira, L.T., Hawkey, J., Meric, G., Vezina, B., Wyres, K.L.,
and Holt, K.E. (2021). Trycycler: consensus long-read assemblies for bacterial genomes.
Genome biology *22*, 266. 10.1186/s13059-021-02483-z.

58.   Li, H. (2016). Minimap and miniasm: fast mapping and de novo assembly for noisy long
sequences. Bioinformatics *32*, 2103-2110. 10.1093/bioinformatics/btw152.

59.   Zimin, A.V., Puiu, D., Luo, M.C., Zhu, T., Koren, S., Marcais, G., Yorke, J.A., Dvorak,
J., and Salzberg, S.L. (2017). Hybrid assembly of the large and highly repetitive genome of
Aegilops tauschii, a progenitor of bread wheat, with the MaSuRCA mega-reads algorithm.
Genome Res *27*, 787-792. 10.1101/gr.213405.116.

60.   Seemann, T. (2014). Prokka: rapid prokaryotic genome annotation. Bioinformatics *30*,
2068-2069. 10.1093/bioinformatics/btu153.

61.   Dehlinger, B., Jurss, J., Lychuk, K., and Putonti, C. (2021). The Dynamic Codon Biaser:
calculating prokaryotic codon usage biases. Microb Genom *7*. 10.1099/mgen.0.000663.

62.   Hall, M.P., Unch, J., Binkowski, B.F., Valley, M.P., Butler, B.L., Wood, M.G., Otto, P.,
Zimmerman, K., Vidugiris, G., Machleidt, T., et al. (2012). Engineered luciferase reporter from a
deep sea shrimp utilizing a novel imidazopyrazinone substrate. ACS Chem Biol *7*, 1848-1857.
10.1021/cb3002478.

63.   Zalacain, M., Gonzalez, A., Guerrero, M.C., Mattaliano, R.J., Malpartida, F., and
Jimenez, A. (1986). Nucleotide sequence of the hygromycin B phosphotransferase gene from
Streptomyces hygroscopicus. Nucleic Acids Res *14*, 1565-1581. 10.1093/nar/14.4.1565.

64.   Cutler, K.J., Stringer, C., Lo, T.W., Rappez, L., Stroustrup, N., Brook Peterson, S.,
Wiggins, P.A., and Mougous, J.D. (2022). Omnipose: a high-precision morphology-independent
solution for bacterial cell segmentation. Nature methods *19*, 1438-1448. 10.1038/s41592-022-
01639-4.

65.   Gallagher, L.A., Bailey, J., and Manoil, C. (2020). Ranking essential bacterial processes
by speed of mutant death. Proc Natl Acad Sci U S A *117*, 18010-18017.
10.1073/pnas.2001507117.

66.   Gallagher, L.A. (2019). Methods for Tn-Seq Analysis in Acinetobacter baumannii.
Methods in molecular biology (Clifton, N.J *1946*, 115-134. 10.1007/978-1-4939-9118-1_12.

67.   Arrieta-Ortiz, M.L., Pan, M., Kaur, A., Pepper-Tunick, E., Srinivas, V., Dash, A.,
Immanuel, S.R.C., Brooks, A.N., Shepherd, T.R., and Baliga, N.S. (2023). Disrupting the ArcA
Regulatory Network Amplifies the Fitness Cost of Tetracycline Resistance in Escherichia coli.
mSystems *8*, e0090422. 10.1128/msystems.00904-22.

68.   Anishchenko, I., Baek, M., Park, H., Hiranuma, N., Kim, D.E., Dauparas, J., Mansoor, S.,
Humphreys, I.R., and Baker, D. (2021). Protein tertiary structure prediction and refinement using
deep learning and Rosetta in CASP14. Proteins *89*, 1722-1733. 10.1002/prot.26194.

69.   Chen, I.A., Chu, K., Palaniappan, K., Ratner, A., Huang, J., Huntemann, M., Hajek, P.,
Ritter, S., Varghese, N., Seshadri, R., et al. (2021). The IMG/M data management and analysis
system v.6.0: new tools and advanced capabilities. Nucleic Acids Res *49*, D751-D763.
10.1093/nar/gkaa939.

70.   Mitchell, A.L., Almeida, A., Beracochea, M., Boland, M., Burgin, J., Cochrane, G.,
Crusoe, M.R., Kale, V., Potter, S.C., Richardson, L.J., et al. (2020). MGnify: the microbiome
analysis resource in 2020. Nucleic Acids Res *48*, D570-D578. 10.1093/nar/gkz1035.
1141

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A

Count

Subject to add'l
MSA improvement

0   200   400   >500
MSA HHblits depth

B

Count

0   200   400   >500
Highest MSA depth

C

Metagenome
(pLDDT)

20   40   60   80   100
HHblits
(pLDDT)

D SEML1_0214

E SEML1_0811

■ *Se* protein
■ FoldSeek hit

Predicted aligned error (Å)

0   161
0   161

0   215
0   215

Figure S1

Figure S2

Figure S3

Figure S4

Figure S5