Ribosomal Frameshifting Selectively Modulates the Biosynthesis, Assembly, and Function of a Misfolded CFTR Variant

Patrick Carmody,[1][†] Francis J. Roushar,[1][†] Austin Tedman,[1] Madeline Herwig,[2] Minsoo Kim,[2,3] Eli F. McDonald,[2] Jennifer Wong-Roushar,[1] Jon-Luc Poirier,[1] Nathan B. Zelt,[1] Ben T. Pockrass,[1] Andrew G. McKee,[1] Charles P. Kuntz,[1] Lars Plate,[2,4] Wesley D. Penn,[1]* and Jonathan P. Schlebach[1]*

[1] Department of Chemistry, Indiana University Bloomington, Bloomington, IN, USA 47401

[2] Department of Chemistry, Vanderbilt University, Nashville, TN, USA 37240

[3] Program in Chemical and Physical Biology, Vanderbilt University, Nashville, TN, USA 37240

[4] Department of Biological Sciences, Vanderbilt University, Nashville, TN, USA 37240

†Contributed Equally

*Corresponding Authors: jschleba@iu.edu, wdpenn@iu.edu

Abstract

The cotranslational misfolding of the cystic fibrosis transmembrane conductance regulator (CFTR) plays a central role in the molecular basis of cystic fibrosis (CF). The misfolding of the most common CF variant (ΔF508) remodels both the translational regulation and quality control of CFTR. Nevertheless, it is unclear how the misassembly of the nascent polypeptide modulates the translation machinery. In this work, we identify a structural motif within the CFTR transcript that stimulates efficient -1 ribosomal frameshifting and triggers the premature termination of translation. Though this motif does not appear to impact the wild-type CFTR interactome, silent mutations that disrupt this RNA structure alter how ΔF508 CFTR interacts with numerous translation and quality control proteins. Moreover, disrupting this RNA structure enhances both the expression and function of ΔF508 CFTR with no impact on wild-type. Finally, we show that disrupting this motif enhances the pharmacological rescue of ΔF508 by Trikafta, which implies ribosomal frameshifting antagonizes the effects of leading CF therapeutics. Together, our results reveal that ribosomal frameshifting selectively reduces the expression and assembly of a misfolded CFTR variant. These findings suggest cotranslational misfolding alters the processivity of translation and potentially the stability of the mRNA transcript through the dynamic modulation of ribosomal frameshifting.

Introduction

Cystic fibrosis (CF) is a disease of protein homeostasis (proteostasis) that arises from the defective biosynthesis, folding, and/ or function of a chloride channel known as the cystic fibrosis transmembrane conductance regulator (CFTR).[1,2] The most common CF mutation (ΔF508) induces CFTR misfolding by decoupling the folding of its subdomains during translation.[3,4] This cotranslational misassembly reaction enhances the retention and degradation of the CFTR protein within the endoplasmic reticulum (ER) and ultimately reduces the trafficking of the functional protein to the plasma membrane. This lapse in protein quality control (QC) coincides with the remodeling of CFTR proteostasis network and changes in the dynamics of CFTR translation.[5-11] Nevertheless, the precise chain of events involved in the crosstalk between the conformational state of nascent CFTR, the activity of the translation machinery, and the interaction of the nascent chain with various components of the cellular proteostasis network is not fully understood.

CFTR QC begins during its cotranslational folding at the ribosome-translocon complex.[12] Vectorial folding of its five subdomains is orchestrated, in part, by positional variations in translation kinetics that are coordinated by rare codons,[9] the relative abundance of certain translation factors,[5,7] and the presence of mRNA secondary structure.[13] Together, these effectors help the ribosome tailor its activity to the kinetic constraints of cotranslational CFTR folding. Indeed, the effects of CF mutations on CFTR folding and assembly are highly sensitive to changes in translational dynamics. For instance, suppressing translation initiation reduces the density of ribosomes on the ΔF508 transcript in a manner that partially rescues its stability, cellular trafficking, and function.[7,11] The cotranslational misfolding of ΔF508 CFTR also appears to modulate translation in *cis* by enhancing ribosome collisions and triggering "preemptive quality control" pathways.[11] Though the later finding suggests the intrinsic activity of the ribosome is sensitive to conformational transitions in the nascent chain, it remains unclear whether cotranslational misfolding events are capable of directly altering translation in real time.

We recently found that cotranslational folding mechanically modulates a translational recoding mechanism known as -1 programmed ribosomal frameshifting (-1PRF) during viral polyprotein biosynthesis.[14,15] These

findings reveal that, under certain circumstances, the ribosome exhibits an enhanced propensity to slip into alternative reading frames in response to conformational transitions in the nascent polypeptide chain.[16] In the following, we evaluate whether similar feedback occurs during CFTR biosynthesis. We first identify a structured segment within the region of the CFTR transcript that stimulates ribosomal frameshifting (RF). We then show that the disruption of this motif selectively remodels the interactome of ΔF508 CFTR in a manner that partially restores its expression and function. These modifications, which appear to have no effect on WT CFTR, also enhance the pharmacological rescue of ΔF508 CFTR by the leading CF therapeutic Trikafta. Our results suggest this RF site acts a QC-mediated translational "kill switch" that selectively promotes the premature termination of translation in response to the cotranslational misfolding of the nascent chain.

Results

Discovery of an Active -1 Ribosomal Frameshift Site in the CFTR Transcript

Our recent findings show that the mechanical forces generated by cotranslational folding can stimulate -1PRF,[14,15] which minimally requires a slippery sequence within the mRNA and an adjacent stem-loop.[17] To identify potential RF sites in the CFTR transcript we first searched for consensus or near-consensus X_1 XXY_4 YYZ_7 slippery sequences, where XXX and YYY are nucleobase triplets and Z can be any nucleobase.[18-20] We identified five consensus X_1 XXY_4 YYZ_7 sites and eight other near-consensus sites distributed throughout the transcript (Figs. 1A & S1). Three of these sites are clustered within the region encoding nucleotide binding domain 2 (NBD2) and are proximal to two predicted RNA stem-loops (Fig. 1A). A uridine rich sequence referred to here as slip-site A (UUU AUU UUU UCU, SSA) lies upstream of the canonical slip-site B (A_1 AAA_4 AAC_7, SSB). SSB is eight nucleotides upstream of a small, predicted stem-loop (SL2). We also identified a third putative non-canonical slip site (A_1 GAA_4 AUA_7, SSC) downstream of SSB that is positioned seven nucleotides upstream of a much longer putative stem-loop (SL2); an ideal spacing for the stimulation of -1PRF.[17]

Given that ribosome collisions can occur during CFTR translation[21] and that they are known to modulate -1PRF,[22] we hypothesized that RF could occur at one or more of these sites. To determine whether efficient -1 RF occurs at any of these positions, we generated a series of bicistronic reporters in which this CFTR segment is flanked by an upstream Renilla luciferase (rLuc, expression control) and a downstream -1 firefly luciferase (fLuc, -1 RF reporter, Fig. 2A).[23] We generated three versions of this reporter that produce fLuc in response to frameshifting at any of these slip-sites (SSABC), or conditionally at B or C (SSBC), or at C only (SSC) by knocking out various -1 frame stop codons downstream of these slip-sites (Fig.

Figure 1. Putative mRNA structures within the CFTR ribosomal frameshift motif. A) The predicted secondary structure downstream of SSB is shown. Bases are colored according to their predicted base pair probabilities as determined by Vienna RNAfold. B) Secondary structure predictions for a construct bearing 14 silent mutations (RF_mut) are shown for the sake of comparison. Mutated bases are enlarged and bases are colored according to their predicted base pair probabilities as determined by Vienna RNAfold in both panels A and B, for reference. The positions of the three putative slip sites and the downstream stop codons in the -1 reading frame are indicated. C) A schematic depicts the layout of the full-length WT and ΔF508 CFTR transcripts bearing the RF_mut modifications.

2A orange). Transient expression of the SSABC reporter in a human bronchial epithelial cell line (CFBE41o-) generates a detectable fLuc frameshift signal 4-fold over the no-insert control baseline. rLuc intensities are similar for all three reporters, suggesting all three reporters exhibit similar expression under these conditions (Fig. 2B). fLuc intensities are also quite similar for all three reporters (Fig. 2C), which implies most frameshifting occurs at SSC. Consistent with this interpretation, we find that the fLuc signal is ablated by mutations that disrupt SSC (SSC_mut, Fig. 2C). We note that we ruled out splicing artifacts[24] by RT-PCR and found that the frameshift signal can also be partially suppressed by stop codons in either upstream of SSC in the 0-frame (5'Ter) or downstream of SSC in the -1 frame (3'Ter, Figs. 2D & S2), which confirms that these signals arise from a genuine RF. By normalizing the rLuc and fLuc signals, we estimate ribosomal frameshifting occurs with an efficiency of 2.9 ± 0.2% at SSC, which is comparable to levels achieved during translation of the canonical HIV gag-pol motif (5.2 ± 0.4%, Fig. 2D). Similar frameshifting also occurs in HEK293T cells (Fig. S3). Together, these results identify an active -1 RF motif within the human CFTR transcript.

Figure 2. Stimulation of ribosomal frameshifting by structural elements within the CFTR transcript. The efficiency of ribosomal frameshifting during the translation of a structured region within the CFTR transcript was assessed in CFBE41o- cells using a bicistronic dual luciferase reporter system. (A) Cartoon depicts the logic of a series of reporter constructs that generate Renilla Luciferase (RLuc) upon translation initiation and a firefly Luciferase (FLuc) in response to a -1 ribosomal frameshift. The positions of the potential slip-sites A (red), B (blue), and C (purple) are shown. Stop codons in the -1 frame are indicated in orange. Putative secondary structures generated by Vienna RNAfold for the 75 bases beginning 5 bases downstream of SSB are shown for reference. (B, C) Dot plots indicate the raw RLuc and FLuc luminescence intensities within the lysates of CFBE41o- cells transiently expressing these constructs, respectively. Results were taken from three biological replicates with 4 transfections per sample each. Central hashes represent the average intensities and whiskers reflect their standard deviation. (D) A bar graph depicts the average -1 ribosomal frameshifting efficiencies for each construct in CFBE41o- cells. Error bars reflect the standard deviation to serve as a measure of precision.

Impact of Ribosomal Frameshifting on the CFTR Interactome

Unlike viral -1PRF events that generate functionally distinct proteins, a -1 frameshift at SSC should only fuse five non-native residues onto nascent CFTR prior to translational termination and truncation in the middle of NBD2 (Fig. 1A). Based on this consideration, we hypothesized that the RF motif serves to promote frameshifting and the premature termination of translation in response to cotranslational CFTR misassembly. To determine whether this motif is involved in assembly, we assessed how this RNA structure influences the interactomes of WT and ΔF508 CFTR. Briefly, we designed a series of mutations to disrupt the slip-sites and weaken the stem-loops within NBD2 while preserving the amino acid sequence of CFTR in the 0-frame (RF$_{mut}$, Fig. 1B). We then incorporated these modifications into the full length CFTR (Fig. 1C) and used affinity purification-mass spectrometry (AP-MS) based interactome profiling to determine the effects of these silent modifications on the interactions formed by WT and ΔF508 CFTR.[5,25,26] In agreement with previous reports,[5,25] the ΔF508 mutation significantly enhances the interaction of CFTR with numerous proteostasis factors (Fig. 3A & Table S1). By comparison, the silent mutations in NBD2 have minimal impact on the WT CFTR interactome (WT-RF$_{mut}$, Fig 3A). In contrast, these same mutations significantly alter the propensity of ΔF508 CFTR to interact with several translation factors and quality control proteins (ΔF508-RF$_{mut}$, Fig. 2A). Consistent with our hypothesis, the disparate effects of this transcript modification on WT and ΔF508 suggests the native RNA structure in NBD2 selectively alters CFTR translation and assembly in response to misfolding. We note that the ΔF508 mutation is ~2.4 kb upstream of the ribosomal frameshift site in NBD2 (Fig. 1C), which suggests these differences likely arise from the effects of the ΔF508 mutation on the nascent CFTR protein rather than perturbation of the mRNA structure.

Figure 3. Impact of ribosomal frameshifting on the CFTR interactome. Affinity purification-mass spectrometry was employed to compare the effects of the silent RF_{mut} mutations on the interactome of WT and ΔF508 CFTR. A) A heatmap indicates the $\log_2$ (fold-change) in interactions of CFTR variants relative to WT. Increases in the association of interactors with CFTR variants relative to WT are indicated in yellow while blue indicates a decrease in relative abundance of interactors. Black indicates no change relative to WT. Scale shown in $\log_2$ fold change over WT abundance. Proteins are organized into 3 clusters according to a hierarchical clustering analysis that groups interactors based on changes in abundance. B) A network map depicts the relationships between interactors in cluster 1, which includes interactors the exhibit the largest changes in the context of ΔF508 RF_{mut} relative to ΔF508. Lines indicate known protein-protein interactions in the String database (human). Line widths indicate the strength of data support. The colors of nodes reflect the identity of sub-clusters from K-means clustering of the interactors within hierarchical cluster 1. The labels for each color summarize the most common classes of proteins within each sub-cluster. Interactors in gray were missing from the String database and were manually added in as isolated nodes.

There are several key differences in the interaction profiles of ΔF508 and ΔF508-RF_{mut} CFTR (Table S1). We
first note that about half of the aberrant interactions formed by ΔF508 CFTR are unaffected by these silent
mutations (see Cluster 1 in Fig. 3B & Clusters 2-3 in Fig. S4-S5). As both genetic constructs encode the same
protein in the 0-frame, we suspect these common interactions may arise from the post-translational effects of
the ΔF508 mutation on the CFTR protein. Nevertheless, ΔF508-RF_{mut} exhibits significantly attenuated
associations with eight ribosomal subunits (Fig. 3B, Table S1), which implies the frameshift motif normally
creates ribosome pile-ups on the ΔF508 transcript. These silent mutations also appear to modulate QC, as
ΔF508 RF_{mut} exhibits both weaker interactions with certain chaperones and co-chaperones (e.g. CANX, HspA6,
BAG2, and DNAJB6) and stronger interactions with others (e.g. HspB1, HspA4, and DNAJA3 & Fig. S5). These
QC modifications also coincide with a decrease in the interaction of ΔF508-RF_{mut} with the UBE3C ubiquitin ligase,
the ER-PHAGY protein RTN3, and various components of the 19S proteasome complex (Table S1), which
suggests the RF motif ultimately modifies CFTR degradation. Finally, we note these silent mutations enhance
the interaction of ΔF508 CFTR with two components of the eIF3 complex (i.e. eIF3c and eIF3e, Cluster 3, Fig.
S4, Table S1), which is known to modify the proteostatic effects of the ΔF508 mutation.[7] Overall, these results
suggest the ribosomal frameshift site in NBD2 acts as a junction where cotranslational misfolding triggers
changes in CFTR translation, QC, and degradation.
156
157 *Impact of Ribosomal Frameshifting on CFTR Expression*
158
159 To determine how these changes in cotranslational assembly impact CFTR proteostasis, we compared the
160 effects of the RF_{mut} modification on ΔF508 and WT CFTR expression. Quantitative western blot analyses reveal

Figure 4. Impact of ribosomal frameshifting on CFTR expression. The expression and maturation of transiently expressed CFTR variants were compared in CFBE41o- and HEK293T cells by western blot and flow cytometry. A) A representative western blot depicting the relative abundance of the mature (band C) and immature (band B) CFTR glycoforms of each indicated variant in CFBE41o- cells is shown. A GAPDH loading control is included for reference. B) A representative western blot depicting the relative abundance of the mature (band C) and immature (band B) CFTR glycoforms of each indicated variant in HEK293T cells is shown. A GAPDH loading control is included for reference. C) A bar graph depicts the average C: B band intensity ratio relative to WT in CFBE41o- cells expressing the indicated variants as determined by western blot (n = 3). Error bars reflect the standard deviation. D) A bar graph depicts the average total CFTR intensity (C+B) relative to WT in CFBE41o- cells expressing the indicated variants as determined by western blot (n = 3). Error bars reflect the standard deviation. E) A bar graph depicts the average intracellular CFTR immunostaining intensity relative to WT among HEK293T cells expressing the indicated variants as determined by flow cytometry (n = 6). Error bars reflect the standard deviation. F) A bar graph depicts the average surface CFTR immunostaining intensity relative to WT among HEK293T cells expressing the indicated variants as determined by flow cytometry (n = 6). Error bars reflect the standard deviation.

that modifications of the RF motif have minimal impact on the maturation of WT or ΔF508 CFTR in both CFBE41o- and HEK293T cells as judged by the relative abundance of the mature (C band) and immature (B band) glycoforms (Figs. 4 A-C). This observation likely reflects the fact that the RF$_{mut}$ constructs generate the same full-length protein as those containing the unmodified RNA structure. Nevertheless, integrated intensities suggest the ΔF508 RF$_{mut}$ transcript generates a modest (~20%), yet statistically insignificant increase in the total CFTR expression relative to the ΔF508 CFTR transcript (Figs. 4 A,B, and D). Consistent with this observation, flow cytometry measurements suggest this modest uptick in expression arises from a 19 ± 11% increase in the intracellular CFTR generated by ΔF508-RF$_{mut}$ relative to the unmodified ΔF508 (p = 0.007, Fig. 4 E). We note that there is no corresponding increase in surface CFTR levels generated by ΔF508-RF$_{mut}$ (Fig. 4F), which may again reflect the fact that this transcript still generates the same unstable ΔF508 CFTR protein. Notably, the intracellular immunostaining intensity of ΔF508-RF$_{mut}$ is indistinguishable from that of WT (Fig. 4E), which suggests this motif reduces the accumulation of ΔF508 CFTR in the ER. We note that all trends were quantitatively consistent in CFBE41o- and HEK293T cells (Figs. 4 & S6). Though these effects are subtle, together these results reveal that the RF motif selectively reduces the expression of ΔF508 CFTR potentially by promoting the premature termination of translation in response to misfolding.

Impact of Ribosomal Frameshifting on CFTR Function

To determine how the RF motif impacts CFTR function, we compared the effects of the RF$_{mut}$ modification on the ΔF508 and WT CFTR-mediated quenching of a cellular halide-sensitive yellow fluorescent protein (hYFP)- a common CFTR activity assay.[27] Briefly, we generated a series of recombinant stable HEK293T cell lines that inducibly express a single CFTR variant off a bicistronic transcript containing a downstream internal ribosomal entry site (IRES) that produces the hYFP sensor and a fluorescent mKate reference protein (Fig. 5A). We then used flow cytometry to track the loss of hYFP intensity relative to that of mKate at the single-cell level following activation of CFTR (Fig. 5B). Half-life values extracted from a global fit of hYFP quenching reveals that the RF$_{mut}$ modification has no impact on the function of WT CFTR (Fig. 5C). This observation is consistent with the limited impact of this modification on the interactome and expression of WT CFTR (Figs. 3A & 4). As expected, ΔF508 CFTR exhibits a sizable increase in quenching half-life relative to WT, which reflects its impaired channel conductance function (Fig. 5C). In contrast to WT, the RF$_{mut}$ mutations decrease the half-life of ΔF508 CFTR-mediated hYFP quenching by 46 ± 6% (Fig. 4 B & C). Given that the disruption of the RF motif does not appear to increase ΔF508 CFTR levels at the plasma membrane (Fig. 4C), the enhanced conductance of ΔF508-RF$_{mut}$ implies that the changes in its interactome enhance the functionality of the mature channel protein. Together, these cumulative findings reveal that the RF motif selectively modulates the assembly of ΔF508 CFTR in a manner that ultimately attenuates its expression and function.

Figure 5. Impact of ribosomal frameshifting on CFTR function. The functional conductance of stably expressed CFTR variants was compared in HEK293T cells by measuring the time-dependent quenching of a halide-sensitive yellow fluorescent protein (hYFP). A) A cartoon depicts a diagram of the stably expressed genetic cassette (top) and a schematic for the CFTR activity assay (bottom). Upon activation, iodide ions flow through the CFTR channel and quench the hYFP fluorophore, which is monitored by the decrease of hYFP intensity relative to that of the fluorescent mKate standard. B) HEK293T cells stably expressing WT (white), WT RF$_{mut}$ (gray), ΔF508 (red), and ΔF508 RF$_{mut}$ (blue) were stimulated with 25 μM forskolin (0.06% DMSO Vehicle) to activate CFTR prior to measurement of the change in cellular hYFP: mKate intensity ratio measurements over time by flow cytometry. Single cell intensity ratios are plotted against the time and the global fits of the decay are shown for reference. C) A bar graph depicts the globally fit half-life for hYFP quenching

Impact of Ribosomal Frameshifting on the Pharmacological Rescue of ΔF508 CFTR

The functional expression of ΔF508 and various other CF variants can be partially restored by Trikafta, an FDA-approved cocktail of two correctors that stabilize the CFTR protein (VX-661 + VX-445) and a potentiator that activates it (VX-770).[28-30] To determine whether ribosomal frameshifting potentially influences the effects of these compounds, we assessed whether the RF motif impacts the pharmacological rescue of ΔF508 CFTR. As expected, flow cytometry-based immunostaining measurements reveal that both WT and ΔF508 exhibit enhanced accumulation at the plasma membrane and within the secretory pathway upon stabilization by VX-661 + VX-445 (Fig. 6 A & B). A western blot analysis also confirms that these correctors enhance the maturation of both variants (Fig. 6C, Fig S7). However, similar gains in expression and maturation were observed for WT-RF$_{mut}$ and ΔF508-RF$_{mut}$ (Fig. 6 A-C, Fig. S7), which demonstrates that the RF motif has minimal impact on ΔF508 expression upon stabilization by corrector molecules. Next, we assessed functional rescue of CFTR conductance using our hYFP quenching assay. For this purpose, the potentiator VX-770 was added in combination with the correctors VX-661 and VX-445 to evaluate the functional rescue by Trikafta. Despite the comparable expression of ΔF508 and ΔF508-RF$_{mut}$ in the presence of correctors, HEK293T cells expressing ΔF508-RF$_{mut}$ that were treated with VX-661 + VX-445 + VX-770 exhibit a is 41 ± 5% decrease in hYFP quenching half-life relative to cells expressing ΔF508 CFTR under the same conditions (Fig. 6D). Together, these results suggest that, while stabilization by correctors minimizes the impact of the RF motif on ΔF508 CFTR expression, disrupting the RNA structure that promotes frameshifting ultimately enhances its functional rescue.

Discussion

The ΔF508 mutation promotes cotranslational CFTR misfolding and degradation in a manner that reduces its functional expression.[1] Nevertheless, its proteostatic effects also stem from aberrant translational dynamics. ΔF508 perturbs over 20 translation factors and RNA processing proteins involved in CFTR biosynthesis.[5] However, it remains unclear how the conformational effects of ΔF508 result in direct modifications of translation and vice versa. In light of the recent discovery that conformational transitions in the nascent chain can stimulate -1PRF,[14,15,31] we searched the CFTR transcript for structural features that might promote ribosomal frameshifting. Our findings identified a slippery sequence (SSC) and stem-loop (SL2) that stimulate ribosomal frameshifting and the premature translational termination (Fig. 2). To determine how this RF motif impacts CFTR expression and function, we designed a series of silent mutations that maintain the native amino acid sequence in the 0-frame while disrupting these putative RNA structures (RF$_{mut}$, Fig. 1). While this modification has minimal impact on WT CFTR, disrupting the RF motif significantly remodels ΔF508 CFTR interactome (Fig. 3). These changes in ΔF508 assembly enhance the accumulation of ΔF508 CFTR within the secretory pathway and increase its functional conductance at the plasma membrane (Figs. 4 & 5). Finally, we show that disrupting the RF motif also

Figure 6. Impact of ribosomal frameshifting on the pharmacological rescue of ΔF508 CFTR. The effects of CFTR modulators on the expression and function of CFTR variants was determined in HEK293T cells. A) A bar graph depicts the average intracellular CFTR variant immunostaining intensity relative to WT among HEK293T cells treated for 16 hours with vehicle (open bars) or 3 μM VX-661 + 3 μM VX-445 (hashes) as determined by flow cytometry (n = 6). B) A bar graph depicts the average intracellular CFTR variant immunostaining intensity relative to WT among HEK293T cells treated for 16 hours with DMSO vehicle (open bars) or 3 μM VX-661 + 3 μM VX-445 (hashes) as determined by flow cytometry (n = 6). C) A bar graph depicts the average CFTR variant C: B band intensity ratios normalized relative to WT in HEK293T cells treated for 16 hours with vehicle (open bars) or 3 μM VX-661 + 3 μM VX-445 (hashes) as determined by western blot (n = 3). D) A bar graph depicts the average hYFP quenching half-life for each variant as determined by flow cytometry (n = 3) in cells treated for 16 hours with vehicle (open bars) or 3 μM VX-661 + 3 μM VX-445 + 3 μM VX-770 (hashes) and pulsed with 25 μM forskolin. Measurements in the presence of vehicle from Figure 5C are shown for reference. Error bars reflect the standard deviation in all cases. All treatments resulted in a final concentration of 0.06% DMSO (v/v).

enhances the pharmacological rescue of ΔF508 CFTR function (Fig. 6). Given that the disruption of this motif has no impact on the assembly, expression, function, or pharmacological profile of WT CFTR (Figs. 2-5), we conclude that this RNA structure stimulates RF and the premature termination of translation in response to cotranslational CFTR misfolding. Together, our findings suggest the RF motif acts as a kill switch that down-regulates the synthesis of defective proteins in real time. Future investigations are needed to determine how this translational regulation influences the biosynthesis and pharmacological response of other known CF variants.

Interactome profiles suggest this RF motif may represent an assembly junction where the conformational state of the nascent chain modulates the interplay between translation and QC. We identify several ΔF508-specific chaperone interactions that are lost and others that are gained upon disruption of the RF motif (Table S1, Fig. 3). Moreover, weakening this RNA structure the context of the ΔF508 transcript significantly reduces the co-immunoprecipitation of several ribosomal subunits (Table S1, Fig. 3), which implies ribosomes typically stall within this region during ΔF508 translation. Based on these observations, we propose that the misassembly of nascent ΔF508 CFTR stimulates ribosomal frameshifting and directs the ribosome to a stop codon in the -1-frame (Fig. 7). Without the prompt intervention of ribosome quality control,[32] this premature termination causes trailing ribosomes to collide (Fig. 7). Such crosstalk may explain the link between cotranslational misfolding and ribosome collisions that occurs during pre-emptive QC.[11] Ribosomal pausing[11] and/or heightened frameshifting during ΔF508 CFTR translation could potentially explain how this mutation enhances associations with UPF1 and other nonsense-mediated decay (NMD) proteins[5] that typically promote the degradation of nonsense transcripts.[33,34] Knocking down UPF1 and other associated RNA binding proteins partially restores ΔF508 expression,[5] which until now has been a perplexing observation given that ΔF508 preserves the native reading frame. Nevertheless, the stimulation of frameshifting by ΔF508 would promote premature termination via stop codons in the -1 frame. Such a link between ribosomal frameshifting and NMD has been previously described,[35] and certain ribosome-associated molecular chaperones have been found to influence -1PRF.[36] Nevertheless, to our knowledge, previous investigations have not established a direct role of ribosomal frameshifting in protein QC. We note that these interpretations remain speculative given that we are unable to compare ribosomal frameshifting efficiencies during CFTR biosynthesis and that our interactome measurements do not reveal a clear link to NMD. Additional investigations are needed to gain mechanistic insights into the interplay between cotranslational misfolding, ribosomal frameshifting, and RNA surveillance pathways.

There are currently few examples of bona fide ribosomal frameshift sites within eukaryotic transcripts and the current evidence for such remains contentious.[24,37] One key line of evidence that has cast doubt on the role of ribosomal frameshifting in eukaryotic gene regulation involves the general lack of the extended coding sequences within the alternative reading frames of eukaryotic genes- a classic indication of translational recoding in viral transcripts. Due to an abundance of stop codons in the -1 reading frame, ribosomal frameshifting in eukaryotes should most often result in premature termination of translation- an outcome that ribosomes have generally evolved to avoid. Nevertheless, premature termination and/ or the destabilization of the transcript may be beneficial under conditions in which the nascent polypeptides begin to misfold. Cells routinely undergo swings in adaptive proteostasis where the products of translation and the pool of molecular chaperones they rely on

Figure 7. Proposed role of ribosomal frameshifting in the selective downregulation of ΔF508 CFTR biosynthesis. A cartoon depicts the proposed role of ribosomal frameshifting in CFTR proteostasis. Ribosomes pause at the ribosomal frameshift site at the point in which MSD1 and MSD2 undergo domain swapping. Synthesis proceeds with modest frameshifting when correct assembly occurs. However, failure to achieve the correct conformation and/ or chaperone interactions stimulates ribosomal frameshifting and the premature termination of termination of translation. Premature termination causes prolonged stalling that leads to collisions with trailing ribosomes and a failure to clear RNA surveillance proteins that trigger mRNA decay.

must be remodeled on short time scales. Based on our findings, we propose that this putative connection between cotranslational misfolding, ribosomal frameshifting, and RNA surveillance pathways provides a means for cells to remodel the transcriptome in response to misassembly in order to prevent the accumulation of misfolded proteins. Additional investigations are needed to determine whether similar QC checkpoints are present in the transcripts of other proteins that are predisposed to misfolding. Such investigations may offer novel mechanistic insights into the molecular basis of a wide array of diseases of aberrant cellular proteostasis.

Materials and Methods

Molecular Biology

All DNA constructs were cloned using the NEBuilder Hifi DNA Assembly System (NEB, cat # E2621X). All site directed mutagenesis reactions were performed using PrimeSTAR HS DNA Polymerase (Takara Bio, cat #R010B). Plasmids used for transient transfections in HEK293T cells were prepared and purified using the Zymopure Midiprep Kit (Zymo Research, cat #D4201). Mutations to create the CFTR ΔPRF constructs were chosen using RNA structure predictions performed by the ViennaRNA RNAfold Web Server.

Cell Culture

HEK293T cells were grown in Dulbecco's modified Eagle's medium (Gibco, Grand Island, NY) containing 10% fetal bovine serum (Corning, Corning, NY) and a penicillin (100U/ml)/streptomycin (100µg/ml) antibiotic supplement (Gibco, Grand Island, NY) in a humidified incubator containing 5% CO_2 at 37°C. HEK293T cells used for interactome measurements were grown under similar conditions but with 1% L-glutamine (200 mM) supplementation. Plasmid DNA constructs were transiently expressed in HEK293T cells using Lipofectamine 3000 (ThermoFisher Scientific, cat #L3000015). Cells were dosed with CFTR modulators a day after transfection when applicable. Two days post-transfection, cells were washed with 1X phosphate buffered saline (AthenaES, Baltimore, MD) and harvested with 1X 0.25% Trypsin-EDTA (Gibco, Grand Island, NY).

CFBE41o- cells were grown in minimal essential medium (Gibco, Grand Island, NY) containing 10% fetal bovine serum (Corning, Corning, NY) and a penicillin (100U/ml)/streptomycin (100µg/ml) (Gibco, Grand Island, NY) antibiotic supplement (Gibco, Grand Island, NY) in a humidified incubator containing 5% CO_2 at 37°C on culture plates coated with PureCol purified bovine collagen (Advanced BioMatrix, Carlsbad, CA). CFBE41o- cells were transfected using Lipofectamine 3000 (ThermoFisher Scientific, cat #L3000015). Two days post-transfection, cells were washed with 1X hepes buffered saline (Gibco, Grand Island, NY) and harvested with 1X 0.25% Trypsin-EDTA (Gibco, Grand Island, NY).

CFTR function measurements were carried out in recombinant stable cells were made from genetically modified HEK293T cells grown in 10 cm dishes in complete media as was previously described.[38,39] Briefly these HEK293T cells contain a Tet-Bxb1-BFP "landing pad" where recombination results in the integration of a plasmid into the gDNA of the cell.[38] These cells were co-transfected with plasmid and a Bxb1 recombinase expression vector using Fugene 6 (Promega, Madison, WI). Doxycycline (2µg/mL) was added one day after transfection and the cells were grown for 3 days at 33°C. The cells were then incubated at 37°C for 24 hours prior to passaging for further experiments.

Mass Spectrometry- Based Interactome Profiling

A pcDNA5 vector containing untagged CFTR with a CMV promoter was used in the interactome profiling experiments. Co-immunoprecipitation (co-IP) of CFTR bound with interactors was carried out as described previously.[25,40] Briefly, cell lysates were normalized by dilution to a common protein concentration at 1 mL total volume and pre-cleared with 4B Sepharose beads (Sigma) at 4°C for 2 h while rocking. Precleared lysates were then immunoprecipitated with Protein G beads covalently crosslinked to 24-1 antibody (6 mg antibody/mL of beads) overnight at 4°C while rocking. Beads were washed three times with TNI buffer, twice with TN buffer, all buffer was removed with a needle and then frozen at –80°C for 1.5 hours. Proteins were then eluted off the beads while shaking at 750 RPM at 37°C for 30-60 min with elution buffer (0.2 M glycine, 0.5% IGEPAL CA-630, pH 2.3). The elution step was repeated once and combined to immediately neutralize with a 10:1 ratio of 1 M ammonium bicarbonate solution.

MS sample preparation of co-IP samples were performed as described previously.[25,26] Briefly, samples were chloroform/methanol precipitated, rinsed with methanol, dried, and reconstituted in 1% Rapigest SF (Waters). Each sample was reduced with 5 mM TCEP (Sigma), alkylated with 10 mM iodoacetamide (Sigma), and digested with 0.5 µg of trypsin (Sequencing Grade, Promega, or Pierce) in 50mM HEPES (pH 8.0) at 37°C with shaking for a least 10 hours. Digested peptides diluted to 60 µL with water, labeled with 40 uL of TMT pro 16plex reagents (Thermo Fisher) for 1 hour, and quenched with 10% w/v ammonium bicarbonate. The sixteen TMT-labeled samples were pooled, acidified with MS-grade formic acid (Sigma), concentrated using a SpeedVac (Thermo Fisher), resuspended in 1500 µL of buffer A (95% water, 4.9% acetonitrile, and 0.1% formic acid), and loaded onto a triphasic MudPIT column. An Exploris 480 (Thermo Fisher) mass spectrometer equipped with an UltiMate3000 RSLCnano System (Thermo Fisher) was used for LC-MS/MS analysis as described previously.[25,26]

Proteome Discoverer 2.4 was used for peptide identification and TMT-based quantification as described previously.[26] MS/MS spectra were searched using SEQUEST against a UniProt human proteome database (released 03/25/2014) using a decoy database of reversed peptide sequences. The following parameters were employed: 0.02 Da fragment mass tolerance, 10 ppm peptide precursor tolerance, six amino acid minimum peptide length, trypsin cleavage with a max of two missed cleavages, static cysteine modification of 57.0215 Da (carbamidomethylation), and static N-terminal and lysine modifications of 304.2071 Da (TMT pro 16plex). Percolator was used to filter SEQUEST search to minimize the peptide false discovery rate to 1% and require a minimum of two peptides per protein identification. Reporter Ion Quantification processing node in Proteome Discoverer 2.4 was used to quantify TMT reporter ion intensities and a summation was performed for peptides in the same protein.

Interactor filtering and data analysis

A total of 5-6 sets of samples per condition (WT, WT-RF$_{mut}$, ΔF508, and ΔF508-RF$_{mut}$) were analyzed via LC-MS/MS over 3 separate mass spectrometry runs. Co-immunoprecipitated interactors were compared against GFP transfection control to obtain log2 fold enrichment and adjusted p-value. To filter for statistically significant interactors of CFTR, log2 fold change over 1 and adjusted p-value below 0.05 were used as cutoffs. Interactors in each condition were filtered and combined to generate a master list. This list was queried against the Contaminant Repository for Affinity Purification (CRAPome 2.0)[41] to remove proteins with a nonspecific detection threshold of over 500 reported experiments. The grouped abundance values of these interactors in each condition were log2 transformed to yield the consensus log2 grouped abundance for each protein. These values were normalized to WT CFTR log2 grouped abundance to standardize protein abundances across conditions.

Hierarchical clustering and network analysis

Hierarchical clustering of proteins in the interactor master list was performed with a custom R script as described previously.[26] Briefly, the log2 grouped abundance that were normalized against WT CFTR were converted to an Euclidean distance matrix and clustered using Ward's minimum variance method. Network plotting of each cluster was performed using the String database (string-db.org) with interaction score set at medium confidence (0.400). Interactors not found in the String database were added manually to the network plots as isolated nodes. K-means clustering was performed for each hierarchical cluster.

Luciferase-Based Ribosomal Frameshifting Measurements

Ribosomal frameshifting measurements were made using a previously described dual luciferase reporter system that was modified to avoid aberrant splicing artifacts.[24] Briefly, all dual luciferase reporters were generated from a pJD2044 plasmid containing firefly and renilla luciferase and a CMV promoter. Potential cryptic splice sites in

the renilla luciferase gene were identified using the human splicing finder webtool and were removed using a
series of synonymous codon mutations (246T>C, 249G>A, 888T>G, and 891G>A).[42] A 138 bp fragment of the
CFTR transcript (3963-4101) was inserted between the 5' renilla luciferase (0-frame) and 3' firefly luciferase (-1-
frame) genes. P2A linkers were inserted both upstream and downstream of the CFTR insert. HEK293T cells
were plated in six well dishes one day prior to transfection of reporter constructs using Lipofectamine 3000
(ThermoFisher Scientific, cat #L3000015). Two days post-transfection, cells expressing dual luciferase reporter
constructs were harvested and lysed using the Passive Lysis Buffer provided in the Dual-Luciferase Reporter
Assay System (Promega, cat #E1980). Relative activities of the firefly and renilla luciferase domains in the
clarified lysate were then measured using the Dual-Luciferase Reporter Assay System on a Synergy Neo2 plate
reader (Biotek, Winooski, VT) in accordance with the manufacturer's instructions. To determine the –1 ribosomal
frameshifting efficiency the firefly: renilla luciferase ratio was normalized relative to that of a construct lacking the
insert as was previously described.[43]
387
388 *Western Blot-Based CFTR Expression Measurements*
389
Western blotting was used to evaluate the relative abundance and maturation of CFTR glycoforms in both
HEK293T and CFBE41o- cells. Cells were grown in 6 cm dishes and transfected with Lipofectamine 3000
(ThermoFisher Scientific, cat #L3000015). Un-tagged CFTR variants were transiently expressed from a pcDNA5
vector featuring a CMV promoter. Cells were passively lysed two days post-transfection by rocking the cells at
4° C in a lysis buffer containing 25 mM Tris (pH 7.6), 150 mM NaCl, 1% Triton X-100, and 2mg/ mL of a protease
inhibitor cocktail (Roche Diagnostics, Indianapolis, IN). Total protein concentrations within the cleared lysates
were then determined using a detergent-compatible Bradford assay (Pierce Biotechnology, Waltham, MA).
Lysates were diluted into SDS-PAGE sample buffer and heated to 37°C for 30 min prior to loading 65µg of total
protein from each sample onto a 7.5% SDS-PAGE gel. Proteins were then separated by electrophoresis and
transferred onto a PVDF membrane overnight at 4°C. Following transfer, the membrane was blocked in a 5%
milk solution in TBST for 2 hours at room temperature. After two hours the lower part of the membrane was cut
to isolate the GAPDH loading control bands. The membranes were incubated in a wash solution containing either
a mouse anti-CFTR antibody (1:1,000 dilution AB217, CFTR Antibody Distribution Program, Chapel Hill, NC) or
a mouse anti-GAPDH antibody (1:10,000 dilution ab8245, Abcam, Cambridge, UK) overnight at 4°C. Both CFTR
and GAPDH containing membranes were then washed three times in TBST prior to incubation in a wash solution
containing an IRDye 680RD-labeled goat anti-mouse secondary antibody (1:10,000 dilution, LI-COR
Biosciences, Lincoln, NE) for two hours. Following incubation, the membranes were washed three more times
in a TBST wash solution and allowed to dry for 30 min prior to imaging using an Odyssey CLx system (LI-COR
Biosciences, Lincoln, NE). Band intensities were analyzed using both Image Studio V. 5.2 software (LI-COR
Biosciences, Lincoln, NE).
410
411 *Flow Cytometry-Based CFTR Expression Measurements*
412
Flow cytometry was used to compare the plasma membrane and intracellular expression levels of CFTR mutants
using a previously described approach.[44] Briefly, CFTR variants were transiently expressed in HEK293T cells
using a pcDNA5 vector featuring a CMV promoter, CFTR variants that were generated in the context of a
previously described triple hemagglutinin (HA) tag in the fourth extra-cellular loop,[45] and an internal ribosome
entry site (IRES)-eGFP cassette. Cells were transfected in 6 cm dishes for 24 hours prior to treatment with either
DMSO (vehicle) or 3µM VX-445 and VX-661. Cells were harvested 48 hours post transfection prior to
immunostaining the plasma membrane CFTR for 30 min. with a DyLight 550-conjugated anti-HA antibody
(ThermoFisher, Waltham, MA). Cells were then fixed, washed, and permeabilized using the Fix and Perm kit
(Invitrogen, Carlsbad, CA). Intracellular CFTR was the immunostained for 30 minutes in the dark using an Alexa
Fluor 647-conjugated anti-HA antibody (Invitrogen, Carlsbad, CA). Cells were washed and filtered prior to
analysis of cellular fluorescence profiles using a BD LSRII flow cytometer (BD Biosciences, Franklin Lakes, NJ).
Forward and side scatter measurements were used to gate for intact single cells. eGFP intensities (488 nm laser,
530/30 nm emission filter) were used to gate for positively transfected cells. Dylight 550 (561 nm laser, 582/15
nm emission filter) and Alexa Fluor 647 (640 nm laser, 670/30 nm emission filter) intensities were then measured
for at least 10,000 positively transfected cells within each biological replicate. Data was analyzed using FlowJo
software (Treestar, Ashland, OR).
429
430
431 *CFTR-Mediated hYFP Quenching Measurements*
432
433 A fluorescent quenching functional assay based on a previously described method[27,46] was used to compare the
434 relative functional conductance of CFTR mutants. A pcDNA5 vector bearing an attB recombination site in place

of its CMV promoter, untagged CFTR cDNA encoding each variant, and an IRES hYFP-P2A-mKate sensor cassette was used to generate a series of stable HEK293T cell lines that inducibly express each CFTR construct as previously described.[38,39] Recombinant HEK293T cells stably expressing CFTR were grown in six well dishes and dosed with either DMSO (vehicle) or 3μM VX-445 and VX-661 48 hours after plating. Following 18 hrs. of treatment with correctors, the cells were harvested with trypsin, washed once in PBS containing 5 mM EDTA (pH 7.4), washed twice in PBS containing 137 mM sodium gluconate (pH 7.4), then resuspended in PBS containing 137 mM sodium gluconate (pH 7.4), 25 μM Forskolin, 50 μM Genestein, and 10 μM VX-770. Cells were incubated in the resuspension media for 10 minutes prior to a final wash and resuspension in PBS containing 137 mM sodium gluconate (pH 7.4) prior to analysis. hYFP quenching was initiated with the addition of 25 mM sodium iodide immediately prior to the analysis of single-cell forward light scattering, side light scattering, hYFP fluorescence (488 nm laser, 530/30 nm emission filter), and mKate fluorescence (561 nm laser, 610/20nm emission filter) intensity values as a function of time using a BD LSRII flow cytometer (BD Biosciences, Franklin Lakes, NJ). Data was analyzed using FlowJo software (Treestar, Ashland, OR). OriginPro 2019 (Origin, Northampton, MA) was used to globally fit the decay of single-cell hYFP: mKate ratios as a function of time using the following exponential decay function:

$$y = y_0 + A_1^{-k_1 \cdot t} \qquad ;$$

where y is the hYFP: mKate fluorescence intensity ratio at time t, y_0 is the baseline hYFP: mKate fluorescence intensity ratio at $t \to \infty$, A_1 is the amplitude of the signal, and k_1 is the observed rate constant for the exponential decay. The value of y_0 was fixed in the global fit to an experimentally determined value for the hYFP: mKate intensity ratio of cells prior to their treatment with sodium iodide. The quenching half-life was calculated from the globally fit parameters using the following equation:

$$t_{1/2} = \frac{\ln(2)}{k_1} \qquad ;$$

where $t_{1/2}$ is the half-life and k_1 is the globally fit rate constant.

Acknowledgements

We thank Suchetana Mukhopadhyay, Jonathan Dinman, and Jeff Brodsky for helpful critiques of our initial findings. We thank Christiane Hassel for technical input and assistance. We also acknowledge the support of the Indiana University Flow Cytometry Core Facility and the Indiana University Center for Genomics and Bioinformatics. This research was supported in part by grant from the National Institute of General Medical Sciences to J. P. S. (R01GM138845).

References

Cheng, S. H. *et al.* Defective intracellular transport and processing of CFTR is the molecular basis of most cystic fibrosis. *Cell* **63**, 827-834 (1990).

Veit, G. *et al.* From CFTR biology toward combinatorial pharmacotherapy: expanded classification of cystic fibrosis mutations. *Mol Biol Cell* **27**, 424-433 (2016).

Du, K., Sharma, M. & Lukacs, G. L. The DeltaF508 cystic fibrosis mutation impairs domain-domain interactions and arrests post-translational folding of CFTR. *Nat Struct Mol Biol* **12**, 17-25 (2005).

Fiedorczuk, K. & Chen, J. Molecular structures reveal synergistic rescue of Delta508 CFTR by Trikafta modulators. *Science* **378**, 284-290 (2022).

Pankow, S. *et al.* F508 CFTR interactome remodelling promotes rescue of cystic fibrosis. *Nature* **528**, 510-516 (2015).

Hutt, D. M., Loguercio, S., Campos, A. R. & Balch, W. E. A Proteomic Variant Approach (ProVarA) for Personalized Medicine of Inherited and Somatic Disease. *J Mol Biol* **430**, 2951-2973 (2018).

490 7 Hutt, D. M., Loguercio, S., Roth, D. M., Su, A. I. & Balch, W. E. Correcting the F508del-CFTR variant by
491 modulating eukaryotic translation initiation factor 3-mediated translation initiation. *J Biol Chem* **293**,
492 13477-13495 (2018).
493

494 8 Veit, G. *et al.* Ribosomal Stalk Protein Silencing Partially Corrects the DeltaF508-CFTR Functional
495 Expression Defect. *PLoS Biol* **14**, e1002462 (2016).
496

497 9 Rauscher, R. *et al.* Positive epistasis between disease-causing missense mutations and silent
498 polymorphism with effect on mRNA translation velocity. *Proc Natl Acad Sci U S A* **118**, e2010612118
499 (2021).
500

501 10 Oliver, K. E. *et al.* Slowing ribosome velocity restores folding and function of mutant CFTR. *J Clin Invest*
502 **129**, 5236-5253 (2019).
503

504 11 Lakshminarayan, R. *et al.* Pre-emptive Quality Control of a Misfolded Membrane Protein by Ribosome-
505 Driven Effects. *Curr Biol* **30**, 854-864 e855 (2020).
506

507 12 Kim, S. J. & Skach, W. R. Mechanisms of CFTR Folding at the Endoplasmic Reticulum. *Front Pharmacol*
508 **3**, 201 (2012).
509

510 13 Bartoszewski, R. A. *et al.* A synonymous single nucleotide polymorphism in DeltaF508 CFTR alters the
511 secondary structure of the mRNA and the expression of the mutant protein. *J Biol Chem* **285**, 28741-
512 28748 (2010).
513

514 14 Harrington, H. R. *et al.* Cotranslational folding stimulates programmed ribosomal frameshifting in the
515 alphavirus structural polyprotein. *J Biol Chem* **295**, 6798-6808 (2020).
516

517 15 Carmody, P. J. *et al.* Coordination of -1 programmed ribosomal frameshifting by transcript and nascent
518 chain features revealed by deep mutational scanning. *Nucl Acids Res* **49**, 12943-12954,
519 doi:10.1093/nar/gkab1172 (2021).
520

521 16 Penn, W. D., Harrington, H. R., Schlebach, J. P. & Mukhopadhyay, S. Regulators of Viral Frameshifting:
522 More Than RNA Influences Translation Events. *Annu Rev Virol* **7**, 219-238 (2020).
523

524 17 Giedroc, D. P. & Cornish, P. V. Frameshifting RNA pseudoknots: structure and mechanism. *Virus Res*
525 **139**, 193-208 (2009).
526

527 18 Brierley, I., Jenner, A. J. & Inglis, S. C. Mutational analysis of the "slippery-sequence" component of a
528 coronavirus ribosomal frameshifting signal. *J Mol Biol* **227**, 463-479 (1992).
529

530 19 Sharma, V. *et al.* Analysis of tetra- and hepta-nucleotides motifs promoting -1 ribosomal frameshifting in
531 Escherichia coli. *Nucl Acids Res* **42**, 7210-7225 (2014).
532

533 20 Bock, L. V. *et al.* Thermodynamic control of -1 programmed ribosomal frameshifting. *Nat Commun* **10**,
534 4598 (2019).
535

536 21 Miller-Vedam, L. E. *et al.* Structural and mechanistic basis of the EMC-dependent biogenesis of distinct
537 transmembrane clients. *Elife* **9**, e62611 (2020).
538

539 22 Smith, A. M., Costello, M. S., Kettring, A. H., Wingo, R. J. & Moore, S. D. Ribosome collisions alter
540 frameshifting at translational reprogramming motifs in bacterial mRNAs. *Proc Natl Acad Sci U S A* **116**,
541 21769-21779 (2019).
542

543 23 Grentzmann, G., Ingram, J. A., Kelly, P. J., Gesteland, R. F. & Atkins, J. F. A dual-luciferase reporter
544 system for studying recoding signals. *RNA* **4**, 479-486 (1998).

Khan, Y. A. *et al.* Evaluating ribosomal frameshifting in CCR5 mRNA decoding. *Nature* **604**, E16-E23, (2022).

McDonald, E. F., Sabusap, C. M. P., Kim, M. & Plate, L. Distinct proteostasis states drive pharmacologic chaperone susceptibility for cystic fibrosis transmembrane conductance regulator misfolding mutants. *Mol Biol Cell* **33**, ar62 (2022).

Kim, M. *et al.* Elexacaftor/VX-445-mediated CFTR interactome remodeling reveals differential correction driven by mutation-specific translational dynamics. *bioRxiv* (2023).

Galietta, L. J., Haggie, P. M. & Verkman, A. S. Green fluorescent protein-based halide indicators with improved chloride and iodide affinities. *FEBS Lett* **499**, 220-224 (2001).

Middleton, P. G. *et al.* Elexacaftor-Tezacaftor-Ivacaftor for Cystic Fibrosis with a Single Phe508del Allele. *New Eng J Med* **381**, 1809-1819 (2019).

Keating, D. *et al.* VX-445-Tezacaftor-Ivacaftor in Patients with Cystic Fibrosis and One or Two Phe508del Alleles. *New Eng J Med* **379**, 1612-1620 (2018).

Veit, G. *et al.* Allosteric folding correction of F508del and rare CFTR mutants by elexacaftor-tezacaftor-ivacaftor (Trikafta) combination. *JCI Insight* **5**, e139983 (2020).

Bhatt, P. R. *et al.* Structural basis of ribosomal frameshifting during translation of the SARS-CoV-2 RNA genome. *Science* **372**, 1306-1313 (2021).

Brandman, O. & Hegde, R. S. Ribosome-associated protein quality control. *Nat Struct Mol Biol* **23**, 7-15 (2016).

Hogg, J. R. & Goff, S. P. Upf1 senses 3'UTR length to potentiate mRNA decay. *Cell* **143**, 379-389 (2010).

Zund, D., Gruber, A. R., Zavolan, M. & Muhlemann, O. Translation-dependent displacement of UPF1 from coding sequences causes its enrichment in 3' UTRs. *Nat Struct Mol Biol* **20**, 936-943 (2013).

Belew, A. T. & Dinman, J. D. Cell cycle control (and more) by programmed -1 ribosomal frameshifting: implications for disease and therapeutics. *Cell Cycle* **14**, 172-178 (2015).

Muldoon-Jacobs, K. L. & Dinman, J. D. Specific effects of ribosome-tethered molecular chaperones on programmed -1 ribosomal frameshifting. *Eukaryot Cell* **5**, 762-770 (2006).

Belew, A. T. *et al.* Ribosomal frameshifting in the CCR5 mRNA is regulated by miRNAs and the NMD pathway. *Nature* **512**, 265-269 (2014).

Matreyek, K. A., Stephany, J. J. & Fowler, D. M. A platform for functional assessment of large variant libraries in mammalian cells. *Nucl Acids Res* **45**, e102 (2017).

Penn, W. D. *et al.* Probing biophysical sequence constraints within the transmembrane domains of rhodopsin by deep mutational scanning. *Sci Adv* **6**, eaay7505 (2020).

Pankow, S., Bamberger, C., Calzolari, D., Bamberger, A. & Yates, J. R., 3rd. Deep interactome profiling of membrane proteins by co-interacting protein identification technology. *Nat Protoc* **11**, 2515-2528 (2016).

Mellacheruvu, D. *et al.* The CRAPome: a contaminant repository for affinity purification-mass spectrometry data. *Nat Methods* **10**, 730-736 (2013).

601   42   Desmet, F. O. *et al.* Human Splicing Finder: an online bioinformatics tool to predict splicing signals. *Nucl*
602   *Acids Res* **37**, e67 (2009).
603

604   43   Jacobs, J. L. & Dinman, J. D. Systematic analysis of bicistronic reporter assay data. *Nucl Acids Res* **32**,
605   e160 (2004).
606

607   44   Roushar, F. J. *et al.* Contribution of Cotranslational Folding Defects to Membrane Protein Homeostasis.
608   *J Am Chem Soc* **141**, 204-215 (2019).
609

610   45   Pedemonte, N. *et al.* Small-molecule correctors of defective DeltaF508-CFTR cellular processing
611   identified by high-throughput screening. *J Clin Invest* **115**, 2564-2571 (2005).
612

613   46   Vijftigschild, L. A., van der Ent, C. K. & Beekman, J. M. A novel fluorescent sensor for measurement of
614   CFTR function by flow cytometry. *Cytometry A* **83**, 576-584 (2013).
615