Extreme mitochondrial reduction in a novel group of free-living metamonads.

Shelby K. Williams[1,2], Jon Jerlström Hultqvist[1,3], Yana Eglit[1,4], Dayana E. Salas-Leiva[1,5], Bruce Curtis[1,2], Russell Orr [6], Courtney W. Stairs[7], Alastair G. B. Simpson[1,4] and Andrew J. Roger[1,2,*]

- Institute for Comparative Genomics, Dalhousie University, Nova Scotia, Canada
– Department of Biochemistry and Molecular Biology, Dalhousie University
– Department of Cell and Molecular Biology, Uppsala University, Sweden
- Department of Biology, Dalhousie University
– Department of Biochemistry, Cambridge University, UK
- Section for Genetics and Evolutionary Biology, Department of Biosciences, University of Oslo, Norway
– Department of Biology, Lund University, Sweden

*Corresponding author: andrew.roger@dal.ca

Abstract

Metamonads are a diverse group of heterotrophic microbial eukaryotes adapted to living in hypoxic environments. All metamonads but one harbour metabolically altered 'mitochondrion-related organelles' (MROs) with reduced functions relative to aerobic mitochondria, however the degree of reduction varies markedly over the metamonad tree. To further investigate metamonad MRO diversity, we generated high quality draft genomes, transcriptomes, and predicted proteomes for five recently discovered free-living metamonads. Phylogenomic analyses place these organisms in a clade sister to the Fornicata – a group of metamonads that includes parasitic and free-living diplomonads and *Carpediemonas*-like organisms. Extensive bioinformatic analyses of the manually curated gene models showed that these organisms have extremely reduced MROs in comparison to other free-living metamonads. Loss of the mitochondrial iron-sulfur cluster (ISC) assembly system in some organisms in this group appears to be linked to the acquisition in their common ancestral lineage of a SUF-like minimal system (SMS) Fe/S cluster pathway through lateral gene transfer (LGT). One of the isolates, named 'RC', appears to have undergone even more drastic mitochondrial reduction losing almost all other detectable MRO-related functions. The extreme mitochondrial reduction observed within this free-living anaerobic protistan clade is unprecedented and demonstrates that mitochondrial functions, under some conditions, can be almost completely lost even in free-living organisms.

Introduction

Many unicellular eukaryote (protist) lineages have independently adapted to low-oxygen conditions. One of their most conspicuous adaptations are highly modified mitochondria referred to as "mitochondrion-related organelles" (MROs) that can function without oxygen. MROs have been best studied in the Metamonada, the largest anaerobic protistan group in the eukaryotic tree of life which encompasses a wide array of organisms with differing lifestyles including parasites, commensals, and free-living marine or freshwater flagellates or amoebae. All metamonad MROs studied to date are functionally reduced compared to aerobic mitochondria with their properties varying markedly across the group (Leger et al., 2017).

The best studied metamonad MROs are hydrogenosomes that occur in the urogenital tract parasite *Trichomonas vaginalis*. The 'hydrogenosomal' ATP-producing pathway uses enzymes typically not found in aerobic mitochondria such as pyruvate:ferredoxin oxidoreductase (PFO) to oxidatively decarboxylate pyruvate to acetyl-CoA and reduce ferredoxin, an acetate:succinate CoA transferase that catalyzes the transfer of CoA onto succinate to produce succinyl-CoA, and an iron-only [FeFe]-hydrogenase that reduces protons to hydrogen gas by re-oxidizing the ferredoxin. ATP is produced from ADP and P_i by the Krebs cycle enzyme succinyl-CoA synthetase by substrate-level phosphorylation, converting succinyl-CoA to succinate in the process. Trichomonad hydrogenosomes also house a variety of other pathways including a mitochondrial-type iron-sulfur cluster (ISC) biogenesis system, amino acid metabolism, and oxygen detoxification (Schneider et al., 2011; Stairs et al., 2015). Even more highly-reduced MROs called 'mitosomes' are found in the metamonad gut parasite *Giardia*. These organelles appear to lack any capacity to generate ATP; instead they participate in Fe/S cluster biogenesis via a mitochondrial-type ISC system (Jedelský et al., 2011). Like trichomonads, *Giardia* catalyzes acetyl-CoA production from pyruvate using PFO. However, this occurs in the cytoplasm. Acetyl-CoA is converted to acetate and ATP is produced by substrate-level phosphorylation catalyzed by the enzyme ADP-forming acetyl-CoA synthetase (ACS; EC 6.2.1.13) (Sánchez et al., 2000). The most extreme form of MRO reduction in Metamonada can be found in the oxymonad gut commensal *Monocercomonoides exilis* that has lost all traces of the organelle, making it the first truly amitochondriate eukaryote discovered (Karnkowska et al., 2016).

More recent investigations of newly discovered free-living metamonads have revealed novel configurations of MRO functions that do not fit into traditionally-defined organelle classes (Müller et al., 2012). For example, the free-living metamonad *Dysnectes brevis* encodes MRO targeted [FeFe]-hydrogenases and associated proteins but lacks organelle-targeted enzymes for substrate-level phosphorylation, suggesting that while the MRO is capable of hydrogen metabolism, ATP is most likely produced in the cytosol (Leger et al., 2017). A similar hydrogen-producing organelle and predicted cytosolic ATP production was reported in a transcriptomic investigation of the free-living metamonad *Barthelona* sp. PAP020 (Yazaki et al., 2019). Recently, the enigmatic *Anaeramoeba* species – recently-discovered distant relatives to parabasalids like *Trichomonas* – were predicted to possess hydrogenosomes that have retained more mitochondrial systems than any other metamonad studied to date (Stairs et al., 2021). The

MRO of the freshwater flagellate *Paratrimastix pyriformis* was recently investigated using spatial proteomic techniques which revealed that the organelle does not participate in either ATP production through extended glycolysis or Fe/S cluster production, and instead produces essential intermediates for the methionine cycle (Zítek et al., 2022). These studies of newly discovered free-living metamonads have provided important insights into the steps by which dramatically distinct MRO-types evolved within the Metamonada.

The multiple independent transitions from aerobically respiring mitochondria to MROs in the eukaryote tree in many cases involved the complete loss of the mitochondrial genome itself, the loss of aerobic metabolism genes encoded in the nucleus, and the gain of genes encoding novel biochemical pathways by lateral gene transfer (LGT) from bacteria, archaea or other anaerobic protists (Roger et al. 2017). For example, the 'hydrogenosomal' metabolism enzymes (Leger et al., 2016; Stairs et al., 2011, 2021), the ability to synthesize rhodoquinone (Stairs et al., 2018) and oxygen defense enzymes (Jiménez-González et al., 2019) all appear to have been acquired by LGT in multiple distinct lineages from bacteria or other protists (Gawryluk & Stairs, 2021; Roger et al., 2017). The most striking examples of the remodeling of mitochondrial functions in anaerobic protists involve the wholesale replacement of the highly-conserved essential mitochondrial ISC system by laterally acquired prokaryotic Fe/S systems. In the Archamoebae, a two-protein FeS biosynthesis pathway (recently classified as the minimal iron sulfur system (MIS) related to the nitrogen fixation system (Garcia et al. 2022)) was acquired from bacteria and, in free-living members of the group, was subsequently duplicated into cytosolic and MRO-functioning versions (Nývltová et al., 2015; Záhonová et al., 2022). A completely independent replacement of the mitochondrial ISC system occurred in the ancestors of the breviate flagellate *Pygsuia biforma* that acquired an archaeal-type SUF system (Stairs et al., 2014), recently classified as the SUF-like minimal system (SMS) pathway (Garcia et al. 2022). The ancestral SmsCB gene (referred to as SufCB in Stairs et al. 2014) was duplicated to create cytosolic and MRO-targeted SmsCB proteins. In the oxymonads, yet another replacement of the ISC system occurred independently by the lateral acquisition of genes encoding a more complex multi-protein SUF pathway. This enabled the ancestral oxymonad to produce Fe/S clusters in the cytosol, easing the evolutionary pressure to keep the canonical ISC system, paving the way for the complete loss of MROs in *Monocercomonoides* (Karnkowska et al., 2016). Phylogenetic analyses revealed that this SUF system is shared by a variety of oxymonads and is likely bacterial in origin (Vacek et al., 2018).

To further investigate the patterns and process of genome evolution associated with adaptation to anaerobiosis in eukaryotes, we investigated a set of five recently-discovered metamonad flagellates: three of these organisms are, as yet, unnamed and one is the recently described *Barthelona* sp. PAP020 (Yazaki et al., 2019). Unlike most previous studies of this kind that rely on incomplete data from transcriptomics, we generated high quality draft genomes, transcriptomes, and predicted proteomes from four of these five organisms. Our phylogenomic analyses demonstrate that these five metamonads form a highly supported clade herein referred to as 'BRC' (Barthelonids and Retortacarps) that emerges as a sister group to the Fornicata (diplomonads, retortamonads and *Carpediemonas*-like organisms (CLOs)). Genome size and coding capacity varies considerably across these organisms with one species possessing the

smallest genome of a free-living eukaryote currently known. The anaerobic metabolisms and MRO functional capacities of these organisms also vary considerably. Unexpectedly, we found yet another instance of the evolutionary acquisition of a Sms system in their common ancestor. While several BRCs retain both the Sms system and core parts of a mitochondrial-type ISC system, all traces of the ISC system have been lost in the others. For one of these species, the MRO proteome appears to be extremely reduced, with only two candidate MRO proteins identifiable. Overall, this work showcases heretofore unknown extreme plasticity in nuclear genome size and the mitochondrial function in free-living eukaryotes.

Results & Discussion

Three of the studied isolates could not immediately be placed into a known taxon and are referred to as "Retortacarp" (RC), "GEM-Retortacarp" (GEMRC), "Retortacarp-like" (RCL). All three "RC" isolates are biflagellated cells with a conspicuous posterior spike, and gently twisted groove down the 'ventral' side (Figure 1A-C). The longer posterior flagellum beats within and above the groove. The posterior part of the groove forms a very broad cytopharynx that opens laterally then narrows and curves anteriorly up the dorsal side of the cell. The "RC" isolates have some similarity to barthelonids, which lack the spike but which are also biflagellates with a thin ventral groove and posterior, curving cytopharynx. The groove and cytopharynx are much subtler in barthelonids, however, and nearly invisible in the very small isolate PCE, the barthelonid for which we resolved the most complete genome assembly (Figure 1D, Yazaki et al. 2019).

Near-complete draft genomes of four novel metamonads

The draft genome assemblies of PCE, RC, RCL and GEMRC were generated using a combination of Oxford Nanopore long-read sequencing for assembly and Illumina short-read sequencing for error-correction post-assembly, followed by decontamination. Genes were predicted for each of these assemblies using an in-house gene prediction pipeline. Briefly, RepeatMasker (Flynn et al., 2020) was used to mask the repetitive regions of the genome. RNA-seq reads were then mapped to the genome. GeneMark-ET (Lomsadze et al., 2014) then utilized this mapping information to generate gene coordinates and intron splice sites, creating a training set of genes. This training set was then used to train AUGUSTUS (Stanke et al., 2004), which generated the gene predictions. PASA (Haas et al., 2003) was then used to validate the coordinates of these gene models.

Many reference-independent methods for assessing genome completeness rely on unassembled sequencing reads that are completely free from contaminated reads. To ensure our reads were thoroughly decontaminated, we subjected the RNA-seq and genomic Illumina-derived read set for the four organisms to the Eukfinder workflow (Zhao, Salas-Leiva & Williams, in prep.). We then utilized Merqury, a k-mer counting tool that estimates genome completeness using genomic Illumina reads, to assess the completeness of the four *de novo* long-read assemblies. Merqury estimated completeness were above 98% for each of the four BRCs sequenced. We also found

that most eukaryotic-classified RNAseq reads mapped to their respective genome, with each of the four assemblies having >99% of their respective RNAseq reads mapping.

In addition, two of the genome assemblies have potential telomeric repeats on the ends of some contigs. PCE possesses three contigs with what appear to be telomeres at one end of the contig (telomere repeat sequence: 5'-TATATGGTCT-3'). The GEMRC assembly has six telomere-to-telomere assembled chromosomes (telomere repeat sequence: 5'-TTAGGG-3') and an additional nine contigs with telomeres at only one end of the contig. To further assess genome completeness, we completed a BUSCO analysis for the four *de novo* genomes. The resulting scores ranged from 52% of BUSCOs identified in PCE to 67% in GEMRC (see Table 1). Finally, the four assemblies are highly contiguous and well covered by both long-read and short-read sequences. See Table 2 for a summary of the *de novo* genomes and their completeness. The *Barthelona* sp. PAP020 genome assembly was based on Illumina reads only and is also listed in Table 2 for comparison. The size of the draft genomes and the number of proteins predicted differ markedly between taxa (Table 2). Notably, at 10.2 Mbp, PCE appears to possess the smallest genome for a free-living eukaryote currently known. It is smaller than the compact genomes of the diplomonad parasite *Giardia intestinalis* (11.7 Mbp, see Xu et al., 2020), the acidophilic alga *Galdieria sulphuraria* (13.7 Mbp, see Schönknecht et al., 2013), *Saccharomyces cerevisiae* (12.07 Mbp, see Goffeau et al., 1996), and the genome of the smallest known free-living eukaryote of the marine alga *Ostreococcus tauri* (12.56 Mbp, see Derelle et al., 2006). While genome reduction is well known to occur in parasitic and endosymbiotic microbes (Manzano-Marín & Latorre, 2016; Xu et al., 2020), it is less common in free-living organisms.

A BUSCO analysis was also performed on the predicted protein sets and transcriptomes of each BRC member to assess the performance of our gene modeling pipeline. Additionally, the predicted protein set for RC was manually curated, to assess improvement of the predicted protein set after curation (see Table 1). These results show that the number of BUSCO genes identified in these predicted proteins sets ranges from 63% complete for PCE to 78% complete for RCL. In general, the BUSCO scores of the predicted protein sets are similar to those from the assembled transcriptomes. In fact, the predicted protein set seems to reduce the fragmentation of identified BUSCO genes relative to the transcriptomes in the case of RC and PCE. These BUSCO scores, including those calculated for the genomes, are comparable to other well-characterized metamonad datasets (Salas-leiva et al., 2021; Stairs et al., 2021). Low BUSCO scores reflect that metamonads generally have divergent genomes and orthologs are either completely missing or too divergent to be detected. Note that the manual curation of RC predicted proteins increased the reported BUSCO number by only one, indicating that the gene prediction pipeline is likely accurately capturing the protein coding content of these genomes. This analysis highlights how BUSCO scores for novel and divergent eukaryotes are often misleading when taken at face value. Instead, in these cases, BUSCO scores are more useful when comparing genome, transcriptome, and proteome completion across a single species, or group of closely related organisms.

A draft genome assembly of PAP020 was also completed. This genome was assembled solely from Illumina reads, and thus is much more fragmented than the genomes of the other

metamonads considered in this study. These data, along with the PAP020 transcriptome (Yazaki et al. 2020) were used to supplement our analyses concerning MRO reconstruction, metabolic pathway inventory, and phylogenomic inference, as discussed below.

The BRCs are sister to, or the deepest branching group within Fornicata

To determine the phylogenetic affinities of the BRCs within the metamonads, we added orthologs from the five draft assemblies to a previously constructed phylogenomic data set of orthologous proteins (Brown et al., 2018; S. Kang et al., 2017). After single-protein alignment manual curation to remove contaminants and paralogs, the final dataset of 174 highly conserved proteins was analyzed by maximum likelihood under the site-heterogeneous model LG+C60+F+Γ (see Figure 1). The resulting tree confirms the placement of *Barthelona* sp. strain PAP020 as previously reported (Yazaki et al., 2019) and placed barthelonid strain PCE as its immediate sister. The other BRCs form a clade with the barthelonid group, with RC and GEMRC branching together to the exclusion of RCL. The 'BRC clade' (barthelonids + the Retortacarp clade) are sister to fornicates with maximum support (i.e. both the fornicate group and the fornicate + BRC clade are supported with 100% non-parametric bootstrapping support).

Members of Fornicata possess a particular cytoskeletal feature, a distinct fibrous arch, that defines the group (Simpson, 2003). As there are no ultrastructural data for the flagellar apparatus cytoskeleton of any BRC member to date, it is unclear whether they should be classified within the Fornicata or remain as a distinct sister group to the fornicates. Regardless, the BRCs 'deep' placement as a sister to Fornicata make them a key group for investigations into the origins and evolutionary trajectories of genes involved in parasitism and MRO reduction within the Metamonada.

The BRCs all possess highly reduced MROs and share the same core anaerobic metabolism

We searched the genomes, transcriptomes, and predicted protein sets of all four BRCs using a variety of bioinformatic approaches (see Experimental Procedures) to recover proteins related to MRO function and anaerobic metabolism. We also utilized a draft short-read Illumina genome assembly of PAP020 in these searches to supplement our comparisons. In short, we utilized several mitochondrial targeting sequence prediction programs to estimate the probability of mitochondrial localization of each predicted protein. We used these predicted proteins as queries against the nr database using BLAST and looked for outputs relating to mitochondrial localizing proteins. We also used BLAST to search through a database of known mitochondrial proteins. Next, we utilized datasets of known metamonad MRO proteins to similarly query the BRCs predicted proteins, as well as the genomes and transcriptomes. This was done through both BLAST search and HMM profile searching. In cases where search results were ambiguous, we utilized appropriate Pfam profiles to search for protein domain information. Finally, we assessed candidate proteins by constructing phylogenetic trees of corresponding MRO proteins with additional cytosolic homologs, to further verify the probable localization of BRC proteins. A summary of the results of these searches can be found as Figure 2. Below we discuss the metabolic pathways that we systematically searched for in the BRC data starting with pathways that, when present, function exclusively in MROs.

We first investigated whether the BRC genomes encoded homologs of the glycine cleavage system (GCS), an exclusively mitochondrial pathway which converts glycine to ammonia and carbon dioxide while also reducing NAD+ to NADH and generating 5,10-methylene-tetrahydrofolate (5,10-CH$_2$-THF) (Kikuchi, 1973). This pathway is commonly retained in MROs of free-living anaerobic protists including metamonads (reviewed in Roger et al., 2017). Except for RC, all BRCs encode the complete GCS (i.e. the T-, P-, L- and H-proteins). Additionally, all BRCs possessing the GCS also have a single serine hydroxymethyltransferase (SHMT) gene. This enzyme produces serine using both glycine and the 5,10-CH2-THF produced by the GCS. GCS and SHMT are found in the MROs of a variety of diverse free-living metamonads including *Paratrimastix pyriformis* (Zítek et al., 2022; Zubáčová et al., 2013) and all of the non-diplomonad fornicates (i.e. CLOs and *Chilomastix cuspidata*; see Leger et al., 2017). Complete absence of the GCS in metamonads is seen in the parasitic diplomonads *Giardia intestinalis* and *Spironucleus salmonicida* (Jerlström-Hultqvist et al., 2013; Leger et al., 2017). Other parasitic metamonads such as *Trichomonas vaginalis* retain some of the GCS subunits (Mukherjee et al., 2006), but have repurposed them to function in a peroxide detoxification pathway (Nývltová et al., 2016). It was particularly unexpected, therefore, that we found neither genes encoding GCS or SHMT in RC making it the only known free-living metamonad completely lacking this pathway, and the only BRC member to have no detectable MRO amino acid metabolism. The MROs within the BRCs are therefore key to understanding the reduction of amino acid metabolism that is usually only ever seen in parasitic metamonads. As previously reported (Yazaki et al., 2019), PAP020 contains a OsmC homolog, but we were unable to detect this enzyme in any of the other BRCs data, including the PAP020 genome. Because this enzyme lacks a mitochondrial targeting sequence (MTS), it is unclear where this protein localizes in PAP020, and if it participates in peroxide detoxification in the PAP020 MRO, as it does in the MRO of *Trichomonas* (Nývltová et al., 2016).

Next, we investigated the highly conserved mitochondrial Fe/S cluster (ISC) pathway that is also exclusively found within mitochondria or MROs. Unexpectedly, amongst all the BRCs, only the barthelonid PCE and PAP020 assemblies encoded *any* detectable ISC system components. These two genomes encode cysteine desulfurase NFS1 (also known as ISCS), the scaffold protein ISCU and a ferredoxin. Notably, they also encode Fe/S cluster containing NuoE and NuoF subunits of complex I, which, like core ISC components, are homologues of which are only known to function in mitochondria and the MROs of other organisms. Surprisingly, with the exceptions of mitochondrial hsp70 (discussed further below) and ferredoxin, no other genes encoding ISC pathway components were detected in the barthelonids or any other of the BRCs (see Supplementary Table 1 for ISC proteins queried). The absence of some of these proteins, like Isd11 and ferredoxin reductase (Arh1), is not unexpected as they have not been detected in most metamonads (Motyčková et al., 2022). The absence of other early ISC pathway proteins implicated in [2Fe-2S] cluster synthesis (e.g. frataxin, Jac1 (HscB) and Mge1 (GrpE)) is more puzzling as these components are usually detected in metamonads (Motyčková et al., 2022). Similarly, some late ISC pathway components implicated in [4Fe-4S] cluster synthesis (Lill & Freibert, 2020) are found in other metamonads (e.g Grx5, Isa2, NFU1 and BolA; Motyčková et al., 2022), but were not detected in any BRCs (Supplementary Table 1). This is strange given that the barthelonids possess mitochondrial NuoF subunits that typically house an [4Fe-4S]

cluster (Ohnishi, 1998). This means that if these NuoF subunits are indeed found within the MROs of the barthelonids, as they are with other eukaryotes, their [4Fe-4S] clusters are being produced by an unknown mechanism.

Despite the lack of ISC system proteins in most BRCs, all BRC members appear to encode a similar partial cytosolic CIA system that includes apoprotein targeting components CIA1, CIA2, and NAR1, the NBP35 scaffold protein, and TAH18, which aids in maturation of Fe/S cluster proteins. This reduced CIA pathway is not unlike the CIA pathway found within *Giardia intestinalis* (Pyrih et al., 2016).

BRCs possess simple archaeal-type SUF-like minimal system (SMS) fusion proteins.

We investigated the possibility that the lack of ISC system components in BRC members may be complemented by the presence of an alternative Fe/S cluster system such as the minimal Fe/S (MIS) or sulfur mobilization (SUF) systems found in a number of other anaerobic protists (Anwar et al., 2014; Stairs et al., 2014; Vacek et al., 2018; Žárský et al., 2021; Garcia et al. 2022). In all of the BRC genomes, we were able to identify a simple archaeal-type SUF-like minimal system (SMS) consisting of a gene encoding an SmsCB fusion protein that was most similar to the simple archaeal-type SmsCB systems found in the breviate *Pygsuia biforma* (Stairs et al., 2014), the anaerobic jakobid *Stygiella incarcerata* (Leger et al., 2016) and gut commensal opilinatan Stramenopiles (Tsaousis et al., 2012; Yubuki et al., 2020). To investigate the origin of this protein, we performed phylogenetic analysis separately for the SmsC and SmsB domains (Supplementary Figures 1 & 2) before concatenating the domains together for the final tree (Figure 3). The BRC SmsCB proteins form a strongly supported clade that is sister group to all archaeal-type SmsCB fusion proteins from anaerobic protists. Consistent with previous reports, the anaerobic protistan clade of SmsCB fusion proteins is sister group to the orthologs from members of the Methanomicrobiales order of Archaea (Tsaousis et al., 2012). This phylogenetic pattern is most easily explained if there were an original transfer and fusion of SmsCB to an anaerobic protist from a methanomicrobiales donor, followed by a series of eukaryote-to-eukaryote LGT events amongst disparate lineages of anaerobic eukaryote lineages. The deeply branching position of the BRC clade amongst the anaerobic protists suggests it is possible that an ancestral BRC was the first recipient of these genes and an early offshoot from the BRC lineage passed the fusion gene to other anaerobic protists, although it is impossible to rule out alternative scenarios. Regardless, this Sms system is phylogenetically distinct, and has a separate origin from, the more complex SUF system found within the metamonad preaxostylans (i.e. oxymonads and *Paratrimastix*) (Vacek et al., 2018) and therefore represents an interesting case of convergence by LGT within the Metamonada. The resulting reduction and loss of the ISC system following acquisition of a Suf-related system is also similar between these two lineages and the breviate *Pygsuia biforma*. The gain of an alternate Fe/S cluster synthesis system in these cases likely 'preadapts' the ancestral organism to the loss of the ISC system in some of the lineages (e.g. an ancestor of GEMRC, RCL and RC). Unlike *P. biforma* and the preaxostylans, PCE and PAP020 retain a few core components of the early ISC system in addition to possessing the novel Sms system, suggesting that their MRO ISC system is at least partially functional. This retention of the ISC proteins may be related to the possession of Fe/S cluster-containing proteins

such as the NuoE and NuoF complex I proteins that are predicted to function in the MRO. We cannot determine the subcellular localization of the Sms system in BRCs bioinformatically because, like many metamonads, BRCs have cryptic (or absent) targeting peptides (see discussion below). However, we hypothesize the SmsCB is cytosolic in BRCs and functions somehow in coordination with the CIA system of these organisms to produce their Fe/S clusters.

Similar to *P. biforma*, the BRCs that completely lack any dedicated ISC components (members of the Retortacarp clade) do not seem to have any protein with identifiable cysteine desulfurase domains as assessed by HMM profile searches and phylogenetic analysis (see Supplementary Figure 3). This is surprising because for most Fe/S cluster synthesis systems, cysteine is the source of sulfur and a cysteine desulfurase is the key enzyme involved in mobilizing it. We suspect that these BRCs may directly utilize sulfide from the environment as the sulfur source for Fe/S clusters, as suggested for methanogenic archaea that possess a simple cysteine-desulfurase-lacking Sms pathway and which also live in sulfidic conditions (Liu et al., 2010). The exact mechanism by which such organisms construct Fe/S clusters is unknown and more work needs to be done to understand their biogenesis in these organisms.

The next highly conserved mitochondrion/MRO proteins that we investigated were the mitochondrial orthologs of the molecular chaperones chaperonin-60 (Cpn60) and hsp70. To distinguish cytosolic Chaperonin Containing TCP-1 (CCT) subunits from mitochondrial Cpn60 (GroEL) homologs and cytosolic and endoplasmic reticiulum (ER) paralogs of hsp70 from their mitochondrial hsp70 (i.e. dnaK orthologs), we used phylogenetic analyses. For RC and RCL, all Cpn60 candidates identified distinctly branched with CCT subunit clades (See Supplementary Figure 4); no bona fide mitochondrial Cpn60s could be identified from these species. GEMRC, PCE, and PAP020 appear to have both CCTs and mitochondrial Cpn60. Phylogenetic analysis of hsp70 proved more difficult, due in part to a greater degree of similarity between paralogous copies of the chaperone. To supplement this analysis, signature insertion-deletions in the multiple alignments were identified that were used in conjunction with well-annotated hsp70 yeast homologs (Nelson et al., 1992) to group sequences into cytosolic/ER versus mitochondrial type hsp70 groups (see Supplementary Figure 5). This information was then checked against the phylogeny for subcellular localization predictions of BRC hsp70 candidates. From this analysis, mitochondrial-type hsp70s were identified in all BRCs except RC. Within MROs, Cpn60 and hsp70 aid in protein translocation into the matrix and help newly translocated proteins adopt their native structure. These chaperones are ubiquitously predicted to function in a variety of MROs, so it is notable that they are missing in some of the BRCs data. For RC, we did not detect any genes encoding MRO chaperones, while RCL appears to contain hsp70 but lacks cpn60. This further highlights the highly reduced nature of the MROs found within these protists, especially within RC.

All mitochondria and MROs have membrane associated proteins, including the TIM/TOM complex proteins involved in protein translocation into the organelle. To search for these, we used HMM profiles as many of these sequences are highly divergent and not detectable by pairwise alignment-based search algorithms. The main channel forming protein in the TOM complex, Tom40, was only identifiable in RCL and GEMRC, while the receptor Tom70 was

identified in PCE, PAP020, and RCL. TIM subunits were generally absent except for Tim17 that was identified in PCE only. Except for RC, all BRC members appeared to have the co-chaperone Pam18 subunit of the presequence translocase-associated motor (PAM) complex in addition to the core mtHsp70 subunit. Pam16 was only identified in GEMRC and Pam17 appeared to be absent. Candidate mitochondrial processing peptidases (MPPs) that remove the MTS were found in RC, GEMRC, and RCL (Supplementary Figure 6). Our phylogenetic analysis shows that all MPP candidates in the BRCs are most closely related to a non-mitochondrial group of fungal metaloproteases, making them unlikely to be true MPP homologs. The divergent nature of MPPs is caused in part by the degeneration and evolution of mitochondrial targeting signals in metamonads and has caused marked differences in the evolution of MPPs found in parasitic lineages (Šmíd et al., 2008). A candidate mitochondrial carrier protein (MCP) was identified in RC; it's substrate specificity could not be determined by phylogenetic analysis (Supplementary Figure 7). Whether this protein functions in the MRO membrane is unclear as MCPs are also known to function in peroxisomes or other subcellular organelles (Mazurek et al., 2010). Finally, a candidate homolog Sam50 was identified in RC and GEMRC, though a phylogenetic analysis could not confirm the validity of this homolog. Orthologs of this candidate Sam50 were not detectable in any of the other BRC genomes, even when this sequence was included in the HMM profile used for searching. To determine if this Sam50 homolog folded into a structure resembling the mitochondrial orthologs, we used AlphaFold 2.0 (Jumper et al., 2021) to predict its structure. The resulting structures (Supplementary Figure 8) show that while these candidate Sam50 proteins possess what could be a intermembrane POTRA domain, the barrels are smaller and adopt a different confirmation compared to Sam50 homologs of yeast and other well characterized examples (Takeda et al., 2021).

GEMRC and RCL contain the most candidate MRO translocation proteins, which is similar to the repertoire found in *Giardia lamblia* (Leger et al., 2017) with the notable exception of the absence of GrpE in all members of BRC. PCE, PAP020 and RC possess a remarkably reduced MRO translocation complex by comparison. This may be due to the highly divergent nature of membrane bound MRO proteins, which makes detecting such proteins challenging (Pyrihová et al., 2018). Alternatively, in the case of RC which lacks all the highly conserved GCS, ISC and protein import/refolding components of MROs, it is unclear if either the candidate Sam50, the MCP and MPP are actually targeted to an organelle at all; it is possible that RC, like the distantly related *Monocercomonoides exilis*, might completely lack a mitochondrial compartment. Further works needs to be undertaken to determine the cellular localization of these key candidate MRO proteins.

MTS prediction is a known challenge in inferring MRO function in anaerobic protists, including metamonads (Jerlström-Hultqvist et al., 2013; Schneider et al., 2011; Tanifuji et al., 2018), due to highly divergent or missing MTSs. It has been suggested that the lack of a long positively charged MTS in metamonads is due to the loss of the proton gradient in the organelles (Garg et al., 2015), though recent analysis of canonical MTSs found in the Anaeramoebae – deep-branching metamonads – suggest that this may not always be the case (Stairs et al., 2021). Previous work has shown that some MRO proteins rely on cryptic internal signals to target proteins to the organelle with high fidelity (Garg et al., 2015; Mentel et al., 2008). Furthermore,

most available MTS prediction software tools were trained on experimentally validated datasets of targeted proteins that are lacking for most protists. For these reasons, making accurate MTS predictions in metamonads is exceedingly difficult. Here, we employed several different MTS prediction software tools to find and annotate these signals in BRCs (see Supplementary Table 1). We found several examples of false positives in RC and GEMRC, as homologs of transposons and nuclear targeting proteins were predicted to possess an MTS by some targeting prediction software with high confidence. Because of this, we considered proteins with MTS predictions from two or more MTS prediction software to be "strongly supported" targeting signals. Based on this criterion, RC lacked any strongly supported signals, while strongly supported signals were rare in all other BRC members. PCE contains strongly supported MTSs predicted on three proteins: an [FeFe]-hydrogenase, the T-protein of the glycine cleavage system (GCST), and on the IscU scaffold protein. PAP020 also contained a strongly supported MTS on its ortholog of IscU, as well as a different component of the glycine cleavage system (GCSP2). Another MTS detected was found on pre-sequence translocated-associated motor protein PAM18 in GEMRC. Interestingly, in RCL we find that a cytosolic CCT-gamma subunit contains a strongly predicted MTS, where in other BRCs, all cytosolic CCT subunits appear to be missing a MTS. However, when aligned to other BRC CCT-gamma subunits, it does not appear to have an obvious N-terminal extension. As RCL has no mitochondrial Cpn60 homologs, but does contain proteins which are homologous to MRO-localized proteins in other metamonads, it is possible that this CCT subunit has replaced the mitochondrial homolog in this species by acquiring an MTS. RCL does not have an additional copy of the gamma subunit of CCT. Follow up localization experiments are needed to deduce the subcellular localization of this particular CCT subunit.

The BRC share a core ATP generation pathway

All BRC members share a similar hypothetical ATP production pathway that is reminiscent of the pathway found in *Giardia* (Jedelský et al., 2011; Sanchez & Müller, 1996). In this pathway, pyruvate:ferredoxin oxidoreductase (PFO) oxidizes pyruvate and combines it with co-enzyme A to produce acetyl-CoA and a molecule of CO_2 while ferredoxin is reduced. Acetyl-CoA is then transformed to acetate by ACS and converting ADP to ATP. Ferredoxin is reoxidized by [FeFe]-hydrogenase (HydA), and the electrons are passed to protons creating hydrogen gas. Additionally, pyruvate-formate lyase was detected in all BRC members, allowing an alternative pathway of acetyl-CoA production from pyruvate (Stairs et al., 2011; see Supplementary Figure 9). Note that there are multiple copies of both HydA and PFO in each BRC member, and some of which possess additional domains encoding ferredoxin or flavodoxin-like proteins (see Supplementary Table 2). No components of typical ATP production pathways found in trichomonad hydrogenosomes, such as acetate:succinyl-CoA transferase or succinyl-CoA synthase (Lahti et al., 1992; Van Grinsven et al., 2008), were detected in any of the BRCs. Though [FeFe]-hydrogenase is present in each BRC member, none of the [FeFe]-hydrogenase maturase proteins (HydG, E or F) were detected. This is notable because these maturases, when present, are always targeted to mitochondria or MROs. This, combined with the lack of MTS detected in any of the proteins from the above pathway, leads us to assume that ATP and H_2 gas are likely produced in the cytosols of the Retortacarp lineages.

However, the pathways in the barthelonids appear to be slightly different in function and likely localization. Both PCE and PAP020 encode NuoE and NuoF NADH-dehydrogenase subunits of complex I that are known to function in the mitochondrial/MRO matrix in eukaryotes. It is widely thought that in anaerobic protists that possess them, NuoE and NuoF likely complex with an MRO-localized HydA to perform simultaneous NADH and ferredoxin (Fd⁻) oxidation by electron confurcating reaction that produces H_2 gas, NAD+ and Fd (Dyall et al., 2004; Stairs et al., 2015). For this reason, and the fact that one of HydA proteins of PCE and PAP020 have a predicted MTS, we infer that both barthelonids likely have MRO-associated hydrogenase activity, as recently suggested for PAP020 by Yazaki et al. (2019). However, unlike the hypothesis of Yazaki and colleagues (see Fig. 4b in Yazaki et al. 2019), we suggest that the reduced ferredoxin may, in fact, get produced within their MROs by PFO activity. If so, it is also possible that at least one of the two ACS homologs in each of these organisms is also MRO-localized and produces ATP within the organelle, though no MTSs are predicted on any of the ACS homologs. Further localization experiments in these barthelonids are needed to test these hypotheses.

Previous studies have shown that some of the enzymes involved in ATP synthesis found within metamonads have a complex phylogenetic history. In particular, ACS is thought to have been transferred into the metamonads multiple times through LGT (Leger et al., 2017). To determine the evolutionary histories of the BRC ACSs, we constructed a phylogenetic tree containing ACSs from a variety of metamonads, which can be found as Supplementary Figure 10. Our results show that, as in previous studies, most metamonad ACSs branch in one of two clades, known as ACS1 and ACS2. The ACS homologs found in the BRCs also form two distinct groups; one appears to be related to ACS2 found within *Carpediemonas membranifera*, *Ergobibamus cyprinoides,* and *Trimastix*. Additional ACS2 homologs from *Monocercomonoides* and *Paratrimastix* were also identified, which also branch with the metamonad ACS2 group. This suggests that ACS2 was inherited in the last common ancestor of Preaxostyla and the fornicates + BRC. The BRC also possess a second ACS homolog which branches separately from other metamonad ACSs. This enzyme seems to be most closely related to ACS found within *Streblomastix*, *Monocercomonoides* and *Blastocystis*, and is distinct from other types of ACS in the tree. As previously reported by Yazaki et al. (2019) in their analysis of the PAP020 transcriptome, this is likely to be a novel homolog of ACS (ACS3) and was probably transferred into metamonads through a LGT event distinct from the event that gave rise to ACS1 and ACS2. Our analysis also suggests that the ACS found within *Spironucleus salmonicida* MROs may not be ACS2, as previously reported (Leger et al., 2017) because, in our analyses, the *S. salmonicida* "ACS2" ortholog did not form a clade with other metamonad ACS2 sequences but instead grouped with several TACK archaea, although the support values for this relationship are low. A diagram summarizing the evolutionary relationships of these proteins, along with other key metabolic processes and proteins mentioned above, can be found as Figure 4. These relationships suggest that there have been at least three, and possibly four, independent transfers of ACS into the metamonads. These transfers occurred at very different points in time along the evolution of various metamonad groups and highlights the role of LGT in the evolution of anaerobic metabolism.

Based on our reconstructions of MRO properties on the metamonad tree, we suggest that the last common ancestor of BRC possessed an already highly reduced MRO compared to the common ancestor of fornicates + BRC (Figure 4). The barthelonids PCE and PAP020 seem to have functionally reduced MROs, but their predicted MRO proteomes retain similar functions to other free-living metamonads such as the GCS and serine metabolism, hydrogen production and Fe/S cluster biogenesis by a simple ISC system. On the other hand, the MROs of RCL and GEMRC appear to be unique amongst metamonads, as we predict that the organelles only function in glycine and serine metabolism and are unlikely to produce hydrogen. For RC, the only proteins typically associated with mitochondria and MROs detected were a candidate MCP and a homolog of Sam50; of these, only the latter is exclusively associated with mitochondrial organelles in eukaryotes. The complete lack of all other proteins homologous to common MRO-localizing proteins could indicate that an MRO may not even exist in this organism, a hypothesis that will be experimentally tested in the future via subcellular localization studies. In any case, this extreme reductive evolutionary path may relate to the acquisition of a simple fused SmsCB gene that was acquired by LGT in a common ancestor of BRC. This acquisition then led to the simplification of the ISC system in the PCE and PAP020 clade, and the outright loss of the ISC system in the Retortacarp clade, as Fe/S proteins for organellar functions were no longer needed.

Regardless of the evolutionary forces involved, the BRCs therefore demonstrate a reduction in MRO function that is independent of the loss seen in other metamonad groups, such as diplomonads and oxymonads (Karnkowska et al., 2016; Leger et al., 2017). This is remarkable given that many metamonads that display a similar loss of MRO functions are parasitic or commensalistic and the nutrient-rich endobiotic lifestyle has often been assumed to be part of the explanation for extreme MRO proteome streamlining (Stairs et al., 2015). Instead, in the case of the free-living BRC members, it appears that gain of a simple SMS system was amongst the first triggers leading to the extreme reductive MRO evolution in this group.

Conclusion

We have generated high quality draft genomes, transcriptomes, and curated predicted proteome sets for a novel group of free-living metamonads. The placement of BRC within the tree of Metamonada as the sister group to the Fornicata make them a key group to study in terms of genome evolution, LGT events, and adaptation to various environmental niches. These data place us several steps forward towards the goal of representing diverse and historically under-sampled microbial eukaryotes in the tree of eukaryotic life.

Our findings demonstrate the diversity of MRO configuration and function amongst the metamonads. As a group, BRCs display a level of MRO reduction not typically seen in free-living metamonads and which is independent of the reduction displayed in the parasitic diplomonads. BRC members share many interesting parallels with the MRO reduction and loss seen in other metamonads, as all BRCs possess a Sms Fe/S cluster system that has, in some cases, replaced the canonical ISC system. This is the second discovered instance of a Suf-like LGT event that has replaced the ISC system in metamonads, which serves to highlight the role of

adaptive LGTs in MRO evolution. Using these findings, we speculate that the last common ancestor of the BRCs possessed both a compact, streamlined genome with a highly reduced MRO, and was markedly different from the last common ancestor of the fornicates. Further work to elucidate the exact subcellular localization of several candidate MRO proteins, such as components of the ATP synthesis pathway and the Sam50-like protein in RC are needed to fully understand the breadth of function in these newly discovered organisms.

Methods

Cell cultivation & microscopy

Established cultures of isolates RC, RCL and GEMRC (Eglit et al. in prep) were maintained in variants of CR media (Gigeroff et al., 2023); RC in 50 mL of 40 ppt CR with 3% LB, GEMRC in 50 mL of 40 ppt CR with 1% LB and 3 sterile wheat grains, RCL in 50 mL of 25 ppt CR with 1% LB and 3 sterile wheat grains. All were inoculated into fresh media every 7-10 days based on culture density, with a second inoculation after 24 hours. Cultures of isolate 'Barthelona sp.' PCE (Yazaki et al. 2019) were maintained in sterile filtered seawater (Bedford Basin, Nova Scotia) supplemented with 3% LB, and kept at 16°C. Cell growth was monitored weekly, and subculturing was done every 3-4 weeks depending on cell density by transferring around 20% of the old culture into the new culture.

Cells were imaged with differential interference contrast optics on a Zeiss Axiovert 200M microscope fitted with a AxioCam M5 camera (Carl Zeiss AG).

Nucleic acid extraction & sequencing

For RC, GEMRC, RCL, and PCE - cell material for DNA and RNA extraction was harvested from 1-2 L of dense cell culture. The cells were collect by centrifugation in 50 mL tubes for 8 min at 2000xg and at 4°C. The pellets were then combined into two 15 mL tubes that were spun again as before. The recovered pellets were resuspended in sterile filtered (0.2 µm) spent culture media and layered over Histopaque, and then subjected to centrifugation at 2000xg for 20 min at room temperature. The protist-containing layer between the Histopaque and the top media was collected, and the procedure was repeated for a total of two rounds of Histopaque separation. Finally, the cells were diluted by 5 mL (10 volumes) of sterile spent-media and pelleted for 8 min, at 2000xg, 4°C. The cells were then resuspended in 2 mL of fresh media.

Genomic DNA was purified using a variety of techniques including traditional phenol-chloroform extraction, CTAB extraction, the QIAGEN Genomic-tip kit, and the QIAGEN MagAttract HMW DNA kit. RNA was extracted using Trizol according to Invitrogen specifications. RNA and DNA were sent to Genome Quebec for library construction and NovaSeq 6000 Illumina sequencing. In the case of the RNA, poly-A selection was performed to enrich for eukaryotic reads. High molecular weight DNA from each species was used to construct a 1D ligation Nanopore library (LSK108 for PCE, LSK109 - EXP-NBD104 for RC and GEMRC, and LSK308 for RCL) and was sequenced using a MinION flowcell (FLO-MIN106 (R9.4) for RC and GEMRC, and FLO-MIN107 (R9.5) for RCL and PCE).

For PAP020, cells were cultured as described in Yazaki et al. (2019). Cells were harvested by pelleting at 1500 rpm for 5 minutes at 4°C and were washed with PBS prior to phenol chloroform DNA extraction. The purified DNA was sequenced on a Illumina Hiseq 2500 PE using the Trueseq library prep.

Genome/transcriptome assembly & gene annotation

For RC, GEMRC, RCL, and PCE, Illumina NovaSeq reads were trimmed using Trimmomatic v0.36 (Bolger et al., 2014). RNASeq reads were assembled using Trinity v2.6.6 (Haas et al., 2013) and checked for multiplexing contamination using WinstonCleaner (https://github.com/kolecko007/WinstonCleaner). Bacterial contamination was removed from the Trinity assemblies using Anvi'o v5 (Eren et al., 2021).

Raw MinION sequencing data was basecalled using Albacore v2.1.3 (Albacore has since been replaced by Guppy – see Wick et al., 2019). These data were assembled using Flye v2.3 (Kolmogorov et al., 2019), Raven v0 (Vaser & Šikić, 2021) Canu v1.7 (Koren et al., 2017) and ABruijn v2.2b (Lin et al., 2016). The resulting assemblies were evaluated for completeness and contiguity by comparing assembly size, contig number, N50 and L50 values, and ALE scores (Clark et al., 2013). The ABruijn assembly was chosen moving forward for RC, RCL and PCE. For GEMRC, the Canu assembly was used. Raw MinION output was used to improve the base-call accuracy of these assemblies using Nanopolish version 0.8.4 (Loman et al., 2015) and Illumina reads were used to error-correct the assemblies using Unicycler version 0.4.3 (Wick et al., 2016). Bacterial contamination was assessed and removed from the resulting contigs using read coverage, GC content, and BLAST search results (Altschul et al., 1990).

Gene predictions were created by using RepeatMasker (Flynn et al., 2020), hisat2 (Kim et al., 2019), GeneMark-ET (Lomsadze et al., 2014), AUGUSTUS (Stanke et al., 2004), and PASA (Haas et al., 2003) as similarly described in Salas-Leiva et al. (2021). The predicted protein set was checked for completeness using the obd9 BUSCO v3.0.1 database (Simao et al., 2015).

To estimate genome completeness we used Merqury v.1.0 (Rhie et al., 2020) with a kmer size of 17 (estimated using Merqury). The Illumina reads used for the Merqury analysis were decontaminated using the Eukfinder workflow (Zhao, Salas-Leiva & Williams, in prep.) before genome completeness estimation. Additionally, RNA-seq mapping was used as a measure of genome completeness, and RNA-seq reads were decontaminated using the same method mentioned used for the DNA Illumina reads.

To construct the PAP020 genome, Illumina reads were assembled using Spades (Prjibelski et al., 2020) with default parameters. Bacterial reads were identified by metaBAT (Kang et al., 2015) and CheckM (Parks et al., 2015). Once the identified bacterial reads were removed, the remain Illumina reads were then used to assemble a new genome. This process was repeated until no bacterial contamination was detected. The final contigs were verified by checking kmer content and were also checked against the PAP020 transcriptome (Yazaki et al., 2019).

Phylogenomic analysis

The phylogenomic dataset was constructed based on a previously published set of 351 highly conserved protein orthologs (Brown et al., 2018), with 177 orthologs excluded due to deep paralogies or insufficient coverage across pertinent species in the analysis. Each of the 174 remaining ortholog alignments were used to construct single-protein phylogenies, and were used to screen out non-orthologous data. The final 174 alignments were realigned using mafft v7.310

einsi, trimmed using BMGE v1.0 (default settings) (Criscuolo & Gribaldo, 2010), and were concatenated together and used to infer a maximum-likelihood tree under the LG+C60+F+gamma model of evolution with IQTree v1.5.5 (Nguyen et al., 2015), using PMSF to calculate non-parametric bootstraps (Wang et al., 2018).

Inventory of anaerobic metabolism

Proteins of interest were searched for in the predicted proteome, transcriptome, and genome of each species using both BLAST and HMM profile searching using Hmmer v3.1b2 (hmmer.org). For each search, queries included both orthologous sequences from closely related metamonads and positively identified orthologs in other BRCs. Proteins relating to mitochondrial protein translocation were searched for using both general HMM profiles (Leger et al., 2017) and profiles tailored for searching in metamonads. The latter profiles were created by retrieving the profiles from Pfam for each protein of interest and then retrieving the best hits using the Pfam profile against the several publicly available metamonad sequencing data databases with hmmsearch. Using an evalue cutoff of <0.01, metamonad sequences were added to the corresponding Pfam seed alignment using the mafft –add function. If there were more than two detected sequences from metamonads, an alignment was made with these sequences and was then used to build a profile to search for more hits. Each candidate protein identified was used to reciprocally search the BLAST database to confirm protein identity. In ambiguous cases, the Pfam and InterPro databases were searched to analyze protein domain structure.

Phylogenies for key proteins identified were constructed to aid in identification in cases with deep paralogy (hsp70 and Cpn60). Phylogenies were also made to identify the origin of enzymes that act in anaerobic metabolism. If available, alignments from Leger et al. 2017 were utilized as a starting point, and we added the corresponding BRC sequences. Where relevant, the top 10 blast hit protein sequences from each BRC homolog were also added to the appropriate alignments. In cases where no starting alignments were available, a preliminary alignment was created by retrieving the top 1000 blast hits and reducing the sequences to 100-120 sequences using CD-HIT v.4.6 with recommended word sizes. These sequences were aligned using mafft einsi with default settings and were trimmed using trimal v1.4.rev15 with the -gappyout setting. These phylogenies were inferred using the LG+C20+F+gamma model of evolution with IQTree.

Mitochondria-related organelle protein localization predictions

Mitochondrial targeted proteins were predicted using the following mitochondrial targeting peptide prediction programs: TargetP v1.1b (Emanuelsson et al., 2007), MitoProt II v1.101 (Claros & Vincens, 1996), and MitoFates v1.1 (Fukasawa et al., 2015)). Additionally, the predicted proteins were used to search both the nr database and a custom database of known mitochondrial proteins.

SMSCB (SUFCB) phylogeny

The SmsCB fusion protein identified in each BRC member was added to previously constructed alignments (Leger et al., 2016) of SmsC and SmsBD. This dataset was aligned using mafft einsi with default settings and was trimmed by hand. The dataset was then concatenated together, and

the phylogeny was inferred using the same methods as the single protein phylogenies mentioned above.

Data availability

Genomes and gene annotations (where applicable) can be found under BioProject ID PRJNA949547.

Citations

Anwar, S., Dikhit, M. R., Singh, K. P., Kar, R. K., Zaidi, A., Sahoo, G. C., Roy, A. K., Nozaki, T., Das, P., & Ali, V. (2014). Interaction between Nbp35 and Cfd1 proteins of cytosolic Fe-S cluster assembly reveals a stable complex formation in Entamoeba histolytica. *PLoS ONE*, *9*(10), 12–14. https://doi.org/10.1371/journal.pone.0108971

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics*, *30*(15), 2114–2120. https://doi.org/10.1093/bioinformatics/btu170

Brown, M. W., Heiss, A. A., Kamikawa, R., Inagaki, Y., Yabuki, A., Tice, A. K., Shiratori, T., Ishida, K. I., Hashimoto, T., Simpson, A. G. B., & Roger, A. J. (2018). Phylogenomics Places Orphan Protistan Lineages in a Novel Eukaryotic Super-Group. *Genome Biology and Evolution*, *10*(2), 427–433. https://doi.org/10.1093/gbe/evy014

Criscuolo, A., & Gribaldo, S. (2010). BMGE (Block Mapping and Gathering with Entropy): A new software for selection of phylogenetic informative regions from multiple sequence alignments. *BMC Evolutionary Biology*, *10*(1). https://doi.org/10.1186/1471-2148-10-210

Derelle, E., Ferraz, C., Rombauts, S., Rouzé, P., Worden, A. Z., Robbens, S., Partensky, F., Degroeve, S., Echeynié, S., Cooke, R., Saeys, Y., Wuyts, J., Jabbari, K., Bowler, C., Panaud, O., Piégu, B., Ball, S. G., Ral, J. P., Bouget, F. Y., … Moreau, H. (2006). Genome analysis of the smallest free-living eukaryote Ostreococcus tauri unveils many unique features. *Proceedings of the National Academy of Sciences of the United States of America*, *103*(31), 11647–11652. https://doi.org/10.1073/pnas.0604795103

Dyall, S. D., Yan, W., Delgadillo-Correa, M. G., Lunceford, A., Loo, J. A., Clarke, C. F., & Johnson, P. J. (2004). Non-mitochondrial complex I proteins in a hydrogenosomal oxidoreductase complex. *Nature*, *431*(7012), 1103–1107. https://doi.org/10.1038/nature02990

Eren, A. M., Kiefl, E., Shaiber, A., Veseli, I., Miller, S. E., Schechter, M. S., Fink, I., Pan, J. N., Yousef, M., Fogarty, E. C., Trigodet, F., Watson, A. R., Esen, Ö. C., Moore, R. M., Clayssen, Q., Lee, M. D., Kivenson, V., Graham, E. D., Merrill, B. D., … Willis, A. D. (2021). Community-led, integrated, reproducible multi-omics with anvi'o. *Nature Microbiology*, *6*(1), 3–6. https://doi.org/10.1038/s41564-020-00834-3

Flynn, J. M., Hubley, R., Goubert, C., Rosen, J., Clark, A. G., Feschotte, C., & Smit, A. F. (2020). RepeatModeler2 for automated genomic discovery of transposable element families. *Proceedings of the National Academy of Sciences of the United States of America*, *117*(17), 9451–9457. https://doi.org/10.1073/pnas.1921046117

Garg, S., Stölting, J., Zimorski, V., Rada, P., Tachezy, J., Martin, W. F., & Gould, S. B. (2015). Conservation of transit peptide-Independent protein import into the mitochondrial and hydrogenosomal matrix. *Genome Biology and Evolution*, *7*(9), 2716–2726. https://doi.org/10.1093/gbe/evv175

Gawryluk, R. M. R., & Stairs, C. W. (2021). Diversity of electron transport chains in anaerobic protists. *Biochimica et Biophysica Acta - Bioenergetics*, *1862*(1), 148334. https://doi.org/10.1016/j.bbabio.2020.148334

Gigeroff, A. S., Eglit, Y., & Simpson, A. G. B. (2023). Characterisation and Cultivation of New Lineages of Colponemids, a Critical Assemblage for Inferring Alveolate Evolution. *Protist*, *174*(2), 125949. https://doi.org/10.1016/j.protis.2023.125949

Goffeau, A. A., Barrell, B. G., Bussey, H., Davis, R. W., Dujon, B., Feldmann, H., Hoheisel, J. D., Jacq, C., Johnston, M., Louis, E. J., Mewes, H. W., Murakami, Y., Philippsen, P., Tettelin, H., & Oliver, S. G. (1996). *Life with 6000 Genes*. *274*(5287).

Haas, B. J., Delcher, A. L., Mount, S. M., Wortman, J. R., Smith, R. K., Hannick, L. I., Maiti, R., Ronning, C. M., Rusch, D. B., Town, C. D., Salzberg, S. L., & White, O. (2003). Improving the Arabidopsis genome annotation using maximal transcript alignment assemblies. *Nucleic Acids Research*, *31*(19), 5654–5666. https://doi.org/10.1093/nar/gkg770

Haas, B. J., Papanicolaou, A., Yassour, M., Grabherr, M., Philip, D., Bowden, J., Couger, M. B., Eccles, D., Li, B., Macmanes, M. D., Ott, M., Orvis, J., Pochet, N., Strozzi, F., Weeks, N., Westerman, R., William, T., Dewey, C. N., Henschel, R., … Regev, A. (2013). De novo transcript sequence recostruction from RNA-Seq: reference generation and analysis with Trinity. In *Nature protocols* (Vol. 8, Issue 8). https://doi.org/10.1038/nprot.2013.084.De

Jedelský, P. L., Doležal, P., Rada, P., Pyrih, J., Šmíd, O., Hrdý, I., Šedinová, M., Marcinčiková, M., Voleman, L., Perry, A. J., Beltrán, N. C., Lithgow, T., & Tachezy, J. (2011). The minimal proteome in the reduced mitochondrion of the parasitic protist Giardia intestinalis. *PLoS ONE*, *6*(2), 15–21. https://doi.org/10.1371/journal.pone.0017285

Jerlström-Hultqvist, J., Einarsson, E., Xu, F., Hjort, K., Ek, B., Steinhauf, D., Hultenby, K., Bergquist, J., Andersson, J. O., & Svärd, S. G. (2013). Hydrogenosomes in the diplomonad Spironucleus salmonicida. *Nature Communications*, *4*, 1–9. https://doi.org/10.1038/ncomms3493

Jiménez-González, A., Xu, F., & Andersson, J. O. (2019). Lateral Acquisitions Repeatedly Remodel the Oxygen Detoxification Pathway in Diplomonads and Relatives. *Genome Biology and Evolution*, *11*(9), 2542–2556. https://doi.org/10.1093/gbe/evz188

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S. A. A., Ballard, A. J., Cowie, A., Romera-Paredes, B., Nikolov, S., Jain, R., Adler, J., … Hassabis, D. (2021). Highly accurate protein structure prediction with AlphaFold. *Nature*, *596*(7873), 583–589. https://doi.org/10.1038/s41586-021-03819-2

Kang, D. D., Froula, J., Egan, R., & Wang, Z. (2015). MetaBAT, an efficient tool for accurately reconstructing single genomes from complex microbial communities. *PeerJ*, *2015*(8), 1–15. https://doi.org/10.7717/peerj.1165

Kang, S., Tice, A. K., Spiegel, F. W., Silberman, J. D., Pánek, T., Cepicka, I., Kostka, M., Kosakyan, A., Alcântara, D. M. C., Roger, A. J., Shadwick, L. L., Smirnov, A., Kudryavtsev, A., Lahr, D. J. G., & Brown, M. W. (2017). Between a Pod and a Hard Test: The Deep Evolution of Amoebae. *Molecular Biology and Evolution*, *34*(9), 2258–2270. https://doi.org/10.1093/molbev/msx162

Karnkowska, A., Vacek, V., Zubáčová, Z., Treitli, S. C., Petrželková, R., Eme, L., Novák, L., Žárský, V., Barlow, L. D., Herman, E. K., Soukal, P., Hroudová, M., Doležal, P., Stairs, C.

W., Roger, A. J., Eliáš, M., Dacks, J. B., Vlček, Č., & Hampl, V. (2016). A eukaryote without a mitochondrial organelle. *Current Biology*, *26*(1), 1274–1284. https://doi.org/10.1016/j.cub.2016.03.053

Kikuchi, G. (1973). *THE GLYCINE CLEAVAGE SYSTEM: COMPOSITION, REACTION MECHANISM, AND PHYSIOLOGICAL SIGNIFICANCE*. *21*(July), 1154–1157.

Kim, D., Paggi, J. M., Park, C., Bennett, C., & Salzberg, S. L. (2019). Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology*, *37*(August). https://doi.org/10.1038/s41587-019-0201-4

Kolmogorov, M., Yuan, J., Lin, Y., & Pevzner, P. A. (2019). Assembly of long, error-prone reads using repeat graphs. *Nature Biotechnology*, *37*(5), 540–546. https://doi.org/10.1038/s41587-019-0072-8

Lahti, C. J., D'Oliveira, C. E., & Johnson, P. J. (1992). β-Succinyl-coenzyme A synthetase from Trichomonas vaginalis is a soluble hydrogenosomal protein with an amino-terminal sequence that resembles mitochondrial presequences. *Journal of Bacteriology*, *174*(21), 6822–6830. https://doi.org/10.1128/jb.174.21.6822-6830.1992

Leger, M. M., Eme, L., Hug, L. A., & Roger, A. J. (2016). Novel Hydrogenosomes in the Microaerophilic Jakobid Stygiella incarcerata. *Molecular Biology and Evolution*, *33*(9), 2318–2336. https://doi.org/10.1093/molbev/msw103

Leger, M. M., Kolisko, M., Kamikawa, R., Stairs, C. W., Kume, K., Čepička, I., Silberman, J. D., Andersson, J. O., Xu, F., Yabuki, A., Eme, L., Zhang, Q., Takishita, K., Inagaki, Y., Simpson, A. G. B., Hashimoto, T., & Roger, A. J. (2017). Organelles that illuminate the origins of Trichomonas hydrogenosomes and Giardia mitosomes. *Nature Ecology and Evolution*, *1*(4), 1–15. https://doi.org/10.1038/s41559-017-0092

Lill, R., & Freibert, S. A. (2020). Mechanisms of Mitochondrial Iron-Sulfur Protein Biogenesis. *Annual Review of Biochemistry*, *89*, 471–499. https://doi.org/10.1146/annurev-biochem-013118-111540

Liu, Y., Sieprawska-Lupa, M., Whitman, W. B., & White, R. H. (2010). Cysteine is not the sulfur source for iron-sulfur cluster and methionine biosynthesis in the methanogenic archaeon Methanococcus maripaludis. *Journal of Biological Chemistry*, *285*(42), 31923–31929. https://doi.org/10.1074/jbc.M110.152447

Lomsadze, A., Burns, P. D., & Borodovsky, M. (2014). *Integration of mapped RNA-Seq reads into automatic training of eukaryotic gene finding algorithm*. *42*(15), 1–8. https://doi.org/10.1093/nar/gku557

Manzano-Marín, A., & Latorre, A. (2016). Snapshots of a shrinking partner: Genome reduction in Serratia symbiotica. *Scientific Reports*, *6*(June), 1–11. https://doi.org/10.1038/srep32590

Mazurek, M. P., Prasad, P. D., Gopal, E., Fraser, S. P., Bolt, L., Rizaner, N., Palmer, C. P., Foster, C. S., Palmieri, F., Ganapathy, V., Stühmer, W., Djamgoz, M. B. A., & Mycielska, M. E. (2010). Molecular origin of plasma membrane citrate transporter in human prostate epithelial cells. *EMBO Reports*, *11*(6), 431–437. https://doi.org/10.1038/embor.2010.51

Mentel, M., Zimorski, V., Haferkamp, P., Martin, W., & Henze, K. (2008). Protein import into

hydrogenosomes of Trichomonas vaginalis involves both N-terminal and internal targeting signals: A case study of thioredoxin reductases. *Eukaryotic Cell*, *7*(10), 1750–1757. https://doi.org/10.1128/EC.00206-08

Motyčková, A., Voleman, L., & Najdrová, V. (2022). The late ISC pathway interactome reveals mitosomal-cytoplasmic crosstalk in Giardia intestinalis. 2005–2003, *8.5.2017*, הארץ.

Mukherjee, M., Brown, M. T., McArthur, A. G., & Johnson, P. J. (2006). Proteins of the glycine decarboxylase complex in the hydrogenosome of Trichomonas vaginalis. *Eukaryotic Cell*, *5*(12), 2062–2071. https://doi.org/10.1128/EC.00205-06

Müller, M., Mentel, M., van Hellemond, J. J., Henze, K., Woehle, C., Gould, S. B., Yu, R.-Y., van der Giezen, M., Tielens, A. G. M., & Martin, W. F. (2012). Biochemistry and Evolution of Anaerobic Energy Metabolism in Eukaryotes. *Microbiology and Molecular Biology Reviews*, *76*(2), 444–495. https://doi.org/10.1128/mmbr.05024-11

Nelson, R. J., Ziegelhoffer, T., Nicolet, C., Werner-Washburne, M., & Craig, E. A. (1992). The translation machinery and 70 kd heat shock protein cooperate in protein synthesis. *Cell*, *71*(1), 97–105. https://doi.org/10.1016/0092-8674(92)90269-I

Nývltová, E., Smutná, T., Tachezy, J., & Hrdý, I. (2016). OsmC and incomplete glycine decarboxylase complex mediate reductive detoxification of peroxides in hydrogenosomes of Trichomonas vaginalis. *Molecular and Biochemical Parasitology*, *206*(1–2), 29–38. https://doi.org/10.1016/j.molbiopara.2016.01.006

Nývltová, E., Stairs, C. W., Hrdý, I., Rídl, J., Mach, J., Paۻes, J., Roger, A. J., & Tachezy, J. (2015). Lateral gene transfer and gene duplication played a key role in the evolution of mastigamoeba balamuthi hydrogenosomes. *Molecular Biology and Evolution*, *32*(4), 1039–1055. https://doi.org/10.1093/molbev/msu408

Ohnishi, T. (1998). Iron-sulfur clusters/semiquinones in Complex I. *Biochimica et Biophysica Acta - Bioenergetics*, *1364*(2), 186–206. https://doi.org/10.1016/S0005-2728(98)00027-9

Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P., & Tyson, G. W. (2015). CheckM: Assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Research*, *25*(7), 1043–1055. https://doi.org/10.1101/gr.186072.114

Prjibelski, A., Antipov, D., Meleshko, D., Lapidus, A., & Korobeynikov, A. (2020). Using SPAdes De Novo Assembler. *Current Protocols in Bioinformatics*, *70*(1), 1–29. https://doi.org/10.1002/cpbi.102

Pyrih, J., Pyrihová, E., Kolísko, M., Stojanovová, D., Basu, S., Harant, K., Haindrich, A. C., Doležal, P., Lukeš, J., Roger, A., & Tachezy, J. (2016). Minimal cytosolic iron-sulfur cluster assembly machinery of Giardia intestinalis is partially associated with mitosomes. *Molecular Microbiology*, *102*(4), 701–714. https://doi.org/10.1111/mmi.13487

Pyrihová, E., Motyčková, A., Voleman, L., Wandyszewska, N., Fišer, R., Seydlová, G., Roger, A., Kolísko, M., & Doležal, P. (2018). A single tim translocase in themitosomes of giardia intestinalis illustrates convergence of protein import machines in anaerobic eukaryotes. *Genome Biology and Evolution*, *10*(10), 2813–2822. https://doi.org/10.1093/gbe/evy215

Rhie, A., Walenz, B. P., Koren, S., & Phillippy, A. M. (2020). Merqury: Reference-free quality,

completeness, and phasing assessment for genome assemblies. *Genome Biology*, *21*(1), 1–27. https://doi.org/10.1186/s13059-020-02134-9

Roger, A. J., Muñoz-Gómez, S. A., & Kamikawa, R. (2017). The Origin and Diversification of Mitochondria. In *Current Biology*. https://doi.org/10.1016/j.cub.2017.09.015

Salas-leiva, D. E., Tromer, E. C., Curtis, B. A., Jerlström-hultqvist, J., Kolisko, M., Yi, Z., Salas-leiva, J. S., Gallot-lavallée, L., Williams, S. K., Kops, G. J. P. L., Archibald, J. M., Simpson, A. G. B., & Roger, A. J. (2021). Genomic analysis finds no evidence of canonical eukaryotic DNA processing complexes in a free-living protist. *Nature Communications*, 1–13. https://doi.org/10.1038/s41467-021-26077-2

Sánchez, L. B., Galperin, M. Y., & Müller, M. (2000). Acetyl-CoA synthetase from the amitochondriate eukaryote Giardia lamblia belongs to the newly recognized superfamily of acyl-CoA synthetases (nucleoside diphosphate-forming). *Journal of Biological Chemistry*, *275*(8), 5794–5803. https://doi.org/10.1074/jbc.275.8.5794

Sanchez, L. B., & Müller, M. (1996). Purification and characterization of the acetate forming enzyme, acetyl-CoA synthetase (ADP-forming) from the amitochondriate protist, Giardia lamblia. *FEBS Letters*, *378*(3), 240–244. https://doi.org/10.1016/0014-5793(95)01463-2

Schneider, R. E., Brown, M. T., Shiflett, A. M., Dyall, S. D., Hayes, R. D., Xie, Y., Loo, J. A., & Johnson, P. J. (2011). The Trichomonas vaginalis hydrogenosome proteome is highly reduced relative to mitochondria, yet complex compared with mitosomes. *International Journal for Parasitology*, *41*(13–14), 1421–1434. https://doi.org/10.1016/j.ijpara.2011.10.001

Schönknecht, G., Chen, W. H., Ternes, C. M., Barbier, G. G., Shrestha, R. P., Stanke, M., Bräutigam, A., Baker, B. J., Banfield, J. F., Garavito, R. M., Carr, K., Wilkerson, C., Rensing, S. A., Gagneul, D., Dickenson, N. E., Oesterhelt, C., Lercher, M. J., & Weber, A. P. M. (2013). Gene transfer from bacteria and archaea facilitated evolution of an extremophilic eukaryote. *Science*, *339*(6124), 1207–1210. https://doi.org/10.1126/science.1231707

Simao, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V, & Zdobnov, E. M. (2015). *Genome analysis BUSCO : assessing genome assembly and annotation completeness with single-copy orthologs*. *31*(June), 3210–3212. https://doi.org/10.1093/bioinformatics/btv351

Simpson, A. G. B. (2003). Cytoskeletal organization, phylogenetic affinities and systematics in the contentious taxon Excavata (Eukaryota). *International Journal of Systematic and Evolutionary Microbiology*, *53*(6), 1759–1777. https://doi.org/10.1099/ijs.0.02578-0

Šmíd, O., Matušková, A., Harris, S. R., Kučera, T., Novotný, M., Horváthová, L., Hrdý, I., Kutějová, E., Hirt, R. P., Embley, T. M., Janata, J., & Tachezy, J. (2008). Reductive evolution of the mitochondrial processing peptidases of the unicellular parasites Trichomonas vaginalis and Giardia intestinalis. *PLoS Pathogens*, *4*(12). https://doi.org/10.1371/journal.ppat.1000243

Stairs, C. W., Eme, L., Brown, M. W., Mutsaers, C., Susko, E., Dellaire, G., Soanes, D. M., Van Der Giezen, M., & Roger, A. J. (2014). A SUF Fe-S cluster biogenesis system in the mitochondrion-related organelles of the anaerobic protist Pygsuia. *Current Biology*, *24*(11),

1176–1186. https://doi.org/10.1016/j.cub.2014.04.033

Stairs, C. W., Eme, L., Muñoz-Gómez, S. A., Cohen, A., Dellaire, G., Shepherd, J. N., Fawcett, J. P., & Roger, A. J. (2018). Microbial eukaryotes have adapted to hypoxia by horizontal acquisitions of a gene involved in rhodoquinone biosynthesis. *ELife*, *7*, 1–23. https://doi.org/10.7554/eLife.34292

Stairs, C. W., Leger, M. M., & Roger, A. J. (2015). Diversity and origins of anaerobic metabolism in mitochondria and related organelles. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *370*(1678). https://doi.org/10.1098/rstb.2014.0326

Stairs, C. W., Roger, A. J., & Hampl, V. (2011). Eukaryotic pyruvate formate lyase and its activating enzyme were acquired laterally from a firmicute. *Molecular Biology and Evolution*, *28*(7), 2087–2099. https://doi.org/10.1093/molbev/msr032

Stairs, C. W., Táborský, P., Salomaki, E. D., Kolisko, M., Pánek, T., Eme, L., Hradilová, M., Vlček, Č., Jerlström-Hultqvist, J., Roger, A. J., & Čepička, I. (2021). Anaeramoebae are a divergent lineage of eukaryotes that shed light on the transition from anaerobic mitochondria to hydrogenosomes. *Current Biology*, *31*(24), 5605-5612.e5. https://doi.org/10.1016/j.cub.2021.10.010

Stanke, M., Steinkamp, R., Waack, S., & Morgenstern, B. (2004). AUGUSTUS: A web server for gene finding in eukaryotes. *Nucleic Acids Research*, *32*(WEB SERVER ISS.), 309–312. https://doi.org/10.1093/nar/gkh379

Takeda, H., Tsutsumi, A., Nishizawa, T., Lindau, C., Busto, J. V., Wenz, L. S., Ellenrieder, L., Imai, K., Straub, S. P., Mossmann, W., Qiu, J., Yamamori, Y., Tomii, K., Suzuki, J., Murata, T., Ogasawara, S., Nureki, O., Becker, T., Pfanner, N., … Endo, T. (2021). Mitochondrial sorting and assembly machinery operates by β-barrel switching. In *Nature* (Vol. 590, Issue 7844). Springer US. https://doi.org/10.1038/s41586-020-03113-7

Tanifuji, G., Takabayashi, S., Kume, K., Takagi, M., Nakayama, T., Kamikawa, R., Inagaki, Y., & Hashimoto, T. (2018). The draft genome of Kipferlia bialata reveals reductive genome evolution in fornicate parasites. *PLoS ONE*, *13*(3), 1–18. https://doi.org/10.1371/journal.pone.0194487

Tsaousis, A. D., De Choudens, S. O., Gentekaki, E., Long, S., Gaston, D., Stechmann, A., Vinella, D., Py, B., Fontecave, M., Barras, F., Lukeš, J., & Roger, A. J. (2012). Evolution of Fe/S cluster biogenesis in the anaerobic parasite Blastocystis. *Proceedings of the National Academy of Sciences of the United States of America*, *109*(26), 10426–10431. https://doi.org/10.1073/pnas.1116067109

Vacek, V., Novak, L. V. F., Treitli, S. C., Taborsky, P., Cepicka, I., Kolısko, M., Keeling, P. J., & Hampl, V. (2018). Fe-S cluster assembly in oxymonads and related protists. *Molecular Biology and Evolution*, *35*(11), 2712–2718. https://doi.org/10.1093/molbev/msy168

Van Grinsven, K. W. A., Rosnowsky, S., Van Weelden, S. W. H., Pütz, S., Van Der Giezen, M., Martin, W., Van Hellemond, J. J., Tielens, A. G. M., & Henze, K. (2008). Acetate:succinate CoA-transferase in the hydrogenosomes of Trichomonas vaginalis: Identification and characterization. *Journal of Biological Chemistry*, *283*(3), 1411–1418. https://doi.org/10.1074/jbc.M702528200

Vaser, R., & Šikić, M. (2021). Time- and memory-efficient genome assembly with Raven. *Nature Computational Science*, *1*(5), 332–336. https://doi.org/10.1038/s43588-021-00073-4

Wick, R. R., Judd, L. M., & Holt, K. E. (2019). Performance of neural network basecalling tools for Oxford Nanopore sequencing. *Genome Biology*, *20*(1), 1–10. https://doi.org/10.1186/s13059-019-1727-y

Xu, F., Jex, A., & Svärd, S. G. (2020). A chromosome-scale reference genome for Giardia intestinalis WB. *Scientific Data*, *7*(1), 1–8. https://doi.org/10.1038/s41597-020-0377-y

Yazaki, E., Kume, K., Shiratori, T., Eglit, Y., Tanifuji, G., Harada, R., Simpson, A., Ishida, K., Hashimoto, T., & Inagaki, Y. (2019). *Barthelonids represent a deep-branching Metamonad clade with mitochondrion-related organelles generating no ATP.* https://doi.org/10.1101/805762

Yubuki, N., Galindo, L. J., Reboul, G., López-García, P., Brown, M. W., Pollet, N., & Moreira, D. (2020). Ancient Adaptive Lateral Gene Transfers in the Symbiotic Opalina-Blastocystis Stramenopile Lineage. *Molecular Biology and Evolution*, *37*(3), 651–659. https://doi.org/10.1093/molbev/msz250

Záhonová, K., Treitli, S. C., Le, T., Škodová-Sveráková, I., Hanousková, P., Čepička, I., Tachezy, J., & Hampl, V. (2022). Anaerobic derivates of mitochondria and peroxisomes in the free-living amoeba Pelomyxa schiedti revealed by single-cell genomics. *BMC Biology*, *20*(1), 1–16. https://doi.org/10.1186/s12915-022-01247-w

Žárský, V., Klimeš, V., Pačes, J., Vlček, Č., Hradilová, M., Beneš, V., Nývltová, E., Hrdý, I., Pyrih, J., Mach, J., Barlow, L., Stairs, C. W., Eme, L., Hall, N., Eliáš, M., Dacks, J. B., Roger, A., & Tachezy, J. (2021). The Mastigamoeba balamuthi Genome and the Nature of the Free-Living Ancestor of Entamoeba. *Molecular Biology and Evolution*, *38*(6), 2240–2259. https://doi.org/10.1093/molbev/msab020

Zítek, J., Füssy, Z., Treitli, S. C., Peña-Diaz, P., Vaitová, Z., Zavadska, D., Harant, K., & Hampl, V. (2022). Reduced mitochondria provide an essential function for the cytosolic methionine cycle. *Current Biology*, 5057–5068. https://doi.org/10.1016/j.cub.2022.10.028

Zubáčová, Z., Novák, L., Bublíková, J., Vacek, V., Fousek, J., Rídl, J., Tachezy, J., Doležal, P., Vlček, Č., & Hampl, V. (2013). The Mitochondrion-Like Organelle of Trimastix pyriformis Contains the Complete Glycine Cleavage System. *PLoS ONE*, *8*(3), 1–9. https://doi.org/10.1371/journal.pone.0055417

Figure 1. 'BRC' is a clade of anaerobic protists that branches as sister to all known Fornicata within the Metamonada. Panels A-D) Differential interference contrast light micrographs of Retortacarp lineages RC (A), GEMRC (B), and RCL (C), as well as *Barthelona* sp. PCE (D) showing overall morphology. Scale bars indicate a length of 10 µm.

E. A maximum likelihood (ML) phylogeny based on 174 concatenated aligned proteins encompassing 46,113 sites. The depicted topology was estimated using IQ-TREE under the LG+C60+F+Γ model and was used as a guide tree for estimating the LG+PMSF(C60)+F+Γ model. Support under the latter model was evaluated by SH-aLRT support percentage, aBayes support and nonparametric bootstrap percentage based on 100 nonparametric bootstrap replicates. The scale bar indicates the expected number of amino acid substitutions per site under the PMSF model. When at least one support value was less than maximal, all three are shown on the branch in the order SH-aLRT support percentage/aBayes/nonparametric bootstrap percentage. A black dot on the branch indicates all three approaches yielded full (i.e. 100/1/100) support. The BRC clade is highlighted in red.

Figure 2. Presence, absence and predicted localization of key MRO and energy metabolism enzymes in 'BRCs' and other metamonads. A schematic phylogeny of the Metamonada is shown on the left based on Fig. 1, with the BRC clade highlighted in red. The pathways depicted include anaerobic extended glycolysis/energy metabolism enzymes, conserved MRO proteins, MRO protein targeting machinery components, and the SmsCB fusion protein. Coloured versus grey circles indicate presence/absence of the protein from predicted proteomes and/or transcriptome data and the colours indicate probable localization. Circles split in half indicate that more than one paralog was present with different predicted localizations. Protein abbreviations: PFO - Pyruvate:ferredoxin oxidoreductase; NuoE/F - Respiratory-chain NADH dehydrogenase 24/51 kDa subunit; HydA – Iron-only hydrogenase; ASCT1B – Acetate:succinyl-CoA transferase B; ASCT1C - Acetate:succinyl-CoA transferase C; SCS - Succinyl-CoA synthase; ACS - Acetyl-CoA synthase; Fdx – Ferredoxin; L – Glycine cleavage system subunit L; H - Glycine cleavage system subunit H; P - Glycine cleavage system subunit P; T - Glycine cleavage system subunit T; SHMT - Serine hydroxymethyltransferaseTom40 - Translocase of outer mitochondrial membrane 40; Tom22 - Translocase of outer mitochondrial membrane 22; Sam50 – Sorting and assembly machinery 50; Tim17/22/23 - Translocase of the inner membrane 17/22/23; Tim44 - Translocase of the inner membrane 44; Pam16 - Presequence translocase-associated motor 16; Pam18 - Presequence translocase-associated motor 18; IscS - Iron-sulfur cluster assembly enzyme cysteine desulfurase; IscU - Iron-sulfur cluster assembly enzyme scaffold; SufCB - Sulfur formation CB fusion protein.

Figure 3. **A phylogeny of the SmsCB fusion protein in a variety of anaerobic eukaryotes (red) and their closest bacterial and archaeal homologs.** The alignment contains the SmsCB fusion protein from the eukaryotes and concatenated SmsC and SmsB proteins from the prokaryotes. This ML tree was estimated under the LG+C60+F+Γ model using IQ-TREE. Ultrafast bootstrap values are displayed on the branches, with dots indicating UFBOOT value over 95%.

Figure 4. **Gain and loss of anaerobic enzymes and MRO proteins over the tree of Metamonada**. Gains (red) and losses (green) of key metabolic pathway components discussed throughout this paper are mapped on a schematic phylogeny of Metamonada (based on Figure 1). Protein abbreviations are: HydG/E/F – Iron hydrogenase maturase G/E/F; ACS - Acetyl-CoA synthase; NuoE/F - Respiratory-chain NADH dehydrogenase 24/51 kDa subunit; GCS – Glycine cleavage system; SCS - Succinyl-CoA synthase; IscS/U - Iron-sulfur cluster assembly enzyme system.

Table 1. BUSCO scores for members of the BRC clade.

	Complete Single Copy	Complete Duplicated	Total Complete	Fragmented	Missing
RC - Genome	182	4	186	15	106
RC - Prediction	215	9	224	7	72
RC - Transcriptome	190	24	214	16	73
RC – Curated Predictions	208	17	225	7	71
GEMRC - Genome	168	9	177	27	99
GEMRC - Prediction	183	14	197	29	77
GEMRC - Transcriptome	139	77	216	9	78
RCL - Genome	182	4	186	16	101
RCL - Prediction	212	10	222	14	67
RCL - Transcriptome	193	27	220	14	69
PCE - Genome	142	3	145	15	143
PCE – Prediction	161	7	168	25	110
PCE - Transcriptome	5	157	162	31	110

Table 2. Summary statistics for the genome, transcriptome, and predicted proteomes of the BRC clade.

	Retortacarp (RC)	GEM-Retortacarp (GEMRC)	Retortacarp-like (RCL)	PEI Cavendish Eelgrass (PCE)	PAP20
*Genome Coverage (Nanopore)	41x	119x	34x	66x	Illumina only
*Genome Coverage (Illumina)	206x	116x	93x	308x	29x
# of Illumina Reads Generated	53.7 M	26.8 M	85.9 M	75.4 M	70.0 M
% RNAseq reads mapping	99.4	99.9	99.6	99.3	-
Merqury Completeness (%)	98.7	98.6	98.4	98.4	-
Genome Size	31.9 Mbp	33.5 Mbp	21.1 Mbp	10.2 Mbp	13.2 Mbp
Contigs (#)	16	21	18	8	1,035
N50	3,724,615	2,551,763	2,069,101	1,587,787	24,218
L50	4	6	4	3	156
%GC	35.8	36.9	36.0	40.3	30.5
%Repetitive	55.4	48.0	28.1	4.46	-
Predicted Proteins (#)	15,224	14,268	9,198	5,064	6,305

*Median contig coverage