Extreme mitochondrial reduction in a novel group of free-living metamonads.

Shelby K. Williams[1,2], Jon Jerlström Hultqvist[1,3], Yana Eglit[1,4], Dayana E. Salas-Leiva[1,5], Bruce
Curtis[1,2], Russell J. S. Orr[6], Courtney W. Stairs[7], Tuğba N. Atalay[1,2], Naomi MacMillan[1,2],
Alastair G. B. Simpson[1,4] and Andrew J. Roger[1,2*]

1 - Institute for Comparative Genomics, Dalhousie University, Nova Scotia, Canada
2 – Department of Biochemistry and Molecular Biology, Dalhousie University
3 – Department of Cell and Molecular Biology, Uppsala University, Sweden
4 - Department of Biology, Dalhousie University
5 – Department of Biochemistry, Cambridge University, UK
6 - Section for Genetics and Evolutionary Biology, Department of Biosciences, University of
Oslo, Norway
7 – Department of Biology, Lund University, Sweden
14
*Corresponding author: andrew.roger@dal.ca

<u>**Summary**</u>

Metamonads are a diverse group of heterotrophic microbial eukaryotes adapted to living in
hypoxic environments. All metamonads but one harbour metabolically altered 'mitochondrion-
related organelles' (MROs) with reduced functions relative to aerobic mitochondria, however the
degree of reduction varies markedly over the metamonad tree. To further investigate metamonad
MRO diversity, we generated high quality draft genomes, transcriptomes, and predicted
proteomes for five recently discovered free-living metamonads. Phylogenomic analyses placed
these organisms in a group we informally named the 'BaSk' (Barthelonids+Skoliomonads) clade,
which emerges as a deeply branching sister group to the Fornicata, a metamonad phylum that
includes parasitic and free-living flagellates. Extensive bioinformatic analyses of the manually
curated gene models showed that these organisms are predicted to have extremely reduced MRO
proteomes in comparison to other free-living metamonads. Loss of the mitochondrial iron-sulfur
cluster (ISC) assembly system in some organisms in this group appears to be linked to the
acquisition in their common ancestral lineage of a SUF-like minimal system (SMS) Fe/S cluster
pathway through lateral gene transfer (LGT). One of the isolates, *Skoliomonas litria*, appears to
have undergone further mitochondrial reduction having lost all other known MRO pathways. No
proteins were confidently assigned to the predicted MRO proteome of this organism suggesting
that the organelle has been lost. The extreme mitochondrial reduction observed within this free-
living anaerobic protistan clade is unprecedented and demonstrates that mitochondrial functions,
under some conditions, may be completely lost even in free-living organisms.

Introduction

Many unicellular eukaryote (protist) lineages have independently adapted to low-oxygen conditions. One of their most conspicuous adaptations are highly modified mitochondria referred to as "mitochondrion-related organelles" (MROs) that can function without oxygen. MROs have been best studied in the Metamonada, a supergroup of anaerobic protists with varied lifestyles including parasites, commensals, and free-living marine or freshwater flagellates and amoebae. All metamonad MROs studied to date are functionally reduced compared to aerobic mitochondria with their properties varying markedly across the group (Leger et al., 2017).

The best studied metamonad MROs are hydrogenosomes that occur in the urogenital tract parasite *Trichomonas vaginalis*. The 'hydrogenosomal' ATP-producing pathway uses enzymes typically not found in aerobic mitochondria such as pyruvate:ferredoxin oxidoreductase (PFO) to oxidatively decarboxylate pyruvate to acetyl-CoA and reduce ferredoxin, an acetate:succinate CoA transferase that catalyzes the transfer of CoA onto succinate to produce succinyl-CoA, and an iron-only [FeFe]-hydrogenase that reduces protons to hydrogen gas by re-oxidizing the ferredoxin. ATP is produced from ADP and P_i by the Krebs cycle enzyme succinyl-CoA synthetase by substrate-level phosphorylation, converting succinyl-CoA to succinate in the process. Trichomonad hydrogenosomes also house a variety of other pathways including a mitochondrial-type iron-sulfur cluster (ISC) biogenesis system, amino acid metabolism, and oxygen detoxification (Schneider et al., 2011; Stairs et al., 2015). Even more highly-reduced MROs called 'mitosomes' are found in the metamonad gut parasite *Giardia intestinalis*. These organelles appear to lack any capacity to generate ATP; instead they participate in Fe/S cluster biogenesis via the mitochondrial-type ISC system (Jedelský et al., 2011). Like trichomonads, *Giardia intestinalis* catalyzes acetyl-CoA production from pyruvate using PFO. However, this occurs in the cytoplasm. Acetyl-CoA is converted to acetate and ATP is produced by substrate-level phosphorylation catalyzed by the enzyme ADP-forming acetyl-CoA synthetase (ACS; EC 6.2.1.13) (Sánchez et al., 2000). The most extreme form of MRO reduction in Metamonada can be found in the oxymonad gut commensal *Monocercomonoides exilis* that has lost all traces of the organelle, making it the first truly amitochondriate eukaryote discovered (Karnkowska et al., 2016).

More recent investigations of newly discovered free-living metamonads have revealed novel configurations of MRO functions that do not fit into traditionally-defined organelle classes (Müller et al., 2012). For example, the free-living metamonad *Dysnectes brevis* encodes MRO targeted [FeFe]-hydrogenases and associated proteins but lacks organelle-targeted enzymes for substrate-level phosphorylation, suggesting that while the MRO is capable of hydrogen metabolism, ATP is most likely produced in the cytosol (Leger et al., 2017). A similar hydrogen-producing organelle and predicted cytosolic ATP production was reported in a transcriptomic investigation of the free-living metamonad *Barthelona* sp. PAP020 (Yazaki et al., 2020). Recently, the enigmatic *Anaeramoeba* species – recently-discovered distant relatives to parabasalids like *Trichomonas* – were predicted to possess hydrogenosomes that have retained more mitochondrial systems than any other metamonad studied to date (Stairs et al., 2021). The MRO of the freshwater flagellate *Paratrimastix pyriformis* was recently investigated using

spatial proteomic techniques which revealed that the organelle does not participate in either ATP
production through extended glycolysis or Fe/S cluster production, and instead produces
essential intermediates for the methionine cycle (Zítek et al., 2022). These studies of newly
discovered free-living metamonads have provided important insights into the steps by which
dramatically distinct MRO-types evolved within the Metamonada.

The multiple independent transitions from aerobically respiring mitochondria to MROs in the
eukaryote tree in most cases have involved the complete loss of the mitochondrial genome itself,
the loss of aerobic metabolism genes encoded in the nucleus, and the gain of genes encoding
novel biochemical pathways by lateral gene transfer (LGT) from bacteria, archaea or other
anaerobic protists (Roger et al. 2017). For example, the 'hydrogenosomal' metabolism enzymes
(Leger et al., 2016; Stairs et al., 2011, 2021), the ability to synthesize rhodoquinone (Stairs et al.,
2018) and oxygen defense enzymes (Jiménez-González et al., 2019) all appear to have been
acquired by LGT in multiple distinct lineages from bacteria or other protists (Gawryluk & Stairs,
2021; Roger et al., 2017). The most striking examples of the remodeling of mitochondrial
functions in anaerobic protists involve the wholesale replacement of the highly-conserved
essential mitochondrial ISC system by laterally acquired prokaryotic Fe/S cluster biogenesis
systems. In the Archamoebae, a two-protein Fe/S biosynthesis pathway (recently classified as the
minimal iron sulfur system (MIS) related to the nitrogen fixation system (Garcia et al., 2022))
was acquired from bacteria and, in free-living members of the group, was subsequently
duplicated into cytosolic and MRO-functioning versions (Nývltová et al., 2015; Záhonová et al.,
2022). A completely independent replacement of the mitochondrial ISC system occurred in the
ancestors of the breviate flagellate *Pygsuia biforma* that acquired an archaeal-type SUF system
(Stairs et al., 2014), recently classified as the SUF-like minimal system (SMS) pathway (Garcia
et al., 2022). This breviate SmsCB gene (referred to as SufCB in Stairs et al. 2014) in *Pygsuia*
biforma was then duplicated to create cytosolic and MRO-targeted SmsCB proteins. In the
oxymonads, yet another replacement of the ISC system occurred independently by the lateral
acquisition of genes encoding a more complex multi-protein SUF pathway (Karnkowska et al.,
2016; Vacek et al., 2018). This enabled the ancestral oxymonad to produce Fe/S clusters in the
cytosol, easing the evolutionary pressure to keep the canonical ISC system, paving the way for
the complete loss of MROs in *Monocercomonoides* (Karnkowska et al., 2016). Phylogenetic
analyses revealed that this SUF system is shared by a variety of preaxostylans and is likely
bacterial in origin (Vacek et al., 2018).

To further investigate the patterns and process of genome and MRO evolution in metamonads,
we sequenced the genomes of five recently-isolated free-living bacterivorous flagellates. Three
of these organisms, the 'skoliomonads' *Skoliomonas litria, Skoliomonas* sp. GEM-RC (GEMRC)
and *Skoliomonas* sp. TZLM3-RCL (RCL) were isolated from sediments of alkaline hypersaline
soda lakes (for detailed descriptions see Eglit et al., 2024) (Figure 1A-C). The other two are
shallow marine-sediment dwelling 'barthelonids' (Yazaki et al. 2020): *Barthelona* sp. strain PCE
(Figure 1D) and *Barthelona* sp. PAP020 (Yazaki et al., 2020). Unlike most previous studies of
this kind that rely on incomplete transcriptomic data, we generated high quality contiguous
genome assemblies, transcriptomes, and predicted proteomes from four of these five organisms.
Our phylogenomic analyses demonstrate that these five metamonads form a highly supported

clade, referred to as 'BaSk' (short for 'Barthelonids+Skoliomonads'), that emerges as a sister
group to Fornicata *sensu stricto* (diplomonads, retortamonads and *Carpediemonas*-like
organisms (CLOs)). Genome size and coding capacity varies considerably across these
organisms, with one possessing the smallest genome of a free-living eukaryote currently known.
Their anaerobic metabolisms and MRO functional capacities also vary considerably.
Unexpectedly, we found yet another instance of the horizontal acquisition of the SmsCB gene in
their common ancestor. While several barthelonids possess both the SmsCB gene and core parts
of a mitochondrial-type ISC system, all traces of the ISC system have been lost in the
skoliomonads. For *Skoliomonas litria*, no MRO proteins could be confidently identified,
suggesting it may lack the organelle altogether.

Results & Discussion

Near-complete draft genomes of four novel metamonads

Near-complete genome assemblies of *Barthelona* sp. PCE, *Skoliomonas litria*, *Skoliomonas* sp.
RCL and *Skoliomonas* sp. GEMRC were generated using a combination of Oxford Nanopore
long-read sequencing for assembly and Illumina short-read sequencing for error-correction post-
assembly. These assemblies were thoroughly decontaminated using the Eukfinder workflow
(Zhao et al., 2023).

To assess the quality of these assemblies we used Merqury, a k-mer counting tool that using
genomic Illumina reads, assesses the completeness of the four *de novo* long-read assemblies
(Rhie et al., 2020). Based on the Merqury analyses, all four assemblies exceeded 98%
completion. Genome completeness was further supported by the fact that, for each of the four
organisms, >99% of their respective eukaryotic-classified RNAseq reads mapped to the
corresponding genome assembly. The combination of long and short-read data afforded
assemblies that were high coverage and comprised of a few relatively large contigs (see Table 1).
A draft genome assembly of *Barthelona* sp. PAP020 was also generated using only Illumina
reads (Table 1). This relatively fragmented assembly was included with other metamonads
considered in this study to supplement our analyses concerning MRO reconstruction, metabolic
pathway inventory, and phylogenomic inference, discussed below.

Gene predictions for these assemblies were generated by an in-house pipeline. To compare the
qualities of the various gene predictions, BUSCO analyses were conducted based on the whole
genomes themselves, the gene predictions and the transcriptome assemblies (Table 2). To assess
potential improvement of the predicted protein set after curation, the gene predictions for
Skoliomonas litria were further manually curated (Table 2). The percent of BUSCO genes
identified amongst the gene predictions (i.e. predicted protein set) ranged from 63.7%
(complete+fragmented) for *Barthelona* sp. PCE to 77.9% (complete+fragmented) for
Skoliomonas sp. RCL. In general, the BUSCO scores of the gene predictions are very similar to
those from the transcriptomes but the BUSCO genes of the former are less fragmented. The
manual curation of *Skoliomonas litria* predicted proteins increased the reported BUSCO number
by only one, indicating that the manual curation did not substantially improve the gene models.
Note that these relatively low BUSCO scores are comparable to other well-characterized

metamonads assemblies which tend to have highly divergent or missing orthologs relative to
most other eukaryotes (Salas-Leiva et al., 2021; Stairs et al., 2021).

The sizes of the draft genomes and the number of proteins predicted differ markedly between
taxa (Table 1). Notably, at 10.2 Mbp, *Barthelona* sp. PCE appears to possess the smallest
genome for a free-living eukaryote currently known. It is smaller than the compact genomes of
the acidophilic alga *Galdieria sulphuraria* (13.7 Mbp, see Schönknecht et al., 2013),
Saccharomyces cerevisiae (12.07 Mbp, see Goffeau et al., 1996), and the genome of the smallest
known free-living eukaryote, the marine alga *Ostreococcus tauri* (12.56 Mbp, see Derelle et al.,
2006). Genome reduction is well known to occur in parasitic and endosymbiotic microbes
(Manzano-Marín & Latorre, 2016; Xu et al., 2020), as well as in free-living organisms, where it
is thought that compact genomes are an efficiency adaptation to nutrient-poor environments
(Giovannoni et al., 2014). However, for heterotrophic phagotrophic protists, like *Barthelona* sp.
PCE, nutrients are obtained predominantly from their bacterial prey. It is unclear whether the
foregoing generalizations hold in this case or what other forces led to genome reduction in the
barthelonid lineage.

Two of the genome assemblies have apparent telomeric repeats on the ends of some contigs.
Barthelona sp. PCE possesses two contigs with what appear to be telomeres at one end of the
contig with an unusual telomere repeat sequence of 5'-TATATGGTCT-3'. Although this
telomeric sequence is quite different from the eukaryotic 'consensus' repeat, other protists across
the eukaryote tree of life have also been reported to have highly divergent telomeric repeats (see
Fulnečková et al., 2013); the significance of these unusual telomeres is unknown. The
Skoliomonas sp. GEMRC assembly has six telomere-to-telomere assembled chromosomes and
an additional nine contigs with telomeres at only one end of the contig. The latter telomeres
have the canonical 5'-TTAGGG-3' repeating sequences.

187   The BaSk taxa are the sister group to known members of Fornicata.

To determine the phylogenetic affinities of the BaSks within the metamonads, we added
orthologs from the five draft assemblies to a previously constructed phylogenomic data set of
orthologous proteins (Brown et al., 2018; S. Kang et al., 2017). After single-protein alignment
manual curation to remove contaminants and paralogs, the final dataset of 174 highly conserved
proteins was analyzed by maximum likelihood under the site-heterogeneous model LG+C60+F+
Γ (see Figure 1E). The resulting tree confirms the placement of *Barthelona* sp. PAP020 as
previously reported (Yazaki et al., 2020) and placed barthelonid strain *Barthelona* sp. PCE as its
immediate sister. The skoliomonads form a clade with the barthelonid group, with *Skoliomonas*
litria and *Skoliomonas* sp. GEMRC branching together to the exclusion of *Skoliomonas* sp. RCL.
The 'BaSk clade' (Barthelonids + Skoliomonads) are sister to fornicates with maximum support
(i.e. both the fornicate group and the fornicate + BaSk clade are supported with 100% non-
parametric bootstrapping support, aBayes support, and approximate likelihood-ratio test support,
all evaluated under the PMSF model).

Members of Fornicata possess a particular cytoskeletal feature, a distinct fibrous arch, that defines the group (Simpson, 2003). As there are no ultrastructural data for the flagellar apparatus cytoskeleton of any BaSk member to date, it is unclear whether they should be classified within the Fornicata or remain as a distinct sister group. Regardless, the BaSks 'deep' phylogenetic placement within the metamonads make them a key group for investigations into the origins and evolutionary trajectories of genes involved in parasitism and MRO reduction within the Metamonada supergroup.

BaSks all possess highly reduced MROs and share the same core anaerobic metabolism.

We searched the genomes, transcriptomes, and predicted protein sets of all four BaSks using a variety of bioinformatic approaches (see Materials and Methods) to recover proteins related to MRO function and anaerobic metabolism. We also utilized a draft short-read Illumina genome assembly of *Barthelona* sp. PAP020 in these searches to supplement our comparisons. Several mitochondrial targeting sequence prediction programs were used to estimate the probability of mitochondrial localization of each predicted protein in each of the five genomes. We used these predicted proteins as queries against the nr database using BLAST and looked for outputs relating to mitochondrial localizing proteins. The predicted proteins from the BaSk genomes were also used as queries in a BLAST search of a database comprised of mitochondrial proteomes from six diverse eukaryote lineages and the best hits from each query were retained and inspected. Next, we utilized datasets of known metamonad MRO proteins to query the predicted protein sets from the BaSk genomes, as well as directly investigate the genomes and transcriptomes. This was done through both BLAST search and HMM profile searching. In cases where search results were ambiguous, we utilized appropriate Pfam profiles to search for protein domain information. Finally, we assessed candidate proteins by constructing phylogenetic trees of corresponding MRO proteins with additional cytosolic homologs, to further verify the probable localization of BaSk proteins. A summary of the results of all of these searches can be found as Figure 2. Below we discuss the systems we investigated in detail by first highlighting the pathways that, when present, function exclusively in MROs and are therefore strong indicators of the presence of the organelle.

We first investigated whether the BaSk genomes encoded homologs of the glycine cleavage system (GCS), an exclusively mitochondrial pathway which converts glycine to ammonia and carbon dioxide while also reducing NAD+ to NADH and generating 5,10-methylene-tetrahydrofolate (5,10-CH$_2$-THF) (Kikuchi, 1973). This pathway is commonly retained in MROs of free-living anaerobic protists including some metamonads (Leger et al., 2017), but is sometimes missing in parasitic lineages (reviewed in Roger et al., 2017). Except for *Skoliomonas litria*, all BaSks encode the complete GCS (i.e. the T-, P-, L- and H-proteins). Additionally, all BaSks possessing the GCS also have a single serine hydroxymethyltransferase (SHMT) gene. This enzyme produces serine using both glycine and the 5,10-CH$_2$-THF produced by the GCS. GCS and SHMT are found in the MROs of a variety of diverse free-living metamonads including *Paratrimastix pyriformis* (Zítek et al., 2022; Zubáčová et al., 2013) and all of the non-diplomonad fornicates (i.e. CLOs and *Chilomastix cuspidata*; see Leger et al., 2017). Complete absence of the GCS in metamonads is seen in the parasitic diplomonads *Giardia intestinalis* and

Spironucleus salmonicida (Jerlström-Hultqvist et al., 2013; Leger et al., 2017). Other parasitic metamonads such as *Trichomonas vaginalis* retain some of the GCS subunits (Mukherjee et al., 2006), but have repurposed them to function in a peroxide detoxification pathway (Nývltová et al., 2016). It was particularly unexpected, therefore, that we found neither genes encoding GCS or SHMT in *Skoliomonas litria,* making it the only known free-living metamonad completely lacking this pathway, and the only BaSk member to have no detectable MRO amino acid metabolism. The MROs within the BaSks are therefore key to understanding the reduction of amino acid metabolism that is usually only ever seen in parasitic metamonads. As previously reported (Yazaki et al., 2020), *Barthelona* sp. PAP020 contains a OsmC homolog, but we were unable to detect this enzyme in any of the other BaSk data, including the *Barthelona* sp. PAP020 genome. Because this enzyme lacks a mitochondrial targeting sequence (MTS), it is unclear where this protein localizes in *Barthelona* sp. PAP020, and if it participates in peroxide detoxification in the *Barthelona* sp. PAP020 MRO, as it does in the MRO of *Trichomonas* (Nývltová et al., 2016).

Next, we investigated the highly conserved mitochondrial Fe/S cluster (ISC) pathway that is also exclusively found within mitochondria or MROs. Unexpectedly, amongst all the BaSks, only the barthelonid PCE and PAP020 assemblies encoded *any* detectable ISC system components. These two genomes encode cysteine desulfurase NFS1 (also known as ISCS), the scaffold protein ISCU and a ferredoxin. Notably, they also encode Fe/S cluster containing NuoE and NuoF subunits of complex I, which, like core ISC components, are homologs which are only known to function in mitochondria and the MROs of other organisms. Surprisingly, with the exceptions of mitochondrial Hsp70 (discussed further below) and ferredoxin, no other genes encoding ISC pathway components were detected in the barthelonids or any other of the BaSks (see Supplementary Table 1 for ISC proteins queried). The absence of some of these proteins, like Isd11 and ferredoxin reductase (Arh1), is not unexpected as they have not been detected in most metamonads (Motyčková et al., 2023). The absence of other early ISC pathway proteins implicated in [2Fe-2S] cluster synthesis (e.g. frataxin, Jac1 (HscB) and Mge1 (GrpE)) is more puzzling as these components are usually detected in metamonads (Motyčková et al., 2023). Similarly, some late ISC pathway components implicated in [4Fe-4S] cluster synthesis (Lill & Freibert, 2020) are found in other metamonads (e.g Grx5, Isa2, NFU1 and BolA; Motyčková et al., 2023), but were not detected in any BaSks (Supplementary Table 1). This is strange given that the barthelonids possess mitochondrial NuoF subunits that typically house an [4Fe-4S] cluster (Ohnishi, 1998). This means that if these NuoF subunits are indeed found within the MROs of the barthelonids, as they are with other eukaryotes, their [4Fe-4S] clusters are being produced by an unknown mechanism.

Despite the lack of ISC system proteins in most BaSks, all BaSk members appear to encode a similar partial cytosolic CIA system that includes apoprotein targeting components CIA1, CIA2, and NAR1, the NBP35 scaffold protein, and TAH18, which aids in maturation of Fe/S cluster proteins. This reduced CIA pathway is not unlike the CIA pathway found within *Giardia intestinalis* (Pyrih et al., 2016).

BaSks possess simple archaeal-type SUF-like minimal system (SMS) fusion proteins.

We investigated the possibility that the lack of ISC system components in BaSk members may be complemented by the presence of an alternative Fe/S cluster system such as the minimal Fe/S (MIS) or sulfur mobilization (SUF) systems found in a number of other anaerobic protists (Anwar et al., 2014; Garcia et al., 2022; Stairs et al., 2014; Vacek et al., 2018; Žárský et al., 2021). In all of the BaSk genomes, we were able to identify a simple archaeal-type SUF-like minimal system (SMS) consisting of a gene encoding an SmsCB fusion protein that was most similar to the simple archaeal-type SmsCB systems found in the breviate *Pygsuia biforma* (Stairs et al., 2014), the anaerobic jakobid *Stygiella incarcerata* (Leger et al., 2016) and gut commensal opilinatan stramenopile *Blastocystis* (Tsaousis et al., 2012; Yubuki et al., 2020). To investigate the origin of this protein, we performed phylogenetic analysis separately for the SmsC and SmsB domains (Supplementary Figures 1 & 2) before concatenating the domains together for the final tree (Figure 3). The BaSk SmsCB proteins form a strongly supported clade that is sister group to all archaeal-type SmsCB fusion proteins from anaerobic protists. Consistent with previous reports, the anaerobic protistan clade of SmsCB fusion proteins is sister group to the orthologs from members of the Methanomicrobiales order of Archaea (Tsaousis et al., 2012). This phylogenetic pattern is most easily explained if there were an original transfer and fusion of SmsCB to an anaerobic protist from a methanomicrobiales donor, followed by a series of eukaryote-to-eukaryote LGT events amongst disparate lineages of anaerobic eukaryote lineages. The deeply branching position of the BaSk clade amongst the anaerobic protists suggests it is possible that an ancestral BaSk was the first recipient of these genes and an early offshoot from the BaSk lineage passed the fusion gene to other anaerobic protists, although it is impossible to rule out alternative scenarios. Regardless, this SMS system is phylogenetically distinct, and has a separate origin from, the more complex SUF system found within the metamonad preaxostylans (i.e. oxymonads and *Paratrimastix pyriformis*) (Vacek et al., 2018) and therefore represents an interesting case of convergence by LGT within the Metamonada. The resulting reduction and loss of the ISC system following acquisition of a SMS system is also similar between these two lineages and the breviate *Pygsuia biforma*. The gain of an alternate Fe/S cluster synthesis system in these cases likely 'preadapts' the ancestral organism to the loss of the ISC system in some of the lineages (e.g. an ancestor of *Skoliomonas* sp. GEMRC, *Skoliomonas* sp. RCL and *Skoliomonas litria*). Unlike *Pygsuia biforma* and the preaxostylans, *Barthelona* sp. PCE and *Barthelona* sp. PAP020 retain a few core components of the early ISC system in addition to possessing the novel SMS system, suggesting that their MRO ISC system is at least partially functional. This retention of the ISC proteins may be related to the possession of Fe/S cluster-containing proteins such as the NuoE and NuoF complex I proteins that are predicted to function in the MRO. Although we cannot definitively determine the subcellular localization of the SmsCB in BaSks bioinformatically, it is likely that this system is cytosolic as in other anaerobic protists such as *Blastocystis* spp. (Tsaousis et al., 2012). If it is indeed cytosolic, it is unclear whether it would function in coordination with the cytosolic CIA Fe/S biogenesis system. One possibility is that the CIA system and the SmsCB system may function in parallel with each responsible for the assembly of Fe/S clusters for different specific subpopulations of apoproteins. Alternatively, the two systems may work together, similar to how the ISC system coordinates with the CIA pathway (Pyrih et al., 2016). In the latter scenario, the SmsCB fusion protein would

provide the CIA pathway with a sulfur-containing intermediate, which the CIA pathway would
then use to assemble Fe/S clusters and insert them into recipient apoproteins.

In any case, the skoliomonads completely lack any dedicated ISC components and do not
possess proteins with identifiable cysteine desulfurase domains as assessed by HMM profile
searches and phylogenetic analysis (see Supplementary Figure 3). This is surprising because for
most Fe/S cluster synthesis systems, cysteine is the source of sulfur and a cysteine desulfurase is
the key enzyme involved in mobilizing it. We suspect that the SmsCB system of these organisms
may directly utilize sulfide from the environment as the sulfur source for Fe/S clusters, as
previously suggested for *Pygsuia biforma* (Stairs et al., 2014). Indeed, direct sulfide utilization
has been demonstrated for methanogenic archaea that possess a simple cysteine-desulfurase-
lacking SMS pathway and which also live in sulfidic conditions (Liu et al., 2010). The exact
mechanism by which such organisms construct Fe/S clusters is unknown and more work needs to
be done to understand their biogenesis in these organisms.

Presence or absence of other conserved mitochondrial proteins in the BaSk taxa.

The next highly conserved mitochondrion/MRO proteins that we investigated were the
mitochondrial orthologs of the molecular chaperones chaperonin-60 (Cpn60) and hsp70. To
distinguish cytosolic Chaperonin Containing TCP-1 (CCT) subunits from mitochondrial Cpn60
(GroEL) homologs and cytosolic and endoplasmic reticulum (ER) paralogs of hsp70 from their
mitochondrial Hsp70 (i.e. dnaK orthologs), we used phylogenetic analyses. For *Skoliomonas*
litria and *Skoliomonas* sp. RCL, all Cpn60 candidates identified distinctly branched with CCT
subunit clades (See Supplementary Figure 4); no bona fide mitochondrial Cpn60s could be
identified from these species. *Skoliomonas* sp. GEMRC, *Barthelona* sp. PCE, and *Barthelona* sp.
PAP020 appear to have both CCTs and mitochondrial Cpn60. Phylogenetic analysis of Hsp70
proved more difficult, due in part to a greater degree of similarity between paralogous copies of
the chaperone. To supplement this analysis, signature insertion-deletions in the multiple
alignments were identified that were used in conjunction with well-annotated Hsp70 yeast
homologs (Nelson et al., 1992) to group sequences into cytosolic/ER versus mitochondrial type
Hsp70 groups (see Supplementary Figure 5). This information was then checked against the
phylogeny for subcellular localization predictions of BaSk Hsp70 candidates. From this analysis,
mitochondrial-type Hsp70s were identified in all BaSks except *Skoliomonas litria*. Within
MROs, Cpn60 and Hsp70 aid in protein translocation into the matrix and help newly translocated
proteins adopt their native structure. These chaperones are ubiquitously predicted to function in a
variety of MROs, so it is notable that they are missing in some of the BaSks data. For
Skoliomonas litria, we did not detect any genes encoding MRO chaperones, while *Skoliomonas*
sp. RCL appears to contain Hsp70 but lacks Cpn60. This further highlights the highly reduced
nature of the MROs found within these protists, especially within *Skoliomonas litria*.

All mitochondria and MROs have membrane associated proteins, including the TIM/TOM
complex proteins involved in protein translocation into the organelle. To search for these, we
used HMM profiles as many of these sequences are highly divergent and not detectable by
pairwise alignment-based search algorithms. The main channel forming protein in the TOM
complex, Tom40, was only identifiable in *Skoliomonas* sp. RCL and *Skoliomonas* sp. GEMRC,

while the receptor Tom70 was identified in *Barthelona* sp. PCE, *Barthelona* sp. PAP020, and *Skoliomonas* sp. RCL. TIM subunits were generally absent except for Tim17 that was identified in *Barthelona* sp. PCE only. Except for *Skoliomonas litria*, all BaSk members appeared to have the co-chaperone Pam18 subunit of the presequence translocase-associated motor (PAM) complex in addition to the core mtHsp70 subunit. Pam16 was only identified in *Skoliomonas* sp. GEMRC and Pam17 appeared to be absent.

In mitochondria, mitochondrial processing peptidases (MPPs) and presequence proteases (PreP) remove the MTS after protein import into the organelles (Garrido et al., 2022). We identified MPP/PreP homologs only in *Skoliomonas litria*, *Skoliomonas* sp. GEMRC, and *Skoliomonas* sp. RCL and none had predicted MTSs. We further investigated their domain structures and phylogenetic position amongst MPP/PreP-related protein families in eukaryotes and prokaryotes. For our phylogenetic analysis we used a recently published multiple alignment of all major families of M16 proteases (Garrido et al., 2022) supplemented with top 20 BLAST hits in the nr database to each of the *Skoliomonas litria* sequences. Our phylogenetic analyses (Supplementary Figure 6A) showed that skoliomonad sequences branch in multiple different positions in the tree, but none are closely related to eukaryote mitochondrial MPPs or PrePs. All but one of these proteins have a four-domain structure that is characteristic of PreP and SPPs but distinct from the two domain structure of the MPP family (Garrido et al., 2022). One *Skoliomonas* sp. RCL sequence has two domains that appear to have evolved separately from the MPP group (Supplementary Figure 6B). Notably, the two *Skoliomonas litria* M16 homologs branch with skoliomonad orthologs in two distinct clades. One of these branches from within a group of entirely bacterial PreP or MAG peptidase sequences and the other is a sister group of uncharacterized homologs from invertebrates (Metazoa) which lack predicted MTSs (except for one homolog in *Branchiostoma* that is the result of a recent duplication). This analysis suggests that none of the M16-related proteases in skoliomonads are orthologs of mitochondrial MPP or PreP; they are likely M16 proteases with separate cytosolic functions.

A candidate mitochondrial carrier protein (MCP) was identified in *Skoliomonas litria* but its substrate specificity could not be determined by phylogenetic analysis (Supplementary Figure 7). Whether this protein functions in the MRO membrane is unclear as MCPs are also known to function in peroxisomes or other subcellular organelles (Mazurek et al., 2010). Finally, a candidate homolog of Sam50 was identified in *Skoliomonas litria* and *Skoliomonas* sp. GEMRC. However, in phylogenetic analyses these proteins did not emerge within the clade of mitochondrial Sam50 orthologs, but instead formed an independent group emerging half-way between mitochondrial Sam50 and bacterial BamA clades (Supplementary Figure 8A). Orthologs of this Sam50/BamA-like protein were not detectable in any of the other BaSk genomes, even when this sequence was included in the HMM profile used for searching. To determine if this Sam50/BamA-like protein folded into a structure resembling the mitochondrial orthologs, we used AlphaFold 2.0 (Jumper et al., 2021) to predict its structure. The resulting structures (Supplementary Figure 8B, C) show that while these candidate Sam50 proteins possess what could be a intermembrane POTRA domain, the beta-barrels are smaller and adopt a different confirmation compared to Sam50 homologs of yeast (Supplementary Figure 9) and other well characterized examples (Takeda et al., 2021).

Among the five MRO predicted proteomes we present here, *Skoliomonas* sp. GEMRC and *Skoliomonas* sp. RCL contain the most candidate MRO translocon proteins, which is similar to the repertoire found in *Giardia intestinalis* (Leger et al., 2017) with the notable exception of the absence of GrpE in all members of BaSk. *Barthelona* sp. PCE, *Barthelona* sp. PAP020 and *Skoliomonas litria* possess a remarkably reduced MRO translocon complex by comparison. This may be due to the highly divergent nature of membrane bound MRO proteins, which makes detecting such proteins challenging (Pyrihová et al., 2018). Alternatively, in the case of *Skoliomonas litria* which lacks all the highly conserved GCS, ISC, protein import/refolding proteins of MROs, it is unclear if the Sam50/BamA-like protein, the MCP and MPP/M16 metalloprotease homologs are targeted to an organelle at all. It is possible that *Skoliomonas litria*, like the distantly related *Monocercomonoides exilis*, might completely lack a mitochondrial compartment and these proteins have taken on roles elsewhere in the cell. Further work needs to be undertaken to determine the cellular localization of these proteins and, if it exists, the function *Skoliomonas litria*'s MRO would carry out in the absence of all known conserved mitochondrial pathways and systems.

MTS prediction is a known challenge in inferring MRO function in anaerobic protists, including metamonads (Jerlström-Hultqvist et al., 2013; Schneider et al., 2011; Tanifuji et al., 2018), due to highly divergent or missing MTSs in these organisms. It has been suggested that the lack of a long positively charged MTS in metamonads is due to the loss of the proton gradient in the organelles (Garg et al., 2015), though recent analysis of canonical MTSs found in the Anaeramoebae – deep-branching metamonads – suggest that this may not always be the case (Stairs et al., 2021). Previous work has shown that some MRO proteins rely on cryptic internal signals to target proteins to the organelle with high fidelity (Garg et al., 2015; Mentel et al., 2008). Furthermore, most available MTS prediction software tools were trained on experimentally validated datasets of targeted proteins that are lacking for most protists. For these reasons, making accurate MTS predictions in metamonads is exceedingly difficult. Here, we employed several different MTS prediction software tools to find and annotate these signals in BaSks (see Supplementary Table 1). We found several examples of false positives in *Skoliomonas litria* and *Skoliomonas* sp. GEMRC, as homologs of transposons and nuclear targeting proteins were predicted to possess an MTS by some targeting prediction software with high confidence. Because of this, we considered proteins with MTS predictions from two or more MTS prediction software to be "strongly supported" targeting signals. Based on this criterion, *Skoliomonas litria* lacked any strongly supported targeting signals, while strongly supported targeting signals were rare in all other BaSk members. *Barthelona* sp. PCE contains strongly supported MTSs predicted on three proteins: an [FeFe]-hydrogenase, the T-protein of the glycine cleavage system (GCST), and on the IscU scaffold protein. *Barthelona* sp. PAP020 also contained a strongly supported MTS on its ortholog of IscU, as well as a different component of the glycine cleavage system (GCSP2). Another MTS detected was found on presequence translocated-associated motor protein PAM18 in *Skoliomonas* sp. GEMRC. Interestingly, in *Skoliomonas* sp. RCL we find that a cytosolic CCT-gamma subunit contains a strongly predicted MTS, where in other BaSks, all cytosolic CCT subunits appear to be missing a MTS. However, when aligned to other BaSk CCT-gamma subunits, it does not appear to have an obvious N-terminal extension. As *Skoliomonas* sp. RCL has no mitochondrial Cpn60 homolog

but does contain proteins which are homologous to MRO-localized proteins in other
metamonads, it is possible (but unprecedented) that this CCT subunit has replaced the
mitochondrial homolog in this species by acquiring an MTS. *Skoliomonas* sp. RCL does not have
an additional copy of the gamma subunit of CCT. Follow up localization experiments are needed
to deduce the subcellular localization of this particular CCT subunit.

The BaSks share a core ATP generation pathway.

All BaSk members share a similar hypothetical ATP production pathway that is reminiscent of
the pathway found in *Giardia intestinalis* (Jedelský et al., 2011; Sanchez & Müller, 1996). In this
pathway, pyruvate:ferredoxin oxidoreductase (PFO) catalyzes the oxidation of pyruvate in the
presence of ferredoxin and CoA to produce acetyl-CoA, CO_2 and reduced ferredoxin. Acetyl-
CoA is then transformed to acetate by ACS and, in the process, converts ADP to ATP.
Ferredoxin is reoxidized by [FeFe]-hydrogenase (HydA) that passes electrons to protons creating
H_2 gas. Notably, pyruvate-formate lyase (PFL) was also detected in all BaSk members, allowing
an alternative pathway of acetyl-CoA production from pyruvate (Stairs et al., 2011; see
Supplementary Figure 10). Note that there are multiple copies of both HydA and PFO in each
BaSk member, some of which possess additional domains encoding ferredoxin or flavodoxin-
like proteins (see Supplementary Table 2). No components of typical ATP production pathways
found in trichomonad hydrogenosomes, such as acetate:succinyl-CoA transferase or succinyl-
CoA synthase (Lahti et al., 1992; Van Grinsven et al., 2008), were detected in any of the BaSks.
Though [FeFe]-hydrogenase is present in each BaSk member, none of the [FeFe]-hydrogenase
maturase proteins (HydG, E or F) were detected. This is notable because these maturases, when
present, are always targeted to mitochondria or MROs. This, combined with the lack of MTS
detected in any of the proteins from the above pathway, leads us to assume that ATP and H_2 gas
are likely produced in the cytosol of the skoliomonads.

However, the pathways in the barthelonids appear to be slightly different in function and likely
localization. Both *Barthelona* sp. PCE and *Barthelona* sp. PAP020 encode NuoE and NuoF
NADH-dehydrogenase subunits of complex I that are known to function in the
mitochondrial/MRO matrix in eukaryotes. It is widely thought that in the anaerobic protists that
possess them, NuoE and NuoF likely form a complex with an MRO-localized HydA to perform
simultaneous NADH and ferredoxin (Fd$^-$) oxidation by an electron confurcating reaction that
produces H_2 gas, NAD+ and Fd (Dyall et al., 2004; Stairs et al., 2015). For this reason, and the
fact that one of the HydA proteins of *Barthelona* sp. PCE and *Barthelona* sp. PAP020 have a
predicted MTS, we infer that both barthelonids likely have MRO-associated hydrogenase
activity, as recently suggested for *Barthelona* sp. PAP020 by Yazaki et al. (2019). However,
unlike the hypothesis of Yazaki and colleagues (see Fig. 4b in Yazaki et al. 2019), we suggest
that the reduced ferredoxin may, in fact, get produced within their MROs by PFO activity. If so,
it is also possible that at least one of the two ACS homologs in each of these organisms is also
MRO-localized and produces ATP within the organelle, though no MTSs are predicted on any of
the ACS homologs. Further localization experiments in these barthelonids are needed to test
these hypotheses.

Previous studies have shown that some of the enzymes involved in ATP synthesis found within metamonads have a complex phylogenetic history. In particular, ACS is thought to have been transferred into the metamonads multiple times through LGT (Leger et al., 2017). To determine the evolutionary histories of the BaSk ACSs, we constructed a phylogenetic tree containing ACSs from a variety of metamonads, which can be found as Supplementary Figure 11. Our results show that, as in previous studies, most metamonad ACSs branch in one of two clades, known as ACS1 and ACS2. The ACS homologs found in the BaSks also form two distinct groups; one appears to be related to ACS2 found within *Carpediemonas membranifera*, *Ergobibamus cyprinoides,* and *Trimastix*. Additional ACS2 homologs from *Monocercomonoides* and *Paratrimastix pyriformis* were also identified, which also branch with the metamonad ACS2 group. This suggests that ACS2 was inherited in the last common ancestor of Preaxostyla and the fornicates + BaSk. The BaSk also possess a second ACS homolog which branches separately from other metamonad ACSs. This enzyme seems to be most closely related to ACS found within *Streblomastix*, *Monocercomonoides* and *Blastocystis*, and is distinct from other types of ACS in the tree. As previously reported by Yazaki et al. (2019) in their analysis of the *Barthelona* sp. PAP020 transcriptome, this is likely to be a novel homolog of ACS (ACS3) and was probably transferred into metamonads through a LGT event distinct from the event that gave rise to ACS1 and ACS2. Our analysis also suggests that the ACS found within *Spironucleus salmonicida* MROs may not be ACS2, as previously reported (Leger et al., 2017) because, in our analyses, the *S. salmonicida* ortholog (denoted ACS* in Figure 4 and Supplementary Figure 11) did not form a clade with other metamonad ACS2 sequences but instead grouped with several TACK archaea, although the support values for this relationship are low. A diagram summarizing the gains and losses of these proteins, along with other key metabolic processes and proteins mentioned above, can be found as Figure 4. These patterns suggest that there have been at least three, and possibly four, independent transfers of ACS into the metamonads. These transfers occurred at very different points in time along the evolution of various metamonad groups and highlights the role of LGT in the evolution of anaerobic metabolism.

To investigate the expression level of these key metabolic enzymes, we mapped RNASeq reads to each gene model, then normalized read counts and compared the fragments per kilobase of transcript per million mapped reads (FPKM) value amongst these gene models (see Supplementary Table 1). We found that some enzymes relating to anaerobic metabolism were amongst the top 100 most highly expressed genes in all four of the BaSk queried, though the specific anaerobic enzymes in this top 100 list differed between each species (see Supplementary Table 3). For *Skoliomonas litria*, *Skoliomonas* sp. GEMRC, and *Skoliomonas* sp. RCL this included two or more copies of genes encoding PFL. For *Skoliomonas litria* and *Skoliomonas* sp. RCL, this also included one copy each of a gene encoding PFO. *Skoliomonas litria*, in addition to PFL, had a copy of an ACS and a malic enzyme gene in its top 100 most highly expressed gene set. *Skoliomonas* sp. GEMRC was the only organism with a SmsCB gene represented its top 100 expressed list; other BaSk members have comparatively lower expression levels of this gene.

Implications for the evolution of MROs within the Metamonada supergroup.

Based on our reconstructions of MRO properties on the metamonad tree, we suggest that the last common ancestor of BaSk possessed an already highly reduced MRO compared to the common ancestor of fornicates + BaSk (Figure 4). The barthelonids PCE and PAP020 seem to have functionally reduced MROs, but their predicted MRO proteomes retain similar functions to other free-living metamonads such as the GCS and serine metabolism, hydrogen production and Fe/S cluster biogenesis by a simple ISC system. On the other hand, the MROs of *Skoliomonas* sp. RCL and *Skoliomonas* sp. GEMRC appear to be unique amongst metamonads, as we predict that the organelles only function in glycine and serine metabolism and are unlikely to produce H_2. For *Skoliomonas litria*, the only proteins typically associated with mitochondria and MROs detected were a candidate MCP and a homolog of Sam50; of these, only the latter is exclusively associated with mitochondrial organelles in eukaryotes. The complete lack of all other proteins homologous to common MRO-localizing proteins could indicate that an MRO may not even exist in this organism. In any case, this extreme reductive evolutionary path may relate to the acquisition of a simple fused SmsCB gene that was acquired by LGT in a common ancestor of BaSk. This acquisition then led to the simplification of the ISC system in the *Barthelona* clade, and the outright loss of the ISC system in the skoliomonads, as Fe/S proteins for organellar functions were no longer needed.

Regardless of the evolutionary forces involved, the BaSks demonstrate reduction in MRO function that is independent of the loss seen in other metamonad groups, such as diplomonads and oxymonads (Karnkowska et al., 2016; Leger et al., 2017). This is remarkable given that many metamonads that display a similar loss of MRO functions are parasitic or commensalistic and the nutrient-rich endobiotic lifestyle has often been assumed to be part of the explanation for extreme MRO proteome streamlining (Karnkowska et al., 2016; Stairs et al., 2015). Instead, in the case of the free-living BaSk members, it appears that gain of a simple SMS system was amongst the first triggers leading to the extreme reductive MRO evolution in this group.

<u>Conclusions</u>

We have generated high quality draft genomes, transcriptomes, and curated predicted proteome sets for a novel group of free-living metamonads. The placement of BaSk within the tree of Metamonada as the sister group to the Fornicata make them a key group to study in terms of genome evolution, LGT events, and adaptation to various environmental niches. These data place us several steps forward towards the goal of representing diverse and historically under-sampled microbial eukaryotes in the tree of eukaryotic life.

Our findings demonstrate the diversity of MRO configuration and function amongst the metamonads. As a group, BaSks display a level of MRO reduction not typically seen in free-living metamonads and which is independent of the reduction displayed in the parasitic diplomonads. BaSk members share many interesting parallels with the MRO reduction and loss seen in other metamonads, as all BaSks possess a SMS Fe/S cluster system that has, in some

cases, replaced the canonical ISC system. This is the second discovered instance of an LGT
event that has replaced the ISC system in metamonads with a SUF-like system and serves to
highlight the role of adaptive LGTs in MRO evolution. Based on these findings, we speculate
that the last common ancestor of the BaSks possessed both a compact, streamlined genome with
a highly reduced MRO, and was markedly different from the last common ancestor of the
fornicates. In many of the BaSk taxa, further work is needed to elucidate the exact subcellular
localization of several candidate MRO proteins including enzymes in the anaerobic ATP
synthesis pathway and to determine the breadth of functions carried out by the MROs in this
newly discovered group. For *Skoliomonas litria*, which lacks all known mitochondrial pathways,
it will be especially important to investigate the localization of its anaerobic ATP synthesis
pathway, its MCP, and the Sam50/BamA-like protein to determine whether this free-living
flagellate has completely dispensed with the organelle.

Materials and Methods

Cell cultivation & microscopy

Established cultures of isolates *Skoliomonas litria*, *Skoliomonas* sp. RCL and *Skoliomonas* sp. GEMRC (Eglit et al. 2024) were maintained in variants of CR media (Gigeroff et al., 2023); *Skoliomonas litria* in 50 mL of 40 ppt CR with 3% LB, *Skoliomonas* sp. GEMRC in 50 mL of 40 ppt CR with 1% LB and 3 sterile wheat grains, *Skoliomonas* sp. RCL in 50 mL of 25 ppt CR with 1% LB and 3 sterile wheat grains. All were inoculated into fresh media every 7-10 days based on culture density, with a second inoculation after 24 hours. Cultures of isolate 'Barthelona sp.' PCE (Yazaki et al. 2019) were maintained in sterile filtered seawater (Northwest Arm, Halifax, Nova Scotia) supplemented with 3% LB, and kept at 16□. Cell growth was monitored weekly, and subculturing was done every 3-4 weeks depending on cell density by transferring around 20% of the old culture into the new culture.

Cells were imaged with differential interference contrast optics on a Zeiss Axiovert 200M microscope fitted with a AxioCam M5 camera (Carl Zeiss AG).

Nucleic acid extraction & sequencing

For *Skoliomonas litria*, *Skoliomonas* sp. GEMRC, *Skoliomonas* sp. RCL, and *Barthelona* sp. PCE: cell material for DNA and RNA extraction was harvested from 1-2 L of dense cell culture. The cells were collected by centrifugation in 50 mL tubes for 8 min at $2000 \times g$ and at 4□. The pellets were then combined into two 15 mL tubes that were spun again as before. The recovered pellets were resuspended in sterile filtered (0.2 µm) spent culture media and layered over Histopaque-1077, and then subjected to centrifugation at $2000 \times g$ for 20 min at room temperature. The protist-containing layer between the Histopaque and the top media was collected, and the procedure was repeated for a total of two rounds of Histopaque separation. Finally, the cells were diluted by 5 mL (10 volumes) of sterile spent-media and pelleted for 8 min, at $2000 \times g$, 4□. The cells were then resuspended in 2 mL of fresh media.

Genomic DNA was purified using a variety of techniques including traditional phenol-chloroform extraction, CTAB extraction, the QIAGEN Genomic-tip kit, and the QIAGEN MagAttract HMW DNA kit. RNA was extracted using Trizol according to Invitrogen specifications. RNA and DNA were sent to Genome Quebec for library construction and NovaSeq 6000 Illumina sequencing. In the case of the RNA, poly-A selection was performed to enrich for eukaryotic reads. High molecular weight DNA from each species was used to construct a 1D ligation Nanopore library (LSK108 for *Barthelona* sp. PCE, LSK109 - EXP-NBD104 for *Skoliomonas litria* and *Skoliomonas* sp. GEMRC, and LSK308 for *Skoliomonas* sp. RCL) and was sequenced using a MinION flowcell (FLO-MIN106 (R9.4) for *Skoliomonas litria* and *Skoliomonas* sp. GEMRC, and FLO-MIN107 (R9.5) for *Skoliomonas* sp. RCL and *Barthelona* sp. PCE).

For *Barthelona* sp. PAP020, cells were cultured as described in Yazaki et al. (2019). Cells were harvested by pelleting at 1500 rpm for 5 minutes at 4□ and were washed with PBS prior to

622 phenol chloroform DNA extraction. The purified DNA was sequenced on an Illumina Hiseq
623 2500 PE using the Trueseq library prep.

Genome/transcriptome assembly, & gene annotation

For *Skoliomonas litria*, *Skoliomonas* sp. GEMRC, *Skoliomonas* sp. RCL, and *Barthelona* sp.
PCE, Illumina NovaSeq reads were trimmed using Trimmomatic v0.36 (Bolger et al., 2014).
RNASeq reads were assembled using Trinity v2.6.6 (Haas et al., 2013) and checked for
multiplexing contamination using WinstonCleaner
(https://github.com/kolecko007/WinstonCleaner). Bacterial contamination was removed from the
Trinity assemblies using Anvi'o v5 (Eren et al., 2021).

Raw MinION sequencing data was basecalled using Albacore v2.1.3 (Albacore has since been
replaced by Guppy – see Wick et al., 2019). These data were assembled using Flye v2.3
(Kolmogorov et al., 2019), Raven v0 (Vaser & Šikić, 2021) Canu v1.7 (Koren et al., 2017) and
ABruijn v2.2b (Lin et al., 2016). The resulting assemblies were evaluated for completeness and
contiguity by comparing assembly size, contig number, N50 and L50 values, and ALE scores
(Clark et al., 2013). The ABruijn assembly was chosen moving forward for *Skoliomonas litria*,
Skoliomonas sp. RCL and *Barthelona* sp. PCE. For *Skoliomonas* sp. GEMRC, the Canu
assembly was used. Raw MinION output was used to improve the base-call accuracy of these
assemblies using Nanopolish version 0.8.4 (Loman et al., 2015) and Illumina reads were used to
error-correct the assemblies using Unicycler version 0.4.3 (Wick et al., 2016). Bacterial
contamination was assessed and removed from the resulting contigs using read coverage, GC
content, and BLAST search results (Altschul et al., 1990). The Eukfinder workflow was used to
remove prokaryotic contaminants from the BaSk genome assemblies (Zhao et al., 2023)

Gene prediction utilized a pipeline described in Salas-Leiva et al. (2021). Briefly, RepeatMasker
(Flynn et al., 2020) was first used to mask the repetitive regions of the genome. RNA-seq reads
were then mapped to the genome using HISAT2 (Kim et al., 2019). GeneMark-ET (Lomsadze et
al., 2014) then utilized this mapping information to generate gene coordinates and intron splice
sites, creating a training set of genes. This training set was then used to train AUGUSTUS
(Stanke et al., 2004), which generated the gene predictions. PASA (Haas et al., 2003) was then
used to validate the coordinates of these gene models. The predicted protein set was checked for
completeness using the obd9 BUSCO v3.0.1 database (Simao et al., 2015).

To estimate genome completeness we used Merqury v.1.0 (Rhie et al., 2020) with a kmer size of
17 (estimated using Merqury). The Illumina reads used for the Merqury analysis were
decontaminated using the Eukfinder workflow (Zhao et al., 2023) before genome completeness
estimation. Additionally, RNA-seq mapping was used as a measure of genome completeness,
and RNA-seq reads were decontaminated using the same method mentioned used for the DNA
Illumina reads.

To construct the *Barthelona* sp. PAP020 genome, Illumina reads were assembled using Spades
(Prjibelski et al., 2020) with default parameters. Bacterial reads were identified by metaBAT
(Kang et al., 2015) and CheckM (Parks et al., 2015). Once the identified bacterial reads were
removed, the remaining Illumina reads were then used to assemble a new genome. This process

was repeated until no bacterial contamination was detected. The final contigs were verified by
checking kmer content and were also checked against the *Barthelona* sp. PAP020 transcriptome
(Yazaki et al., 2020).

Gene expression analysis

666 Expression of predicted protein-coding genes was quantified by mapping RNASeq reads to their
667 respective long-read genome using HISAT2 (Kim et al., 2019). Using this mapping information
668 and gene model coordinates, the expression level was normalized using Cufflinks (Cuffquant and
669 Cuffnorm with default settings)(Trapnell et al., 2010) to compute a FPKM value for each gene.

Phylogenomic analysis

The phylogenomic dataset was constructed based on a previously published set of 351 highly
conserved protein orthologs (Brown et al., 2018), with 177 orthologs excluded due to deep
paralogies or insufficient coverage across pertinent species in the analysis. Each of the 174
remaining ortholog alignments were used to construct single-protein phylogenies, and were used
to screen out non-orthologous data. The final 174 alignments were realigned using mafft v7.310
einsi, trimmed using BMGE v1.0 (default settings) (Criscuolo & Gribaldo, 2010), and were
concatenated together and used to infer a maximum-likelihood tree under the
LG+C60+F+gamma model of evolution with IQTree v1.5.5 (Nguyen et al., 2015). A PMSF
model based on this tree and mixture model (Wang et al., 2018) was estimated and used to
calculate branch supports using non-parametric bootstrapping, approximate Bayes (aBayes;
Anisimova et al., 2011) and approximate likelihood-ratio support (aLRT; Guindon et al., 2010).

Inventory of MRO and anaerobic metabolism proteins

Proteins of interest were searched for in the predicted proteome, transcriptome, and genome of
each species using both NCBI BLAST (using blastp, tblastn and blastx) and HMM profile
searching using Hmmer v3.1b2 (hmmer.org). For each search, queries included both orthologous
sequences from closely related metamonads and, once identified by these methods, the orthologs
from other BaSks. The protein sets of the BaSk taxa were also used as queries in BLAST
searches against the mitochondrial proteomes from *Andalucia godoyi* (Genbank accession
VRVR00000000; see Gray et al., 2019), *Tetrahymena thermophila* (predicted protein Genbank
accession EAR80512 to EAS07932, see Eisen et al., 2006; Smith et al., 2007), *Arabidopsis*
thaliana (NCBI Bioproject PRJNA10719 and SAMN03081427; see Cheng et al., 2017; The
Arabidopsis Genome Iniative, 2000), *Homo sapiens* (International Human Genome Sequencing
Consortium, 2001; mitochondrial proteins from UniProt (GO:0005739)), *Acanthamoeba*
castellanii (NCBI Bioproject PRJNA599339 and PRJNA487265; see Gawryluk et al., 2014;
Matthey-Doret et al., 2022), and *Saccharomyces cerevisiae* (https://www.yeastgenome.org/; see
Engel et al., 2014). The top five hits for each query sequence were examined manually. Proteins
relating to mitochondrial protein translocation were searched for using both general HMM
profiles (Leger et al., 2017) and profiles tailored for searching in metamonads. The latter profiles
were created by retrieving the profiles from Pfam for each protein of interest and then retrieving
the best hits using the Pfam profile against the several publicly available metamonad sequencing
data databases with hmmsearch. Using an evalue cutoff of <0.01, metamonad sequences were

added to the corresponding Pfam seed alignment using the mafft –add function. If there were
more than two detected sequences from metamonads, an alignment was made with these
sequences and was then used to build a profile to search for more hits. Each candidate protein
identified was used to reciprocally search the BLAST database to confirm protein identity. In
ambiguous cases, the Pfam and InterPro databases were searched to analyze protein domain
structure.

Phylogenies for key proteins identified were constructed to aid in identification in cases with
deep paralogy (Hsp70 and Cpn60). Phylogenies were also made to identify the origin of enzymes
that act in anaerobic metabolism. Alignments from Leger et al. 2017 or Tsaousis et al., 2014
were utilized as a starting point, and we added the corresponding BaSk sequences. Where
relevant, the protein sequences of the top 10 BLAST hits from each BaSk homolog were also
added to the appropriate alignments. In cases where no starting alignments were available, a
preliminary alignment was created by retrieving the top 1000 blast hits and reducing the
sequences to 100-120 sequences using CD-HIT v.4.6 with recommended word sizes. These
sequences were aligned using mafft einsi with default settings and were trimmed using trimal
v1.4.rev15 (Capella-Gutiérrez et al., 2009) with the -gappyout setting. These phylogenies were
inferred using the LG+C20+F+gamma model of evolution with IQ-TREE.

Mitochondrion-related organelle protein localization predictions

Putative mitochondrial targeted proteins and mitochondrial targeting sequences (MTS) were
predicted using the following mitochondrial targeting peptide prediction programs: TargetP
v1.1b (Emanuelsson et al., 2007), MitoProt II v1.101 (Claros & Vincens, 1996), and MitoFates
v1.1 (Fukasawa et al., 2015)). Additionally, the predicted proteins were used to search both the
nr database and a custom database of known mitochondrial proteins.

Protein structure prediction and alignment

The structure of candidate Sam50/BamA-like proteins in *Skoliomonas litria* and *Skoliomonas* sp.
GEMRC were predicted using the ColabFold v1.5.2 server (Mirdita et al., 2022). The β-barrel
portion of the *Skoliomonas litria* Sam50/BamA-like protein was aligned to the yeast Sam50
protein structure (Takeda et al., 2021) using the PyMOL "cealign" function.

SMSCB (SUFCB) phylogeny

The SmsCB fusion protein identified in each BaSk member was added to previously constructed
alignments (Leger et al., 2016) of SmsC and SmsBD. This dataset was aligned using mafft einsi
with default settings and was trimmed by hand. The dataset was then concatenated together, and
the phylogeny was inferred using the same methods as the single protein phylogenies mentioned
above.

M16 protease phylogeny

The MPP candidate proteins identified in the skoliomonads were added to the previously
constructed alignments of M16 proteases (Garrido et al., 2022), which include SPP, PreP, MPP-
alpha and MPP-beta sequences. In addition, we used the protein sequences of the two MPP

candidates of *Skoliomonas litria* to BLAST search nr, and added the top 20 hits for each
candidate to the alignments. This dataset was aligned using mafft einsi with default settings and
was trimmed by trimal as above. After trimming, we removed the duplicated protease domains in
four domain proteins, relative to their position when aligned to two domain proteins. This
alignment was then inferred using the same methods as the single protein phylogenies described
above. The domain structures displayed to the right of the phylogeny represents the predicted
protein domains of the skoliomonads (in red) and other major clades within the tree (in grey) as
predicted by the InterProScan online tool (see https://www.ebi.ac.uk/interpro/), which utilizes the
InterPro 98.0 database (Paysan-Lafosse et al., 2023).

Data availability

Genomes and gene annotations (where applicable) can be found under NBCI BioProject ID
PRJNA949547.

Acknowledgments

The majority of work was supported by Foundation grant (FRN-142349) from the Canadian
Institutes for Health Research (CIHR) awarded to Andrew J. Roger. Shelby K. Williams was
supported by a graduate scholarship from the Killam Foundation.

Citations

Anisimova, M., Gil, M., Dufayard, J. F., Dessimoz, C., & Gascuel, O. (2011). Survey of branch support methods demonstrates accuracy, power, and robustness of fast likelihood-based approximation schemes. *Systematic Biology*, *60*(5), 685–699. https://doi.org/10.1093/sysbio/syr041

Anwar, S., Dikhit, M. R., Singh, K. P., Kar, R. K., Zaidi, A., Sahoo, G. C., Roy, A. K., Nozaki, T., Das, P., & Ali, V. (2014). Interaction between Nbp35 and Cfd1 proteins of cytosolic Fe-S cluster assembly reveals a stable complex formation in Entamoeba histolytica. *PLoS ONE*, *9*(10), 12–14. https://doi.org/10.1371/journal.pone.0108971

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics*, *30*(15), 2114–2120. https://doi.org/10.1093/bioinformatics/btu170

Brown, M. W., Heiss, A. A., Kamikawa, R., Inagaki, Y., Yabuki, A., Tice, A. K., Shiratori, T., Ishida, K. I., Hashimoto, T., Simpson, A. G. B., & Roger, A. J. (2018). Phylogenomics Places Orphan Protistan Lineages in a Novel Eukaryotic Super-Group. *Genome Biology and Evolution*, *10*(2), 427–433. https://doi.org/10.1093/gbe/evy014

Capella-Gutiérrez, S., Silla-Martínez, J. M., & Gabaldón, T. (2009). trimAl: A tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*, *25*(15), 1972–1973. https://doi.org/10.1093/bioinformatics/btp348

Cheng, C. Y., Krishnakumar, V., Chan, A. P., Thibaud-Nissen, F., Schobel, S., & Town, C. D. (2017). Araport11: a complete reannotation of the Arabidopsis thaliana reference genome. *Plant Journal*, *89*(4), 789–804. https://doi.org/10.1111/tpj.13415

Criscuolo, A., & Gribaldo, S. (2010). BMGE (Block Mapping and Gathering with Entropy): A new software for selection of phylogenetic informative regions from multiple sequence alignments. *BMC Evolutionary Biology*, *10*(1). https://doi.org/10.1186/1471-2148-10-210

Derelle, E., Ferraz, C., Rombauts, S., Rouzé, P., Worden, A. Z., Robbens, S., Partensky, F., Degroeve, S., Echeynié, S., Cooke, R., Saeys, Y., Wuyts, J., Jabbari, K., Bowler, C., Panaud, O., Piégu, B., Ball, S. G., Ral, J. P., Bouget, F. Y., … Moreau, H. (2006). Genome analysis of the smallest free-living eukaryote Ostreococcus tauri unveils many unique features. *Proceedings of the National Academy of Sciences of the United States of America*, *103*(31), 11647–11652. https://doi.org/10.1073/pnas.0604795103

Dyall, S. D., Yan, W., Delgadillo-Correa, M. G., Lunceford, A., Loo, J. A., Clarke, C. F., & Johnson, P. J. (2004). Non-mitochondrial complex I proteins in a hydrogenosomal oxidoreductase complex. *Nature*, *431*(7012), 1103–1107. https://doi.org/10.1038/nature02990

Eglit, Y., Williams, S. K., Roger, A. J., & Simpson, A. G. B. (2024). *Characterisation of Skoliomonas gen. nov., a haloalkaliphilic anaerobe related to barthelonids (Metamonada)*. https://doi.org/10.1101/2024.03.12.584707

Eisen, J. A., Coyne, R. S., Wu, M., Wu, D., Thiagarajan, M., Wortman, J. R., Badger, J. H., Ren, Q., Amedeo, P., Jones, K. M., Tallon, L. J., Delcher, A. L., Salzberg, S. L., Silva, J. C.,

Haas, B. J., Majoros, W. H., Farzad, M., Carlton, J. M., Smith, R. K., … Orias, E. (2006). Macronuclear genome sequence of the ciliate Tetrahymena thermophila, a model eukaryote. *PLoS Biology*, *4*(9), 1620–1642. https://doi.org/10.1371/journal.pbio.0040286

Engel, S. R., Dietrich, F. S., Fisk, D. G., Binkley, G., Balakrishnan, R., Costanzo, M. C., Dwight, S. S., Hitz, B. C., Karra, K., Nash, R. S., Weng, S., Wong, E. D., Lloyd, P., Skrzypek, M. S., Miyasato, S. R., Simison, M., & Cherry, J. M. (2014). The Reference Genome Sequence of Saccharomyces cerevisiae: Then and Now. *G3: Genes, Genomes, Genetics*, *4*(3), 389–398. https://doi.org/10.1534/g3.113.008995

Eren, A. M., Kiefl, E., Shaiber, A., Veseli, I., Miller, S. E., Schechter, M. S., Fink, I., Pan, J. N., Yousef, M., Fogarty, E. C., Trigodet, F., Watson, A. R., Esen, Ö. C., Moore, R. M., Clayssen, Q., Lee, M. D., Kivenson, V., Graham, E. D., Merrill, B. D., … Willis, A. D. (2021). Community-led, integrated, reproducible multi-omics with anvi'o. *Nature Microbiology*, *6*(1), 3–6. https://doi.org/10.1038/s41564-020-00834-3

Flynn, J. M., Hubley, R., Goubert, C., Rosen, J., Clark, A. G., Feschotte, C., & Smit, A. F. (2020). RepeatModeler2 for automated genomic discovery of transposable element families. *Proceedings of the National Academy of Sciences of the United States of America*, *117*(17), 9451–9457. https://doi.org/10.1073/pnas.1921046117

Fulnečková, J., Ševčíková, T., Fajkus, J., Lukešová, A., Lukeš, M., Vlček, Č., Lang, B. F., Kim, E., Eliáš, M., & Sýkorová, E. (2013). A broad phylogenetic survey unveils the diversity and evolution of telomeres in eukaryotes. *Genome Biology and Evolution*, *5*(3), 468–483. https://doi.org/10.1093/gbe/evt019

Garcia, P. S., D'Angelo, F., Ollagnier de Choudens, S., Dussouchaud, M., Bouveret, E., Gribaldo, S., & Barras, F. (2022). An early origin of iron–sulfur cluster biosynthesis machineries before Earth oxygenation. *Nature Ecology and Evolution*, *6*(10), 1564–1572. https://doi.org/10.1038/s41559-022-01857-1

Garg, S., Stölting, J., Zimorski, V., Rada, P., Tachezy, J., Martin, W. F., & Gould, S. B. (2015). Conservation of transit peptide-Independent protein import into the mitochondrial and hydrogenosomal matrix. *Genome Biology and Evolution*, *7*(9), 2716–2726. https://doi.org/10.1093/gbe/evv175

Garrido, C., Wollman, F. A., & Lafontaine, I. (2022). The Evolutionary History of Peptidases Involved in the Processing of Organelle-Targeting Peptides. *Genome Biology and Evolution*, *14*(7), 1–18. https://doi.org/10.1093/gbe/evac101

Gawryluk, R. M. R., Chisholm, K. A., Pinto, D. M., & Gray, M. W. (2014). Compositional complexity of the mitochondrial proteome of a unicellular eukaryote (Acanthamoeba castellanii, supergroup Amoebozoa) rivals that of animals, fungi, and plants. *Journal of Proteomics*, *109*, 400–416. https://doi.org/10.1016/j.jprot.2014.07.005

Gawryluk, R. M. R., & Stairs, C. W. (2021). Diversity of electron transport chains in anaerobic protists. *Biochimica et Biophysica Acta - Bioenergetics*, *1862*(1), 148334. https://doi.org/10.1016/j.bbabio.2020.148334

Gigeroff, A. S., Eglit, Y., & Simpson, A. G. B. (2023). Characterisation and Cultivation of New Lineages of Colponemids, a Critical Assemblage for Inferring Alveolate Evolution. *Protist*,

174(2), 125949. https://doi.org/10.1016/j.protis.2023.125949

Giovannoni, S. J., Cameron Thrash, J., & Temperton, B. (2014). Implications of streamlining theory for microbial ecology. *ISME Journal*, *8*(8), 1553–1565. https://doi.org/10.1038/ismej.2014.60

Goffeau, A. A., Barrell, B. G., Bussey, H., Davis, R. W., Dujon, B., Feldmann, H., Hoheisel, J. D., Jacq, C., Johnston, M., Louis, E. J., Mewes, H. W., Murakami, Y., Philippsen, P., Tettelin, H., & Oliver, S. G. (1996). Life with 6000 Genes. *Science*, *274*(5287). 10.1126/science.274.5287.546

Gray, M. W., Burger, G., Derelle, R., Klimes, V., Leger, M. M., Sarrasin, M., Vlcek, C., Roger, A. J., Elias, M., & Franz Lang, B. (2019). The draft nuclear genome sequence and predicted mitochondrial proteome of Andalucia godoyi, a protist with the most gene-rich and bacteria-like mitochondrial genome. *Current Biology*, *3*(20), 1–35. doi.org/10.1186/s12915-020-0741-6

Guindon, S., Dufayard, J. F., Lefort, V., Anisimova, M., Hordijk, W., & Gascuel, O. (2010). New algorithms and methods to estimate maximum-likelihood phylogenies: Assessing the performance of PhyML 3.0. *Systematic Biology*, *59*(3), 307–321. https://doi.org/10.1093/sysbio/syq010

Haas, B. J., Delcher, A. L., Mount, S. M., Wortman, J. R., Smith, R. K., Hannick, L. I., Maiti, R., Ronning, C. M., Rusch, D. B., Town, C. D., Salzberg, S. L., & White, O. (2003). Improving the Arabidopsis genome annotation using maximal transcript alignment assemblies. *Nucleic Acids Research*, *31*(19), 5654–5666. https://doi.org/10.1093/nar/gkg770

Haas, B. J., Papanicolaou, A., Yassour, M., Grabherr, M., Philip, D., Bowden, J., Couger, M. B., Eccles, D., Li, B., Macmanes, M. D., Ott, M., Orvis, J., Pochet, N., Strozzi, F., Weeks, N., Westerman, R., William, T., Dewey, C. N., Henschel, R., … Regev, A. (2013). De novo transcript sequence recostruction from RNA-Seq: reference generation and analysis with Trinity. In *Nature protocols* (Vol. 8, Issue 8). https://doi.org/doi.org/10.1038/nprot.2013.084

International Human Genome Sequencing Consortium. (2001). Initial sequencing and analysis of the human genome. *Nature*, *412*(6846), 565–566. https://doi.org/10.1038/35087627

Jedelský, P. L., Doležal, P., Rada, P., Pyrih, J., Šmíd, O., Hrdý, I., Šedinová, M., Marcinčiková, M., Voleman, L., Perry, A. J., Beltrán, N. C., Lithgow, T., & Tachezy, J. (2011). The minimal proteome in the reduced mitochondrion of the parasitic protist Giardia intestinalis. *PLoS ONE*, *6*(2), 15–21. https://doi.org/10.1371/journal.pone.0017285

Jerlström-Hultqvist, J., Einarsson, E., Xu, F., Hjort, K., Ek, B., Steinhauf, D., Hultenby, K., Bergquist, J., Andersson, J. O., & Svärd, S. G. (2013). Hydrogenosomes in the diplomonad Spironucleus salmonicida. *Nature Communications*, *4*, 1–9. https://doi.org/10.1038/ncomms3493

Jiménez-González, A., Xu, F., & Andersson, J. O. (2019). Lateral Acquisitions Repeatedly Remodel the Oxygen Detoxification Pathway in Diplomonads and Relatives. *Genome Biology and Evolution*, *11*(9), 2542–2556. https://doi.org/10.1093/gbe/evz188

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S. A. A., Ballard, A. J., Cowie, A., Romera-Paredes, B., Nikolov, S., Jain, R., Adler, J., … Hassabis, D. (2021). Highly accurate protein structure prediction with AlphaFold. *Nature*, *596*(7873), 583–589. https://doi.org/10.1038/s41586-021-03819-2

Kang, D. D., Froula, J., Egan, R., & Wang, Z. (2015). MetaBAT, an efficient tool for accurately reconstructing single genomes from complex microbial communities. *PeerJ*, *2015*(8), 1–15. https://doi.org/10.7717/peerj.1165

Kang, S., Tice, A. K., Spiegel, F. W., Silberman, J. D., Pánek, T., Cepicka, I., Kostka, M., Kosakyan, A., Alcântara, D. M. C., Roger, A. J., Shadwick, L. L., Smirnov, A., Kudryavtsev, A., Lahr, D. J. G., & Brown, M. W. (2017). Between a Pod and a Hard Test: The Deep Evolution of Amoebae. *Molecular Biology and Evolution*, *34*(9), 2258–2270. https://doi.org/10.1093/molbev/msx162

Karnkowska, A., Vacek, V., Zubáčová, Z., Treitli, S. C., Petrželková, R., Eme, L., Novák, L., Žárský, V., Barlow, L. D., Herman, E. K., Soukal, P., Hroudová, M., Doležal, P., Stairs, C. W., Roger, A. J., Eliáš, M., Dacks, J. B., Vlček, Č., & Hampl, V. (2016). A eukaryote without a mitochondrial organelle. *Current Biology*, *26*(1), 1274–1284. https://doi.org/10.1016/j.cub.2016.03.053

Kikuchi, G. (1973). The Glycine Cleavage System: Composition, Reaction Mechanism, And Physiological Significance. *Molecular and Cellular Biochemistry*, *21*(July), 1154–1157.

Kim, D., Paggi, J. M., Park, C., Bennett, C., & Salzberg, S. L. (2019). Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology*, *37*(August). https://doi.org/10.1038/s41587-019-0201-4

Kolmogorov, M., Yuan, J., Lin, Y., & Pevzner, P. A. (2019). Assembly of long, error-prone reads using repeat graphs. *Nature Biotechnology*, *37*(5), 540–546. https://doi.org/10.1038/s41587-019-0072-8

Lahti, C. J., D'Oliveira, C. E., & Johnson, P. J. (1992). β-Succinyl-coenzyme A synthetase from Trichomonas vaginalis is a soluble hydrogenosomal protein with an amino-terminal sequence that resembles mitochondrial presequences. *Journal of Bacteriology*, *174*(21), 6822–6830. https://doi.org/10.1128/jb.174.21.6822-6830.1992

Leger, M. M., Eme, L., Hug, L. A., & Roger, A. J. (2016). Novel Hydrogenosomes in the Microaerophilic Jakobid Stygiella incarcerata. *Molecular Biology and Evolution*, *33*(9), 2318–2336. https://doi.org/10.1093/molbev/msw103

Leger, M. M., Kolisko, M., Kamikawa, R., Stairs, C. W., Kume, K., Čepička, I., Silberman, J. D., Andersson, J. O., Xu, F., Yabuki, A., Eme, L., Zhang, Q., Takishita, K., Inagaki, Y., Simpson, A. G. B., Hashimoto, T., & Roger, A. J. (2017). Organelles that illuminate the origins of Trichomonas hydrogenosomes and Giardia mitosomes. *Nature Ecology and Evolution*, *1*(4), 1–15. https://doi.org/10.1038/s41559-017-0092

Lill, R., & Freibert, S. A. (2020). Mechanisms of Mitochondrial Iron-Sulfur Protein Biogenesis. *Annual Review of Biochemistry*, *89*, 471–499. https://doi.org/10.1146/annurev-biochem-013118-111540

Liu, Y., Sieprawska-Lupa, M., Whitman, W. B., & White, R. H. (2010). Cysteine is not the sulfur source for iron-sulfur cluster and methionine biosynthesis in the methanogenic archaeon Methanococcus maripaludis. *Journal of Biological Chemistry*, *285*(42), 31923–31929. https://doi.org/10.1074/jbc.M110.152447

Lomsadze, A., Burns, P. D., & Borodovsky, M. (2014). Integration of mapped RNA-Seq reads into automatic training of eukaryotic gene finding algorithm. *Nucleic Acids Research*, *42*(15), 1–8. https://doi.org/10.1093/nar/gku557

Manzano-Marín, A., & Latorre, A. (2016). Snapshots of a shrinking partner: Genome reduction in Serratia symbiotica. *Scientific Reports*, *6*(June), 1–11. https://doi.org/10.1038/srep32590

Matthey-Doret, C., Colp, M. J., Escoll, P., Thierry, A., Moreau, P., Curtis, B., Sahr, T., Sarrasin, M., Gray, M. W., Franz Lang, B., Archibald, J. M., Buchrieser, C., & Koszul, R. (2022). Chromosome-scale assemblies of Acanthamoeba castellanii genomes provide insights into Legionella pneumophila infection–related chromatin reorganization. *Genome Research*, *32*(9), 1698–1710. https://doi.org/10.1101/gr.276375.121

Mazurek, M. P., Prasad, P. D., Gopal, E., Fraser, S. P., Bolt, L., Rizaner, N., Palmer, C. P., Foster, C. S., Palmieri, F., Ganapathy, V., Stühmer, W., Djamgoz, M. B. A., & Mycielska, M. E. (2010). Molecular origin of plasma membrane citrate transporter in human prostate epithelial cells. *EMBO Reports*, *11*(6), 431–437. https://doi.org/10.1038/embor.2010.51

Mentel, M., Zimorski, V., Haferkamp, P., Martin, W., & Henze, K. (2008). Protein import into hydrogenosomes of Trichomonas vaginalis involves both N-terminal and internal targeting signals: A case study of thioredoxin reductases. *Eukaryotic Cell*, *7*(10), 1750–1757. https://doi.org/10.1128/EC.00206-08

Mirdita, M., Schütze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S., & Steinegger, M. (2022). ColabFold: making protein folding accessible to all. *Nature Methods*, *19*(6), 679–682. https://doi.org/10.1038/s41592-022-01488-1

Motyčková, A., Voleman, L., Najdrová, V., Arbonová, L., Benda, M., Dohnálek, V., Janowicz, N., Malych, R., Šuťák, R., Ettema, T. J. G., Svärd, S., Stairs, C. W., & Doležal, P. (2023). Adaptation of the late ISC pathway in the anaerobic mitochondrial organelles of Giardia intestinalis. *PLoS Pathogens*, *19*(10 October), 1–24. https://doi.org/10.1371/journal.ppat.1010773

Mukherjee, M., Brown, M. T., McArthur, A. G., & Johnson, P. J. (2006). Proteins of the glycine decarboxylase complex in the hydrogenosome of Trichomonas vaginalis. *Eukaryotic Cell*, *5*(12), 2062–2071. https://doi.org/10.1128/EC.00205-06

Müller, M., Mentel, M., van Hellemond, J. J., Henze, K., Woehle, C., Gould, S. B., Yu, R.-Y., van der Giezen, M., Tielens, A. G. M., & Martin, W. F. (2012). Biochemistry and Evolution of Anaerobic Energy Metabolism in Eukaryotes. *Microbiology and Molecular Biology Reviews*, *76*(2), 444–495. https://doi.org/10.1128/mmbr.05024-11

Nelson, R. J., Ziegelhoffer, T., Nicolet, C., Werner-Washburne, M., & Craig, E. A. (1992). The translation machinery and 70 kd heat shock protein cooperate in protein synthesis. *Cell*, *71*(1), 97–105. https://doi.org/10.1016/0092-8674(92)90269-I

Nývltová, E., Smutná, T., Tachezy, J., & Hrdý, I. (2016). OsmC and incomplete glycine
decarboxylase complex mediate reductive detoxification of peroxides in hydrogenosomes of
Trichomonas vaginalis. *Molecular and Biochemical Parasitology*, *206*(1–2), 29–38.
https://doi.org/10.1016/j.molbiopara.2016.01.006

Nývltová, E., Stairs, C. W., Hrdý, I., Rídl, J., Mach, J., Pa□es, J., Roger, A. J., & Tachezy, J.
(2015). Lateral gene transfer and gene duplication played a key role in the evolution of
mastigamoeba balamuthi hydrogenosomes. *Molecular Biology and Evolution*, *32*(4), 1039–
1055. https://doi.org/10.1093/molbev/msu408

Ohnishi, T. (1998). Iron-sulfur clusters/semiquinones in Complex I. *Biochimica et Biophysica*
Acta - Bioenergetics, *1364*(2), 186–206. https://doi.org/10.1016/S0005-2728(98)00027-9

Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P., & Tyson, G. W. (2015). CheckM:
Assessing the quality of microbial genomes recovered from isolates, single cells, and
metagenomes. *Genome Research*, *25*(7), 1043–1055. https://doi.org/10.1101/gr.186072.114

Paysan-Lafosse, T., Blum, M., Chuguransky, S., Grego, T., Pinto, B. L., Salazar, G. A., Bileschi,
M. L., Bork, P., Bridge, A., Colwell, L., Gough, J., Haft, D. H., Letunić, I., Marchler-Bauer,
A., Mi, H., Natale, D. A., Orengo, C. A., Pandurangan, A. P., Rivoire, C., … Bateman, A.
(2023). InterPro in 2022. *Nucleic Acids Research*, *51*(D1), D418–D427.
https://doi.org/10.1093/nar/gkac993

Prjibelski, A., Antipov, D., Meleshko, D., Lapidus, A., & Korobeynikov, A. (2020). Using
SPAdes De Novo Assembler. *Current Protocols in Bioinformatics*, *70*(1), 1–29.
https://doi.org/10.1002/cpbi.102

Pyrih, J., Pyrihová, E., Kolísko, M., Stojanovová, D., Basu, S., Harant, K., Haindrich, A. C.,
Doležal, P., Lukeš, J., Roger, A., & Tachezy, J. (2016). Minimal cytosolic iron-sulfur
cluster assembly machinery of Giardia intestinalis is partially associated with mitosomes.
Molecular Microbiology, *102*(4), 701–714. https://doi.org/10.1111/mmi.13487

Pyrihová, E., Motyčková, A., Voleman, L., Wandyszewska, N., Fišer, R., Seydlová, G., Roger,
A., Kolísko, M., & Doležal, P. (2018). A single tim translocase in themitosomes of giardia
intestinalis illustrates convergence of protein import machines in anaerobic eukaryotes.
Genome Biology and Evolution, *10*(10), 2813–2822. https://doi.org/10.1093/gbe/evy215

Rhie, A., Walenz, B. P., Koren, S., & Phillippy, A. M. (2020). Merqury: Reference-free quality,
completeness, and phasing assessment for genome assemblies. *Genome Biology*, *21*(1), 1–
27. https://doi.org/10.1186/s13059-020-02134-9

Roger, A. J., Muñoz-Gómez, S. A., & Kamikawa, R. (2017). The Origin and Diversification of
Mitochondria. In *Current Biology*. https://doi.org/10.1016/j.cub.2017.09.015

Salas-Leiva, D. E., Tromer, E. C., Curtis, B. A., Jerlström-hultqvist, J., Kolisko, M., Yi, Z.,
Salas-leiva, J. S., Gallot-lavallée, L., Williams, S. K., Kops, G. J. P. L., Archibald, J. M.,
Simpson, A. G. B., & Roger, A. J. (2021). Genomic analysis finds no evidence of canonical
eukaryotic DNA processing complexes in a free-living protist. *Nature Communications*, 1–
13. https://doi.org/10.1038/s41467-021-26077-2

Sánchez, L. B., Galperin, M. Y., & Müller, M. (2000). Acetyl-CoA synthetase from the

amitochondriate eukaryote Giardia lamblia belongs to the newly recognized superfamily of acyl-CoA synthetases (nucleoside diphosphate-forming). *Journal of Biological Chemistry*, *275*(8), 5794–5803. https://doi.org/10.1074/jbc.275.8.5794

Sanchez, L. B., & Müller, M. (1996). Purification and characterization of the acetate forming enzyme, acetyl-CoA synthetase (ADP-forming) from the amitochondriate protist, Giardia lamblia. *FEBS Letters*, *378*(3), 240–244. https://doi.org/10.1016/0014-5793(95)01463-2

Schneider, R. E., Brown, M. T., Shiflett, A. M., Dyall, S. D., Hayes, R. D., Xie, Y., Loo, J. A., & Johnson, P. J. (2011). The Trichomonas vaginalis hydrogenosome proteome is highly reduced relative to mitochondria, yet complex compared with mitosomes. *International Journal for Parasitology*, *41*(13–14), 1421–1434. https://doi.org/10.1016/j.ijpara.2011.10.001

Schönknecht, G., Chen, W. H., Ternes, C. M., Barbier, G. G., Shrestha, R. P., Stanke, M., Bräutigam, A., Baker, B. J., Banfield, J. F., Garavito, R. M., Carr, K., Wilkerson, C., Rensing, S. A., Gagneul, D., Dickenson, N. E., Oesterhelt, C., Lercher, M. J., & Weber, A. P. M. (2013). Gene transfer from bacteria and archaea facilitated evolution of an extremophilic eukaryote. *Science*, *339*(6124), 1207–1210. https://doi.org/10.1126/science.1231707

Simao, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V, & Zdobnov, E. M. (2015). Genome analysis BUSCO□: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics*, *31*(June), 3210–3212. https://doi.org/10.1093/bioinformatics/btv351

Simpson, A. G. B. (2003). Cytoskeletal organization, phylogenetic affinities and systematics in the contentious taxon Excavata (Eukaryota). *International Journal of Systematic and Evolutionary Microbiology*, *53*(6), 1759–1777. https://doi.org/10.1099/ijs.0.02578-0

Smith, D. G. S., Gawryluk, R. M. R., Spencer, D. F., Pearlman, R. E., Siu, K. W. M., & Gray, M. W. (2007). Exploring the Mitochondrial Proteome of the Ciliate Protozoon Tetrahymena thermophila: Direct Analysis by Tandem Mass Spectrometry. *Journal of Molecular Biology*, *374*(3), 837–863. https://doi.org/10.1016/j.jmb.2007.09.051

Stairs, C. W., Eme, L., Brown, M. W., Mutsaers, C., Susko, E., Dellaire, G., Soanes, D. M., Van Der Giezen, M., & Roger, A. J. (2014). A SUF Fe-S cluster biogenesis system in the mitochondrion-related organelles of the anaerobic protist Pygsuia. *Current Biology*, *24*(11), 1176–1186. https://doi.org/10.1016/j.cub.2014.04.033

Stairs, C. W., Eme, L., Muñoz-Gómez, S. A., Cohen, A., Dellaire, G., Shepherd, J. N., Fawcett, J. P., & Roger, A. J. (2018). Microbial eukaryotes have adapted to hypoxia by horizontal acquisitions of a gene involved in rhodoquinone biosynthesis. *ELife*, *7*, 1–23. https://doi.org/10.7554/eLife.34292

Stairs, C. W., Leger, M. M., & Roger, A. J. (2015). Diversity and origins of anaerobic metabolism in mitochondria and related organelles. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *370*(1678). https://doi.org/10.1098/rstb.2014.0326

Stairs, C. W., Roger, A. J., & Hampl, V. (2011). Eukaryotic pyruvate formate lyase and its activating enzyme were acquired laterally from a firmicute. *Molecular Biology and*

Evolution, *28*(7), 2087–2099. https://doi.org/10.1093/molbev/msr032

Stairs, C. W., Táborský, P., Salomaki, E. D., Kolisko, M., Pánek, T., Eme, L., Hradilová, M.,
Vlček, Č., Jerlström-Hultqvist, J., Roger, A. J., & Čepička, I. (2021). Anaeramoebae are a
divergent lineage of eukaryotes that shed light on the transition from anaerobic
mitochondria to hydrogenosomes. *Current Biology*, *31*(24), 5605-5612.e5.
https://doi.org/10.1016/j.cub.2021.10.010

Stanke, M., Steinkamp, R., Waack, S., & Morgenstern, B. (2004). AUGUSTUS: A web server
for gene finding in eukaryotes. *Nucleic Acids Research*, *32*(WEB SERVER ISS.), 309–312.
https://doi.org/10.1093/nar/gkh379

Takeda, H., Tsutsumi, A., Nishizawa, T., Lindau, C., Busto, J. V., Wenz, L. S., Ellenrieder, L.,
Imai, K., Straub, S. P., Mossmann, W., Qiu, J., Yamamori, Y., Tomii, K., Suzuki, J.,
Murata, T., Ogasawara, S., Nureki, O., Becker, T., Pfanner, N., … Endo, T. (2021).
Mitochondrial sorting and assembly machinery operates by β-barrel switching. In *Nature*
(Vol. 590, Issue 7844). Springer US. https://doi.org/10.1038/s41586-020-03113-7

Tanifuji, G., Takabayashi, S., Kume, K., Takagi, M., Nakayama, T., Kamikawa, R., Inagaki, Y.,
& Hashimoto, T. (2018). The draft genome of Kipferlia bialata reveals reductive genome
evolution in fornicate parasites. *PLoS ONE*, *13*(3), 1–18.
https://doi.org/10.1371/journal.pone.0194487

The Arabidopsis Genome Iniative. (2000). Analysis of the genome sequence of the Flowering
plant Arabidopsis thaliana. *Nature*, *408*, 796–815.
https://doi.org/10.1134/S1022795411020074

Trapnell, C., Williams, B. A., Pertea, G., Mortazavi, A., Kwan, G., Van Baren, M. J., Salzberg,
S. L., Wold, B. J., & Pachter, L. (2010). Transcript assembly and quantification by RNA-
Seq reveals unannotated transcripts and isoform switching during cell differentiation.
Nature Biotechnology, *28*(5), 511–515. https://doi.org/10.1038/nbt.1621

Tsaousis, A. D., De Choudens, S. O., Gentekaki, E., Long, S., Gaston, D., Stechmann, A.,
Vinella, D., Py, B., Fontecave, M., Barras, F., Lukeš, J., & Roger, A. J. (2012). Evolution of
Fe/S cluster biogenesis in the anaerobic parasite Blastocystis. *Proceedings of the National*
Academy of Sciences of the United States of America, *109*(26), 10426–10431.
https://doi.org/10.1073/pnas.1116067109

Tsaousis, A. D., Gentekaki, E., Eme, L., Gaston, D., & Roger, A. J. (2014). Evolution of the
cytosolic iron-sulfur cluster assembly machinery in Blastocystis species and other microbial
eukaryotes. *Eukaryotic Cell*, *13*(1), 143–153. https://doi.org/10.1128/EC.00158-13

Vacek, V., Novák, L. V. F., Treitli, S. C., Táborský, P., Čepička, I., Kolísko, M., Keeling, P. J.,
& Hampl, V. (2018). Fe-S cluster assembly in oxymonads and related protists. *Molecular*
Biology and Evolution, *35*(11), 2712–2718. https://doi.org/10.1093/molbev/msy168

Van Grinsven, K. W. A., Rosnowsky, S., Van Weelden, S. W. H., Pütz, S., Van Der Giezen, M.,
Martin, W., Van Hellemond, J. J., Tielens, A. G. M., & Henze, K. (2008). Acetate:succinate
CoA-transferase in the hydrogenosomes of Trichomonas vaginalis: Identification and
characterization. *Journal of Biological Chemistry*, *283*(3), 1411–1418.
https://doi.org/10.1074/jbc.M702528200

Vaser, R., & Šikić, M. (2021). Time- and memory-efficient genome assembly with Raven.
Nature Computational Science, *1*(5), 332–336. https://doi.org/10.1038/s43588-021-00073-4

Wick, R. R., Judd, L. M., & Holt, K. E. (2019). Performance of neural network basecalling tools
for Oxford Nanopore sequencing. *Genome Biology*, *20*(1), 1–10.
https://doi.org/10.1186/s13059-019-1727-y

Xu, F., Jex, A., & Svärd, S. G. (2020). A chromosome-scale reference genome for Giardia
intestinalis WB. *Scientific Data*, *7*(1), 1–8. https://doi.org/10.1038/s41597-020-0377-y

Yazaki, E., Kume, K., Shiratori, T., Eglit, Y., Tanifuji, G., Harada, R., Simpson, A., Ishida, K.,
Hashimoto, T., & Inagaki, Y. (2020). Barthelonids represent a deep-branching Metamonad
clade with mitochondrion-related organelles generating no ATP. *Proc. R. Soc.*
https://doi.org/10.1101/805762

Yubuki, N., Galindo, L. J., Reboul, G., López-García, P., Brown, M. W., Pollet, N., & Moreira,
D. (2020). Ancient Adaptive Lateral Gene Transfers in the Symbiotic Opalina-Blastocystis
Stramenopile Lineage. *Molecular Biology and Evolution*, *37*(3), 651–659.
https://doi.org/10.1093/molbev/msz250

Záhonová, K., Treitli, S. C., Le, T., Škodová-Sveráková, I., Hanousková, P., Čepička, I.,
Tachezy, J., & Hampl, V. (2022). Anaerobic derivates of mitochondria and peroxisomes in
the free-living amoeba Pelomyxa schiedti revealed by single-cell genomics. *BMC Biology*,
20(1), 1–16. https://doi.org/10.1186/s12915-022-01247-w

Žárský, V., Klimeš, V., Pačes, J., Vlček, Č., Hradilová, M., Beneš, V., Nývltová, E., Hrdý, I.,
Pyrih, J., Mach, J., Barlow, L., Stairs, C. W., Eme, L., Hall, N., Eliáš, M., Dacks, J. B.,
Roger, A., & Tachezy, J. (2021). The Mastigamoeba balamuthi Genome and the Nature of
the Free-Living Ancestor of Entamoeba. *Molecular Biology and Evolution*, *38*(6), 2240–
2259. https://doi.org/10.1093/molbev/msab020

Zhao, D., Salas-leiva, D. E., Williams, S. K., & Dunn, K. A. (2023). Eukfinder□: a pipeline to
retrieve microbial eukaryote genomes from metagenomic sequencing data. *BioRxiv*.
https://doi.org/https://doi.org/10.1101/2023.12.28.573569

Zítek, J., Füssy, Z., Treitli, S. C., Peña-Diaz, P., Vaitová, Z., Zavadska, D., Harant, K., & Hampl,
V. (2022). Reduced mitochondria provide an essential function for the cytosolic methionine
cycle. *Current Biology*, 5057–5068. https://doi.org/10.1016/j.cub.2022.10.028

Zubáčová, Z., Novák, L., Bublíková, J., Vacek, V., Fousek, J., Rídl, J., Tachezy, J., Doležal, P.,
Vlček, Č., & Hampl, V. (2013). The Mitochondrion-Like Organelle of Trimastix pyriformis
Contains the Complete Glycine Cleavage System. *PLoS ONE*, *8*(3), 1–9.
https://doi.org/10.1371/journal.pone.0055417

1119

1120

1121 **Tables & Figures**

1122 Table 1. Summary statistics for the genomes, transcriptomes, and predicted proteomes of the

1123 BaSk clade.

	Skoliomonas litria	*Skoliomonas* sp. GEMRC	*Skoliomonas* sp. RCL	*Barthelona* sp. PCE	*Barthelona* sp. PAP020
*Genome Coverage (Nanopore)	41x	119x	34x	66x	Illumina only
*Genome Coverage (Illumina)	206x	116x	93x	308x	29x
# of Illumina Reads Generated	53.7 M	26.8 M	85.9 M	75.4 M	70.0 M
% RNAseq reads mapping	99.4	99.9	99.6	99.3	-
Merqury Completeness (%)	98.7	98.6	98.4	98.4	-
Genome Size	31.9 Mbp	33.5 Mbp	21.1 Mbp	10.2 Mbp	13.2 Mbp
Contigs (#)	16	21	18	8	1,035
N50	3,724,615	2,551,763	2,069,101	1,587,787	24,218
L50	4	6	4	3	156
%GC	35.8	36.9	36.0	40.3	30.5
%Repetitive	55.4	48.0	28.1	4.46	-
Predicted Proteins (#)	15,224	14,268	9,198	5,064	6,305

1124   *Median contig coverage

1125

1126

1127

1128   Table 2. BUSCO scores for members of the BaSk clade.

	Complete Single Copy	Complete Duplicated	Total Complete	Fragmented	Missing
S. litria - Genome	178	4	182 (60.1%)	15 (4.9%)	106 (35.0%)
S. litria - Prediction	215	9	224 (73.9%)	7 (2.3%)	72 (23.8%)
S. litria - Transcriptome	190	24	214 (70.6%)	16 (5.3%)	73 (24.1%)
S. litria – Curated Predictions	208	17	225 (74.3%)	7 (2.3%)	71 (23.4%)
S. sp. GEMRC - Genome	168	9	177 (58.4%)	27 (8.9%)	99 (32.7%)
S. sp. GEMRC - Prediction	183	14	197 (65.0%)	29 (9.6%)	77 (25.4%)
S. sp. GEMRC - Transcriptome	139	77	216 (71.3%)	9 (3.0%)	78 (25.7%)
S. sp. RCL - Genome	182	4	186 (61.4%)	16 (5.3%)	101 (33.3%)
S. sp. RCL - Prediction	212	10	222 (73.3%)	14 (4.6%)	67 (22.1%)
S. sp. RCL - Transcriptome	193	27	220 (72.6%)	14 (4.6%)	69 (22.8%)
B. sp. PCE - Genome	142	3	145 (47.9%)	15 (4.9%)	143 (47.2%)
B. sp. PCE – Prediction	161	7	168 (55.4%)	25 (8.3%)	110 (36.3%)
B. sp. PCE - Transcriptome	5	157	162 (53.5%)	31 (10.2%)	110 (36.3%)

1129

1130

1131

1132

1133

1134

1135

1136

1137

1138

Figure 1. 'BaSk' is a clade of anaerobic protists that branches as sister to all known Fornicata
within the Metamonada. Panels A-D) Differential interference contrast light micrographs of
skoliomonad lineages *Skoliomonas litria* (A), *Skoliomonas* sp. GEMRC (B), and *Skoliomonas* sp.
RCL (C), as well as *Barthelona* sp. PCE (D) showing overall morphology (imaging and methods
described in Eglit et al. 2024). Scale bars indicate a length of 10 μm. (E) A maximum likelihood
(ML) phylogeny based on 174 concatenated aligned proteins encompassing 46,113 sites. The
depicted topology was estimated using IQ-TREE under the LG+C60+F+Γ model and was used as
a guide tree for estimating the LG+PMSF(C60)+F+Γ model. Support under the latter model was
evaluated by SH-aLRT support percentage, aBayes support and nonparametric bootstrap
percentage based on 100 nonparametric bootstrap replicates. The scale bar indicates the
expected number of amino acid substitutions per site under the PMSF model. When at least
one support value was less than maximal, all three are shown on the branch in the order SH-
aLRT support percentage/aBayes/nonparametric bootstrap percentage. A black dot on the
branch indicates all three approaches yielded full (i.e. 100/1/100) support. The BaSk clade is
highlighted in red.

1154

1155

Figure 2. Presence, absence and predicted localization of key MRO and energy metabolism enzymes in 'BaSks' and other metamonads. A schematic phylogeny of the Metamonada is shown on the left based on Fig. 1, with the BaSk clade highlighted in red. The pathways depicted include anaerobic extended glycolysis/energy metabolism enzymes, conserved MRO proteins, MRO protein targeting machinery components, and the SmsCB fusion protein. Coloured versus grey circles indicate presence/absence of the protein from predicted proteomes and/or transcriptome data and the colours indicate probable localization. Circles split in half indicate that more than one paralog was present with different predicted localizations. Protein abbreviations: PFO - Pyruvate:ferredoxin oxidoreductase; NuoE/F - Respiratory-chain NADH dehydrogenase 24/51 kDa subunit; HydA – Iron-only hydrogenase; ASCT1B – Acetate:succinyl-CoA transferase B; ASCT1C - Acetate:succinyl-CoA transferase C; SCS - Succinyl-CoA synthase; ACS - Acetyl-CoA synthase; Fdx – Ferredoxin; L – Glycine cleavage system subunit L; H - Glycine cleavage system subunit H; P - Glycine cleavage system subunit P; T - Glycine cleavage system subunit T; SHMT - Serine hydroxymethyltransferaseTom40 - Translocase of outer mitochondrial membrane 40; Tom22 - Translocase of outer mitochondrial membrane 22; Sam50 – Sorting and assembly machinery 50; Tim17/22/23 - Translocase of the inner membrane 17/22/23; Tim44 - Translocase of the inner membrane 44; Pam16 - Presequence translocase-associated motor 16; Pam18 - Presequence translocase-associated motor 18; IscS - Iron-sulfur cluster assembly enzyme cysteine desulfurase; IscU - Iron-sulfur cluster assembly enzyme scaffold; SufCB - Sulfur formation CB fusion protein.

1200 **Figure 3. A phylogeny of the SmsCB fusion protein in a variety of anaerobic eukaryotes (red)**
1201 **and their closest bacterial and archaeal homologs.** The alignment contains the SmsCB fusion
1202 protein from the eukaryotes and concatenated SmsC and SmsB proteins from the prokaryotes.
1203 This ML tree was estimated under the LG+C60+F+Γ model using IQ-TREE. Ultrafast bootstrap
1204 values are displayed on the branches, with dots indicating UFBOOT value over 95%.

1205

1206

Figure 4. Gain and loss of anaerobic enzymes and MRO proteins over the tree of Metamonada. Gains (red) and losses (blue) of key metabolic pathway components discussed throughout this paper are mapped on a schematic phylogeny of Metamonada (based on Figure 1). Protein abbreviations are: HydG/E/F – Iron hydrogenase maturase G/E/F; ACS - Acetyl-CoA synthase; NuoE/F - Respiratory-chain NADH dehydrogenase 24/51 kDa subunit; GCS – Glycine cleavage system; SCS - Succinyl-CoA synthase;IscS/U - Iron-sulfur cluster assembly enzyme system. The *Spironucleus* ACS is denoted as ACS* as an ACS "type" could not be confidently assigned.

1214

1215   Supplementary tables 1-3: Available for download from Dryad

1216   https://datadryad.org/stash/share/SNhoYuY19An041SeN194bwM_K3WIqqOO4TwltqK1mn0

1217

1218

1219   Supplementary figures:

1220

1221   Supp. Figure 1. **A phylogeny of the SmsC protein from eukaryotes, bacteria, and archaea.**
1222   The ML tree was estimated using IQ-TREE under the LG+C60+F+Γ model of evolution.
1223   Ultrafast bootstrap values are displayed on the branches. For eukaryotic homologs, the N-
1224   terminal portion of the protein of the eukaryote SmsCB fusion that aligned with prokaryotic
1225   SmsC was used in the alignment. BaSk sequences are bolded and highlighted in red.

1226

Supp. Figure 2. **A phylogeny of SmsBD protein from eukaryotes, bacteria, and archaea,** The
ML tree was estimated using IQ-TREE under the LG+C20+F+Γ model of evolution. Ultrafast
bootstrap values are displayed on the branches. For eukaryotic homologs, the C-terminal portion
of the protein of the eukaryote SmsCB fusion that aligned with prokaryotic SmsB was used in
the alignment. BaSk sequences are bolded and highlighted in red.

1232

Supp. Figure 3. **Phylogenetic relationships amongst several different types of cysteine desulfurase enzymes and several enzymes that share structural similarities with them.** The ML tree was estimated using IQ-TREE with the LG+C20+F+Γ evolutionary model. Ultrafast bootstrap values are displayed on the branches. Several families of related proteins have been collapsed into wedges. BaSk sequences are bolded and highlighted in red. Abbreviations are: MCS – Molybdenum cofactor sulfurase; SmsS – SUF-like minimal system; CsdA - Cysteine sulfinate desulfinase A; MisS - Minimal iron-sulfur system; IscS - Iron-sulfur cluster assembly enzyme; SelD – Selenide water dikinase

Supp. Figure 4. **Phylogenetic relationships amongst chaperonin 60 (cpn60) homologs in eukaryotes.** These ML trees were estimated using IQ-TREE with the LG+C20+F+Γ evolutionary model. Ultrafast bootstrap values are displayed on the branches. BaSk sequences are bolded and highlighted in red. (A) Both cytosolic (purple) and mitochondrial (orange) homologs are included in this tree. The major groups of proteins are displayed as wedges. (B) A reconstruction of the mitochondrial (purple), bacterial (orange), and choroplasatic (green) Cpn60 subunit GroEL.

1272

Supp. Figure 5. **Phylogenetic relationships amongst heat shock protein 70 (hsp70) paralogs in eukaryotes and prokaryotes.** (A) The ML tree was estimated using IQ-TREE with the LG+C20+F+Γ evolutionary model. Ultrafast bootstrap values are displayed on the branches. Cytoplasmic and ER (purple) and mitochondrial (green) homologs are labeled according to indel mapping patterns in the alignment. BaSk sequences are bolded and highlighted in red. (B) An example of indel patterns within the alignment used to assign probable localizations of hsp70 homologs in the tree. Amino acid residues are colour coded.

1280

1281

Supp. Figure 6. **Estimated phylogenetic relationships amongst a variety of M16**
metallopeptidases. This tree was estimated using the M16 peptidase database previously
constructed by Garrido and colleagues (2022) and was estimated using IQ-TREE under the
LG+C20+F+Γ evolutionary model. Ultrafast bootstrap values are displayed on the branches.
Major groups of proteins have been collapsed into wedges. To the right of the tree, the domain
structure of each protein/protein group is displayed, as annotated by InterPro. Eukaryotes – purple;
bacteria – green; skoliomonads – red.

Supp. Figure 7. **Estimated phylogenetic relationships between members of the mitochondrial carrier family (MCP) with candidate MCP sequences from 'BaSks'.** Where possible, MCP sequences are labeled with their family subtype designation according to the NCBI nr database. This phylogenetic tree was constructed using IQ-TREE and the LG+C20+F+Γ model of evolution. The *Skoliomonas litria* candidate MCP is highlighted in red. Values on the branches represent ultrafast bootstrap values.

1306

1307

Supp. Figure 8. **The phylogeny of Sam50/BamA family in eukaryotes and prokaryotes and predicted structures of the BaSk homolog.** (A) A phylogenetic reconstruction of the Sam50 candidate proteins in *Skoliomonas litria* and *Skoliomonas* sp. GEMRC, and their relationship to Sam50 (purple) and BamA (green) homolog proteins in eukaryotes and prokaryotes. (B) Alphafold2 structure prediction using the sequence of the candidate Sam50 protein found in *Skoliomonas litria* (C) Alphafold2 structure prediction using the sequence of the candidate Sam50 protein found in *Skoliomonas* sp. GEMRC. The protein regions resembling "POTRA" domains in other Sam50 proteins are highlighted with a red circle in each candidate.

1316

1317

1318   Supp. Figure 9. **A structural alignment of the β-barrel portions of the *Skoliomonas litria***
1319   **candidate Sam50 protein and Sam50 from yeast.** The *Skoliomonas litria* protein is shown in
1320   blue, aligned to the β-barrel domain of the yeast Sam50, which is shown in green. The missing
1321   sections of the β-barrel in *Skoliomonas litria*, when compared to the yeast model, is highlighted
1322   by a red circle. This structural alignment was produced using the PyMOL "cealign" function.

1323

1324

1325

1326

1327 Supp. Figure 10. **A schematic of the hypothesized ATP production pathway in the BaSk**
1328 **clade.** Hyd – Iron-only hydrogenase; ME – Malic enzyme; PFO – Pyruvate:ferredoxin oxidoreductase; ACS –
1329 Acetyl-CoA synthase; Fdx – Ferredoxin; PFL – pyruvate-formate lyase.

1330

1331

1332

Supp. Figure 11. **A phylogenetic estimation of the relationship amongst ACS protein**
sequences in a variety of prokaryotes and eukaryotes, including metamonads. ACS1 is
represented in orange, ACS2 in green, and the newly detected 'ACS3' in purple. BaSk sequences
are bolded and highlighted in red. Major groupings of each ortholog are collapsed into wedges.
This phylogeny was constructed using IQ-TREE and the LG+C20+F+Γ model of evolution.
Ultrafast bootstrap values are indicated on the branches.

1339