Impacts of global food supply on biodiversity via land use and climate change

Abstract

Land-use change is currently the greatest driver of biodiversity change, with climate change

predicted to match or surpass its impacts by mid-century. The global food system is a key driver of

both these anthropogenic pressures, thus the development of sustainable food systems will be

critical to halting and reversing biodiversity loss. Previous studies of the biodiversity footprint of

food tend to focus on land use alone. We use the multi-regional input-output model EXIOBASE to

estimate the impacts of biodiversity embedded within the global food system. We build on prior

analyses, calculating the impacts of both agricultural land-use and greenhouse gas (GHG) emission

footprints for the same two metrics of biodiversity: local species richness and rarity-weighted

species richness. Our biodiversity models capture regional variation in the sensitivity of biodiversity

both to land-use differences and to climate change. We find that the footprint of land area does not

capture the biodiversity impact embedded within trade that is provided by our metric of land-driven

species richness change, and that our metric of rarity-weighted richness places a greater emphasis

on the biodiversity costs in Central and South America. We show that the climate contributions of

food production are accumulating rapidly over time, particularly in North America and Western

Europe. The measures we present are simple to calculate and could be incorporated into decision

making and environmental impact assessments by governments and businesses.

Keywords

1. Introduction

Anthropogenic pressures continue to drive biodiversity loss despite increasing conservation

efforts[1]. Land use is currently the greatest driver of biodiversity change[1,2], and has thus been the

focus of much previous conservation research. However, the impacts of climate change on

biodiversity are expected to increase considerably[3], and, by 2070, climate may match or surpass land

use as the greatest driver of biodiversity change[4,5]. Agricultural land used for food production covers

an estimated 38-55% of Earth's habitable land[6,7], while the global food system is responsible for 21-

37% of anthropogenic greenhouse gas (GHG) emissions[8-10]. The development of sustainable food

systems will be critical in halting/reversing land-driven[11]. and climate-driven biodiversity loss[12].

Increasingly, food grown in one country is traded internationally to satisfy demand

elsewhere[13-15]. The associated biodiversity impact of this international consumption is driven by

upstream economic activities that are often geographically distant from the locations of biodiversity

loss[16]. Reducing the environmental impacts of food therefore requires both supply-side and

demand-side changes[16].

Understanding the separate contributions of agricultural land use and greenhouse gas

emissions to biodiversity loss is an important step toward developing a sustainable international

trade in food. A strategy which considers land use alone might well differ from one that incorporates

GHG-driven biodiversity loss. Importantly, the land and GHG footprint per tonne of crops arising

from N_2O tend to be negatively correlated – intensive farming uses less land than extensive

farming, but has higher GHG emissions (N_2O) arising from higher fertilizer use[17]. Moreover, land use

and climate differ in their spatial impacts on biodiversity loss. Land use affects biodiversity that is

local to the land conversion. In contrast, climate change affects biodiversity globally, no matter the

location of the emissions source.

Environmentally-extended multi-regional input-output models (EEMRIOs) are used to link

downstream environmental impacts to upstream drivers, allowing the 'footprints' of commodities to

be followed back through complex supply chains[18]. The footprint of a hamburger, for example,

would contain impacts not only from cattle but from cattle feed, fertilizer, machinery, water,

packaging, fuel etc.[19]. Summing the impacts associated with each upstream product would, in

theory, give the hamburger's total footprint. However, in reality this simple addition process is

impossible given the complications of tracking supply chains, double counting of recycled products,

and infinite loops, for example electricity production requiring water which requires electricity to

pump it. EEMRIOs resolve these problems by using input-output tables to infer production recipes,

allocating environmental costs to sectors to avoid double counting and approximating infinite sums

with the Leontief inverse matrix[19]. Regional production data and their associated environmental

costs are combined with information on international trade, allowing the calculation of a variety of

consumption footprints, e.g. land use[20], GHG emissions[21] and water[22].

EEMRIO models have been used to follow the effects of international trade along supply

chains, to estimate biodiversity footprints. Lenzen et al.[23] performed the first global biodiversity

footprint analysis, using the IUCN Red List to count species' threats within trade regions. This

method assumes that species are equally threatened across their ranges, and excludes non-

threatened species. Other methods use a similar philosophy to calculate footprints based on

biodiversity threat hotspots[24], bird ranges and the number of individual birds lost from an area[16]. An

alternative method, derived from land-use data, uses the countryside species-area relationship to

estimate the potential number of extinctions caused by land conversion and international

trade[15,25,26]. Since many of the extinctions have yet to be realized, it is unclear how these extinctions

would be allocated to different actors across time[16]. Wilting et al.[27] take a step further, deriving

biodiversity footprints from GHG emissions as well as land use, using the biodiversity metric Mean

Species Abundance (although climate impacts are based on model predictions of changes in species

richness, so land-use and climate impacts are not entirely comparable). The GLOBIO biodiversity

model[28], on which Wilting et al.'s[27] analysis was based, assumes that effects of land use and climate

change are even across all terrestrial regions; in reality, biodiversity tends to be more sensitive to

land use and climate change in tropical regions[29,30].

We build on these prior analyses, calculating the impacts of agricultural land use and GHG-

emission footprints using models that directly output the same two metrics of terrestrial

biodiversity: change in local species richness and local rarity-weighted species richness, relative to an

unimpacted baseline. Species richness, although easy to measure, captures only one of the many

dimensions of biodiversity, and does not always decline with global biodiversity loss[31]. Rarity-

weighted richness gives greater weight to species with small geographic range size (range size

correlates with species extinction risks[32]) and so declines if rare species are replaced by more

common ones. Furthermore, we use biodiversity models that allow us for the first time to capture

regional variation in the sensitivity of biodiversity both to land-use differences and to climate

change[29]. We base our biodiversity metrics on local measures of biodiversity in the relevant

agricultural areas as opposed to a value averaged across an entire trade region, meaning that we

account for the wide variation in species richness that occurs within regions.

We examine the international production and consumption footprints of food-related

commodities in terms of: a) land area; b) species richness (land-driven and GHG-driven); and c)

rarity-weighted species richness (land-driven and GHG-driven). We calculate footprints for 33 food-

related products that span food's journey from field to farm gate to household to waste in 49

regions (44 countries and 5 "rest of world" regions). We ask whether the different footprints would

give the same conclusions with respect to sustainable production, consumption and trade. We

identify the regions and food groups with the highest biodiversity footprints, examining the

contributions of land use and GHG emissions to these footprints. We explore biodiversity footprints

per km^2 and per capita for each region and look at the proportion of regions' footprints which are

imported. Our analysis provides a detailed examination into the pathways by which regions'

consumption of particular food-related products affects biodiversity worldwide, giving insight into

the trade-offs between land use and GHG emissions, and into priorities for demand-side changes.

2. Methods

2.1 EEMRIO analysis

Input-output analysis is a top-down approach that uses sectoral transactions data (either in

financial terms or units of product) to account for the complex interdependencies of industries. An

input-output table can be environmentally extended by adding information on exchanges with the

environment, e.g. GHG emissions or land use, by each industry sector. EEMRIO analysis traces the

production and supply chain of traded goods and services and their associated materials back to the

source of primary extraction, thus capturing the direct and indirect environmental pressures

associated with a country's final consumption[19]. Full details of the calculations underlying EEMRIO

analysis can be found in Kitzes[19] and Miller and Blair[18].

We used the standard environmentally-extended Leontief model to calculate the effects of

consumption on biodiversity loss

$$E = f \cdot (I - A)^{-1} \cdot Y$$

114   (Equation 1)

Where, for *i* regions and *m* economic sectors:

E is a $(1 \times i)$ matrix, representing the environmental impacts associated with the final

demand of each region (GHG emissions, in kg CO_2-eq, and agricultural area, in km^2).

f is the $(1 \ x \ im)$ direct intensity vector, representing the environmental pressures (e.g. area

of land, mass of CO_2 emissions) associated with one unit (€1M) of production for each

product sector in each region.

I is the $(im \times im)$ identity matrix.

A is the $(im \times im)$ technical coefficient matrix which gives the amount of input (€1M) that

every sector must receive from every other sector in order to produce one unit (€1M) of

output.

Y is the $(im \times i)$ matrix of final demand given in monetary terms (€1M).

The direct intensity vector f allows the MRIO to be extended to include environmental costs

of production and consumption. Sections 2.2 and 2.3 describe our environmental 'characterisation

factors' based on species richness, rarity-weighted richness, land use and GHG emissions. The vector

relating to land use is a sparse vector populated in the entries for production activities that involve

use of cropland or pasture.

We use the EEMRIO database EXIOBASE3.8.1 (product by product version)[33], because of its

superior balance of sectoral and regional disaggregation relative to other MRIOs. EXIOBASE3

provides a harmonized time-series of MRIO tables and environmental extensions over the period

1995-2022, covering 200 products, 44 countries (EU countries and other major economies), plus 5

'Rest of World' (RoW) regions. We chose 2011 as our year for analysis since land use data are

available up to 2011 only and spatial crop data were not available after this date (see below).

Our analysis focuses on the production and consumption of the 33 product-sectors

associated with food (Supplementary Table 1) in the year 2011, including fertiliser production and

food waste processing. 'Hotel and restaurant services' was excluded since it encompasses energy

and materials, for example, as well as food. Full details of EXIOBASE's land-use and emissions

accounts are given in Stadler et al.[33]. Our land-driven biodiversity footprints are based on

agricultural land, and exclude impacts of forestry, infrastructure or other uses. Emissions from crops

derive from nitrogen and phosphorous fertilisers, while emissions from livestock derive from the

animals, the manure excreted and the cultivation of feed crops. Impacts of fertiliser refer to the

production of the fertiliser only and impacts of food waste do not include the production impacts of

the wasted food. For simplicity, we regard land use as a one-off cost that results in a change in

biodiversity and we assume that once land has been converted, it can be used repeatedly without

the biodiversity cost increasing. In contrast, GHG-driven biodiversity loss accrues year on year. GHG

emissions that are released during land conversion are not considered, because, without detailed

land-history knowledge, we cannot estimate the proportion of emissions that have dissipated since

conversion, nor apportion food-production emissions across years. To put this gap in our coverage of

GHGs into context, direct emissions from agriculture contribute 5.1-6.1 Pg CO_2-eq,/yr while the

clearing of native land for agriculture contributes around 5.9 (SD 2.9) Pg CO_2-eq/yr[34].

2.2 Characterisation factors for land-driven biodiversity impacts

Maps for a) the 49 EXIOBASE trade regions, b) the six biomes used in Newbold et al.[29] (but

here also including boreal forest within the temperate-forest grouping), and c) the 15 EXIOBASE

agricultural land-use categories were identified, and map masks for each unique 3-way combination

were generated. Shapefiles mapping the borders of individual countries[35] were aggregated and

rasterised to match EXIOBASE's 49 regions. Biomes were obtained from The Nature Conservancy[36]

and were aggregated into tropical forest, temperate/boreal forest, tropical grassland,

temperate/montane grassland, Mediterranean, and drylands (tundra, mangroves, flooded

grasslands, inland water and rock & ice were excluded; Supplementary Table 2). Cropland data were

obtained from the Spatial Production Allocation Model (SPAM)[37] and from EarthStat[38] and pasture

data from EarthStat[39]. All spatial data were reprojected to an equal-area Behrmann grid. A summary

schematic is given in Figure 1a and further details of the preparation of datasets are given in the

Supplementary Information.

a)

b)

Figure 1. Schematics summarising the characterisation factor calculations. Characterisation factors for a) land-driven biodiversity loss and b) GHG-driven biodiversity loss. The methods allow for biodiversity gain which would be represented as a negative loss.

Mapped spatial estimates of species richness and rarity-weighted richness for terrestrial vertebrates were obtained by stacking species geographical distributions at a 10-km equal-area resolution. Extent-of-occurrence distribution maps were obtained from the IUCN[40], BirdLife International[41], Meiri et al. [42], and Roll et al. [43]. We selected only areas where species are extant or probably extant, and resident during the breeding season. As in Etard et al. [44], we further excluded areas outside the known elevational limits of species[40]. Species richness was calculated as the number of species occurring within a grid cell. Rarity-weighted species richness weights were calculated as the inverse of a species' estimated range size, and rarity-weighted richness as the sum of these weights across species occurring within a grid cell. Mean species richness and rarity-

weighted richness were calculated for each land-use type within each biome within each trade

region.

We estimated the sensitivity of biodiversity to land use using models of the PREDICTS

database[45,46], following the methods given in Newbold et al.[29] (see Supplementary Information). The

PREDICTS database contains 3.25 million samples of biodiversity in six different land uses from 26,

114 locations and 47,044 different species. For the purposes of this study, we modelled biodiversity

differences among four land-use classes: primary vegetation, secondary vegetation, agriculture

(including both plantations and herbaceous cropland) and pasture, allowing the response of

biodiversity (i.e. species richness and rarity-weighted richness) to these land-use categories to vary

among the six broad biome groupings listed above. For further details, see the Supplementary

Information.

Characterisation factors for species richness were calculated as follows. We used the species

richness map to estimate the mean species richness $S_{i,j,k}$ that would occur in undisturbed land

across the area covered by each EXIOBASE land-use type i, within each biome j and each EXIOBASE

trade region k. The change in species richness $\Delta S_{i,j,k}$ due to land-use was calculated as:

$$\Delta S_{i,j,k} = S_{i,j,k} \times P_{j,l}$$

(Equation 2)

where $P_{j,l}$ is the proportional change in species richness in PREDICTS land-use type l and biome j,

compared to primary vegetation in that biome (see Supplementary Table 3 for the correspondence

between EXIOBASE and PREDICTS land-use types, and Supplementary Tables 4 and 5 for values of

$P_{j,l}$). The change in species richness $\Delta S_{i,k}$ within each EXIOBASE land-use type i and each region k

was calculated as:

$$\Delta S_{i,k} = \sum_{j=1}^{J_{i,k}} (\Delta S_{i,j,k} \times B_{i,j,k})$$

207   (Equation 3)

208   where $J_{i,k}$ is the number of biomes covered by the EXIOBASE land use type i within the region k and

209    $B_{i,j,k}$ is the proportion of the EXIOBASE land-use type i within the region k that is covered by the

210   biome j. The characterisation factor $CF_{i,k}$ was then calculated as:

$$CF_{i,k} = \Delta S_{i,k} \times A_{i,k}$$

(Equation 4)

where $A_{i,k}$ is the area of agricultural land used to produce €1M of product in land-use type i in

region k). Characterisation factors for rarity-weighted richness were calculated using the same

method, but using the map of rarity-weighted species richness, and corresponding modelled

estimates of land-use sensitivity (for characterisation factor values, see Supplementary Tables 6 and

7).

The characterisation factors for species richness have units of number of species $\times$ km^2 and

can be thought of as the count of the species lost, with this loss extending over a certain area (i.e.,

the area of land use given in EXIOBASE). Rarity-weighted richness characterisation factors are less

intuitive, but can be thought of as having a unit of species range fractions $\times$ km^2.

2.3 Characterisation factors for GHG-driven biodiversity loss

We calculated GHG-driven biodiversity loss in a three-step process. First, we calculated the

GHG emissions associated with each region's 2011 final demand for the 33 different food-related

products. Second, we calculated the warming that would result from these emissions. Third, we

calculated the biodiversity loss that would result from this warming. A summary schematic is given in

Figure 1b.

EXIOBASE gives emissions for six GHGs: carbon dioxide (CO_2), methane (CH_4), nitrous oxide

(N_2O), and three fluorinated gases. We calculated footprints using the first three gases only. Their

contribution to warming via food-related emissions is much greater than that of the fluorinated

gases. We summed methane emissions over the four categories given by EXIOBASE ('Combustion –

air', 'non-combustion', 'agriculture' and 'waste').

We calculated the increase in global surface temperature due to GHG emissions using the

Global Temperature Change Potential (GTP), a metric designed to be an intuitive measure of climate

response[47]. The GTP is defined as the ratio between the global mean surface temperature change at

a given future time horizon following an emission (pulse or sustained) of a compound relative to a

reference gas[47]. We chose a 20-year time horizon, to capture warming due to the relatively short-

lived methane emissions, and to represent a time that is tangible to today's policy and decision-

makers. GTP values in units of degrees of warming (°C) per kilogram of emissions were taken from

the AR5 IPCC report[48], Table 8.A.1 (CO_2 6.84×10^{-16} °C /kg, CH_4 4.62×10^{-14} °C /kg, N_2O 1.89×10^{-13}

°C /kg). The warming in °C that would result by 2031 $\Delta T_{p,k}$ due to the emissions $E_{g,p,k}$ associated

with each gas g due to the production of €1M of each product p in each region k was calculated as:

$$\Delta T_{p,k} = \sum_g (E_{g,p,k} \times C_g)$$

(Equation 5)

where C_g is the GTP value for each gas g.

We estimated the sensitivity of biodiversity to climate change based on future projections of

changes in the distributions of terrestrial vertebrates under the Representative Concentration

Pathways (RCP) climate-change scenarios[5]. It is necessary to use future projections rather than

observed responses of species to climate change, because we do not yet have enough data for a

wide range of species from locations that have experienced extensive historical climate change[29].

Full details are given in the Supplementary Information.

To account for the non-uniformity of warming across the globe, we calculated a grid of

spatial weighting factors to up or down weight temperature change depending on the difference

between the projected local and mean global temperature change. The projected temperature

change ΔT_x in each cell x between 2011 and 2031 under RCP8.5 was calculated by averaging the

projections of four different climate models (GFDL-ESM2M, HadGEM2-ES, IPSL-CM5A-LR, MIROC5[49]).

The temperature weighting factor F_x for each cell x was calculated as

$$F_x = \Delta T_x / \overline{\Delta T_x}$$

257 (Equation 6)

258 where $\overline{\Delta T_x}$ is the mean global temperature change. The fractional loss in species richness $\Delta S_{x,p,k}$ in

259 2031 due to $\Delta T_{p,k}$, the temperature increase from emissions from food production of each product p

260 in each region k, was calculated as:

$$\Delta S_{x,p,k} = F_x \times H_x \times \Delta T_{p,k}$$

(Equation 7)

where H_x is the grid estimating sensitivity of species richness to climate change.

This fractional loss in species richness $\Delta S_{x,p,k}$ was then multiplied by our species richness

grid and by the area of each grid cell to give a characterisation factor for GHG-driven biodiversity

loss, $CF_{p,k}$, with a unit of species $\times$ km^2 that is comparable to our metric for land-driven biodiversity

loss. This characterisation factor was calculated as:

$$CF_{p,k} = \Delta S_{x,p,k} \times S_x \times A_x$$

(Equation 8)

where S_x is the species richness in undisturbed vegetation under a natural climate in grid cell x and

A_x is the area of grid cell x. The same basic method was used to calculate the characterisation

factors for change in rarity-weighted richness (see Supplementary Information for details). The GHG-

driven characterisation factors for change in species and rarity-weighted richness are directly

correlated (for characterisation factor values, see Supplementary Tables 8 and 9.)

273   Global maps illustrating the proportional richness changes that would occur due to land-

driven and GHG-driven processes are given in Supplementary Figure 3. Our metrics of land-driven

and GHG-driven biodiversity loss are modelled and presented in the same units, and hence are

comparable. However, there are differences in the way that the two drivers impact biodiversity, and

in the methods we used to model these impacts. We consider land use as a one-off cost that has

already occurred, but GHG emissions as repeated annual costs that occur 20 years after emission.

We would expect the biodiversity loss caused by a single year of emissions to be much lower than

that caused by the one-off cost of total land conversion. We calculate the ratio of land-driven to

GHG-driven biodiversity loss for products and regions. We can crudely approximate a ratio of 100,

for example, to mean that within a century, assuming the same emissions repeat annually, the

global biodiversity loss caused by emissions from a region's food production will equal the

biodiversity loss due to total land conversion in that region. In reality, this may be an underestimate

since we do not consider the emissions released by land conversion.

2.4 Per land area and per capita footprints

288   The regions within EXIOBASE have unequal land areas and human populations. To make a

fairer comparison of footprints between regions, we converted production footprints to footprint

per km^2 and consumption footprints to footprint per capita. Per-km^2 footprints were calculated

using EXIOBASE trade regions' areas (Supplementary Table 10), which were calculated in R from ESRI

country shapefiles[35]. Per-capita footprints were calculated using population data for the year 2011

for the EXIOBASE trade regions (Supplementary Table 11), which were obtained from

CountryEconomy.com[50] (Taiwan) and the World Bank[51] (all other regions).

3. Results

3.1 Regional production and consumption-based footprints for total food

For all footprints, regions with a high production-based footprint tend to have a high
consumption-based footprint. Our land-driven biodiversity footprints show a different picture from a
simple land-area footprint. Important information is missed if the area of agricultural land alone is
used as a proxy for the impact of food production on biodiversity (Figure 2). Differences between
land-associated footprints are driven in part by differences in species richness between regions'
agricultural areas but also by our characterisation factors allowing for biome-specific sensitivity to
land use. Furthermore, footprints also vary according to the measure of biodiversity used. The
regions with the greatest land-area footprints in 2011 were Rest of World (RoW) Africa, China, and
RoW Asia & Pacific (Figure 2a). However, while RoW Africa also has the highest land-driven species
richness footprint, Brazil and RoW Central & South America (RoW CS America) have the second and
third highest species richness footprints, respectively (Figure 2b). RoW CS America hosts many
narrow-ranged species, which is reflected in its high rarity-weighted richness footprint (Figure 2c).
Decisions regarding land use and food production should therefore consider not only the land area
involved but also the land-driven loss of different facets of biodiversity (see Supplementary Figure 4
and Supplementary Table 12 for the footprints of individual products within regions).

GHG-driven biodiversity loss shows a different pattern from the land-driven footprints
(Figure 2) with China, the US and India having the highest production-based footprints. RoW Africa is
notable for having a much lower GHG-driven footprint relative to its land-driven biodiversity
footprint. The ratio of land-driven to GHG-driven biodiversity loss varied by region from 54 for rarity-
weighted richness production footprint in Russia to the order of 10,000 for production in RoW
Africa, for example. Finding ratios as low as 50-100 is concerning as it shows that direct emissions
from a single year of a region's food production will cause biodiversity loss equivalent to 1-2% of the
biodiversity loss caused by that region's total historic land use. Furthermore, we substantially
underestimate biodiversity losses from GHG emissions since our analysis does not include emissions
from land clearance.

Some regions are net importers of land-driven biodiversity loss but net exporters of GHG-
driven biodiversity loss, e.g. China, or *vice versa*, e.g. Australia. RoW Asia & Pacific, RoW CS America,
Brazil, Mexico and South Africa are all net exporters of both land-driven and GHG-driven biodiversity
losses whilst Russia, RoW Middle East and Japan are net importers (Figure 2).

Figure 2. Total production and consumption-based footprints of food-related products. Footprints relate to a) land area, b) land-driven species richness (SR) loss, c) land-driven rarity-weighted richness (RWR) loss, d) GHG-driven biodiversity loss (right-hand axis – species richness; left hand axis – rarity-weighted richness), e) the ratio of land-driven species richness loss to GHG-driven species richness loss and f) the ratio of land-driven rarity-weighted richness loss to GHG-driven rarity-weighted richness loss. Black bars show production-based and grey bars consumption-based footprints. Regions which are in the highest ten for any footprint are shown.

3.2 Production-based footprints for world regions aggregated by food-related sector

Production-based footprints vary considerably among food-related groups and, within food-
related groups, among world regions (Figure 3). As for total food footprints, land-driven biodiversity
footprints do not mirror land-area footprints, and the relative size of biodiversity footprints is
affected by the richness metric. Using animal-derived products as an example, Asia & Pacific has a
land footprint 60% higher than CS America's (and Africa's, Figure 3a), but its land-driven species-
richness footprint is less than half (Figure 3b), and its rarity-weighted richness footprint less than a
third that of CS America (Figure 3c). This result reflects Asia & Pacific's lower average natural species
richness compared to CS America's and the lower biodiversity sensitivity of some of its biomes to
land use.

Again, GHG-driven and land-driven biodiversity footprints show different patterns, with
tropical regions no longer having consistently higher footprints. Although Asia & Pacific has the
highest GHG-driven footprints for all food groups but waste, production in Western Europe and
North America also drives high GHG-driven biodiversity loss. With land-driven to GHG-driven ratios
in the order of 100, Western and Eastern Europe's animal-derived and 'other food' products stand
out as areas for which emissions reductions would have a large benefit for biodiversity alongside
reductions in land area used for agriculture.

Animal-derived products tend to have higher footprints than plant-derived (i.e. crops),
although Asia & Pacific is a notable exception for both land-driven and GHG-driven biodiversity
footprints, in part due to production of paddy rice in India and fruit/vegetables/nuts in China (for
individual product footprints see Supplementary Table 13). GHG-driven biodiversity loss from
fertiliser production and the processing of food waste is much lower than that stemming from the
production of food itself. Land-driven biodiversity loss from these two food-related groups is very
low since their production uses very little agricultural land.

The 'Other Food' category has the highest production-based GHG-driven biodiversity
footprint for almost all world regions (Figure 3d). The ratio of land-driven:GHG-driven footprints for
this category is in the order of 100 for non-tropical world regions (Figure 3e,f), indicating that
reducing GHG emissions from this sector would be a priority for reducing biodiversity loss. Other
Food includes 'Food products not elsewhere classified (nec)' (coded in EXIOBASE as 'OFOD') and
beverages ('BEVR'). 'Food products nec' contributes the vast majority of the Other Food footprint
and covers a broad range of processed foods such as soups, sandwiches and sauces. The EXIOBASE
land-use types that contribute most to the Food Products nec land-driven footprints are cereal

grains nec, cattle pasture, wheat, oil seeds, raw milk pasture and fruit/vegetables/nuts

(Supplementary Figure 5).

Figure 3. Production-based footprints of aggregated food-related groups within world

regions. Footprints relate to a) land area, b) land-driven species richness (SR) loss, c) land-driven

rarity-weighted richness (RWR) loss, d) GHG-driven biodiversity loss (right-hand axis – species

richness; left-hand axis – rarity-weighted richness), e) the ratio of land-driven species richness loss to

GHG-driven species richness loss and f) the ratio of land-driven rarity-weighted richness loss to GHG-

driven rarity-weighted richness loss.

3.3 Per-area and per-capita footprints

Taiwan stands out as having the highest per-area land-related production-based footprints

(Figure 4). The agriculturally dense but biodiversity poor UK is an example of a region with a

relatively high per-area land-use footprint but lower per-area land-driven biodiversity footprints

(Figure 4). In contrast, the species-rich Brazil has a lower per-area land-use footprint but a relatively

higher per-area land-driven species richness footprint. Some regions (e.g. Brazil, India, South Africa)

have both high total production-based footprints (Figure 2) and high per-area production-based

footprints (Figure 4). The other regions with high per-area production tend to be smaller regions

with wide agricultural coverage, for example the Netherlands and Belgium. Small agricultural regions

have the highest per-area production-based GHG-driven footprints, for example Taiwan, Belgium

and Malta.

394   Figure 4. Total per-area production-based footprints of food-related products. Footprints
relate to a) land use, b) land-driven species richness (SR) loss, c) land-driven rarity-weighted richness
(RWR) loss and d) GHG-driven biodiversity loss (right-hand axis - species richness; left-hand axis -
rarity-weighted richness). Regions which are in the highest ten for any footprint are shown.

Australia is notable for having extremely high per-capita land-driven consumption-based

footprints (Figure 5), driven particularly by animal-based products (notably 'Products of Cattle'), but

also by plant-based products and 'Other Food' (Supplementary Figure 7). Luxembourg also has

consistently high per-capita consumption-based footprints, probably in part due its highly affluent

population[51]. In contrast to Australia, Luxembourg's consumption-based footprint is primarily driven

by plant-derived products (fruit/vegetables/nuts, oil seeds and crops nec (i.e., coffee, cocoa,

spices)'). China, India and RoW Asia & Pacific are not within the top ten highest per-capita

consumption-based footprints for any category, despite having high total consumption due to their

large populations (Figure 2). However, some regions with high total biodiversity consumption also

have high per-capita footprints (Australia, Brazil, Japan, Mexico, Russia, the United States, RoW

Africa, RoW CS America and South Africa.) Again, we see contrasts between different footprint

types. For example, Brazil's per-capita land-driven species richness consumption-based footprint is

much larger than RoW CS America's, but the latter's per-capita rarity-weighted footprint is almost

twice that of Brazil. Regions with relatively low species richness can have high per-capita land-driven

biodiversity consumption-based footprints as a result of importing food.

The products underpinning high per-capita consumption-based GHG-driven biodiversity

footprints vary among regions. Belgium's high footprint, for example, is largely driven by other foods

nec (e.g. pizza, ready-meals, sauces) and dairy products, Canada's by beverages and dairy products,

Estonia's by dairy products, raw milk and other foods nec, Luxembourg's by fruit/vegetable/nuts,

crops nec (i.e., coffee, cocoa, spices) and chemical fertiliser, and South Africa's by

fruit/vegetable/nuts and other foods nec (see also Supplementary Figures 7 & 14 for further

breakdowns).

Figure 5. Total per-capita consumption-based footprints of food-related products. Footprints

are in terms of a) land use, b) land-driven species richness (SR) loss, c) land-driven rarity-weighted

richness (RWR) loss and d) GHG-driven biodiversity loss (right-hand axis – species richness; left-hand

axis – rarity-weighted richness). Regions which are in the highest ten for any footprint are shown.

3.4 Biodiversity loss embedded in export/import trade

We found a near-reversal of net trade flows for land-driven versus GHG-driven biodiversity

footprints (Figure 6, Supplementary Figure 8). The United Kingdom, Germany, Russia, Japan and

RoW Middle East are all net importers of land-driven biodiversity loss but net exporters of GHG-

driven biodiversity loss. The different land-driven footprints also show different messages. For

example, RoW CS America exports a similar land-area footprint to that exported by RoW Asia &

Pacific but approximately four times RoW Asia & Pacific's exported rarity-weighted richness

footprint. RoW Africa is a net exporter of land-driven species richness impacts but a net importer of

land-driven rarity-weighted richness impacts. The three greatest net importers of both land-driven

biodiversity footprints are RoW Middle East, Russia and the United States.

.

Figure 6. Regions' net import/export footprints for food-related products. Footprints are in terms of a) land use, b) land-driven species richness (SR) loss, c) land-driven rarity-weighted richness (RWR) loss and d) GHG-driven richness loss. Regions which are in the highest five for any of the import (+ve)/export (-ve) footprints are shown.

Understanding the percentage of a region's footprint that is imported is important in devising policies to reduce footprints. Imports make up 5% or less of land-driven species richness footprints for highly biodiverse regions such as Mexico and RoW CS America (Figure 7); reducing

domestic consumption in these regions might actually lower their consumption-based footprints. In

contrast, the regions with the highest percentages of imported footprints are European.

Luxembourg, the Republic of Ireland and the United Kingdom import 98%, 88% and 82% of their

land-driven species richness footprints respectively. These high percentages will in part be due to

the regions' relatively low domestic food production footprints but may also reflect imports grown in

highly biodiverse regions, which could potentially be sourced more sustainably.

Figure 7. The percentage of a region's food-related footprint that is imported. Footprints are

in terms of a) land use, b) land-driven species richness (SR) loss, c) land-driven rarity-weighted

richness (RWR) loss and d) GHG-driven richness loss. Regions which are in the 5 lowest or highest

percentage for any footprint are shown.

10%, 15% and 9% respectively of land-driven species richness, land-driven rarity-weighted

richness and GHG-driven richness footprints are embedded within trade between world regions

(Supplementary Figure 10), with patterns differing between the footprint types, and between animal

and plant-derived foods (Supplementary Figures 10-12). Although the total footprints of animal-

derived products are higher than plant-derived, a greater percentage of the plant-derived footprint

is traded between world regions (approximately 8%, 15%, 6% (animal-derived) versus 17%, 21%,

12% (plant-derived) of the total footprint for land-driven species richness, land-driven rarity-

weighted richness and GHG-driven species richness respectively). These results are discussed in

more detail in the Supplementary Information.

4. Discussion

Our metric of land-driven species richness change captures the higher biodiversity footprint

of products from more species-rich regions and from biomes in which biodiversity is more sensitive

to land-use change. The metric thus provides a more accurate footprint of the biodiversity impact

embedded within food than would be obtained using land area alone. Weighting richness by species'

range area further changes assessment of embedded biodiversity footprint. Rarity-weighted richness

places a greater emphasis on the biodiversity costs in areas containing a higher proportion of

narrow-ranged species, which are of greater conservation concern than wider-ranging species,

notably parts of CS America. Our analysis also shows the need to consider GHG-driven biodiversity

loss, particularly in assessing longer term future impacts of food production. We find that the

climate contributions of food production are accumulating rapidly over time – in some regions,

annual GHG-driven production footprints are as high as 1% of the total historic land-use footprint,

meaning that just a decade's worth of emissions (not including land conversion emissions) will add

an additional 10% to the biodiversity loss that has already occurred due to wholesale conversion of

land to agriculture.

4.1 Comparison with other studies

Our study incorporated the novel combination of: (i) allowing sensitivity to land use to vary

by land-use type and biome; (ii) allowing for the natural variation of species richness across political

regions; and (iii) allowing for spatial variation in future temperature change and in species'

responses to that temperature change. The models we used to capture land-use and climate impacts

output the same biodiversity metrics, allowing us to compare these two major drivers of biodiversity

change. Comparing the results of our study with those of previous analyses of the embedded

biodiversity footprints of food is not straightforward since studies generally differ in resolution with

respect to regions and/or products. Where comparisons can be made, we find both common ground

and divergence. In line with Marques et al. [26], we show that production in Africa, CS America and

Asia & Pacific regions has the highest overall biodiversity footprints, and that cattle products have a

particularly high impact on biodiversity globally. Marques et al. [26] use a biodiversity metric of

number of impending extinctions of bird species and, in common with our land-driven species

richness metric, identify almost equally high impacts of cattle in Africa as in CS America. However, it

is only by using our rarity-weighted richness measure that the particularly high cost to narrow-

ranged species from cattle products in CS America is revealed. Chaudhary and Kastner[15] use a metric

of the number of species committed to extinction. Their study had greater spatial disaggregation

than ours meaning we cannot compare regions included within our RoW regions. Nevertheless, our

results broadly support each other with respect to highlighting particularly high consumption

footprints in India, China, Brazil and the US. However, Chaudhary and Kastner found Indonesia to

have the second highest consumption footprint, whereas it does not appear in our top ten

consumption footprints for any metric. The relative magnitude of the regions with the greatest

consumption footprints in Chaudhary et al.'s [52] study differs from our results. These discrepancies

are likely to result from the biodiversity metric used as well as differences in trade models, and

further support our finding that different biodiversity metrics lead to different conclusions. Sun et al.

[53] also use a metric of the number of species committed to extinction at a greater spatial

disaggregation than our study. Their results show strong agreement with ours with respect to

consumption per capita, both studies finding high footprints in Central America, Brazil, South Africa

and Australia. Lenzen et al. [23] look at the biodiversity footprint, as measured by number of threats to

species, of all commodities, not just those associated with food. Both Lenzen et al.'s [23] and our

studies find the US, Germany, the UK and Japan to be among the greatest importers of embedded

biodiversity but our study has Russia as a major importer of embedded biodiversity whereas Lenzen

et al. [23] find it to be one of the greatest net exporters.

Our estimates of the impacts of food-related GHGs on biodiversity are much lower than

those of Wilting et al. [27] which is at least in part due to differences in methodology. We predicted the

temperature rise in 2031 due to an emissions pulse in 2011 using the global temperature potential

(GTP). Wilting et al. [27] used the integrated GTP over a 100 year time horizon, thus summing the

warming that occurred in every year following the emission pulse, as opposed to calculating the

temperature of the hundredth year only. We would therefore expect Wilting et al.'s [27] GHG-driven

footprint to be two orders of magnitude greater than ours, as was the case. Wilting et al. [27] analysed

the land-driven and GHG-driven biodiversity loss from all economic sectors using an indicator of

Mean Species Abundance and estimated that GHGs contributed to an average of 18% of the

biodiversity loss associated with food although contributions varied by region from 7% (Africa) to

45% (rest of Oceania). Accounting for the different metrics of global warming used, our results were

relatively similar, with our GHG-driven footprints estimated to contribute from the order of 0.01% –

1% of biodiversity loss for food products and up to 10% for fertiliser. Wilting et al.'s [27] study further

differed from ours since it used a different biodiversity metric and MRIO, and assumed agricultural

land was evenly distributed across regions. Our studies support each other in suggesting current

GHG-driven emissions from the food-related products will play a very significant role in future

biodiversity loss. Reducing emissions associated with food is therefore a high priority, particularly in

Europe, North America and Asia & Pacific.

4.2 Policy implications

Our study shows that measurements of trade-related impacts of biodiversity differ

considerably depending on the metric used. Whilst detailed discussion of the merits and relevance

of different biodiversity metrics may be appropriate in the scientific literature, it is unlikely to be

helpful to businesses and policy makers who require succinct and readily interpretable biodiversity

footprints. This leads to the question of which measure(s) should be recommended to the public and

private sectors to aid them in reducing their environmental footprints. What might the

consequences be of one metric being chosen over others? Our study suggests that a metric of land

area, whilst easy to measure and interpret, will over inflate the cost of biodiversity in biodiversity-

poor areas and under-estimate it in biodiversity-rich areas, and thus is not recommended when the

focus is biodiversity conservation. Species richness is preferable in this case, and is still relatively

easy to interpret, particularly if used in relative rather than absolute terms. However, species

richness has been criticised as a biodiversity metric, since it does not decline if rare species are

replaced with common species (see Hillebrand et al. [31]). Weighting for species rarity overcomes this,

and highlights the higher losses of smaller-ranged species from CS America. Using rarity-weighted

richness over a simple species richness measure would likely lead to different decisions with respect

to sustainable trading.

We must also consider whether GHG-driven biodiversity loss adds value or unnecessary

complexity to footprint estimates. Our study has shown that it is fairly simple to calculate GHG-

driven biodiversity loss and to aggregate it with the land-driven impact. The annual GHG-driven

biodiversity impact of food produced in the US and Western Europe is not insignificant and, in

livestock sectors, is equivalent to approximately 1% of land-driven biodiversity loss. Importantly, this

equates to 10% of the biodiversity impact of complete land conversion within the timespan of just

one decade. For some individual products, at a level of disaggregation finer than that used in this

study, the ratio of GHG to land-driven impacts could be even higher. Including the GHG-driven losses

in estimates of the food industry's impact on biodiversity would therefore seem worthwhile.

The biodiversity footprints that we present could be used to inform policies and trade

agreements targeted at reducing biodiversity loss. Policies which encourage a reduction of high-

income countries' consumption of beef, for example, could be used to target biodiversity loss in

Australia and CS America, regions which dominate exported biodiversity footprint embedded in

cattle products. Trade deals which lead to an increase in imports from these regions (e.g. the 2021

UK-Australia trade deal, the European Union's 'New Green Deal') will likely lead to an increase in

biodiversity loss and offshoring of yet more of Europe's environmental footprint[54] unless strict and

enforceable environmental regulations are included in the agreements. Decisions on future trade

deals must take embedded environmental costs into account and use appropriate metrics of

biodiversity. If impacts are based on land use alone they will severely underestimate biodiversity

impacts in CS America.

The examination of production-based footprints per-area (which facilitate comparisons of

production between different-sized regions) allows for an increased understanding of the

implications of regional/national agricultural policies. Regions with high per-area production-based

footprints are targets for policies that reduce the environmental impact of farming. There is

considerable overlap of high per-area land-driven and GHG-driven production impacts and it is

interesting that the Netherlands and Belgium, both less species-rich temperate countries, have some

of the highest per-area production-based footprints, even for rarity-weighted richness (Figure 4).

Taiwan is notable for all three of its per-area land-driven production-based footprints being

extremely high (Figure 4), with impacts being driven by a variety of products including

fruit/vegetables/nuts, animal products nec (e.g., eggs, honey), fish, pork products, sugar, and food

products nec (e.g., pizza, soups, sauces) (Supplementary Figure 6). India stands out as having very

high paddy rice production per area for all footprints (Supplementary Figure 6).

Per-capita consumption-based footprints (which facilitate comparisons of consumption

between regions with different population sizes) also allow for targeting of agricultural and trade

policies. We found China, the US and India had the highest overall GHG-driven biodiversity

footprints, but once population size was considered, only the US remained in the top ten per-capita

footprints. Identifying the products that lead to regions' high per-capita footprints is the first step

towards reducing those footprints. For example, extremely high per-capita consumption-based

footprints from raw milk and dairy products contribute to Estonia's place in the top ten for GHG-

driven biodiversity loss (see Figure 5 and Supplementary Figure 7). Whilst we recognise that

changing cultural norms can be very difficult, reducing Estonia's per-capita footprint would be

relatively straight forward from a policy perspective, since it involves only one food group and could

perhaps be tackled via a combination of dietary shifts, source-shifting and change in dairy-farming

practices. Australia, in contrast, has particularly high per-capita land-driven consumption-based

footprints for a wide variety of products spanning cereal grains nec, fruit/vegetables/nuts, sugar

cane/beet, all products associated with cattle and other meat animals (e.g. sheep), rice, food

products nec and beverages. Reducing these footprints will likely require a more complex suite of

strategies.

We found that some regions import over 90% of their embedded biodiversity footprint,

further indicating the need for sustainable trade as a measure to halt biodiversity loss. Regions with

a high proportion of imported footprint coupled with high per-capita consumption-based footprints

are targets for reducing their imports and/or the biodiversity footprint embedded within those

imports[16]; Estonia, Luxembourg, the Netherlands and Norway all fall into this category (Figures 5 and

7). Conversely, regions with a low percentage of imported footprint but high per-capita consumption

such as Brazil, Mexico and South Africa should be lowering their domestic footprint in order to

reduce their biodiversity impact.

The main net importers of land-driven biodiversity footprint tend to be net exporters of

GHG-driven biodiversity footprint, and vice versa. The emissions associated with cropland largely

arise from fertiliser use. Low fertiliser application will be associated with low emissions but lead to a

smaller yield and therefore a larger land-use footprint. On the other hand, heavy fertiliser use will

result in more emissions but smaller land use. This negative correlation is exacerbated since areas

that use more fertiliser tend to be areas of lower species richness such as N America and W

Europe[55].

4.3 Limitations

There are several limitations of MRIOs which must be considered in interpreting our results:
sectoral and regional aggregation[56,57], an assumption of price homogeneity[58,59], limitations in data
reporting[60], and the age of input data[61]. The original EXIOBASE 3 data end in 2011 and although
more recent trade data are now available, land use is still limited to 2011
(https://zenodo.org/record/5589597#.YnkHvOjMK3A). Our results therefore reflect the biodiversity
impact of food production over a decade ago, and will likely underestimate the current impact due
to the expansion of agricultural land[62] in the intervening period.

In common with other footprinting studies, e.g.[25,26], our analysis only measures biodiversity
impacts caused by direct land-use change, and does not explicitly consider the impacts of
agricultural intensification. Neither does our study explicitly account for indirect biodiversity loss
that might occur through loss of ecosystem function, e.g. soil impoverishment. Nevertheless, the
PREDICTS database does sample gradients of agricultural intensity and ecosystem degradation, and
so these factors should be captured implicitly[29].

While our study highlights that the consumption of cattle products contributes considerably
to land-driven biodiversity loss, it is important to be aware that there is high variability in pasture
maps stemming from the different definitions, methods and underlying datasets used in their
generation[63]. However, the Ramankutty et al.[39] dataset that we used is one of the best available,
aligning well with other datasets[63].

4.4 Conclusions

We have presented new methods to estimate the land-driven and GHG-driven biodiversity
impacts embedded within the production, consumption and trade of food-related commodities. We
find that land-driven footprints differ depending on the metric used, and that GHG-driven
biodiversity impacts are more intense in regions of higher income and lower species-richness. The
measures we present are simple to calculate and could be incorporated into decision making and
environmental impact assessments by governments and businesses. Our consistent metric for
biodiversity impacts allows us to present multiple aspects of land-driven and GHG-driven
biodiversity footprints, enabling insight into priorities for reducing biodiversity costs via both global
and regional production and consumption.

References

653   **References**

654   1   IPBES. Global assessment report on biodiversity and ecosystem services of the
655   Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. (Bonn,
656   Germany, 2019).

657   2   Maxwell, S. L., Fuller, R. A., Brooks, T. M. & Watson, J. E. M. The ravages of guns, nets and
658   bulldozers. *Nature* **536**, 143-145 (2016).

659   3   Schipper, A. M. *et al.* Projecting terrestrial biodiversity intactness with GLOBIO 4. *Glob Chang*
660   *Biol* **26**, 760-771 (2020). https://doi.org:10.1111/gcb.14848

661   4   Di Marco, M. *et al.* Projecting impacts of global climate and land-use scenarios on plant
662   biodiversity using compositional-turnover modelling. *Glob Chang Biol* **25**, 2763-2778 (2019).
663   https://doi.org:10.1111/gcb.14663

664   5   Newbold, T. Future effects of climate and land-use change on terrestrial vertebrate
665   community diversity under different scenarios. *Proc Biol Sci* **285** (2018).
666   https://doi.org:10.1098/rspb.2018.0792

667   6   Ellis, E. C., Klein Goldewijk, K., Siebert, S., Lightman, D. & Ramankutty, N. Anthropogenic
668   transformation of the biomes, 1700 to 2000. *Global Ecology and Biogeography*, no-no
669   (2010). https://doi.org:10.1111/j.1466-8238.2010.00540.x

670   7   FAOStat.   (License: CC BY-NC-SA 3.0 IGO., 2022).

671   8   Mbow, C. *et al.* Food Security. (2019).

672   9   Rosenzweig, C., N Tubiello, F., Sandalow, D., Benoit, P. & N Hayek, M. Finding and fixing food
673   system emissions: the double helix of science and policy. *Environmental Research Letters* **16**
674   (2021). https://doi.org:10.1088/1748-9326/ac0134

675   10   Crippa, M. *et al.* Food systems are responsible for a third of global anthropogenic GHG
676   emissions. *Nature Food* **2**, 198-209 (2021).

677   11   Leclere, D. *et al.* Bending the curve of terrestrial biodiversity needs an integrated strategy.
678   *Nature* **585**, 551-556 (2020). https://doi.org:10.1038/s41586-020-2705-y

679   12   Lynch, J., Cain, M., Frame, D. & Pierrehumbert, R. Agriculture's contribution to climate
680   change and role in mitigation is distinct from predominantly fossil CO_2-emitting sectors.
681   *Frontiers in Sustainable Food Systems* **4**, 518039 (2021).

682   13   Kastner, T., Erb, K.-H. & Haberl, H. Rapid growth in agricultural trade: effects on global area
683   efficiency and the role of management. *Environmental Research Letters* **9**, 034015 (2014).

684   14   FAO. *The State of Agricultural Commodity Markets 2020. Agricultural markets and*
685   *sustainable development: Global value chains, smallholder farmers and digital innovations.*
686   (FAO, 2020).

687   15   Chaudhary, A. & Kastner, T. Land use biodiversity impacts embodied in international food
688   trade. *Global Environmental Change* **38**, 195-204 (2016).
689   https://doi.org:10.1016/j.gloenvcha.2016.03.013

690   16   Kitzes, J. *et al.* Consumption-Based Conservation Targeting: Linking Biodiversity Loss to
691   Upstream Demand through a Global Wildlife Footprint. *Conserv Lett* **10**, 531-538 (2017).
692   https://doi.org:10.1111/con4.12321

693   17   Balmford, A. *et al.* The environmental costs and benefits of high-yield farming. *Nature*
694   *Sustainability* **1**, 477-485 (2018). https://doi.org:10.1038/s41893-018-0138-5

695   18   Miller, R. E. & Blair, P. D. *Input-output Analysis: Foundations and Extension.* (Cambridge
696   University Press, 2009).

697   19   Kitzes, J. An introduction to environmentally-extended input-output analysis. *Resources* **2**,
698   489-503 (2013).

699   20   Roux, N., Kastner, T., Erb, K.-H. & Haberl, H. Does agricultural trade reduce pressure on land
700   ecosystems? Decomposing drivers of the embodied human appropriation of net primary
701   production. *Ecological Economics* **181** (2021).
702   https://doi.org:10.1016/j.ecolecon.2020.106915

21 Peters, G. & Hertwich, E. G. CO_2 embodied in international trade with implications for global
climate policy. *Environmental Science and Technology* **42**, 1401-1407 (2008).
22 Hoekstra, A. Y. & Mekonnen, M. M. The water footprint of humanity. *Proceedings of the*
National Academy of Sciences **109**, 3232-3237 (2012).
23 Lenzen, M. *et al.* International trade drives biodiversity threats in developing nations. *Nature*
486, 109-112 (2012).
24 Moran, D. & Kanemoto, K. Identifying species threat hotspots from global supply chains.
Nature Ecology and Evolution **1**, 0023 (2017).
25 Chaudhary, A., Verones, F., de Baan, L. & Hellweg, S. Quantifying Land Use Impacts on
Biodiversity: Combining Species-Area Models and Vulnerability Indicators. *Environ Sci*
Technol **49**, 9987-9995 (2015). https://doi.org:10.1021/acs.est.5b02507
26 Marques, A. *et al.* Increasing impacts of land use on biodiversity and carbon sequestration
driven by population and economic growth. *Nat Ecol Evol* **3**, 628-637 (2019).
https://doi.org:10.1038/s41559-019-0824-3
27 Wilting, H. C., Schipper, A. M., Bakkenes, M., Meijer, J. R. & Huijbregts, M. A. Quantifying
Biodiversity Losses Due to Human Consumption: A Global-Scale Footprint Analysis. *Environ*
Sci Technol **51**, 3298-3306 (2017). https://doi.org:10.1021/acs.est.6b05296
28 Alkemade, R. *et al.* GLOBIO3: A Framework to Investigate Options for Reducing Global
Terrestrial Biodiversity Loss. *Ecosystems* **12**, 374-390 (2009).
https://doi.org:10.1007/s10021-009-9229-5
29 Newbold, T., Oppenheimer, P., Etard, A. & Williams, J. J. Tropical and Mediterranean
biodiversity is disproportionately sensitive to land-use and climate change. *Nat Ecol Evol* **4**,
1630-1638 (2020). https://doi.org:10.1038/s41559-020-01303-0
30 Martins, I. S. & Pereira, H. M. Improving extinction projections across scales and habitats
using the countryside species-area relationship. *Sci Rep* **7**, 12899 (2017).
https://doi.org:10.1038/s41598-017-13059-y
31 Hillebrand, H. *et al.* Biodiversity change is uncoupled from species richness trends:
Consequences for conservation and monitoring. *Journal of Applied Ecology* **55**, 169-184
(2018). https://doi.org:10.1111/1365-2664.12959
32 Chichorro, F., Juslén, A. & Cardoso, P. A review of the relation between species traits and
extinction risk. *Biological Conservation* **237**, 220-229 (2019).
33 Stadler, K. *et al.* EXIOBASE 3: Developing a time series of detailed environmentally extended
multi-regional input-output tables. *Journal of Industrial Ecology*
https://doi.org/10.1111/jiec.12715 (2018).
34 Smith, P. & Gregory, P. J. Climate change and sustainable food production. *Proc Nutr Soc* **72**,
21-28 (2013). https://doi.org:10.1017/S0029665112002832
35 ESRI. (2015).
36 The Nature Conservancy. (2019).
37 International Food Policy Research Institute. *MapSPAM* (ed International Food Policy
Research Institute) (Harvard Dataverse, 2019).
38 Monfreda, C., Ramankutty, N. & Foley, J. A. Farming the planet: 2. Geographic distribution of
crop areas, yields, physiological types, and net primary production in the year 2000. *Global*
Biogeochemical Cycles **22**, GB1022 (2008). https://doi.org:doi: 10.1029/2007GB002947
39 Ramankutty, N., Evan, A. T., Monfreda, C. & Foley, J. A. Farming the planet: 1. Geographic
distribution of global agricultural lands in the year 2000. *Global Biogeochemical Cycles* **22**,
GB1003 (2008). https://doi.org:doi: 10.1029/2007GB002952
40 IUCN, I. U. f. C. o. N. (2020).
41 BirdLife International. (2020).
42 Meiri, S. *et al.* (2017).

43 Roll, U. *et al.* The global distribution of tetrapods reveals a need for targeted reptile
conservation. *Nat Ecol Evol* **1**, 1677-1682 (2017). https://doi.org:10.1038/s41559-017-0332-
2
44 Etard, A., Morrill, S. & Newbold, T. Global gaps in trait data for terrestrial vertebrates. *Global*
Ecology and Biogeography **29**, 2143-2158 (2020).
https://doi.org:https://doi.org/10.1111/geb.13184
45 Hudson, L. N. *et al.* The database of the PREDICTS (Projecting Responses of Ecological
Diversity In Changing Terrestrial Systems) project. *Ecol Evol* **7**, 145-188 (2017).
https://doi.org:10.1002/ece3.2579
46 Hudson, L. N. *et al.*   (ed Natural History Museum) (https://doi.org/10.5519/0066354, 2016).
47 Shine, K. P., Fuglestvedt, J. S., Hailemariam, K. & Stuber, N. Alternatives to the global
warming potential for comparing climate impacts of emissions of greenhouse gases. *Climate*
Change **68**, 281-302 (2005).
48 Myhre, G. *et al.* Anthropogenic and Natural Radiative Forcing. (Cambridge University Press,
Cambridge, United Kingdom and New York, NY, USA, 2013).
49 Hijmans, R., Cameron, S. E., Parra, J. L., Jones, P. G. & Jarvis, A. Very high resolution
interpolated climate surfaces for global land areas. *International Journal of Climatology* **25**,
1965-1978 (2005). https://doi.org:doi.org/10.1002/joc.1276
50 CountryEconomy.com.
https://countryeconomy.com/demography/population/taiwan?year=2011 2021).
51 World Bank.   (2021).
52 Chaudhary, A., Carrasco, L. R. & Kastner, T. Linking national wood consumption with global
biodiversity and ecosystem service losses. *Sci Total Environ* **586**, 985-994 (2017).
https://doi.org:10.1016/j.scitotenv.2017.02.078
53 Sun, Z., Behrens, P., Tukker, A., Bruckner, M. & Scherer, L. Shared and environmentally just
responsibility for global biodiversity loss. *Ecological Economics* **194** (2022).
https://doi.org:10.1016/j.ecolecon.2022.107339
54 Fuchs, R., Brown, C. & Rounsevell, M. Europe's Green Deal offshores environmental damage
to other nations. *Nature* **586**, 671-673 (2020).
55 Mueller, N. D. *et al.* Closing yield gaps through nutrient and water management. *Nature* **490**,
254-257 (2012). https://doi.org:10.1038/nature11420. 490:254-257
56 Steen-Olsen, K., Owen, A., Hertwich, E. G. & Lenzen, M. Effects of sector aggregation on CO_2
multipliers in multiregional input-output analyses. *Economic Systems Research* **26**, 284-302
(2014).
57 Lenzen, M. Aggregation versus disaggregation in input-output analysis of the environment.
Economic Systems Research **23**, 73-89 (2011).
58 Merciai, S. & Heijungs, R. Balance issues in monetary input-output tables. *Ecological*
Economics **102**, 69-74 (2014).
59 de Koning, A. *et al.* Effect of aggregation and disaggregation on embodied material use of
products in input-output analysis. *Ecological Economics* **116**, 289-299 (2015).
60 Tukker, A. *et al.* Towards robust, authoritative assessments of environmental impacts
embodied in trade: Current state and recommendations. *Journal of Industrial Ecology* **22**
(2018).
61 Chandrakumar, C., McLaren, S., Malik, A., Ramilan, T. & Lenzen, M. Understanding New
Zealand's consumption-based greenhouse gas emissions: an application of multi-regional
input-output analysis. *The International Journal of Life Cycle Assessment* **25**, 1323-1332
(2020).
62 FAO. How to Feed the World in 2050. (Rome, Italy, 2009).
63 Oliveira, J. *et al.* Choosing pasture maps: An assessment of pasture land classification
definitions and a case study of Brazil. *International Journal of Applied Earth Observation and*
Geoinformation **93** (2020). https://doi.org:10.1016/j.jag.2020.102205