Impacts of global food supply on biodiversity via land use and climate change

Abstract

Land-use change is currently the greatest driver of biodiversity change, with climate change

predicted to match or surpass its impacts by mid-century. The global food system is a key driver of

both these anthropogenic pressures, thus the development of sustainable food systems will be

critical to halting and reversing biodiversity loss. Previous studies of the biodiversity footprint of

food tend to focus on land use alone. We use the multi-regional input-output model EXIOBASE to

estimate the impacts of biodiversity embedded within the global food system. We build on prior

analyses, calculating the impacts of both agricultural land-use and greenhouse gas (GHG) emission

footprints for the same two metrics of biodiversity: local species richness and rarity-weighted

species richness. Our biodiversity models capture regional variation in the sensitivity of biodiversity

both to land-use differences and to climate change. We find that the footprint of land area does not

capture the biodiversity impact embedded within trade that is provided by our metric of land-driven

species richness change, and that our metric of rarity-weighted richness places a greater emphasis

on the biodiversity costs in Central and South America. We find that methane emissions are

responsible for 70% of the overall GHG-driven biodiversity footprint and that, in several regions,

emissions from a single year's food production cause biodiversity loss equivalent to 2% or more of

that region's total historic land use. The measures we present are simple to calculate and could be

incorporated into decision making and environmental impact assessments by governments and

businesses.

Keywords

Multi-regional input-output model; impacts embodied in trade; biodiversity footprint; consumption-

based accounting; agriculture; food system sustainability

1. Introduction

Anthropogenic pressures continue to drive biodiversity loss despite increasing conservation

efforts[1]. Land use is currently the greatest driver of biodiversity change[1,2], and has thus been the

focus of much previous conservation research. However, the impacts of climate change on

biodiversity are expected to increase considerably[3], and, by 2070, climate may match or surpass land

use as the greatest driver of biodiversity change[4,5]. Agricultural land used for food production covers

an estimated 38-55% of Earth's habitable land[6,7], while the global food system is responsible for 21-

37% of anthropogenic greenhouse gas (GHG) emissions[8-10]. The development of sustainable food

systems will be critical in halting/reversing land-driven[11]. and climate-driven biodiversity loss[12].

Increasingly, food grown in one country is traded internationally to satisfy demand

elsewhere[13-15]. The associated biodiversity impact of this international consumption is driven by

upstream economic activities that are often geographically distant from the locations of biodiversity

loss[16]. Reducing the environmental impacts of food therefore requires both supply-side and

demand-side changes[16].

Understanding the separate contributions of agricultural land use and greenhouse gas

emissions to biodiversity loss is an important step toward developing a sustainable international

trade in food. A strategy which considers land use alone might well differ from one that incorporates

GHG-driven biodiversity loss. Importantly, the land and GHG footprint per tonne of crops arising

from N_2O tend to be negatively correlated – intensive farming uses less land than extensive

farming, but has higher GHG emissions (N_2O) arising from higher fertilizer use[17]. Moreover, land use

and climate differ in their spatial impacts on biodiversity loss. Land use affects biodiversity that is

local to the land conversion. In contrast, climate change affects biodiversity globally, no matter the

location of the emissions source.

Environmentally-extended multi-regional input-output models (EEMRIOs) are used to link

downstream environmental impacts to upstream drivers, allowing the 'footprints' of commodities to

be followed back through complex supply chains[18]. The footprint of a hamburger, for example,

would contain impacts not only from cattle but from cattle feed, fertilizer, machinery, water,

packaging, fuel etc.[19]. Summing the impacts associated with each upstream product would, in

theory, give the hamburger's total footprint. However, in reality this simple addition process is

impossible given the complications of tracking supply chains, double counting of recycled products,

and infinite loops, for example electricity production requiring water which requires electricity to

pump it. EEMRIOs resolve these problems by using input-output tables to infer production recipes,

allocating environmental costs to sectors to avoid double counting and approximating infinite sums

with the Leontief inverse matrix[19]. Regional production data and their associated environmental

costs are combined with information on international trade, allowing the calculation of a variety of

consumption footprints, e.g. land use[20], GHG emissions[21] and water[22].

EEMRIO models have been used to follow the effects of international trade along supply

chains, to estimate biodiversity footprints. Lenzen et al.[23] performed the first global biodiversity

footprint analysis, using the IUCN Red List to count species' threats within trade regions. This method assumes that species are equally threatened across their ranges, and excludes non-threatened species. Other methods use a similar philosophy to calculate footprints based on biodiversity threat hotspots[24], bird ranges and the number of individual birds lost from an area[16]. An alternative method, derived from land-use data, uses the countryside species-area relationship to estimate the potential number of extinctions caused by land conversion and international trade[15,25,26]. Since many of the extinctions have yet to be realized, it is unclear how these extinctions would be allocated to different actors across time[16]. Wilting et al.[27] take a step further, deriving biodiversity footprints from GHG emissions as well as land use, using the biodiversity metric Mean Species Abundance (although climate impacts are based on model predictions of changes in species richness, so land-use and climate impacts are not entirely comparable). The GLOBIO biodiversity model[28], on which Wilting et al.'s[27] analysis was based, assumes that effects of land use and climate change are even across all terrestrial regions; in reality, biodiversity tends to be more sensitive to land use and climate change in tropical regions[29,30].

We build on these prior analyses, introducing three novel aspects. (i) We calculate the biodiversity impacts of agricultural land use and GHG-emission footprints using models that directly output metrics of terrestrial biodiversity change in the same units, allowing the drivers' impacts to be compared and splitting emissions into carbon dioxide (CO_2), methane (CH_4) and nitrous dioxide (N_2O). (ii) We consider change in local rarity-weighted species richness relative to an unimpacted baseline in addition to local species richness. Species richness, although easy to measure, captures only one of the many dimensions of biodiversity, and does not always decline with global biodiversity loss[31]. Rarity-weighted richness gives greater weight to species with small geographic range size (range size correlates with species extinction risks[32]) and so declines if rare species are replaced by more common ones. (iii) We use biodiversity models that allow us for the first time to capture regional variation in the sensitivity of biodiversity both to land-use differences and to climate change[29]. We base our biodiversity metrics on local measures of biodiversity in the relevant agricultural areas as opposed to a value averaged across an entire trade region, meaning that we account for the wide variation in species richness that occurs within regions.

We examine the international production and consumption footprints of food-related commodities in terms of: a) land area; b) species richness (land-driven and GHG-driven); and c) rarity-weighted species richness (land-driven and GHG-driven). We calculate footprints for 33 food-related products that span food's journey from field to farm gate to household to waste in 49 regions (44 countries and 5 "rest of world" regions). We ask whether the different footprints would give the same conclusions with respect to sustainable production, consumption and trade. We

identify the regions and food groups with the highest biodiversity footprints, examining the

contributions of land use and GHG emissions to these footprints. We explore biodiversity footprints

per km^2 and per capita for each region and look at the proportion of regions' footprints which are

imported. Our analysis provides a detailed examination into the pathways by which regions'

consumption of particular food-related products affects biodiversity worldwide, giving insight into

the trade-offs between land use and GHG emissions, and into priorities for demand-side changes.

2. Methods

2.1 EEMRIO analysis

Input-output analysis is a top-down approach that uses sectoral transactions data (either in

financial terms or units of product) to account for the complex interdependencies of industries. An

input-output table can be environmentally extended by adding information on exchanges with the

environment, e.g. GHG emissions or land use, by each industry sector. EEMRIO analysis traces the

production and supply chain of traded goods and services and their associated materials back to the

source of primary extraction, thus capturing the direct and indirect environmental pressures

associated with a country's final consumption[19]. Full details of the calculations underlying EEMRIO

analysis can be found in Kitzes[19] and Miller and Blair[18].

We used the standard environmentally-extended Leontief model to calculate the effects of

consumption on biodiversity loss

$$E = f \cdot (I - A)^{-1} \cdot Y$$

118   (Equation 1)

Where, for *i* regions and *m* economic sectors:

E is a $(1 \times i)$ matrix, representing the environmental impacts associated with the final

demand of each region (CO_2, CH_4 and N_2O emissions, in kg, and agricultural area, in km^2).

f is the $(1 \times im)$ direct intensity vector, representing the environmental pressures (e.g. area

of land, mass of CO_2 emissions) associated with one unit (€1M) of production for each

product sector in each region.

I is the $(im \times im)$ identity matrix.

A is the $(im \times im)$ technical coefficient matrix which gives the amount of input (€1M) that
every sector must receive from every other sector in order to produce one unit (€1M) of
output.

Y is the $(im \times i)$ matrix of final demand given in monetary terms (€1M).

The direct intensity vector f allows the MRIO to be extended to include environmental costs
of production and consumption. Sections 2.2 and 2.3 describe our environmental 'characterisation
factors' based on species richness, rarity-weighted richness, land use and GHG emissions. The vector
relating to land use is a sparse vector populated in the entries for production activities that involve
use of cropland or pasture.

We use the EEMRIO database EXIOBASE3.8.1 (product by product version)[33], because of its
superior balance of sectoral and regional disaggregation relative to other MRIOs. EXIOBASE3
provides a harmonized time-series of MRIO tables and environmental extensions over the period
1995-2022, covering 200 products, 44 countries (EU countries and other major economies), plus 5
'Rest of World' (RoW) regions. We chose 2011 as our year for analysis since land use data are
available up to 2011 only and spatial crop data were not available after this date (see below).

Our analysis focuses on the production and consumption of the 33 product-sectors
associated with food (Supplementary Table 1) in the year 2011, including fertiliser production and
food waste processing. 'Hotel and restaurant services' was excluded since it encompasses energy
and materials, for example, as well as food. Full details of EXIOBASE's land-use and emissions
accounts are given in Stadler et al.[33]. Our land-driven biodiversity footprints are based on agricultural
land, and exclude impacts of forestry, infrastructure or other uses. Emissions from crops derive from
nitrogen and phosphorous fertilisers, while emissions from livestock derive from the animals, the
manure excreted and the cultivation of feed crops. Impacts of fertiliser refer to the production of the
fertiliser only and impacts of food waste do not include the production impacts of the wasted food.
For simplicity, we regard land use as a one-off cost that results in a change in biodiversity and we
assume that once land has been converted, it can be used repeatedly without the biodiversity cost
increasing. In contrast, GHG-driven biodiversity loss accrues year on year. GHG emissions that are
released during land conversion are not considered, because, without detailed land-history
knowledge, we cannot estimate the proportion of emissions that have dissipated since conversion,
nor apportion food-production emissions across years. To put this gap in our coverage of GHGs into
context, direct emissions from agriculture contribute 5.1-6.1 Pg CO_2-eq,/yr while the clearing of
native land for agriculture contributes around 5.9 (SD 2.9) Pg CO_2-eq/yr[34].

2.2 Characterisation factors for land-driven biodiversity impacts

Maps for a) the 49 EXIOBASE trade regions, b) the six biomes used in Newbold et al.[29] (but here also including boreal forest within the temperate-forest grouping), and c) the 15 EXIOBASE agricultural land-use categories were identified, and map masks for each unique 3-way combination were generated. Shapefiles mapping the borders of individual countries[35] were aggregated and rasterised to match EXIOBASE's 49 regions. Biomes were obtained from The Nature Conservancy[36] and were aggregated into tropical forest, temperate/boreal forest, tropical grassland, temperate/montane grassland, Mediterranean, and drylands (tundra, mangroves, flooded grasslands, inland water and rock & ice were excluded; Supplementary Table 2). Cropland data were obtained from the Spatial Production Allocation Model (SPAM)[37] and from EarthStat[38] and pasture data from EarthStat[39]. All spatial data were reprojected to an equal-area Behrmann grid. A summary schematic is given in Figure 1a and further details of the preparation of datasets are given in the Supplementary Information.

a)

b)

Figure 1. Schematics summarising the characterisation factor calculations. Characterisation

factors for a) land-driven biodiversity loss and b) GHG-driven biodiversity loss. The methods allow for

biodiversity gain which would be represented as a negative loss.

Mapped spatial estimates of species richness and rarity-weighted richness for terrestrial

vertebrates were obtained by stacking species geographical distributions at a 10-km equal-area

resolution. Extent-of-occurrence distribution maps were obtained from the IUCN[40], BirdLife

International[41], Meiri et al.[42], and Roll et al.[43]. We selected only areas where species are extant or

probably extant, and resident during the breeding season. As in Etard et al.[44], we further excluded

areas outside the known elevational limits of species[40]. Species richness was calculated as the

number of species occurring within a grid cell. Rarity-weighted species richness weights were

calculated as the inverse of a species' estimated range size, and rarity-weighted richness as the sum

of these weights across species occurring within a grid cell. Mean species richness and rarity-

weighted richness were calculated for each land-use type within each biome within each trade
region.

We estimated the sensitivity of biodiversity to land use using models of the PREDICTS
database[45,46], following the methods given in Newbold et al.[29] (see Supplementary Information). The
PREDICTS database contains 3.25 million samples of biodiversity in six different land uses from 26,
114 locations and 47,044 different species. For the purposes of this study, we modelled biodiversity
differences among four land-use classes: primary vegetation, secondary vegetation, agriculture
(including both plantations and herbaceous cropland) and pasture, allowing the response of
biodiversity (i.e. species richness and rarity-weighted richness) to these land-use categories to vary
among the six broad biome groupings listed above. For further details, see the Supplementary
Information.

Characterisation factors for species richness were calculated as follows. We used the species
richness map to estimate the mean species richness $S_{i,j,k}$ that would occur in undisturbed land
across the area covered by each EXIOBASE land-use type i, within each biome j and each EXIOBASE
trade region k. The change in species richness $\Delta S_{i,j,k}$ due to land-use was calculated as:

$$\Delta S_{i,j,k} = S_{i,j,k} \times P_{j,l}$$

204   (Equation 2)

205   where $P_{j,l}$ is the proportional change in species richness in PREDICTS land-use type l and biome j,
206   compared to primary vegetation in that biome (see Supplementary Table 3 for the correspondence
207   between EXIOBASE and PREDICTS land-use types, and Supplementary Tables 4 and 5 for values of
208     $P_{j,l}$). The change in species richness $\Delta S_{i,k}$ within each EXIOBASE land-use type i and each region k
209   was calculated as:

$$\Delta S_{i,k} = \sum_{j=1}^{J_{i,k}} (\Delta S_{i,j,k} \times B_{i,j,k})$$

210
211   (Equation 3)

212   where $J_{l,k}$ is the number of biomes covered by the EXIOBASE land use type i within the region k and
213     $B_{i,j,k}$ is the proportion of the EXIOBASE land-use type i within the region k that is covered by the
214   biome j. The characterisation factor $CF_{i,k}$ was then calculated as:

$$CF_{i,k} = \Delta S_{i,k} \times A_{i,k}$$

215   (Equation 4)

where $A_{i,k}$ is the area of agricultural land used to produce €1M of product in land-use type i in

region k). Characterisation factors for rarity-weighted richness were calculated using the same

method, but using the map of rarity-weighted species richness, and corresponding modelled

estimates of land-use sensitivity (for characterisation factor values, see Supplementary Tables 6 and

7).

The characterisation factors for species richness have units of number of species $\times$ km^2 and

can be thought of as the count of the species lost, with this loss extending over a certain area (i.e.,

the area of land use given in EXIOBASE). Rarity-weighted richness characterisation factors are less

intuitive, but can be thought of as having a unit of species range fractions $\times$ km^2.

2.3 Characterisation factors for GHG-driven biodiversity change

We calculated GHG-driven biodiversity change in a three-step process. First, we calculated

the GHG emissions (split into carbon dioxide (CO_2), methane (CH_4) and nitrous oxide (N_2O))

associated with each region's production of the 33 different food-related products in 2011, using

EXIOBASE. Second, we calculated the warming that would result from these emissions. Third, we

calculated the biodiversity change that would result from this warming, allowing for non-uniformity

of warming across the globe. A summary schematic is given in Figure 1b.

EXIOBASE gives emissions for six GHGs: carbon dioxide, methane, nitrous oxide, and three

fluorinated gases. We calculated footprints using the first three gases only since their contribution to

warming via food-related emissions is much greater than that of the fluorinated gases. We summed

methane emissions over the four categories given by EXIOBASE ('Combustion – air', 'non-

combustion', 'agriculture' and 'waste'). Footprints were calculated using the environmentally-

extended Leontief model described in section 2.1.

We calculated the increase in global surface temperature due to GHG emissions using the

Global Temperature Change Potential (GTP), a metric designed to be an intuitive measure of climate

response[47]. The GTP is defined as the ratio between the global mean surface temperature change at

a given future time horizon following an emission (pulse or sustained) of a compound relative to a

reference gas[47]. We chose a 20-year time horizon, to capture warming due to the relatively short-

lived methane emissions, and to represent a time that is tangible to today's policy and decision-

makers. GTP values in units of degrees of warming (°C) per kilogram of emissions were taken from

the AR5 IPCC report[48], Table 8.A.1 (CO_2 6.84 $\times$ 10^{-16} °C /kg, CH_4 4.62 $\times$ 10^{-14} °C /kg, N_2O 1.89 $\times$ 10^{-13}

°C /kg). The warming in °C that would result by 2031 $\Delta T_{p,k}$ due to the emissions $E_{g,p,k}$ associated

with each gas g due to the production of €1M of each product p in each region k was calculated as:

$$\Delta T_{p,k} = \sum_g (E_{g,p,k} \times C_g)$$

(Equation 5)

where C_g is the GTP value for each gas g.

We estimated the sensitivity of biodiversity to climate change based on future projections of

changes in the distributions of terrestrial vertebrates under the Representative Concentration

Pathways (RCP) climate-change scenarios[5]. It is necessary to use future projections rather than

observed responses of species to climate change, because we do not yet have enough data for a

wide range of species from locations that have experienced extensive historical climate change[29].

Full details are given in the Supplementary Information.

To account for the non-uniformity of warming across the globe, we calculated a grid of

spatial weighting factors to up or down weight temperature change depending on the difference

between the projected local and mean global temperature change. The projected temperature

change ΔT_x in each cell x between 2011 and 2031 under RCP8.5 was calculated by averaging the

projections of four different climate models (GFDL-ESM2M, HadGEM2-ES, IPSL-CM5A-LR, MIROC5[49]).

The temperature weighting factor F_x for each cell x was calculated as

$$F_x = \Delta T_x / \overline{\Delta T_x}$$

263 (Equation 6)

264 where $\overline{\Delta T_x}$ is the mean global temperature change. The fractional loss in species richness $\Delta S_{x,p,k}$ in

265 2031 due to $\Delta T_{p,k}$, the temperature increase from emissions from food production of each product p

266 in each region k, was calculated as:

$$\Delta S_{x,p,k} = F_x \times H_x \times \Delta T_{p,k}$$

(Equation 7)

where H_x is the grid estimating sensitivity of species richness to climate change.

This fractional loss in species richness $\Delta S_{x,p,k}$ was then multiplied by our species richness

grid and by the area of each grid cell to give a characterisation factor for GHG-driven biodiversity

loss, $CF_{p,k}$, with a unit of species × km^2 that is comparable to our metric for land-driven biodiversity

loss. This characterisation factor was calculated as:

$$CF_{p,k} = \Delta S_{x,p,k} \times S_x \times A_x$$

273   (Equation 8)

where S_x is the species richness in undisturbed vegetation under a natural climate in grid cell x and

A_x is the area of grid cell x. The same basic method was used to calculate the characterisation

factors for change in rarity-weighted richness (see Supplementary Information for details). The GHG-

driven characterisation factors for change in species and rarity-weighted richness are directly

correlated (for characterisation factor values, see Supplementary Tables 8 and 9.)

Global maps illustrating the proportional richness changes that would occur due to land-

driven and GHG-driven processes are given in Supplementary Figure 3. Our metrics of land-driven

and GHG-driven biodiversity loss are modelled and presented in the same units, and hence are

comparable. However, there are differences in the way that the two drivers impact biodiversity, and

in the methods we used to model these impacts. We consider land use as a one-off cost that has

already occurred, but GHG emissions as repeated annual costs that occur 20 years after emission.

We would expect the biodiversity loss caused by a single year of emissions to be much lower than

that caused by the one-off cost of total land conversion. We calculate the ratio of land-driven to

GHG-driven biodiversity loss for products and regions. We can crudely approximate a ratio of 100,

for example, to mean that within a century, assuming the same emissions repeat annually, the

global biodiversity loss caused by emissions from a region's food production will equal the

biodiversity loss due to total land conversion in that region. In reality, this may be an underestimate

since we do not consider the emissions released by land conversion.

2.4 Per land area and per capita footprints

The regions within EXIOBASE have unequal land areas and human populations. To make a

fairer comparison of footprints between regions, we converted production footprints to footprint

per km^2 and consumption footprints to footprint per capita. Per-km^2 footprints were calculated

using EXIOBASE trade regions' areas (Supplementary Table 10), which were calculated in R from ESRI

country shapefiles[35]. Per-capita footprints were calculated using population data for the year 2011

for the EXIOBASE trade regions (Supplementary Table 11), which were obtained from

CountryEconomy.com[50] (Taiwan) and the World Bank[51] (all other regions).

3. Results

 3.1 Regional production and consumption-based footprints for total food

For all footprints, regions with a high production-based footprint tend to have a high

consumption-based footprint. Our land-driven biodiversity footprints show a different picture from a

simple land-area footprint. Important information is missed if the area of agricultural land alone is

used as a proxy for the impact of food production on biodiversity (Figure 2). Differences between

land-associated footprints are driven in part by differences in species richness between regions'

agricultural areas but also by our characterisation factors allowing for biome-specific sensitivity to

land use. Furthermore, footprints also vary according to the measure of biodiversity used. The

regions with the greatest land-area footprints in 2011 were Rest of World (RoW) Africa, China, and

RoW Asia & Pacific (Figure 2a). However, while RoW Africa also has the highest land-driven species

richness footprint, Brazil and RoW Central & South America (RoW CS America) have the second and

third highest species richness footprints, respectively (Figure 2b). RoW CS America hosts many

narrow-ranged species, which is reflected in its high rarity-weighted richness footprint (Figure 2c).

Decisions regarding land use and food production should therefore consider not only the land area

involved but also the land-driven loss of different facets of biodiversity (see Supplementary Figure 5

and Supplementary Table 12 for the footprints of individual products within regions).

India and China are the regions responsible for the highest GHG-driven biodiversity loss,

followed by RoW Africa (Figure 2). However, compared to India and China, RoW Africa has a much

lower GHG-driven footprint relative to its land-driven biodiversity footprint. We expected the GHG-

driven footprint to be much lower than the land-driven footprint since it relates to the global

biodiversity loss caused by a single year of emissions whereas our land-driven footprint relates to

the historic conversion of all agricultural land used in 2011. The ratio of land-driven to GHG-driven

biodiversity loss varied by region from 16 for rarity-weighted richness production footprint in Russia

to 855 for production in RoW C&S America, with several regions, including China, India and RoW

Asia, having ratios around 50. Finding ratios of 50 or lower is concerning as it shows that direct

emissions from a single year of a region's food production will cause biodiversity loss equivalent to

2% or more of the biodiversity loss caused by that region's total historic land use. Furthermore, we

substantially underestimate biodiversity losses from GHG emissions since our analysis does not

include emissions from land clearance.

Some regions are net importers of land-driven biodiversity loss but net exporters of GHG-

driven biodiversity loss, e.g. India, or *vice versa*, e.g. Indonesia. RoW Asia & Pacific, RoW CS America,

Australia and Mexico are all net exporters of both land-driven and GHG-driven biodiversity meaning

their international exports are harming both their domestic biodiversity and, via climate change,

global biodiversity. China, the United States, Russia and RoW Middle East are net importers of both

footprints (Figure 2). The top ten footprints stem from regions with very large land areas and/or

populations and, aside from Russia, do not include regions in continental Europe.

Methane emissions account for 70% of the total GHG-driven biodiversity footprint from

food-related products, compared to 42% of the GHG-driven footprint of all EXIOBASE's products.

Carbon dioxide contributed to 18% of food's total footprint versus 54% of the footprints of all

products and nitrous oxide contributed to 12% of food's footprint and 4% of the footprints of all

products. Food-related products account for 23% of the total emissions in EXIOBASE and the total

GHG-driven footprint of all food-related products was approximately 1% of the land-driven richness

footprints.

Figure 2. Total production-based (blue bars) and consumption-based (red bars) footprints of

food-related products. Footprints relate to a) land area, b) land-driven species richness (SR) loss, c)

land-driven rarity-weighted richness (RWR) loss, d) GHG-driven biodiversity loss split by emissions

type: carbon dioxide (dark blue/red), methane (mid blue/red), nitrous oxide (light blue/red) (right-

hand axis – species richness; left hand axis – rarity-weighted richness), e) the ratio of land-driven

species richness loss to total GHG-driven species richness loss and f) the ratio of land-driven rarity-

weighted richness loss to total GHG-driven rarity-weighted richness loss. Regions which are in the

highest ten for one or more footprints are shown.

3.2 Production-based footprints for world regions aggregated by food-related sector

Production-based footprints vary considerably among food-related groups and, within food-

related groups, among world regions (Figure 3). As for total food footprints, land-driven biodiversity

footprints do not mirror land-area footprints, and the relative size of biodiversity footprints is

affected by the richness metric. Using animal-derived products as an example, Asia & Pacific has a

land footprint 60% higher than CS America's (and Africa's, Figure 3a), but its land-driven species-

richness footprint is less than half (Figure 3b), and its rarity-weighted richness footprint less than a

third that of CS America (Figure 3c). This result reflects Asia & Pacific's lower average natural species

richness compared to CS America's and the lower biodiversity sensitivity of some of its biomes to

land use.

Again, GHG-driven and land-driven biodiversity footprints show different patterns, with

tropical regions no longer having consistently higher footprints. Production in Western Europe and

North America also drives relatively high GHG-driven biodiversity loss, particularly in the 'Other

Food' sector although Asia & Pacific has the highest GHG-driven footprints across all food groups

(Figure 3d). Asia & Pacific's ratio of land-driven to GHG-driven ratios ranges from around 40-100

(Figures 3e, f) which is extremely concerning given the high land-driven biodiversity footprints this

region has. If emissions continue at this rate their impact on global biodiversity will, in under a

century, equal that already caused by land conversion in Asia & Pacific.

Animal-derived products tend to have much higher footprints than plant-derived (i.e. crops),

although Asia & Pacific is a notable exception for both land-driven and GHG-driven biodiversity

footprints, in part due to production of paddy rice, wheat and other cereals in India (for individual

product footprints see Supplementary Figure 5 and Supplementary Table 13). As would be expected,

land-driven biodiversity loss from fertiliser production and the processing of food waste is much

lower than that stemming from the production of food itself since the processes use very little

agricultural land.

In many instances, the 'Other Food' category has higher production-based GHG-driven

biodiversity footprints than the footprints from crops (Figures 3a-d). Other Food includes 'Food

products not elsewhere classified (nec)' (coded in EXIOBASE as 'OFOD') and beverages ('BEVR').

'Food products nec' contributes the vast majority of the Other Food footprint and covers a broad

range of processed foods such as soups, sandwiches and sauces. The EXIOBASE land-use types that

contribute most to the Food Products nec land-driven footprints are cereal grains nec, cattle

pasture, wheat, oil seeds, raw milk pasture and fruit/vegetables/nuts (Supplementary Figure 6).

Figure 3. Production-based footprints of aggregated food-related groups within world

regions. Footprints relate to a) land area, b) land-driven species richness (SR) loss, c) land-driven

rarity-weighted richness (RWR) loss, d) GHG-driven biodiversity loss split by emissions type: carbon

dioxide (dark shade), methane (mid shade), nitrous oxide (light shade) (right-hand axis – species

richness; left-hand axis – rarity-weighted richness), e) the ratio of land-driven species richness loss to

total GHG-driven species richness loss and f) the ratio of land-driven rarity-weighted richness loss to

total GHG-driven rarity-weighted richness loss.

3.3 Per-area and per-capita footprints

Taiwan stands out as having the highest per-area land-related production-based footprints

(Figure 4a-c). The agriculturally dense but biodiversity poor UK is an example of a region with

relatively lower per-area land-driven biodiversity footprints than per-area land-use footprints (Figure

4). In contrast, the species-rich Brazil has a lower per-area land-use footprint but a relatively higher

per-area land-driven species richness footprint. Some regions (e.g. Brazil, India, Mexico) have both

high total production-based footprints (Figure 2) and high per-area production-based footprints

(Figure 4). The other regions with high per-area production tend to be smaller regions with wide

agricultural coverage, for example the Netherlands and Belgium. Small agricultural regions have the

highest per-area production-based GHG-driven footprints, for example Taiwan, the Netherlands and

Malta. The relative contributions of the different GHGs differs between regions, for example nitrous

oxide contributing a larger proportion of the Netherlands' and Belgium's GHG-driven footprint than

Malta's (Figure 4d). For per-area footprints for all products and regions see Supplementary Figure 7

and Supplementary Table 14.

Figure 4. Total per-area production-based footprints of food-related products. Footprints

relate to a) land use, b) land-driven species richness (SR) loss, c) land-driven rarity-weighted richness

(RWR) loss and d) GHG-driven biodiversity loss split by emissions type: carbon dioxide (black),

methane (dark grey), nitrous oxide (light grey) (right-hand axis - species richness; left-hand axis -

rarity-weighted richness). Regions which are in the highest ten for one or more footprints are

shown.

Australia is notable for having extremely high land-driven consumption-based footprints per

capita (Figure 5), driven particularly by animal-based products (notably 'Products of Cattle'), but also

by plant-based products and 'Other Food' (Supplementary Figure 8, Supplementary Tables 15, 16).

Luxembourg also has consistently high per-capita consumption-based footprints, probably in part

due its highly affluent population[51]. In contrast to Australia, Luxembourg's consumption-based

footprint is primarily driven by plant-derived products (fruit/vegetables/nuts, oil seeds and other

crops, i.e., coffee, cocoa, spices). Despite having high total consumption due to their large

populations (Figure 2), China, India and RoW Asia & Pacific are not within the top ten highest per-

capita consumption-based footprints for any category. However, some regions with high total

biodiversity consumption also have high per-capita footprints: Australia, Brazil, Mexico, Russia, the

United States, RoW Africa and RoW CS America. Again, we see contrasts between different footprint

types. For example, Brazil's land-driven species richness consumption-based footprint per capita is

much larger than RoW CS America's, but the latter's per-capita rarity-weighted footprint is almost

twice that of Brazil (Figures 5b,c). (Brazil has particularly high per-capita species richness footprints

relative to RoW CS America for cattle, processed cattle, vegetable oils and dairy, see Supplementary

Figure 8). Regions with relatively low species richness can have high per-capita land-driven

biodiversity consumption-based footprints as a result of importing food.

The products underpinning high per-capita consumption-based GHG-driven biodiversity

footprints vary among regions (Figure 5d). Belgium's high footprint, for example, is largely driven by

other foods (e.g. pizza, ready-meals, sauces) and dairy products, Estonia's by dairy products, raw

milk, cattle and other foods, Luxembourg's by fruit/vegetable/nuts, other crops (i.e., coffee, cocoa,

spices), cattle and chemical fertiliser, and the Republic of Ireland's by processed cattle products and

food waste disposed of in land fill. Some regions, e.g. Australia, Luxembourg and Lithuania, have

high footprints across multiple products and at least one animal-based product is associated with

particularly high consumption footprints for all regions in Figure 5, except Taiwan (< ~0.0004 species

x km^2 per capita for every animal-based product) (see also Supplementary Figure 8 and

Supplementary Tables 15, 16 for further breakdowns). Relative consumption of the different GHGs

differs across regions, carbon dioxide contributing relatively highly to South Africa's footprint but

relatively little to Brazil's, for example.

Figure 5. Total consumption-based footprints per-capita of food-related products. Footprints are in terms of a) land use, b) land-driven species richness (SR) loss, c) land-driven rarity-weighted richness (RWR) loss and d) GHG-driven biodiversity loss split by emissions type: carbon dioxide (black), methane (dark grey), nitrous oxide (light grey) (right-hand axis – species richness; left-hand axis – rarity-weighted richness). Regions which are in the highest ten for one or more footprints are shown.

3.4 Biodiversity loss embedded in export/import trade

We found a near-reversal of net imports (i.e. imports minus exports, or consumption-based

footprint minus production-based footprint) for land-driven versus GHG-driven biodiversity

footprints (Figure 6, Supplementary Figure 9). The United States, the United Kingdom, Germany,

Russia, Japan, China and RoW Middle East ⬚ regions that tend to be relatively biodiversity poor and

highly industrialised in their land use systems ⬚ are all net importers of land-driven biodiversity loss

but net exporters of GHG-driven biodiversity loss. The different land-driven footprints also show

different messages. For example, RoW CS America exports a similar land-area footprint to that

exported by RoW Asia & Pacific but approximately four times RoW Asia & Pacific's exported rarity-

weighted richness footprint. RoW Africa is a net exporter of land-driven species richness impacts but

a net importer of land-driven rarity-weighted richness impacts. The three greatest net importers of

both land-driven biodiversity footprints are RoW Middle East, Russia and the United States.

477   .

Figure 6. Regions' net import footprints for food-related products. Footprints are in terms of a) land use, b) land-driven species richness (SR) loss, c) land-driven rarity-weighted richness (RWR) loss and d) GHG-driven richness loss. Regions which are in the highest five (net importers) or lowest five (net exporters) for the net import footprints are shown.

Understanding the percentage of a region's footprint that is imported is important in devising policies to reduce footprints. Imports make up 5% or less of land-driven species richness footprints for highly biodiverse regions such as Brazil and Mexico (Figure 7b, Supplementary Figure

10); reducing domestic consumption in these regions might actually lower their consumption-based

footprints. In contrast, the regions with the highest percentages of imported footprints are

European. Luxembourg, the Republic of Ireland and the United Kingdom import 98%, 88% and 83%

of their land-driven species richness footprints respectively. These high percentages will in part be

due to the regions' relatively low domestic food production footprints but may also reflect imports

grown in highly biodiverse regions, which could potentially be sourced more sustainably.

Luxembourg, Ireland and Norway import high percentages of their GHG-driven biodiversity

footprints (69%, 65% and 48% respectively) (Figure 7d) and also have particularly high per capita

consumption-based GHG-driven footprints (Figure 5d).

Figure 7. The percentage of a region's food-related footprint that is imported. Footprints are in terms of a) land use, b) land-driven species richness (SR) loss, c) land-driven rarity-weighted richness (RWR) loss and d) GHG-driven richness loss. Regions which are in the 5 lowest or highest percentage for one or more footprints are shown.

We find that 10%, 15% and 8% respectively of land-driven species richness, land-driven rarity-weighted richness and GHG-driven richness footprints are embedded within trade between world regions (Supplementary Figure 11), with patterns differing between the footprint types, and

between animal and plant-derived foods (Supplementary Figures 11-13). Although the total
footprints of animal-derived products are higher than plant-derived, a greater percentage of the
plant-derived footprint is traded between world regions: approximately 8%, 15%, 7% (animal-
derived) versus 17%, 21%, 11% (plant-derived) of the total footprint for land-driven species richness,
land-driven rarity-weighted richness and GHG-driven species richness respectively. These results are
discussed in more detail in the Supplementary Information (Supplementary Figure 13).

4. Discussion

Our metric of land-driven species richness change captures the higher biodiversity footprint
of products from more species-rich regions and from biomes in which biodiversity is more sensitive
to land-use change. The metric thus provides a more accurate footprint of the biodiversity impact
embedded within food than would be obtained using land area alone. Weighting richness by species'
range area further changes assessment of embedded biodiversity footprint. Rarity-weighted richness
places a greater emphasis on the biodiversity costs in areas containing a higher proportion of
narrow-ranged species, which are of greater conservation concern than wider-ranging species,
notably parts of CS America. Our analysis also shows the need to consider GHG-driven biodiversity
loss, particularly in assessing longer term future impacts of food production. We find that the
climate contributions of food production are accumulating rapidly over time – in several regions,
annual GHG-driven production footprints are as high as 2% of the total historic land-use footprint,
meaning that just a decade's worth of emissions (not including land conversion emissions) will add
an additional 20% to the biodiversity loss that has already occurred due to wholesale conversion of
land to agriculture. 70% of the total GHG-driven biodiversity footprint of food-related products
stems from methane.

4.1 Comparison with other studies

Our study incorporated the novel combination of: (i) allowing sensitivity to land use to vary
by land-use type and biome; (ii) allowing for the natural variation of species richness across political
regions; and (iii) allowing for spatial variation in future temperature change and in species'
responses to that temperature change. The models we used to capture land-use and climate impacts
output the same biodiversity metrics, allowing us to compare these two major drivers of biodiversity
change. Comparing the results of our study with those of previous analyses of the embedded
biodiversity footprints of food is not straightforward since studies generally differ in resolution with
respect to regions and/or products. Where comparisons can be made, we find both common ground

and divergence. In line with Marques et al.[26], we show that production in Africa, CS America and Asia

& Pacific regions has the highest overall biodiversity footprints, and that cattle products have a

particularly high impact on biodiversity globally. Marques et al.[26] use a biodiversity metric of number

of impending extinctions of bird species and, in common with our land-driven species richness

metric, identify almost equally high impacts of cattle in Africa as in CS America. However, it is only by

using our rarity-weighted richness measure that the particularly high cost to narrow-ranged species

from cattle products in CS America is revealed. Chaudhary and Kastner[15] use a metric of the number

of species committed to extinction. Their study had greater spatial disaggregation than ours

meaning we cannot compare regions included within our RoW regions. Nevertheless, our results

broadly support each other with respect to highlighting particularly high consumption footprints in

India, China, Brazil and the US. However, Chaudhary and Kastner found Indonesia to have the second

highest consumption footprint, whereas it does not appear in our top ten consumption footprints

for any metric. The relative magnitude of the regions with the greatest consumption footprints in

Chaudhary et al.'s[52] study differs from our results. These discrepancies are likely to result from the

biodiversity metric used as well as differences in trade models, and further support our finding that

different biodiversity metrics lead to different conclusions. Sun et al.[53] also use a metric of the

number of species committed to extinction at a greater spatial disaggregation than our study. Their

results show strong agreement with ours with respect to consumption per capita, both studies

finding high footprints in Central America, Brazil, South Africa and Australia. Lenzen et al.[23] look at

the biodiversity footprint, as measured by number of threats to species, of all commodities, not just

those associated with food. Both Lenzen et al.'s[23] and our studies find the US, Germany, the UK and

Japan to be among the greatest importers of embedded biodiversity but our study has Russia as a

major importer of embedded biodiversity whereas Lenzen et al.[23] find it to be one of the greatest

net exporters.

Our estimates of the impacts of food-related GHGs on biodiversity are lower than those of

Wilting et al.[27] which is at least in part due to differences in methodology. We predicted the

temperature rise in 2031 due to an emissions pulse in 2011 using the global temperature potential

(GTP). Wilting et al.[27] used the integrated GTP over a 100 year time horizon, thus summing the

warming that occurred in every year following the emission pulse, as opposed to calculating the

temperature of the hundredth year only. Wilting et al.[27] analysed the land-driven and GHG-driven

biodiversity loss from all economic sectors and estimated that GHGs contributed to an average of

18% of the biodiversity loss associated with food although contributions varied by region from 7%

(Africa) to 45% (rest of Oceania). Given the different metric and time horizons of global warming

used, we would expect our results to differ from Wilting et al.'s[27], with our GHG-driven footprints

estimated to contribute from the order of 0.1% – 6% of biodiversity loss for food products and up to

16% for fertiliser. Wilting et al.'s[27] study further differed from ours since it used a different

biodiversity metric and MRIO, and assumed agricultural land was evenly distributed across regions.

Our studies support each other in suggesting current GHG-driven emissions from the food-related

products will play a very significant role in future biodiversity loss. Reducing emissions associated

with food is therefore a high priority, particularly in Europe, North America and Asia & Pacific.

4.2 Policy implications

Our study shows that measurements of trade-related impacts of biodiversity differ

considerably depending on the biodiversity metric used. Whilst detailed discussion of the merits and

relevance of different biodiversity metrics may be appropriate in the scientific literature, it is unlikely

to be helpful to businesses and policy makers who require succinct and readily interpretable

biodiversity footprints. This leads to the question of which measure(s) should be recommended to

the public and private sectors to aid them in reducing their environmental footprints. What might

the consequences be of one metric being chosen over others? Our study suggests that a metric of

land area, whilst easy to measure and interpret, will over inflate the cost of biodiversity in

biodiversity-poor areas and under-estimate it in biodiversity-rich areas, and thus is not

recommended when the focus is biodiversity conservation. Species richness is preferable in this

case, and is still relatively easy to interpret, particularly if used in relative rather than absolute terms.

However, species richness has been criticised as a biodiversity metric, since it does not decline if rare

species are replaced with common species (see Hillebrand et al.[31]). Weighting for species rarity

overcomes this, and highlights the higher losses of smaller-ranged species from CS America. Using

rarity-weighted richness over a simple species richness measure would likely lead to different

decisions with respect to sustainable trading.

We must also consider whether GHG-driven biodiversity loss adds value or unnecessary

complexity to footprint estimates. Our study has shown that it is fairly simple to calculate GHG-

driven biodiversity loss and to aggregate it with the land-driven impact. The annual GHG-driven

biodiversity impact of livestock sectors is equivalent to approximately 5% of land-driven biodiversity

loss in Asia, the US and Western Europe. Importantly, this equates to 50% of the biodiversity impact

of complete land conversion within the timespan of just one decade. For some individual products,

the ratio of GHG to land-driven impacts is even higher. Including the GHG-driven losses in estimates

of the food industry's impact on biodiversity would therefore seem worthwhile, , particularly since

the bulk of emissions are from methane. Methane's short lifetime means that (assuming methane

sinks are constant), reducing methane emissions actually leads to global cooling but on the flipside,

increased emissions lead to substantial rapid warming[54]. Reducing emissions from food in the

immediate-term would therefore be an extremely effective route to reducing near-term warming.

Our results add to the already weighty evidence, e.g.[55-57], in support of policies that assist farmers to

transition away from livestock and nudge consumers toward a more plant-based diet.

The biodiversity footprints that we present could be used to inform policies and trade

agreements targeted at reducing biodiversity loss. Policies which encourage a reduction of high-

income countries' consumption of beef, for example, could be used to target biodiversity loss in

Australia and CS America, regions which dominate exported biodiversity footprint embedded in

cattle products. Trade deals which lead to an increase in imports from these regions (e.g. the 2021

UK-Australia trade deal, the European Union's 'New Green Deal') will likely lead to an increase in

biodiversity loss and offshoring of yet more of Europe's environmental footprint[58] unless strict and

enforceable environmental regulations are included in the agreements. Decisions on future trade

deals must take embedded environmental costs into account and use appropriate metrics of

biodiversity. If impacts are based on land use alone they will severely underestimate biodiversity

impacts in CS America.

The examination of production-based footprints per-area (which facilitate comparisons of

production between different-sized regions) allows for an increased understanding of the

implications of regional/national agricultural policies. Regions with high per-area production-based

footprints are targets for policies that reduce the environmental impact of farming. There is

considerable overlap of high per-area land-driven and GHG-driven production impacts and it is

interesting that the Netherlands and Belgium, both less species-rich temperate countries, have some

of the highest per-area production-based footprints, even for rarity-weighted richness (Figure 4).

Taiwan is notable for all three of its per-area land-driven production-based footprints being

extremely high (Figure 4), with impacts being driven by a variety of products including

fruit/vegetables/nuts, other animal products (e.g., eggs, honey), fish, pork products, sugar, and

other food products (e.g., pizza, soups, sauces) (Supplementary Figure 7). India stands out as having

very high paddy rice production per area for all footprints (Supplementary Figure 7).

Per-capita consumption-based footprints (which facilitate comparisons of consumption

between regions with different population sizes) also allow for targeting of agricultural and trade

policies. We found China, India and RoW Africa had the highest overall GHG-driven biodiversity

footprints, but once population size was considered, only RoW Africa remained in the top ten per-

capita footprints. Identifying the products that lead to regions' high per-capita footprints is the first

step towards reducing those footprints. For example, extremely high per-capita consumption-based

footprints from cattle, raw milk and dairy products contribute to Estonia's place in the top ten for

GHG-driven biodiversity loss (see Figure 5 and Supplementary Figure 8). Whilst we recognise that

changing cultural norms can be very difficult, reducing Estonia's per-capita footprint would be

relatively straight forward from a policy perspective, since it involves only one food group and could

perhaps be tackled via a combination of dietary shifts, source-shifting and change in dairy-farming

practices. Australia, in contrast, has particularly high per-capita land-driven consumption-based

footprints for a wide variety of products spanning fruit/vegetables/nuts, all products associated with

cattle and other meat animals (e.g. sheep), rice, other food products and beverages. Reducing these

footprints will likely require a more complex suite of strategies.

We found that some regions import over 90% of their embedded biodiversity footprint,

further indicating the need for sustainable trade as a measure to halt biodiversity loss. Regions with

a high proportion of imported footprint coupled with high per-capita consumption-based footprints

are targets for reducing their imports and/or the biodiversity footprint embedded within those

imports[16]; Estonia, Luxembourg, the Netherlands and Norway all fall into this category (Figures 5 and

7). Conversely, regions with a low percentage of imported footprint but high per-capita consumption

such as Brazil, Mexico and South Africa should be lowering their domestic footprint in order to

reduce their biodiversity impact.

The main net importers of land-driven biodiversity footprint tend to be net exporters of

GHG-driven biodiversity footprint, and vice versa. The emissions associated with crop production

largely arise from fertiliser use. Low fertiliser application will be associated with low emissions but

lead to a smaller yield and therefore a larger land-use footprint. On the other hand, heavy fertiliser

use will result in more emissions but smaller land use. This negative correlation is exacerbated since

areas that use more fertiliser tend to be areas of lower species richness such as N America and W

Europe[59].

4.3 Limitations

There are several limitations of MRIOs which must be considered in interpreting our results:

sectoral and regional aggregation[60,61], an assumption of price homogeneity[62,63], limitations in data

reporting[64], and the age of input data[65]. The original EXIOBASE 3 data end in 2011 and although

more recent trade data are now available, land use is still limited to 2011

(https://zenodo.org/record/5589597#.YnkHvOjMK3A). Our results therefore reflect the biodiversity

impact of food production over a decade ago, and will likely underestimate the current impact due

to the expansion of agricultural land[66] in the intervening period.

675   In common with other footprinting studies, e.g.[25,26], our analysis only measures biodiversity

676   impacts caused by direct land-use change, and does not explicitly consider the impacts of

677   agricultural intensification. Neither does our study explicitly account for indirect biodiversity loss

678   that might occur through loss of ecosystem function, e.g. soil impoverishment. Nevertheless, the

679   PREDICTS database does sample gradients of agricultural intensity and ecosystem degradation, and

680   so these factors should be captured implicitly[29].

681   While our study highlights that the consumption of cattle products contributes considerably

682   to land-driven biodiversity loss, it is important to be aware that there is high variability in pasture

683   maps stemming from the different definitions, methods and underlying datasets used in their

684   generation[67]. However, the Ramankutty et al.[39] dataset that we used is one of the best available,

685   aligning well with other datasets[67].

687   *4.4 Conclusions*

688   We have presented new methods to estimate the land-driven and GHG-driven biodiversity

impacts embedded within the production, consumption and trade of food-related commodities. We

find that land-driven footprints differ depending on the metric used, and that GHG-driven

biodiversity impacts are driven largely by methane emissions and contribute a higher proportion of

the total footprint in regions of higher income and lower species-richness. The measures we present

are simple to calculate and could be incorporated into decision making and environmental impact

assessments by governments and businesses. Our consistent metric for biodiversity impacts allows

us to present multiple aspects of land-driven and GHG-driven biodiversity footprints, enabling

insight into priorities for reducing biodiversity costs via both global and regional production and

consumption.

Data availability

The EXIOBASE 3.8.1 pxp data that were used in our study are publicly available here:
https://zenodo.org/record/4588235#.ZGT8NnbMK3B

References

704   1   IPBES. Global assessment report on biodiversity and ecosystem services of the
705   Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. (Bonn,
706   Germany, 2019).
707   2   Maxwell, S. L., Fuller, R. A., Brooks, T. M. & Watson, J. E. M. The ravages of guns, nets and
708   bulldozers. *Nature* **536**, 143-145 (2016).
709   3   Schipper, A. M. *et al.* Projecting terrestrial biodiversity intactness with GLOBIO 4. *Glob Chang*
710   *Biol* **26**, 760-771 (2020). https://doi.org:10.1111/gcb.14848
711   4   Di Marco, M. *et al.* Projecting impacts of global climate and land-use scenarios on plant
712   biodiversity using compositional-turnover modelling. *Glob Chang Biol* **25**, 2763-2778 (2019).
713   https://doi.org:10.1111/gcb.14663
714   5   Newbold, T. Future effects of climate and land-use change on terrestrial vertebrate
715   community diversity under different scenarios. *Proc Biol Sci* **285** (2018).
716   https://doi.org:10.1098/rspb.2018.0792
717   6   Ellis, E. C., Klein Goldewijk, K., Siebert, S., Lightman, D. & Ramankutty, N. Anthropogenic
718   transformation of the biomes, 1700 to 2000. *Global Ecology and Biogeography*, no-no
719   (2010). https://doi.org:10.1111/j.1466-8238.2010.00540.x
720   7   FAOStat.   (License: CC BY-NC-SA 3.0 IGO., 2022).
721   8   Mbow, C. *et al.* Food Security. (2019).
722   9   Rosenzweig, C., N Tubiello, F., Sandalow, D., Benoit, P. & N Hayek, M. Finding and fixing food
723   system emissions: the double helix of science and policy. *Environmental Research Letters* **16**
724   (2021). https://doi.org:10.1088/1748-9326/ac0134
725   10   Crippa, M. *et al.* Food systems are responsible for a third of global anthropogenic GHG
726   emissions. *Nature Food* **2**, 198-209 (2021).
727   11   Leclere, D. *et al.* Bending the curve of terrestrial biodiversity needs an integrated strategy.
728   *Nature* **585**, 551-556 (2020). https://doi.org:10.1038/s41586-020-2705-y
729   12   Lynch, J., Cain, M., Frame, D. & Pierrehumbert, R. Agriculture's contribution to climate
730   change and role in mitigation is distinct from predominantly fossil CO2-emitting sectors.
731   *Frontiers in Sustainable Food Systems* **4**, 518039 (2021).
732   13   Kastner, T., Erb, K.-H. & Haberl, H. Rapid growth in agricultural trade: effects on global area
733   efficiency and the role of management. *Environmental Research Letters* **9**, 034015 (2014).
734   14   FAO. *The State of Agricultural Commodity Markets 2020. Agricultural markets and*
735   *sustainable development: Global value chains, smallholder farmers and digital innovations.*
736   (FAO, 2020).
737   15   Chaudhary, A. & Kastner, T. Land use biodiversity impacts embodied in international food
738   trade. *Global Environmental Change* **38**, 195-204 (2016).
739   https://doi.org:10.1016/j.gloenvcha.2016.03.013
740   16   Kitzes, J. *et al.* Consumption-Based Conservation Targeting: Linking Biodiversity Loss to
741   Upstream Demand through a Global Wildlife Footprint. *Conserv Lett* **10**, 531-538 (2017).
742   https://doi.org:10.1111/con4.12321
743   17   Balmford, A. *et al.* The environmental costs and benefits of high-yield farming. *Nature*
744   *Sustainability* **1**, 477-485 (2018). https://doi.org:10.1038/s41893-018-0138-5
745   18   Miller, R. E. & Blair, P. D. *Input-output Analysis: Foundations and Extension.* (Cambridge
746   University Press, 2009).
747   19   Kitzes, J. An introduction to environmentally-extended input-output analysis. *Resources* **2**,
748   489-503 (2013).
749   20   Roux, N., Kastner, T., Erb, K.-H. & Haberl, H. Does agricultural trade reduce pressure on land
750   ecosystems? Decomposing drivers of the embodied human appropriation of net primary
751   production. *Ecological Economics* **181** (2021).
752   https://doi.org:10.1016/j.ecolecon.2020.106915

21 Peters, G. & Hertwich, E. G. CO_2 embodied in international trade with implications for global
climate policy. *Environmental Science and Technology* **42**, 1401-1407 (2008).
22 Hoekstra, A. Y. & Mekonnen, M. M. The water footprint of humanity. *Proceedings of the*
National Academy of Sciences **109**, 3232-3237 (2012).
23 Lenzen, M. *et al.* International trade drives biodiversity threats in developing nations. *Nature*
486, 109-112 (2012).
24 Moran, D. & Kanemoto, K. Identifying species threat hotspots from global supply chains.
Nature Ecology and Evolution **1**, 0023 (2017).
25 Chaudhary, A., Verones, F., de Baan, L. & Hellweg, S. Quantifying Land Use Impacts on
Biodiversity: Combining Species-Area Models and Vulnerability Indicators. *Environ Sci*
Technol **49**, 9987-9995 (2015). https://doi.org:10.1021/acs.est.5b02507
26 Marques, A. *et al.* Increasing impacts of land use on biodiversity and carbon sequestration
driven by population and economic growth. *Nat Ecol Evol* **3**, 628-637 (2019).
https://doi.org/10.1038/s41559-019-0824-3
27 Wilting, H. C., Schipper, A. M., Bakkenes, M., Meijer, J. R. & Huijbregts, M. A. Quantifying
Biodiversity Losses Due to Human Consumption: A Global-Scale Footprint Analysis. *Environ*
Sci Technol **51**, 3298-3306 (2017). https://doi.org:10.1021/acs.est.6b05296
28 Alkemade, R. *et al.* GLOBIO3: A Framework to Investigate Options for Reducing Global
Terrestrial Biodiversity Loss. *Ecosystems* **12**, 374-390 (2009).
https://doi.org/10.1007/s10021-009-9229-5
29 Newbold, T., Oppenheimer, P., Etard, A. & Williams, J. J. Tropical and Mediterranean
biodiversity is disproportionately sensitive to land-use and climate change. *Nat Ecol Evol* **4**,
1630-1638 (2020). https://doi.org:10.1038/s41559-020-01303-0
30 Martins, I. S. & Pereira, H. M. Improving extinction projections across scales and habitats
using the countryside species-area relationship. *Sci Rep* **7**, 12899 (2017).
https://doi.org/10.1038/s41598-017-13059-y
31 Hillebrand, H. *et al.* Biodiversity change is uncoupled from species richness trends:
Consequences for conservation and monitoring. *Journal of Applied Ecology* **55**, 169-184
(2018). https://doi.org/10.1111/1365-2664.12959
32 Chichorro, F., Juslén, A. & Cardoso, P. A review of the relation between species traits and
extinction risk. *Biological Conservation* **237**, 220-229 (2019).
33 Stadler, K. *et al.* EXIOBASE 3: Developing a time series of detailed environmentally extended
multi-regional input-output tables. *Journal of Industrial Ecology*
https://doi.org/10.1111/jiec.12715 (2018).
34 Smith, P. & Gregory, P. J. Climate change and sustainable food production. *Proc Nutr Soc* **72**,
21-28 (2013). https://doi.org/10.1017/S0029665112002832
35 ESRI.   (2015).
36 The Nature Conservancy.   (2019).
37 International Food Policy Research Institute.   *MapSPAM* (ed International Food Policy
Research Institute) (Harvard Dataverse, 2019).
38 Monfreda, C., Ramankutty, N. & Foley, J. A. Farming the planet: 2. Geographic distribution of
crop areas, yields, physiological types, and net primary production in the year 2000. *Global*
Biogeochemical Cycles **22**, GB1022 (2008). https://doi.org:doi: 10.1029/2007GB002947
39 Ramankutty, N., Evan, A. T., Monfreda, C. & Foley, J. A. Farming the planet: 1. Geographic
distribution of global agricultural lands in the year 2000. *Global Biogeochemical Cycles* **22**,
GB1003 (2008). https://doi.org:doi:10.1029/2007GB002952
40 IUCN, I. U. f. C. o. N.   (2020).
41 BirdLife International.   (2020).
42 Meiri, S. *et al.*   (2017).

802	43	Roll, U. *et al.* The global distribution of tetrapods reveals a need for targeted reptile
803		conservation. *Nat Ecol Evol* **1**, 1677-1682 (2017). https://doi.org:10.1038/s41559-017-0332-
804		2
805	44	Etard, A., Morrill, S. & Newbold, T. Global gaps in trait data for terrestrial vertebrates. *Global*
806		*Ecology and Biogeography* **29**, 2143-2158 (2020).
807		https://doi.org:https://doi.org/10.1111/geb.13184
808	45	Hudson, L. N. *et al.* The database of the PREDICTS (Projecting Responses of Ecological
809		Diversity In Changing Terrestrial Systems) project. *Ecol Evol* **7**, 145-188 (2017).
810		https://doi.org:10.1002/ece3.2579
811	46	Hudson, L. N. *et al.* (ed Natural History Museum) (https://doi.org/10.5519/0066354, 2016).
812	47	Shine, K. P., Fuglestvedt, J. S., Hailemariam, K. & Stuber, N. Alternatives to the global
813		warming potential for comparing climate impacts of emissions of greenhouse gases. *Climate*
814		*Change* **68**, 281-302 (2005).
815	48	Myhre, G. *et al.* Anthropogenic and Natural Radiative Forcing. (Cambridge University Press,
816		Cambridge, United Kingdom and New York, NY, USA, 2013).
817	49	Hijmans, R., Cameron, S. E., Parra, J. L., Jones, P. G. & Jarvis, A. Very high resolution
818		interpolated climate surfaces for global land areas. *International Journal of Climatology* **25**,
819		1965-1978 (2005). https://doi.org:doi.org/10.1002/joc.1276
820	50	CountryEconomy.com.
821		*https://countryeconomy.com/demography/population/taiwan?year=2011* 2021).
822	51	World Bank. (2021).
823	52	Chaudhary, A., Carrasco, L. R. & Kastner, T. Linking national wood consumption with global
824		biodiversity and ecosystem service losses. *Sci Total Environ* **586**, 985-994 (2017).
825		https://doi.org:10.1016/j.scitotenv.2017.02.078
826	53	Sun, Z., Behrens, P., Tukker, A., Bruckner, M. & Scherer, L. Shared and environmentally just
827		responsibility for global biodiversity loss. *Ecological Economics* **194** (2022).
828		https://doi.org:10.1016/j.ecolecon.2022.107339
829	54	Cain, M. *et al.* Improved calculation of warming-equivalent emissions for short-lived climate
830		pollutants. *NPJ climate and atmospheric science* **2**, 29 (2019).
831	55	Kozicka, M. *et al.* Feeding climate and biodiversity goals with novel plant-based meat and
832		milk alternatives. *Nature Communications* **14**, 5316 (2023).
833	56	Dimbleby, H. National Food Strategy Independent Review: The Plan. (2021).
834	57	Ivanovich, C. C., Sun, T., Gordon, D. R. & Ocko, I. B. Future warming from global food
835		consumption. *Nature Climate Change* **13**, 297-302 (2023).
836	58	Fuchs, R., Brown, C. & Rounsevell, M. Europe's Green Deal offshores environmental damage
837		to other nations. *Nature* **586**, 671-673 (2020).
838	59	Mueller, N. D. *et al.* Closing yield gaps through nutrient and water management. *Nature* **490**,
839		254-257 (2012). https://doi.org:10.1038/nature11420. 490:254-257
840	60	Steen-Olsen, K., Owen, A., Hertwich, E. G. & Lenzen, M. Effects of sector aggregation on CO_2
841		multipliers in multiregional input-output analyses. *Economic Systems Research* **26**, 284-302
842		(2014).
843	61	Lenzen, M. Aggregation versus disaggregation in input-output analysis of the environment.
844		*Economic Systems Research* **23**, 73-89 (2011).
845	62	Merciai, S. & Heijungs, R. Balance issues in monetary input-output tables. *Ecological*
846		*Economics* **102**, 69-74 (2014).
847	63	de Koning, A. *et al.* Effect of aggregation and disaggregation on embodied material use of
848		products in input-output analysis. *Ecological Economics* **116**, 289-299 (2015).
849	64	Tukker, A. *et al.* Towards robust, authoritative assessments of environmental impacts
850		embodied in trade: Current state and recommendations. *Journal of Industrial Ecology* **22**
851		(2018).

65   Chandrakumar, C., McLaren, S., Malik, A., Ramilan, T. & Lenzen, M. Understanding New
Zealand's consumption-based greenhouse gas emissions: an application of multi-regional
input-output analysis. *The International Journal of Life Cycle Assessment* **25**, 1323-1332
(2020).
66   FAO. How to Feed the World in 2050. (Rome, Italy, 2009).
67   Oliveira, J. *et al.* Choosing pasture maps: An assessment of pasture land classification
definitions and a case study of Brazil. *International Journal of Applied Earth Observation and*
Geoinformation **93** (2020). https://doi.org:10.1016/j.jag.2020.102205